Microbiome-based correction for random errors in nutrient profiles derived from self-reported dietary assessments

Tong Wang[1], Yuanqing Fu[2,3,4], Menglei Shuai[2,3,5], Ju-Sheng Zheng[2,3,4], Lu Zhu[6], Andrew T. Chan[1,4,7,8,9], Qi Sun[1,10,11], Frank B. Hu[1,10,11], Scott T. Weiss[1], Yang-Yu Liu[1,12,]*

[1]*Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02115, USA*
[2]*School of Life Sciences, Westlake University, Hangzhou, China*
[3]*Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, China*
[4]*Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, China*
[5]*Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA*
[6]*Department of Epidemiology, University of Iowa College of Public Health, Iowa City, IA 52242, USA*
[7]*Clinical and Translational Epidemiology Unit, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA*
[8]*Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA*
[9]*Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA*
[10]*Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA*
[11]*Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA*
[12]*Center for Artificial Intelligence and Modeling, The Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA*
* Correspondence: yyl@channing.harvard.edu

Abstract

Since dietary intake is challenging to directly measure in large-scale cohort studies, we often rely on self-reported instruments (e.g., food frequency questionnaires, 24-hour recalls, and diet records) developed in nutritional epidemiology. Those self-reported instruments are prone to measurement errors, which can lead to inaccuracies in the calculation of nutrient profiles. Currently, few computational methods exist to address this problem. In the present study, we introduce a deep-learning approach --- **M**icrobiom**e**-based nu**t**rient p**r**of**i**le **c**orrector (METRIC), which leverages gut microbial compositions to correct random errors in self-reported dietary assessments using 24-hour recalls or diet records. We demonstrate the excellent performance of METRIC in minimizing the simulated random errors, particularly for nutrients metabolized by gut bacteria in both synthetic and three real-world datasets. Further research is warranted to examine the utility of METRIC to correct actual measurement errors in self-reported dietary assessment instruments.

Introduction

An unhealthy diet can increase the risk of many diseases[1,2,3,4]. For instance, excess intake of sugar or saturated fat could elevate the risk of coronary heart disease[5–8]. The investigation of the association between poor dietary habits and chronic diseases requires an accurate assessment of dietary intake in large population samples. Typically, epidemiologic studies rely on self-reported instruments such as food frequency questionnaires (FFQ)[9], Automated Self-Administered 24-hour Dietary Assessment Tool (ASA24)[10], and 7-Day Dietary Record (7DDR)[11], to collect diet data in large populations. However, self-reported instruments are subject to both random and systematic measurement errors[12–15]. While random errors are largely due to day-to-day individual variations in dietary intakes, systematic errors can be caused by under- or over-reporting and inaccurate estimation of portion sizes[16]. Measurement errors in the assessed food intake will be naturally propagated forward to the computation of the nutrient profile, causing inaccuracies in the assessed nutrient profile (Fig. 1a). Correcting such errors in the nutrient profile is crucial for improving the quality of nutritional epidemiology research. Several methods, such as regression calibrations[17,18] and cumulative averages[19] using repeated dietary assessments, have been used to correct for random errors of habitual diets in nutritional studies. These methods involve using a regression model to map the habitual dietary intake measured by a less accurate method (such as FFQ) to a more accurate habitual dietary measurement method (such as the average of 7DDRs). However, it is important to note that these methods are designed specifically to address measurement errors in habitual dietary intake and are incapable of correcting random measurement errors in single-day dietary assessments.

Signal reconstruction from corrupted or incomplete measurements is a common problem in various fields of research. For example, image denoising (i.e., removing noise from a noisy image to restore the true image) is an important topic in computer vision. The recent advance in deep neural networks has led to new methods that learn to map corrupted images to unobserved clean ones[20,21]. However, training those methods typically requires clean images, which are difficult to obtain in many cases. To overcome this challenge, a new class of methods that only leverages noisy images in training has been developed[22,23]. One noticeable example is Noise2Noise[22], which restores clean images by only being trained on corrupted ones. When the noise is random with a mean of zero, its performance is comparable and sometimes even better than other methods

trained using clean images[22]. The Noise2Noise's underlying principle involves training

the model on pairs of noisy images as both input and output. This strategy forces the

neural network to predict the average of the corrupted images, thereby statistically

converging the prediction of Noise2Noise towards the clean image due to the zero-mean

nature of the noise[22].

Inspired by the success of Noise2Noise[22], we hypothesize that we can correct the

random error in the assessed nutrient profile derived from the self-reported dietary

assessment without using clean data (i.e., the ground truth dietary intake). Note that we

focus on correcting the nutrient profile instead of the food profile (or the original dietary

assessment) because the prevalence of zero values in the food profile (i.e., no

consumption for many food items) renders a big challenge for machine learning tasks,

while the derived nutrient profile usually has non-zero values. Also, we focus on the

correction for random errors with zero means instead of systematic bias/errors with non-

zero means because effectively correcting the latter requires the ground truth dietary

intake, which is typically not available.

Our key idea is to incorporate gut microbial compositions as part of the input of a

deep-learning model and to infer the true nutrient profile based on the assessed nutrient

profile and the measured gut microbial composition (Fig. 1b). This idea is based on the

knowledge that many dietary constituents reaching the large intestine fuel the growth of

gut microbes[24–26]. One example is the growth of gut microbes enabled by anaerobic

fermentation of indigestible polysaccharides[24,25]. As a result, the gut microbial

composition is linked to dietary intake. Advances in sequencing technologies have made

it possible to determine the gut microbial composition quickly and accurately. It has been

shown that fecal bacteria and metabolites can be used as biomarkers for predicting

whether a few food items are introduced as dietary interventions (e.g., avocado) for

healthy adults in dietary intervention studies[27,28]. Hence, using objective microbial

biomarkers to infer dietary intake might complement the self-reported dietary assessment

and thus reduce measurement error in dietary intake. In this work, we developed a deep-

learning method: **M**icrobiom**e**-based nu**t**rient p**r**of**i**le **c**orrector (METRIC). We

demonstrated the effectiveness of METRIC in removing randomly added noise to both

synthetic and real data.

Results

Overview of METRIC

We aim to infer the true nutrient profile based on the assessed one and the measured gut microbial composition. A naive way to do this is to train a machine-learning model with the assessed nutrient profiles and microbial compositions as the input and the true nutrient profiles as the output. However, this is not feasible because such training requires the true nutrient profiles that are not easily available. To address this issue, we developed METRIC that does not rely on the true nutrient profile during its training but still can remove random errors in the assessed nutrient profile during testing (Fig. 2). During the training, we generated the corrupted nutrient profiles by adding random noise to the assessed nutrient profiles, and then trained METRIC to remove the added noise by taking the corrupted nutrient profiles and the measured microbial compositions as its input and generating the assessed nutrient profiles as its output (Fig. 2a). We introduced the corrupted nutrient profiles to avoid METRIC copying the assessed nutrient profiles directly to the output, thus forcing METRIC to remove the added noise. Note that the training of METRIC did not use true nutrient profiles.

The architecture of METRIC is a neural network that consists of three hidden layers, in addition to its input and output layers (Fig. 2b). Each hidden layer has a fixed dimension of 256. The link weights in the neural network are initialized using the Xavier Initialization. The training loss is the mean squared error. The predictive performance is assessed by the mean Pearson correlation coefficient between true and predicted nutrient concentrations averaged across all nutrients $\bar{\rho}$. One unique feature is the skip connection that adds the corrupted nutrient profiles directly to the final output of neural networks, which previously has been shown to enhance the training and predictive performance for deep neural networks[29]. Similar to the existence of the skip connection in Noise2Noise[22], the introduction of the skip connection enables the neural networks to adjust the prediction based on the corrupted nutrient profile, ensuring that output variables would not deviate too much from the corrupted nutrient profile. More details about the architecture can be found in the Methods section. Overall, METRIC is a generic "denoiser" that learns to remove any random noise added to the nutrient profile. With this generic ability to remove random noise, the well-trained METRIC should be able to remove random measurement errors by generating predictions closer to the true nutrient

profiles when it takes the assessed nutrient profiles and microbial compositions as its
input (Fig. 2c).

We split each dataset into two non-overlapping parts: a training and a test set.
METRIC was trained on the training set and then used to generate predictions for the test
set. The correction performance is measured by comparing the predicted nutrient profiles
with the "true" nutrient profiles in the test set (how we obtain the "true" nutrient profiles for
each scenario will be explained in separate sections below). To measure the predictive
performance of each nutrient, we adopted the Pearson Correlation Coefficient ρ between
its predicted and true corrected values.

METRIC can reduce measurement errors in nutrient profiles of synthetic data

We first validated METRIC using synthetic data for which we know the ground truth. We
used the Microbial Consumer-Resource Model (MiCRM)[30] to generate three types of data:
(1) *true* nutrient profiles (i.e., the ground-truth nutrient consumption), (2) *assessed*
nutrient profiles (i.e., the true nutrient profiles with random noise added to mimic
measurement errors), and (3) *corrupted* nutrient profiles (i.e., the assessed nutrient
profiles with artificially added random noise). MiCRM simulates the process of nutrient
consumption by microbes and the following microbial growth[30]. Note that the random
noise added to assessed and corrupted nutrient profiles are different. For simplicity, we
only considered the nutrient consumption in MiCRM and did not model the nutrient
production because most dietary nutrients cannot be produced by microbes. We created
different samples by randomly sampling nutrient fluxes and then running the community
assembly until we achieved a steady state. We consider sampled ground-truth nutrient
fluxes as true nutrient profiles and the steady-state microbial abundances as microbial
compositions. Gaussian noise $N(0, \sigma^2)$ with the mean of zero and standard deviation σ
is added to true nutrient profiles to create assessed nutrient profiles. More details about
MiCRM, the generation of synthetic data, and added noise can be found in the Methods
section.

We generated 250 samples by simulating the assembly process for 250
independent communities with 20 nutrients and 20 bacterial species. We trained METRIC
on 200 samples and tested it on the remaining 50 samples. We measured how assessed
values and corrected values (i.e., predicted values on the test set) respectively correlate
with true values. For each nutrient, we measured ρ between its corrected values from

predictions and its true concentration (denoted as ρ_c). Similarly, we calculated ρ between

its assessed concentrations and its true concentration (denoted as ρ_a). As the standard

deviation σ of the Gaussian noise increases, METRIC starts to correct the introduced

noise, represented by the switch from negative values of $(\rho_c - \rho_a)$ to positive values in

Fig. 3a. It is natural to expect that the correction of nutrient profiles with weaker noises

does not work ($\rho_c < \rho_a$ for small σ in Fig. 3a), because the correction is not necessary

when the measurement error is small (e.g., when $\rho_a > 0.8$). We focus on the case of $\sigma =$

1.5 from now on. The difference between the two metrics (i.e., $\rho_c - \rho_a$) reflects the

correction performance. The nutrient in Fig. 3d-e is slightly corrected ($\rho_c - \rho_a = 0.01$),

while the nutrient in Fig. 3f-g is strongly corrected ($\rho_c - \rho_a = 0.09$). Most nutrients have a

better alignment between their corrected values and true values (Fig. 3h).

METRIC mitigates measurement errors added to nutrient profiles in real data

Next, we tested METRIC on three real-world datasets. The first dataset, MCTS

(MiCrobiome dieT Study), comes from a unique study that investigated the influence of

diets on gut microbial composition[31]. It is unique because a large number of samples

(n=210) of paired nutrient profiles and microbial compositions were collected. The nutrient

profiles were calculated from ASA24 (see Methods). Different from the availability of true

nutrient profiles in synthetic data, the true nutrient profiles are not available for real data.

To deal with this issue, we treated the nutrient profiles derived from ASA24 as the "true"

nutrient profiles and added random noise (Gaussian noise with the mean of zero and

standard deviations σ) to them as "assessed" nutrient profiles. As σ increases, METRIC

starts to better correct the introduced noise (Fig. 4a). We focus on the case of $\sigma = 1.0$

from now on. We found that carotene has a large ρ_a and is not improved by METRIC

($\rho_a = 0.99$ versus $\rho_c = 0.97$ in Figs. 4b-c). By contrast, fiber has a small ρ_a and is strongly

improved ($\rho_a = 0.35$ versus $\rho_c = 0.58$ for Figs. 4f-g). We believe that the large correction

in the total fiber intake was due to most fibers being digested by gut microbes[24,25]. Overall,

nutrients with smaller ρ_a have better correction performance, and nutrients with large ρ_a

rarely improved (Fig. 4h). The mean correction performance averaged over all nutrients

is $(\overline{\rho_c} - \overline{\rho_a}) = 0.079$.

We also tried to run METRIC without using the microbial composition, finding that

the correction performance ($\overline{\rho_c} - \overline{\rho_a} = 0.067$) is worse than that with included microbial

composition (Supplementary Fig. 1). We also computed the sensitivity which is defined

as the ratio between the reduction in ρ of nutrient α and the perturbation amount of

species i (Supplementary Fig. 2), which could detect some possible interactions between

nutrients and species. For example, the sensitivity of monounsaturated fatty acids

towards *Bacteroides uniformis* is large (~0.025, larger than 99.98% of the inferred

sensitivity values), and is supported by the previously observed reduction of

monounsaturated fatty acids by *Bacteroides uniformis*[32]. In addition, our sensitivity

analysis revealed that the top three microbial taxa linked to fiber content correction, which

exhibit the highest sensitivity values, are well-documented as fiber degraders in the

literature: *Bacteroides plebeius*[33], *Parabacteroides sp*[34], and *Bacteroides sp*[35].

Then, we applied METRIC to the second dataset MLVS (Men's Lifestyle

Validation Study)[36,37]. Specifically, we utilized the composition of gut microbiomes and

the one-day dietary assessment of the 7-day dietary records (7DDRs). The 7DDRs are

widely recognized to be the most reliable estimation of dietary intake because participants

are required to measure and report gram weights for foods both before they start eating

and after they finish, thereby enabling the calculation of the actual food consumption

based on the difference in weight[38]. To guarantee the usefulness of gut microbial

composition in correcting dietary assessment, we required the identification of paired

microbial compositions and dietary assessments with matching dates. In MLVS, a total

of 599 paired samples with matching dates were found. Similarly, as we did for the MCTS

dataset, we regarded the nutrient profile derived from the 7DDRs as the "true" nutrient

profile and added varying levels of Gaussian noise to it as the "assessed" nutrient profile.

Then we trained METRIC on 80% of the data and tested it on the remaining 20%.

Consistent with our previous findings, the trained METRIC exhibits an ability to correct

the nutrient profile (Fig. 5a), especially for large σ. For the case of $\sigma = 1.0$, the mean

correction performance ($\overline{\rho_c} - \overline{\rho_a}$) is 0.072 (Fig. 5h). Across all nutrients, dietary fiber was

the strongest corrected nutrient (Figs. 5f-g).

Finally, we applied METRIC to the third dataset WE-MACNUTR (Westlake N-of-

1 Trials for Macronutrient Intake)[39]. WE-MACNUTR is a dietary intervention study that

implemented a 'complete feeding' strategy, providing three isocaloric meals per day to

28 participants over a span of 72 days. Each participant completed high-fat, low-

carbohydrate and low-fat, high-carbohydrate diets in a randomized sequence, with a 6-

day wash-out period between them. Since the diets were completely controlled and well-

documented, the nutrient profile derived from this dataset closely reflects the true nutrient

profile. In WE-MACNUTR, we found 317 paired samples with both microbial compositions

and nutrient profiles. Considering the nutrient profile from the complete feeding as the

true nutrient profile, we introduced varying levels of noise (Gaussian noise $N(0, \sigma^2)$ with

different standard deviations σ) to create the assessed nutrient profile. Like our earlier

results, METRIC can correct the nutrient profile when σ is large (Fig. 6a). When $\sigma = 1.0$,

the mean correction performance ($\overline{\rho_c} - \overline{\rho_a}$) is 0.118 (Fig. 6h) and dietary fiber again

exhibits a substantial correction (Figs. 6f-g).

To provide a more representative picture of METRIC's robustness and

effectiveness across different noise levels, we also investigated situations where the

correction is less effective for all three datasets (standard deviation of the noise $\sigma = 0.5$;

Supplementary Figs. 3-5). Although the overall correction performance ($\bar{\rho}_c - \bar{\rho}_a$) is weak

when $\sigma = 0.5$, the correction still works well for nutrients with smaller ρ_a (e.g., dietary

fibers). We also evaluated the predictive performance using a more quantitative metric,

the mean absolute error. We found that the overall pattern in correction performance

measured by mean absolute error aligns with that measured by Pearson correlation

across datasets (Supplementary Figs. 6-8). For the MLVS and WE-MACNUTR datasets,

we also tried to evaluate METRIC's correction performance without using the microbial

composition, finding that the correction performance is comparable to that achieved when

the microbial composition is included (Supplementary Figs. 9&10). This implies that

METRIC can still be leveraged to correct nutrient profiles even in the absence of gut

microbial compositions.

We capitalized on the longitudinal nature of the MCTS dataset to explore whether

increasing temporal offsets between microbiome and diet data impacts the correction

efficiency of our method. Specifically, we increased the offset by aligning the diet of day

t with the microbiome of day $t + \Delta t$ and subsequently correcting nutrient profiles. Our

analysis reveals that the correction performance progressively decreases as the offset

Δt deviates from 1 day (Supplementary Fig. 11). This serves as a validation of METRIC,

as it indicates that microbiome-diet relationships are causal.

Given the absence of ground-truth nutrient profiles for direct validation, we instead

conducted an indirect validation of our method by checking if the noise level of real-life

nutrient profiles is within the regime where we can remove the random measurement

errors well. We found that the correction performance of METRIC is great when the mean

Pearson correlation coefficient $\bar{\rho}_a$ is below 0.8 (Figs. 4-6). Due to the lack of ground-truth

nutrient profiles to directly quantify the noise level, we can only approximate this indirectly

by reflecting the nutrient variability using the multiple-day 7DDRs in MLVS. Specifically,

we calculated the Pearson correlation coefficient ρ between concentrations of a nutrient
derived from one 7DDR and its average values obtained from multiple 7DDRs for seven
consecutive days (Supplementary Fig. 12a). We found that the mean Pearson correlation
coefficient $\bar{\rho}$ is 0.77, which is below 0.8. Additionally, 62.0% of 329 nutrients have $\rho < 0.8$.
A similar analysis on the dataset MCTS revealed that $\bar{\rho} = 0.63$ and 95.0% of nutrients
have $\rho < 0.8$ (Supplementary Fig. 12b). The WE-MACNUTR dataset was not analyzed
due to the absence of dietary assessments. These findings across both datasets confirm
that the approximated noise levels are within a range where METRIC is effective at
correcting random measurement errors.

We also examined the impact of noise with a non-zero mean by introducing the
Gaussian noise $N(\mu, \sigma^2)$. For the three datasets we used, we set $\sigma = 1$ as this is the
regime where our method, METRIC, consistently shows strong correction performance.
We then gradually increased the mean of the noise μ from 0.0 to 2.0 to create the
assessed nutrient profile. When applying METRIC to remove the noise, we observed that
its correction performance diminishes to zero as μ increases (Supplementary Figs. 13-
15), indicating that METRIC can remove the random error but not the systematic drift or
shift.

Discussion

We presented a deep-learning method, METRIC, to correct random simulated
measurement errors in the nutrient profile. The method relies on the assessed nutrient
profile and microbial composition to learn how to infer the true nutrient profile. First, we
validated its performance on synthetic data where we directly modeled true and assessed
nutrient profiles. Then we applied METRIC to three distinct real clinical datasets with
added noise to the nutrient profile and found that it can correct the nutrient profile well,
especially for nutrients with large errors or metabolized by gut microbes. Similar to the
class of computational methods that denoise images even without clean targets[22,23],
METRIC offers a significant advantage by being capable of correcting the random
measurement error without using the true nutrient profiles during the training. This
attribute makes our method particularly useful in real-life scenarios where only assessed
nutrient profiles are available, without access to true nutrient profiles.

We admit that METRIC has several limitations. First, without using clean targets
(i.e., the ground truth dietary intake), METRIC is only capable of removing random

measurement errors that have zero means. It cannot remove noise with a non-zero mean,

just like Noise2Noise[22]. Thus, METRIC cannot correct the systematic bias/drift/error (with

a non-zero mean) in nutrient profiles. Effectively correcting the systematic bias requires

both assessed and true nutrient profiles to discern the consistent deviation between them.

However, it is very unlikely that both assessed and true nutrient profiles are available to

measure the real systematic bias in real-world data. If both types of nutrient profiles are

available, it remains to be investigated whether it is possible to design a regression model

that predicts true nutrient profiles based on assessed nutrient profiles, thereby achieving

the task of removing the systematic bias. Second, METRIC uses microbial compositions

of fecal samples to correct nutrient profiles derived from ASA24 or 7DDR, which were

collected exactly one day prior to fecal sample collection. We do not expect that microbial

compositions of fecal samples collected from a particular time point could be used to

correct nutrient profiles derived from FFQ, where subjects report how often each food

item was consumed over a specified period, typically the past month or year. To clearly

demonstrate this point, we leveraged the FFQ, 7DDR, and gut microbiome data from

MLVS to compare the performance of using gut microbial compositions to predict the

7DDR- or FFQ-based nutrient profiles. We found that the predictability of FFQ-based

nutrient profiles is much worse than that of 7DDR-based nutrient profiles (Supplementary

Fig. 16). Consequently, we believe that METRIC cannot be used to correct random errors

in FFQ-based nutrient profiles.

We recognize the challenges of applying a model trained on one dataset to

another, particularly when biases like sequencing errors or differences in nutrient

databases are present. However, training the method on one dataset and generating

predictions for the other is possible if both datasets were collected and processed in the

same way. For example, in the PRISM and NLIBD studies, two Inflammatory Bowel

Disease cohorts—one from Boston (n=155) and an external validation cohort from the

Netherlands (n=65)—were collected using the same protocols. This standardization

enables accurate predictions of disease status in the NLIBD cohort by a machine learning

method trained on the PRISM data[40]. Similarly, the deep-learning model mNODE trained

on PRISM data showed great performance in predicting fecal metabolome based on gut

microbial compositions from the NLIBD cohort[41].

There are other methods to improve the accuracy of dietary intake measurement

over traditional self-reported dietary assessments. For instance, digital documentation of

meals through taking photos can be used to improve the accuracy of dietary assessment,

though the validity of such technologies is yet to be established. In addition, nutritional

biomarkers such as DNA barcodes for plants[42] and metabolite biomarkers[28,43,44] have

been utilized to improve the assessment of food intake. A more accurate reflection of

nutrient consumption can be obtained by using objective measurements of microbial

composition or metabolomic profile, which can complement self-reported dietary

assessment tools. Currently, an active research direction in the field of precision nutrition

is to identify microbial and metabolite biomarkers for dietary intake. Although several

studies attempted to predict the presence of food items based on fecal bacteria and

metabolites[27,28], the analyses were limited to several food items and no connection to

nutrient profiles was examined. Further research is warranted to examine the utility of

METRIC to correct actual measurement errors in self-reported dietary assessments.

Methods

Datasets. The first dataset we used comes from a study that investigated the association

between diet and the gut microbiome[31]. The study has paired 24-h food records and fecal

shotgun metagenomes from 34 healthy human subjects collected daily over 17 days, with

210 paired samples in total. The second dataset is from MLVS[36,37] with 599 paired gut

microbial compositions and 7DDRs. The third dataset is from WE-MACNUTR[39], a dietary

intervention study that provides three meals per day for all participants. The study

involved 30 participants, with 2 participants withdrawing from the trial early and not

included in the final analysis. It has 317 paired gut microbial compositions and true

nutrient profiles of complete feeding with matching dates. For all machine learning tasks,

the same five random 80/20 train-test splits were utilized to guarantee a fair comparison

of methods.

The generation of nutrient profiles based on dietary assessments. The generation

of nutrient profiles generated from dietary assessments involves several steps and relies

on comprehensive food composition databases such as USDA's Food and Nutrient

Database for Dietary Studies (FNDDS)[45] or Harvard Food Composition Database

(HFDB)[46], etc. Each food item consumed is matched with a corresponding item in a food

composition database. These databases provide detailed information on the nutrient

content of food items, including macronutrients (e.g., fats, proteins, carbohydrates) and

micronutrients (e.g., vitamins and minerals). By multiplying the amount of each food

consumed by its nutrient content, the total intake of each nutrient is calculated to obtain

the nutrient profiles. For example, the MCTS dataset utilized ASA24 to collect dietary

intake data, where documented foods were assigned to nutrients based on the USDA's

FNDDS 2011-2012[45]. In the case of the MLVS dataset, nutrient intake was derived from

7DDRs using the Nutrition Data System for Research[47], which provides a comprehensive

food and nutrient database managed by the University of Minnesota Nutrition

Coordinating Center. For the WE-MACNUTR dataset, which focuses on a Chinese

population, nutrient profiles were derived from the foods directly provided to participants

using the electronic version of China Food Composition Tables (Standard Edition)[48].

The generation of synthetic data. Similar to the traditional consumer-resource

model[49,50] and Microbial Consumer-Resource Model[30], we simulated the process of

nutrient consumption by microbes and the following microbial growth. For simplicity, we

assumed a global pool with 20 microbial species and 20 nutrients in total. To mimic the

difference in nutrient consumption profiles across individuals, we randomly assigned flux

for each sample: to create a unique sample, the supply rate of each nutrient is randomly

drawn from a uniform distribution $\mathcal{U}[0, 1]$. We recorded the assigned supply rates of all

nutrients as the true nutrient profile. The Gaussian noise with a mean of 0 and standard

deviation of 0.5 is added to true nutrient profile to create assessed nutrient profiles with

measurement errors. After the assignment of nutrient fluxes, we simulated the community

assembly dynamics until the system reaches a steady state. The steady-state microbial

relative abundances are recorded as the microbial composition for the sample. Then we

repeated this procedure of generating samples 250 times, forming the synthetic dataset

we used in this study. The 250 samples are split in the 80/20 ratio as the training/test set.

The overall population dynamics for the concentration of nutrient N_α and

abundance of microbial species M_i can be written as follows:

400
$$\frac{dN_\alpha}{dt} = -\sum_i a_{i\alpha} M_i N_\alpha - \delta N_\alpha,$$

401
$$\frac{dM_i}{dt} = \frac{\sum_\gamma a_{i\gamma} M_i N_\gamma}{Y} - \delta M_i,$$

where $a_{i\alpha}$ is the consumption rate of nutrient α by the species i, δ is the dilution rate, and

Y is the yield. For simplicity, we assumed $\delta = 0.1$ and $Y = 1$. The $a_{i\alpha}$ is assumed to be

non-zero with a probability of 50%. If $a_{i\alpha}$ is non-zero, its value is drawn from the uniform

distribution $\mathcal{U}[0, 10]$. Eventually, each $a_{i\alpha}$ is divided by the number of nutrients that can

be consumed by the species i, avoiding the outgrowth of generalists.

METRIC. The core of METRIC is the MLP (Multilayer Perceptron).

- 408 • Data processing: The Centered Log-Ratio transformation is applied to microbial
- 409   relative abundances, and the log transformation is applied to the nutrient profiles.
- 410 • Model detail: METRIC has 3 hidden layers in the middle, sandwiched by input and
- 411   output variables. Each hidden layer has a fixed hidden layer dimension of 256.
- 412   The Xavier Initialization is used to initialize the weights in the neural network. A
- 413   skip connection from corrupted nutrient profiles is introduced to add to the final
- 414   layer of MLP. Specifically, the final prediction is the sum of (1) the corrupted input
- 415   from the skip connection multiplied by a weight parameter α and (2) the final layer
- 416   of MLP multiplied by $(1 - \alpha)$. The optimal value for α is chosen based on the five-
- 417   fold cross-validation results on the training set.
- 418 • Training method: The Adam optimizer[51] is used for the gradient descent. The
- 419   training loss is the mean squared error. The training stops when the mean
- 420   Pearson Correlation Coefficient of nutrients $\bar{\rho}$ on the validation/test set starts to
- 421   decrease within the past 10 epochs.
- 422 • Activation function: ReLU (Rectified Linear Unit).
- 423 **Statistics.** To calculate correlation throughout the study, we used Pearson's correlation
- 424 coefficient. All simulations and analyses were performed using standard numerical and
- 425 scientific computing libraries such as NumPy and SciPy in the Python programming
- 426 language (version 3.7.1) and Jupyter Notebook (version 6.1).
- 427
- 428 **Data availability.** We only used the data collected by existing studies. Instructions for
- 429 downloading sequencing data and dietary intakes analyzed in this work can be found in
- 430 the literature exploring MCTS (MiCrobiome dieT Study)[31], MLVS (Men's Lifestyle
- 431 Validation Study)[36,37], and WE-MACNUTR (Westlake N-of-1 Trials for Macronutrient
- 432 Intake)[39]. To facilitate the data downloading, the URLs to those datasets are also
- 433 provided in our GitHub repository (https://github.com/wt1005203/METRIC).
- 434
- 435 **Code availability.** All code for simulations used in this manuscript can be found at
- 436 https://github.com/wt1005203/METRIC.
- 437

- 438 **Acknowledgements.** We acknowledge grants from the National Institutes of Health
- 439 (R01AI141529, R01HD093761, R35CA253185, RF1AG067744, UH3OD023268,
- 440 U19AI095219, U01HL089856, U01-152905, and U01-167552) and Cancer Grand

Challenges Team PROSPECT. We thank Walter Willett, Eric Rimm, and Lorelei Mucci

for valuable discussions.

Author contributions. T.W. and Y.-Y.L. designed the project. T.W. performed all the

numerical calculations and data analysis. T.W. processed the real data with assistance

from Y.Fu. All authors analyzed the results. T.W. and Y.-Y.L. wrote the paper. All authors

edited and approved the paper.

Correspondence. Correspondence and requests for materials should be addressed to

Y.-Y. L. (email: yyl@channing.harvard.edu).

Competing Interests. The authors declare no competing interests.

References

1.   Afshin, A. *et al.* Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet* **393**, 1958–1972 (2019).

2.   Mozaffarian, D. *et al.* Global Sodium Consumption and Death from Cardiovascular Causes. *New England Journal of Medicine* **371**, 624–634 (2014).

3.   Hu, F. B. Diet strategies for promoting healthy aging and longevity: An epidemiological perspective. *Journal of Internal Medicine* **n/a**,.

4.   Donaldson, M. S. Nutrition and cancer: A review of the evidence for an anti-cancer diet. *Nutrition Journal* **3**, 19 (2004).

5.   Hu, F. B. & Willett, W. C. Optimal Diets for Prevention of Coronary Heart Disease. *JAMA* **288**, 2569–2578 (2002).

6.   Mensink, R. P., Zock, P. L., Kester, A. D. & Katan, M. B. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. *The American Journal of Clinical Nutrition* **77**, 1146–1155 (2003).

7.   DiNicolantonio, J. J., Lucan, S. C. & O'Keefe, J. H. The Evidence for Saturated Fat and for Sugar Related to Coronary Heart Disease. *Progress in Cardiovascular Diseases* **58**, 464–472 (2016).

8.   Malik, V. S. & Hu, F. B. The role of sugar-sweetened beverages in the global epidemics of obesity and chronic diseases. *Nat Rev Endocrinol* **18**, 205–218 (2022).

9.   McNutt, S., Zimmerman, T. P. & Hull, S. G. Development of food composition databases for food frequency questionnaires (FFQ). *Journal of Food Composition and Analysis* **21**, S20–S26 (2008).

10.   Sharpe, I. *et al.* Automated Self-Administered 24-H Dietary Assessment Tool (ASA24) recalls for parent proxy-reporting of children's intake (> 4 years of age): a feasibility study. *Pilot and Feasibility Studies* **7**, 123 (2021).

11.   Hebert, J. R. *et al.* Development and testing of a seven-day dietary recall. *Journal of Clinical Epidemiology* **50**, 925–937 (1997).

12.   Westerterp, K. R. & Goris, A. H. C. Validity of the assessment of dietary intake: problems of misreporting. *Current Opinion in Clinical Nutrition & Metabolic Care* **5**, 489 (2002).

13.   Poslusna, K., Ruprich, J., Vries, J. H. M. de, Jakubikova, M. & Veer, P. van't. Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *British Journal of Nutrition* **101**, S73–S85 (2009).

14.   Dao, M. C. *et al.* Dietary assessment toolkits: an overview. *Public Health Nutrition* **22**, 404–418 (2019).

15.   Ravelli, M. N. & Schoeller, D. A. Traditional Self-Reported Dietary Instruments Are Prone to Inaccuracies and New Approaches Are Needed. *Frontiers in Nutrition* **7**, (2020).

16.   Willett, W. Correction for the Effects of Measurement Error. in *Nutritional Epidemiology* (ed. Willett, W.) 0 (Oxford University Press, 2012). doi:10.1093/acprof:oso/9780199754038.003.0012.

17.   Rosner, B., Willett, W. C. & Spiegelman, D. Correction of logistic regression relative risk estimates and confidence intervals for systematic within-person measurement error. *Statistics in Medicine* **8**, 1051–1069 (1989).

18.   Spiegelman, D., McDermott, A. & Rosner, B. Regression calibration method for correcting measurement-error bias in nutritional epidemiology. *The American Journal of Clinical Nutrition* **65**, 1179S-1186S (1997).

19.   Hu, F. B. *et al.* Dietary Fat and Coronary Heart Disease: A Comparison of
Approaches for Adjusting for Total Energy Intake and Modeling Repeated Dietary
Measurements. *American Journal of Epidemiology* **149**, 531–540 (1999).
20.   Schmidt, U. & Roth, S. Shrinkage Fields for Effective Image Restoration. in
2774–2781 (2014).
21.   Chen, Y. & Pock, T. Trainable Nonlinear Reaction Diffusion: A Flexible
Framework for Fast and Effective Image Restoration. *IEEE Transactions on Pattern
Analysis and Machine Intelligence* **39**, 1256–1272 (2017).
22.   Lehtinen, J. *et al.* Noise2Noise: Learning Image Restoration without Clean Data.
in *Proceedings of the 35th International Conference on Machine Learning* 2965–2974
(PMLR, 2018).
23.   Krull, A., Buchholz, T.-O. & Jug, F. Noise2Void - Learning Denoising From
Single Noisy Images. in 2129–2137 (2019).
24.   Donia, M. S. & Fischbach, M. A. Small molecules from the human microbiota.
Science **349**, 1254766 (2015).
25.   Koh, A., De Vadder, F., Kovatcheva-Datchary, P. & Bäckhed, F. From Dietary
Fiber to Host Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites. *Cell*
165, 1332–1345 (2016).
26.   Santhiravel, S. *et al.* The Impact of Plant Phytochemicals on the Gut Microbiota
of Humans for a Balanced Life. *International Journal of Molecular Sciences* **23**, 8124
(2022).
27.   Shinn, L. M. *et al.* Fecal Bacteria as Biomarkers for Predicting Food Intake in
Healthy Adults. *The Journal of Nutrition* **151**, 423–433 (2021).
28.   Shinn, L. M. *et al.* Fecal Metabolites as Biomarkers for Predicting Food Intake
by Healthy Adults. *The Journal of Nutrition* nxac195 (2022) doi:10.1093/jn/nxac195.
29.   He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image
Recognition. in *2016 IEEE Conference on Computer Vision and Pattern Recognition
(CVPR)* 770–778 (IEEE, Las Vegas, NV, USA, 2016). doi:10.1109/CVPR.2016.90.
30.   Marsland, R. *et al.* Available energy fluxes drive a transition in the diversity,
stability, and functional structure of microbial communities. *PLoS Comput Biol* **15**,
e1006793 (2019).
31.   Johnson, A. J. *et al.* Daily Sampling Reveals Personalized Diet-Microbiome
Associations in Humans. *Cell Host & Microbe* **25**, 789-802.e5 (2019).
32.   López-Almela, I. *et al.* Bacteroides uniformis combined with fiber amplifies
metabolic and immune benefits in obese mice. *Gut Microbes* **13**, 1–20 (2021).
33.   Um, C. Y. *et al.* Grain, Gluten, and Dietary Fiber Intake Influence Gut Microbial
Diversity: Data from the Food and Microbiome Longitudinal Investigation. *Cancer
Research Communications* **3**, 43–53 (2023).
34.   Wei, W. *et al.* Parabacteroides distasonis uses dietary inulin to suppress NASH
via its metabolite pentadecanoic acid. *Nat Microbiol* **8**, 1534–1548 (2023).
35.   Yang, J. *et al.* High Soluble Fiber Promotes Colorectal Tumorigenesis Through
Modulating Gut Microbiota and Metabolites in Mice. *Gastroenterology* **166**, 323-337.e7
(2024).
36.   Mehta, R. S. *et al.* Stability of the human faecal microbiome in a cohort of adult
men. *Nat Microbiol* **3**, 347–355 (2018).
37.   Li, J. *et al.* Interplay between diet and gut microbiome, and circulating
concentrations of trimethylamine N-oxide: findings from a longitudinal cohort of US
men. *Gut* **71**, 724–733 (2022).
38.   Yue, Y. *et al.* Reproducibility and validity of diet quality scores derived from
food-frequency questionnaires. *The American Journal of Clinical Nutrition* **115**, 843–853
(2022).

39.   Ma, Y. *et al.* Individual Postprandial Glycemic Responses to Diet in n-of-1 Trials:
Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR). *The Journal of*
Nutrition **151**, 3158–3167 (2021).
40.   Franzosa, E. A. *et al.* Gut microbiome structure and metabolic activity in
inflammatory bowel disease. *Nat Microbiol* **4**, 293–305 (2019).
41.   Wang, T. *et al.* Predicting metabolomic profiles from microbial composition
through neural ordinary differential equations. *Nat Mach Intell* **5**, 284–293 (2023).
42.   Reese, A. T. *et al.* Using DNA Metabarcoding To Evaluate the Plant Component
of Human Diets: a Proof of Concept. *mSystems* **4**, e00458-19 (2019).
43.   Guasch-Ferré, M., Bhupathiraju, S. N. & Hu, F. B. Use of Metabolomics in
Improving Assessment of Dietary Intake. *Clinical Chemistry* **64**, 82–98 (2018).
44.   Brennan, L., Hu, F. B. & Sun, Q. Metabolomics Meets Nutritional Epidemiology:
Harnessing the Potential in Metabolomics Data. *Metabolites* **11**, 709 (2021).
45.   FNDDS : USDA ARS. https://www.ars.usda.gov/northeast-area/beltsville-md-
bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-
group/docs/fndds/#.
46.   Colditz, G. A., Manson, J. E. & Hankinson, S. E. The Nurses' Health Study: 20-
Year Contribution to the Understanding of Health Among Women. *Journal of Women's*
Health **6**, 49–62 (1997).
47.   Feskanich, D., Sielaff, B. H., Chong, K. & Buzzard, I. M. Computerized collection
and analysis of dietary intake information. *Computer Methods and Programs in*
Biomedicine **30**, 47–57 (1989).
48.   Yang, Y. & Wang, Z. China food composition tables standard edition. *Peking*
University Medical Press: Beijing, China (2018).
49.   MacArthur, R. Species packing and competitive equilibrium for many species.
Theoretical Population Biology **1**, 1–11 (1970).
50.   Chesson, P. MacArthur's consumer-resource model. *Theoretical Population*
Biology **37**, 26–38 (1990).
51.   Kingma, D. P. & Ba, J. Adam: A Method for Stochastic Optimization. Preprint at
https://doi.org/10.48550/arXiv.1412.6980 (2017).
585
586

587
Figure 1: **The idea of inferring the true nutrient profile based on the assessed**
nutrient profile and microbial composition. a, The typical dietary assessment such as
ASA24 (Automated Self-Administered 24-hour Dietary Assessment Tool) often induces
a measurement error in the food intake due to unreliable mental recall. Such a random
measurement error or systematic error/bias in the food intake is carried to the nutrient
profile when the food profile is converted to the nutrient profile. **b,** Our goal in this study
is to remove the random part of measurement error. More specifically, we would like to
infer true nutrient profiles based on assessed nutrient profiles and microbial compositions.

Figure 2: **The architecture and workflow of METRIC to infer true nutrient profiles.**
For simplicity, we used a hypothetical example with n=3 training samples and 2 samples
in the test set. For each sample, there are N_s microbial species and N_n nutrients. Across
panels, microbial species and their relative abundances are colored blue. Nutrients and
their amounts are colored red. The corrupted nutrient profiles are created by adding
different types of random noise (i.e., Gaussian, Uniform, etc.) to the assessed nutrient
profiles. Icons associated with assessed/corrupted nutrient profiles are bounded by solid
black/dashed lines. Icons associated with true nutrient profiles are bounded by solid
green lines. **a,** During the training of METRIC, the method takes corrupted nutrient
profiles and microbial compositions as the input and learns to infer assessed nutrient
profiles. **b,** Similar to multilayer perceptrons, METRIC has several hidden layers in the
middle. The skip connection provides the corrupted nutrient profile directly to one layer
before the final output, enabling it to skip the propagation through the hidden layers. The
skipped corrupted nutrient profile multiplied by the weight parameter α and the final
hidden layer (the bottom grey nodes) multiplied by $(1 - \alpha)$ add up as the final output (the
bottom red nodes). **c,** The well-trained METRIC is applied to the test set to generate
predictions for nutrient profiles whose values are compared to true nutrient profiles.

Figure 3: **METRIC can correct the measurement error in assessed nutrient profiles**
on synthetic data from MiCRM[30]**.** The Pearson's Rank Correlation Coefficient ρ is
adopted to evaluate the correlation across various types of nutrient profiles. All
corrected/true values shown are the log of nutrient concentrations. **a,** ρ_c (i.e., ρ between
corrected and true values) and ρ_a (i.e., ρ between assessed and true values) decrease
as the standard deviation of added Gaussian noise σ increases. All following panels
focus on the case of σ=1.5. **b,** The correlation between assessed values and true values
of log concentrations of one nutrient among different samples. **c,** The correlation between
corrected values (predictions of METRIC) and true values of log concentrations of the
same nutrient among different samples. Similar comparisons for another two nutrients
are shown in **d-e** and **f-g**. **e,** The correction performance of all nutrients is measured by
$(\rho_c - \rho_a)$.

Figure 4: **METRIC can correct the measurement error in assessed nutrient profiles**
on real data from MCTS[31]**.** The Pearson's Rank Correlation Coefficient ρ is adopted to
evaluate the correlation across various types of nutrient profiles. All nutrient
concentrations are in the unit of grams. All corrected/true values shown are the log of
nutrient concentrations. **a,** ρ_c (i.e., ρ between corrected and true values) and ρ_a (i.e., ρ
between assessed and true values) decrease as the standard deviation of added
Gaussian noise σ increases. All following panels focus on the case of σ=1.0. **b,** The
correlation between assessed values and true values of log concentrations of carotene
among different samples. **c,** The correlation between corrected values (predictions of
METRIC) and true values of log concentrations of carotene among different samples. **d-**
e, The similar comparison for octadecanoic acid shows a modest correction. **f-g,** The
similar comparison for fiber shows a strong correction. **h,** The correction performance for
all nutrients is measured by ($\rho_c - \rho_a$).

Figure 5: **METRIC can correct the measurement error in assessed nutrient profiles from MLVS**[36,37]**.** The Pearson's Rank Correlation Coefficient ρ is adopted to evaluate the correlation across various types of nutrient profiles. All nutrient concentrations are in the unit of grams. All corrected/true values shown are the log of nutrient concentrations. **a,** ρ_c (i.e., ρ between corrected and true values) and ρ_a (i.e., ρ between assessed and true values) decrease as the standard deviation of added Gaussian noise σ increases. All following panels focus on the case of σ=1.0. **b,** The correlation between assessed values and true values of log concentrations of fructose among different samples. **c,** The correlation between corrected values (predictions of METRIC) and true values of log concentrations of fructose among different samples. **d-e,** The similar comparison for monounsaturated fatty acids shows a modest correction. **f-g,** The similar comparison for dietary fiber shows a strong correction. **h,** The correction performance for all nutrients is measured by ($\rho_c - \rho_a$). Nutrient names are not added due to lack of space.

658
659

Figure 6: **METRIC can correct the measurement error in assessed nutrient profiles from WE-MACNUTR**[39]**.** The Pearson's Rank Correlation Coefficient ρ is adopted to evaluate the correlation across various types of nutrient profiles. **a,** ρ_c (i.e., ρ between corrected and true values) and ρ_a (i.e., ρ between assessed and true values) decrease as the standard deviation of added Gaussian noise σ increases. All nutrient concentrations are in the unit of grams. All corrected/true values shown are the log of nutrient concentrations. All following panels focus on the case of σ=1.0. **b,** The correlation between assessed values and true values of log concentrations of magnesium among different samples. **c,** The correlation between corrected values (predictions of METRIC) and true values of log concentrations of magnesium among different samples. **d-e,** The similar comparison for threonine shows a modest correction. **f-g,** The similar comparison

671 for dietary fiber shows a strong correction. **h,** The correction performance for all nutrients
672 is measured by $(\rho_c - \rho_a)$. Nutrient names are not added due to lack of space.