Effects of deafening on vocal production learning

in the Egyptian fruit-bat

Julie E. Elie[1*], Sandra E. Muroy[2,3], Daria Genzel[1], Tong Na[4], Lisa A. Beyer[5], Donald L. Swiderski[5],

Yehoash Raphael[5], Michael M. Yartsev[1,6*]

[1]Department of Bioengineering, University of California, Berkeley; Berkeley, CA 94708, USA.

[2]Department of Molecular and Cell Biology, University of California, Berkeley; Berkeley, CA 94708, USA.

[3]Department of Integrative Biology, University of California, Berkeley; Berkeley, CA 94708, USA.

[4]Department of Electrical Engineering and Computer Sciences, University of California, Berkeley;

Berkeley, CA 94708, USA.

[5]Department of Otolaryngology–Head and Neck Surgery, Michigan Medicine, Kresge Hearing Research

Institute; University of Michigan, 1150 Medical Center Drive, Ann Arbor, MI, 48109-5616, USA.

[6]Helen Wills Neuroscience Institute, University of California, Berkeley; Berkeley, CA 94708, USA.

* Corresponding authors: Julie E Elie, julie.elie@berkeley.edu; Michael M Yartsev,

myartsev@berkeley.edu

Abstract

Some species have evolved the ability to use the sense of hearing to modify existing vocalizations, or even create new ones. This ability corresponds to various forms of vocal production learning that are all possessed by humans, and independently displayed by distantly related species. Among mammals, a few species, including the Egyptian fruit-bat, would possess such vocal production learning abilities. Yet the necessity of an intact auditory system for the development of the Egyptian fruit-bat typical vocal repertoire has not been tested. Here we addressed this gap by eliminating pups' sense of hearing at birth and assessing its effects on vocal production in adulthood. The deafening treatment enabled us to both causally test these bats vocal learning ability and discern learned from innate aspects of their vocalizations. Leveraging wireless individual audio recordings from freely interacting adults, we show that a subset of the Egyptian fruit-bat vocal repertoire necessitates auditory feedback. Intriguingly, these affected vocalizations belong to different acoustic groups in the vocal repertoire of males and females. These findings open the possibilities for targeted studies of the mammalian neural circuits that enable sexually dimorphic forms of vocal learning.

Significance

Vocal production learning is the rare capacity amongst animals where hearing is used to modify or create new vocalizations. The Egyptian fruit-bat is believed to possess this capacity, yet whether they need audition to achieve a mature vocal repertoire is unknown. Furthermore, a systematic causal examination of learned and innate aspects of the entire repertoire has never been performed in a vocal learner. Here, we addressed these major gaps directly by abolishing hearing in Egyptian fruit-bats at birth and investigating its effects on the vocal production in adulthood. Leveraging simultaneous individual wireless audio-recordings from freely interacting

42   adult bats, we identify the subset of learned vocalizations and provide evidence that vocal learning

43   is sexually dimorphic in that species.

45   **Introduction**

Vocalizations constitute the building blocks of the acoustic code used for vocal

communication. The efficiency and complexity of a vocal communication system depends on

the species ability to produce a diversity of such building blocks. While this diversity can be

supported by hard wired motor programs that are innate (e.g., motor programs of male mouse

song[1]), the sound diversity is also enhanced in some species by their ability to modify existing

calls or create new ones[2–7]. This ability is defined as vocal production learning and has been

examined in a wide variety of species[2,4,8]. Yet, experiments discerning innate versus learned

vocalizations in the entire repertoire of a single species are missing.

The Egyptian fruit-bat has been proposed as one of the species capable of vocal

production learning after several studies demonstrated how manipulation of either the social or

physical environment affects the acoustic characteristics of their communication calls[9–11].

However, it remains unknown whether all or only a subset of their repertoire is acquired through

learning. Furthermore, the acoustic characteristics of the voice of individuals can easily be

mistaken for an effect of vocal learning when comparing the vocal behaviors of two groups of

animals (one control and one experimental). Previous conclusions on vocal learning abilities are

weakened by the lack of reliable identification of vocalizing individuals[9–11]. Combined, these

gaps limit thorough exploration of the neural mechanisms of vocal production learning in this

species, although cutting-edge technologies for measuring brain activity are readily available[12–

15]

Here we reasoned that because vocal production learning heavily depends on the sense

of hearing, the main sensory feedback that the nervous system can use to change or create

new motor programs[16], abolishing this feedback during development can begin revealing what

aspects of the repertoire are learned and which are innate. Therefore, we conducted the first

causal study of vocal production learning in the Egyptian fruit-bat by establishing a method for

deafening pups shortly after birth and testing the impact of such manipulation on the vocal

behavior of each of these bats when they became adults.

Results

Deafening at birth and recording of vocalizations at adulthood.

The acoustic isolation of newborn pups to any bat vocalization was ensured by single-

housing pregnant mothers before pup birth in acoustically isolated chambers. Deafening of the

pups was then chemically achieved shortly after birth (Figure 1A; n=10 bats). Five pups

received an injection regimen of the deafening antibiotic kanamycin, while five pups received

control saline injections (Methods). In each experimental group of five, there were three females

and two males. The extent of hearing loss incurred was verified both functionally, using auditory

brainstem recordings (Figure 1B, Supplementary Figure 1; Methods), and histologically, by

examining the cochlear epithelium (Figure 1C, Supplementary Figure 1; Methods). These

approaches revealed profound deafness in the kanamycin treated bats and normal hearing in

the saline treated bats.

Because the ontogeny and maturation of the Egyptian fruit-bat vocal behavior is

unknown, we focused our investigation of the vocalizations of hearing and deaf bats when they

were fully mature adults (Figure 1A and 1D, Methods). By then, the bats had lived for years in a

communal social environment where they were equally exposed to conspecific vocalizations.

Over a seven week period, the deaf and hearing adults were co-housed and daily equipped with

wireless audiologger collars, that enabled to detect all vocalizations from the bat repertoire and

to unambiguously assign the vocalizer identity to each vocalization[13]. Additionally, an ambient

microphone further ensured high quality audio recordings of the vocalizations. After manual

curation, we obtained thousands of calls from both hearing (n=16,937) and deaf bats

(n=11,154).

No effect of deafening on the whole repertoire when controlling for identity.

Deafening did not affect the vocal activity of bats (Supplementary Figure 2) and we

obtained a similar number of vocalizations from both deafened and hearing bats (>1,000

vocalizations on average per individual). To explore if deafening was affecting the acoustic

features of vocalizations, we used both a classical bioacoustical approach (28 predefined

acoustic features measured on each call) and a data driven approach (Figure 1D). In the latter

approach, we used the Modulation Power Spectrum[17] (MPS) to describe the spectro-temporal

modulations of vocalizations. The resulting MPS of each vocalization was used both in UMAP

projections of all vocalizations and in regularized linear classifiers that quantified the level of

discrimination between calls emitted by deaf and hearing bats (Regularized Permutation

Discriminant Function Analysis, RPDFA, see Methods). Unexpectedly, with regards to previous

reports[9–11], when we considered all the recorded vocalizations, none of the two approaches

revealed any effect of deafness when the identity of the vocalizing bats was accounted for

(Supplementary Figure 3, all 28 linear mixed-effects models on acoustic features with p>0.05).

The performance of the RPDFA at classifying call MPS between deaf and hearing bats was

60.4% ± 1% and not significantly different from the chance level of classifying any random two

groups of five bats, 62.4% ± 1.3% (Supplementary Figure 4A and 4B; note that considering the

voice characteristics of individuals raises the chance level higher than 50%).

Heterogeneity in vocal space coverage according to sex and hearing ability.

However, we next considered the possibility that not all vocalizations necessarily had to

depend on auditory feedback. That is, some could be innate whereas others could be learned.

Therefore, considering all vocalizations together, as was done in previous studies, may result in

masking the effect of deafening. Indeed, linear measures of acoustic proximity revealed that

bats were not homogeneously occupying the vocal space (Figure 2). First, we found that males

covered a larger vocal space than females (vocalizations with large values of Sex Dissimilarity;

Figure 2A and Supplementary Figure 5A, 5B and 5D) and that for each sex, some vocalizations

were distinct between deaf and hearing bats (Supplementary figure 6A-6C; performance of the

RPDFA against identity controlled chance level: all male calls, 67.4 ± 1.2% vs 64.5 ± 0.8%; all

female calls, 62.0 ± 0.6% vs 58.6 ± 1.3%). Second, a small proportion of the repertoire of each

sex was idiosyncratic to the hearing ability of bats (Figure 2B and 2D). In males, these

distinctive vocalizations represented 23.9 ± 3.2% of hearing bats repertoire and could not be

matched with similar calls from deaf bats, indicating that hearing males produced vocalizations

that deaf males did not (Figure 2B, Supplementary Figure 7). In females, the few distinctive

vocalizations represented 4.5 ± 1.5% of deaf bats repertoire and did not have a match with

vocalizations produced by hearing females, indicating that these deaf females were producing a

few atypical vocalizations (Figure 2D, Supplementary Figure 7). These vocalizations belonged

to regions of the vocal space that grouped the loudest, more tonal, and higher pitch

vocalizations in the repertoire (Supplementary Figure 8).

To further quantify and identify the spectro-temporal differences between deaf and

hearing bats' vocalizations throughout the repertoire, we parsed the vocal space into four

groups (Figure 2C, Supplementary Figure 9). This exact number of groups optimized both

maximum division and cluster quality (Supplementary Figure 10). Deafness affected

vocalizations in two of the four groups (Supplementary Figure 4; performance of the RPDFA

against the identity-controlled chance level: Group 1, 73.2 ± 3.7% vs 66.8 ± 3.2%; Group 4, 85.9

± 1.8% vs 76.9 ± 2.5%). Group 4 corresponded to loud and high pitch vocalizations that females

rarely produced (Figure 2A, Supplementary Figure 5C, 8 and 9) and only male calls were

different between hearing and deaf individuals in this acoustic group (Supplementary figure 6G-

6I; performance of the RPDFA against identity controlled chance level: Male Group 4 calls, 89.5

± 2.7% vs 79.2 ± 2.1%; Female Group 4, 62 ±10.4% vs 59.7 ± 9.2%). Surprisingly, although

both sexes produced calls that belonged to acoustic Group 1 (Supplementary Figure 5C), only

female calls were distinct between hearing and deaf bats (Supplementary figure 6D-6F;

performance of the RPDFA against identity controlled chance level: Male Group 1 calls, 69.9 ±

5.4% vs 65.7 ± 3.9%; Female Group 1, 76.4 ± 1.9% vs 61.2 ± 3.1%). Combined, these findings

suggest that a subset of the Egyptian fruit bat repertoire depended on intact auditory feedback

in a sexually dimorphic manner.

Effect of deafening on the spectro-temporal patterns of a subset of male and

female calls.

The acoustic differences in the vocalizations of deaf and hearing bats were a complex

combination of both spectral and temporal parameters. The atypical vocalizations from deaf

females in acoustic group1 were characterized by complex patterns of pitch modulations and

decreased formants power (Supplementary figure 11). However, these differences were not

captured by the classical bioacoustical approach (Supplementary figure 12). In contrast, the

pattern of spectro-temporal modulations that characterized hearing male calls from acoustic

group 4 was clear. Hearing males produced simple downsweep calls that were never matched

by deaf males who produced calls with more upsweeps (Figure 3, Supplementary figure 13).

Furthermore, deaf male calls had higher spectral content (i.e. more harmonics) than hearing

male calls as measured by spectral features such as the third spectral quartile (Figure 3,

Supplementary figure 13).

Discussion

In this study, we assessed the necessity of hearing for the development of a normal

vocal repertoire in Egyptian fruit bats. The acoustic analysis demonstrated that the repertoire of

this species consists of a large majority of innate vocalizations (>75% for males and >95% for

females). Auditory input is not necessary to establish, fine-tune or maintain the neural motor

programs controlling the production of these calls. However, some vocal features were not

produced by males that lacked auditory input (downsweep calls with filtered higher harmonics)

and atypical spectro-temporal patterns emerged in a few calls from deaf females, indicating that

neural motor programs for these calls might be plastic and require an intact auditory input to

properly mature. According to the source-filter theory that was first developed to explain human

voice production[18,19] and then applied to animal vocalizations[20–22], the upsweeps and

downsweeps of the fundamental frequency are controlled by the vocal source (the larynx), while

the filtering of call harmonics and formants are achieved by change in the shape of the vocal

tract (e.g., vocalizing with open or closed mouth). Under this theory, the results of the present

acoustic analyses support the view that male Egyptian fruit-bats need to hear themselves,

and/or other bats, in order to properly control both their larynx and upper vocal tract to produce

the downsweep calls. The hypothesis of a fine motor control of the larynx is consistent with our

recent findings demonstrating a direct monosynaptic pathway from the vocal-motor cortex to the

brainstem center innervating the larynx[12]. These later findings are, so far, the strongest

evidence of the existence of this direct connection in mammals, which has been suggested as a

central anatomical hallmark of the neural circuitry that enables vocal production learning[23–27].

Previous experiments, such as pups reared in social isolation, or manipulation of the

acoustic environment of both young and adults, had shown effects when considering together

all vocalizations in the Egyptian fruit bat repertoire[9–11]. Deafening would also affect all the

vocalizations in the adult repertoire of the pale spear-nosed bat, *Phyllostomus discolor*[28]. We

cannot exclude that in our experiment, deaf bats could have compensated for their lack of

hearing by using somatosensation as a sensory feedback and social responses of their hearing

companions as guiding cues to produce most of the vocalizations in their repertoire.

Somatosensation is indeed a feedback that is used along with audition to guide vocal production

in humans and songbirds alike[29–31]. Nonetheless, even after years of communal living, males

could not properly produce a significant proportion of their vocal repertoire in the absence of

auditory feedback. These results pave the way to targeted studies of these male Egyptian fruit

bat vocalizations, with the possibility for comparing the neural mechanisms of production

between learned and innate vocalizations in a mammal brain. Furthermore, the observed

difference between males and females opens new opportunities to investigate how the sex of an

individual might interfere with the maturation of vocal behavior in a mammalian brain that is

particularly suitable for neurophysiological research.

Methods

Subjects and housing

All experimental procedures were approved by the Institutional Animal Care and Use Committee

of the University of California, Berkeley. Nine wild-caught adult female Egyptian fruit-bats

(*Rousettus Aegyptiacus*) and their ten pups (five females and five males) were used for this

study. Each female gave birth to one pup except for one female that gave birth twice. The

progenitor female bats were imported from Israel and housed in our lab colony at the University

of California Berkeley for at least six months before being included in the experiments. Female

bats were isolated in soundproof booths when they were highly pregnant (individual cage size:

10x12x20 inches). This protocol has been previously used to ensure that pups would not be

exposed to any vocalization including their mother's vocalizations[10]. Indeed, an isolated mother

does not vocalize even in presence of its pup[10]. Each female was maintained with its pup in

their soundproof booth until the end of the deafening protocol (see below) of the pup. After the

end of the injection protocol (post-natal day 56), the young bats were moved to a collective

housing room and housed together with their mothers and male bats from the colony in larger

cages (21 x 32 x 35.8 inches). Each cage contained one saline pup and one kanamycin pup

with their respective mothers and two adult males. At sexual maturity (7 months old), the young

bats were separated from their mothers and housed in single sex cages with age matched

unmanipulated bats.

When reaching 3-4 years of age, all ten experimental bats were grouped together in a large

cage (180x60x60 cm) and isolated from the rest of the colony in a soundproof room for two

months over which their vocalizations were recorded as detailed below.

In all cages, bats were maintained on a reverse day-light cycle set at 12h/12h. Recordings of

vocalizations were conducted during the dark period.

Deafening protocol

Half of the pups followed a treatment by kanamycin (powder from Alfa Aesar diluted in PBS, pH

7.4, Gibco 10010-02B) while the other half were kept as control subjects and injected with saline

solution (PBS, pH 7.4, Gibco 10010-02B). Starting on post-natal day 32 (PND32) and for 21

days, the pups were subcutaneously injected twice daily (8 hrs apart). The dosage of kanamycin

was set to increase stepwise over the 21 days (6 days at 300 mg/kg, 5 days at 400 mg/kg, 5

days at 500 mg/kg and 5 days at 600 mg/kg). The weight of the young bats was measured at

most 4 days before a change of dosage or at least on the eve of a change of dosage to correctly

adjust dosage. Control subjects were injected with the same volume of saline solution as the

volume they would have been injected with if they were kanamycin subjects.

Audiogram

Eight months after the end of the acoustic recordings, the audiogram of each experimental bat
was established by measuring their auditory brain response (ABR) to tone pips of 2.5 ms (Hann
window) spanning the auditory range expected for that species (1.5 - 90kHz, 25 logarithmically
spaced levels of frequencies) and presented at various levels of intensity (30-110 dB SPL, 9
steps of 10dB SPL), following the same protocol as Linnenschmidt and Wiegrebe (2019)[32].
Briefly, bats were isolated and fasted from 1h and up to 2h before inducing anesthesia by
subcutaneous injection of a mix of Ketamine (22 mg/kg), DexaMedetomidine (0.09 mg/kg) and
Midazolam (0.45 mg/kg). Once the bat was deeply anesthetized, it was placed laying on its
stomach on a heating pad in a soundproof room, facing the loudspeaker (Scanspeak
R2604/833000) at 5 cm, with an oxygen delivery tube placed right under its nose (flow rate
~300mL/min). The animal temperature and breathing rate were monitored every 15 min during
the procedure. The ABR was measured as the difference of potential between a signal
subdermal electrode inserted at the base of the head in the neck of the animal and a reference
subdermal electrode inserted subcutaneously atop of the head between the two ears. The
whole system was grounded by an alligator clip placed on the lower part of one ear. The
electrode signals were amplified by a TDT headstage (RA4Li, Tucker Davis Technologies Inc.,
Fl, USA) and further amplified and digitized by an audio interface (Fireface UC, RME,
Haimhausen, Germany; sampling rate 192 kHz). After bandpass filtering between 100 Hz and 3
kHz and downsampling 10 times, the electrophysiological signals were stored for offline
analysis. The 225 (nine amplitude levels and 25 frequency steps) tone pips were each
presented 256 times at a repetition rate of 44Hz. Using previously used matlab algorithm[32], tone
pips were generated at 192kHz, digitally equalized in magnitude and phase to compensate for
the loudspeaker impulse response, converted to analog using the same audio interface as the

one used for processing the ABR signal, and amplified using one channel of an audio amplifier

(AVR 347, Harman Kardon, Stamford, CT, USA).

For each of the 256 presentations of a tone pip, the ABR signal was extracted in two

windows of 7 ms, one aligned to the tone pips (response window) and the second one randomly

placed between 0 and 23ms after stimulus onset (background noise window). From these 256

pairs of snippets of signal, we generated one tone response (the average electrophysiological

signal over all response windows) and one background response (the average

electrophysiological signal over all background windows) and measured the RMS of each. We

bootstrap the calculation of the RMS background response for each stimulus by repeating 1000

times the selection of random background noise windows in the 256 ABR time series. The exact

p-value of obtaining a tone response RMS higher than the background response RMS values

was then calculated as the number of background response RMS values equal or higher than

the tone response RMS for that stimulus. We could then draw the audiogram for each bat as the

lowest sound amplitude, for each carrier frequency tested, at which the RMS of the ABR signal

was significant at $p=0.05$ (see Supplementary Figure 14).

Histology

Cochlear extraction

After ABR measurements, the bats still under anesthesia were euthanized by a lethal

intraperitoneal injection of sodium pentobarbital. The body was then transcardially perfused with

200mL of phosphate buffer saline (PBS) solution (pH=7.4) spiked with 0.5mL of Heparin (1,000

USP units/mL) followed by 250 mL of fixative (3.7% formaldehyde PBS solution, pH=7.4). The

temporal bones were removed from the skull and the inner ears isolated. The oval and round

window were opened to facilitate flow of fixative solution for rapid fixation. Inner ear tissues were

shipped to the University of Michigan for histological analyses.

The cochleae with their surrounding bone were decalcified in 5% EDTA until the bone was soft

to facilitate dissection of the sensory area, the organ of Corti. The tissues were rinsed in PBS

and further dissection was done to remove the bone, stria vascularis and tectorial membrane.

Phalloidin staining

The cochleae were permeabilized with 0.3% Triton X-100 in PBS for 15 minutes followed by

incubation with Alexa Fluor 488 phalloidin (Invitrogen A12379) for 40 minutes, all at room

temperature. After rinsing with PBS, the cochleae were further dissected into segments of the

basal turn of the cochlea (higher frequencies area) and the apical turn (lower frequencies).

Segments were then mounted on slides with Prolong Gold (Invitrogen P36930). The slides were

stored in the dark at 4°C until viewed with fluorescence microscopy.

Imaging of the organ of Corti

The samples were imaged with a Leica DMRB epifluorescence microscope. Images were

captured using a CCD cooled SPOT-RT monochrome digital camera. Digital images were

processed with Adobe Photoshop.

<u>Audio recordings</u>

When the animals reached 3-4 years old, the ten experimental bats were recorded for 35 days

over the course of seven weeks, for 5-7 hours per day during their night cycle. For acclimation

purposes, the bats were all housed in the recording cage starting 16 days before the first day of

audio recording and were housed in that recording cage for the entire duration of the recording

period (7 weeks). The cage was a rectangular prism (180 x 60 x 60 cm) that had two sides

made of plexiglass, thereby permitting clear remote visual monitoring of bats behavior via two

cameras (Flea 3 FLIR). The remaining sides of the enclosure were made of plastic mesh,

allowing bats to easily perch and crawl on the surfaces. The enclosure was placed in an

acoustically shielded room and all recording sessions were conducted during the dark cycle

under red LED light. All bats were equipped daily with audio data-loggers to record and identify

their vocalizations. Water was given *ad libitum* and the daily diet supply was placed into the

cage at the beginning of each recording session to increase the diversity of behavioral contexts.

The audio recordings of each individual bat vocalizations were performed using both an audio

data-logger and an ambient ultrasonic microphone (Earthworks, M50) centered 20 cm above

the cage ceiling and connected to the main computer unit via an analog to digital convertor

(MOTU, 896mk3). The audio collected from the ambient microphone was recorded throughout

the session (sampling rate of 192 kHz) using an in-house Matlab GUI (VocOperant;

https://github.com/julieelie/operant_bats). As previously described (Rose et al 2021) the audio

data-logger consists in a single-axis, low mass, piezo-ceramic accelerometers (BU-27135,

Knowles Electronics, sensitivity 0-10kHz) that is mounted on a flexible rubber necklace placed

against the throat of the subject, in a way that does not restrict normal behavior but enables the

detection of laryngeal vibrations produced during vocalizations. The signal of the accelerometer

is recorded, digitized at a sampling rate of 50 Hz, and saved on removable SD cards with a

wireless audio data logging device ("audiologgers"; Audio Logger AL1 or AL2, Deuteron

Technologies) mounted on the necklace on the back of the subject. The audiologger and its

lithium-polymer battery are encapsulated in a house-made 3D-printed plastic casing to prevent

damage to the electronics. All audiologgers were controlled and synchronized by a single

transceiver. The Egyptian fruit bats in our experiment weighed more than 110 gr and could fly

with ease while equipped with their audiologger. The synchronization between the microphone

recording and the transceiver controlling the audiologgers was achieved using transistor-

transistor logic (TTL) pulses generated by an UltraSoundGate Player 216H (Avisoft

Bioacoustics) and sent via coaxial cables. The experimenter could monitor the behavior of the

animals in an ante-chamber via the video cameras and the ambient microphone.

Vocalizations detection and acoustic analysis

From all the days of recording, we sampled 13 days for acoustical analysis. These days were

chosen such as to optimize their spread across the 8 weeks of recording and optimize the

recording conditions (no audiologger breaking incident).

Vocalization detection

Acoustic data logged as voltage traces on the SD cards of the audiologgers were extracted into

Matlab files and aligned across bats simultaneously recorded using a custom-made Matlab

code (https://github.com/NeuroBatLab/LoggerDataProcessing/). Potential vocalizations were

then detected and segmented from these piezo recordings using an in-house series of Matlab

scripts (https://github.com/NeuroBatLab/SoundAnalysisBats). The whole process consisted in

three major steps: 1) detection of sound events on audiologger signals, 2) automatic

classification between vocalizations and noise, 3) manual curation of potential vocalizations.

Supplementary Figure 15 presents a full flow chart of the approach taken for the detection of

vocalizations in audiologger recordings.

1. Detection: To focus on the detection of vocalizations emitted by the bat wearing the

collar, the signal of each audiologger was first band-passed between 1 and 5kHz. As

previously described, this frequency range is not contaminated by airborne vocalizations

from other bats standing close to the collar wearer (Rose et al, 2021). After determining

a noise threshold from sections of silence during the recording session for that

audiologger (three times the RMS of silence sections), potential vocalizations were

detected by threshold crossing on the amplitude envelope (RMS, sampling frequency

1kHz) and any sound event above threshold for longer than 7ms was kept.

2. Automatic step of data sorting between actual vocalizations and noise: Sound events

closer than 50ms were merged as a single sound sequence and a battery of 20 acoustic

measurements (pre-defined acoustic features, PAFs) were applied on them (see Elie

and Theunissen, 2016, for mathematical definitions): RMS, maximum amplitude, the five

first momentum of the amplitude envelope taken as a distribution (mean, standard

deviation, kurtosis, skewness, and entropy), the five first momentum of the frequency

spectrum taken as a distribution, the three quartiles of the frequency spectrum, the mean

pitch saliency, and four parameters that pertain to the sound as recorded from the

ambient microphone (RMS, mean and maximum of the amplitude envelope, and

maximum value of cross-correlation between the microphone and the audiologger

signals). Potential vocalizations among the detected sound events were then identified

using these 20 acoustic parameters as input to a support vector machine (SVM) trained

on the data of two sessions manually sorted between vocalizations and noise by an

expert (JEE). This data was obtained from a total of 100 min of recording of 8 bats (800

min of audiologger data) and contained 24,633 sound elements, of which 1,377 were

vocalizations. The automatic sorting was set to be very conservative of vocalizations by

setting the threshold on the posterior probability of a vocalization at 0.002, with the

drawback of yielding many false discoveries (percentage of vocalizations correctly

identified in 10-fold cross validation: 98.3%; percentage of noise miss-classified as

vocalizations in 10-fold cross validation: 7.5%).

After the manual curation (see below) of a reasonable dataset (five days), an extra step

of automatic detection could be added to the pipeline to further filter out sequences

devoid of vocalization. We trained a neural network on the manual annotations obtained

from these five days of recording (4,936 vocalizations). The network was implemented

using the PyTorch library and the architecture was composed of ResNet-50 with a

sigmoid output layer. For each 100ms non-overlapping window of the audiologger signal,

391	we bandpass filtered the signal between 100Hz and 5kHz, and calculated its
392	spectrogram using *soundsig* from the BioSound library
393	(https://github.com/theunissenlab/soundsig) with the following parameters: sampling
394	rate=1kHz; frequency spacing=100Hz; dynamic range=50dB. The spectrogram was then
395	colorized using the color scale in *soundsig*. The training label for each 100ms window
396	was 1 if there existed a vocalization in that window and 0 otherwise. The 3-channel color
397	spectrograms and corresponding binary labels were used as inputs for the network. To
398	take into account the large imbalance between noise and vocalization elements in the
399	dataset, each training batch was set to contain 5 noise elements for each vocalization
400	element. We used a weighted binary cross-entropy loss function with penalty=7.5 to
401	further address the imbalance within each training batch. 20 percent of the data was
402	held out for validation. We randomly initialized the network weights and trained the
403	network for 10 epochs using the Adam optimizer with learning rate=1e-3. A sound
404	sequence on a given audiologger was considered as worthy of manual examination if
405	the maximum of all posterior probability values (one value per 100ms window) was
406	higher than 0.9. At this threshold of maximum posterior probability value, the trained
407	neural network achieved high performance of correct classification of sequences
408	containing vocalizations (98.6%) with low values of false discovery (5.4% of noise
409	sequences detected as containing vocalizations) enabling to greatly speed up the
410	vocalization discovery procedure. The data from 8 more days of recording were
411	evaluated by the trained neural network.
412	
413	3. Manual curation: to further eliminate noise and check the identity of the bat producing
414	vocalizations, each potential vocalization was aurally and visually scrutinized by trained
415	students and an expert (JEE) based on the inspection of the spectrograms of its signal
416	as recorded from the ambient microphone and from the audiologgers of all the bats

(*manual annotation* in Supplementary figure 15). Vocal utterances had to have a

minimum duration of 7ms to be considered as vocalizations and those spaced by less

than 50ms were grouped together. A final step of audio quality evaluation performed by

an expert eliminated vocalizations that were contaminated by other vocalizations or

noise on the microphone recording.

Acoustic features and modulation power spectrum (MPS)

28 predefined acoustic features (PAFs) were chosen to describe both the temporal and the

spectral content of vocalizations. For each vocalization, PAFs were calculated from both the

audiologger recording and from the ambient microphone recording. The signal captured by the

piezo sensor of the audiologger was particularly suitable to measure pitch or amplitude related

acoustic features because it is consistently placed very close to the larynx of the animal, the

vocal source. The PAFs calculated from the audiologger signal were: duration, pitch saliency (a

measure of how tonal vs noisy a sound is), fundamental frequency, coefficient of variation (CV)

of the fundamental frequency, the five first momentum of the sound temporal envelope taken as

a distribution (mean, standard deviation, kurtosis, skewness and entropy), the amplitude

periodicity frequency (frequency peak in the amplitude power spectrum), the amplitude

periodicity power index (ratio of the power at the amplitude periodicity frequency with the signal

RMS), the five first momentum of the sound power spectrum taken as a distribution (mean,

standard deviation, kurtosis, skewness and entropy) and the first, second and third spectral

quartiles. The PAFs calculated from the microphone signal were: the five first momentum of the

sound spectrum taken as a distribution (mean, standard deviation, kurtosis, skewness and

entropy) and the first, second and third spectral quartiles. All these parameters except the two

amplitude periodicity measurements were calculated using *soundsig* from the BioSound Python

library (https://github.com/theunissenlab/soundsig). The two amplitude periodicity

measurements were calculated using custom Matlab code.

The modulation power spectrum (MPS) of each vocalization was calculated from the

microphone signal using *soundsig* from the BioSound Python library

(https://github.com/theunissenlab/soundsig). The MPS is the 2D fast Fourier transform of the

sound spectrogram. To ensure that each vocalization MPS could be calculated at the same time

resolution, we ensured that each vocalization was isolated within a 100 ms window. After

bandpassing the microphone signal between 0.3 kHz and 50 kHz, the spectrogram was

calculated at a 1 kHz sampling rate using a frequency spacing of 100Hz. The spectrogram was

then z-scored and the MPS calculated using a window bin of 100ms. The MPS range was

focused on values with the most energy: the range of spectro-temporal modulations was

restricted from 0 to 3 cycles/kHz for the frequency modulations and from -150 Hz to 150 Hz for

the temporal modulations (see example MPS in Supplementary Figure 9). To account for the

various dynamic range across the spectro-temporal modulations, the MPS of each vocalization

was normalized by dividing with the average MPS over all vocalizations in the dataset to obtain

the Normalized Modulation Power Spectrum (NMPS, Supplementary Figure 9).

UMAP projection and acoustic clustering

The NMPS of the 28,091 vocalizations were reduced to fewer dimensions by applying the

algorithm UMAP (Uniform manifold approximation and projection) setting the number of

neighbors to 10, the minimum distance to 0.051 and the target weight to 0.55 (Matlab code

provided by the Herzenberg Lab at Stanford University, CA, USA). The NMPS as represented

by the UMAP values were then used as input to a hierarchical agglomerative clustering (HAC)

algorithm (*linkage* function of Matlab using the Ward method and the Euclidean metric) to divide

the acoustic space covered by the vocalizations into groups. The optimal number of dimensions

for the UMAP, the optimal tree cut and optimal number of acoustic groups were found by

measuring the average and median silhouette value over UMAPs of various dimensions and

various levels of tree cutting in the HAC (Supplementary Figure 10). Four acoustic groups in a

3D UMAP were found to be optimal numbers.

Vocal dispersion

Vocal dispersion is calculated for each individual bat as the generalized variance of its

vocalizations using the first 100 dimensions of the PCA on the NMPS of vocalizations from the

whole dataset. The first 100 dimensions explained 90% of the variance. The result depicted on

Supplementary Figure 5A, showing a larger vocal dispersion for males as compared to females,

was robust across all the number of dimensions tested (from 10 to 100 with a step of 10 and

from 100 to 300 with a step of 20).

Acoustic dissimilarity (distance to opposite sex and distance to opposite hearing ability)

The dissimilarity was calculated for each vocalization emitted by bat i as the distance to the

vocalizations from the nearest target bats j ($j \neq i$) in the space of the principal component

analysis (PCA) calculated on NMPS of vocalizations (Figure 2). Dimensions of the PCA had to

explain at least 1% of the variance to be included in this measure to ensure that the measure

would not be noise sensitive. As such the first 5 dimensions were retained. To represent how

males and females differentially occupy the acoustic space (Figure 2A), the target bats were the

opposite sex bats. To represent which vocalizations were affected by deafening (Figure 2B and

2D), the target bats were the opposite treatment but same sex bats. For each vocalization, the

acoustic dissimilarity is the distance to the closest of the target bats beyond what would be

expected given the local density of bats vocalizations. The distance from one particular

vocalization of bat i to the vocalizations of a j target bat was calculated as the average

Euclidean distance to the nearest 10 vocalizations of target bat j. The local density of bat

vocalizations (null distance) was the distance to any bat for the calculation of the sex

dissimilarity, and the distance to any same-sex bat for the calculation of hearing/deaf

dissimilarity. For a vocalization emitted by a male, the sex dissimilarity is the difference between

the average Euclidean distance to the 10 nearest vocalizations from the closest female bat, with

the average Euclidean distance to the 10 nearest vocalizations from any closest bat. For a

vocalization emitted by a hearing male, the hearing/deaf dissimilarity is the difference between

the average Euclidean distance to the 10 nearest vocalizations from the closest deaf male bat,

with the average Euclidean distance to the 10 nearest vocalizations from the closest male bat

(hearing or deaf). The proportion of the repertoire that was affected by deafening was estimated

as the proportion of vocalizations that displayed a hearing/deaf dissimilarity value larger than

the null distance to any same sex bats (Supplementary Figure 7).

Regularized Permutation Discriminant Function Analysis

The performance of classification of NMPS of vocalizations according to the hearing ability of

the emitter bat (Deaf and Hearing) was determined using a regularized permutation discriminant

function analysis (RPDFA). For each comparison (set of data), this procedure consisted in a

regularization achieved by a principal component analysis (PCA) on the NMPS, followed by a

Linear Fisher Discriminant Function Analysis (DFA). The performance of the DFA was

estimated in 10-fold cross-validation fixing the prior probability at 0.5. To find out the optimal

number of PCA dimensions to consider in the DFA, the performance of the classifier was

measured across a range of numbers of dimensions up to the number of dimensions that

explained 90% of the variance in the PCA (step of 10 up to 100 dimensions; step of 20 between

100 and 500 dimensions; step of 100 dimensions above). The optimal number of dimensions of

the PCA to consider in the DFA was found as the first value where the increase in classification

performance weighted by the increase of percentage of variance explained by the change of

number of dimensions is smaller than half the average standard deviation of the performance

across all values of dimensions tested (see Supplementary Figures 4, 6, 16 and 17). The

performance of the classifier at this optimized number of PCA dimensions was retained and

reported in the article. The chance level for the classification performance using the optimized

number of PCA dimensions was determined by running a permutation procedure where the

treatment of individuals was randomized across bats. Depending on the number of subjects in

the dataset there were 126 (ten male and female bats), 9 (six female bats) or 2 (four male bats)

different possible permutations. To determine if the odds of correct classification was higher in

the non-permuted treatment case than in the permutation cases, we perform a right tailed

Fisher's exact test for each permutation.

For display and interpretation purposes, the DFA at the optimal number of PCA dimensions was

recalculated without cross-validation to obtain the discriminant function as shown in Figure 3

and Supplementary Figure 11.

Data availability

531 The database of calls will be made available on FigShare upon publication.

Code availability

534 All the code used for call detection, curation and analyses is available under Github.

References

1.   Mahrt, E. J., Perkel, D. J., Tong, L., Rubel, E. W. & Portfors, C. V. Engineered Deafness
Reveals That Mouse Courtship Vocalizations Do Not Require Auditory Experience. *Journal of*
Neuroscience **33**, 5573–5583 (2013).
2.   Tyack, P. L. A taxonomy for vocal learning. *Philosophical Transactions of the Royal*
Society B: Biological Sciences **375**, 20180406 (2020).
3.   Tyack, P. L. Vocal Learning and Auditory-Vocal Feedback. in *Vertebrate Sound*

Production and Acoustic Communication (eds. Suthers, R. A., Fitch, W. T., Fay, R. R. & Popper,
A. N.) 261–295 (Springer International Publishing, 2016). doi:10.1007/978-3-319-27721-9_9.
4.   Janik, V. M. & Slater, P. J. B. The different roles of social learning in vocal
communication. *Animal Behaviour* **60**, 1–11 (2000).
5.   Fischer, J., Wegdell, F., Trede, F., Dal Pesco, F. & Hammerschmidt, K. Vocal
convergence in a multi-level primate society: insights into the evolution of vocal learning.
Proceedings of the Royal Society B: Biological Sciences **287**, 20202531 (2020).
6.   Ruch, H., Zürcher, Y. & Burkart, J. M. The function and mechanism of vocal
accommodation in humans and other primates. *Biological Reviews* **93**, 996–1013 (2018).
7.   Wirthlin, M. *et al.* A Modular Approach to Vocal Learning: Disentangling the Diversity of a
Complex Behavioral Trait. *Neuron* **104**, 87–99 (2019).
8.   Janik, V. M. & Knörnschild, M. Vocal production learning in mammals revisited.
Philosophical Transactions of the Royal Society B: Biological Sciences **376**, 20200244 (2021).
9.   Prat, Y., Azoulay, L., Dor, R. & Yovel, Y. Crowd vocal learning induces vocal dialects in
bats: Playback of conspecifics shapes fundamental frequency usage by pups. *PLOS Biology* **15**,
1–14 (2017).
10.   Prat, Y., Taub, M. & Yovel, Y. Vocal learning in a social mammal: Demonstrated by
isolation and playback experiments in bats. *Science Advances* **1**, e1500019 (2015).
11.   Genzel, D., Desai, J., Paras, E. & Yartsev, M. M. Long-term and persistent vocal
plasticity in adult bats. *Nature Communications* **10**, 3372 (2019).
12.   Wirthlin, M. E. *et al. Vocal learning-associated convergent evolution in mammalian
proteins and regulatory elements.* http://biorxiv.org/lookup/doi/10.1101/2022.12.17.520895
(2022) doi:10.1101/2022.12.17.520895.
13.   Rose, M. C., Styr, B., Schmid, T. A., Elie, J. E. & Yartsev, M. M. Cortical representation
of group social communication in bats. *Science* **374**, eaba9584 (2021).
14.   Yartsev, M. M. & Ulanovsky, N. Representation of Three-Dimensional Space in the
Hippocampus of Flying Bats. *Science* **340**, 367–372 (2013).
15.   Liberti, W. A., Schmid, T. A., Forli, A., Snyder, M. & Yartsev, M. M. A stable hippocampal
code in freely flying bats. *Nature* **604**, 98–103 (2022).
16.   Nottebohm, F. The Origins of Vocal Learning. *The American Naturalist* **106**, 116–140
(1972).
17.   Singh, N. C. & Theunissen, F. E. Modulation spectra of natural sounds and ethological
theories of auditory processing. *The Journal of the Acoustical Society of America* **114**, 3394–
3411 (2003).
18.   Fant, G. The Relations between Area Functions and the Acoustic Signal. *Phonetica* **37**,
55–86 (1980).
19.   Fant, Gunnar. Acoustic theory of speech production: with calculations based on X-ray
studies of Russian articulations. (1971).
20.   Vannoni, E. & McElligott, A. G. Individual Acoustic Variation in Fallow Deer (Dama
dama) Common and Harsh Groans: A Source-Filter Theory Perspective. *Ethology* **113**, 223–
234 (2007).
21.   Beckers, G. J. L., Nelson, B. S. & Suthers, R. A. Vocal-Tract Filtering by Lingual
Articulation in a Parrot. *Current Biology* **14**, 1592–1597 (2004).
22.   Taylor, A. M. & Reby, D. The contribution of source–filter theory to mammal vocal

communication research. *Journal of Zoology* **280**, 221–236 (2010).

23.   Jarvis, E. D. Evolution of vocal learning and spoken language. *Science* **366**, 50–54

(2019).

24.   Simonyan, K. The laryngeal motor cortex: its organization and connectivity. *Current*

Opinion in Neurobiology **28**, 15–21 (2014).

25.   Jürgens, U. Neural pathways underlying vocal control. *Neuroscience & Biobehavioral*

Reviews **26**, 235–258 (2002).

26.   Fitch, W. T. The Biology and Evolution of Speech: A Comparative Analysis. *Annu. Rev.*

Linguist. **4**, 255–279 (2018).

27.   Kuypers, H. G. J. M. Corticobulbar connexions to the Pons and lower brain-stem in man:

an anatomical study. *Brain* **81**, 364–388 (1958).

28.   Lattenkamp, E. Z. *et al.* The vocal development of the pale spear-nosed bat is

dependent on auditory feedback. *Philosophical Transactions of the Royal Society B: Biological*

Sciences **376**, 20200253 (2021).

29.   Nasir, S. M. & Ostry, D. J. Speech motor learning in profoundly deaf adults. *Nature*

Neuroscience **11**, 1217–1222 (2008).

30.   Tremblay, S., Shiller, D. M. & Ostry, D. J. Somatosensory basis of speech production.

Nature **423**, 866–869 (2003).

31.   McGregor, J. N. *et al.* Shared mechanisms of auditory and non-auditory vocal learning in

the songbird brain. *eLife* **11**, e75691 (2022).

32.   Linnenschmidt, M. & Wiegrebe, L. Ontogeny of auditory brainstem responses in the bat,

Phyllostomus discolor. *Hearing Research* **373**, 85–95 (2019).

Acknowledgements

The authors deeply thank the undergraduate students from UC Berkeley who worked on the first

manual curation step of the bats audio recordings: Yousuf Hashmi, Sun Wee Yun, Nicole Wang,

Maggie McDonough, Leo Huang, Jason Huang, Christopher Lung, Aunesha Bhowmick, Anya

Cutter, Isha Yeleswarapu, Geoffrey Li, Ethan Lee, Dorian Andrei and Chirstina Yao.

We also would like to thank the undergraduate students from UC Berkeley who participated in

establishing the automatic algorithm for bat call detection: Shunyu Wang, Richard Ma, Jonathan

Wang and Chase Adams. We thank Yartsev lab members for discussions and thoughts on the

project, L. Wiegrebe for his help on the initial project design and sharing of ABR protocol &

software, and OLAC staff for support with animal husbandry and care. **Funding:** this research

was supported by NIH (DP2-DC016163), the New York Stem Cell Foundation (NYSCF-R-NI40),

the Alfred P. Sloan Foundation (FG-2017-9646), the Brain Research Foundation (BRFSG-2017-

626   Author contributions

Conceptualization: MMY, SEM, JEE

Methodology: MMY, SEM, JEE, YR, LAB, DLS

Resources: MMY, YR

Data Curation: JEE

Investigation: SEM, JEE, DG, LAB

Software: JEE, TN

Formal acoustical analysis: JEE

Formal histological analysis: LAB, DLS, YR

Funding acquisition: MMY

Supervision: JEE, MMY

Writing – original draft: JEE

Writing – review & editing: JEE, MMY, LAB, TN, SEM, DG, YR, DLS

640   Competing interests

The authors have no competing interests to declare.

Figure 1 Experimental design testing the necessity of hearing from birth for the normal

development of the vocal repertoire in adult Egyptian fruit-bats

A. Pregnant females gave birth and cared for their young in complete isolation from the rest of

the colony for two months. Their ten pups were randomly assigned to a treatment by kanamycin

or saline (control). Mothers and pups were then reunited with the rest of the colony for 3-4

years. Well after reaching adulthood, the ten manipulated bats were housed together and their

adult vocal behavior was recorded for seven weeks. Finally, their audiograms were measured

and their cochlea inspected for histological damage.

B. Audiograms obtained from measuring the Auditory Brainstem Response of all bats (n=10) to

tone pips covering the species auditory range. Dots depict individual values, while lines show

the average fitted with a spline. Note that responses of the auditory nerve were absent (values

ceiled at 120dB) or only detectable for very loud sounds (>90dB) in kanamycin treated bats,

while the expected audiogram was obtained for saline treated bats.

C. Representative examples of the hearing sensory epithelium (organ of Corti) at two levels of

the cochlea (apical turn, first and second columns; basal turn, third column) in two bats stained

with phalloidin (first row, female; second row, male). Scale bars: 200µm in first column, 25µm in

other columns; grey square: area shown in second column. Note the intact epithelium (OHC,

outer hair cells; IHC, inner hair cells) both in the apical and basal cochlear areas of the saline

bat, as opposed to the non-sensory flat epithelium in the kanamycin bat. Hair cells are absent

throughout the organ of Corti (see white arrows) except for a few cells in kanamycin female bats

(Supplementary Figure 1).

D. Vocalizations are recorded both from an environment microphone and a piezo sensor placed

against the throat of the bat (wireless audiologger). Two approaches are used to find the effect

of deafening on bats' vocalizations. Upper panel: the data driven approach is using the

modulation power spectrum of vocalizations as input to a classifier that is trained and tested in

cross-validation to discriminate between hearing and deaf bats' calls. The permutation test gives

the correct chance level taking into account any acoustic differences due to bat identity.

The second approach test the effect of deafening by fitting a Linear Mixed-Effect model

controlling for the identity of the bats on each of 28 acoustic features of the vocalizations

(Supplementary Figure 3 and methods).

<u>Figure 2 Effect of sex and deafness through the vocal space</u>

Projections of vocalizations as represented by their MPS in a 3-dimension UMAP, the vocal

space.

A. Dissimilarity of bat vocalizations to opposite sex vocalizations. The color of the points indicate

the sex of the emitter (male, blue, and female, red) and the intensity of the color the linear PCA

distance to the closest ten vocalizations produced by an opposite sex bat (see methods). Note

that vocalizations with UMAP2<-2 are almost exclusively produced by males.

B and D Dissimilarity of male (B) or female (D) vocalizations to vocalizations from a same-sex

bat with opposite hearing ability. Male (B) or female (D) vocalizations are color coded according

to the hearing ability of the emitter (hearing, blue, and deaf, red). The intensity of the color

corresponds to the linear PCA distance to the closest ten vocalizations produced by a male (B)

or a female (D) of opposite hearing ability (methods). The arrowhead on the color scale

indicates the threshold over which vocalizations were considered idiosyncratic of the bat hearing

ability (dissimilarity value larger than expected on average to any same sex calls; see

Supplementary figure 8 and methods).

C. Parsing of the vocal space into four groups according to the results of a Hierarchical

Agglomerative Clustering algorithm.

Figure 3 Effect of deafness on male calls in acoustic group 4

A. Coefficients of the acoustic Group 4 vocalizations of each male bat when projected on the

first linear acoustic axis of the PCARDFA that best discriminates between deaf (M1 and M2)

and hearing (M3 and M4) bats. Each circle in the scatter plot represents a vocalization. The

circle color indicates the value of the Third Spectral Quartile of the vocalization. The three

images on the left show the average MPS of male vocalizations in acoustic Group 4 (middle)

and the spectro-temporal changes in the MPS (discriminant function) that best characterize

vocalizations from deaf males (top) and vocalizations from hearing males (bottom). Note that

while on average male vocalizations in acoustic Group 4 are characterized by pitch values of

1c/kHz or 0.5 c/kHz corresponding respectively to 1kHz and 2kHz (black arrows), the time

varying pitch modulations are different between deaf and hearing males: hearing males produce

calls with more downsweeps (black circle) and deaf males with more upsweeps (gray circle).

The black arrows in the scatter plot point to the example calls shown in B.

B. Spectrogram of two example calls from acoustic Group 4, per male bat. Note that hearing

male vocalizations are short downsweep vocalizations with a low Third Spectral Quartile (black

dotted lines) while the deaf male vocalizations are longer and more pitch modulated

vocalizations (presence of upsweeps) with high Third Spectral Quartile (black dotted lines). The

audio file of each of these examples can be found as supplementary materials

(Group4_DeafMaleBatM2_1.wav, Group4_DeafMaleBatM2_2.wav,

Group4_DeafMaleBatM1_1.wav, Group4_DeafMaleBatM1_2.wav,

Group4_HearingMaleBatM3_1.wav, Group4_HearingMaleBatM3_2.wav,

Group4_HearingMaleBatM4_1.wav, Group4_HearingMaleBatM4_2.wav)