Active and abundant alphaproteobacterial nitric oxide transforming enzymes in a marine oxygen

deficient zone

Claire E. Elbon[1], Frank J. Stewart[2], and Jennifer B. Glass[1*]

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA;

[2]Department of Microbiology & Cell Biology, Montana State University, Bozeman, MT, USA

*Corresponding author: Jennifer B. Glass, Jennifer.Glass@eas.gatech.edu

Abstract

Marine oxygen deficient zones (ODZs) are portions of the ocean where intense nitrogen loss

occurs primarily via denitrification and anammox. Despite many decades of study, the identity of

the microbes that catalyze nitrogen loss in ODZs are still being elucidated. Intriguingly, high

transcription of genes in the same family as nitric oxide dismutase from *Methylomirabilota* have

been reported in the anoxic core of ODZs. Here, we show that the most abundantly transcribed

nod genes in the Eastern Tropical North Pacific ODZ belong to *Rhodospirillaceae*

(Alphaproteobacteria)*,*   rather   than   *Methylomirabilota*   as   previously   assumed.

Gammaproteobacteria and Planctomycetia also transcribe *nod*, but at lower relative abundance

than *Rhodospirillaceae* in the upper ODZ. The *Rhodospirillaceae* are likely methylotrophs that

oxidize methanol as a source of electrons for aerobic respiration; additional electrons may come

from sulfide oxidation. Molecular oxygen for aerobic respiration may originate from nitric oxide

dismutation via cryptic oxygen cycling. The *Rhodospirillaceae* also transcribe multiheme

cytochrome (here named *ptd*) genes for a putative porin-cytochrome protein complex of

unknown function, potentially involved in extracellular reduction electron transfer. Our results

implicate *Rhodospirillaceae* as a significant player in marine nitrogen loss and highlight its

potential in one-carbon, nitrogen, and sulfur metabolism in ODZs.

Significance statement

In marine oxygen deficient zones, microbes transform bioavailable nitrogen to gaseous nitrogen,

with nitric oxide as a key intermediate. The Eastern Tropical North Pacific contains the world's

largest oxygen deficient zone, but the identity of the microbes transforming nitric oxide remain

unknown. Here, we show that highly transcribed nitric oxide dismutase *(nod)* genes belong to

Rhodospirillaceae (Alphaproteobacteria). These *Rhodospirillaceae* perform aerobic respiration,

using oxygen potentially sourced from nitric oxide dismutase, and possess a novel porin-

cytochrome protein complex with unknown function. Gammaproteobacteria and Planctomycetia

transcribe *nod* at lower levels. Our results pinpoint the microbes mediating a key step in marine

nitrogen loss and reveal an unexpected metabolism for marine Alphaproteobacteria.

Introduction

Marine oxygen deficient zones (ODZs) contribute up to half of the ocean's nitrogen loss (DeVries et al., 2013) and are a major source of marine emissions of the potent greenhouse gas nitrous oxide (N_2O) (Yang et al., 2020). The primary source of the N_2O at the oxic-anoxic interface and in anoxic waters in ODZs is denitrification (Babbin et al., 2015; Frey et al., 2020). The microbial enzyme responsible for N_2O production during denitrification is nitric oxide reductase (Nor), which uses electrons from cytochrome *c* (cNor) or quinol (qNor), to reduce nitric oxide (NO) to N_2O (Wasser et al., 2002; Zumft, 2005; Kraft et al., 2011). In the qNor family, there are *bona fide* qNor enzymes and NO dismutase (NOD). NOD proteins lack the quinol-binding site, seemingly preventing the enzyme from taking up external electrons; instead, NOD is theorized to disproportionate NO into dinitrogen and O_2 in methane-oxidizing *Methylomirabilota* bacteria (Ettwig et al., 2010; Ettwig et al., 2012) and alkane-oxidizing gammaproteobacterium HdN1 (Zedelius et al., 2011).

The Eastern Tropical North and South Pacific (ETNP and ETSP) ODZs are the world's largest and second largest ODZs, and the subjects of extensive microbial ecology studies. Abundant NO reductase-like genes and transcripts in the ETNP and ETSP ODZ cluster in the same enzyme subfamily as NOD (Dalsgaard et al., 2014; Ganesh et al., 2014; Padilla et al., 2016; Fuchsman et al., 2017). Due to the similarity of ODZ Nod proteins to those of *Methylomirabilota* (NC10), it was initially presumed that ODZ bacteria also used Nod proteins to disproportionate NO into N_2 and O_2 for use in intra-aerobic methane oxidation (Dalsgaard et al., 2014; Padilla et al., 2016; Thamdrup et al., 2019). However, Fuchsman et al. (2017) found that the peak of *nod* gene abundance in the ETNP ODZ correlates with a peak of modeled N_2O production (Babbin et al., 2015) and does not correlate with abundance of methane monooxygenase genes, suggesting

that Nod proteins in the ETNP ODZ are potentially an important source of N_2O, and are unlikely

to be involved in methane oxidation. The plausibility that Nod proteins can reduce NO to N_2O is

supported by a study of a novel eukaryotic denitrification pathway in foraminifera

(*Globobulimina* spp.) that produces N_2O while expressing Nod (Woehle et al., 2018). Yet, the

phylogenetic identity and metabolic context of marine Nod proteins, which are a key biological

source of either N_2O or O_2+N_2 in marine ODZs, remain unresolved.

In this study, we sought to determine the identity, metabolism, and environmental niche

of the ODZ organism responsible for the highly transcribed *nod* genes first discovered in Padilla

et al. (2016). We found that the most abundantly transcribed *nod* genes in the ETNP ODZ belong

to Alphaproteobacteria related to *Rhodospirillaceae*. Significant transcription of *nod* genes was

limited to waters with <1 μM O_2. These *nod*-transcribing alphaproteobacteria also transcribe

genes involved in aerobic respiration, which was unexpected given that they inhabit anoxic

waters, as well as genes involved in oxidation of formaldehyde, likely indicating methylotrophy.

Genes encoding multi-heme cytochrome proteins potentially implicated in nitrogen or iron

cycling were also transcribed.

Results

Transcribed* nod *sequences in the ETNP ODZ belong to Alphaproteobacteria,

Gammaproteobacteria, and Planctomycetia. Our reanalysis of highly transcribed *nod* genes

("ETNP 2014 Stn10 150m" and "ETNP 2013 Stn6 300m") in the ETNP ODZ (Padilla et al.,

2016) shows that these genes belong to Alphaproteobacteria rather than a member of

Methylomirabilota as previously assumed. Querying the Nod amino acid sequences from Padilla

et al. (2016) against ETNP ODZ metagenomes in the IMG-JGI database returned multiple 100%

identity matches, including a gene co-occurring on a scaffold (Ga0066848_10003785) with

hypothetical genes with 100% identity to *Rhodospirillaceae* metagenome-assembled genomes

(MAGs) from the ETNP ODZ (Uzun et al., 2020) **(Table S1)**. Binning of ETNP ODZ

metagenomes Ga0066848 (ETNP201310SV72) and Ga0066829 (ETNP201306SV43) placed the

two contigs with the most highly transcribed *nod* genes into MAGs assigned to

Alphaproteobacteria (GTDB taxonomy: UBA11136 sp002686135; GTDB species

representative: *Rhodospirillaceae* bacterium isolate ARS27) with 97% average nucleotide

identity. Querying the Nod amino acid sequences from Padilla et al. (2016) against NCBI's non-

redundant protein database returned matches to Alphaproteobacteria/*Rhodospirillaceae* MAGs

from low-oxygen marine settings (ETNP, Saanich Inlet, and the Black Sea; 78-80% identity), the

marine magnetotactic alphaproteobacterium *Magnetovibrio blakemorei* MV-1 (75% identity),

Gammproteobacterium HdN1 (66% identity), and *Methylomirabilota* spp. (66% identity; **Table**

S2).

To glean additional insights into the evolutionary relationships of ODZ *nod* genes, we

gathered Nod amino acid sequences from cultured organisms and large ODZ metagenome

datasets (ETNP and Saanich Inlet), and created a Nod phylogeny **(Figure 1A; Table S3)**. The

topology was generally consistent with a previous phylogeny from Fuchsman et al. (2017), with

additional taxonomic data from MAGs in the TARA oceans dataset further constraining Nod

placement (Tully et al., 2018). Six unique Nod ODZ protein sequences (two of which were

present in multiple metagenomes) clustered with Planctomycetia (OTU I in Fuchsman et al.

(2017), hereafter "Planctomycetia-type Nod"), and were primarily found in free-living cells (0.2-

1.6 micron, "FL"). Four unique ODZ Nod sequences clustered with marine

Gammaproteobacteria (OTU II in Fuchsman et al. (2017), hereafter "Gamma-type Nod"); these

sequences were monophyletic with a cluster of gammaproteobacterial Nod cluster sequences

from sewage sludge, including gammproteobacterium HdN1 (Ehrenreich et al., 2000) and other

wastewater gammproteobacteria. Multiple ETNP ODZ metagenomes contained Gamma-type

Nod sequences identical to those of Gammaproteobacteria NP964 (MBP20251). Several ODZ

Nod sequences, all from the particle fraction (>1.6 micron, "PF") clustered with marine

Deltaproteobacteria in a clade monophyletic with *Methylomirobalis*, Deltaproteobacteria, and

Acidobacteria MAGs from groundwater. As expected based on the binning and BLAST results,

the Nod sequence from Padilla et al. (2016) clustered phylogenetically with marine

alphaproteobacteria (OTU III in Fuchsman et al. (2017), hereafter "Alpha-type Nod"); this clade

contained three unique sequences, all of which were present in multiple metagenomes and all

from the free-living fraction, and one of which was identical to that of *Rhodospirillaceae*

NP1106 (MBV28360). Intriguingly, two ODZ sequences clustered in the eukaryotic

Globobulimina clade.

We investigated gene neighborhoods surrounding ODZ *nod* genes in the three main

phylogenetic clusters of ODZ sequences: Planctomycetia-type Nod, Gamma-type Nod, and

Alpha-type Nod. Whereas "unknown Nor-related" marine *Bacteroidota* sequences were located

on an operon with other *nor* genes, there was no consistent gene neighborhood for *nod* sequences

(Figure 1B). Planctomycetia-type *nod* genes were not located in the vicinity of any genes with

recognizable related function. Gamma-type *nod* gene neighborhoods contained ferredoxins and

cytochrome b_{561} genes for electron transport. Upstream of the Alpha-type *nod* in

Rhodospirillaceae NP1106 is a cluster of formylmethanofuran dehydrogenase genes (*fmd/fwd*)

used in C1 metabolism via tetrahydromethanopterin/methanofuran-linked reactions.

Immediately upstream or downstream of *nod* genes, helix-turn-helix transcriptional

regulators were common **(Figure 1B)**. Neighboring Gamma-type and *Methylomirobalis nod*

genes, LuxR-type regulators were common; these regulators have diverse functions and their

potential connection to Nod remains unclear. Neighboring Alpha-type and Bacteroidota (e.g.

Cecembia calidifontis) *nod* genes, Rrf2-type regulators were present. The protein NsrR in the

Rrf2 family regulates global cellular response to NO toxification by directly sensing NO with an

iron-sulfur cluster (Bodenmiller and Spiro, 2006; Tucker et al., 2010). The presence of this

NsrR-like regulator suggests that Nod in marine *Rhodospirillaceae* and Bacteroidota may be

involved in nitrosative stress response and NO detoxification.

Alphaproteobacterial* nod *is highly transcribed in anoxic waters. We assessed

transcription of Alpha, Gamma-, and Planctomycetia-type *nod* genes from the oxycline to upper

ODZ (secondary nitrite maximum) using ETNP ODZ metatranscriptomes from an onshore

station with a shallower oxycline (P1; **Figure 1C**) and an offshore station with a deeper oxycline

(P2; **Figure 1D**) (Mattes et al., 2022). In both oxyclines, transcription was low (4-10 reads per

kilobase per million mapped reads (RPKM), n=8) for all three *nod* types (**Figure 1C, D**). Below

the oxyclines, *nod* transcripts began to rise and were highest at the secondary nitrite maxima,

with Alpha-type (184-274 RPKM, n=4) > Gamma-type (55-95 RPKM, n=4) > Planctomycetia-

type (13-19 RPKM, n=4; **Table S4**).

Alphaproteobacteria transcribe genes for formate metabolism, aerobic respiration, and

a multiheme cytochrome complex. To glean insight into potential roles for Nod in cellular

context, we sought to reconstruct the electron transport chain of the alphaproteobacterium that

most highly transcribed *nod* genes (Alphaproteobacterium MAG ETNP2013_S06_300m_15 and

Alphaproteobacterium MAG ETNP2013_S10_300m_22, 69% and 73% estimated completeness,

respectively) at the secondary nitrite maximum. In both MAGs, *nod* was in the top three most

transcribed genes in the ETNP ODZ (~44,000 FPKM; **Table S5)**, after a bacterial nucleoid

DNA-binding protein and a potassium gated channel protein. In addition to *nod,* we found that

genes for formaldehyde oxidation via tetrahydromethanopterin/methanofuran-linked reactions,

including   formylmethanofuran   dehydrogenase   (*fwd/fmd*)   and   formylmethanofuran-­

tetrahydromethanopterin N-formyltransferase (*ftr*), were transcribed in both MAGs **(Table S5)**.

Both MAGs also transcribed NAD-dependent formate dehydrogenase **(Table S5)**. Thus, the

alphaproteobacterium appears to be capable of conversion of formaldehyde to formate and use of

formate as a source of electrons for NADH:ubiquinone oxidoreductase (Complex I; **Figure 2)**.

The source of formaldehyde is likely methanol oxidation, as pyrroloquinoline quinone (PQQ)-

dependent ethanol/methanol dehydrogenases were found in *Rhodospirillaceae* MAGs from low-

oxygen marine settings **(Table S6)**. Methane monooxygenase genes were not found in the partial

Rhodospirillaceae MAGs, precluding our ability to rule out the possibility of these genes in the

missing portions of the genomes. The *Rhodospirillaceae* PQQ-dependent dehydrogenase genes

contained the motif DYDG **(Table S6)**, which is characteristic of the lanthanide-containing form

of the enzymes rather than calcium form (Keltjens et al., 2014).

A full aerobic electron transport chain (Complex I, II, III, and IV) and F0F1-type ATP

synthase were transcribed in both bins **(Table S5)**. Complex IV (cytochrome c oxidase) was type

A1 according to the Sousa et al. (2012) classification, and the *cox* operon in the GTDB species

representative Rhodospirallaceae ARS27 was subtype b (COX2-COX1-CtaB-CtaG_Cox11-

COX3-DUF983-SURF1-CtaA1-M32-Tsy-M16B)   according   to   the   Geiger   et   al.   (2023)

classification. Sulfur oxidation genes, including flavocytochrome c sulfide dehydrogenase

(FccAB), sulfane hydrogenase (SoxCD), and carrier protein SoxYZ, were also transcribed, as

were numerous transposes **(Table S5)**.

Genes for a multiheme cytochrome complex were transcribed in both bins. To our

knowledge, this putative operon has not previously been described. Hereafter, we designate it the

ptdABCDEFG operon for its sequence of p̲enta/t̲etra/d̲eca-heme proteins, interspersed with other

conserved proteins. *ptdAB* genes are highly transcribed in our *Rhodospirillaceae* MAGs, but it is

unclear if the rest of the operon is also highly transcribed, because it was truncated in our MAGs'

scaffolds. The *ptd* gene cluster consists of a penta-heme protein with a C-terminal beta-sandwich

(PtdA), a porin (PtdB), a FAD/NAD(P)-binding oxidoreductase (PtdC), a periplasmic tetra-heme

protein (PtdD), a cyclic nucleotide-binding domain protein with two 4Fe-4S clusters (PtdE), a

cytoplasmic transmembrane ferric reductase-like protein (PtdF), and a periplasmic deca-heme

protein (PtdG; **Figure 2; Tables S7, S8**). The function of this complex is unknown, but the

presence of genes encoding a porin and multiple multiheme proteins resembles porin-

cytochrome protein complexes involved in extracellular reduction electron transfer during Fe(III)

and Mn(IV) reduction (Richardson et al., 2012; Shi et al., 2014). PtdA has a homolog to a penta-

heme cytochrome c_{552} protein of unknown function in a thermophilic purple sulfur

gammproteobacterium (Chen et al., 2019) and is in the same COG family (COG3303) as formate

dependent nitrite reductase, NrfA. *ptdABCDEFG* genes were prevalent in Alphaproteobacteria,

Gammaproteobacteria, Nitrospirales, and Planctomycetes MAGs from marine or high salinity

environments **(Figure 3)**.

Discussion

This study illuminates the previously ambiguous identity of the microorganisms that make the

dominant nitric oxide-transforming protein (Nod) in the world's largest ODZ, the Eastern

Tropical North Pacific. Extensive horizontal gene transfer of *nod* genes between microbial

genomes is evident from the lack of conservation of gene neighborhood and patchy phylogeny

(Fuchsman et al., 2017), which may be mediated by viral infection (Gazitúa et al., 2021). We

found that the most transcriptionally active *nod* genes in the ETNP upper ODZ belong to

Alphaproteobacteria related to *Rhodospirillaceae*. Alpha-type *nod* transcript abundances (~200

RPKM) are similar to those of dissimilatory nitrate reductase (*narG*) in the ODZ (Tsementzi et

al., 2016). The *nod*-transcribing *Rhodospirillaceae* are likely methylotrophs, and transcribe genes

for formaldehyde oxidation, likely as a source of electrons to the respiratory chain via NAD

reduction by formate dehydrogenase. Sulfide may be used as a supplemental electron donor

and/or may be concomitantly oxidized for detoxification (Callbeck et al., 2021; Schmitz et al.,

2023).

Our discovery of a putative porin-cytochrome complex (*ptd* operon) in marine bacteria

was unexpected. Porin-cytochrome complexes have been best studied for their role in

extracellular electron transport, particularly for respiratory metal reduction and oxidation

(Richardson et al., 2012; Shi et al., 2014). It is conceivable that the Ptd complex is involved in

iron reduction in ODZs; there is iron reduction at the secondary nitrite maximum and it is

hypothesized to be bacterially mediated, but the microbes involved have yet to be determined

(Moffett et al., 2007; Glass et al., 2015). Alternatively, the presence of *ptdABCDEFG* genes in

numerous nitrite-oxidizing bacteria (Nitrospirales) could imply the involvement of these genes in

nitrogen cycling; PtdA was in the same COG family as formate-dependent nitrite reductase

(Simon et al., 2000) and PtdC is similar to a flavohemoprotein with predicted nitric oxide

dioxygenase activity, also annotated as hydroxylamine oxidoreductase-linked cytochrome. The

function of PtdABCDEFG remains completely unknown and requires future biochemical

characterization.

On the other end of the electron transport chain, high transcription of a heme/copper

terminal oxidase suggests that O_2 is being used as the terminal election acceptor in *nod*-

transcribing *Rhodospirillaceae*. The transcribed heme/copper oxidase is A1-type (low O_2

affinity), also present in mitochondria, and adapted for high O_2 concentrations. Low O_2 affinity

A1-type heme/copper oxidases are transcribed in other anoxic environments (Berg et al., 2022).

Because ODZs have extremely low concentrations of molecular oxygen below the oxycline, O_2

for aerobic respiration may be generated *in situ* and rapidly consumed. Given that the function of

Nod is proposed to be dismutation of two NO molecules into N_2 and O_2 (Ettwig et al., 2010), it is

possible that the O_2 source for aerobic respiration in *Rhodospirillaceae* is NO dismutation,

although other sources of O_2 in anoxic waters are also conceivable (Garcia-Robledo et al., 2017).

The physiological uses of Gamma-type and Planctomycetia-type Nod may be different from

Alpha-type Nod, although this remains to be investigated.

The source of NO, the presumed substrate for Nod, may be generated in the same

organism using Nod, or generated by a different organism (or chemical pathway). Nitric oxide

was positively correlated with nitrite in the Eastern Tropical South Pacific ODZ, and was only

detectable when O_2 was <1-2 μM (Lutterbeck et al., 2018). In the Eastern Tropical North Pacific,

NO concentration and turnover rates were elevated at O_2 <100 μM (Ward and Zafiriou, 1988).

Both studies suggest that the NO in ODZs likely originates from nitrification or nitrifier

denitrification, while genomic analyses indicate that the copper-containing nitrite reductase

(*nirK*) in SAR11 bacteria (presumably performing denitrification) may be a key source of NO

(Fuchsman et al., 2017). Because most ODZ denitrifiers specialize in only one of the three steps

(NO$_2^-$ reduction, NO reduction, and N$_2$O reduction) (Zhang et al., 2023), and known nitrite

reductases were not identified in our MAGs, existing data indicate that the NO that is used as a

substrate for *Rhodospirillaceae* Nod is not generated *in vivo*. (Only 4 out of 32 *nod*-containing

MAGs contained a nitrite reductase gene: two Gammaproteobacteria MAGs contained *nirK*, one

Myxococcota MAG contained *nirS*, and one Scalindua MAG contained *nirS*). It is also possible

that another uncharacterized enzyme produces NO.

This study suggests that marine alphaproteobacteria (*Rhodospirillaceae*) are actively

reducing NO under anoxia, as implied by their abundant transcription of *nod* genes. While we

can be fairly certain that the substrate for Nod is NO, the products of this enzyme (N$_2$O vs.

N$_2$+O$_2$) remain uncertain. Nod is theorized to disproportionate NO into N$_2$ and O$_2$ in methane-

oxidizing *Methylomirabilota* bacteria (Ettwig et al., 2010; Ettwig et al., 2012), but no

biochemical characterizations of Nod have been published to date, and foraminifera expressing

Nod produce N$_2$O (Woehle et al., 2018). The apparent lack of other denitrification genes in *nod*-

transcribing *Rhodospirillaceae* is consistent with the observation that denitrification in ODZ is

largely divided into distinct microbial taxa (Dalsgaard et al., 2014; Fuchsman et al., 2017; Zhang

et al., 2023). For example, although nitrate reductase (*narG*) genes are widely distributed

amongst ODZ microbes (Zhang et al., 2023), SAR11 bacteria appear to dominate in *narG*

transcriptional activity (Tsementzi et al., 2016). Our finding that the transcription of *nod* is

catalyzed primarily by marine alphaproteobacteria implies that this taxa contributes significantly

to marine nitrogen loss.

METHODS

Nod *phylogeny and gene neighborhood*. Amino acid sequences of highly transcribed *nod* genes

"ETNP 2014 Stn10 150m" and "ETNP 2013 Stn6 300m" were acquired from the authors of

Padilla et al. (2016) (see Table S2 for sequences). These sequences were used for BLASTP

searches of ODZ metagenomes in the IMG-JGI database and the non-redundant protein (nr)

database in NCBI. Sequences (n=53, 731 gap-free sites) were aligned using the MAFFT online

server with the L-INS-i method (Katoh et al., 2019). A phylogeny was generated with 1000

bootstraps using model LG+I+G4 using W-IQ-Tree (Trifinopoulos et al., 2016). The phylogeny

was visualized using FigTree v.1.4.4, and the fasta file (Trimmed_NOD_tree) is available as a

supplemental figure. Gene neighborhoods were generated using the EFI Gene Neighborhood

Tool (Zallot et al., 2019) with single sequence BLAST of the UniProt database using the amino

acid sequence Ga0066848_100037855 (JGI IMG) as the Nod query with an e-value cutoff of 10^{-5}

and with 10 genes upstream and downstream the gene of interest.

Transcription of* nod *genes in ETNP ODZ depth profiles. Magic Basic Local Alignment

Search Tool (Boratyn et al., 2019) was used to search ETNP ODZ metatranscriptomes

(PRJNA727903; Mattes et al. (2022)) using representative nucleotide sequences for

Planctomycetia-like   (Ga0066826_100064333   (JGI   IMG)),   Gamma-like

(PBRC01000062.1:19833-22205 (NCBI)), and Alpha-like (Ga0066848_100037855 (JGI IMG))

nod genes. Default parameters were used except for the score threshold (18). Read hits were

normalized to reads per kilobase million (RPKM).

Metagenomic binning. Binning of metagenome-assembled genomes (MAGs) and

metatranscriptomic mapping to MAGs was performed using the KBase platform (Arkin et al.,

2018). Assemblies for the ETNP ODZ metagenomes containing Alpha-type *nod* genes

(ETNP201310SV72 (GOLD Analysis Project ID Ga0066848; stn10 300m) and

ETNP201306SV43 (GOLD Analysis Project ID Ga0066829; stn6 300m) were imported from

JGI IMG into Kbase. Metagenomic assemblies were binned into MAGs using MaxBin2 v2.2.4

(Wu et al., 2016). The two MAGs containing *nod* genes (MAG ETNP2013_S10_300m_22 from

ETNP201310SV72 and ETNP2013_S06_300m_15 from ETNP201306SV43) were selected for

further analysis. Average nucleotide identity was calculated using FastANI (Jain et al., 2018).

MAG taxonomy was evaluated by GTDB-Tk v2.3.2 (Chaumeil et al., 2022). MAGs were

annotated with RASTtk v1.073 (Brettin et al., 2015).

295   ***Mapping transcripts to metagenomic bins***. Metatranscriptomic fragments were imported

from the depth with highest *nod* transcription at the secondary nitrite maximum (NCBI run

SRR14460584). Fragments were aligned to MAGs using Bowtie2 (Langmead and Salzberg,

2012). Transcripts were assembled from RNA-seq read alignments using Cufflinks v2.2.1

(Trapnell et al., 2012). Fragment hits were normalized to fragments per kilobase million

(FPKM), which differ from the above RPKM because fragments are derived from paired-end

RNA-seq data where there can be two reads corresponding to a single fragment.

302   ***Cellular localization and heme numbers***. Cellular locations of Ptd proteins were

predicted using PSORTb v3.0.3 analysis (Yu et al., 2010). Numbers of heme-binding motifs per

protein were identified by counting CXXCH sequences. Ptd gene neighborhoods was generated

using the EFI Gene Neighborhood Tool (Zallot et al., 2019) with single sequence BLAST of the

UniProt database using the amino acid sequence Ga0066848_100031354 (JGI IMG) as the PtdA

query, with an e-value cutoff of 10^{-5} and with 10 genes upstream and downstream the gene of

interest.

Data availability

The Kbase bioinformatic pipeline and MAGs are at https://narrative.kbase.us/narrative/106999.

MAGs were also deposited into BioProject PRJNA375524 (ETNP201306SV43) and BioProject

PRJNA375542 (ETNP201310SV72).

Acknowledgements

We thank Laura Bristow for helpful discussions. We thank Cory Padilla, Anthony Bertagnolli,

and Neha Sarode for sharing previous data. We acknowledge funding from an NSF Graduate

Research Fellowship to CEE, the Simons Foundation, and NSF Awards 2022991 and 2054927 to

FJS.

Figure 1. Marine Nod clades, gene neighborhoods, and depth profiles of transcription. (A)

Maximum likelihood phylogeny of nitric oxide dismutase (Nod) amino acid sequences in marine

(blue) and select terrestrial (brown) taxa. Branch support was evaluated using 1000 rapid

bootstrap replicates, with bootstrap values shown for deep branches. The tree is drawn to scale,

with branch lengths in number of substitutions per site. Bold sequences represent those present in

multiple ETNP ODZ metagenomes (see **Table S3** for duplicate accession numbers). "PF"

indicates genes from the particle fraction (> 1.6 micron fraction) of filters. "FL" indicates genes

from the free-living fraction (0.2-1.6 micron) collected on Sterivex filters. The most highly

transcribed ETNP ODZ sequence is indicated with an asterisk. The qNor sequence *Geobacillus*

stearothermophilus was used as the outgroup. (B) Gene neighborhoods surrounding *nod* genes in

select taxa. GenBank contigs: *Cecembia calfifontis* SGXG01000001, *Scalindua japonica*

BAOS01000045, Gammaproteobacteria NP964 PBRC01000062, Gammaproteobacterium HdN1

FP929140, Deltaproteobacteria NZCL01000067, *Candidatus* Methylomirabilis oxyfera

FP565575, and *Rhodospirillaceae* NP1106 PCBZ01000014. Unlabeled gray genes are

hypothetical. (C) Oxygen and nitrite concentrations, and *nod* transcripts (reads per kilobase per

million mapped reads (RKPM)) with depth in ETNP ODZ P1 (onshore) and P2 (offshore) sites

(Mattes et al., 2022).

Figure 2. Schematic of the electron transport chain in *nod*-containing ODZ Rhodospiralleace. Enzymes were included based on presence and transcriptional activity of metagenome-assembled genomes (MAGs) assigned to Alphaproteobacteria (GTDB taxonomy: UBA11136 sp002686135; see text). The color of each protein is chosen according to transcriptional activity and represented from 0-100, 100-1,000, 1,000-10,000, and 10,000-100,000 FPKM in gradient from lighter to darker blue (Table S5). Heme proteins are indicated

by red circular hemes with the cartoon number corresponding to the number of actual hemes

present on each protein. Hypothetical Ptd proteins are labelled A, B, C, D, E, F, and G, and

location within the cell is determined using Psort bacterial localization prediction tool (Table

S8). ETC complexes I-V found in *Rhodospirillaceae* MAGs are labelled with proposed

interactions between formate oxidation and complex I NADH electron transfer. Highly

transcribed NOD protein and predicted O_2 generation is shown as feeding into A1 type CCO

complex IV reduction. Additional electrons for CytC and the ETC are proposed to come from

sulfur oxidation carried out by the flavocytochrome *c* sulfide dehydrogenase (FccAB, FCC), and

sulfane-sulfur dehydrogenase (SoxCD) with the multi-enzyme carrier complex (SoxYZ).

Figure 3. Gene neighborhoods of petaheme-tetraheme-decaheme genes from select organisms. Depicted heme spacing is approximate.

References

Arkin, A.P., Cottingham, R.W., Henry, C.S., Harris, N.L., Stevens, R.L., Maslov, S. et al. (2018) KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* **36**: 566-569.

Babbin, A.R., Bianchi, D., Jayakumar, A., and Ward, B.B. (2015) Rapid nitrous oxide cycling in the suboxic ocean. *Science* **348**: 1127-1129.

Berg, J.S., Ahmerkamp, S., Pjevac, P., Hausmann, B., Milucka, J., and Kuypers, M.M.M. (2022) How low can they go? Aerobic respiration by microorganisms under apparent anoxia. *FEMS Microbiol Rev* **46**.

Bodenmiller, D.M., and Spiro, S. (2006) The yjeB (nsrR) gene of Escherichia coli encodes a nitric oxide-sensitive transcriptional regulator. *J Bacteriol* **188**: 874-881.

Boratyn, G.M., Thierry-Mieg, J., Thierry-Mieg, D., Busby, B., and Madden, T.L. (2019) Magic-BLAST, an accurate RNA-seq aligner for long and short reads. *BMC Bioinformatics* **20**: 405.

Brettin, T., Davis, J.J., Disz, T., Edwards, R.A., Gerdes, S., Olsen, G.J. et al. (2015) RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* **5**: 8365.

Callbeck, C.M., Canfield, D.E., Kuypers, M.M.M., Yilmaz, P., Lavik, G., Thamdrup, B. et al. (2021) Sulfur cycling in oceanic oxygen minimum zones. *Limnology and Oceanography* **66**: 2360-2392.

Chaumeil, P.A., Mussig, A.J., Hugenholtz, P., and Parks, D.H. (2022) GTDB-Tk v2: memory friendly classification with the genome taxonomy database. *Bioinformatics* **38**: 5315-5316.

Chen, J.H., Yu, L.J., Boussac, A., Wang-Otomo, Z.Y., Kuang, T., and Shen, J.R. (2019) Properties and structure of a low-potential, penta-heme cytochrome c(552) from a thermophilic purple sulfur photosynthetic bacterium Thermochromatium tepidum. *Photosynth Res* **139**: 281-293.

Dalsgaard, T., Stewart, F.J., Thamdrup, B., De Brabandere, L., Revsbech, N.P., Ulloa, O. et al. (2014) Oxygen at nanomolar levels reversibly suppresses process rates and gene expression in anammox and denitrification in the oxygen minimum zone off northern Chile. *mBio* **5**: e01966.

DeVries, T., Deutsch, C., Rafter, P.A., and Primeau, F. (2013) Marine denitrification rates determined from a global 3-D inverse model. *Biogeosciences* **10**: 2481-2496.

Ehrenreich, P., Behrends, A., Harder, J., and Widdel, F. (2000) Anaerobic oxidation of alkanes by newly isolated denitrifying bacteria. *Arch Microbiol* **173**.

Ettwig, K.F., Speth, D.R., Reimann, J., Wu, M.L., Jetten, M.S., and Keltjens, J.T. (2012) Bacterial oxygen production in the dark. *Frontiers in microbiology* **3**: 273.

Ettwig, K.F., Butler, M.K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M.M.M. et al. (2010) Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* **464**: 543-548.

Frey, C., Bange, H.W., Achterberg, E.P., Jayakumar, A., Löscher, C.R., Arévalo-Martínez, D.L. et al. (2020) Regulation of nitrous oxide production in low-oxygen waters off the coast of Peru. *Biogeosciences* **17**: 2263-2287.

Fuchsman, C.A., Devol, A.H., Saunders, J.K., McKay, C., and Rocap, G. (2017) Niche partitioning of the
N cycling microbial community of an offshore oxygen deficient zone. *Frontiers in microbiology* **8**: 2384.

Ganesh, S., Parris, D.J., DeLong, E.F., and Stewart, F.J. (2014) Metagenomic analysis of size-fractionated
picoplankton in a marine oxygen minimum zone. *ISME J* **8**: 187-211.

Garcia-Robledo, E., Padilla, C.C., Aldunate, M., Stewart, F.J., Ulloa, O., Paulmier, A. et al. (2017)
Cryptic oxygen cycling in anoxic marine zones. *Proc Natl Acad Sci U S A* **114**: 8319-8324.

Gazitúa, M.C., Vik, D.R., Roux, S., Gregory, A.C., Bolduc, B., Widner, B. et al. (2021) Potential virus-
mediated nitrogen cycling in oxygen-depleted oceanic waters. *The ISME Journal* **15**: 981-998.

Geiger, O., Sanchez-Flores, A., Padilla-Gomez, J., and Esposti, M.D. (2023) Multiple approaches of
cellular metabolism define the bacterial ancestry of mitochondria. *Science Advances* **9**: eadh0066.

Glass, J.B., Kretz, C.B., Ganesh, S., Ranjan, P., Seston, S.L., Buck, K.N. et al. (2015) Meta-omic
signatures of microbial metal and nitrogen cycling in marine oxygen minimum zones. *Frontiers in*
Microbiology **6**: 998.

Jain, C., Rodriguez, R.L., Phillippy, A.M., Konstantinidis, K.T., and Aluru, S. (2018) High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* **9**: 5114.

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019) MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. *Briefings in bioinformatics* **20**: 1160-1166.

Keltjens, J.T., Pol, A., Reimann, J., and Op den Camp, H.J. (2014) PQQ-dependent methanol
dehydrogenases: rare-earth elements make a difference. *Appl Microbiol Biotechnol* **98**: 6163-6183.

Kraft, B., Strous, M., and Tegetmeyer, H.E. (2011) Microbial nitrate respiration–genes, enzymes and
environmental distribution. *Journal of biotechnology* **155**: 104-117.

Langmead, B., and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357-
359.

Lutterbeck, H.E., Arévalo-Martínez, D.L., Löscher, C.R., and Bange, H.W. (2018) Nitric oxide (NO) in
the oxygen minimum zone off Peru. *Deep Sea Research Part II: Topical Studies in Oceanography* **156**:
148-154.

Mattes, T.E., Burke, S., Rocap, G., and Morris, R.M. (2022) Two metatranscriptomic profiles through
low-dissolved-oxygen waters (DO, 0 to 33 μM) in the Eastern Tropical North Pacific Ocean.
Microbiology Resource Announcements **11**: e01201-01221.

Moffett, J.W., Goepfert, T.J., and Naqvi, S.W.A. (2007) Reduced iron associated with secondary nitrite
maxima in the Arabian Sea. *Deep Sea Research Part I: Oceanographic Research Papers* **54**: 1341-1349.

Padilla, C.C., Bristow, L.A., Sarode, N., Garcia-Robledo, E., Ramírez, E.G., Benson, C.R. et al. (2016)
NC10 bacteria in marine oxygen minimum zones. *The ISME journal* **10**: 2067-2071.

Richardson, D.J., Butt, J.N., Fredrickson, J.K., Zachara, J.M., Shi, L., Edwards, M.J. et al. (2012) The
'porin-cytochrome' model for microbe-to-mineral electron transfer. *Mol Microbiol* **85**: 201-212.

Schmitz, R.A., Peeters, S.H., Mohammadi, S.S., Berben, T., van Erven, T., Iosif, C.A. et al. (2023)
Simultaneous sulfide and methane oxidation by an extremophile. *Nat Commun* **14**: 2974.

Shi, L., Fredrickson, J.K., and Zachara, J.M. (2014) Genomic analyses of bacterial porin-cytochrome
gene clusters. *Front Microbiol* **5**: 657.

Simon, J., Gross, R., Einsle, O., Kroneck, P.M., Kroger, A., and Klimmek, O. (2000) A NapC/NirT-type
cytochrome c (NrfH) is the mediator between the quinone pool and the cytochrome c nitrite reductase of
Wolinella succinogenes. *Mol Microbiol* **35**: 686-696.

Sousa, F.L., Alves, R.J., Ribeiro, M.A., Pereira-Leal, J.B., Teixeira, M., and Pereira, M.M. (2012) The
superfamily of heme–copper oxygen reductases: Types and evolutionary considerations. *Biochimica et*
Biophysica Acta (BBA) - Bioenergetics **1817**: 629-637.

Thamdrup, B., Steinsdóttir, H.G.R., Bertagnolli, A.D., Padilla, C.C., Patin, N.V., Garcia-Robledo, E. et
al. (2019) Anaerobic methane oxidation is an important sink for methane in the ocean's largest oxygen
minimum zone. *Limnology and Oceanography* **64**: 2569-2585.

Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R. et al. (2012) Differential gene and
transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**: 562-
578.

Trifinopoulos, J., Nguyen, L.T., von Haeseler, A., and Minh, B.Q. (2016) W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* **44**: W232-235.

Tsementzi, D., Wu, J., Deutsch, S., Nath, S., Rodriguez, R.L., Burns, A.S. et al. (2016) SAR11 bacteria
linked to ocean anoxia and nitrogen loss. *Nature* **536**: 179-183.

Tucker, N.P., Le Brun, N.E., Dixon, R., and Hutchings, M.I. (2010) There's NO stopping NsrR, a global
regulator of the bacterial NO stress response. *Trends Microbiol* **18**: 149-156.

Tully, B.J., Graham, E.D., and Heidelberg, J.F. (2018) The reconstruction of 2,631 draft metagenome-
assembled genomes from the global oceans. *Scientific data* **5**: 1-8.

Uzun, M., Alekseeva, L., Krutkina, M., Koziaeva, V., and Grouzdev, D. (2020) Unravelling the diversity
of magnetotactic bacteria through analysis of open genomic databases. *Sci Data* **7**: 252.

Ward, B.B., and Zafiriou, O.C. (1988) Nitrification and nitric oxide in the oxygen minimum of the eastern
tropical North Pacific.

Wasser, I.M., De Vries, S., Moënne-Loccoz, P., Schröder, I., and Karlin, K.D. (2002) Nitric oxide in
biological denitrification: Fe/Cu metalloenzyme and metal complex NO x redox chemistry. *Chemical*
Reviews **102**: 1201-1234.

Woehle, C., Roy, A.S., Glock, N., Wein, T., Weissenbach, J., Rosenstiel, P. et al. (2018) A novel
eukaryotic denitrification pathway in foraminifera. *Curr Biol* **28**: 2536-2543 e2535.

Wu, Y.W., Simmons, B.A., and Singer, S.W. (2016) MaxBin 2.0: an automated binning algorithm to
recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**: 605-607.

Yang, S., Chang, B.X., Warner, M.J., Weber, T.S., Bourbonnais, A.M., Santoro, A.E. et al. (2020) Global
reconstruction reduces the uncertainty of oceanic nitrous oxide emissions and reveals a vigorous seasonal
cycle. *Proc Natl Acad Sci U S A* **117**: 11954-11960.

Yu, N.Y., Wagner, J.R., Laird, M.R., Melli, G., Rey, S., Lo, R. et al. (2010) PSORTb 3.0: improved
protein subcellular localization prediction with refined localization subcategories and predictive
capabilities for all prokaryotes. *Bioinformatics* **26**: 1608-1615.

Zallot, R., Oberg, N., and Gerlt, J.A. (2019) The EFI web resource for genomic enzymology tools:
leveraging protein, genome, and metagenome databases to discover novel enzymes and metabolic
pathways. *Biochemistry* **58**: 4169-4182.

Zedelius, J., Rabus, R., Grundmann, O., Werner, I., Brodkorb, D., Schreiber, F. et al. (2011) Alkane
degradation under anoxic conditions by a nitrate-reducing bacterium with possible involvement of the
electron acceptor in substrate activation. *Environmental microbiology reports* **3**: 125-135.

Zhang, I.H., Sun, X., Jayakumar, A., Fortin, S.G., Ward, B.B., and Babbin, A.R. (2023) Partitioning of
the denitrification pathway and other nitrite metabolisms within global oxygen deficient zones. *ISME*
Commun **3**: 76.

Zumft, W.G. (2005) Nitric oxide reductases of prokaryotes with emphasis on the respiratory, heme–
copper oxidase type. *Journal of Inorganic Biochemistry* **99**: 194-215.
480