Novel Alphaproteobacteria transcribe genes for nitric oxide transformation at high levels in a

marine oxygen deficient zone

Claire E. Elbon[1], Frank J. Stewart[2], and Jennifer B. Glass[1*]

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA;

[2]Department of Microbiology & Cell Biology, Montana State University, Bozeman, MT, USA

*Corresponding author: Jennifer B. Glass, Jennifer.Glass@eas.gatech.edu

Abstract

Marine oxygen deficient zones (ODZs) are portions of the ocean where intense nitrogen loss

occurs primarily via denitrification and anammox. Despite many decades of study, the identity of

the microbes that catalyze nitrogen loss in ODZs are still being elucidated. Intriguingly, high

transcription of genes in the same family as the nitric oxide dismutase (*nod*) gene from

Methylomirabilota has been reported in the anoxic core of ODZs. Here, we show that the most

abundantly transcribed *nod* genes in the Eastern Tropical North Pacific ODZ belong to a new

order (UBA11136) of Alphaproteobacteria, rather than *Methylomirabilota* as previously

assumed. Gammaproteobacteria and Planctomycetia also transcribe *nod*, but at lower relative

abundance than UBA11136 in the upper ODZ. The *nod*-transcribing Alphaproteobacteria likely

use formaldehyde and formate as a source of electrons for aerobic respiration, with additional

electrons possibly from sulfide oxidation. They also transcribe multiheme cytochrome (here

named *ptd*) genes for a putative porin-cytochrome protein complex of unknown function,

potentially involved in extracellular electron transfer. Molecular oxygen for aerobic respiration

may originate from nitric oxide dismutation via cryptic oxygen cycling. Our results implicate

Alphaproteobacteria order UBA11136 as a significant player in marine nitrogen loss and

highlight their potential in one-carbon, nitrogen, and sulfur metabolism in ODZs.

Significance statement

In marine oxygen deficient zones, microbes transform bioavailable nitrogen to gaseous nitrogen,

with nitric oxide as a key intermediate. The Eastern Tropical North Pacific contains the world's

largest oxygen deficient zone, but the identity of the microbes transforming nitric oxide remain

unknown. Here, we show that highly transcribed nitric oxide dismutase *(nod)* genes belong to

Alphaproteobacteria of the novel order UBA11136, which lacks cultivated isolates. These

Alphaproteobacteria show evidence for aerobic respiration, using oxygen potentially sourced

from nitric oxide dismutase, and possess a novel porin-cytochrome protein complex with

unknown function. Gammaproteobacteria and Planctomycetia transcribe *nod* at lower levels. Our

results pinpoint the microbes mediating a key step in marine nitrogen loss and reveal an

unexpected predicted metabolism for marine Alphaproteobacteria.

Introduction

Marine oxygen deficient zones (ODZs) contribute up to half of the ocean's nitrogen loss (DeVries et al., 2013) and are a major source of marine emissions of the potent greenhouse gas nitrous oxide (N_2O) (Yang et al., 2020). The primary source of the N_2O at the oxic-anoxic interface and in anoxic waters in ODZs is denitrification (Babbin et al., 2015; Frey et al., 2020). The microbial enzyme responsible for N_2O production during denitrification is nitric oxide reductase (Nor), which uses electrons from cytochrome *c* (cNor) or quinol (qNor), to reduce nitric oxide (NO) to N_2O (Wasser et al., 2002; Zumft, 2005; Kraft et al., 2011). In the qNor family, there are *bona fide* qNor enzymes and NO dismutase (NOD). NOD proteins lack the quinol-binding site, seemingly preventing the enzyme from taking up external electrons; instead, NOD is theorized to disproportionate NO into dinitrogen and O_2 in methane-oxidizing *Methylomirabilota* bacteria (Ettwig et al., 2010; Ettwig et al., 2012) and in the alkane-oxidizing gammaproteobacterium HdN1 (Zedelius et al., 2011).

The Eastern Tropical North and South Pacific (ETNP and ETSP) ODZs are the world's largest and second largest ODZs, and the subjects of extensive microbial ecology studies. Abundant NO reductase-like genes and transcripts in the ETNP and ETSP ODZ cluster in the same enzyme subfamily as NOD (Dalsgaard et al., 2014; Ganesh et al., 2014; Padilla et al., 2016; Fuchsman et al., 2017). Due to the similarity of ODZ Nod proteins to those of *Methylomirabilota* (NC10), it was initially presumed that ODZ bacteria also used Nod proteins to disproportionate NO into N_2 and O_2 for use in intra-aerobic methane oxidation (Dalsgaard et al., 2014; Padilla et al., 2016; Thamdrup et al., 2019). However, Fuchsman et al. (2017) found that the peak of *nod* gene abundance in the ETNP ODZ correlates with a peak of modeled N_2O production (Babbin et al., 2015) and does not correlate with abundance of methane monooxygenase genes, suggesting

that Nod proteins in the ETNP ODZ are potentially an important source of N_2O, and are unlikely

to be involved in methane oxidation. The plausibility that Nod proteins can reduce NO to N_2O is

supported by a study of a novel eukaryotic denitrification pathway in foraminifera

(*Globobulimina* spp.) that produces N_2O while expressing Nod (Woehle et al., 2018). Yet, the

phylogenetic identity and metabolic context of marine Nod proteins, which are a key biological

source of either N_2O or O_2+N_2 in marine ODZs, remain unresolved.

In this study, we sought to determine the identity, predicted metabolism, and

environmental niche of the ODZ organism responsible for the highly transcribed *nod* genes first

discovered in Padilla et al. (2016). We found that the most abundantly transcribed *nod* genes in

the ETNP ODZ belong to Alphaproteobacteria in the novel order UBA11136. Significant

transcription of *nod* genes was limited to waters with <1 μM O_2. These *nod*-transcribing

alphaproteobacteria also transcribe genes involved in aerobic respiration, which was unexpected

given that they inhabit anoxic waters, as well as genes involved in oxidation of formaldehyde,

likely indicating methylotrophy. Genes encoding multi-heme cytochrome proteins potentially

implicated in nitrogen or iron cycling were also transcribed.

Results

Transcribed* nod *sequences in the ETNP ODZ belong to Alphaproteobacteria,

Gammaproteobacteria, and Planctomycetia. Our reanalysis of highly transcribed *nod* genes in

the ETNP ODZ (Padilla et al., 2016) shows that these genes belong to Alphaproteobacteria rather

than a member of *Methylomirabilota* as previously assumed. Querying the Nod amino acid

sequences from Padilla et al. (2016) against ETNP ODZ metagenomes in the IMG-JGI database

returned multiple 100% identity matches, including a *nod* gene (Ga0066848_100037855) on a

scaffold with hypothetical genes with 100% identity to Alphaproteobacteria metagenome-

assembled genomes (MAGs) from the ETNP ODZ (Uzun et al., 2020) **(Table S1)**. We binned

previously sequenced ETNP ODZ metagenomes Ga0066848 (ETNP201310SV72) and

Ga0066829 (ETNP201306SV43) (Ruiz-Perez et al., 2021) into MAGs. Contigs with the most

highly transcribed *nod* genes were present in two Alphaproteobacteria MAGs (GTDB taxonomy:

UBA11136 sp002686135; species representative: Rhodospirillaceae bacterium isolate ARS27)

with 97% average nucleotide identity. Querying the Nod amino acid sequences from Padilla et

al. (2016) against NCBI's non-redundant protein database returned matches to other MAGs

assigned to Alphaproteobacteria order UBA11136 from low-oxygen marine settings (ETNP,

Saanich Inlet, and the Black Sea; 78-80% identity), the marine magnetotactic

alphaproteobacterium *Magnetovibrio blakemorei* MV-1 (75% identity), Gammaproteobacterium

HdN1 (66% identity), and *Methylomirabilota* spp. (66% identity; **Table S2**).

To glean additional insights into evolutionary relationships, we updated a previous Nod

phylogeny (Hu et al., 2019) with additional amino acid sequences from marine MAGs (Tully et

al., 2018; Cabello-Yeves et al., 2021; Lin et al., 2021) and ETNP ODZ metagenomes (Ruiz-

Perez et al., 2021), subdivided in free-living cells (0.2-1.6 micron) and cells from the particle

fraction (>1.6 micron; **Figure 1A; Table S3**). The Nod topology was generally consistent with a

previous phylogeny from Fuchsman et al. (2017), with additional taxonomic data from MAGs in

the TARA oceans dataset further constraining Nod placement (Tully et al., 2018). As expected

based on the binning and BLAST results, the Nod sequence from Padilla et al. (2016)

(Ga0066848_100037855) clustered phylogenetically with marine Alphaproteobacteria (OTU III

in Fuchsman et al. (2017), hereafter "Alpha-type Nod"); this clade contained three unique

sequences, all of which were present in multiple metagenomes and all from the free-living

fraction, and one of which was identical to that of Rhodospirillaceae NP1106 (MBV28360). Four

unique ODZ Nod sequences clustered with marine Gammaproteobacteria (OTU II in Fuchsman

et al. (2017), hereafter "Gamma-type Nod"); these sequences were monophyletic with a cluster

of Gammaproteobacteria Nod cluster sequences from sewage sludge, including

Gammaproteobacterium HdN1 (Ehrenreich et al., 2000) and other wastewater

Gammaproteobacteria. Multiple ETNP ODZ metagenomes contained Gamma-type Nod

sequences identical to those of Gammaproteobacteria NP964 (MBP20251). Gamma-type Nod

had ~70% identity to Alpha-type Nod. Several ODZ Nod sequences, all from the particle fraction

("PF") clustered with marine Deltaproteobacteria in a clade of monophyletic *nod* genes from

groundwater Methylomirabilis, Deltaproteobacteria, and Acidobacteria MAGs (~65% identity to

Alpha-type Nod). Six unique Nod ODZ protein sequences (two of which were present in

multiple metagenomes) clustered with Planctomycetia (OTU I in Fuchsman et al. (2017),

hereafter "Planctomycetia-type Nod"), and were primarily found in free-living cells ("FL") and

had ~40% identity to Alpha-type Nod. Intriguingly, two ODZ sequences clustered in the

eukaryotic *Globobulimina* clade (~50% identity to Alpha-type Nod). Viral Nod sequences from

Saanich Inlet (~55% identity to Alpha-Nod) clustered with the viral Nod sequence previously

reported by Gazitúa et al. (2021) from the Eastern Tropical South Pacific ODZ (St16 OMZ 317E

– viral).

We investigated gene neighborhoods surrounding ODZ *nod* genes in the three main

phylogenetic clusters of ODZ sequences: Planctomycetia-type Nod, Gamma-type Nod, and

Alpha-type Nod. Whereas "unknown Nor-related" marine Bacteroidota sequences were located

on an operon with other *nor* genes, there was no consistent gene neighborhood for *nod* sequences

(Figure 1B). Planctomycetia-type *nod* genes were not located in the vicinity of any genes with

recognizable related function. Gamma-type *nod* gene neighborhoods contained ferredoxins and

cytochrome b_{561} genes for electron transport. Upstream of the Alpha-type *nod* in

Rhodospirillaceae NP1106 is a cluster of formylmethanofuran dehydrogenase genes (*fmd/fwd*)

used in C1 metabolism via tetrahydromethanopterin/methanofuran-linked reactions.

Immediately upstream or downstream of *nod* genes, helix-turn-helix transcriptional

regulators were common (**Figure 1B**). Neighboring Gamma-type and *Methylomirabilis nod*

genes, LuxR-type regulators were common; these regulators have diverse functions and their

potential connection to Nod remains unclear. Neighboring Alpha-type and Bacteroidota (e.g.

Cecembia calidifontis) *nod* genes, Rrf2-type regulators were present. The protein NsrR in the

Rrf2 family regulates global cellular response to NO toxification by directly sensing NO with an

iron-sulfur cluster (Bodenmiller and Spiro, 2006; Tucker et al., 2010). The presence of this

NsrR-like regulator suggests that Nod in marine Alphaproteobacteria and Bacteroidota may be

involved in nitrosative stress response and NO detoxification.

Alphaproteobacterial* nod *is highly transcribed in anoxic waters. We assessed

transcription of Alpha, Gamma-, and Planctomycetia-type *nod* genes from the oxycline to upper

ODZ (secondary nitrite maximum) using ETNP ODZ metatranscriptomes from an onshore

station with a shallower oxycline (P1; **Figure 1C**) and an offshore station with a deeper oxycline

(P2; **Figure 1D**) (Mattes et al., 2022). In both oxyclines, transcription was low (4-10 reads per

kilobase per million mapped reads (RPKM), n=8) for all three *nod* types (**Figure 1C, D**). Below

the oxyclines, *nod* transcripts began to rise and were highest at the secondary nitrite maxima,

with Alpha-type (184-274 RPKM, n=4) > Gamma-type (55-95 RPKM, n=4) > Planctomycetia-

type (13-19 RPKM, n=4; **Table S4**).

MAGs with highly transcribed* nod *gene represent a new order of Alphaproteobacteria.

In order to assess the phylogeny of the *nod*-containing Alphaproteobacteria MAGs, we

constructed an alphaproteobacterial phylogeny using the conserved protein NADH ubiquinone

oxidoreductase subunit L (NuoL) as in Cevallos and Degli Esposti (2022), with additional

representation of order UBA11136 including our MAG ETNP2013_S10_300m_22 (**Figure 2**).

MAG ETNP2013_S06_300m_15 was not included in the phylogeny because its *nuoL* gene was

truncated. UBA11136 is situated in the phylogeny near other orders found in ODZs.

Alphaproteobacteria transcribe genes for formate metabolism, aerobic respiration, and

a multiheme cytochrome complex. To glean insight into potential roles for Nod in cellular

context, we sought to reconstruct the electron transport chain of the Alphaproteobacteria with the

most highly transcribed *nod* genes (Alphaproteobacterium MAG ETNP2013_S10_300m_22 and

Alphaproteobacterium MAG ETNP2013_S06_300m_15, 73% and 69% estimated completeness,

respectively) at the secondary nitrite maximum. Of total metagenomic reads, 0.38% map to

ETNP2013_S10_300m_22 and 0.39% map to ETNP2013_S06_300m_15. In both MAGs, *nod*

was in the top three most transcribed genes in the ETNP ODZ (~44,000 FPKM; **Table S5**), after

a bacterial nucleoid DNA-binding protein and a potassium gated channel protein. In addition to

nod,   we   found   that   genes   for   formaldehyde   oxidation   via

tetrahydromethanopterin/methanofuran-linked   reactions,   including   formylmethanofuran

dehydrogenase   (*fwd/fmd*)   and   formylmethanofuran--tetrahydromethanopterin   N-

formyltransferase (*ftr*), were transcribed in both MAGs (**Table S5**). Both MAGs also transcribed

NAD-dependent formate dehydrogenase (**Table S5**). Thus, the alphaproteobacterium appears to

be capable of conversion of formaldehyde to formate and use of formate as a source of electrons

for NADH:ubiquinone oxidoreductase (Complex I; **Figure 3**). The source of formaldehyde is

likely methanol oxidation, as pyrroloquinoline quinone (PQQ)-dependent ethanol/methanol

dehydrogenases were found in Alphaproteobacteria MAGs from low-oxygen marine settings

(Table S6). Methane monooxygenase genes were not found in the partial Alphaproteobacteria

MAGs, precluding our ability to rule out the possibility of these genes in the missing portions of

the genomes. The Alphaproteobacteria PQQ-dependent dehydrogenase genes contained the motif

DYDG **(Table S6)**, which is characteristic of the lanthanide-containing form of the enzymes

rather than calcium form (Keltjens et al., 2014).

A full aerobic electron transport chain (Complex I, II, III, and IV) and F0F1-type ATP

synthase were transcribed in both bins **(Figure 3; Table S5)**. Complex IV (cytochrome c

oxidase) was type A1 according to the Sousa et al. (2012) classification, and the *cox* operon in

the GTDB species representative Rhodospirallaceae ARS27 was subtype b (COX2-COX1-CtaB-

CtaG_Cox11-COX3-DUF983-SURF1-CtaA1-M32-Tsy-M16B) according to the Geiger et al.

(2023) classification. Sulfur oxidation genes, including flavocytochrome c sulfide dehydrogenase

(FccAB), sulfane hydrogenase (SoxCD), and carrier protein SoxYZ, were also transcribed, as

were numerous transposes **(Figure 3; Table S5)**.

Genes for a multiheme cytochrome complex were transcribed in both bins. To our

knowledge, this putative operon has not previously been described. Hereafter, we designate it the

ptdABCDEFG operon for its sequence of penta/tetra/deca-heme proteins, interspersed with other

conserved proteins. *ptdAB* genes are highly transcribed in our Alphaproteobacteria MAGs, but it

is unclear if the rest of the operon is also highly transcribed, because it was truncated in our

MAGs' scaffolds. The *ptd* gene cluster consists of a penta-heme protein with a C-terminal beta-

sandwich (PtdA), a porin (PtdB), a FAD/NAD(P)-binding oxidoreductase (PtdC), a periplasmic

tetra-heme protein (PtdD), a cyclic nucleotide-binding domain protein with two 4Fe-4S clusters

(PtdE), a cytoplasmic transmembrane ferric reductase-like protein (PtdF), and a periplasmic

deca-heme protein (PtdG; **Figure 3; Tables S7, S8)**. The function of this complex is unknown,

but the presence of genes encoding a porin and multiple multiheme proteins resembles porin-

cytochrome protein complexes involved in extracellular reduction electron transfer during Fe(III)

and Mn(IV) reduction (Richardson et al., 2012; Shi et al., 2014). PtdA has a homolog to a penta-

heme cytochrome c_{552} protein of unknown function in a thermophilic purple sulfur

gammaproteobacterium (Chen et al., 2019) and is in the same COG family (COG3303) as

formate dependent nitrite reductase, NrfA. *ptdABCDEFG* genes were prevalent in

Alphaproteobacteria, Gammaproteobacteria, Nitrospirales, and Planctomycetes MAGs from

marine or high salinity environments **(Figure 4; Table S7)**.

Discussion

This study predicts the previously ambiguous identity of the microorganisms that make the

dominant nitric oxide-transforming protein (Nod) in the world's largest ODZ, the Eastern

Tropical North Pacific. Extensive horizontal gene transfer of *nod* genes between microbial

genomes is evident from the lack of conservation of gene neighborhood and patchy phylogeny

(Fuchsman et al., 2017), which may be mediated by viral infection (Gazitúa et al., 2021). We

found that the most transcriptionally active *nod* genes in the ETNP upper ODZ belong to the

novel Alphaproteobacteria order UBA11136. Alpha-type *nod* transcript abundances (~200

RPKM) are similar to those of dissimilatory nitrate reductase (*narG*) in the ODZ (Tsementzi et

al., 2016). The *nod*-transcribing Alphaproteobacteria are also transcribing genes for

formaldehyde oxidation, likely as a source of electrons to the respiratory chain via NAD

reduction by formate dehydrogenase. Sulfide may be used as a supplemental electron donor

and/or may be concomitantly oxidized for detoxification (Callbeck et al., 2021; Schmitz et al.,

2023).

Our discovery of a putative porin-cytochrome complex (*ptd* operon) in marine bacteria

was unexpected. Porin-cytochrome complexes have been best studied for their role in

extracellular electron transport, particularly for respiratory metal reduction and oxidation

(Richardson et al., 2012; Shi et al., 2014). It is conceivable that the Ptd complex is involved in

iron reduction in ODZs; there is iron reduction at the secondary nitrite maximum and it is

hypothesized to be bacterially mediated, but the microbes involved have yet to be determined

(Moffett et al., 2007; Glass et al., 2015). Alternatively, the presence of *ptdABCDEFG* genes in

numerous nitrite-oxidizing bacteria (Nitrospirales) could imply the involvement of these genes in

nitrogen cycling; PtdA was in the same COG family as formate-dependent nitrite reductase

(Simon et al., 2000) and PtdC is similar to a flavohemoprotein with predicted nitric oxide

dioxygenase activity, also annotated as hydroxylamine oxidoreductase-linked cytochrome. The

function of PtdABCDEFG remains completely unknown and requires future biochemical

characterization.

On the other end of the electron transport chain, high transcription of a heme/copper

terminal oxidase suggests that O_2 is being used as the terminal election acceptor in *nod*-

transcribing Alphaproteobacteria MAGs. The transcribed heme/copper oxidase is A1-type (low

O_2 affinity), also present in mitochondria, and adapted for high O_2 concentrations. Low O_2

affinity A1-type heme/copper oxidases are transcribed in other anoxic environments (Berg et al.,

2022). Because ODZs have extremely low concentrations of molecular oxygen below the

oxycline, O_2 for aerobic respiration may be generated *in situ* and rapidly consumed. Given that

the function of Nod is proposed to be dismutation of two NO molecules into N_2 and O_2 (Ettwig et

al., 2010), it is possible that the O_2 source for aerobic respiration in the UBA11136 MAGs is NO

dismutation, although other sources of O_2 (e.g. *in situ* photosynthesis, mixing) in anoxic waters

are also conceivable (Garcia-Robledo et al., 2017). The physiological uses of Gamma-type and

Planctomycetia-type Nod may be different from Alpha-type Nod, although this remains to be

investigated.

The source of NO, the presumed substrate for Nod, may be generated in the same

organism using Nod, or generated by a different organism (or chemical pathway). Nitric oxide

was positively correlated with nitrite in the Eastern Tropical South Pacific ODZ, and was only

detectable when O_2 was <1-2 µM (Lutterbeck et al., 2018). In the Eastern Tropical North Pacific,

NO concentration and turnover rates were elevated at O_2 <100 µM (Ward and Zafiriou, 1988).

Both studies suggest that the NO in ODZs likely originates from nitrification or nitrifier

denitrification, while genomic analyses indicate that the copper-containing nitrite reductase

(*nirK*) in SAR11 bacteria (presumably performing denitrification) may be a key source of NO

(Fuchsman et al., 2017). Because most ODZ denitrifiers specialize in only one of the three steps

(NO_2^- reduction, NO reduction, and N_2O reduction) (Zhang et al., 2023), and known nitrite

reductases were not identified in our MAGs, existing data indicate that the NO that is used as a

substrate for alphaproteobacterial Nod is not generated *in vivo*. (Only 4 out of 32 *nod*-containing

MAGs contained a nitrite reductase gene: two Gammaproteobacteria MAGs contained *nirK*, one

Myxococcota MAG contained *nirS*, and one Scalindua MAG contained *nirS*). It is also possible

that another uncharacterized enzyme produces NO.

This study suggests that marine Alphaproteobacteria from order UBA11136 are actively

reducing NO under anoxia, as implied by their abundant transcription of *nod* genes. While there

is strong evidence that the substrate for Nod in ODZs is NO based on its abundance, the products

of this enzyme (N_2O vs. N_2+O_2) remain uncertain. Nod is theorized to disproportionate NO into

N_2 and O_2 in methane-oxidizing *Methylomirabilota* bacteria (Ettwig et al., 2010; Ettwig et al.,

2012), but no biochemical characterizations of Nod have been published to date, and

foraminifera expressing Nod produce N_2O (Woehle et al., 2018). The apparent lack of other

denitrification genes in *nod*-transcribing Alphaproteobacteria is consistent with the observation

that denitrification in ODZs is largely divided into distinct microbial taxa (Dalsgaard et al., 2014;

Fuchsman et al., 2017; Zhang et al., 2023). For example, although nitrate reductase (*narG*) genes

are widely distributed amongst ODZ microbes (Zhang et al., 2023), SAR11 bacteria appear to

dominate in *narG* transcriptional activity (Tsementzi et al., 2016). Our finding that the

transcription of *nod* is catalyzed primarily by marine Alphaproteobacteria implies that this taxa

contributes significantly to marine nitrogen loss.

METHODS

Nod *phylogeny and gene neighborhood*. Amino acid sequences of highly transcribed *nod* genes

"ETNP 2014 Stn10 150m" and "ETNP 2013 Stn6 300m" were acquired from the authors of

Padilla et al. (2016) (see Table S2 for sequences). These sequences were used for BLASTP

searches of ODZ metagenomes in the IMG-JGI database and the non-redundant protein (nr)

database in NCBI. Sequences (n=53, 731 gap-free sites) were aligned using the MAFFT online

server with the L-INS-i method (Katoh et al., 2019). A phylogeny was generated with 1000

bootstraps using model LG+I+G4 using W-IQ-Tree (Trifinopoulos et al., 2016). The phylogeny

was visualized using FigTree v.1.4.4, and the fasta file (Nod_alignment) is available as a

supplemental dataset. Gene neighborhoods were generated using the EFI Gene Neighborhood

Tool (Zallot et al., 2019) with single sequence BLAST of the UniProt database using the amino

acid sequence Ga0066848_100037855 (JGI IMG) as the Nod query with an e-value cutoff of 10^{-}

^{5} and with 10 genes upstream and downstream the gene of interest.

Transcription of* nod *genes in ETNP ODZ depth profiles. Magic Basic Local Alignment

Search Tool (Boratyn et al., 2019) was used to search ETNP ODZ metatranscriptomes

(PRJNA727903; Mattes et al. (2022)) using representative nucleotide sequences for

Planctomycetia-like   (Ga0066826_100064333   (JGI   IMG)),   Gamma-like

(PBRC01000062.1:19833-22205 (NCBI)), and Alpha-like (Ga0066848_100037855 (JGI IMG))

nod genes. Default parameters were used except for the score threshold (18). Read hits were

normalized to reads per kilobase million (RPKM).

***Metagenomic binning*.** Binning of metagenome-assembled genomes (MAGs) was

performed using the KBase platform (Arkin et al., 2018). ETNP ODZ metagenomes were

collected in 2013 and sequenced by Joint Genome Institute (JGI) using an Illumina HiSeq 2500

as described in Ruiz-Perez et al. (2021). Assemblies for the ETNP ODZ metagenomes (Ruiz-

Perez et al., 2021) containing Alpha-type *nod* genes (ETNP201310SV72 (GOLD Analysis

Project ID Ga0066848; stn10 300m) and ETNP201306SV43 (GOLD Analysis Project ID

Ga0066829; stn6 300m) were imported from JGI IMG into Kbase. Metagenomic assemblies

were binned into MAGs using MaxBin2 v2.2.4 (Wu et al., 2016). The two MAGs containing *nod*

genes   (MAG   ETNP2013_S10_300m_22   from   ETNP201310SV72   and

ETNP2013_S06_300m_15 from ETNP201306SV43) were selected for further analysis. Average

nucleotide identity was calculated using FastANI (Jain et al., 2018). MAG taxonomy and

genome quality was evaluated by GTDB-Tk v2.3.2 (Chaumeil et al., 2022). MAGs were

annotated with RASTtk v1.073 (Brettin et al., 2015). Metagenomic reads were mapped to MAGs

using Bowtie2 (Langmead and Salzberg, 2012).

Alphaproteobacterial NuoL phylogeny. Alphaproteobacterial NADH ubiquinone

oxidoreductase subunit L (NuoL) and mitochondrial ND5 marker proteins (n=320) were aligned

as in Cevallos and Degli Esposti (2022) with additional representation of UBA11136. A

maximum likelihood phylogeny with 1000 bootstraps was constructed using IQ-tree (Nguyen et

al., 2015) using the LG+F model with ultrafast bootstrap (Hoang et al., 2017). Taxonomic names

and clades are from Cevallos and Esposti (2022) and GTDB Release 08-RS214. The fasta file

(NuoL_alignment) is available as a supplemental dataset. Alphaproteobacteria MAGs containing

nod genes (Table S2) were classified as belonging to order UBA11136 using GTDB-Tk v2.3.2

(Chaumeil et al., 2022).

Mapping transcripts to metagenomic bins. Metatranscriptomic mapping to MAGs was

performed using the KBase platform (Arkin et al., 2018). RNA-seq data (Mattes et al., 2022)

were imported from the depth with highest *nod* transcription, the secondary nitrite maximum

(126 meters, NCBI run SRR14460584) and aligned to MAGs using the Bowtie2 (Langmead and

Salzberg, 2012) app in KBase. The Cufflinks v2.2.1 (Trapnell et al., 2012) app in Kbase was

then used to assemble the aligned RNA-seq data into a set of transcripts and to calculate the

relative abundances of the transcripts expressed in fragments per kilobase of exon per million

fragments mapped (FPKM).

Cellular localization and heme numbers. Cellular locations of Ptd proteins were

predicted using PSORTb v3.0.3 analysis (Yu et al., 2010). Numbers of heme-binding motifs per

protein were identified by counting CXXCH sequences. Ptd gene neighborhoods was generated

using the EFI Gene Neighborhood Tool (Zallot et al., 2019) with single sequence BLAST of the

UniProt database using the amino acid sequence Ga0066848_100031354 (JGI IMG) as the PtdA

query, with an e-value cutoff of 10^{-5} and with 10 genes upstream and downstream the gene of

interest.

Data availability

The Kbase bioinformatic pipeline and MAGs are at https://narrative.kbase.us/narrative/106999.

Original metagenomic reads are available at BioProject PRJNA375524 (ETNP201306SV43,

SAMN06344130) and BioProject PRJNA375542 (ETNP201310SV72, SAMN06344148). MAG

Alphaproteobacteria bacterium ETNP2013_S06_300m_15 was deposited into BioProject

PRJNA375524 (BioSample SAMN38229257 , WGS Accession JAZDBU000000000) and

Alphaproteobacteria bacterium ETNP2013_S10_300m_22 was deposited into BioProject

PRJNA375542 (BioSample SAMN38228782, WGS Accession JAZDCE000000000). All

ETNP ODZ datasets used in this manuscript are listed in **Table S9**.

Acknowledgements

We thank Laura Bristow for helpful discussions. We thank Mauro Degli Esposti for assistance

with the NADH ubiquinone oxidoreductase subunit L phylogeny. We thank Cory Padilla,

Anthony Bertagnolli, and Neha Sarode for sharing previous data. We acknowledge funding from

an NSF Graduate Research Fellowship to CEE, the Simons Foundation, and NSF

Awards 2022991 and 2054927 to FJS.

Figure 1. Marine Nod clades, gene neighborhoods, and depth profiles of transcription. (A)

Maximum likelihood phylogeny of nitric oxide dismutase (Nod) amino acid sequences in marine

(blue) and select terrestrial (brown) taxa, primarily from marine MAGs (Tully et al., 2018;

Cabello-Yeves et al., 2021; Lin et al., 2021) and ETNP ODZ metagenomes (Ruiz-Perez et al.,

2021). Branch support was evaluated using 1000 rapid bootstrap replicates, with bootstrap values

shown for deep branches. The tree is drawn to scale, with branch lengths in number of

substitutions per site. Bold sequences represent those present in multiple ETNP ODZ

metagenomes (see **Table S3** for duplicate accession numbers). "PF" indicates genes from the

particle fraction (>1.6 micron fraction) of filters. "FL" indicates genes from the free-living

fraction (0.2-1.6 micron) collected on Sterivex filters. The most highly transcribed ETNP ODZ

sequence is indicated with an asterisk. The qNor sequence *Geobacillus stearothermophilus* was

used as the outgroup. (B) Gene neighborhoods surrounding *nod* genes in select taxa. GenBank

contigs: *Cecembia calfifontis* SGXG01000001, *Scalindua japonica* BAOS01000045,

Gammaproteobacteria NP964 PBRC01000062, Gammaproteobacterium HdN1 FP929140,

Deltaproteobacteria NZCL01000067, *Candidatus* Methylomirabilis oxyfera FP565575, and

Rhodospirillaceae NP1106 PCBZ01000014. Unlabeled gray genes are hypothetical. (C, D)

Oxygen and nitrite concentrations (circles), and *nod* transcripts (squares, as reads per kilobase

per million mapped reads (RKPM)) with depth in ETNP ODZ P1 (onshore) and P2 (offshore)

sites (Mattes et al., 2022).

Figure 2. Phylogeny of Alphaproteobacteria showing order UBA11136, with *nod*-containing MAGs. The phylogeny was constructed using the Alphaproteobacterial phylogenetic marker NADH ubiquinone oxidoreductase subunit L as in Cevallos and Esposti (2022). Order UBA11136 is expanded and *nod*-containing MAGs are bolded. Taxonomic names are from Cevallos and Esposti (2022) and GTDB Release 08-RS214. The scale bar represents amino acid substitutions per site. The full phylogeny is shown in **Supplementary Figure S1**.

Figure 3. Schematic of the electron transport chain in *nod*-containing ODZ Alphaproteobacteria. Enzymes were included based on presence and transcriptional activity of metagenome-assembled genomes (MAGs) assigned to Alphaproteobacteria (GTDB taxonomy: UBA11136 sp002686135; see text). The color of each protein is chosen according to transcriptional activity and represented from 0-100, 100-1,000, 1,000-10,000, and 10,000-100,000 FPKM in gradient from lighter to darker blue (Table S5). Heme proteins are indicated

by red circular hemes with the cartoon number corresponding to the number of actual hemes

present on each protein. Hypothetical Ptd proteins are labelled A, B, C, D, E, F, and G, and

location within the cell is determined using Psort bacterial localization prediction tool (Table

S8). ETC complexes I-V found in Alphaproteobacteria MAGs are labelled with proposed

interactions between formate oxidation and complex I NADH electron transfer. Highly

transcribed NOD protein and predicted O_2 generation is shown as feeding into A1 type CCO

complex IV reduction. Additional electrons for CytC and the ETC are proposed to come from

sulfur oxidation carried out by the flavocytochrome c sulfide dehydrogenase (FccAB, FCC), and

sulfane-sulfur dehydrogenase (SoxCD) with the multi-enzyme carrier complex (SoxYZ).

Figure 4. Gene neighborhoods of petaheme-tetraheme-decaheme genes from select organisms. Depicted heme spacing is approximate. All organisms are from saline environments (seawater, marine sediment, or saline spring).

References

Arkin, A.P., Cottingham, R.W., Henry, C.S., Harris, N.L., Stevens, R.L., Maslov, S. et al. (2018) KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* **36**: 566-569.

Babbin, A.R., Bianchi, D., Jayakumar, A., and Ward, B.B. (2015) Rapid nitrous oxide cycling in the suboxic ocean. *Science* **348**: 1127-1129.

Berg, J.S., Ahmerkamp, S., Pjevac, P., Hausmann, B., Milucka, J., and Kuypers, M.M.M. (2022) How low can they go? Aerobic respiration by microorganisms under apparent anoxia. *FEMS Microbiol Rev* **46**: 1–14.

Bodenmiller, D.M., and Spiro, S. (2006) The *yjeB* (*nsrR*) gene of *Escherichia coli* encodes a nitric oxide-sensitive transcriptional regulator. *J Bacteriol* **188**: 874-881.

Boratyn, G.M., Thierry-Mieg, J., Thierry-Mieg, D., Busby, B., and Madden, T.L. (2019) Magic-BLAST, an accurate RNA-seq aligner for long and short reads. *BMC Bioinf* **20**: 405.

Brettin, T., Davis, J.J., Disz, T., Edwards, R.A., Gerdes, S., Olsen, G.J. et al. (2015) RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* **5**: 8365.

Cabello-Yeves, P.J., Callieri, C., Picazo, A., Mehrshad, M., Haro-Moreno, J.M., Roda-Garcia, J.J. et al. (2021) The microbiome of the Black Sea water column analyzed by shotgun and genome centric metagenomics. *Environ Microbiome* **16**: 5.

Callbeck, C.M., Canfield, D.E., Kuypers, M.M.M., Yilmaz, P., Lavik, G., Thamdrup, B. et al. (2021) Sulfur cycling in oceanic oxygen minimum zones. *Limnol Oceanogr* **66**: 2360-2392.

Cevallos, M.A., and Degli Esposti, M. (2022) New Alphaproteobacteria thrive in the depths of the ocean with oxygen gradient. *Microorganisms* **10**: 455.

Chaumeil, P.A., Mussig, A.J., Hugenholtz, P., and Parks, D.H. (2022) GTDB-Tk v2: memory friendly classification with the genome taxonomy database. *Bioinformatics* **38**: 5315-5316.

Chen, J.H., Yu, L.J., Boussac, A., Wang-Otomo, Z.Y., Kuang, T., and Shen, J.R. (2019) Properties and structure of a low-potential, penta-heme cytochrome c(552) from a thermophilic purple sulfur photosynthetic bacterium Thermochromatium tepidum. *Photosynth Res* **139**: 281-293.

Dalsgaard, T., Stewart, F.J., Thamdrup, B., De Brabandere, L., Revsbech, N.P., Ulloa, O. et al. (2014) Oxygen at nanomolar levels reversibly suppresses process rates and gene expression in anammox and denitrification in the oxygen minimum zone off northern Chile. *mBio* **5**: e01966.

DeVries, T., Deutsch, C., Rafter, P.A., and Primeau, F. (2013) Marine denitrification rates determined from a global 3-D inverse model. *Biogeosciences* **10**: 2481-2496.

Ehrenreich, P., Behrends, A., Harder, J., and Widdel, F. (2000) Anaerobic oxidation of alkanes by newly isolated denitrifying bacteria. *Arch Microbiol* **173**.

Ettwig, K.F., Speth, D.R., Reimann, J., Wu, M.L., Jetten, M.S., and Keltjens, J.T. (2012) Bacterial
oxygen production in the dark. *Front Microbiol* **3**: 273.

Ettwig, K.F., Butler, M.K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M.M.M. et al. (2010)
Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* **464**: 543-548.

Frey, C., Bange, H.W., Achterberg, E.P., Jayakumar, A., Löscher, C.R., Arévalo-Martínez, D.L. et al.
(2020) Regulation of nitrous oxide production in low-oxygen waters off the coast of Peru. *Biogeosciences*
17: 2263-2287.

Fuchsman, C.A., Devol, A.H., Saunders, J.K., McKay, C., and Rocap, G. (2017) Niche partitioning of the
N cycling microbial community of an offshore oxygen deficient zone. *Front Microbiol* **8**: 2384.

Ganesh, S., Parris, D.J., DeLong, E.F., and Stewart, F.J. (2014) Metagenomic analysis of size-fractionated
picoplankton in a marine oxygen minimum zone. *ISME J* **8**: 187-211.

Garcia-Robledo, E., Padilla, C.C., Aldunate, M., Stewart, F.J., Ulloa, O., Paulmier, A. et al. (2017)
Cryptic oxygen cycling in anoxic marine zones. *Proc Natl Acad Sci U S A* **114**: 8319-8324.

Gazitúa, M.C., Vik, D.R., Roux, S., Gregory, A.C., Bolduc, B., Widner, B. et al. (2021) Potential virus-
mediated nitrogen cycling in oxygen-depleted oceanic waters. *ISME J* **15**: 981-998.

Geiger, O., Sanchez-Flores, A., Padilla-Gomez, J., and Esposti, M.D. (2023) Multiple approaches of
cellular metabolism define the bacterial ancestry of mitochondria. *Sci Adv* **9**: eadh0066.

Glass, J.B., Kretz, C.B., Ganesh, S., Ranjan, P., Seston, S.L., Buck, K.N. et al. (2015) Meta-omic
signatures of microbial metal and nitrogen cycling in marine oxygen minimum zones. *Front Microbiol* **6**:
998.

Hoang, D.T., Chernomor, O., Haeseler, A.v., Minh, B.Q., and Vinh, L.S. (2017) UFBoot2: improving the
ultrafast bootstrap approximation. *Mol Biol Evol* **35**: 518–522

Hu, Q.-Q., Zhou, Z.-C., Liu, Y.-F., Zhou, L., Mbadinga, S.M., Liu, J.-F. et al. (2019) High microbial
diversity of the nitric oxide dismutation reaction revealed by PCR amplification and analysis of the nod
gene. *Int Biodeterior Biodegradation* **143**: 104708.

Jain, C., Rodriguez, R.L., Phillippy, A.M., Konstantinidis, K.T., and Aluru, S. (2018) High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* **9**: 5114.

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019) MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. *Brief Bioinform* **20**: 1160-1166.

Keltjens, J.T., Pol, A., Reimann, J., and Op den Camp, H.J. (2014) PQQ-dependent methanol
dehydrogenases: rare-earth elements make a difference. *Appl Microbiol Biotechnol* **98**: 6163-6183.

Kraft, B., Strous, M., and Tegetmeyer, H.E. (2011) Microbial nitrate respiration–genes, enzymes and
environmental distribution. *J Biotech* **155**: 104-117.

Langmead, B., and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357-
359.

Lin, H., Ascher, D.B., Myung, Y., Lamborg, C.H., Hallam, S.J., Gionfriddo, C.M. et al. (2021) Mercury
methylation by metabolically versatile and cosmopolitan marine bacteria. *ISME J* **15**: 1810-1825.

Lutterbeck, H.E., Arévalo-Martínez, D.L., Löscher, C.R., and Bange, H.W. (2018) Nitric oxide (NO) in
the oxygen minimum zone off Peru. *Deep Sea Res Part II Top Stud Oceanogr* **156**: 148-154.

Mattes, T.E., Burke, S., Rocap, G., and Morris, R.M. (2022) Two metatranscriptomic profiles through
low-dissolved-oxygen waters (DO, 0 to 33 μM) in the Eastern Tropical North Pacific Ocean. *Microbiol*
Resour Announc **11**: e01201-01221.

Moffett, J.W., Goepfert, T.J., and Naqvi, S.W.A. (2007) Reduced iron associated with secondary nitrite
maxima in the Arabian Sea. *Deep Sea Res Part I Oceanogr Res Pap* **54**: 1341-1349.

Nguyen, L.T., Schmidt, H.A., von Haeseler, A., and Minh, B.Q. (2015) IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**: 268-274.

Padilla, C.C., Bristow, L.A., Sarode, N., Garcia-Robledo, E., Ramírez, E.G., Benson, C.R. et al. (2016)
NC10 bacteria in marine oxygen minimum zones. *ISME J* **10**: 2067-2071.

Richardson, D.J., Butt, J.N., Fredrickson, J.K., Zachara, J.M., Shi, L., Edwards, M.J. et al. (2012) The
'porin-cytochrome' model for microbe-to-mineral electron transfer. *Mol Microbiol* **85**: 201-212.

Ruiz-Perez, C.A., Bertagnolli, A.D., Tsementzi, D., Woyke, T., Stewart, F.J., and Konstantinidis, K.T.
(2021) Description of Candidatus Mesopelagibacter carboxydoxydans and Candidatus Anoxipelagibacter
denitrificans: Nitrate-reducing SAR11 genera that dominate mesopelagic and anoxic marine zones. *Syst*
Appl Microbiol **44**: 126185.

Schmitz, R.A., Peeters, S.H., Mohammadi, S.S., Berben, T., van Erven, T., Iosif, C.A. et al. (2023)
Simultaneous sulfide and methane oxidation by an extremophile. *Nat Commun* **14**: 2974.

Shi, L., Fredrickson, J.K., and Zachara, J.M. (2014) Genomic analyses of bacterial porin-cytochrome
gene clusters. *Front Microbiol* **5**: 657.

Simon, J., Gross, R., Einsle, O., Kroneck, P.M., Kroger, A., and Klimmek, O. (2000) A NapC/NirT-type
cytochrome c (NrfH) is the mediator between the quinone pool and the cytochrome c nitrite reductase of
Wolinella succinogenes. *Mol Microbiol* **35**: 686-696.

Sousa, F.L., Alves, R.J., Ribeiro, M.A., Pereira-Leal, J.B., Teixeira, M., and Pereira, M.M. (2012) The
superfamily of heme–copper oxygen reductases: Types and evolutionary considerations. *Biochim Biophys*
Acta Bioenerg **1817**: 629-637.

Thamdrup, B., Steinsdóttir, H.G.R., Bertagnolli, A.D., Padilla, C.C., Patin, N.V., Garcia-Robledo, E. et
al. (2019) Anaerobic methane oxidation is an important sink for methane in the ocean's largest oxygen
minimum zone. *Limnol Oceanogr* **64**: 2569-2585.

Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R. et al. (2012) Differential gene and
transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**: 562-
578.

Trifinopoulos, J., Nguyen, L.T., von Haeseler, A., and Minh, B.Q. (2016) W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* **44**: W232-235.

Tsementzi, D., Wu, J., Deutsch, S., Nath, S., Rodriguez, R.L., Burns, A.S. et al. (2016) SAR11 bacteria
linked to ocean anoxia and nitrogen loss. *Nature* **536**: 179-183.

Tucker, N.P., Le Brun, N.E., Dixon, R., and Hutchings, M.I. (2010) There's NO stopping NsrR, a global
regulator of the bacterial NO stress response. *Trends Microbiol* **18**: 149-156.

Tully, B.J., Graham, E.D., and Heidelberg, J.F. (2018) The reconstruction of 2,631 draft metagenome-
assembled genomes from the global oceans. *Sci Data* **5**: 1-8.

Uzun, M., Alekseeva, L., Krutkina, M., Koziaeva, V., and Grouzdev, D. (2020) Unravelling the diversity
of magnetotactic bacteria through analysis of open genomic databases. *Sci Data* **7**: 252.

Ward, B.B., and Zafiriou, O.C. (1988) Nitrification and nitric oxide in the oxygen minimum of the eastern
tropical North Pacific. *Deep Sea Res Part A Oceanogr Res Pap* **35**: 1127-1142.

Wasser, I.M., De Vries, S., Moënne-Loccoz, P., Schröder, I., and Karlin, K.D. (2002) Nitric oxide in
biological denitrification: Fe/Cu metalloenzyme and metal complex NO_x redox chemistry. *Chem Rev* **102**:
1201-1234.

Woehle, C., Roy, A.S., Glock, N., Wein, T., Weissenbach, J., Rosenstiel, P. et al. (2018) A novel
eukaryotic denitrification pathway in foraminifera. *Curr Biol* **28**: 2536-2543 e2535.

Wu, Y.W., Simmons, B.A., and Singer, S.W. (2016) MaxBin 2.0: an automated binning algorithm to
recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**: 605-607.

Yang, S., Chang, B.X., Warner, M.J., Weber, T.S., Bourbonnais, A.M., Santoro, A.E. et al. (2020) Global
reconstruction reduces the uncertainty of oceanic nitrous oxide emissions and reveals a vigorous seasonal
cycle. *Proc Natl Acad Sci U S A* **117**: 11954-11960.

Yu, N.Y., Wagner, J.R., Laird, M.R., Melli, G., Rey, S., Lo, R. et al. (2010) PSORTb 3.0: improved
protein subcellular localization prediction with refined localization subcategories and predictive
capabilities for all prokaryotes. *Bioinformatics* **26**: 1608-1615.

Zallot, R., Oberg, N., and Gerlt, J.A. (2019) The EFI web resource for genomic enzymology tools:
leveraging protein, genome, and metagenome databases to discover novel enzymes and metabolic
pathways. *Biochemistry* **58**: 4169-4182.

Zedelius, J., Rabus, R., Grundmann, O., Werner, I., Brodkorb, D., Schreiber, F. et al. (2011) Alkane
degradation under anoxic conditions by a nitrate☐reducing bacterium with possible involvement of the
electron acceptor in substrate activation. *Environ Microbiol Rep* **3**: 125-135.

Zhang, I.H., Sun, X., Jayakumar, A., Fortin, S.G., Ward, B.B., and Babbin, A.R. (2023) Partitioning of
the denitrification pathway and other nitrite metabolisms within global oxygen deficient zones. *ISME*
Commun **3**: 76.

Zumft, W.G. (2005) Nitric oxide reductases of prokaryotes with emphasis on the respiratory, heme–
copper oxidase type. *J Inorg Biochem* **99**: 194-215.

542