# Distributed representations of innate behaviors in the hypothalamus do not predict specialized functional centers

Stefanos Stagkourakis[1, *], Giada Spigolon[2], Markus Marks[3], Michael Feyder[4], Joseph Kim[1], Pietro Perona[3], Marius Pachitariu[5, #], David J. Anderson[1, 6, #, *]

[1] Division of Biology and Biological Engineering 156-29, Tianqiao and Chrissy Chen Institute for Neuroscience, California Institute of Technology, Pasadena, California 91125, USA

[2] Biological Imaging Facility, California Institute of Technology, Pasadena, California 91125, USA

[3] Division of Engineering and Applied Science, California Institute of Technology, Pasadena, California 91125, USA

[4] Division of Psychiatry and Behavioral Sciences, Stanford University, Palo Alto, California 94305, USA

[5] Howard Hughes Medical Institute, Janelia Research Campus, Ashburn, VA, USA

[6] Howard Hughes Medical Institute, California Institute of Technology, 1200 East California Blvd, Pasadena, California 91125, USA

[#] Joint supervision

* Corresponding authors: Stefanos Stagkourakis, David J. Anderson

E-mail: stefanos.stagkourakis@gmail.com, wuwei@caltech.edu

## Summary

Survival behaviors are orchestrated by hardwired circuits located in deep subcortical brain regions, most prominently the hypothalamus. Artificial activation of spatially localized, genetically defined hypothalamic cell populations is known to trigger distinct behaviors, suggesting a nucleus-centered organization of behavioral control. However, no study has investigated the hypothalamic representation of innate behaviors using unbiased, large-scale single neuron recordings. Here, using custom silicon probes, we performed recordings across the rostro-caudal extent of the medial hypothalamus in freely moving animals engaged in a diverse array of social and predator defense ("fear") behaviors. Nucleus-averaged activity revealed spatially distributed generic "ignition signals" that occurred at the onset of each behavior, and did not identify sparse, nucleus-specific behavioral representations. Single-unit analysis revealed that social and fear behavior classes are encoded by activity in distinct sets of spatially distributed neuronal ensembles spanning the entire hypothalamic rostro-caudal axis. Individual ensemble membership, however, was drawn from neurons in 3-4 adjacent nuclei. Mixed selectivity was identified as the most prevalent mode of behavior representation by individual hypothalamic neurons. Encoding models indicated that a significant fraction of the variance in single neuron activity is explained by behavior. This work reveals that innate behaviors are encoded in the hypothalamus by activity in spatially distributed neural ensembles that each span multiple neighboring nuclei, complementing the prevailing view of hypothalamic behavioral control by single nucleus-restricted cell types derived from perturbational studies.

## Introduction

Brains evolved to create a series of abstractions that represent objects or other sensory cues in an organism's environment, assess their value, predict impending changes in their features, and transform those signals into adaptive behavioral responses. To deconstruct this complex process, two main experimental strategies have been taken to either: 1) record neural activity during behavior, or 2) manipulate neural activity and observe changes in behavior. The first strategy starts with measurements of neuronal activity and correlates those data with manipulations and measurements of sensory features or task variables (*1-5*). The behavioral readouts typically involve artificial tasks that require intensive training and are highly reproducible across trials. In mammals, this strategy has been employed to study the cortex, hippocampus, or other evolutionarily recent areas (*6-10*), and their influence on cognitive functions.

The second strategy starts with surgical, electrical, neurochemical, or genetically-based perturbation experiments to identify brain regions and cell types whose activation, inhibition, or destruction can promote or suppress particular behaviors. These manipulations are targeted to pre-selected areas based on hypotheses, assumptions, or prior knowledge. This strategy has dominated the study of internal states, motivation, reward, and emotion (*11-15*). The behavioral readouts typically comprise innate behaviors or simple forms of conditioning based on such instincts. The study of such functions has typically focused on the so-called "limbic system," including evolutionarily ancient regions such as the extended amygdala (*16, 17*), hypothalamus (*18, 19*), and midbrain (*20, 21*).

The two above-mentioned strategies can yield fundamentally different views of the mechanisms underlying brain functions of interest. Although many studies in practice employ both strategies sequentially, prevailing views in a field are often strongly shaped by the approach that was initially undertaken (*22-24*). The "recording first" strategy has led to an emphasis on high-dimensional population codes, neural dynamics, and manifolds in cortico-hippocampal state-space. Nevertheless, the field is increasingly employing perturbation experiments to test hypotheses based on observational studies. In

contrast, the "perturbation first" strategy has led to an emphasis in the hypothalamus field on hard-wired neural circuits that causally control specific innate behaviors in a manner that is highly deterministic and low-dimensional (*25-27*). However, large-scale recording experiments in this field have been relatively rare.

It remains unclear, therefore, whether these different views of how cortical and limbic systems control behavior reflect true biological differences, or rather differences in the evolution of these fields according to the initial experimental strategies used at their inception. To help resolve this discrepancy, here we have performed large-scale recordings from neural populations located along the rostro-caudal axis of the medial hypothalamus, using custom-designed silicon probes in freely moving mice engaged in solitary, social, or predator-defense behaviors. Our results reveal that the neural coding of such instinctive behaviors is, surprisingly, more distributed, dynamic, and high-dimensional than would have been expected from perturbation experiments. These results invite a re-evaluation of views of neural coding principles in cortical vs. limbic areas and highlight the importance of integrating results from observational and perturbational studies to arrive at a comprehensive understanding of brain function.

# Results

**Recording single neurons across multiple sites of the hypothalamus during behavior**

We sought to acquire synchronized large-scale electrophysiological and video recording data from freely moving mice engaging in diverse innate behaviors. To record the activity of hundreds of neurons simultaneously throughout the medial hypothalamus, we used a custom-fabricated silicon probe (NeuroNexus, Inc.) that allows chronic extracellular electrophysiological recordings (with single-unit resolution) from multiple nuclei (Fig. 1A). The eight-shank 256-channel silicon probe (Fig. 1B) was designed to provide simultaneous access to twenty hypothalamic subregions (nuclei or subnuclei), which are spatially distributed across the rostro-caudal extent of the hypothalamus (Fig. 1E). The probe covers ~2.3mm of brain tissue (Fig. 1A, lower), and we recorded ~400 units per animal (n=6 animals) for a total of 2,347 units.

To map shank trajectories, the back side of the probe shanks was coated with a red lipophilic dye that could be traced post-hoc. Intact iDISCO-cleared mouse brains were volumetrically imaged using light-sheet microscopy (*28*), and probe coordinates were mapped to the Allen Common Coordinate Framework (*29*) (Fig. 1, C and D). A unique electrode map was generated for each mouse by manually mapping electrodes to computationally sectioned sagittal sections of each mouse brain (Fig. S1 and S2). A schematic map of the areas recorded in each mouse is shown in Fig. 1E. Based on the electrode maps, neurons were allocated to specific hypothalamic brain areas (Fig. 1F, single-neuron activity colored according to the location of each neuron). Spike-sorting was performed using Kilosort (*30*), and basic analysis of neuron waveforms (Fig. 1G), and quality control of recordings was performed using pre-existing pipelines (see also Materials and Methods, section: *in vivo* electrophysiology). Neural data were analyzed in relation to behavior either a) with single neuron resolution (Fig. 1H – upper part), or b) by averaging the activity of cells per nucleus (Fig. 1H – lower part). This experimental design enabled us to test whether the encoding of behavior in the hypothalamus conforms to anatomical nuclear boundaries

and examine the extent to which neurons from the same nucleus are highly correlated or have distinct

activity patterns across diverse behaviors. (Fig. 1I).

To investigate the representation of behavior in the hypothalamus, we grouped behaviors into three

hierarchical levels (Fig 1I, right). At the highest level, we defined three broad behavioral categories which

we refer to as "classes": the solitary class includes actions such as self-grooming, and rearing. The social

class includes behaviors such as sniffing a male or female conspecific, or attacking a male conspecific.

The fear class includes behaviors such as sniffing a predator or exhibiting an escape response. At the

second level, we distinguished two "domains" within the "social" class: intermale and male-female social

behaviors (Fig. 1I, right; green and violet, respectively). Finally, at the third and lowest level, we annotated

43 distinct actions among the different classes and domains (Fig. 3B).

To ensure that the behaviors of interest would be expressed during the neural data recording

sessions, animals (male Swiss-Webster mice) were pre-screened for robust performance in the relevant

behavioral assays (Fig. 2A, a) solitary, b) social and c) fear). This pre-screening step reinforced the

expression of social behaviors (*11, 12*), and the display of fear behaviors during the recording sessions

(Fig. S3). Following the probe implantation, animals were allowed to recover and were habituated to the

experimental setup in the absence of the behavioral assays (Fig. 2B).

As previous work has shown that the neural firing rates often correlate with motion-related variables

in some brain areas (*31-33*), we assessed this question in our mesoscale hypothalamic recordings. Self-

motion variables and distance between animals did not correlate with mean single unit firing rates, either

during whole recording sessions (Fig. S4, A and B), or during specific behavior actions (Fig. S5, A-D).

Partial least squares (PLS) regression (Fig. S5, E-G) provided further evidence that single unit activity

also does not correlate with motion variables (Fig. S5Ed). However, the PLS regression revealed that

single-unit activity was correlated with consummatory and, to a lesser extent, appetitive social and fear

behaviors (Fig. S5, Fd and Gd).

**Spatially distributed and behavior non-specific neural "ignition signals"**

Next, we sought to address whether activity during specific appetitive and consummatory behaviors, including intermale social behaviors (Fig. 2, C and D), male-female social behaviors (Fig. 2, E and F), and predatory defensive behaviors (Fig. 2, G and H), is restricted anatomically or rather spatially distributed, based on averaging units recorded from each of the twenty nuclei. Throughout the entire hypothalamus we observed a synchronous burst of activity near the onset of all behaviors except for predator-induced freezing (Fig. 2, C-G). This spiking burst that occurred at all behavior onsets following the introduction of an intruder (example behavior raster plot shown in Fig. S6A) appeared consistently across behavioral episodes (Fig. S6, B and C). It had an onset of -200 msec prior to the initiation of behavior (Fig. S6E), average duration of 1.5 sec (Fig. S6F), and always ended before the termination of the behavior (Fig. S6G). This response, which we refer to here as an "ignition signal," encompassed ~25-35% of recorded units across the entire hypothalamus, except for freezing behavior which engaged only 7% of units (Fig. 2D-H, gray/black pie charts). The ignition signal was observed across multiple episodes of a given behavior directed towards the same intruder or predator (Fig. S4, C and D, and Fig. S6, A-C), indicating that it did not simply represent a novelty or neophobic response. However, the activity levels and proportion of the neurons recruited across behaviors markedly differed across nuclei (Fig. 2, C, E and G – right bar plots). Each nucleus was differentially activated during behaviors of interest (Fig. 2, D, F and H). Together, these findings indicate that ignition signals, while globally distributed, exhibit quantitative biases in their anatomic distribution that likely reflect behavior-specific responses in different hypothalamic nuclei (Fig. 2, D, F and H).

The near-uniform expression of ignition signals likely accounts for the lack of statistical significance in decoding behaviors from nucleus-averaged neural activity (SVM decoding models, Fig. S4E). An analysis of activity at single-unit resolution – discussed in the next section, however, could help to address whether behavior-specific modes of activity exist at the level of individual neurons.

**Single neuron activity primarily encodes distinct behavior classes**

To investigate the representation of behavior by single neurons in the hypothalamus, we annotated 43 behavioral actions among the different classes and domains (functional models discussed in Fig. 1I, right). Using these annotations, the first approach we used was to compute the single neuron activity-based behavior correlation matrix (Fig. 3, Aa-Ad). Using rastermap(*34*), we sorted the row and column indices of the behavior correlation matrix to maximize similarity between consecutive rows and columns, which allowed us to visually evaluate the differences in neural representations across behaviors (Fig. 3B). Following averaging of the behavior correlation matrix across six mice, we observed a prominent segregation between social and predatory-defensive behavioral classes (note the separation of red-["fear"] vs. green- and purple-tagged [social] behaviors in Fig. 3B). This effect was also prominent in the behavior correlation matrices of individual mice (Fig. S7, except for mouse #4). Overall, this analysis revealed that social and fear behavioral classes can be broadly segregated based on neuron population activity. Male- and female-directed social behaviors were also generally separated, but not as prominently. Related individual actions had correlated neural patterns to some extent within domains (Fig. 3B, sorting of behavior actions reflects similarity in their encoding).

The results of the behavior correlation matrix suggested that social and fear behaviors are encoded differently by hypothalamic neurons. However, this population-level analysis cannot distinguish whether individual behaviors are encoded by the same set of neurons that exhibit behavior class-specific activity patterns or whether distinct groups of neurons encode distinct behavior classes. To address this question, we next used a more targeted method, focusing on only two representative behavioral actions from each of the four behavioral domains of interest (total of eight behavioral actions, Fig. 3C). This approach could 1) confirm or disprove the segregation of social vs. fear behaviors, and 2) if the former is the case, indicate whether this segregation reflects different neuron groups active in the two behavioral classes, or whether it is driven by different patterns of activity exhibited by the same neurons.

In the solitary domain, the self-grooming (sg) and rearing (r) behavioral actions were included. Representative behavioral actions in the intermale domain were male intruder sniffing (ms), and attack

(ma), while in the male-female domain female intruder sniffing (fs) and mounting (fm) were included.

Lastly, in the predatory fear domain, the rat sniffing (rs) and escape response (re) behavior actions were

included. The timepoints of three episodes per behavior were extracted from the experimental design

illustrated in Fig. 3Aa. Specifically, solitary behavior episodes were extracted from experiment (a),

female-oriented behavior episodes were extracted from experiments (a) and (b), male-oriented behavior

episodes were extracted from experiments (c) and (d), and predator-defence behavior episodes were

extracted from experiment (e). Fig. S8A shows the activity of single units across the eight above-

mentioned behavioral actions, across three consecutive behavioral episodes. The activity of single units

was sorted from high to low within each brain area.

To test whether separate groups of recorded units exhibit distinct behavior-correlated activity profiles,

we performed $k$-means clustering of the single neuron activity (single neuron activity is shown in Fig.

S8A). The number of clusters, 41, was selected using the elbow method (*35, 36*); 41 clusters explained

70% of the single neuron activity data variance. The average activity of each of the 41 clusters during the

8 different domain-specific actions was plotted for each of the three consecutive episodes (Fig. 3C).

Agglomerative hierarchical linkage clustering (ALC) of these 41 clusters revealed five groups (i.e.,

dendrogram branches, Fig. 3C, right, threshold for separating branches of the dendrogram was defined

using the elbow method). This analysis suggested that the social and fear behavior classes are encoded

by functionally distinct groups of hypothalamic neurons (Fig. 3C, highlighted as blue [social] and red

[predatory fear] clusters in the dendrogram). This finding was also qualitatively supported by performing

non-negative matrix factorization (NMF) on the data shown in Fig. S8A. However, with NMF, the

distinction between social- vs. fear-correlated ensemble activity was less prominent (Fig. S9, A and B).

The weaker segregation of behavioral classes through NMF likely reflects that NMF allows neurons to

contribute to multiple factors (components).

While the $k$-means clusters associated with social or fear behavior classes were well separated,

neurons exhibited multi-action activity, also known as mixed selectivity, within each class-specific cluster.

To quantify this finding, we classified a cluster as active during a specific action if its activity (spiking rate)

was $\geq$ 2σ above its baseline activity. Accordingly, 56% of all clusters were identified to exhibit mixed

selectivity within their respective behavior classes. Only seven percent of hypothalamic clusters exhibited

mixed selectivity within a single behavioral domain (male- or female-directed behavior; Fig. 3E), while

27% of clusters showed prominent activity during only a single action (Fig. 3F). While the activity of any

single cluster exhibited adaptation across episodes (Fig. S8B) there was an overall positive correlation

between episodes (Fig. S8C) and a high-fidelity score (a measure of a cluster being identified as active

in the same behavior(s) across episodes, Fig. S8D). Thus, the representation of behavioral actions by

different clusters is reproducible across episodes, but not rigidly fixed.

**Encoding of behavior is spatially distributed, but individual ensembles are**

**anatomically clustered**

The foregoing analysis revealed that distinct neuron clusters encode social and fear behavior classes.

We next sought to determine whether individual activity-defined ensembles are anatomically localized

and whether they respect classical cytoarchitectonically defined nuclear boundaries.

We found that a) functionally similar clusters were distributed broadly across the hypothalamic rostro-

caudal axis (Fig 4A), and b) that most clusters were composed of cells that originated from a few

neighboring hypothalamic regions (Fig. 4B). Re-sorting these clusters within each branch of the

dendrogram according to their position along the rostro-caudal axis (Fig. 4C), revealed that each behavior

class was broadly represented along this axis (Fig. 4C).

The number of regions contributing to single clusters was typically four areas (Fig. 4d), suggesting

that most hypothalamic clusters consist of cells that originate from four neighbouring hypothalamic

regions. The same observation occurred when quantifying the composition of clusters that were identified

as "social", female-specific, and predatory-fear relevant (Fig. 4D – inset). We investigated whether

increasing the number of clusters in *k*-means leads to a reduced number of regions per cluster (Fig. 4E).

However, we found an asymptote at an average number of four nuclei per cluster which did not further

decrease even when the number of clusters was increased up to 120 (Fig. 4E). The smallest number of clusters that reached the asymptotic value of 4 regions per cluster was 41, and this model explained a majority of the variance in the data (70%). A qualitatively similar result emerged when analyzing factors identified by NMF, with closely related factors composed of neurons at different positions along the rostro-caudal axis, and heavily weighted neurons in each factor were present in neighboring hypothalamic regions (Fig. S9, D and E).

**Behavior explains a quarter of the variance of hypothalamic neuron activity**

The above approaches suggest that: 1) distinct populations of neurons represent social and fear behaviors, 2) the encoding of each behavioral class is represented by a set of clusters comprised of units defined by correlated neuronal activity, which are spatially distributed across the rostro-caudal axis, and 3) individual clusters typically contained neurons from ca. four neighboring nuclei, therefore exhibiting a strong local anatomical bias in their composition. However, since most of the units in these functional clusters exhibited mixed selectivity, it remained uncertain whether specific behavior actions could be decoded from neural activity.

To determine whether individual behavioral actions could be predicted from single-unit activity, we trained decoders based on neural activity using two different machine learning algorithms (gradient boosted trees (*37*), results plotted in Fig. S10, and support vector machines (*23*), results plotted in Fig. S11). A subset of appetitive and consummatory representative behavior actions per behavioral domain were selected to address the above point. The results indicated that most, but not all, actions within each of the four different behavioral domains could be decoded from neural data significantly better than chance. In the case of male-directed attack, decoder performance was high, reaching a shuffled subtracted median F1 score of ~0.5 (Fig. S10B).

The foregoing analysis suggested that behavioral actions could be decoded from neural activity but left open the question of which features of these behaviors were encoded.  To address this question, we

next examined how well the firing patterns of individual units could be predicted from a weighted sum of

different behavioral actions and features (features included animal's velocity, acceleration, distance, and

angle between animals). These features were computed based on pose estimation using DeepLabCut

(*38*).

For this purpose, we trained generalized linear models (GLMs) (*39, 40*) to predict the activity of each

neuron based on a weighted sum of selected predictor variables (design of GLMs shown in Fig. 5A). This

approach can: 1) identify the overall variance explained by specific predictors (Fig. 5B), 2) reveal the

influence of all annotated predictors on single neuron activity, by inspection of their coefficients/weights

(Fig. 5C), and 3) indicate whether neurons that belong to the same cluster (based on the similarity of their

behavior weights – Fig. 5C, see dendrogram colored branches) show any spatial bias (Fig. 5, D and E).

First, we sought to identify the percentage of variance explained by the four high-level behavior

classes. Only 8% of the variance of single neuron activity was accounted for by the four behavioral

domains (Fig. S12A). Next, we examined the percent of variance explained by forty behavioral actions,

annotated across the four behavioral domains of interest. Behavioral actions alone explained 23% of the

variance of hypothalamic neuron activity ($R^2$ median = 0.23, Fig. 5B).  Training GLM models with predictor

variables being the behavioral actions, plus motion signals (the animals' velocity and acceleration), and

proxies of sensory cues (distance and angle between animals), did not lead to an increase in the variance

explained (Fig. S12, C-E).

Many different behaviors had positive and negative weights on the activity pattern of each single-unit

(Fig. 5C), highlighting mixed selectivity as a prime mode of the hypothalamic encoding of behavior, and

providing further support to the results of the *k*-means (Fig. 3D) and NMF analysis (Fig. S9A). Application

of agglomerative clustering (ALC) to the matrix of GLM-derived behavior weights yielded 24 clusters of

neurons (the elbow method was used to select the number of clusters). Each cluster (Fig. 5C) contained

a unique combination of neurons whose activity was influenced by a specific set of behavior weights (Fig.

5C), with a near-uniform contribution of neurons from each of the six recorded mice (Fig. S13A). Sorting

these neurons based on their relative spatial location, however, did not generate clusters in a pattern

similar to that identified by ALC, suggesting that the neurons comprising the clusters are not restricted to

single hypothalamic nuclei (Fig. S13B). To investigate whether the cells comprising the ALC-generated

neuron clusters exhibit any restriction in their anatomic distribution, we measured the average

contribution of each nucleus to each cluster (Fig. 5D, rows). This revealed that clusters indeed exhibited

a spatial bias, with >80% of the cells comprising each cluster originating from ~four nearby regions (Fig.

5, D and E) – a finding consistent with the *k*-means-revealed cluster spatial bias (Fig. 4D). Also similar to

the *k*-means analysis, we found that GLM-based clusters exhibited spatial bias and mixed selectivity (Fig.

3D and 4C), which we quantified using a spatial localization index (single nucleus restricted vs. broadly

distributed) and using encoding entropy respectively (the degree of behavior specific vs. mixed selectivity

coding; see Materials and Methods section: GLM output – functional vs. spatial distribution of clusters).

## Discussion

The hypothalamus is traditionally viewed as a "low-dimensional" system in which innate social, defensive, and feeding behaviors are controlled by bulk neural activity flowing through specific pathways connecting functionally specialized anatomical subdivisions and cell types (*41-43*). In contrast, the cortico-hippocampal system is thought to operate using high-dimensional, dynamic neural codes involving complex spatio-temporal patterns of population activity (*44-47*). Here, we addressed whether this difference is truly biological or reflects the fact that most recording studies in the hypothalamus have used low-dimensional recording methods (e.g., fiber photometry) applied to anatomically restricted regions pre-selected based on perturbational studies. We report here what is, to our knowledge, the first unbiased high-dimensional single-unit recording of neural activity along the rostro-caudal axis of the medial hypothalamus during social and predator-defense behaviors. Our results suggest that behavioral coding in the hypothalamus is more high-dimensional, spatially distributed, dynamic, and variable than previously appreciated.

Several unexpected findings have emerged from this study. First, we observe that the initiation of these innate behaviors is (except for freezing) preceded by a hypothalamus-wide, synchronous burst of peri-onset activity, which we call an "ignition signal." Second, we find that functionally distinct subpopulations of neurons (defined by unbiased *k*-means clustering) encode social (mating and fighting) vs. predator defense behavioral classes. Specific behavioral actions or male- vs. female-directed social behaviors are less well-discriminated by such clusters. While these clusters are reproducibly activated across multiple episodes of each behavioral class, inter-episodic variability exists in the recruitment and activity level of the neurons comprising each cluster. Nevertheless, despite this variability, many behavioral actions could be decoded from single-unit activity with high accuracy. Third, we find that the subpopulations representing social and defense behaviors are broadly distributed across the rostro-caudal length of the hypothalamus rather than spatially localized to a single nucleus or region. Fourth, we find that cluster membership draws from neurons distributed in neighboring nuclei, typically four.

Nevertheless, behavior cannot be decoded from bulk neural activity based on anatomical location alone.

Finally, we find that a substantial fraction (median 23%) of the single-neuron variance in activity can be

explained by behavioral actions and, to a lesser degree, by behavioral domains. We consider the median

$R^2$ of 0.23 as a high value, as other studies which used brain wide recordings during stereotyped

behavioral tasks in head-fixed mice reported a cross-validated variance explained in single neuron

activity in the range of 5-15% (*48, 49*).

The present study paints an overall picture that challenges prevailing views of behavioral encoding

by the hypothalamus. We note that if these data had provided the very first, foundational view of

hypothalamic behavioral encoding, they would not have directed subsequent functional perturbations to

specific single nuclei such as VMH (*11, 13, 22, 23, 50-55*) or MPOA (*14, 56-60*). Instead, they would

have prompted the study of behavior encoding units across several neighboring nuclei. They would have

shifted the focus from seeking the identification of behavior-specific cell types to understanding the

properties conferred by the prime mode of activity we report here, i.e., mixed selectivity (*61-65*).

Is there any way to reconcile these two seemingly opposing views of behavioral control by the

hypothalamus? There are several possible explanations. One possibility is that prior, low-dimensional,

anatomically restricted approaches have missed what is suggested by our silicon probe recordings: i.e.,

that the "units" of hypothalamic computations are anatomically distributed, and that mixed-selectivity is

the prime mode of hypothalamic behavioral encoding. Recent studies suggest that even in the absence

of behavior action-tuned cells, behavior can still be decoded with high accuracy, with mixed selectivity

providing a basis for the encoding of numerous and diverse features in a computationally efficient manner

(*61-64*). For example, non-linear mixed selectivity enables the integration of diverse lines of information

in single neurons that can elicit behavior actions based on a threshold mechanism.

A second possibility is that our silicon probe recordings have missed anatomically restricted behavior-

specific subsets. This could be due to sparse sampling (we recorded on average only ~20 units per

nucleus per animal) and/or to variability in the activity of individual units across episodes and animals

during a given behavior, perhaps reflecting representational drift (*66, 67*) or stochastic recruitment of

individual units (*68*). This variability may explain, in part, why a recent multi-site coarse recording of bulk

calcium activity from Esr1$^+$ neurons across the hypothalamus yielded reproducible, spatially distinct

patterns of activity during male- vs. female-directed social behaviors, in specific nuclei previously

identified as causally relevant to these behaviors by perturbation studies (*69*). Alternatively, Esr1$^+$

neurons may represent a "privileged," more behavior-specific subpopulation than the "anonymous" units

sampled here. A third possibility (not mutually exclusive with the second) is that the intrinsic variability in

the performance or sequence of specific behavioral actions during a given activity (e.g., a male-male

social interaction) may confound efforts to correlate a high-granularity annotation of these actions (43)

with high-dimensional neural data (~2400 units from 6 animals). Finally, we note that causation and

representation are not the same thing: not all neural representations during a given behavior necessarily

reflect a causal role for those neurons; for example, some activated units may represent efference copies

of motor signals, or GABAergic neurons that serve to inhibit competing behaviors.

Further work is required to distinguish these possibilities and to arrive at an understanding of

behavioral coding and function that reconciles and integrates the two views of hypothalamic computations

provided by recording and perturbation strategies, respectively. At the very least, our data indicate that

the application to the hypothalamus of unbiased, large-scale, anonymous single-unit recordings reveals

a surprisingly complex pattern of spatio-temporal activity that does not easily map to specific nuclei or

single behavior actions. It suggests that the prevailing view of the hypothalamus as a low-dimensional

control system needs to be revised, or alternatively that the signals that control behavior are, to a large

extent, "hidden" in the population code.

# References

1. X. D. Lin *et al.*, High-throughput mapping of brain-wide activity in awake and drug-responsive vertebrates. *Lab on a chip* **15**, 680-689 (2015).
2. E. Litvina *et al.*, BRAIN Initiative: Cutting-Edge Tools and Resources for the Community. *Journal of Neuroscience* **39**, 8275-8284 (2019).
3. M. G. Perich, K. Rajan, Rethinking brain-wide interactions through multi-region 'network of networks' models. *Current opinion in neurobiology* **65**, 146-151 (2020).
4. B. J. Edelman, E. Macé, Functional ultrasound brain imaging: Bridging networks, neurons, and behavior. *Current Opinion in Biomedical Engineering* **18**, (2021).
5. T. Schrodel, R. Prevedel, K. Aumayr, M. Zimmer, A. Vaziri, Brain-wide 3D imaging of neuronal activity in Caenorhabditis elegans with sculpted light. *Nature methods* **10**, 1013-1020 (2013).
6. S. Brodt *et al.*, Fast track to the neocortex: A memory engram in the posterior parietal cortex. *Science* **362**, 1045-+ (2018).
7. F. Chersi, N. Burgess, The Cognitive Architecture of Spatial Navigation: Hippocampal and Striatal Contributions. *Neuron* **88**, 64-77 (2015).
8. C. F. Doeller, C. Barry, N. Burgess, Evidence for grid cells in a human memory network. *Nature* **463**, 657-U687 (2010).
9. S. F. Grieco, T. C. Holmes, X. M. Xu, Probing neural circuit mechanisms in Alzheimer's disease using novel technologies. *Mol Psychiatr*, (2023).
10. D. F. Tomé, S. Sadeh, C. Clopath, Coordinated hippocampal-thalamic-cortical communication crucial for engram dynamics underneath systems consolidation. *Nature communications* **13**, (2022).
11. S. Stagkourakis, G. Spigolon, G. Liu, D. J. Anderson, Experience-dependent plasticity in an innate social behavior is mediated by hypothalamic LTP. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 25789-25799 (2020).
12. S. Stagkourakis *et al.*, A neural network for intermale aggression to establish social hierarchy. *Nature neuroscience* **21**, 834-+ (2018).
13. D. Lin *et al.*, Functional identification of an aggression locus in the mouse hypothalamus. *Nature* **470**, 221-226 (2011).
14. T. Karigo *et al.*, Distinct hypothalamic control of same- and opposite-sex mounting behaviour in mice (vol 589, pg 258, 2021). *Nature* **589**, E9-E9 (2021).
15. P. S. Kunwar *et al.*, Ventromedial hypothalamic neurons control a defensive emotion state. *eLife* **4**, (2015).
16. A. Adhikari *et al.*, Basomedial amygdala mediates top-down control of anxiety and fear. *Nature* **527**, 179-+ (2015).
17. J. Kim, X. Y. Zhang, S. Muralidhar, S. A. LeBlanc, S. Tonegawa, Basolateral to Central Amygdala Neural Circuits for Appetitive Behaviors. *Neuron* **93**, 1464-+ (2017).
18. N. Scott, M. Prigge, O. Yizhar, T. Kimchi, A sexually dimorphic hypothalamic circuit controls maternal care and oxytocin secretion. *Nature* **525**, 519-+ (2015).
19. S. Stagkourakis, G. Spigolon, G. Liu, D. J. Anderson, Experience-dependent plasticity in an innate social behavior is mediated by hypothalamic LTP. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 25789-25799 (2020).
20. D. Mahadevia *et al.*, Dopamine promotes aggression in mice via ventral tegmental area to lateral septum projections. *Nature communications* **12**, (2021).
21. A. L. Falkner *et al.*, Hierarchical Representations of Aggression in a Hypothalamic-Midbrain Circuit. *Neuron* **106**, 637-648 e636 (2020).
22. H. Lee *et al.*, Scalable control of mounting and attack by Esr1+ neurons in the ventromedial hypothalamus. *Nature* **509**, 627-632 (2014).

23.  R. Remedios *et al.*, Social behaviour shapes hypothalamic neural ensemble representations of conspecific
sex. *Nature* **550**, 388-+ (2017).
24.  A. Nair *et al.*, An approximate line attractor in the hypothalamus encodes an aggressive state. *Cell* **186**,
178-193 e115 (2023).
25.  A. Asaba *et al.*, Prelimbic cortex responds to male ultrasonic vocalizations in the presence of a male
pheromone in female mice. *Frontiers in neural circuits* **16**, (2022).
26.  K. K. Ishii, K. Touhara, Neural circuits regulating sexual behaviors via the olfactory system in mice.
*Neuroscience research* **140**, 59-76 (2019).
27.  T. Itakura *et al.*, A single vomeronasal receptor promotes intermale aggression through dedicated
hypothalamic neurons. *Neuron* **110**, 2455-2469 e2458 (2022).
28.  N. Renier *et al.*, iDISCO: A Simple, RapidMethod to Immunolabel Large Tissue Samples for Volume Imaging.
*Cell* **159**, 896-910 (2014).
29.  Q. X. Wang *et al.*, The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. *Cell* **181**,
936-+ (2020).
30.  N. A. Steinmetz *et al.*, Neuropixels 2.0: A miniaturized high-density probe for stable, long-term brain
recordings. *Science* **372**, 258-+ (2021).
31.  M. Vinck, R. Batista-Brito, U. Knoblich, J. A. Cardin, Arousal and Locomotion Make Distinct Contributions
to Cortical Activity Patterns and Visual Encoding. *Neuron* **86**, 740-754 (2015).
32.  M. C. Dadarlat, M. P. Stryker, Locomotion Enhances Neural Encoding of Visual Stimuli in Mouse V1. *Journal
of Neuroscience* **37**, 3764-3775 (2017).
33.  A. J. Christensen, J. W. Pillow, Reduced neural activity but improved coding in rodent higher-order visual
cortex during locomotion. *Nature communications* **13**, 1676 (2022).
34.  Z. L. Stringer C., Syeda A., Du F., Kesa M., & Pachitariu M., Rastermap: a discovery method for neural
population recordings. *bioRxiv*, (2023).
35.  C. Goutte, P. Toft, E. Rostrup, F. Å. Nielsen, L. K. Hansen, On clustering fMRI time series. *NeuroImage* **9**,
298-310 (1999).
36.  R. L. Thorndike, Who Belongs in the Family? *Psychometrika* **18**, 267-276 (1953).
37.  G. Viejo, T. Cortier, A. Peyrache, Brain-state invariant thalamo-cortical coordination revealed by non-linear
encoders. *PLoS computational biology* **14**, (2018).
38.  A. Mathis *et al.*, DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.
*Nature neuroscience* **21**, 1281-1289 (2018).
39.  A. I. Weber, J. W. Pillow, Capturing the Dynamical Repertoire of Single Neurons with Generalized Linear
Models. *Neural computation* **29**, 3260-3289 (2017).
40.  I. M. Park, M. L. R. Meister, A. C. Huk, J. W. Pillow, Encoding and decoding in parietal cortex during
sensorimotor decision-making. *Nature neuroscience* **17**, 1395-1403 (2014).
41.  L. Mei, T. Osakada, D. Lin, Hypothalamic control of innate social behaviors. *Science* **382**, 399-404 (2023).
42.  J. D. Hahn, G. Fink, M. R. Kruk, B. G. Stanley, Editorial: Current Views of Hypothalamic Contributions to the
Control of Motivated Behaviors. *Frontiers in systems neuroscience* **13**, (2019).
43.  C. Dulac, L. A. O'Connell, Z. Wu, Neural control of maternal and paternal behaviors. *Science* **345**, 765-770
(2014).
44.  J. Basu, S. A. Siegelbaum, The Corticohippocampal Circuit, Synaptic Plasticity, and Memory. *Cold Spring
Harb Perspect Biol* **7**, (2015).
45.  M. C. Aoi, V. Mante, J. W. Pillow, Prefrontal cortex exhibits multidimensional dynamic encoding during
decision-making. *Nature neuroscience* **23**, 1410-1420 (2020).
46.  A. Sarel *et al.*, Natural switches in behaviour rapidly modulate hippocampal coding. *Nature* **609**, 119-127
(2022).
47.  J. Kim, A. Joshi, L. Frank, K. Ganguly, Cortical-hippocampal coupling during manifold exploration in motor
cortex. *Nature* **613**, 103-110 (2023).

48.   N. A. Steinmetz, P. Zatka-Haas, M. Carandini, K. D. Harris, Distributed coding of choice, action and
engagement across the mouse brain. *Nature* **576**, 266-+ (2019).
49.   W. E. Allen *et al.*, Thirst regulates motivated behavior through modulation of brainwide neural population
dynamics. *Science* **364**, 253-+ (2019).
50.   A. Kennedy *et al.*, Stimulus-specific hypothalamic encoding of a persistent defensive state. *Nature* **586**,
730-734 (2020).
51.   T. Yang *et al.*, Social Control of Hypothalamus-Mediated Male Aggression. *Neuron* **95**, 955-+ (2017).
52.   K. K. Ishii *et al.*, A Labeled-Line Neural Circuit for Pheromone-Mediated Sexual Behaviors in Mice. *Neuron*
**95**, 123-+ (2017).
53.   A. L. Falkner, L. Grosenick, T. J. Davidson, K. Deisseroth, D. Y. Lin, Hypothalamic control of male aggression-
seeking behavior. *Nature neuroscience* **19**, 596-+ (2016).
54.   H. Lee *et al.*, Scalable control of mounting and attack by Esr1+ neurons in the ventromedial
hypothalamus. *Nature* **509**, 627-+ (2014).
55.   C. F. Yang *et al.*, Sexually Dimorphic Neurons in the Ventromedial Hypothalamus Govern Mating in Both
Sexes and Aggression in Males. *Cell* **153**, 896-909 (2013).
56.   D. Liu *et al.*, A Hypothalamic Circuit Underlying the Dynamic Control of Social Homeostasis. *bioRxiv*,
(2023).
57.   S. Stagkourakis *et al.*, A Neuro-hormonal Circuit for Paternal Behavior Controlled by a Hypothalamic
Network Oscillation. *Cell* **182**, 960-+ (2020).
58.   L. Mei, R. Z. Yan, L. P. Yin, R. M. Sullivan, D. Y. Lin, Antagonistic circuits mediating infanticide and maternal
care in female mice. *Nature*, (2023).
59.   S. X. Zhang *et al.*, Hypothalamic dopamine neurons motivate mating through persistent cAMP signalling.
*Nature* **597**, 245-+ (2021).
60.   J. Kohl, C. Dulac, Neural control of parental behaviors. *Current opinion in neurobiology* **49**, 116-122 (2018).
61.   S. Kira, H. Safaai, A. S. Morcos, S. Panzeri, C. D. Harvey, A distributed and efficient population code of
mixed selectivity neurons for flexible navigation decisions. *Nature communications* **14**, 2121 (2023).
62.   M. T. Kaufman *et al.*, The implications of categorical and category-free mixed selectivity on
representational geometries. *Current opinion in neurobiology* **77**, 102644 (2022).
63.   W. J. Johnston, S. E. Palmer, D. J. Freedman, Nonlinear mixed selectivity supports reliable neural
computation. *PLoS computational biology* **16**, e1007544 (2020).
64.   I. S. Grunfeld, E. Likhtik, Mixed selectivity encoding and action selection in the prefrontal cortex during
threat assessment. *Current opinion in neurobiology* **49**, 108-115 (2018).
65.   M. Rigotti *et al.*, The importance of mixed selectivity in complex cognitive tasks. *Nature* **497**, 585-590
(2013).
66.   C. E. Schoonover, S. N. Ohashi, R. Axel, A. J. P. Fink, Representational drift in primary olfactory cortex.
*Nature* **594**, 541-546 (2021).
67.   P. Zhao *et al.*, Accelerated social representational drift in the nucleus accumbens in a model of autism.
*bioRxiv*, (2023).
68.   C. Haimerl, C. Savin, E. P. Simoncelli, Flexible information routing in neural populations through stochastic
comodulation. *Advances in Neural Information Processing Systems 32 (Nips 2019)* **32**, (2019).
69.   Z. Guo *et al.*, Neural dynamics in the limbic system during male social behaviors. *Neuron*, (2023).
70.   C. Segalin *et al.*, The Mouse Action Recognition System (MARS) software pipeline for automated analysis
of social behaviors in mice. *eLife* **10**, (2021).
71.   J. J. Jun *et al.*, Fully integrated silicon probes for high-density recording of neural activity. *Nature* **551**, 232-
+ (2017).
72.   S. M. Lundberg, S. I. Lee, A Unified Approach to Interpreting Model Predictions. *Adv Neur In* **30**, (2017).
73.   T. Q. Chen, C. Guestrin, XGBoost: A Scalable Tree Boosting System. *Kdd'16: Proceedings of the 22nd Acm*
*Sigkdd International Conference on Knowledge Discovery and Data Mining*, 785-794 (2016).

**Acknowledgments**

Members of the Anderson laboratory are thanked for their advice and discussion during the preparation of this manuscript. Light-sheet imaging was performed in the Biological Imaging Facility, with the support of the Caltech Beckman Institute and the Arnold and Mabel Beckman Foundation. Special thanks to the Director of the Biological Imaging Facility at Caltech, Dr. Andres Collazo, for providing crucial experimental support and advice. We also thank Jung-Sook Chang and Xiaolin Da for experimental support, C. Chiu for laboratory management, G. Mancuso, L. Chavarria and D. R. Navarrete for administrative assistance. S.S. received support from the Wenner-Gren Foundations and by NIH K99MH131754. D.J.A. is an investigator of the Howard Hughes Medical Institute. This work was supported in part by NIH grants NS123916, MH1223612, and MH070053 to D.J.A. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. All authors support an inclusive, diverse, and equitable conduct of research.

**Author contributions**

S.S. conceived the study, designed and performed all experiments, wrote hardware-operating and analysis code, analyzed data, and generated the first draft of the manuscript. G.S. contributed to brain clearing, light-sheet microscopy, and silicon probe registration to the Allen Common Coordinate Framework. M.M. performed the decoding analysis using gradient boosted trees. P.P. designed decoding approaches. M.F. wrote and provided code for image registration for light-sheet microscopy data. J.K. contributed to the analysis. M.P. supervised S.S.. D.J.A. supervised S.S., contributed to the study design and analysis, and wrote the manuscript. All authors reviewed the manuscript.

**Competing interests**

The authors declare no competing financial interests.

# Main Figures

**Fig. 1. Approach for single neuron recordings across the hypothalamus using custom extracellular electrophysiology silicon probes in freely moving mice.** (**A**) Top - outbred Swiss Webster male mice were chronically implanted with a custom silicon probe. Bottom - design, and specifics of the eight-shank 256-channel custom silicon probe manufactured by NeuroNexus. (**B**) Stereoscope-acquired image of the shank-tips of the probe containing the microelectrodes. (**C**) Post-hoc identification of probe shanks and electrode placement through volumetric light-sheet imaging. (**D**) Registration of the

probes to the Allen Common Coordinate Framework across all recorded mice. (**E**) Target coordinates include twenty, individually colored subregions of the hypothalamus. (**F**) Following electrode mapping and spike sorting, single neurons were assigned to their respective locations / hypothalamic areas. (**G**) Example waveforms of two recorded neurons. (**H**) Example spike data across thirteen neurons of VMHvl (top), and illustration of their average activity (bottom). (**I**) Spatial and functional plausible models, combination of which could explain the hypothalamic encoding of behavior.

Abbreviations: DAQ: data acquisition system, MPOAv: medial preoptic area – ventral subdivision. MPOAc: medial preoptic area – central subdivision. MPOAd: medial preoptic area – dorsal subdivision. MPNv: medial preoptic nucleus – ventral subdivision. MPNc: medial preoptic nucleus – central subdivision. RCHr: retrochiasmatic area – rostral subdivision. RCHc: retrochiasmatic area – caudal subdivision. AHNr: anterior hypothalamic nucleus – rostral subdivision. AHNce: anterior hypothalamic nucleus – central subdivision. AHNca: anterior hypothalamic nucleus – caudal subdivision. TUr: tuberal nucleus – rostral subdivision. TUc: tuberal nucleus – caudal subdivision. VMHvlR: ventromedial hypothalamic nucleus ventrolateral segment – rostral subdivision. VMHvlC: ventromedial hypothalamic nucleus ventrolateral segment – caudal subdivision. VMHcR: ventromedial hypothalamic nucleus central segment – rostral subdivision. VMHcC: ventromedial hypothalamic nucleus central segment – caudal subdivision. DMH: dorsomedial hypothalamic nucleus. PMvR: ventral premammillary nucleus – rostral subdivision. PMvC: ventral premammillary nucleus – caudal subdivision. PMd: dorsal premammillary nucleus. See also Fig. S1 and S2.

**Fig. 2. Neural ignition signals dominate the peri-onset interval of all instinctual behaviors, apart from freezing to the presence of a predator.** (**A**) Behavioral screening of outbred Swiss Webster mice was used to identify individual mice with prominent behavioral expressions in all of the three behavior paradigms of interest used to assess mating with a female conspecific, aggression against a male conspecific and fear behaviors in the presence of a predator (Long-Evans [LE] rat). Only mice that exhibited appetitive and consummatory behaviors across the three behavior domains (i.e., mating, aggression, and fear), across all behavior screening sessions were selected for chronic silicon probe implant surgery. (**B**) Schematic illustrating the timeline following implantation of the silicon probe. Animals were allowed to fully recover for two weeks in their home cage and were given access to a running wheel. Four habituation sessions were performed (one daily), 4 days prior to the start of the neural recordings

and behavior experiments. Habitation sessions included exposure to the exact experimental conditions,

in the absence of the stimulus (conspecific, or predator). (**C**) Average behavior signal and mean neural

activity at the peri-onset interval of an appetitive and a consummatory intermale social behavior (top).

Average neural activity per nucleus (heatmap), participation index (% of neurons activated above 2 SD

from baseline activity), and average firing rate of the activated neurons (bottom, n = 6 mice, neural data

averaged across three consecutive behavior instances per mouse). (**D**) Small black and gray pie plot:

Black – total % of neurons activated, Gray: total % of neurons that did not raise their activity above 2σ.

Larger colored pie plot illustrates the spatial coordinates of the neurons that became active (raised their

activity >2σ), during male anogenital sniffing (left) and intermale attack (right, n = 6 mice, neural data

averaged across three consecutive behavior instances). (**E** and **F**) Same as panels (C) and (D) for an

appetitive and a consummatory male-female social behavior. (**G** and **H**) Same as panels (C) and (D) for

two predatory fear behaviors.

In box-and-whisker plots, center lines indicate medians, box edges represent the interquartile range, and

whiskers extend to the minimal and maximal values. See also Fig. S3, S4, and S5. Abbreviations: Partic.

Idx.: participation index, sp/s: Firing rate (spikes / sec). Abbreviations of brain areas are described on the

legend of Fig. 1.

**Fig. 3. Encoding of social and fear behavior classes by functionally non-overlapping hypothalamic neurons.** (**A**) Analysis design to generate a behaviors correlation matrix. (**Aa**) Experimental conditions. (**Ab**) Single-unit neural activity during the behavioral paradigms shown in Aa. (**Ac**) Z-scored activity of all recorded neurons across solitary-, social-, and fear-behaviors (data from an example mouse). (**Ad**) Multiplication of the matrix shown in ac with its transposed version, results to the behavior covariance matrix, which is then transformed to a behavior correlation matrix. (**B**) Correlation matrix of behavior (average across n = 6 mice). Correlation matrices of behavior of the individual mice can be found in Fig. S7. (**C**) Mean activity of *k*-means clusters explaining 70% of the variance of the single unit data (Extended Data Figure 8). Clusters are sorted using agglomerative linkage clustering to generate a dendrogram whose branches have been assigned semantic labels, based on cluster activity being higher than two standard deviations in specific behavioral classes. (**D**) Left: Mean activity of a *k*-means cluster selected from the forty-one clusters identified and shown in (Fig. 3c), exhibiting mixed selectivity, i.e., activity in diverse behavior domains – in this case, in male-male and male-female behaviors. Right: Portion of clusters showing mixed-selectivity / multi-domain activity. (**E**) Left: Mean activity of a *k*-means cluster selected from the forty-one clusters identified (Fig. 3C), showing mixed selectivity within a behavioral domain – in this case, in intermale behaviors. Right: Portion of clusters showing mixed selectivity within a behavioral domain (alone / intermale / male-female / and fear behaviors). (**F**) Left: Mean activity of a *k*-means cluster selected from the forty-one clusters identified (Fig. 3c), showing behavior-specific tuning – in this case, in escape response to a predator. Right: Portion of clusters showing behavior action-specific tuning.

Abbreviations: sg: self-grooming, r: rearing, ms: male intruder anogenital sniffing, ma: intermale attack, fs: female intruder anogenital sniffing, fm: mounting female intruder, rs: sniffing predator, re: escape response to a predator. See also Fig. S7, S8, and S9.

**Fig. 4. Encoding of behavior classes is spatially distributed, but single clusters are spatially localized.** (**A**) Functional clusters of single neuron activity (panel from Fig. 3c – left). (**B**) Spatial composition of each of the 41 functional clusters shown in panel (A). (**C**) Re-sorting the spatial composition of each cluster, to present clusters composed of most rostral regions, to clusters composed of most caudal regions. (**D**) Histogram of hypothalamic area contribution per cluster. Inset: Histogram of hypothalamic area contribution to clusters that are active in specific behaviors. (**E**) Spatial composition of clusters based on the choice of *k* (number of clusters selected for *k*-means clustering), reveals that the average composition of a cluster by four regions is a stable solution.

Abbreviations: sg: self-grooming, r: rearing, ms: male intruder anogenital sniffing, ma: intermale attack, fs: female intruder anogenital sniffing, fm: mounting female intruder, rs: sniffing predator, re: escape response to a predator. See also Fig. S9, B-D.

**Fig. 5. Solitary, social and fear behaviors explain a quarter of hypothalamic neuron activity variance and suggest a spatial organization of functional neuron clusters.** (**A**) Analysis design using generalized linear models (GLMs) to study the encoding of behavior by non-genetically defined mesoscale hypothalamic populations. (**B**) Variance explained (cross-validated, shuffled data $R^2$ subtracted, $R^2$) of single-neuron activity (n=6 mice, 2347 neurons) based on forty-two instinctual behaviors. (**C**) Agglomerative linkage clustering of behavior GLM-generated coefficients into twenty-four neural dendrogram branches/clusters. (**D**) Average neuron counts per brain area across the hypothalamic rostro-caudal axis across all neural branches, reveals a spatial bias of neural branch composition. (**E**) Histogram of hypothalamic area contribution per cluster. (**F**) Spatial vs. functional characteristics of neuron clusters.

In box-and-whisker plots, center lines indicate medians, box edges represent the interquartile range, and whiskers extend to the minimal and maximal values. See also Fig. S11, S12, and S13.

# Extended Data Figures

see also *Fig. S2* for the mapping of probe recording sites located in shanks 1-4

**Fig. S1. Mapping of recording sites across the medial-caudal hypothalamus.** (**A**) Left: Computationally sectioned coronal view of the mouse brain that includes the eighth shank of the chronic silicon probe implant. Center and Right: Mapping the shank and the electrode-containing area of the

shank to the Allen Common Coordinate Framework (example section/mouse). (**B**) Mapping the overlap

of all recording electrodes of the eighth shank across all probes/mice to the caudal hypothalamic areas

(PMd: dorsal premammillary nucleus, and PMv: ventral premammillary nucleus, n=6 mice). (**C**) Left:

Computationally sectioned coronal view of the mouse brain that includes the seventh shank of the chronic

silicon probe implant. Center and Right: Mapping the shank and the electrode-containing area of the

shank to the Allen Common Coordinate Framework (example section/mouse). (**D**) Mapping the overlap

of all recording electrodes of the seventh shank across all probes/mice to the caudal hypothalamic areas

(PMd: dorsal premammillary nucleus, and PMv: ventral premammillary nucleus, n=6 mice). (**E**) Left:

Computationally sectioned coronal view of the mouse brain that includes the sixth shank of the chronic

silicon probe implant. Center and Right: Mapping the shank and the electrode-containing area of the

shank to the Allen Common Coordinate Framework (example section/mouse). (**F**) Mapping the overlap

of all recording electrodes of the sixth shank across all probes/mice to the medial-caudal hypothalamic

areas (DMH: dorsomedial hypothalamic nucleus, VMHc: ventromedial hypothalamic nucleus central

segment, VMHvl: ventromedial hypothalamic nucleus ventrolateral segment, and tuberal nucleus, n=6

mice). (**G**) Left: Computationally sectioned coronal view of the mouse brain that includes the fifth shank

of the chronic silicon probe implant. Center and Right: Mapping the shank and the electrode-containing

area of the shank to the Allen Common Coordinate Framework (example section/mouse). (**H**) Mapping

the overlap of all recording electrodes of the fifth shank across all probes/mice to the medial-caudal

hypothalamic areas (DMH: dorsomedial hypothalamic nucleus, VMHc: ventromedial hypothalamic

nucleus central segment, VMHvl: ventromedial hypothalamic nucleus ventrolateral segment, and tuberal

nucleus, n=6 mice).

Figure relevant to Fig. 1.

**Fig. S2. Mapping of recording sites across the medial-rostral hypothalamus.** (**A**) Left:
Computationally sectioned coronal view of the mouse brain that includes the fourth shank of the chronic

silicon probe implant. Center and Right: Mapping the shank and the electrode-containing area of the

shank to the Allen Common Coordinate Framework (example section/mouse). (**B**) Mapping the overlap

of all recording electrodes of the fourth shank across all probes/mice to the caudal hypothalamic areas

(RCH: retrochiasmatic area, and AHN: anterior hypothalamic nucleus, n=6 mice). (**C**) Left:

Computationally sectioned coronal view of the mouse brain that includes the third shank of the chronic

silicon probe implant. Center and Right: Mapping the shank and the electrode-containing area of the

shank to the Allen Common Coordinate Framework (example section/mouse). (**D**) Mapping the overlap

of all recording electrodes of the third shank across all probes/mice to the medial-rostral hypothalamic

areas (RCH: retrochiasmatic area and AHN: anterior hypothalamic nucleus, n=6 mice). (**E**) Left:

Computationally sectioned coronal view of the mouse brain that includes the second shank of the chronic

silicon probe implant. Center and Right: Mapping the shank and the electrode-containing area of the

shank to the Allen Common Coordinate Framework (example section/mouse). (**F**) Mapping the overlap

of all recording electrodes of the second shank across all probes/mice to the medial-rostral hypothalamic

areas (MPN: medial preoptic nucleus, n=6 mice). (**G**) Left: Computationally sectioned coronal view of the

mouse brain that includes the first shank of the chronic silicon probe implant. Center and Right: Mapping

the shank and the electrode-containing area of the shank to the Allen Common Coordinate Framework

(example section/mouse). (**H**) Mapping the overlap of all recording electrodes of the first shank across

all probes/mice to the rostral hypothalamic areas (MPOA: medial preoptic area, n=6 mice).

Figure relevant to Fig. 1.

**Fig. S3. Quantification of behavior in the predatory fear assay.** (**A**) Experimental design of a custom

predatory fear assay, with four compartments. R: compartment containing the predator (rat), with a single,

clear, and perforated side facing the compartment IZ, i.e., interaction compartment. TZ: is the transition

compartment that links compartments R and SZ, the latter furthest away from the predator. (**B**) Timeline

of the behavioral experiments with the fear condition flanked by two periods where the implanted mouse

is alone in the arena. (**C**) Snapshots illustrating four manually annotated behaviors, specifically;

approaching the predator, interacting with the predator, escape response to a predator (a high-velocity

activity bout towards the safe zone compartment), and freezing behavior (observed in the freezing

compartment). (**D**) Frequency distribution of velocity during the three conditions shown in panel (B). (**E**)

Freezing bout duration in the three conditions presented in in panel (B). (**F**) Number of crosses of the

interaction zone across the three conditions presented in panel (B). (**G**) Distance to interaction zone

among the three conditions presented in panel (B).

$*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$. Statistical comparisons of the data illustrated in panel

(D) were performed by the Kolmogorov-Smirnov test, while comparisons of the data shown in panels (E-

G) were performed through One-Way ANOVA and Dunnett's post-hoc test was used to correct for

multiple comparisons. In box-and-whisker plots, center lines indicate medians, box edges represent the

interquartile range, and whiskers extend to the minimal and maximal values. Figure relevant to Fig. 2.

**Fig. S4. Motion variables including distance between animals and animal's velocity, do not**
**correlate with the mean firing rate of hypothalamic neurons, ignition signals occur across**
**behavioral action episodes, and behavior actions are not decodable based on nucleus-averaged**
**neural activity.** (**A**) Example time series data of behavior, single neuron activity, mean firing rate,
distance between animals, and velocity of recorded mouse. (**B**) Average behavior signal and mean neural
activity at the peri-onset interval of predator-directed anogenital sniffing, in three behavioral episodes
(top). Average neural activity per nucleus (heatmap), participation index (% of neurons activated above
2 SD from baseline activity), and average firing rate of the activated neurons (bottom, n = 6 mice, in three
behavior episodes per mouse). (**C**) Average behavior signal and mean neural activity at the peri-onset
interval of escape responses in the presence of a predator, in three behavioral episodes (top). Average
neural activity per nucleus (heatmap), participation index (% of neurons activated above 2 SD from
baseline activity), and average firing rate of the activated neurons (bottom, n = 6 mice, in three behavior
episodes per mouse). (**D**) Correlation analysis of motion variables and average firing rate (n=6 mice, data
pooled across mice). (**E**) SVM decoder F1 score on behavior actions of interest based on nucleus-
averaged neural activity (n = 6 mice).

Abbreviations: mFR: mean firing rate all recorded neurons, dist.: distance between animals, vel.: velocity,
sp/s: spikes per second, nrm.: data normalized between 0 and 1 values, 0.2 s gsf: 200 millisecond
gaussian smoothing filter applied on the spike and behavior data prior to performing linear regression.
Figure relevant to Fig. 2.

**Fig. S5. Consummatory instinctual behaviors have stronger representation in hypothalamic neural data, as opposed to appetitive behaviors and motion-related variables.** (**A**) Landscape of neural activity vs. velocity of implanted mouse and distance between animals (pooled data across six

mice). (**Ba**) Plotting of mean firing rate vs. velocity of implanted mouse and distance between animals during the duration of intermale attack instances. (**Bb**) Linear regression between mean firing rate and velocity, and mean firing rate and distance between animals during the duration of intermale attack instances (pooled data across six mice). (**Ca**) Plotting of mean firing rate vs. velocity of implanted mouse and distance between animals during the duration of mounting a female conspecific. (**Cb**) Linear regression between mean firing rate and velocity, and mean firing rate and distance between animals during the duration of mounting a female conspecific (pooled data across six mice). (**Da**) Plotting of mean firing rate vs. velocity of implanted mouse and distance between animals during instances of freezing in the predatory-fear assay. (**Db**) Linear regression between mean firing rate and velocity, and mean firing rate and distance between animals during instances of freezing in the predatory-fear assay (pooled data across six mice). (**Ea**) Analysis design and input to partial least-squares regression, to assess the correlation between partial least-square components between neural data and motion-related variables. (**Eb**) Neural trajectories of PLS components in the PLS-space (example mouse). (**Ec**) Left: Variance explained by PLS components of the single-unit neural data. Right: Variance explained by PLS components of the motion-related variables (n=6 mice). (**Ed**) Fit between neural and motion-variable PLS components, indicative of a poor link between the two types of data (n=6 mice). (**Fa**) Analysis design and input to partial least-squares regression, to assess the correlation between partial least-square components between neural data and consummatory social and fear behaviors. (**Fb**) Neural trajectories of PLS components in the PLS-space (example mouse). (**Fc**) Left: Variance explained by PLS components of the single-unit neural data. Right: Variance explained by PLS components of the consummatory social and fear behaviors (n=6 mice). (**Fd**) Fit between neural and consummatory instinctual behavior-based PLS components, indicative of a link between the two types of data (n=6 mice). (**Ga**) Analysis design and input to partial least-squares regression, to assess the correlation between partial least-square components between neural data and appetitive social and fear behaviors. (**Gb**) Neural trajectories of PLS components in the PLS-space (example mouse). (**Gc**) Left: Variance explained by PLS components of the single-unit neural data. Right: Variance explained by PLS components of the

appetitive social and fear behaviors (n=6 mice). (**Gd**) Fit between neural and appetitive social and fear

behavior-based PLS components, indicative of a link between the two types of data (n=6 mice).

Figure relevant to Fig. 2.

**Fig. S6. "Ignition signals" occur across social and fear behaviors, across behavioral episodes,**

**typically 200 msec prior to behavior onset, and have an average duration of 1.5 seconds.** (**A**)

Example behavior raster plot indicating the repetitive nature of specific intermale behaviors following the

introduction of a male intruder in the resident-intruder assay. (**B**) Ignition signals (highlighted in light blue)

occur across behavior action episodes of approaching a male conspecific (n = 6 mice). (**C**), Ignition

signals (highlighted in light blue) occur across behavior action episodes of attacking a male conspecific

(n = 6 mice). (**D**) Characteristics of ignition signals include onset, offset and duration relevant to behavior.

E, Quantification of the onset of ignition signals relevant to the onset of behavior (n = 6 mice, neural data

averaged across three consecutive behavior instances per mouse). (**F**) Quantification of the ignition

signal duration (n = 6 mice, neural data averaged across three consecutive behavior instances per

mouse). (**G**) Quantification of the ignition signal offset subtracted behavior offset (n = 6 mice, neural data

averaged across three consecutive behavior instances per mouse).

In box-and-whisker plots, center lines indicate medians, box edges represent the interquartile range, and

whiskers extend to the minimal and maximal values. Abbreviations of brain areas are described in the

legend of Figure 1. Statistical comparisons of the data shown in panel (E-G) were performed through

One-Way ANOVA and Tukey's post-hoc test was used to correct for multiple comparisons. In box-and-

whisker plots, center lines indicate medians, box edges represent the interquartile range, and whiskers

extend to the minimal and maximal values. Figure relevant to Fig. 2.

**Fig. S7. Correlation matrix of behavior for each of the recorded six mice, with behavior actions sorted based on similarity.** Note the relative segregation of fear behavior actions (marked in red)

800    towards the end of the row labels in each matrix, while other behaviors are intermingled. Figure relevant

801    to Fig. 3, A and B.

**A** Single-neuron activity across behavior actions, and behavior action episodes

**Behavior domains**

| core behaviors | intermale behaviors | male-female behaviors | predatory fear behav. |
|---|---|---|---|
| sg: self-grooming | ms: sniffing male intruder | fs: sniffing female intruder | rs: sniffing predator |
| r: rearing | ma: attacking male intruder | fm: mounting fe male intruder | re: escape response |

**Brain areas**

● PMd ● PMvC ● PMvR ● TUc ● TUr ● VMHvlC ● DMH ● VMHvlR ● VMHcC
● VMHcR ● AHNca ● AHNce ● AHNr ● RCHc ● RCHr ● MPNc ● MPNv
● MPOAd ● MPOAc ● MPOAv

**B** intermale domain cluster

attack cluster

fear cluster

● first behavioural action episodes
● second behavioural action episodes
● third behavioural action episodes

**C**

**D**

**Fig. S8. Single unit activity across four behavior domains, eight behaviors of interest, and three**

**behavior episodes.** (**A**) Single unit activity in eight behaviors of interest during three consecutive

behavioral episodes/instances (n=6 mice, 2347 neurons). (**B**) Selected clusters from panel c Fig. 3,

indicating the activity of each cluster across the first, second and third behavioral action episodes. (**C**)

Correlation coefficient of average cluster activity across behavioral episodes (relevant to Fig. 3C). (**D**)

Fidelity score of cluster identity across behavior episodes; a measure that quantifies the degree to which

a cluster is active to the exact same behavior(s). A cluster considered active during the expression of any

single behavior, scales its activity above the threshold of two standard deviations (as compared to its

baseline activity).

ns: statistically not significant, *$P < 0.05$. Statistical comparisons of the data shown in panels (B, C) were

performed through One-Way ANOVA and Dunnett's post-hoc test was used to correct for multiple

comparisons. In box-and-whisker plots, center lines indicate medians, box edges represent the

interquartile range, and whiskers extend to the minimal and maximal values. Figure relevant to Fig. 3.

**NNMF component activity across behavior actions, and behavior action episodes**

**A**

Episode #1  Episode #2  Episode #3

Encoded behavior classes

Social

Fear

behavioral onset

Firing rate (spikes/s)

NNMF components (#) [sorted by similarity]

sg  r  ms  ma  fs  fm  rs  re

20 s

**Behavior domains**

| solitary behaviors | intermale behaviors | male-female behaviors | predatory fear behav. |
|---|---|---|---|
| sg: self-grooming | ms: sniffing male intruder | fs: sniffing female intruder | rs: sniffing predator |
| r: rearing | ma: attacking male intruder | fm: mounting fe male intruder | re: escape response |

**B**

*k-means* (quantification of data shown in Fig. 3**c**)  NNMF (quantification of data shown in panel **a**)

s *vs* f: ****  s *vs* f: ****  s *vs* f: ****  s *vs* f: ****

Cumulative frequency (#)

Activity of social vs fear clusters during social behaviors

Activity of social vs fear clusters during fear behaviors

Activity of social vs fear clusters during social behaviors

Activity of social vs fear clusters during fear behaviors

Firing rate (norm.)

**C**  **D**  **E**  50 cells

Rostral  Rostral

Hypothalamic brain areas  Hypothalamic brain areas

mice (9-1)

Caudal  Caudal

Neurons (norm.)

1  0

Neuron weights (norm.)

240 neurons

NNMF components (#)  NNMF components (#)

raw data  smoothened data (20 cells wide filter)

Brain areas

● PMd ● PMvC ● PMvR ● TUc ● TUr ● VMHvlC ● DMH ● VMHvlR ● VMHcC ● VMHcR ● AHNca ● AHNce ● AHNr ● RCHc ● RCHr ● MPNc ● MPNv ● MPOAd ● MPOAc ● MPOAv

**Fig. S9. Non-negative matrix factorization (NMF) identifies ensembles with behavior class specificity and spatial bias.** (**A**) NMF matrix factorization of the single neuron activity data shown in Extended Data Fig. 8, generates 39 NMF components (collectively explaining 79% of the variance of the single unit data, a number selected using the elbow method). Activity of components occurs somewhat similarly to the *k*-means clusters identified and shown in Fig. 3, during the social and fear behavior classes. (**B**) Cumulative frequency of recorded firing rates in "social" or "fear" clusters identified in the *k*-means (Fig. 3) and NMF (present figure) analysis respectively. Note that cluster activity is significantly different between conditions, but magnitude of effect is smaller in the NMF analysis. (**C**) Histogram (left) and raster plot (right) showing neuron origin from the six recorded mice. (**D**) Neuron weights of the NMF components (raw data, n=6 mice). (**E**) Neuron weights of the NMF components (smoothened data [Gaussian smoothing filter with twenty cells size], n=6 mice).

****$P < 0.0001$. Statistical comparisons were performed through the Kolmogorov-Smirnov test. Abbreviations of brain areas presented in the legend of Figure 1. Figure relevant to Fig. 3 and 4.

**A**

**B**

**Fig. S10. Solitary, social and fear behavior actions can be decoded from hypothalamic single**

**neuron activity using gradient boosted trees decoding models.** (**A**) Schematic of the analysis

pipeline using gradient-boosted trees for binary decoding of behavior based on single unit activity. (**B**)

Shuffled F1-score subtracted, k-fold cross-validated F1-score output of decoding behavior based on the

gradient boosted trees decoders (box plots are made of points from the six recorded mice – each mouse

contributing a single point unless a behavior was not expressed by an individual). In box-and-whisker

plots, center lines indicate medians, box edges represent the interquartile range, and whiskers extend to

the minimal and maximal values.

ns: statistically not significant, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. In box-and-whisker

plots, center lines indicate medians, box edges represent the interquartile range, and whiskers extend to

the minimal and maximal values. See also Fig. S11. Figure relevant to Fig. 3 and 5.

**Fig. S11. Intermale and male-female social behavior actions can be decoded with high accuracy by support vector machines decoding models.** (**A**) Example accuracy of three decoders on the test data of three social behaviors, and activity projected on the SVM hyperplane. (**B**) Decoder accuracy on behaviors of interest (n = 6 mice). (**C**) ROC curves on the observed and shuffled data (n = 6 mice). (**D**) Quantification of the area under the curve of the ROC curves shown in panel (**C**).

*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. Statistical comparisons of the data shown in panel (D) were performed against each group's shuffled control using paired two-tailed $t$-test. In box-and-whisker plots, center lines indicate medians, box edges represent the interquartile range, and whiskers extend to the minimal and maximal values. See also Fig. S10. Figure relevant to Fig. 3 and 5.

**Fig. S12. Single neuron activity variance explained by behavioral domains, motion variables, and sensory proxy signals.** (**A**) Variance explained (cross-validated, shuffled data $R^2$ subtracted, $R^2$) of single-neuron activity (n=6 mice, 2347 neurons) based on the four behavioral domains of interest, solitary, male intruder in arena, female intruder in arena, and predator in arena. Output of Poisson-GLM. (**B**) Behavioral domain coefficients on single neurons recorded across all mice (n=6 mice). (**C**) Variance explained (cross-validated, shuffled data $R^2$ subtracted, $R^2$) of single-neuron activity (n=6 mice, 2347 neurons) based on forty-two annotated behaviors. (**D**) Variance explained (cross-validated, shuffled data $R^2$ subtracted, $R^2$) of single-neuron activity (n=6 mice, 2347 neurons) based on forty-two instinctual behaviors and four motion variables, i.e., animal's velocity, animal's acceleration, distance and angle between animals. (**E**) Difference in the variance explained by behavior vs. behavior + motion variables. Histogram generated by subtraction of each neuron's $R^2$ generated via GLMs trained and tested only using annotated behaviors as predictor variables, against the neuron's $R^2$ generated via GLMs trained

865    and tested using annotated behaviors and motion variables as predictors. The peak around $R^2$=0, is

866    suggestive that motion variables explain little if any of the variance of hypothalamic single neuron activity.

867    Figure relevant to Fig. 5.

**Fig. S13. Contibution of neurons per mouse to the composition of clusters shown in Fig. 5C, and sorting of neurons based on their anatomical coordinates across the rostro-caudal hypothalamic axis (also related to Fig. 5C).** (**A**) Neuron contribution from each of the recorded mice to the neural branches presented in Fig. 5C. (**B**) Sorting neurons based on their spatial coordinates (as compared to the agglomerative linkage clustering-based sorting using behavior coefficients shown in Fig. 5C).

# Materials And Methods

**Experimental animals**

All experiments involving the use of living animals or their tissues adhered strictly to the guidelines set by the National Institutes of Health (NIH). They were conducted under the approval of both the Institutional Animal Care and Use Committee and the Institutional Biosafety Committee at the California Institute of Technology (Caltech). Swiss Webster (CFW) mice (Crl: CFW [SW]), C56BL/6NCrl mice, and Long-Evans (Crl:LE) rats were purchased from Charles River US. In all behavioral and resident-intruder assays, wild-type mice from the C56BL/6NCrl strain were used (sourced from Charles River Laboratories). Experimental animals were provided with additional enrichment and running wheels in their cages, were maintained in good health, and were monitored by the local veterinarian. Housing conditions for the mice were carefully monitored. Ventilated micro-isolator cages were maintained in an environment with stable temperature (23°C) and humidity (55-60%). A reverse light-dark cycle of 12 hours each was used, and animals were provided unlimited access to food and water. The cages were cleaned and changed once a week on a specific day and no experiments were conducted within two days of a cage change.

**Selection of animals for experiments**

*In vivo* electrophysiology data were collected from six Swiss-Webster mice (out of 11 implanted animals generated in a single cohort) and are reported here. Mice were selected for combined electrophysiology and behavior experiments based on the following criteria: (1) recovered within two days to pre-implantation baseline behaviors. (2) expressed the majority of instinctual behaviors on the experimental days; and (3) identified single neurons did not show drift across experimental sessions.

**Surgical methods**

All surgical and behavioral procedures adhered to the standards set by the National Institutes of Health and received approval from the Institutional Animal Care and Use Committee at Caltech. Male Swiss

Webster mice, aged between 3-4 months, were used for neural recordings during behavior. Male mice were used to avoid potential confounding effects of the estrous cycle in female mice.

Prior to surgery, mice were provided with medicated food cups (MediGel CPF, Clear H20 74-05-5022), containing carprofen, for two days. Throughout the surgical procedure, mice were kept anesthetized with isoflurane. The initial steps included removing the skin and clearing any underlying connective tissue. The tissue surrounding the skull edges was securely adhered with Vetbond (Santa Cruz Biotechnology, cat. no. sc-361931), and the skull surface was prepared by being scraped, dried, and textured.

A custom boomerang-shaped titanium head bar was affixed to the occipital bone of the skull, and a hole for the ground screw was carefully drilled, placing the screw above the right prefrontal cortex. Optibond Solo Plus (Kerr, cat No. 31514) was applied to the skull and cured under UV light. Charisma (Net32, cat. No. 66000085) was employed to establish a base for the implant and to reinforce the area around the ground screw.

With the help of a dental drill, eight 100 μm holes were made above the somatosensory cortex. These holes were designed to accommodate the eight shanks of the silicon probe, aimed at penetrating the deep brain. The entire silicon probe assembly, including the probe and headstage, was placed in the stereotax. Post-application of DiD' (ThermoFisher Scientific, cat. no. D7757) on the posterior to the electrode surface of each shank, the probe was gently lowered into the brain at a speed of 6 μm/sec resulting to an insertion time of fifteen minutes. Higher insertion speeds negatively impacted post-surgery recovery and neuron yield per mouse.

The precise coordinates for probe insertion and the angle of attack were adapted for each mouse accommodating idiosyncratic differences in cortical vasculature identified post-drilling, and enabling the recording of the same deep brain structures. The ground wire was soldered to the ground screw and Metabond cement was used to secure the probe and headstage to the skull. The entire assembly was wrapped in Kapton Tape (ULINE S-7595), followed by a recovery period typically extending to two weeks. Post-surgery, mice were individually housed, and additional enrichment elements were provided. Daily monitoring was performed to evaluate locomotion and the expression of behaviors (self-grooming, rearing

and digging). Between the second and third weeks, mice were acclimatized in test arenas to examine

their natural movement and the manifestation of thigmotaxis in the open field test, among other behaviors.

**Behavioral tests**

All behavioral tests were performed during the dark phase and under dim red light between 2 h after the

initiation of the dark phase and three hours before the initiation of the light phase. Mice were acclimated

to the testing behavior room for a minimum of one hour before testing with *ad libitum* access to food and

water. Videos were recorded using the GS3-U3-51S5C-C 2/3" FLIR Grasshopper®3 USB 3.0 Color

Camera, at thirty frames per second. Video acquisition was triggered through a TTL signal provided to

the *trigger*-in GPIO port of the camera, by the neural data acquisition software Allego – NeuroNexus'

proprietary software for data acquisition. Timestamps for frames were collected and recorded as an

auxiliary signal through the software Allego and NeuroNexus' data acquisition box (SmartBox Pro).

**Resident–intruder (RI) test**

The resident's cage was not cleaned for a minimum of 2 days before the behavioral test. A single male

or female intruder was introduced per testing session. A ten-minute habituation period was allowed and

recorded, and was followed by the introduction of a freely moving intruder for ten minutes or an intruder

within a perforated cage. Video and neural data acquisition continued for ten minutes following the

removal of the intruder. Intruders were of the C57BL/6N background, and were either intact males, or

intact females in the estrus phase of the estrous cycle (stage of cycle was defined by inspection of vaginal

smears, and the cycle of individual female intruders was followed for three consecutive days prior to the

day of the experiment). Behaviors during the RI test were video recorded using a top view camera, and

were manually annotated using BENTO (*70*) (https://github.com/neuroethology/bentoMAT).

**Predatory fear (PF) behavioral assay**

A custom four-chamber arena was used to present a caged-predator. All but the front sides of the cage of the predator were opaque white colored, with the exception of the front side that was clear and the wall was perforated evenly with square holes five square millimetres in size placed fifteen millimetres apart (center to center). This design aims to allow diffusion of pheromonal cues that may affect the behavior of pray animals. Other than the chamber containing the predator, the arena includes three other chambers: i) the interaction zone chamber that allows the mouse to sniff the predator and interact with the cage before, during and after the presence of the rat in the arena. ii) The transition zone chamber that allows the mouse to move between the proximal and furthest away chambers from the rat, and iii) the safe zone chamber, that places a large distance between the mouse and the predator. The validation parameters of this custom experimental design are presented in extended data figure four.

**Behavior annotations and quantification of behavior**

Behavior videos were manually annotated using BENTO (*70*) (freely available software package: https://github.com/neuroethology/bentoMAT). A total of forty-three behaviors during the resident intruder and predatory fear assays were annotated. The behavioral videos were loaded into MATLAB and a trained individual blind to the experimental design annotated the videos frame-by-frame for all behavior actions. Behavior annotations were then further edited by an expert (author: S.S.) who was not blind to the experiment conditions. Imaging frames were spatially downsampled by a factor of two, a modification that we recognize to not influence identification of the behaviors of interest and that facilitates smoother processing of the videos by BENTO (*70*) in MATLAB. Behavior features including velocity, acceleration, distance and angle between animals were calculate using DeepLabCut (*38*), and custom MATLAB routines.

***In vivo* electrophysiology**

*In vivo* electrophysiology recordings were performed in freely moving mice, using chronic silicon probe

implants. All extracellular recordings were conducted across the rostro-caudal hypothalamic axis of the

left hemisphere. Recordings were performed using a 256-channel custom-built for this specific project

silicon probe, the A8x32-Poly2-7.5mm-25s-325-177 NeuroNexus probe, now commercially available.

This specific probe includes eight shanks evenly spaced by 325 μm, and each shank at its lower end

contains two columns of electrodes, each electrode with 25 μm distance from the next. The recording

area of each electrode was designed to 25 μm$^2$. The length of each shank was 7.5 mm to enable the

probe to reach the model ventral nuclei of the deep brain in the hypothalamus.

Electrophysiology data was collected with Allego (proprietary software of NeuroNexus, recently updated

to Radiens [https://www.neuronexus.com/products/software/radiens/]), at 20 kHz sampling frequency.

Kilosort 2.5 was used for spike sorting (*30*), and we refined its output with manual curation of the results

through phy: https://github.com/kwikteam/phy. Neurons that exhibited drift or units that had multiunit

activity were excluded. To assess the neural recordings and perform basic related analysis prior to

exploring the data relevant to behavior, we used MATLAB code that can be found here:

https://github.com/cortex-lab/spikes. The quality of recordings was assessed through quantification of the

event rate, i.e., the frequency of spikes that are higher than the background noise level. We considered

an 'event' to be any occurrence that matches in time (less than 1 millisecond) and space (50 millimetres

in radius) and that has a strength that is at least six times greater than the median absolute deviation

(MAD) (*71*). Secondly, for unit quality assessment we computed and plotted the signal-to-noise ratio

(SNR) for each event. Event detection (spikes), spike amplitudes, and firing rates were further inspected

with custom MATLAB routines (https://github.com/jenniferColonell/Neuropixels_evaluation_tools).

1003

**Whole-brain Imaging and Anatomical Registration of Probe Electrodes**

Brains that had been cleared were imaged using the LaVision BioTec UltraMicroscope II in dibenzylether

at two different wavelengths: 488 nm (which creates green autofluorescence) and 632 nm (which allows

for the visualization of electrode tracks through the DiD dye, a fixable, lipophilic far-red dye that remains

stable even in dehydrated conditions). The imaging was done using an Olympus MVPLAPO 2X/0.5

objective, a 5 μm step size, approximately 5 μm pixel size (at 0.8X magnification), and a single light sheet

focus. The imaging only included the hemisphere of the brain where the electrode tracks were located,

and it was carried out in a sagittal orientation.

Images obtained from the green and far-red channels were subsequently downsampled to a 25 μm

resolution by means of averaging. The green channel was then matched to the Allen Brain Atlas CCFv3

autofluorescence atlas with a 25 μm resolution using a two-step process: an initial affine transformation

followed by a non-linear transformation carried out in Elastix. Previous studies have demonstrated that

this method has an average registration error of about 4 voxels (or 100 μm). The same transformation

process was also applied to the red channel.

Electrode tracks were traced in the registered space using a custom R code that is based on AllenCCF

(https://github.com/cortex-lab/allenCCF). This was done after electrophysiological recordings were made

following the placement of a chronic implant that targeted the full anterior-posterior length of the

hypothalamus.

For the final analysis, brain regions were organized into twenty distinct groups based on the Allen Brain

Atlas hierarchy. The specific location of each unit was determined based on where that unit produced the

highest amplitude on the electrode. The location along each electrode was transformed into the Allen

Brain Atlas space according to the orientation and position of the traced electrode track for each

recording. The final part of allocating each electrode to a hypothalamic region was performed manually,

to further improve the output of the earlier-mentioned process. For each mouse, a unique spatial map of

electrodes/hypothalamic brain areas was defined and was later used to assign individual extracellular

units to specific anatomical regions.

1030

**_k_-means clustering**

The variance explained by a clustering solution in the context of the $k$-means algorithm reflects the

proportion of the total variance in the dataset that is "captured" by the centroids of the clusters. In other

words, it indicates how well the centroids represent the data. For computing the variance explained is we

used the sum of squared distances between data points and their assigned cluster centroids. The more

the variance explained, the closer the data points are to their respective centroids, and hence the better

the clustering solution. Specifically, we calculated within-cluster sum of squares (WCSS). After

performing *k*-means clustering we obtained the cell indices, which provide the cluster assignments for

each data point, and the centroids for the clusters. The sum of squared distances of each point to its

cluster centroid gives us the WCSS. We then iterated over all data point. For each point, we calculated

the squared Euclidean distance between the point and its assigned centroid and summed these squared

distances. We then computed the total sum of squares (TSS). i.e., the sum of squared distances of each

point from the overall mean of the data. The variance explained by the clustering solution can be found

by subtracting the WCSS from the TSS and then dividing by the TSS. Note that as the number of clusters

*k* increases, the WCSS tends to decrease (each point can be closer to its centroid), but there's usually a

diminishing return. This is often visualized with an "elbow plot," where the rate of decrease in WCSS

slows down after a certain number of clusters, indicating a potential good choice for k.

1048

**GLM output – functional vs. spatial distribution of clusters**

Neuronal cluster data was analyzed to determine its relationship with four predefined behavioral

categories. For each cluster, its activity was averaged across specific row ranges corresponding to each

category. Negative activity was disregarded, and data was normalized using the z-score to mitigate outlier

effects. The entropy was computed for each cluster based on its normalized activity distribution across

behavioral categories. Entropy was utilized to quantify the distribution of each neuronal cluster's activity

across the four behavioral categories. After averaging the activity for each category and normalizing

these averages to form a probability distribution, the entropy was computed using the formula for a

discrete probability distribution. This measure provided insights into whether a cluster's activity

predominantly aligned with specific behavioral categories (low entropy) or was distributed across multiple

categories (high entropy). For spatial analysis, a 'center of mass' approach was employed to gauge the

distribution of each cluster's activity across 20 assumed brain regions. Both the functional (behavioral

category-based) and spatial distributed-ness of each cluster were then visualized using a scatter plot.

1062

**Decoding Analysis**

Gradient boosted trees: neuronal data (spike rates, binned at 100 ms) was smoothened through

convolution with a Gaussian kernel of 10 timesteps (1 second). The data to the decoder was then made

recurrent by creating a temporal window of size ten at each given timestamp. The data was then flattened

before decoding. The data was analyzed using five-fold cross-validation. Each behavior was decoded

individually based on the activity of all neurons. Folds were created by determining the onsets of the

behavior and then creating a temporal window of the length of the behavior before its onset. All events

were concatenated, and cross-validation was performed on the resulting data. The decoding was

performed using gradient-boosted trees (*72, 73*) (depth of 3, 50 estimators, lr = 0.01). Performance was

evaluated by calculating the cross validate, shuffled data subtracted F1-score per behavior. For each

behavior, we calculated the maximum performance within a specific temporal jitter (10 timesteps). We

then averaged performances across animals.

Support Vector Machines: To assess if single neuron activity holds information about specific social

actions, we trained a unique binary linear decoder for each behavior action, to predict whether or not the

behavior was taking place. To ensure the behavior's representations weren't mixed with the

representations of the intruder's gender, we created separate decoders for behaviors directed towards

males and females. During times when the behavior of interest was not being exhibited, we under-

sampled the negative training data to balance the number of positive and negative training examples.

We then used these binary labels alongside their matching activity vectors to train a linear SVM decoder.

1082

1083      **Code Availability**

1084      The code used for analysis in this study is available upon request from the first author

1085      ([stefanos.stagkourakis@gmail.com](mailto:stefanos.stagkourakis@gmail.com)).