araucan **regulates butterfly wing iridescence by coordinating scale**

structure and pigmentation

ABSTRACT

Butterfly wings exhibit a remarkable diversity of structural iridescent colors, yet the

genetic regulation of iridescence remains poorly understood. Here, we show that the

Iroquois-complex transcription factor gene *araucan* plays a role in modulating wing

scale iridescence in the common buckeye butterfly, *Junonia coenia*. Using CRISPR-

Cas9 knockouts, we demonstrate that loss of *araucan* function causes dorsal wing

scales to shift from golden-brown to blue iridescence, and eyespot center scales to shift

from saturated purple-violet to dull grey-brown. These phenotypes are associated with

changes in thickness of the lower lamina in scale cells, a structural feature known to

influence thin-film interference, along with reduction of pigmentation in ground scales.

Together, these results identify *araucan* as a single transcription factor that determines

coloration by simultaneously regulating both photonic architecture and light-absorbing

pigmentation, and show that these effects can be regulated in pattern-specific manner.

Our findings provide a framework for linking gene regulation to the spatially coordinated

evolution of structural and pigmentary components of color.

Keywords

structural coloration, iridescence, butterfly wing patterns, *Iroquois*-complex, evo-devo

INTRODUCTION

Coloration in animals serves as a powerful model for understanding how gene regulatory networks give rise to complex, adaptive traits. Color is most commonly generated by light-reflecting pigments, by structural features that alter light reflectance spectra, or through a combination of both. Although structural coloration is widespread across the animal kingdom—particularly in insects—its genetic and developmental bases remain poorly understood (1). Butterfly wings, with their nanostructured scales and vivid structural organization that produce blue, purple, and green hues, offer a useful system for uncovering how genes shape the physical traits that underlie these colors.

Over the past decade advances in genomic mapping and CRISPR-Cas9 gene editing have enabled the discovery and functional validation of key genes and regulatory elements involved in butterfly wing pigmentation and patterning. Comparative studies and insights from model insects such as *Drosophila melanogaster* have further expanded the developmental genetic framework for wing pattern determination. However, the genetic basis of structural coloration remains less well resolved, with only a handful of causative genes identified across a few butterfly species. These include *optix*, a regulator of blue iridescence in *Junonia coenia* (2,3) and silver scale development in *Speyeria mormonia* (4); *bric-a-brac* and *doublesex*, which control UV-reflecting scales in *Colias spp.* (5) and *Zerene spp.* (6), and *antennapedia*, *apterousA*, *doublesex*, *ultrabithorax*, *optix*, and *hr38*, which regulate silver reflectance in *Bicyclus anynana* (7,8). Of these examples, *optix* is the only gene thus far characterized to play a role in iridescent coloration.

Here, we investigate the role of *araucan/caupolican* (also known as *iroquois2*), an *Iroquois*-complex transcription factor gene, in the wing patterning of the common buckeye butterfly *J. coenia*. Most hexapods, including butterflies, possess two *Iroquois* genes—one orthologous to *D. melanogaster's mirror* (*iroquois1*) and one to *araucan/caupolican* (*iroquois2*). In *D. melanogaster iroquois2* has undergone a lineage-specific duplication, giving rise to the paralogs *araucan* and *caupolican* (9–11). The

three *Iroquois* genes (*araucan*, *caupolican*, and *mirror*) in *D. melanogaster* regulate

multiple developmental processes, including dorsal notum patterning, longitudinal wing

vein specification, and posterior wing domain (alula) identity (12–16). Recent findings

suggest that *mirror* specifies a far posterior domain in butterfly wings—a role likely

ancestral in broad winged insects but lost in *D. melanogaster* and other calyptrate flies

due to evolutionary reductions in wing complexity (11). Despite its expression in the

basal part of the wing in *Heliconius* butterflies (17), the function of *araucan/caupolican*

(hereafter *araucan*) in butterflies has thus far remained untested.

In this study, we used CRISPR-Cas9 mutagenesis and gene expression analyses to

investigate the role of *araucan* in wing coloration in *J. coenia*. We show that *araucan* is

a regulator of both wing scale ultrastructure, including the iridescence-determining lower

lamina thickness, and scale pigmentation. Our findings identify *araucan* as a novel

regulator of coloration in butterflies that functions via a process of coordinated pigment

and structural effects. This work provides the first functional evidence for this gene

outside of *D. melanogaster*, highlighting its evolutionary flexibility and specialized role in

modulating scale-based structural color traits.

MATERIALS AND METHODS

Identification of *Iroquois*-complex genes

We previously identified the *Iroquois*-complex genes in *J. coenia* (11). For this study, we

used an updated annotation (18), which resulted in revised gene model identifiers for

the *Iroquois*-complex. We identified the *J. coenia* gene *JC_g12213* as the single

ortholog of *D. melanogaster araucan/caupolican* paralogs (figure S1). For clarity and

consistency, we refer to *JC_g12213* as *araucan* in *J. coenia*, with the understanding

that it is equally related to both the paralogs in *D. melanogaster*. The updated

phylogeny and sequence information used for orthology assignments are provided in

figures S1 and file S1, respectively. To ensure reproducibility and compatibility with

future genome and annotation releases, we have provided both the predicted amino

acid and coding DNA sequences of *araucan* in the supplementary materials (file S1).

CRISPR-Cas9 mutagenesis of *J. coenia araucan*

To assess the function of *araucan* in *J. coenia* we generated CRISPR-Cas9 somatic

mosaic deletions targeting the *araucan* coding region. We used a previous protocol

where early embryos are injected with single guide RNAs (sgRNAs) and Cas9 enzyme

(19). Mutations induced in early syncytial embryos typically only affect a subset of

nuclei, resulting in mosaic knockout (mKO) animals.

We designed two sgRNAs – 5'-CTGGGTACGATTTAGCAGCC-3' and 5'-

CTCCTGCTCCTTGTCCTCGT-3' – to target the exon of *araucan* containing the

conserved homeobox domain in *J. coenia* (figure S1a). Possible off target sgRNAs were

ruled out by blasting against the *J. coenia* genome assembly. sgRNAs were

synthesized by Integrated DNA Technologies, Inc. (Coralville, IA, USA), rehydrated in

1x Tris-EDTA buffer, and mixed with Alt-R S.p. Cas9 Nuclease/Nickase (Integrated

DNA Technologies, Inc., Coralville, IA, USA) to a final concentration of 200ng/ul sgRNA

and 500ng/ul Cas9.

We collected eggs by placing *Plantago lanceolata* leaves in a cage with gravid females

for 1-2 hours, and then prepared eggs for microinjection by submersing them in 5%

benzalkonium chloride (Sigma-Aldrich, St Louis, MO, USA) for 30s followed by a wash

with distilled water. Prepared eggs were then lined up on double sided tape on glass

slides and sgRNA and Cas9 mixture was injected into the eggs using 0.5-mm

borosilicate needle (Sutter Instruments, Novato, CA, USA) and a PLI-100 Picoinjector

(Harvard Apparatus, Holliston, MA, USA). Injections were completed within 2-3hrs after

oviposition, and larvae from injected eggs were raised on artificial diet at 27°C, 70%

humidity, and a 16hr : 8hr light : dark cycle. Adults were frozen upon emergence, and

phenotyped and imaged using a Keyence VHX-7000 microscope.

Genotyping CRISPR mutants

We genotyped *J. coenia* mosaic knockout (mKO) mutants to confirm Cas9-induced

deletions in the *araucan* gene. Mutants "mKO A" and "mKO 3" were selected based on

their blue cover scale phenotypes (figure S2), while "mKO M" was chosen for its

symmetrically disrupted eyespot phenotype (figure S4). We extracted DNA (OMEGA

BIO-TEK, Norcross, GA, USA) from the thoracic wing muscles of butterflies that showed

obvious wing pattern anomalies compared to wild types. We used PCR to amplify DNA

from around the potential cut sites using the primers 5'-ATACTAAACCCTGCACCCGC-

3' and 5'-ACGTACTCACAAGCGAACGT-3'. We ran amplified DNA from wild type and

mutant individuals on an agarose gel and purified (Monarch DNA Extraction and

Purification kit, new England Biolabs, MA, USA) and cloned the PCR products into

JM109 competent cells using pGEM®-T Easy Vector Systems (Promega Corporation,

Madison, WI, USA). Transformed cells were amplified and Sanger sequenced at the

Genomics Facility, Cornell Institute of Biotechnology, Ithaca, NY, USA.

Micro-spectrophotometry of wing scales

We mounted individual wing scales (n = 3) from wild-type and *araucan* mutant *J. coenia*

butterflies on glass slides and measured their normal-incidence reflectance spectra

using a mercury-xenon light source (Thorlabs, New Jersey, USA) connected to a

uSight-2000-Ni micro-spectrophotometer (Technospex Pte. Ltd., Singapore). We used a

Nikon TU Plan Fluor 100× objective (numerical aperture = 0.9) to collect spectra and

calibrated the setup with a dielectric mirror. For each measurement, we averaged 10

scans with an integration time of 100 ms. We recorded three measurements from

different locations on each of three scales per genotype and averaged them to obtain a

representative reflectance spectrum.

To measure absorbance, we used the same optical setup with a 20× objective (NA =

0.5). We mounted individual scales (n = 5) submerged in clove oil (Hayashi Pure

Chemical Ind., Ltd.) between a slide and coverslip to match the refractive index of chitin

and used a clove oil-covered transparent area of the slide as a reference. For each

genotype, we measured three locations on each of five scales and averaged the data to

generate absorbance spectra.

We performed all spectral analysis and visualization using the pavo 2 package in R

(20). To test for significant differences in spectral values between wild-type and *araucan*

mutant scales, we conducted independent two-sample t-tests at each 50 nm increment

from 350 to 700 nm. We applied the Benjamini-Hochberg False Discovery Rate (FDR)

method to adjust p-values for multiple comparisons and considered adjusted p-values <

0.05 as significant.

Scanning electron microscope imaging of wing scales

We used scanning electron microscopy (SEM) to assess structural differences between

wild-type and *araucan* mutant scales. We mounted individual scales (n = 5) from mutant

mosaic clones and contrasting wild-type regions on carbon tape and sputter-coated

them with platinum (JEOL JFC-1600) for 30 s at 30 mA. We imaged the samples using

a FEI Versa 3D with the following parameters: voltage 10 kV, current 32 pA. To obtain

transverse-sections of wing scales, we used Focused Ion Beam (FIB) milling with the

gallium ion beam of the FEI Versa 3D, applying a beam voltage of 8 kV, beam current of

25 pA, and a tilt of 52°. We performed the FIB-SEM in the Electron Microscopy Facility,

National University of Singapore.

We obtained the transverse section of the scales shown in Figure 1c by cryo-fracturing

the scales following a previously described protocol (21). We then mounted whole and

cryo-fractured scales of interest on carbon tape and sputter-coated them with an 8 nm

layer of gold using a Cressington 208HR Sputter Coater. We acquired the image on an

FEI Teneo LV SEM using the Everhard Thornley detector and beam parameters of 5.00

kV/6.3 pA with a 10 µs dwell time. We used cryo-fracturing only to generate the image

used in figure 1c. All measurements for laminar thickness reported were done on scales

that were milled using FIB and imaged using SEM.

Measurement of scale ultrastructure and statistical analysis

We quantified scale ultrastructure by measuring three parameters: lower lamina

thickness (LL), inter-ridge distance (d_{RR}), and crossrib ($d_{CR\text{-}CR}$) spacing (figures 1c-d).

We performed all measurements in Fiji using the line tool (22). Lower lamina thickness

was corrected for tilted perspective (measured thickness/sin 52°) (23). For each

phenotype (i.e. wild type or *araucan* mutant), we analyzed five individual scales per
wing region and scale type. From each scale, we recorded 10 measurements of lower
lamina thickness, 25 measurements of inter-ridge distance, and 50 measurements of
crossrib spacing. We collected measurements from cover and ground scales in the
forewing eyespot center, and from cover scales in the dorsal non-eyespot region.

We analyzed scale ultrastructure data using linear mixed-effects (LME) models
implemented in the R package nlme version 3.1 (24) which accounts for hierarchical
data structures. For each geometric parameter (lower lamina thickness, inter-ridge
distance, and crossrib spacing), we treated scale type as a fixed effect and included
scale identity nested within individual butterfly as a random effect. To address
heteroscedasticity among scale types, we applied the varIdent() function. We selected
the best-fitting model based on the Akaike Information Criterion (AIC). We conducted
pairwise comparisons of group means using the multcomp R package. We report p-
values and 95% confidence intervals for all pairwise comparisons in Tables 1 and 2. A
summary of the fixed effects from the linear mixed-effects models, including degrees of
freedom, F-values, and p-values, is provided in Supplementary Tables S2 and S3. Raw
measurement data are also included to support independent reanalysis (see Data
Availability).
203
Hybridization chain reaction fluorescent *in situ* hybridization of *araucan*
We designed a hybridization chain reaction (HCR) probe set targeting the coding
sequence of *J. coenia araucan* using 19 pairs of 25 bp probes (50 bp per site) spanning
the annotated transcript (file S2). We ordered the probes as oligonucleotides from
Genewiz (Azenta Life Sciences, Morrisville, NC). We confirmed probe specificity using
BLAST against the most recent *J. coenia* genome assembly, ensuring no off-target
binding sites.
211
212 For HCR experiments, we dissected last instar larval imaginal discs, as well as pupal
213 wings at 13% (first day post-pupation), 25% (second day post-pupation) and 45% (third
214 day post-pupation) of pupal development. We used *araucan* probes paired with B1

initiators and visualized them with Alexa Fluor 647-labeled hairpins. As a positive

control, we used *spalt* probes labelled with B3 initiators (file S2) and visualized them in

the 546 nm channel. For figure S6 we performed HCR of *optix* with B5 initiator (file S2).

We performed co-detection of *araucan* and spalt expression in wings from one side of

each individual, while using the contralateral wings as negative controls. In these

controls, we omitted the probes but followed all other HCR steps to assess background

fluorescence. We performed the HCR following the protocol described by Bruce et al

(dx.doi.org/10.17504/protocols.io.bunznvf6). To capture for pattern specific expressions

across the entire wing tissue we used a Keyence BZ-X800 fluorescence microscope in

the laboratory of Dr. Anyi Mazo-Vargas (Department of Biological Sciences, Duke

University).

RESULTS

araucan knockouts result in iridescent blue scales via laminar thickening

We injected 758 eggs with Cas9 and *araucan* sgRNAs, of which 179 hatched and 117

survived to adulthood. Twenty-two of the adults that emerged had wing color mutations.

We observed two distinct classes of mutations, both of which involved iridescent

coloration: (1) blue iridescence in mutant clones across the dorsal wing surface (n = 13),

and (2) a shift from purple-violet to grey-brown specifically in eyespot centers (n = 9).

Genotyping confirmed that individuals from both mutant classes had lesions at the

sgRNA sites in the *araucan* coding region (figure S1c).

The thirteen individuals that showed mosaic clones of blue iridescence showed the

phenotype exclusively on dorsal wing surfaces (figure 1a; hindwings: figure S2;

forewings: figure. S3). Since iridescent blue colored scales can be artificially selected

for in populations of *J. coenia* (3), we assessed individuals from our wild type colony for

similar iridescent scales to rule out the possibility of phenotypes being due to natural

variation. Out of the individuals randomly observed (n = 32), we did not observe any

blue iridescent scales on the dorsal surface of the hind wings, which supports that the

blue iridescence in the mKOs is a result of knockdown of *araucan* (chi-squared test, *p* <

0.005). This conclusion was also supported by the fact that individuals with iridescent

patches of scales presented this phenotype asymmetrically (figures S2 and S3),

consistent with cell lineage mosaicism (2), whereas natural variation in iridescence is

typically symmetrical.

The shift from golden-brown scales in the wild type to vivid blue in mutants was

quantified using reflectance spectrophotometry. Mutant scales exhibited increased

reflectance in the ultraviolet and blue wavelengths (< 500 nm), while wild-type scales

reflected more light in longer wavelengths (> 500 nm), consistent with their golden-

brown appearance (figure 1b). To determine whether this observed color change was

structural or pigmentary in origin, we immersed individual scales in oil, which matches

the refractive index of chitin and thereby suppresses structural coloration. As in previous

studies, absorbance spectra collected under these refractive index-matched conditions

provide a measure of the light transmitted through the scale and served as a proxy for

pigment content, as differences in pigment type and concentration result in distinct

patterns of wavelength-specific light absorption (3,7,21,25–28). We observed minimal

differences in absorbance of visible wavelengths between mutant and wild-type scales

(figure 1b; table S2), indicating that the shift in coloration is primarily due to structural

changes rather than differences in pigmentation.

To investigate the ultrastructural basis of the iridescent blue coloration in *araucan*

mutant scales, we measured the thickness of the lower lamina—a key structural

determinant of scale color in butterflies of the *Junonia* genus, where a thicker lower

lamina causes a shift towards bluer hues (3). We quantified laminar thickness in both

wild-type brown scales and blue mutant scales to assess whether this relationship held

true in our *araucan* knockouts. SEM imaging after FIB milling to get the transverse

section of scales revealed that mutant cover scales had significantly thicker lower

laminae compared to wild type (figure 1c; LME model, AIC-selected best fit, $p < 2 \times$

10^{-16}; tables 1, S1). The mean corrected lower lamina thickness in wild-type scales was

96.7 ± 10.2 nm, while mutant scales averaged 152.0 ± 18.9 nm, supporting the

hypothesis that the observed color shift is structurally mediated.

To assess whether additional ultrastructural features contribute to the iridescent color

shift observed in *araucan* mutants we quantified ridge spacing and crossrib distances in

cover scales. These geometric features determine the size of the scale "windows" that

determine the amount of incident light that interacts with underlying scale

nanostructures (figure 1d). Mean ridge-to-ridge spacing was comparable between wild-

type brown and mutant blue scales (WT: 1921 ± 90 nm; mutant: 1921 ± 73 nm; figure

1d; tables 1, S1), and crossrib distance was also unchanged (WT: 923 ± 167 nm;

mutant: 914 ± 160 nm; figure 1d; tables 1, S1). These results suggest that the shift to

blue structural coloration in the *araucan* mutants is not due to changes in ridge or

crossrib geometry, but instead is primarily driven by increased lower lamina thickness.

araucan knockouts eliminate purple-violet iridescence in forewing eyespot

centers

Nine of the recovered *araucan* mKOs exhibited a striking phenotype affecting the scales

in the forewing eyespot centers. In wild-type *J. coenia*, these eyespot centers exhibit a

vivid purple-violet iridescence (figures 2a and S4). In contrast, *araucan* mutants lacked

this strong purple-violet coloration, which was replaced by a muted grey-brown hue,

occasionally with faint traces of pink or violet iridescence (figures 2a and S4). The loss

of purple-violet iridescence was asymmetrical in all but two individuals (figure S4),

validating the mosaic nature of the knockout phenotype. Notably, eyespot centers on

the hindwings were unaffected.

The iridescent purple-violet coloration of *J. coenia* eyespot centers arises from the

combined optical properties of both cover scales and underlying ground scales. The

ground scales enhance the overall saturation of the reflected color when stacked

beneath the more translucent cover scales (highlighted by yellow dashed outlines in

figure 2a). Because this mechanism is consistent with observations in other butterfly

species exhibiting purple-violet coloration (29,30), we characterized the optical

properties of each scale type within the eyespot center. Specifically, we measured

reflectance spectra and refractive index-matched absorbance spectra of both cover and

ground scales in the forewing eyespot center to identify which scale type and optical

mechanism was contributing to the observed color shift.

Spectral analysis showed that both scale types, cover and ground, were altered in the

araucan mutants differently, resulting in the observed color shift. Mutant cover scales

exhibited increased reflectance above 500 nm, corresponding to a duller grey-brown

coloration, whereas wild-type cover scales reflected more light in the ultraviolet to violet

wavelengths (380–450 nm), producing the observed purple-violet hue (figures 2a, b). In

contrast, reflectance measurements of ground scales revealed no significant differences

between wild-type and mutant individuals (figure 2c). To assess pigment contributions

independent of structural effects, we measured absorbance in a refractive index–

matched medium. Under these conditions, both wild-type and mutant cover scales

showed minimal light absorption in the ultraviolet and red regions, with no significant

differences between groups, indicating that most incident light was transmitted through

the cover scales (figure 2b). However, mutant ground scales showed a different pattern

across the ultraviolet and visible spectrum (300–700 nm) where they exhibited

significantly lower absorbance compared to wild-type scales, suggesting reduced light-

absorbing pigmentation in the mutants (figure 2c).

To identify structural components in the scales that might have been altered, we

characterized lamina thickness, ridge spacing, and crossrib distances in both cover and

ground scales. We found that the lower lamina was significantly thicker in the eyespot

scales of wild-type individuals compared to *araucan* mutants in both scale types. In

cover scales, the lower lamina thickness averaged 151 ± 16.9 nm in wild types and 112

± 13.1 nm in the mutants (LME model, AIC-selected best fit, $p < 2e\text{-}16$; figure 2d; tables

2 and S3). A similar pattern was observed in ground scales (146 ± 13.9 nm in wild types

vs 108 ± 10.3 nm in mutants, $p < 2e\text{-}16$; figure 2e; tables 2 and S3).

Ridge spacing was significantly narrower in mutants versus wild-type scales, for both

scale types. In cover scales, ridge spacing was 2123 ± 82.9 nm in wild types and 1805

± 62.9 nm in mutants ($p < 2e\text{-}16$; figure 2d; tables 2 and S3), and in ground scales,

1470 ± 116 nm in wild types versus 1188 ± 74 in mutants (p < 2e-16 figure 2e; tables 2

and S3). In contrast, the distance between crossribs was significantly greater in *araucan*

mutants for both scale types. In cover scales, mutants had crossrib spacing of 1202 ±

322 nm compared to 946 ± 217 in wild types (p < 2e-16; figure 2d; tables 2 and S3); and

in ground scales, mutants had 663 ± 103 nm of spacing vs 631 ± 79.3 nm in wild types

(p = 1.42e-05; figure 2e; tables 2 and S3). Together, these optical and structural

analyses reveal that *araucan* knockouts disrupt the pigmentation of ground scales and

the nanostructural organization of both cover and ground scales in the eyespot centers.

In combination, these changes likely underlie the loss of saturated purple-violet

iridescence and its replacement with muted hues in *araucan* mutants.

Dynamic spatial and temporal expression of *araucan* in the developing wing

Our mKOs demonstrate that *araucan* is necessary for normal structural coloration of *J.*

coenia wing scales and appears to play a role in determining scale structure itself.

Accordingly, we hypothesized that we should observe *araucan* expression in wings

during scale morphogenesis. Because *Iroquois*-complex genes—including *araucan*—

exhibit pattern-specific expression in the *D. melanogaster* wing disc (12,13), we further

sought to ask whether *araucan* is expressed in a pattern-specific manner in the

developing *J. coenia* wings, including in regions affected in mutants (e.g., eyespot

centers), or in other defined wing domains across development. To characterize spatial

transcriptomic patterns, we performed HCR *in situ* hybridization for *araucan* mRNA at

key developmental stages: (1) in last-instar wing discs, immediately during and after the

transcription factor Spalt defines future eyespot domains (31–33); and (2) in pupal

wings at two time points -13% pupal development (PD) , when eyespot centers have

been established, and 25% PD, when scale precursors differentiate into scale cells (34).

In last-instar discs, *araucan* transcripts are enriched in the posterior wing domain

(vannus) bounded by the 2A vein (figure 3a). Additional signal appears in an anterior–

proximal region and in enlarged cells arrayed along the trachea that mark future veins

and the wing margin (figures 3a and S5b). At ~13% PD, the posterior-domain

expression persists and *araucan* is detected in large cells interspersed along the veins

of both forewings and hindwings (figure 3b). By ~25% PD, posterior-domain signal

contracts to a narrow stripe flanking the 2A vein, while expression becomes broadly

peppered across both wings without aligning to obvious adult color elements (figure 3c).

This expression, along with the expression in the veins, persists into later pupal

development (figure S5e). Across stages co-stained with the eyespot marker *spalt*, we

did not detect *araucan* expression obviously upregulated in eyespots (figure S5),

despite peppered expression spanning those regions in the stages sampled after 13 %

pupal development (figure 3c).

Together, these data show that *araucan* transitions from early domain- and vein-

associated patterns to a later dispersed distribution in developing scale cells. This later

expression pattern mirrors the spatially scattered scale iridescence phenotypes in mKO

patches and is consistent with a role for *araucan* in scale-cell differentiation and

morphogenesis occurring at the assayed stage.

DISCUSSION

Region specific control of iridescence by *araucan*

In this study, we investigated the role of the *Iroquois*-complex gene *araucan* in butterfly

wing patterning. Our findings reveal that *araucan* is a key regulator of structural

coloration in *J. coenia*, modulating iridescence in two spatially and functionally distinct

ways: (1) by suppressing blue iridescence across the dorsal wing surface, and (2) by

promoting purple-violet iridescence in forewing eyespot center.

Structural coloration in butterfly wings arises from complex interactions between

incident light and the ultrastructure of wing scales, as reviewed by Thayer and Patel

(35). One well-characterized mechanism is thin-film interference, in which the lower

lamina of the scale functions as a reflective surface, producing a range of iridescent

hues depending on its thickness and the angle of incident light (29,35–37). Previous

work in *Junonia spp*. has shown that the thickness of the lower lamina within individual

cover scales reliably predicts spectral reflectance: thin laminae generate golden-brown

hues as seen in wild type *J. coenia*, whereas thicker laminae produce blue or teal hues

as seen in *optix* knockouts of *J. coenia* or species in the genus that naturally exhibit

blue-green iridescence (3). In our *araucan* knockouts, we observed an increase in lower

lamina thickness in dorsal cover scales, resulting in a shift from brown to blue structural

coloration. This optical shift occurred without detectable changes in scale pigmentation,

or other scales indicating that *araucan* suppresses iridescence in non-eyespot regions

by constraining laminar thickness.

In contrast, the eyespot centers of wild-type *J. coenia* display purple-violet iridescence

that emerges from the combined optical effects of stacked cover and ground scales as

shown in other nymphalid butterflies (29,30,36). The ultrastructure and pigmentation of

the cover scales modulate the amount of incident light reflected or transmitted, while

those of the underlying ground scales influence the extent of light that is scattered back.

Together, these layers generate interference patterns that either enhance or dampen

the resulting color signal. In *araucan* mutants, we observed increased laminar thickness

in both cover and ground scales, accompanied by a marked reduction in ground scale

pigmentation. We infer that the loss of pigment decreases light absorbance, thereby

increasing backscatter, which—when combined with structural modifications in the

cover scales—produces a desaturated, muted tan-grey hue in place of the saturated

purple-violet seen in wild-type eyespots. This enhanced backscatter likely contributes to

the shimmering or hazy appearance of iridescence observed in some mutant individuals

(figure S4). In addition, differences in ridge spacing and crossrib distances may further

influence optical properties, either directly or by influencing pigmentation levels as seen

before (27). Together, these findings demonstrate that *araucan* regulates multiple

dimensions of scale morphology and pigmentation that collectively shape the optical

characteristics of forewing eyespot center coloration.

The opposing effects of *araucan* in eyespot versus non-eyespot scales underscore a

broader principle in butterfly wing development: regulatory genes often function in a

region-specific, context-dependent manner. The divergent roles of *araucan* reinforce the

idea that eyespots represent modular, semi-independent developmental units capable

of evolving specialized scale properties—including iridescence—without altering

patterning in adjacent regions. This spatial modularity enables flexible evolutionary

innovation within the wing, raising important questions about the regulatory architecture

controlling *araucan* expression across the wing.

Gene regulatory dynamics of *araucan*

We were intrigued to observe that *araucan* had a very dynamic pattern of transcription

over developmental time, ranging from vein- and posterior domain-related patterns in

early development, to a whole-wing scattered pattern later in pupal development that is

consistent with iridescence phenotypes. The early posterior domain expression pattern

was of particular interest because it resembles the expression of *mirror*–the sister gene

of *araucan* which has been shown to play a role in posterior domain determination (11).

Curiously, we did not observe any domain-related or otherwise *mirror*-like phenotypes in

our *araucan* mutants and it therefore remains unclear whether this posterior *araucan*

expression pattern has any functional relevance. It is interesting, however, to speculate

whether the parallel expression patterns of *araucan* and *mirror* may reflect ancestral

regulatory architecture that predates the divergence of these genes. In any case, the

complex and dynamic expression patterns of *araucan* over developmental time suggest

this gene is likely responding to a varied progression of upstream trans-regulatory

signals.

Our CRISPR mKOs and expression data raise the question of where *araucan* sits within

the gene regulatory networks that pattern the wing and specify scale-cell fates. One of

the two major mKO classes we observe was a blue shift in the iridescence of dorsal

wing scales, consistent with increased lower-lamina thickness in cover scales. A very

similar effect was previously reported for *optix* mKOs in *J. coenia*. Indeed, the blue shift

effect observed in *araucan* knockouts could reasonably be described as a subset of

optix knockout effects. Although we did not detect marked co-expression of *optix* and

araucan in the same cells at the timepoints we sampled (figure S6), this does not rule

out participation in a shared pathway: transient or low-abundance transcripts,

development staging, or non-cell-autonomous interactions could all evade overlap in our

datasets. We therefore propose two working models that explain the similar effects of

these genes: (1) a shared pathway in which *optix* and *araucan* act in series or converge

on common ultrastructural effectors to modulate lower-lamina thickness (yielding similar

blue-shift phenotypes), and (2) parallel pathways that independently alter scale

ultrastructure and pigment deposition but produce convergent phenotypes. Future work

should assess to what degree *optix* and *araucan* may work together to determine scale

cell structural coloration.

In addition to structural blue shifting of dorsal wing scales, *araucan* had a different, yet

highly specific effect on the structural and pigmentary coloration of eyespot centers.

How might *araucan* be deployed for such a different, spatially restricted eyespot

function? In *J. coenia*, mKO of *spalt* eliminates all eyespots (38), consistent with *spalt*

acting as a master regulator of eyespot formation. Because the *araucan* phenotype

tracks *spalt* expression patterns, one plausible model is that *spalt* functions upstream of

araucan in eyespot centers. The absence of detectable *spalt–araucan* co-expression in

our HCR in situs does not rule this out: *araucan* eyespot transcription may be below our

detection threshold or expressed at later stages of wing development, which makes

sense for a gene that regulates terminal stages of scale maturation. A similar detection–

phenotype mismatch is familiar from other wing transcription factors (e.g., *optix*), where

in situ hybridization patterns from earlier stages of development (figure S6) contrast with

broader phenotypic effects of knockouts. This all suggests that gene expression

patterns during later stages of development, which are challenging to image for

technical reasons, are important for structural and pigmentary color determination (2).

The forewing-specific *araucan* eyespot phenotype also lead us to speculate that

Ultrabithorax, a selector gene for hindwing identity (39–42), may play an upstream role

in regulating *araucan* in this context.

Conclusion

Overall, our results identify *araucan* as a region-specific regulator of structural coloration

in *J. coenia*, suppressing blue iridescence across the dorsal wing while promoting

saturated purple–violet at the forewing eyespot focus. These effects map to coordinated

changes in scale ultrastructure—especially lower-lamina thickness—and, in eyespots,

to coupled shifts in optical layering with ground-scale pigmentation. Together, the data

reveal how a single transcription factor can differentially tune photonic architecture

across the wing, and reinforce eyespots as modular, context-dependent developmental

units. While the precise placement of *araucan* within color pattern gene regulatory

networks remains unresolved, our findings establish a framework for dissecting the

regulatory logic that links gene activity to the coordinated evolution of cell structure and

pigment synthesis.

Data availability

The raw SEM images used for ultrastructural measurements, associated data tables,

raw spectral data from scale micro-spectrophotometry, genome assembly and

annotation files, are available through Dryad (DOI: DOI: 10.5061/dryad.rfj6q57n6).

Conflict of Interest

The authors declare no conflict of interest.

REFERENCES

1. Finet C. Developmental genetics of cuticular micro- and nano-structures in insects. Current Opinion in Insect Science. 2024 Oct 1;65:101254.

2. Zhang L, Mazo-Vargas A, Reed RD. Single master regulatory gene coordinates the evolution and development of butterfly color and iridescence. Proc Natl Acad Sci U S A. 2017 Oct 3;114(40):10707–12.

3. Thayer RC, Allen FI, Patel NH. Structural color in Junonia butterflies evolves by tuning scale lamina thickness. Tautz D, Desplan C, Jiggins C, editors. eLife. 2020 Apr 7;9:e52187.

4. Livraghi L, Hanly JJ, Loh LS, Henry A, Keck CMT, Shirey VM, et al. Genetic basis of an adaptive polymorphism controlling butterfly silver iridescence. Current Biology 2025 May 5;35(9):2154-63.

5. Ficarrotta V, Hanly JJ, Loh LS, Francescutti CM, Ren A, Tunström K, et al. A genetic switch for male UV iridescence in an incipient species pair of sulphur butterflies. Proc Natl Acad Sci U S A. 2022 Jan 18;119(3):e2109255118.

6. Rodriguez-Caro F, Fenner J, Bhardwaj S, Cole J, Benson C, Colombara AM, et al. Novel Doublesex Duplication Associated with Sexually Dimorphic Development of Dogface Butterfly Wings. Molecular Biology and Evolution. 2021 Nov 1;38(11):5021–33.

7. Prakash A, Finet C, Banerjee TD, Saranathan V, Monteiro A. Antennapedia and optix regulate metallic silver wing scale development and cell shape in Bicyclus anynana butterflies. Cell Reports. 2022 Jul 5;40(1).

8. Prakash A, Dion E, Banerjee TD, Monteiro A. The molecular basis of scale development highlighted by a single-cell atlas of Bicyclus anynana butterfly pupal forewings. Cell Reports. 2024 May 28;43(5).

9. Kerner P, Ikmi A, Coen D, Vervoort M. Evolutionary history of the iroquois/Irx genes in metazoans. BMC Evolutionary Biology. 2009 Apr 15;9(1):74.

10. Setton EVW, Ballesteros JA, Blaszczyk PO, Klementz BC, Sharma PP. A taxon-restricted duplicate of Iroquois3 is required for patterning the spider waist. PLOS Biology. 2024 Aug 29;22(8):e3002771.

11. Chatterjee M, Yu XY, Brady NK, Hatto GC, Amendola CA, Reed RD. mirror determines the far posterior domain in butterfly wings. eLife. 2025 Jun 25;13:RP96904.

12. Gómez-Skarmeta JL, Corral RD del, Calle-Mustienes E de la, Ferrés-Marcó D, Modolell J. araucan and caupolican, Two Members of the Novel Iroquois Complex,

Encode Homeoproteins That Control Proneural and Vein-Forming Genes. Cell. 1996
Apr 5;85(1):95–105.

13. Gómez-Skarmeta JL, Modolell J. araucan and caupolican provide a link between
compartment subdivisions and patterning of sensory organs and veins in the
Drosophila wing. Genes Dev. 1996 Nov 15;10(22):2935–45.

14. Kehl BT, Cho KO, Choi KW. mirror, a Drosophila homeobox gene in the iroquois
complex, is required for sensory organ and alula formation. Development. 1998 Apr
1;125(7):1217–27.

15. Cavodeassi F, Modolell J, Gómez-Skarmeta JL. The Iroquois family of genes: from
body building to neural patterning. Development. 2001 Aug 1;128(15):2847–55.

16. Ikmi A, Netter S, Coen D. Prepatterning the Drosophila notum: The three genes of
the iroquois complex play intrinsically distinct roles. Developmental Biology. 2008
May 15;317(2):634–48.

17. Hines HM, Papa R, Ruiz M, Papanicolaou A, Wang C, Nijhout HF, et al.
Transcriptome analysis reveals novel patterning and pigmentation genes underlying
Heliconius butterfly wing pattern variation. BMC Genomics. 2012 Jun 29;13(1):288.

18. Fandino RA, Brady NK, Chatterjee M, McDonald JMC, Livraghi L, van der Burg
KRL, et al. The ivory lncRNA regulates seasonal color patterns in buckeye
butterflies. Proceedings of the National Academy of Sciences. 2024 Oct
8;121(41):e2403426121.

19. Zhang L, Reed RD. A Practical Guide to CRISPR/Cas9 Genome Editing in
Lepidoptera. In: Sekimura T, Nijhout HF, editors. Diversity and Evolution of Butterfly
Wing Patterns: An Integrative Approach. 2017 Aug 30 (pp. 155-172). Singapore:
Springer Singapore.

20. Maia R, Gruson H, Endler JA, White TE. 2: New tools for the spectral and spatial
analysis of colour in r. Methods in Ecology and Evolution. 2019;10(7):1097–107.

21. Ren A, Day CR, Hanly JJ, Counterman BA, Morehouse NI, Martin A. Convergent
Evolution of Broadband Reflectors Underlies Metallic Coloration in Butterflies.
Frontiers in Ecology and Evolution. 2020 Jun 30;8:206.

22. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.
Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012
Jul;9(7):676–82.

23. Villinger C, Gregorius H, Kranz C, Höhn K, Münzberg C, von Wichert G, et al.
FIB/SEM tomography with TEM-like resolution for 3D imaging of high-pressure
frozen cells. Histochem Cell Biol. 2012 Oct;138(4):549–56.

24. Pinheiro JC, Bates DM. Mixed-effects models in S and S-PLUS. New York, NY:
Springer New York; 2000 May 5.

25. Matsuoka Y, Monteiro A. Melanin Pathway Genes Regulate Color and Morphology
of Butterfly Wing Scales. Cell Reports. 2018 Jul 3;24(1):56–65.

26. Finet C, Ruan Q, Bei YY, You En Chan J, Saranathan V, Yang JKW, et al. Multi-
scale dissection of wing transparency in the clearwing butterfly Phanus vitreus.
Journal of The Royal Society Interface. 2023 May 31;20(202):20230135.

27. Banerjee TD, Finet C, Seah KS, Monteiro A. Optix regulates nanomorphology of
butterfly scales primarily via its effects on pigmentation. Frontiers in Ecology and
Evolution. 2024 May 1;12:1392050.

28. Jessop AL, Pirih P, Wang L, Patel NH, Clode PL, Schröder-Turk GE, et al.
Elucidating nanostructural organization and photonic properties of butterfly wing
scales using hyperspectral microscopy. Journal of The Royal Society Interface.
2024 Sep 11;21(218):20240185.

29. Giraldo MA, Yoshioka S, Liu C, Stavenga DG. Coloration mechanisms and
phylogeny of Morpho butterflies. Journal of Experimental Biology. 2016 Dec
15;219(24):3936–44.

30. Siddique RH, Vignolini S, Bartels C, Wacker I, Hölscher H. Colour formation on the
wings of the butterfly Hypolimnas salmacis by scale stacking. Sci Rep. 2016 Nov
2;6(1):36204.

31. Reed RD, Chen PH, Frederik Nijhout H. Cryptic variation in butterfly eyespot
development: the importance of sample size in gene expression studies. Evolution &
Development. 2007;9(1):2–9.

32. Oliver JC, Tong XL, Gall LF, Piel WH, Monteiro A. A Single Origin for Nymphalid
Butterfly Eyespots Followed by Widespread Loss of Associated Gene Expression.
PLOS Genetics. 2012 Aug 16;8(8):e1002893.

33. Reed RD, Selegue JE, Zhang L, Brunetti CR. Transcription factors underlying wing
margin color patterns and pupal cuticle markings in butterflies. EvoDevo. 2020 May
27;11(1):10.

34. Dinwiddie A, Null R, Pizzano M, Chuong L, Leigh Krup A, Ee Tan H, et al. Dynamics
of F-actin prefigure the structure of butterfly wing scales. Developmental Biology.
2014 Aug 15;392(2):404–18.

35. Thayer RC, Patel NH. A meta-analysis of butterfly structural colors: their color range,
distribution and biological production. Journal of Experimental Biology. 2023 Nov
8;226(21):jeb245940.

36. Stavenga DG, Leertouwer HL, Wilts BD. Coloration principles of nymphaline
butterflies – thin films, melanin, ommochromes and wing scale stacking. Journal of
Experimental Biology. 2014 Jun 15;217(12):2171–80.

37. Wasik BR, Liew SF, Lilien DA, Dinwiddie AJ, Noh H, Cao H, et al. Artificial selection
for structural color on butterfly wings and comparison with natural evolution.
Proceedings of the National Academy of Sciences. 2014 Aug 19;111(33):12109–14.

38. Zhang L, Reed RD. Genome editing in butterflies reveals that spalt promotes and
Distal-less represses eyespot colour patterns. Nat Commun. 2016 Jun 15;7:11769.

39. Weatherbee SD, Frederik Nijhout H, Grunert LW, Halder G, Galant R, Selegue J, et
al. Ultrabithorax function in butterfly wings and the evolution of insect wing patterns.
Current Biology. 1999 Feb 11;9(3):109–15.

40. Tendolkar A, Pomerantz AF, Heryanto C, Shirk PD, Patel NH, Martin A.
Ultrabithorax is a micromanager of hindwing identity in butterflies and moths.
Frontiers in Ecology and Evolution. 2021 Mar 18;9:643661.

41. Tong X, Hrycaj S, Podlaha O, Popadic A, Monteiro A. Over-expression of
Ultrabithorax alters embryonic body plan and wing patterns in the butterfly Bicyclus
anynana. Developmental Biology. 2014 Oct 15;394(2):357–66.

42. Tendolkar A, Mazo-Vargas A, Livraghi L, Hanly JJ, Van Horne KC, Gilbert LE, et al.
Cis-regulatory modes of Ultrabithorax inactivation in butterfly forewings. Chipman
AD, Desplan C, editors. eLife. 2024 Jan 23;12:RP90846.

Table 1. Comparison of wild-type and *araucan* mutant non-eyespot cover scale

ultrastructural features.

Scale type	Factor	Comparison	Estimate	Std error	Z-value	Pr(>|z|)	95% CI
cover	Lower-lamina thickness	WT - mutant == 0	-55.41	2.93	-18.89	<2e-16 ***	[-61.16 ; -49.66]
cover	Distance ridge-ridge	WT - mutant == 0	0.54	0.54	10.13	0.05	[-19.32 ; 20.39]
cover	Distance crossrib-crossrib	WT - mutant == 0	8.61	14.25	0.6	0.54	[-19.31 ; 36.54]

640   **Table 2.** Comparison of wild-type and *araucan* mutant cover and ground scale

641   ultrastructural features collected from eyespot center.

Scale type	Factor	Comparison	Estimate	Std error	Z-value	Pr(>|z|)	95% CI
cover	Lower-lamina thickness	WT - mutant == 0	38.92	2.61	14.89	<2e-16 ***	[33.79 ; 44.04]
cover	Distance ridge-ridge	WT - mutant == 0	317.75	6.83	46.51	<2e-16 ***	[304.36 ; 331.14]
cover	Distance crossrib-crossrib	WT - mutant == 0	-256.73	24.15	-10.63	<2e-16 ***	[-304.06 ; -209.40]
ground	Lower-lamina thickness	WT - mutant == 0	38.46	2.45	15.72	<2e-16 ***	[33.67 ; 43.25]
ground	Distance ridge-ridge	WT - mutant == 0	282.4	6.87	41.08	<2e-16 ***	[268.93 ; 295.88]
ground	Distance crossrib-crossrib	WT - mutant == 0	-32.52	7.49	-4.34	1.42e-05 ***	[-47.20 ; -17.83]

642
643

 FIGURES

646
Figure 1. *araucan* knockouts cause blue iridescence in dorsal hindwings by increasing

lower lamina thickness of cover scales. (a) *araucan* mKO produces patches of blue

iridescent scales in the dorsal hindwing. *Left*: Wild-type wing showing uniform golden-

brown color. Adjacent are magnified views of individual cover scales from the same

wing, with the abaxial surface (*top*) and adaxial surface (*bottom*) shown. Right: *araucan*

mKO hindwing showing a mosaic patch of blue iridescent scales (*dashed line*). Adjacent

are magnified views of mutant cover scales, revealing blue iridesence on both the

abaxial (*top*) and adaxial (*bottom*) surfaces. Scale bar = 50 μm. Additional images of

contralateral wings showing mutant blue iridescent scales are in figures S2 and S3. (b)

Reflectance and absorbance spectra of cover scales confirm that the observed color

shift is structural, not pigment-based. Solid lines indicate mean spectra; shaded regions

represent standard deviation; asterisks mark wavelengths with significant differences

between groups ($n_{reflectance}$ = 3; $n_{absorbance}$ = 5; **p-adj* < 0.01; table S2). (c) *Left:* SEM

images of transverse sections from isolated cover scales, prepared by cryo-fracturing,

show that wild-type scales (*top*) have a thinner lower lamina (LL; *grey shaded region*)

compared to mutant scales (*bottom*). *Right*: Violin-box plot quantifies lower lamina

thickness from FIB-milled isolated cover scales (n = 5; ***$p < 2 \times 10^{-16}$; tables 1, S1). (d)

Additional cover scale measurements: ridge-to-ridge spacing (d_{R-R}, *top left*), distance

between crossribs (d_{CR-CR}, *top right*), and area of the windows (*teal*) - through which

incident light goes into the scale (*bottom*, ns – not significant; tables 1, S1).

668
669
670

Figure 2. *araucan* knockouts diminish purple-violet iridesence by altering cover scale

reflectance, ground scale pigmentation, and the ultrastructure of both scale types in

forewing eyespot centers. (a) On both dorsal (*left*) and ventral (*middle*) surfaces,

forewing eyespot centers of wild-type individuals (*top*) exhibit distinct purple-violet

iridescence, whereas *araucan* knockouts (*bottom*) show muted grey-brown scales

(figure S4). Magnified images of the eyespot center (*right*) highlight the stacking

arrangement of cover (c.s; *white border*) and ground scales (g.s; *yellow arrows where*

exposed and yellow dashed boundaries where overlaid by cover scales). Overlaid

ground scales highlight saturation of overall color (*yellow dashed boundary*). Abwing

surfaces of isolated cover and ground scales (*far right*) of wild-type (*top*) and mutant

(*bottom*) scales. (b) Spectrophotometry of isolated cover scales and (c) ground scales

from eyespot centers in wild-type and mutant forewings. Reflectance spectra (*left*) and

absorbance spectra (*right*). Solid lines represent mean spectra; shaded areas indicate

standard deviation; asterisks denote wavelengths with significant differences between

groups ($n_{reflectance}$ = 3; $n_{absorbance}$ = 5; ***p-adj < 0.001, **p-adj < 0.01; Table S4). (d–e)

Violin-box plots quantifying ultrastructural features in cover scales (d) and ground scales

(e) (n = 5; ***$p < 2 \times 10^{-16}$; Tables 2, S3).

689
690

691
Figure 3. Spatiotemporal dynamics of *araucan* transcripts in late larval and early

pupal wings. (a–b) In the 5th-instar larva (a) and ~13% pupal development (b),

araucan signal (yellow) is enriched in the far-posterior vannus domain, bounded

anteriorly by the 2A vein in both wings. Signal is also detected in cells along pre-vein

tracheae and at the wing margin (white arrows). (c) By ~25% pupal development,

araucan persists along the 2A vein and appears in scattered cells across the wing blade

in a pattern-nonspecific manner (yellow dots). In the forewing (*top*), this includes the

eyespot focus (*inset*). In the hindwing, this recaptures the mosaic clones seen in the

mKOs (*inset*). [Scale bar = 50 μm]

SUPPLEMENTARY INFORMATION

Figure S1. (a) Map of the *araucan* locus in *J. coenia,* annotated with the location of sgRNAs. The primers shown were used to validate deletions in mutants. Refer to Materials and Methods for the sequences of sgRNAs and genotyping primers. (b) Maximum likelihood phylogeny of *J. coenia*, *Heliconius erato lativitta*, *Tribolium castaneum*, *Apis mellifera*, and *D. melanogaster Iroquois* complex genes show that *JC_g12213* is the *araucan*/*caupolican* ortholog in *J. coenia* and *JC_g12219* is the ortholog of *mirror*. Re-analysed already published data (Chatterjee et al. 2025), using new gene model identifiers. (c) CRISPR-Cas9 targeting of *araucan* produces deletions at the sgRNA sites. Wild type and mutant sequences are shown. Green boxes indicate sgRNA binding sites. Purple boxes denote PAM sites Knockouts mKO "A" and mKO "3" show blue dorsal iridescence phenotypes (Figure S2). mKO M has eyespot focus phenotype (Figure S2).

725 **Figure S2.** *araucan* mKOs displaying mutant patches (dashed lines) of iridescent blue

726 scales (arrows) in hindwings. Contrasting lateral wings from individual butterflies are

727 shown to highlight the asymmetrical nature of the mosaic phenotypes.

Figure S3. *araucan* mKOs displaying mutant patches (dashed lines) of iridescent blue

scales (arrows) in a subset of mutant forewings. Contrasting lateral wings from

individual butterflies are shown to highlight the asymmetrical nature of the mosaic

phenotypes.

Figure S4. Loss of purple-violet iridescence in forewing eyespot foci following *araucan*
mosaic knockout (mKO). (a) Contralateral forewings from representative individuals
showing the elimination of purple-violet iridescence in dorsal (*top*) and ventral (*bottom*)
eyespot foci. In most cases, both forewings exhibited the mutant phenotype, with
exceptions noted in specimens "mKO D" and "mKO L." Blue arrows indicate wild-type
foci with saturated iridescent coloration; white arrows indicate mutant foci displaying
altered or absent iridescence. (b) Enlarged views of eyespot foci highlighting the
phenotypic effects of *araucan* mKOs. Mutant scales show a range of altered hues, from
muted grey to shimmering pink-purple, compared to the vivid purple-violet observed in
wild type. Insets show scale-level differences in the dorsal (*top*) and ventral (*bottom*)
surfaces.
783
784
785

Figure S5. **Comparative expression of** *araucan* **and** *spalt* **during wing**
development. Each panel shows three columns: left, *araucan* mRNA (yellow); middle,
spalt mRNA (magenta); right, merge. (a–b) 5th-instar larval wings: *araucan* is enriched
in the posterior domain and in discrete cells along pre-vein tracheae (white arrows).
spalt marks future eyespot domains (red arrows). No overlap between *araucan* and
spalt is detected at this stage. (c) ~13% pupal wings: the larval pattern persists—
araucan expression remains to the posterior wing and large cells along the veins (white
arrows), while *spalt* marks eyespot fields (red arrows) with no detectable co-expression
in the eyespot focus (inset). (d) ~25% pupal wings: *spalt* is broadly expressed across
scale-forming cells, whereas *araucan* is retained along the posterior/2A domain and
scattered cells across the wing; with no distinct difference in the focus (inset). (e) ~45%
pupal development: Only hindwing showing expression patterns of *araucan* and *spalt*
similar to earlier stage with no overlap. White circles highlight large, interspersed cells
along the Cu2 vein. [scale bar = 50 µm]

Figure S6. **Optix mRNA expression in early to mid-pupal development (PD)**

captures only a subset of large CRISPR knockout effects in adults. Strong *optix*

expression in the wing conjugation scales across development (white arrowhead).

Pattern specific expression in the eyespots and parafocal elements (magenta arrow)

peaks at 16% pupal development and gradually lowers in 25% and 43% development.

Optix mKO phenotypes in *J. coenia* adults results in large effects across the wing (2),

despite early pupal spatial expression. *Optix* mKO CRISPR images taken from Zhang

and Mazo-Vargas et. al (2).

854 **TABLES**

856 **Table S1.** Comparison of wild-type and *araucan* mutant non-eyespot cover scale

857 ultrastructural features.

scale type	factor	numDF	denDF	F-value	P-value
cover	lower lamina thickness	1	94	356.72	<.0001
cover	distance ridge-ridge	1	244	0.003	0.96
cover	distance crossrib-crossrib	1	494	0.37	0.55

862   **Table S2.** Significance table of reflectance and absorbance spectra measured at

863   different wavelengths.

Spectral type	Region -scale type	Wavelengths(nm)	p_value	p_adjusted	Significance
Absorbance	Non-eyespot-cover	350	0.0422	0.0562	ns
Absorbance	Non-eyespot-cover	400	0.5401	0.5401	ns
Absorbance	Non-eyespot-cover	450	0.0897	0.1026	ns
Absorbance	Non-eyespot-cover	500	0.02717234	0.043475744	*
Absorbance	Non-eyespot-cover	550	0.0137932 24	0.02758644 9	*
Absorbance	Non-eyespot-cover	600	0.0010344 15	0.00827531 7	**
Absorbance	Non-eyespot-cover	650	0.0033162 06	0.01287740 1	*
Absorbance	Non-eyespot-cover	700	0.0048290 25	0.01287740 1	*
Reflectance	Non-eyespot-cover	350	0.058198	0.066512	ns
Reflectance	Non-eyespot-cover	400	0.000189	0.001511	**
Reflectance	Non-eyespot-cover	450	0.012675	0.020281	*
Reflectance	Non-eyespot-cover	500	0.798358	0.798358	ns

Reflectance	Non-eyespot-cover	550	0.032901	0.043868	*
Reflectance	Non-eyespot-cover	600	0.002676	0.005351	**
Reflectance	Non-eyespot-cover	650	0.002317	0.005351	**
Reflectance	Non-eyespot-cover	700	0.002275	0.005351	**

869 **Table S3.** Comparison of wild-type and *araucan* mutant cover and ground scale

870 ultrastructural features collected from eyespot center.

scale type	factor	numDF	denDF	F-value	P-value
cover	lower lamina thickness	1	94	221.66	<.0001
cover	distance ridge-ridge	1	244	2163.07	<.0001
cover	distance crossrib-crossrib	1	494	113.02	<.0001
ground	lower lamina thickness	1	98	247.29	<.0001
ground	distance ridge-ridge	1	244	1687.48	<.0001
ground	distance crossrib-crossrib	1	494	18.84	<.0001

874 **Table S4.** Significance table of reflectance and absorbance spectra measured at

875 different wavelengths.

Spectral type	Region - scale type	Wavelengths(nm)	p_value	p_adjusted	Significance
Reflectance	Eyespot-ground	350	0.42529807 6	0.4252981	ns
Reflectance	Eyespot-ground	400	0.1537366 35	0.175699	ns
Reflectance	Eyespot-ground	450	0.0805972 06	0.1289555	ns
Reflectance	Eyespot-ground	500	0.0359267 62	0.0958047	ns
Reflectance	Eyespot-ground	550	0.0985159 12	0.1313545	ns
Reflectance	Eyespot-ground	600	0.0245332 69	0.0958047	ns
Reflectance	Eyespot-ground	650	0.0028847 28	0.0230778	*
Reflectance	Eyespot-ground	700	0.0540729 1	0.1081458	ns
Reflectance	Eyespot-cover	350	0.2271713 85	0.2596244	ns
Reflectance	Eyespot-cover	400	0.0155021 89	0.0206696	*
Reflectance	Eyespot-cover	450	0.0082181 4	0.013149	*
Reflectance	Eyespot-cover	500	0.6769728 61	0.6769729	ns
Reflectance	Eyespot-cover	550	0.0002504 99	0.0006858	***

Reflectance	Eyespot-cover	600	0.00286324 3	0.005726 5	**
Reflectance	Eyespot-cover	650	0.0002571 88	0.000685 8	***
Reflectance	Eyespot-cover	700	8.9404E-05	0.000685 8	***
Absorbance	Eyespot-ground	350	0.3086519 17	0.308651 9	ns
Absorbance	Eyespot-ground	400	8.29698E-05	0.000167 3	***
Absorbance	Eyespot-ground	450	0.0001068 84	0.000171	***
Absorbance	Eyespot-ground	500	0.0001584 22	0.000181 1	***
Absorbance	Eyespot-ground	550	8.3648E-05	0.000167 3	***
Absorbance	Eyespot-ground	600	5.60723E-05	0.000167 3	***
Absorbance	Eyespot-ground	650	0.0001479 18	0.000181 1	***
Absorbance	Eyespot-ground	700	1.30136E-05	0.000104 1	***

FILES

File S1. mRNA and amino acid sequences of genes used in this study, along with their

D. melanogaster orthologs. These sequences were used to construct the gene family

phylogeny shown in figure S1b and to design probes for HCR *insitu* hybridization.

File S2. *Sequences of araucan, spalt* and *optix HCR-probes used for in situ*
hybridization.