1

dFLASH; dual FLuorescent transcription factor Activity Sensor for Histone integrated live-cell reporting and high-content screening

Authors: Timothy P. Allen[1], Alison E. Roennfeldt[1], Moganalaxmi Reckdharajkumar[1], Miaomiao Liu[3], Ronald J. Quinn[3#], Darryl L. Russell[2#], Daniel J. Peet[1#], Murray L. Whitelaw[1,4#] & David C. Bersten[1,2*]

Affiliations: 1. School of Biological Sciences, University of Adelaide, Adelaide, South Australia, Australia 2. Robinson Research Institute, University of Adelaide, South Australia, Australia. 3. Griffith Institute for Drug Discovery, Griffith University, Brisbane, Australia 4. ASEAN Microbiome Nutrition Centre, National Neuroscience Institute, Singapore 169857, Singapore. *Corresponding author. #labs that contributed to the work

Abstract

Live-cell reporting of regulated transcription factor (TF) activity has a wide variety of applications in synthetic biology, drug discovery, and functional genomics. As a result, there is high value in the generation of versatile, sensitive, robust systems that can function across a range of cell types and be adapted toward diverse TF classes. Here we present the dual FLuorescent transcription factor Activity Sensor for Histone integrated live-cell reporting (dFLASH), a modular sensor for TF activity that can be readily integrated into cellular genomes. We demonstrate readily modified dFLASH platforms that homogenously, robustly, and specifically sense regulation of endogenous Hypoxia Inducible Factor (HIF) and Progesterone receptor (PGR) activities, as well as regulated coactivator recruitment to a synthetic DNA-Binding Domain- Activator Domain fusion proteins. The dual-colour nuclear fluorescence produced normalised dynamic live-cell TF activity sensing with facile generation of high-content screening lines with strong signal:noise ratios and reproducible screening capabilities (Z' = 0.68-0.74). Finally, we demonstrate the utility of this platform for functional genomics applications by using CRISPRoff to modulate the HIF regulatory pathway, and utility for drug screening by using high content imaging in a bimodal design to isolate activators and inhibitors of the HIF pathway from a ~1600 natural product library.

35
36

Introduction

Cells integrate biochemical signals in a variety of ways to mediate effector function and alter gene expression. Transcription factors (TF) sit at the heart of cell signalling and gene regulatory networks, linking environment to genetic output[1,2]. TF importance is well illustrated by the consequences of their dysregulation within disease, particularly cancer where TFs drive pathogenic genetic programs[3-5]. As a result, there is widespread utility in methods to manipulate and track TF activity in basic biology and medical research, predominantly using TF responsive reporters. Recent examples include enhancer activity screening[6] by massively parallel reporter assay, discovery and characterisation of transcription effector domains [7,8] and CRISPR-based functional genomic screens that use reporter gene readouts to understand transcriptional regulatory networks[2,9]. Beyond the use in discovery biology TF reporters are increasingly utilised as sensors and actuators in engineered synthetic biology applications such as diagnostics and cellular therapeutics. For example, synthetic circuits that utilise either endogenous or synthetic TF responses have exploited to engineer cellular biotherapeutics[10]. In particular, the synthetic Notch receptor (SynNotch) in which programable extracellular binding elicits synthetic TF signalling has been exploits to enhance tumour-specific activation of CAR-T cells, overcome cancer immune suppression, or provide precise tumour target specificity [11-14].

Fluorescent reporter systems are now commonplace in many studies linking cell signalling to TF function and are particularly useful to study single cell features of gene expression, such as stochastics and heterogeneity[15], or situations where temporal recordings are required. In addition, pooled CRISPR/Cas9 functional genomics screens rely on the ability to select distinct cell pools from a homogenous reporting parent population. Screens to select functional gene regulatory elements or interrogate chromatin context in gene activation also require robust reporting in polyclonal pools[16]. Many of the current genetically encoded reporter approaches, by nature of their design, are constrained to particular reporting methods or applications [9,17]. For example, high content arrayed platforms are often incompatible with flow cytometry readouts and vice versa. As such there is a need to generate modular, broadly applicable platforms for robust homogenous reporting of transcription factor and molecular signalling pathways.[2].

Here we address this by generating a versatile, high-performance sensor of signal regulated TFs. We developed a reporter platform, termed the dual FLuorescent TF Activity Sensor for Histone integrated live-cell reporting (dFLASH), that enables lentiviral mediated genomic integration of a TF responsive reporter coupled with an internal control. The well-defined hypoxic and steroid receptor signalling pathways were targeted to demonstrate that the composition of the modular dFLASH cassette is critical to robust enhancer-driven reporting. dFLASH acts as a dynamic sensor of targeted endogenous pathways as well as synthetic TF chimeras in polyclonal pools by temporal high-content imaging and flow cytometry. Routine isolation of homogenously responding reporter lines enabled robust high content image-based screening (Z' = 0.68-0.74) for signal regulation of endogenous and synthetic TFs, as well as demonstrating utility for functional genomic investigations with CRISPRoff. Array-based temporal high content imaging with a hypoxia response element dFLASH

85 successfully identified novel regulators of the hypoxic response pathway, illustrating
86 the suitability of dFLASH for arrayed drug screening applications. This shows the
87 dFLASH platform allows for intricate interrogation of signalling pathways and
88 illustrates its value for functional gene discovery, evaluation of regulatory elements or
89 investigations into chemical manipulation of TF regulation.
90

Results

Design of versatile dFLASH, a dual fluorescent, live cell sensor of TF activity

To fulfil the need for a modifiable fluorescent sensor cassette that can be integrated into chromatin and enable robust live-cell sensing that is adaptable for any nominated TF, applicable to high content imaging (HCI) and selection of single responding cells from polyclonal pools via image segmentation or flow cytometry (**Figure 1c**) a lentiviral construct with enhancer regulated expression of *Tomato*, followed by independent, constitutive expression of *d2EGFP* as both selectable marker and an internal control was constructed (**Figure 1a, b**). Three nuclear localisation signals (3xNLS) integrated in each fluorescent protein ensured nuclear enrichment to enable single cell identification by nuclear segmentation, with accompanying image-based quantification of normalised reporter outputs using high content image analysis, or single-cell isolation using FACS in a signal dependent or independent manner. The enhancer insertion cassette upstream of the minimal promoter driving Tomato expression is flanked by restriction sites, enabling alternative enhancer cloning (**Figure 1a**). The sensor response to endogenous signal-regulated TF pathways was first assessed by inserting a Hypoxia Inducible Factor (HIF) enhancer pathway. HIF-1 is the master regulator of cellular adaption to low oxygen tension and has various roles in several diseases[18-20]. To mediate its transcriptional program, the HIF-1α subunit heterodimerises with Aryl Hydrocarbon Nuclear Translocator (ARNT), forming an active HIF-1 complex. At normoxia[4], HIF-1a is post-translationally downregulated through the action of prolyl hydroxylase (PHD) enzymes and the Von Hippel Lindau (VHL) ubiquitin ligase complex[21]. Additionally, the C-terminal transactivation domain of HIF-1α undergoes asparaginyl hydroxylation mediated by Factor Inhibiting HIF (FIH), which blocks binding of transcription coactivators CBP/p300[22]. These hydroxylation processes are repressed during low oxygen conditions, enabling rapid accumulation of active HIF-1α. HIF-1α stabilisation at normoxia[4] was artificially triggered by treating cells with the hypoxia mimetic dimethyloxalylglycine (DMOG), which inhibits PHDs and FIH, thereby inducing HIF-1α stabilisation, activity and hypoxic gene expression[23]. The well characterised regulation and disease relevance of HIF-1α made it an ideal TF target for prototype sensor development.

Optimisation of dFLASH sensors

Initially, we tested FLASH constructs with a hypoxia response element (HRE) enhancer that contains repeats of HIF enhancers (RCGTG)[24] from endogenous target genes (HRE-FLASH), controlling expression of either nuclear mono (m) or tandem dimer (td)Tomato observing no DMOG induced Tomato expression in stable HEK293T cell lines (mnucTomato or tdnucTomato, **Supp Figure 1a,b**). Given the HIF response element has been validated previously[24], the response to HIF-1α was optimised by altering the reporter design, all of which utilised the smaller mnucTomato (vs tdnucTomato) to contain transgene size. We hypothesised that transgene silencing, chromosomal site-specific effects or promoter enhancer coupling/interference may result in poor signal induced reporter activity observed in initial construct designs. As such we optimised the downstream promoter, the reporter composition and incorporated a 3xNLS d2EGFP internal control from the constitutive promoter to monitor chromosomal effects and transgene silencing.

139
Dual FLASH (dFLASH) variants incorporated three variations of the downstream
promoter (EF1a, PGK and PGK/CMV) driving *3xNLS EGFP (nucEGFP)* and 2A
peptide linked hygromycin (detailed in **Supp Figure 1c**) in combination with alternate
reporter transgenes that it expressed mnucTomato alone, or mnucTomato-Herpes
Simplex Virus Thymidine Kinase (HSVtK)-2A-Neomycin resistance (Neo). Stable
HEK293T and HepG2 HRE-dFLASH cells lines with these backbones were generated
by lentiviral transduction and hygromycin selection. The reporter efficacy of dFLASH
variant cell lines was subsequently monitored by high content imaging 48 hours after
DMOG induction (**Supp Figure 1d, e)**. The downstream composite PGK/CMV or PGK
promoters, enabled the strong DMOG induced Tomato or Tomato/GFP expression
dramatically outperforming EF1a (**Figure 1b** and **Supp Figure 1d**). The composite
PGK/CMV provided bright, constitutive nucEGFP expression in both HepG2 and
HEK293T cells which was unchanged by DMOG, whereas nucEGFP controlled by the
PGK promoter was modestly increased (~2.5 fold) by DMOG (**Supp Figure 1E**).
Substitution of the mnucTomato with the longer mnucTomato-HSVtK-Neo reporter
had no effect on DMOG induced reporter induction in EF1a containing HRE-dFLASH
cells, still failing to induce tomato expression (**Supp Figure 1f**). Whereas, CMV/PGK
containing dFLASH sensors maintained DMOG induction when either the
mnucTomato or the mnucTomato/HSVtK/Neo reporters were utilised (**Supp Figure
1g, h**). Although, mnucTomato without HSVtK, and Neo produced lower absolute
mnucTomato fluorescence and a smaller percentage of cells responding to DMOG,
albeit with lower background. Taken together these findings indicate that certain
backbone compositions prevented or enabled robust activation of the enhancer driven
cassette, similar to the suppression of an upstream promoter by a downstream,
contiguous promoter previously described[25,26] suggesting that the 3' EF1a promoter
results in poorly functioning multi-cistronic synthetic reporter designs[27]. Consequently,
the PGK/CMV backbone and the mnucTomato/HSVtK/Neo reporter from **Supp Figure
1** was chosen as the optimised reporter design (HRE-dFLASH). To confirm that the
HRE element was conferring HIF specificity a no response element dFLASH construct
in HEK293T cells treated with DMOG produced no change in either mnucTomato or
nucEGFP compared to vehicle-treated populations (**Supp Figure 2a**). This result,
together with the robust induction of Tomato in response to DMOG (**Figure 2D**, **Supp
Figure 1f**, **1h**), confirms HIF enhancer driven reporter activation to respond robustly
to induction of the HIF pathway (subsequently labelled dFLASH-HIF).
174
To validate the high inducibility and nucEGFP independence of dFLASH was not
specific to the HIF pathway, we generated a Gal4 responsive dFLASH construct
(Gal4RE-dFLASH), using Gal4 responsive enhancers[22,28]. HEK293T cells were
transduced with Gal4RE-dFLASH and a dox-inducible expression system to express
synthetic Gal4DBDtransactivation domain fusion protein. To evaluate Gal4RE-
dFLASH we expressed Gal4DBD fused with a compact VPR (miniVPR), a strong
transcriptional activator[29] (**Supp Figure 2b, 3a-c**). We observed ~25% of the
polyclonal population was highly responsive to doxycycline treatment (**Supp Figure
2b**), with a ~14-fold change in Tomato expression relative to nucEGFP by HCI (**Supp
Figure 3c**) demonstrating our optimised dFLASH backbone underpins a versatile
reporting platform.
186

dFLASH senses functionally distinct TF activation pathways

Following the success in utilising dFLASH to respond to synthetic transcription factors and HIF signalling, we explored the broader applicability of this system to sense other TF activation pathways. We chose the Progesterone Receptor (PGR), a member of the 3-Ketosteriod receptor family that includes the Androgen, Glucocorticoid and Mineralocorticoid receptors, as a functionally distinct TF pathway with dose-dependent responsiveness to a progestin steroids to assess the adaptability of dFLASH performance. Keto-steroid receptors act through a well-described mechanism which requires direct ligand binding to initiate homodimerization via their Zinc finger DNA binding domains, followed by binding to palindromic DNA consensus sequences. PGR is the primary target of progesterone (P4, or a structural mimic R5020) and has highly context dependant roles in reproduction depending on tissue type[30,31,32]. We inserted PGR-target gene enhancers sequences containing the canonical NR3C motif (ACANNNTGT[31]) into dFLASH, conferring specificity to the ketosteroid receptor family to generate PRE-dFLASH (**Figure 2b,** see **Methods**).

A chimeric TF system was also established with Gal4DBD fusion proteins to create a synthetic reporter to sense the enzymatic activity of oxygen sensor Factor Inhibiting HIF (FIH). This sensor system termed SynFIH for its ability to synthetically sense FIH activity) contained Gal4DBD-HIFCAD fusion protein expressed in a doxycycline-dependant manner, in cells harbouring stably integrated Gal4RE-dFLASH. FIH blocks HIF transactivation through hydroxylation of a conserved asparagine in the HIF-1α C-terminal transactivation domain (HIFCAD), preventing recruitment of the CBP/p300 co-activator complex[22]. As FIH is a member of the 2-oxoglutarate dioxygenase family, like the PHDs which regulate HIF post-translationally, it is inhibited by DMOG (**Figure 2C**), allowing induction of SynFIH-dFLASH upon joint Dox and DMOG signalling (**Supp Figure 3d,3e**). dFLASH-based sensors for PGR and Gal4DBD-HIFCAD generated in the optimised backbone used for dFLASH-HIF (**Figure 2a-c**). For the PGR sensor we transduced T47D cells with PRE-dFLASH, as these have high endogenous PGR expression, while for the FIH-dependant system we generated HEK293T cells with Gal4RE-dFLASH and the GAL4DBD-HIFCAD system (dFLASH-synFIH).

Stable polyclonal cell populations were treated with their requisite chemical regulators and reporter responses analysed by either flow cytometry or temporal imaging using HCI at 2hr intervals for 38 hours (**Figure 2**). Flow cytometry revealed all three systems contain a population that strongly induced nucTomato and maintained nucEGFP (**Supp Figure 2C-E**). In HEK293T cells, ~20% of dFLASH-synFIH and ~50% of dFLASH-HIF population induced Tomato fluorescence substantially relative to untreated controls (**Figure 2d, Figure 2f**). The ~20% reporter response to inhibition of dFLASH-synFIH (**Supp Figure 2e. Figure 2f**) is comparable with what was observed for GalRE-dFLASH response to Gal4DBD-miniVPR expression after equivalent selection (**Supp Figure 2b**). The PGR reporter in T47D cells showed ~50% of the population substantively induced Tomato (**Figure 2e, Supp Figure 2d)**. The presence of considerable responsive populations for FIH, PGR, and HIF sensors, reflected in the histograms of the EGFP positive cells (**Figure 2d-f**) indicated that isolation of a highly responsive clone or subpopulations can be readily achievable for a range of transcription response types. Importantly, the induction of dFLASH-synFIH

235   by Dox/DMOG co-treatment was ablated and displayed high basal Tomato levels in
236   FIH knockout dFLASH-synFIH cells (**Supp Figure 3e**), indicating that the dFLASH-
237   synFIH specifically senses FIH enzymatic activity.
238
All dFLASH systems showed consistent signal-dependant increases in reporter
activity out to 38 hours by temporal HCI enabling polyclonal populations of dFLASH to
track TF activity (**Figure 2g-i**). PRE-dFLASH was more rapidly responsive to R5020
ligand induction (~6 hours, **Figure 2h**) than dFLASH-HIF and dFLASH-synFIH to
DMOG or Dox/DMOG treatment, respectively (~10 hours, **Figure 2g, i**). Treatment of
PRE-dFLASH with estrogen (E2), which activates the closely related Estrogen
Receptor which binds a distinct consensus DNA site to the PGR, or the hypoxia
pathway mimetic DMOG, failed to produce a response on PRE-dFLASH (**Figure 2h**).
This indicates that the PRE enhancer element is selective for the ketosteroid receptor
family (also see below), and that enhancer composition facilitates pathway specificity.
We also observed a signal-dependant change in EGFP expression by flow cytometry
in the T47D PRE-dFLASH reporter cells (**Supp Figure 2g**) but did not observe a
significant change in EGFP expression for HEK293T or HEPG2 dFLASH-HIF (**Supp
Figure 1C, Supp Figure 2c**) or in HEK293T dFLASH-synFIH cells (**Supp Figure 2h**),
with only a small change with Gal4RE-dFLASH with Gal4DBD-miniVPR (**Supp Figure
2b**). While this change in T47D cells was not detected in the in other cellular contexts
(see below), it highlights that care needs to be taken in confirming the utility of the
constitutive nucEGFP as an internal control in certain scenarios.
257

Monoclonal dFLASH cell lines confer robust screening potential in live cells

The observed heterogenous expression of dFLASH within polyclonal cell pools is
useful in many assay contexts but reduces efficiency in arrayed high content screening
experiments and incompatible with pooled isolation of loss of function regulators.
Therefore, monoclonal lines were derived to increase reliability of induction and
consistency of reporting across the population (**Figure 3**). from the HEK293T and
HepG2 dFLASH-HIF, T47D and BT474 PRE-dFLASH and HEK293T dFLASH-synFIH
cell lines (**Figure 3a, Supp Figure 4**). The isolated mcdFLASH-synFIH and
mcdFLASH-HIF lines also demonstrated constitutive signal insensitive nucEGFP
expression (**Supp Figure 4a,b,i**). While the T47D PRE-mcdFLASH showed a small
increase in nucEGFP in response to R5020, this did not preclude the use in
normalisation of high content imaging experiments (see below).
270

No change in EGFP in BT474 PRE-mcdFLASH cell, indicates that strong
transactivation leading to promoter read through or cell-type specific effects may be at
play. Flow cytometry of monoclonal dFLASH cell lines with their cognate ligand
inducers (DMOG (**Figure 3b**), R5020 (**Figure 3f**) or Dox/DMOG (**Figure 3j**)) revealed
robust homogeneous induction of mnucTomato in all cell lines. Using temporal high
content imaging we also found that that, clonally derived lines displayed similar signal
induced kinetics as the polyclonal reporters although they displayed higher signal to
noise and increased consistency (**Figure 3, Supp Figure 4J**). Using physiologically
relevant concentrations of steroids or steroid analogs (10nM-35nM), the PRE-
mcdFLASH lines selectively respond to R5020 (10nM) and not E2 (35nM), DHT
(10nM), Dexamethasone (Dex, 10nM) or Retinoic acid (RA, 10nM) (**Figure 3g, Supp
Figure 4j**). In addition, dose response curves of R5020 mediated Tomato induction

indicate that PRE-mcdFLASH line responds to R5020 with an EC_{50} ~200pM, in
agreement with orthogonal methods[33]. (**Supp Figure 4f, h**). This suggests that the
PRE-mcdFLASH responds sensitively and selectively to PGR selective agonist
R5020, with the potential for high-content screening for modulators of *PGR* activity.
As such, we term this line mcdFLASH-PGR from herein, for its specific ability to report
on PGR activity at physiological steroid concentrations.
289
The temporal HCI of populations (**Figure 2 and Figure 3**) were imaged every 2hrs
and do not inherently provide single-cell temporal dynamics of transcriptional
responses. Using clonally derived mcdFLASH-PGR or mcdFLASH-HIF lines we also
imaged transcriptional responses to R5020 or DMOG, respectively every 15mins
(**Supp Video 1 and 2**). Demonstrating that high temporal resolution imaging has the
potential to monitor transcriptional dynamics in single cells, facilitated by the dual
fluorescent nature of dFLASH. Taken together this indicated that clonal lines displayed
improved signal to noise and assay consistency, possibly enabling high content
screening experiments.
299
Typically, high-content screening experiments require high in-plate and across plate
consistency, therefore we evaluated mcdFLASH lines (HIF-1α, PGR, FIH) across
multiple plates and replicates. System robustness was quantified with the Z' metric[34]
accounting for fold induction and variability between minimal and maximal dFLASH
outputs. Signal induced nucEGFP normalised mnucTomato fluorescence levels
across replicates from independent plates was highly consistent (Z' 0.68-0.74) and
robust (9.3-11.8 fold) (**Figure 3 d, h, l**)), while the signal induced changes in activity
for mcdFLASH-HIF and mcdFLASH-FIH were driven by increased mnucTomato, with
minimal changes in nucEGFP (**Figures 3e and 3m**). Despite the changes previously
observed in nucEGFP, mcdFLASH-PGR in T47D cells provided an equivalent reporter
response to the other systems, with mnucTomato-driven responses (**Figure 3h, i**) As
a result, monoclonal mcdFLASH cell lines represent excellent high-throughput
screening systems routinely achieving Z' scores > 0.5. Importantly, the induction of
the mcdFLASH lines (HEK293T and HepG2 mcdFLASH-HIF, T47D mcdFLASH-PGR
and HEK293T mcdFLASH-SynFIH) was stable over extended passaging (months),
enabling protracted large screening applications.
316
dFLASH-HIF genetic sensor CRISPR-perturbations of the HIF pathway
The robust signal window and high Z' score of mcdFLASH-HIF cell line, coupled with
facile analysis by flow cytometry and HCI, indicate that the reporter system is
amenable to functional genomic screening. We utilised the recently developed
CRISPRoffv2.1 system[35] to stably repress expression of VHL, which mediates post-
translational downregulation of the HIF-1α pathway [36,37]. We generated stable
mcdFLASH-HIF cells expressing a guide targeting the VHL promoter and
subsequently introduced CRISPRoffv2.1 from either a lentivirus driven by an EF1a or
SFFV promoter (**Figure 4c, d**). Cells were then analysed by flow cytometry 5- or 10-
days post selection to determine if measurable induction of mcdFLASH-HIF reporter
was able to indicate the loss of HIF-1α regulation through VHL knockdown (Supp
Figure 6, **Figure 4a**). As expected, mcdFLASH-HIF/sgVHL cells expressing
CRISPRoffv2.1 from either promoter induced the mcdFLASH-HIF reporter in ~35% by
5 days and the majority of cells (~60%) by 10 days as compared to parental cells.

Demonstration that mcdFLASH-HIF is responsive to CRISPRi/off perturbations of key
regulators of the HIF pathway illustrates the potential for the dFLASH platform to
provide a readout for CRISPR screens at-scale in a larger format including genome-
wide screens.
335
dFLASH facilitates bimodal screening for small molecule discovery
Manipulation of the HIF pathway is an attractive target in several disease states, such
as in chronic anaemia[38] and ischemic disease[39] where its promotion of cell adaption
and survival during limiting oxygen is desired. Conversely, within certain cancer
subtypes[40,41] HIF signalling is detrimental and promotes tumorigenesis. Therapeutic
agents for activation of HIF-α signalling through targeting HIF-α regulators were
initially discovered using *in vitro* assays. However, clinically effective inhibitors of HIF-
1α signalling are yet to be discovered[42]. The biological roles for HIF-1α and closely
related isoform HIF-2α, which share the same canonical control pathway, can be
disparate or opposing in different disease contexts requiring isoform selectivity for
therapeutic intervention[43]. To confirm that HIF-1α is the sole isoform regulating
mcdFLASH-HIF in HEK293T cells[44] tagging endogenous HIF-1α and HIF-2α with a
tandem HA-3xFLAG epitope in HEK293T cells, provided for a direct comparison by
immunoblot[45] and determined HIF-1α is the heavily predominant isoform (**Supp
Figure 5a**). Furthermore, there was no change in DMOG induced mnucTomato
expression in HEK293T mcdFLASH-HIF cells when co-treated for up to 72 hours with
the selective HIF2a inhibitor PT-2385 (**Supp Figure 5b)**, consistent with the minimal
detection of HIF-2α via immunoblot. This confirmed that our HEK293T dFLASH-HIF
cell line specifically reports on HIF-1 activity and not HIF-2, indicating that it maybe
useful for identification of drugs targeting the HIF-1α pathway.
356
dFLASH-HIF facilitates multiple measurements across different treatment regimens
and time points, enabling capture of periodic potentiated and attenuated HIF signalling
during a single experiment. Having validated the robust, consistent nature of
mcdFLASH-HIF, we exploited its temporal responsiveness for small molecule
discovery of activators or inhibitors of HIF-1α signalling in a single, bimodal screening
protocol. To test this bimodal design, we utilised a natural product library of 1595
compounds containing structures that were unlikely to have been screened against
HIF-1α prior. We first evaluated library compounds for ability to activate the reporter
after treatment for 36 hours (**Figure 5a**) or 24 hours (**Figure 5d**). The selection of two
different screening time points was to minimise any potential toxic effects of
compounds at the later time points. Consistency of compound activity between the
two screens was assessed by Pearson correlations (**Supp Figure 7i**, R = 0.79, p <
2.2×10^{-16}). Lead compounds were identified by their ability to increase
mnucTomato/nucEGFP (**Figure 5b, c**) and mnucTomato MFI more than 2SD
compared with vehicle controls, with less than 2SD decrease in nucEGFP (21/1595
compounds (1.3%) each expt; **Supp Figure 7a, e**) and an FDR adjusted P score <0.01
across both screens (3/1595 (0.18%) compounds; **Supp Figure 7b, f**). After imaging
of reporter fluorescence to determine these compounds ability to activate HIF-1α we
then treated the cells with 1mM DMOG and imaged after a further 36-hour (**Figure 5c**)
and 24-hour (**Figure 5f**) period. Again, consistency of compound activity was
assessed by Person correlation (**Supp Figure 7j, F,** R = 0.62, p < 2.2×10^{-16}). Lead
compounds were defined as those exhibiting a decrease in mnucTomato MFI >2SD

from DMOG-treated controls in each screen without changing nucEGFP >2SD relative
to the DMOG-treated controls (26/1595 compounds (1.3%) (36hr treatment) and
13/1595 compounds (<1%) (24hr treatment); **Supp Figure 7c, g**), and decrease in
mnucTomato/nucEGFP >2SD with an FDR adjusted P score < 0.01 (3/1595
compounds (0.18%) across both expt; **Supp Figure 7d, h**).
384
dFLASH identified novel and known compounds that alter HIF TF activity.
We confirmed that several of the identified compounds from the pilot screen impacted
HIF-1α. From the activator screening, we identified RQ200674 (**Figure 6d**) as a
previously unreported HIF-1α stabilising compound. RQ200674 increased reporter
activity 2-fold in repeated assays (**Figure 6d**) and stabilised endogenously tagged
HIF-1α at normoxia in HEK293T cells (**Supp Figure 8b**). Mechanistically, RQ200674
had weak iron chelation activity in an *in vitro* chelation assay (**Figure 6e**), suggesting
it intersects with the HIF-1α pathway by sequestering iron similar to other reported HIF
stabilisers. In the inhibitor compound dataset, Celastarol and Flavokawain B
downregulated the reporter at several concentrations (**Supp figure 9b, c**). Celastarol
is a previously reported HIF-1α inhibitor[46-48] and Flavokawain B is a member of the
chalcone family which has previously exhibited anti-HIF-1α activity[49]. RQ500235 was
identified as a HIF-1 inhibitor by mcdFLASH-HIF screening. Dose dependant inhibition
of mcdFLASH-HIF (**Figure 6a**) correlated with a dose-dependent decrease in protein
expression by immunoblot (**Figure 6C**). We observed significant (p=0.0139)
downregulation of HIF-1α transcript levels (**Figure 6D**) and were unable to rescue
HIF-1α protein loss with proteasomal inhibition (**Supp Figure 9d**), indicating
RQ500235 was decreasing HIF-1α at the RNA level. More broadly however, the
identification of these compounds by mcdFLASH-HIF in the bimodal set up
demonstrates successful application of the dFLASH platform to small molecule
discovery efforts for both gain and loss of TF function.
406

Discussion

We designed and optimised dFLASH to offer a versatile, robust live-cell reporting platform that is applicable across TF families and allows for facile high-throughput applications. We validated dFLASH against three independent signal-responsive TFs, two with endogenous signalling pathways (dFLASH-PRE for Progesterone receptors; dFLASH-HRE for hypoxia induced transcription factors) and a synthetic system for a hybrid protein transcriptional regulator (dFLASH- Gal4RE). Each dFLASH construct produced robustly detected reporter activity by temporal high-content imaging and FACS after signal stimulation for its responsive TF (**Figure 2,3**). The use of previously validated enhancer elements for HIF[24] and synthetic gal4 DNA binding domains[22,28] demonstrated that dFLASH can be adapted toward both endogenous and synthetic pathways displaying highly agonist/activator-specific responses, indicating utility in dissecting and targeting distinct molecular pathways. mcdFLASH lines to distinct pathways produced highly consistent (Z' = 0.68-0.74) signal induced Tomato induction measured by high content imaging suggesting dFLASH is ideally suited to arrayed high-throughput screening (**Figure 3**). In addition, mcdFLASH lines also displayed homogenous signal induced reporter induction by flow cytometry indicating that pooled high content screening would also be possible.

Indeed, reporter systems like dFLASH have been increasingly applied to functional genomic screens which target specific transcriptional pathways[9,50-52]. Proof of principle with CRISPRoff mediated downregulation of the core HIF protein regulator, VHL produced distinct tomato expressing cell pools (**Figure 4**), demonstrating genetic perturbations of endogenous TF signalling pathways. The robust induction of the dFLASH-HIF reporter upon VHL knockdown in the majority of cells indicates that whole genome screening would also be successful[9,17,50,53].

Using the HIF-1α specific reporter line, mcdFLASH-HIF, the application of high-content screening was exemplified. This approach was successful in discovering a novel activator and novel inhibitor of the HIF pathway, as well as previously identified inhibitory compounds. This ratified dFLASH as a reporter platform for arrayed-based screening and demonstrates the utility of the linked nucEGFP control for rapid hit bracketing. The novel inhibitor RQ500235 was shown to downregulate HIF-1α transcript levels, like another HIF-1α inhibitor PX-478[54,55]. As PX-478 has demonstrated anti-cancer activity in several cell lines [55,56] and preserved β-cell function in diabetic models[54], a future similar role may exist for an optimised analogue of RQ500235.

The dFLASH system is characterised by some distinct advantages which may enable more precise dissection of molecular pathways. The ability to control for cell-to-cell fluctuations and to decouple generalised or off-target effects on reporter function may aid the precision necessary for large drug library or genome-wide screening applications[57]. In addition, dFLASH, unlike many other high-throughput platforms can be used to screen genetic or drug perturbations of temporal transcriptional dynamics or as used here at multiple time points. In addition, the results indicate that dFLASH is ideally suited to array-based functional genomics approaches[58] allowing for multiplexing with other phenotypic or molecular outputs[59,60 2,61].

The dFLASH approach has some limitations. The fluorescent nature of dFLASH limits the chemical space by which it can screen due to interference from auto-fluorescent compounds. In addition, we acknowledge that fluorescent proteins require O_2 for their activity and this limits the use mnucTomato as a readout of hypoxia. Also, while the backbone design has been optimised for a robust activation of a variety of transcription response pathways the mechanistic underpinning of this is unclear and could be further improved, providing insights into the sequence and architectural determinants of enhancer activation in chromatin. In addition to the strong effect of the dFLASH downstream promoter on upstream enhancer activity It is clear that either the distance between contiguous promoter/enhancer or the sequence composition of the linker has a functional consequence on enhancer induction.

The incorporation of robust native circuits such as those described here (Hypoxia or Progesterone) has the potential to allow the manipulation or integration of these pathways into synthetic biology circuitry for biotherapeutics. In these cases, it is critical that robust signal to noise, is achieved for these circuits to effectively function in biological systems. Further, the use of a synthetic approach to 'sense' FIH enzymatic activity through the HIF-CAD:P300/CBP interaction opens up the possibility that other enzymatic pathways that lack effective *in vivo* activity assay may also be adapted. We also envisage that dFLASH could be adapted to 2-hybrid based screens as a compliment to other protein-protein interaction approaches.

The ability to temporally track TF regulated reporters in populations and at the single-cell level enable dFLASH to be used to understand dynamics of transcriptional responses as has been used to dissect mechanisms of synthetic transcriptional repression[7,8] or understand notch ligand induced synthetic transcriptional dynamics[62]. For instance [62] utilised synthetic reporter circuits to delineate how diverse notch ligands induce different signalling dynamics between responsive cell populations. The large dynamic range of the dFLASH-PGR and HIF reporter lines in conjunction with the high proportion of cells induced in polyclonal pools (**Figure 2**) also suggests dFLASH as a candidate system for forward activity-based enhancer screening. These approaches have been applied to dissect enhancer activity or disease variants with other similar systems such as lentiviral-compatible Massively Parallel Reporter Assays (LentiMPRA)[63,64]. However, the use of the internal control normalisation provided by dFLASH may be useful in separating chromosomal from enhancer driven effects in forward enhancer screens.

Given dFLASH has robust activity in both pooled and arrayed formats, it offers a flexible platform for these investigations. dFLASH can be used to sense endogenous and synthetic transcription factor activity and represents a versatile, stable, live-cell reporter system of a broad range of applications.

499   **Main Figures**

500
501
Figure 1. Summary of dFLASH LV-REPORT construction, utility, and validation
(**a**) The dFLASH system utilises the LV-REPORT construct has a cis-element multiple
cloning site for enhancer insertion, followed by a minimal promoter that drives a
transcription factor (TF) dependant cassette that encodes three separate expression
markers; a nuclear Tomato fluorophore with a 3x C-terminal nuclear localisation signal
(NLS), Herpes Simplex Virus Thymidine Kinase (HSVtK) for negative selection, and
Neomycin (Neo) for positive selection are separated by a 2A self-cleaving peptide
(2A). This is followed by a downstream promoter that drives an independent cassette
encoding EGFP with a 3x N-terminal NLS, and a Hygromycin selection marker
separated by a 2A peptide. (**b**) This design allows for initial identification of the EGFP
fluorophore in nuclei, independent of signal. Expression of the Tomato fluorophore is
highly upregulated in a signal-dependant manner. Images shown are monoclonal
HEK293T dFLASH-HIF cells. Populations were treated for 48 hours ±DMOG induction
of HIF-1α and imaged by HCI. (**c**) This system can be adapted to a range of different
applications. This includes (clockwise) flow cytometry, arrayed screening in a high
throughput setting with high content imaging, isolation of highly responsive clones or
single cells from a heterogenous population or temporal imaging of pooled or individual
cells over time.
520

523
524
525

Figure 2. dFLASH provides sensitive readouts to three distinct TF pathways
(**a-c**) Three distinct enhancer elements enabling targeting of three different signalling aspects. (**a**) Hypoxic response elements (HRE) provide a read out for HIF-1α induction; (**b**) Progesterone response elements (PRE) derived from progesterone receptor target genes facilitates progestin induction; (**c**) Gal4 response elements (GalRE) enable targeting of synthetic transcription factors to dFLASH such as a GAL4DBD-HIFCAD fusion protein that provides a FIH-dependant reporter response. Flow cytometry histograms of (**d, f**) HEK293T and (**e**) T47D cells were treated for 48 hours with (**d**) 1mM DMOG or 0.1% DMSO (Ctrl), (**e**) 100nM R5020 or Ethanol (Ctrl), (**f**) 1μg/mL Doxycycline (Dox) and DMOG or Dox and 0.1% DMSO (Ctrl) for the FIH sensor prior to flow cytometry for Tomato expression among polyclonal population of reporter cells. (**g-i**) Reporter populations from **d-e** were temporally imaged for 38 hours using HCI directly after treatment with (**g**) 0.5mM DMOG or 0.1% DMSO, (4 replicates) (**h**) 100nM R5020, 35nM E2, 0.5mM DMOG and 0.1% Ethanol (EtOH) (4 replicates), (**i**) 0.1% DMSO, 1mM DMOG, 100ng/mL Dox and DMSO, or Dox and DMOG (4 replicates)

HIF response Pathway
HEK293T dFLARE-HRE

PGR response Pathway
T47D dFLARE-PRE

Synthetic FIH Sensor
HEK293T dFLARE-synFIH

Figure 3. Derivation of robust, screen-ready dFLASH clonal lines

(**a**) Schematic for derivation and assessment of robustness for clonal lines of (**b-e**) HEK239T dFLASH-HIF (mcdFLASH-HIF), (**f-i**) T47D dFLASH-PGR (mcdFLASH-PGR) and (**j- m**) HEK293T dFLASH-synFIH (mcdFLASH-synFIH) were analysed by flow cytometry, temporal HCI over 38 hours and for inter-plate robustness by mock multi-plate high throuput screening with HCI. (**b-e**) mcdFLASH-HIF was (**b**) treated with DMOG for 48 hours and assessed for Tomato induction by flow cytometry relative to vehicle controls with fold change between populations displayed and (**c**) treated with vehicle or 0.5mM DMOG and imaged every 2 hours for 38 hours by HCI (mean ±sem, 8 replicates). (**d-e**) mcdFLASH-HIF was treated for 48 hours with 1mM DMOG or vehicle (6 replicates/plate, n = 10 plates) by HCI in a high throughput screening setting (HTS-HCI) for (**d**) normalised dFLASH expression and (**e**) Tomato MFI alone. (**f – i**) T47D mcdFLASH-PGR was (**f**) assessed after 48 hours of treatment with 100nM R5020 by flow cytometry for Tomato induction and (**g**) treated with 10nM R5020, 35nM E2, 10nM DHT and vehicle then imaged every 2 hours for 38 hours by temporal HCI for normalised dFLASH expression (mean ±sem, 8 replicates). (**h-i**) T47D mcdFLASH-PGR was assessed by HTS-HCI at 48 hours (24 replicates/plate, n = 5 plates) for (**h**) dFLASH normalised expression and (**i**) Tomato MFI alone. (**j**) HEK293T dFLASH-synFIH was assessed, with 200ng/mL and or Dox and 1mM DMOG by flow cytometry for dFLASH Tomato induction (**k**) mcdFLASH-synFIH was treated with 100ng/mL Dox, 1mM DMOG and relevant vehicle controls and assessed for reporter induction by temporal HCI (mean ±sem 4 replicates). (**l-m**) mcdFLASH-synFIH cells were treated with 200ng/mL Dox (grey), 1mM DMOG (red), vehicle (pink) or Dox and DMOG (orange) and assessed by HTS-HCI after 48 hours (24 replicates/plate, n = 3 plates) for (**l**) normalised dFLASH expression or (**m**) Tomato MFI induction between Dox and Dox and DMOG treated populations. Dashed lines represent 3SD from relevant vehicle (+3SD) or requisite ligand treated population (-3SD). Fold change for flow cytometry and HTS-HCI (FC) is displayed. Z' was calculated from all analysed plates by HTS-HCI. Z' for all plates analysed was > 0.5.

582
583
584
Figure 4. Near homogenous activation of mcdFLASH-HIF by CRISPRoff
knockdown of VHL.
(**a**) Clonal (1) mcdFLASH-HIF lines derived post-hygromycin (HygroB) selection were
transduced first with the (2) sgRNA vector targeting *VHL* transcriptional start site,
followed by puromycin selection (Puro). This pool was subsequently transduced by
the (3) CRISPRoffv2.1 virus and selected for with blasticidin (BlastS) prior to flow
cytometry (on day 5 and 10 post Blasticidin selection). (**b**) The (1) dFLASH vector with
the HRE enhancer was transduced as were 2 variants of the CRISPRoffv2.1 vector
with either (**a**) EF1α promoter or (**b**) SFFV promoter driving the dCas9 expression
cassette. (**c, d**) Flow cytometry for dFLASH-HIF induction in response to the
CRISPRoffv2.1 VHL knockdown relative to parental line (Ctrl) with (**c**) EF1a or (**d**)
SFFV expression constructs after 10 days of selection.
597
598

Figure 5. Bimodal small molecule screening of the HIF signalling pathway with dFLASH-HIF identifies positive and negative regulators

(**a, d**) HEK293T mcdFLASH-HIF cells were treated with a 1595 compound library and incubated for (**a**) 36 hours or (**d**) 24 hours prior to the first round of HCI. Normalised dFLASH activity for (**b**) the 36-hour time point or the (**e**) 24-hour time point is shown. (**b, e**) Compounds that changed EGFP >±2SD are shown in grey and excluded as hits. Compounds that increase Tomato/EGFP >2SD from the vehicle controls (dashed line) are highlighted in red. After activation screen, the compound wells were then treated with 1mM DMOG for (**a, c**) 36 hours or (**d, f**) 24 hours prior to another round of HCI. (**c, f**) Compounds that decreased dFLASH activity greater than 2SD from DMOG controls (dashed line) are shown in red. Compounds that changed EGFP >±2SD are shown in grey and excluded as hits. Normalised dFLASH output was normalised by Z scoring for all analysed wells.

Figure 6. **Identification of how novel activator and inhibitor of HIF-1α impact reporter activity**

(**a, b**) Identified inhibitor RQ500235 from the bimodal screen (**a**) represses the dFLASH-HIF's normalised (black) and Tomato (magenta) expression in a dose dependant manner (n=2) at the same concentrations by which (**b**) RQ500235 decreases HIF-1α protein by immunoblot in HEK293T cells with HIF-1α.HA-Flag endogenously tagged. (**c**) HIF-1α transcript is significantly decreased (*p =0.0139) in HEK293T cells treated for 6 hours with RQ500235 (n =3). Significance assessed with students t-test (*p < 0.05, p = 0.0139). (**d**) Identified dFLASH-HIF activator RQ200674 from the bimodal screen recapitulated activating activity at 50μM against HEK293T dFLASH-HIF (n = 2). (**e**) *in vitro* iron chelation assay of RQ200674 displays weak chelating activity at 236μM from line of best fit (n = 3) compared to dipyridyl.

640
641

643 **Supplementary Figures**

644
645

646
647
648

Supplementary Figure 1. Optimised dFLASH design enables robust HIF sensor.

(**a-b**) HEK293T cells with HRE-dFLASH constructs without EGFP and (**a**) expressing monomeric Tomato or (**b**) dimeric Tomato were treated with and without DMOG for 48 hours and quantified by FACS. Tomato MFI >200AU was used to compare induction (black line). (**c-e**) HEK293T and HEPG2 cells were transduced with HRE-dFLASH reporters that had different downstream promoters controlling EGFP or Tomato cassette composition and treated for 48 hours with and without 1mM DMOG then imaged using HCI. The (**d**) Tomato EGFP MFI ratio and (**e**) EGFP MFI for each backbone was then compared (Data from three independent biological replicates).

(**f**) HEK293T cells transduced with reporter constructs containing the PGK/CMV or EF1a promoters were compared for DMOG induction by HCI after 48 hours of DMOG treatment (Data from three independent biological replicates). Significance was assessed with a Two-Way ANOVA (**** $p < 0.001$, ns = not significant). (**h,g**) HEK293T cells with the HRE enhancer and different dFLASH backbone compositions for (**f**) dFLASH with the EF1a promoter, (**g**) PGK/CMV dFLASH with Tomato alone as the upstream cassette or (**h**) dFLASH-HIF were treated for 48-hours DMOG or Dox prior to FACS.

667
668
669

Supplementary Figure 2. dFLASH provides a TF-responsive, versatile reporter platform in heterogenous pools.

(**a-b**) HEK293T cells were transduced with (**a**) dFLASH with no enhancer and treated with 1mM DMOG or 0.1% DMSO (Ctrl) or (**b**) GalRE-dFLASH and Gal4DBD-miniVPR and treated with milliQ (Ctrl) or 1μg/mL Dox after 48 hours prior to FACS. Dot plots of populations' Tomato and EGFP intensity with or without activating ligand and histograms comparing EGFP and Tomato MFI between treated populations are shown. (**c-h**) Dot plots and EGFP histograms for (**c, f**) dFLASH-HIF, (**d, g**) dFLASH-synFIH, and (**e, h**) dFLASH-PR polyclonal pools for **Figure 2a-c** ligand induction.

Supplementary Figure 3. Synthetic transcription factors drive a strong response from the GalRE reporter and can respond to endogenous pathways.
(**a**) GAL4DBD-miniVPR is expressed from an independent dox-dependent construct that produces the transcription factor, enabling it to interact with GalRE-dFLASH. (**b**,**c**) HEK293T GalRE-dFLASH cells were transduced with GAL4DBD-miniVPR expression construct and were treated with and without doxycycline for 48 hours prior to HCI for (**b**) Tomato expression (top panels) and EGFP expression (bottom panels). (**c**) Fluorescence intensity was also quantified for treated populations (n=3, mean ±sem). FC is Fold change between the populations. (**d, e**) To confirm HEK293T dFLASH-synFIH system was FIH dependent, (**d**) GalRE-dFLASH and the GAL4DBD-HIFCAD transduced into HEK293T cells with FIH knocked out. (**e**) FIH KO cells were compared with wildtype HEK293T dFLASH-synFIH (WT) in a 200ng/mL dox background for DMOG-dependent reporter induction by HCI (n=3). (**c, e**) Significance was assessed by t-test with Welch's correction (ns = not significant, *** $p < 0.001$, ****$p < 0.0001$).

HIF response Pathway

HEK293T dFLASH-HRE

HepG2 dFLASH-HRE

PGR response Pathway

T47D dFLASH-PRE

BT474 dFLASH-PRE

Synthetic FIH Sensor

HEK293T dFLASH-synFIH

Supplementary Figure 4. Clonal dFLASH reporter lines enables improved reporting across different cell types.

(**a-c**) Flow cytometry of clonal lines derived with the dFLASH-HIF reporter in (**a**) HEK293T (**Figure 3b**) and (**b**) HepG2 cells after 48 hours of 0.5mM DMOG or DMSO stimulation. (**d-h**) dFLASH-PGR functionality was assessed by flow cytometry in (**d**) T47D (**Figure 3f**) and (**e, g**) BT474 cells after 48 hours of vehicle or R5020 stimulation. (**f,g**) T47D dFLASH-PGR cells were treated with increasing concentrations of levels of R5020 (0.01-100nM, 8 replicates per group) and (**f**) imaged over 38 hours with temporal HCI or (**g**) imaged at 48 hours to determine sensitivity to R5020. (**i**) Comparison of the induction of the T47D mcdFLASH-PGR line to different steroids (10nM R5020, 35nM E2, 10nM DHT, 10nM Dex, 10nM RA) by HCI after 48 hours of treatment. (**g**) and (**i**) are the mean±sem of min/max normalised Tomato/GFP (within each expt) from n = 3 independent experiments (24 replicates), except Dex and RA (n=2 (16 replicates)).(**j, k**) Clonally derived HEK293T dFLASH-synFIH cells were (**j**) analysed by flow cytometry after 48 hours of 200ng/mL Dox and DMSO (Ctrl) or Dox and DMOG treatment (**Figure 3k**) and (**k**) the clonally derived line (mcDMSO/mcDMOG) temporal HCI result (**Figure 3l**) was compared with induction from the polyclonal line (pcDMSO/pcDMOG) (**Figure 2j**).

Supplementary Figure 5. HIF-1α is the predominant isoform that affects the reporter in HEK293T cells

(**a**) Monoclonal HEK293T cells with endogenously tagged HIF-1α or HIF-2α were treated with hypoxia (<1% O_2) for 16 hours prior to blotting for factor expression against the HA tag. Representative of three independent experiments. (**b**) mcdFLASH-HIF cells were imaged and quantified over 72-hour period with 1mM DMOG or vehicle and with or without 10μM of the HIF-2α antagonist (PT-2385).

a.

Supplementary Figure 6. CRISPRoff VHL knockdown induces mcdFLASH-HIF
reporter lines.
(**a**) HEK293T cells were first transduced with dFLASH-HRE and a clonal reporting line
was derived after hygromycin (HygroB) treatment. This line was in turn transduced
with the VHL sgRNA vector and selected with puromycin (Puro). This line was then
transduced with the CRISPRoffv2.1 vector and selected with blasticidin S (Blast) and
populations were subjected to flow cytometry after 5 days of selection or 10 days for
reporter expression. (**b-d**) dot plots for dFLASH expression from the (**b**) non-
CRISPRoff parental line, (**c**) EF1a-CRISPRoffv2.1 transduced and (**d**) SFFVp-
CRISPRoffv2.1 population after 5 or 10 days of blasticidin selection from **Figure 4**.
759
760
761

765
766
767
768
Supplementary Figure 7. Hits selection and assessment of bimodal screen
reproducibility between independant screens.
Compound-induced dFLASH-HIF reporter activity was used to score as hits from the
(**a-d**) 36-hour or the (**e-h**) 24-hour bimodal screens for reporter activation or inhibition
against Tomato MFI and adjusted P scores. Lines indicate cut offs for hit criteria and
compounds that changes EGFP $>\pm$2SD are shown in grey and hits are shown in red
for each metric. (**i, j**) Pearson correlations of the Tomato/EGFP between the 36-hour
and the 24-hour screens for (**i**) reporter activation (R = 0.62, p < 2.2×10^{-16}) or (**j**)
reporter inhibition (R = 0.62, p < 2.2×10^{-16}) for all 1595 compounds screened. Line
indicates line of best fit, grey boundary is 95% confidence interval.
779

780
781
782
783

Supplementary Figure 8. Rescreening of activator hits from 1595 compound
small molecule screen reveals RQ200674 that causes normoxic stabilisation of
HIF-1α

 (**a**) The 11 top performing hits from the activator screens, including RQ200674
(**Figure 6d**) were rescreened against HEK293T mcdFLASH-HIF at 3 concentrations
10μM, 25μM and 50μM compared and were compared for Tomato/GFP and Tomato
MFI dFLASH induction against vehicle and DMOG treated populations for reporter
induction (n=2). (**b**) Immunoblot of HEK293T with endogenously tagged HIF-
1α.HAFLAG and treated with vehicle (DMSO) and 1mM DMOG controls or 100μM
and 200μM of RQ200674 for 18 hours. Representative of 2 independent experiments.

794
795
796
797

Supplementary Figure 9. Flavokawain B, Celastarol and RQ500235 decrease dFLASH-HIF and proteasomal inhibition doesn't rescue RQ500235 impact on HIF-1α.

(**a-c**) The 18 top inhibitory compounds, including (**b**) Flavokawain B (RQ100976) and (**c**) Celastarol (RQ000155) and RQ500235 (**Figure 6a**) were screened against dFLASH-HIF at three concentrations (10, 25 and 50μM), in a DMOG background at 24 hours for inhibitory activity against vehicle and DMOG controls. (**d**) Immunoblot of HEK293T cells with endogenously tagged HIF-1α.HAFLAG treated with DMOG for 12 hours, 50μM RQ500235 for 6 hours and treated and MG132 for 3 hours with relevant vehicle controls. Representative of 2 independent experiments.

859
860
861
Supplementary Movie 1. Single cell temporal dynamics of HEK293T
mcdFLASH-HIF cells
HEK293T mcdFLASH-HIF cells were seeded at 1×10^5 cells/dish in Poly-D-Lysine
coated plates overnight prior to imaging with spinning disk confocal microscopy at
40x magnification. Cells were imaged every 15 min for 48 hours for Tomato
(Magenta) and EGFP (Green) expression. Time stamps are given in top left.
868
869

870
871
872
873
Supplementary Movie 2. Single cell temporal dynamics of T47D mcdFLASH-
PGR cells
T47D mcdFLASH-PGR cells were seeded at 5×10^5 cells/dish in Poly-D-Lysine
coated plates overnight prior to imaging with spinning disk confocal microscopy at
40x magnification. Cells were imaged every 15 min for 48 hours for Tomato
(Magenta) and EGFP (Green) expression. Time stamps are given in top left.
880
881
882
883
884 **Methods:**
885
Plasmid Construction. cDNAs were amplified using the Phusion polymerase (NEB)
and assembled into ClaI/NheI digested pLV410 digested backbone by Gibson
assembly[31]. Sequence verified LV-REPORT plasmid sequences and constructs are
listed in **Supplementary Table 1**. Briefly, the plasmids contained an upstream multiple
cloning sites followed by a minimal promoter (derived from the pTRE3G minimal
promoter) and then followed by a reporter construct mnucTomato/HSVtk-2a-Neo or
other variants). This was then followed by a constitutive promoter (EF1a, PGK or
PGK/CMV) driving the expression or hygromycinR cassette with or without a 2a linked
d2nucEGFP (**Supplementary Figure 1C**).
895

To improve the performance of our previously reported lentiviral inducible expression systems[65], the PGK promoter in Tet-On3G IRES Puro was replaced by digestion with MluI/NheI and insertion of either EF1a-Tet-On3G-2A-puro, EF1a-Tet-On3G-2A-BlastR or EF1a-Tet-On3G-2A-nucTomato using Phusion polymerase (NEB) amplified PCR products from existing plasmids. Plasmids were cloned by Gibson isothermal assembly and propagated in DB3.1 cells (Invitrogen). We also generated a series of constitutive lentiviral plasmids as part of this work pLV-EgI-BlastR (EF1a-Gateway-IRES-BlastR), pLV-EgI-ZeoR (EF1a-Gateway-IRES-ZeoR), pLV-EgI-HygroR (EF1a-Gateway-IRES-HygroR), pLV-SFFVp-gI-BlastR (SFFVp-Gateway-IRES-BlastR), pLV-SV40p-gI-BlastR (SV40p-Gateway-IRES-BlastR). These plasmids were constructed by isothermal assembly of G-Blocks (IDT DNA) or PCR fragments, propagated in ccbD competent cells, sequence verified and deposited with Addgene (**Supplementary Table 1**).

The Lentiviral backbone expression construct pLV-TET2BLAST-GtwyA was then using to insert expression constructs cloned into pENTR1a by LR Clonase II enzyme recombination (Cat#11791020, Thermo). GAL4DBD-HIFCAD (727-826aa) and the GAL4DBD[28] were cloned into pENTR1a by ScaI/EcoRV or KpnI/EcoRI respectively. The miniVPR sequence[29] was cloned into the pENTR1a-GAL4DBD construct at the EcoRI and NotI sites. The pENTR1a vectors were then Gateway cloned into the pLV-TET2PURO-GtwyA vector. pENTR1a-CRISPRoffv2.1 was generated by inserting an EcoRI/NotI digested CRISPRoff2.1 (CRISPRoff-v2.1 was a gift from Luke Gilbert, Addgene #167981) into pENTR1a plasmid. pLV-SFFVp-CRISPRofv2.1-IRES-BLAST and pLV-EF1a-CRISPRofv2.1-IRES-BLAST were generated by pENTR1a by LR Clonase II enzyme recombination (Cat#11791020, Thermo). All Lentiviral plasmids were propagated in DH5a without any signs of recombination.

Enhancer element cloning. The 12x HRE enhancer from hypoxic response target genes (PGK1, ENO1 and LDHA) was liberated from pUSTdS-HRE12-mCMV-lacZ[24] with XbaI/SpeI and cloned into AvrII digested pLV-REPORT plasmids. Progesterone responsive pLV-REPORT-PRECat PRECat was cloned by isothermal assembly of a G-Block (IDT-DNA) containing enhancer elements from 5 PGR target gene enhancers (Zbtb16, Fkbp5, Slc17a11, Erfnb1, MT2)[66] into AscI/ClaI digested pLV-REPORT(PGK/CMV). Gal4 response elements (5xGRE) were synthesised (IDT DNA) with ClaI/AscI overhangs and cloned into Cla/AscI digested pLV-REPORT(PGK/CMV). Sequences are in **Supplementary Table 2.**

Mammalian cell culture and ligand treatment. HEK293T (ATCC CRL-3216), HEPG2 (ATCC HB-8065) line were grown in Dulbecco's Modified Eagle Medium (DMEM high glucose) + pH 7.5 HEPES (Gibco), 10% Foetal Bovine Serum (Corning 35-076-CV or Serana FBS-AU-015), 1% penicillin-streptomycin (Invitrogen) and 1% Glutamax (Gibco). T47D (ATCC HTB-133) or BT474 (ATCC HTB-20) were grown in RPMI 1640 (ATCC modified) (A1049101 Gibco) with 10% Foetal Bovine Serum (Fisher Biotech FBS-AU-015) and 1% penicillin-streptomycin[67]. Cells were maintained at 37^0C and at 5% CO_2.Clonal lines were isolated by either limiting dilution or FACS single cell isolation into 96 wells trays. Resultant monoclonal populations were evaluated for single colony formation or assessed by HCI or FACS. Ligand treatments were done 24 hours after seeding of cells in requisite plate or vessel. Standard

concentrations and solvent, unless specified otherwise, are 200ng/mL Doxycycline
(Sigma, H_2O), 0.5mM or 1mM DMOG (Cayman Scientific, DMSO), 100nM R5020
(Perkin-Elmer NLP004005MG, EtoH), 35nM Estradiol (E2, Sigma E2758, EtOH),
10nM all-trans retinoic acid (RA, Sigma #R2625), 10nM Dihydrotestosterone (DHT,
D5027), 10nM Dexamethasone (Dex, Sigma D4902), 10μM PT-2385 (Abcam,
DMSO).
950
Lentiviral Production & stable cell line production. Near confluent HEK293T cells
were transfected with either psPAX2 (Addgene #12260) and pMD2.G (Addgene
#12259) or pCMV-dR8.2 dvpr (Addgene #8455), pRSV-REV (Addgene; #12253) and
pMD2.G along with the Lentivector (described above) and PEI (1μg/μl,
polyethyleneimine) (Polysciences, USA), Lipofectamine 2000, or Lipofectamine 3000
at a 3μl:1μg ratio with DNA. Media changed 1-day post-transfection to complete media
or Optimem. Virus was harvested 1-2 days post-transfection, then viral media was
filtered (0.45μM or 0.22μM, Sartorius) before the target cell population was transduced
at a MOI < 1. Cells were incubated with virus for 48 hours prior media being exchanged
for antibiotic containing complete media. Standard antibiotic concentrations were
140μg/mL hygromycin (ThermoFisher Scientific #10687010), 1μg/mL Puromycin
(Sigma; #P8833) or 10μg/mL Blasticidin S (Sigma; CAT#15205).
963
Generation of CRISPR knockout or knockdown cell lines. Generation of CRISPR
knockout guides and plasmids against FIH has been previously described[68]. These
guides were transfected into HEK293T cells and with PEI at a 3μg:1μg ratio then
clonally isolated as above. Knockouts were confirmed with PCR amplification and
sanger sequencing coupled with CRISPR-ID[69]. FIH knockouts were selected via serial
dilution and confirmation of knockout by sequencing and T7E1 assay. The VHL sgRNA
guides were selected from the Dolcetto CRISPRi library[70] with BsmBI compatible
overhangs (**Supplementary Table 3**). These oligos were annealed, phosphorylated
then ligated into BsmBI-digested pXPR050 (Addgene#9692), generating XPR-050-
VHL. Monoclonal HEK293T LV-REPORT-12xHRE cell lines were transduced with the
XPR-050-sgVHL virus, and stable cell lines selected with Puromycin. Subsequently,
LV-SFFVp-CRISPRoffv2.1-IRES-BlastR or LV-EF1a-CRISPRoffv2.1-IRES-BlastR
virus was infected into HEK293T LV-REPORT-12xHRE/XPR-050-sgVHL stable cells
and selected with Blasticidin S (15μg/ml) for 5 days. FACS was used to assess
activation of the dFLASH-HRE reporter in parental (dFLASH-HRE/sgVHL) or
CRISPRoffv2.1 expressing cells at day 5 or day 10 after Blasticidin S addition.
980
CRISPR knock-in of tags to endogenous HIF-1α and HIF-2α. CRISPR targeting
constructs clones targeting adjacent to the endogenous HIF-1α and HIF-2α stop
codons[71]. Constructs were cloned into px330 by ligating annealed and phosphorylated
oligos with BbsI digested px330, using hHIF-1α and hHIF-2α CTD sgRNA
(**Supplementary Table 3**). Knock-in of HA-3xFlag epitopes into the endogenous HIF-
1α or HIF-2α loci in HEK293T cells was achieved by transfection with 0.625 μg of
pNSEN, 0.625μg of pEFIRES-puro6, 2.5μg of px330-sgHIF-α CTD, and 1.25μg of
ssDNA HDR template oligo containing flanking homology to CRISPR targeting site the
tag insertion and a PAM mutant into ~0.8x10^6 cells using PEI (3:1). 48 hours after
transfection, the medium was removed from cells and replaced with fresh medium
supplemented with 2 μg/ml puromycin for 48 hours and the cell medium was changed

to fresh medium without puromycin. 48 hours later cells were seeded by limiting
dilution into 96-well plates at an average of 0.5 cells/well. Correct integration was
identified by PCR screening using HIF-1α and HIF-2α gDNA screening primers
(**Supplementary Table 4).** Positive colonies reisolated as single colonies by limiting
dilution. Isolated HIF-1α and HIF-2α tag insertions were confirmed by PCR, sanger
sequencing and western blotting.
998

High Content Imaging (HCI). Cells were routinely seeded at 1×10^4 to 5×10^4 cells per
well in black walled clear bottom 96 well plates (Costar Cat#3603), unless otherwise
stated. Cell populations were imaged in media at the designated time points at 10x
magnification and 2x2 binning using the ArrayScan^{TM} XTI High Content Reader
(ThermoFisher). Tomato MFI and EGFP MFI was imaged with an excitation source of
560/25nm and 485/20nm respectively. Individual nuclei were defined by nuclear EGFP
expression, nuclear segmentation and confirmed to be single cells by isodata
thresholding. Nuclei were excluded from analysis when they couldn't be accurately
separated from neighbouring cells and background objects, cells on image edges and
abnormal nuclei were also excluded. EGFP and Tomato intensity was then measured
for each individual nucleus from at least 2000 individual nuclei per well. Fixed
exposure times were selected based on 10-35% peak target range. Quantification of
the images utilised HCS Studio^{TM} 3.0 Cell Analysis Software (ThermoFisher). For
assessment of high throughput robustness of each individual reporting line in a high
throughput setting (HTS-HCI), replicate 96 well plates were seeded for the HIF (10
plates), PGR (5 plates) and synFIH (3 plates) monoclonal reporter lines and imaged
as above at 48 hours. For the HIF line, each plate had 6 replicates per treatment
(vehicle or DMOG) per plate. For the PGR, 24 replicates per treatment, either vehicle
or R5020 per plate were present with edge wells excluded. 24 replicates per treatment
were also used for synFIH, with system robustness assessed between the
DOX/DMSO and DOX/DMOG treatment groups. Z' and fold change (FC) for the
Tomato/EGFP ratio for each individual plate was then calculated as per [34]:

$$Z' = 1 - \frac{(3\sigma_{c+} - 3\sigma_{c-})}{|\mu_{c+} - \mu_{c-}|}$$

Z' for every plate across each system was confirmed to be >0.5. Overall robustness
of each system is the average of every individual Z' and FC for each system. For
temporal high content imaging, HIF, PGR and synHIF lines were seeded in plates and
treated with requisite ligands immediately prior to HCI. Four treatment replicates per
plate were used to assess the polyclonal population. 4 treatments per plate were used
to assess the synFIH monoclone (DOX, DMSO, DOX/DMSO, DOX/DMOG), with
100ng/μL Doxycycline utilised, and 8 treatments per plate (vehicle, DMOG or R5020)
were used to assess the PGR and HIF monoclonal lines. Plates were humidified and
maintained at 37^{0}C, 5% CO_2 throughout the imaging experiment. Plates were then
imaged every 2 hours for 40-48 hours. At every timepoint, a minimum 2000 nuclei
were resampled from each well population.
1033

1034   **T47D mcdFLASH-PGR R5020 Dose response curve EC50 calculation.** T47D
1035   mcdFLASH-PGR cells were treated with increasing doses of 0.01-100nM R5020 and
1036   quantified by HCI after 48hrs. Tomato/GFP values were min/max normalised within
1037   each experiment (n = 3) and the EC50 constant and curve fitted using the drc R
1038   package from [72].

1039

Bimodal small molecule screen to identify activators or inhibitors of the hypoxic
response pathway. Library of natural and synthetic compounds was supplied by Prof.
Ronald Quinn and Compounds Australia, available by request. 5mM of each of the
1595 compounds were spotted in 1μL DMSO into Costar Cat#3603 plates and stored
at -80^{o}C prior to screening. Plates were warmed to 37^{o}C prior to cell addition.
Monoclonal HIF HEK293T reporter cells were seeded at 0.5×10^4 cells per well across
20 Costar Cat#3603 plates pre-spiked with 5mM of compound in 1uL of DMSO in
100uL. On each plate, 4 wells were treated with matched DMSO amounts to
compound wells as were four 1mM DMOG controls. Plates were then imaged using
HCI (described above) at 36 hrs or 24 hours for reporter activation. Wells were then
treated with 100uL of 2mM DMOG (for 1mM DMOG final, 200uL media final). 4 vehicle
and 8 DMOG-treated controls (excluding the initial controls from the activator screen)
were used for the inhibitor screen. Cells were imaged again 36 hours (Screen 1) or 24
hours (Screen 2) after treatment with 1mM DMOG in the compound wells. Data was
Z scored and control wells were used to establish gating for abnormal expression of
Tomato and EGFP fluorophores. For the activator screen, compounds within +/- 2SD
EGFP MFI of vehicle wells were counted as having unchanged transcriptional effects.
Compounds with Tomato/EGFP ratio greater than +2SD of vehicle controls was
counted as a putative hit. For the inhibitor screen, compounds within +/- 2SD EGFP
MFI of DMOG controls were counted as having unchanged GFP expression and
Compounds with Tomato/EGFP ratio lower than -2SD from the DMOG control were
considered putative inhibitors. To correct for false positives within each screen, Z
scored compounds were converted to their respective P score and adjusted with a [73]
correction. Pearson correlations were then used to compare compound expression
between screens with the base R package (4.4.0). Putative activators and inhibitors
identified in the screens were re-spotted at 1mM, 2.5mM and 5mM in 1μL of DMSO in
Costar Cat#3603 96 well trays. Activators were rescreened by HCI after 24 hours
against 1×10^4 cells HIF reporter monoclones in biological duplicate against with
vehicle and 1mM DMOG controls in 100μL. Inhibitors were rescreened by HCI after
24 hours in duplicate against 1×10^4 cells HIF reporter monoclones with 1mM DMOG
to compound wells. Final compound concentrations were 10μM, 25μM and 50μM
respectively and Tomato MFI and Tomato/EGFP ratio for each compound was
assessed.

1073

Reverse Transcription and Real Time PCR. Cells were seeded in 60mm dishes at
8×10^4 cells per vessel overnight before treatment for 48 hours with 1mM DMOG or
0.1% DMSO. Cells were lysed in Trizol (Invitrogen), and RNA was purified with Qiagen
RNAEasy Kit, DNaseI treated and reverse transcribed using M-MLV reverse
transcriptase (Promega). cDNA was then diluted for real time PCR. Real-time PCR
used primers specific for *HIF-1α*, and human RNA Polymerase 2 (*POLR2A*)
(**Supplementary Table 4**). All reactions were done on a StepOne Plus Real-time PCR
machine utilising SYBER Green, and data analysed by 'QGene' software. Results are
normalised to *POLR2A* expression. RT-qPCR was performed in triplicate and single
amplicons were confirmed via melt curves.

1084

Flow cytometry analysis and sorting (FACS). Prior to flow cytometry, cells were
trypsinised, washed in complete media and resuspended in resuspended in flow

cytometry sort buffer (Ca^{2+}/Mg^{2+}-free PBS, 2%FBS, 25mM HEPES pH 7.0) for cell
sorting) prior to cell sorting or flow cytometry analysis buffer (Ca^{2+}/Mg^{2+} free PBS,
2%FBS, 1mM EDTA, 25mM HEPES pH 7.0) for analysis followed by filtration through
a 40μM nylon cell strainer (Corning Cat#352340. Cell populations were kept on ice
prior to sorting. Flow cytometry was performed either using the BD Biosciences BD
LSRFortessa or the BD Biosciences FACS ARIA2 sorter within a biosafety cabinet
and aseptic conditions, using an 85μM nozzle. Cell populations were gated by FSC-
W/FSC-H, then SSC-W/SSC-H, followed by SSC-A/FSC-A to gate cells. EGFP
fluorescence was measured by a 530/30nm detector, and the Tomato fluorescence
was determined with the 582/15nm detector. A minimum of 10,000 cells were sorted
for all FACS-based analysis. Data is presented as $\log_{10}$ intensity for both fluorophores.
Tomato induction was gated from the top 1% of the negative control population. Cell
counts for histograms are normalised to mode unless stated otherwise. FACS
analysis was done on FlowJo^{TM} v10.9.1 software (BD Life Sciences)[74].
1101
Time Lapse Spinning Disc Confocal Microscopy. HEK293T mcdFLASH-HIF and
T47D mcdFLASH-PGR cells were seeded at 1×10^5 or 5×10^5 cells per dish
respectively, onto 50μg/mL poly-D-lysine μ-Dish 35 mm, high Glass Bottom dishes
(Ibidi, #81158) in FluoroBrite DMEM (Gibco, A1896701)/10% FBS/ 1% Pens/1%
Glutamax/10mM HEPES pH7.9 and incubated overnight at 37°C with 5% CO2 prior
imaging. Cells were treatment with either 0.5mM DMOG (mcdFLASH-HIF) or 100nM
R05020 (mcdFLASH-PGR) immediately prior to imaging with a CV100 cell voyager
spinning disk confocal Tomato (561 nm, 50% laser, 400ms exposure and 20% gain)
and EGFP (488 nm, 50% laser, 400ms exposure and 20% gain) fluorescence for 48
hours post treatment with 15min imaging intervals. Images were captured at 40x with
an objective lens with a ~30μm Z stack across multiple fields of view. Maximum
projected intensity images were exported to Image J for analysis and movie creation.
1114
Cell Lysis and Immunoblotting. Cells were washed in ice-cold PBS and lysates were
generated by resuspending cells in either cell lysis buffer (20mM HEPES pH 8.0,
420mM $NaCl_2$, 0.5% NP-40, 25% Glycerol, 0.2mM EDTA, 1.5mM $MgCl_2$, 1mM DTT,
1x Protease Inhibitors (Sigma)) (**Supp Figure 4**) or urea lysis buffer (6.7M Urea,
10mM Tris-Cl pH 6.8, 10% glycerol, 1% SDS, 1mM DTT) (**Figure 6, Supp Figure 8,
9**). Quantification of protein levels was done by Bradford Assay (Bio-Rad). Lysates
were separated on a 7.5% SDS-PAGE gel and transferred to nitrocellulose via
TurboBlot (Bio-Rad). Primary Antibodies used were anti-HIF1α (BD Biosciences #),
anti-HA (HA.11, Biolegend #16B12), anti-Tubulin (Serotec #MCA78G), anti-GAPDH
(Sigma #G8796), anti-ARNT (Proteintech #14105-1-AP). Primary antibodies were
detected using horseradish peroxidase conjugated secondary antibodies (Pierce
Bioscience #). Blots were visualised via chemiluminescence and developed with
Clarity Western ECL Blotting substrates (Bio-Rad).
1128
***In vitro* iron chelation activity assay.** Chelation of iron for RQ200674 was
measured by a protocol adapted from [75] for use in 96 well plate format. 0.1mM
$FeSO_4$ (50μL) and 50μL of RQ200674, Dipyridyl (positive control) or DMOG
solutions were incubated for 1hr at room temperature prior to addition of 100μL of
0.25mM Ferrozine (Sigma) and incubated for a further 10 minutes. Absorbance was
measured at 562nM. Chelation activity was quantified as:

$$\text{Chelation activity} = \frac{(A_{control} - A_x)}{A_{control}} \times 100$$

Where $A_{control}$ is absorbance of control reactions without RQ200674, DP or DMOG
and A_x is absorbance of solutions with compound.

Statistical Analysis. All data in graphs were presented as a mean $\pm$ sem unless
otherwise specified. Significance was calculated by a Two-Way ANOVA with Tukey
multiple comparison or unpaired t-test with Welches correction where appropriate
using Graphpad PRISM (version 9.0.0). All statistical analysis is from three
independent biological replicates

Figure Creation. Schematics and diagrams were created with BioRender
(BioRender.com) and graphs were made either with ggplot package in R[76] and
GraphPad PRISM (version 9.0.0).

Data Availability. Source data are provided with this paper. Additional data, including
full construct sequences, are available from corresponding authors upon request.
Constructs not available on Addgene can be requested from corresponding authors.

Acknowledgements. We thank Nicholas Smith, Alexander Pace, and members of
our laboratories for critical feedback and helpful discussions. We also wish to
acknowledge Adelaide Microscopy and the AHMS and SAHMRI Flow Cytometry
facilities for technical assistance. We acknowledge Compounds Australia
(www.compoundsaustralia.com) for their provision of specialized compound
management and logistics research services to the project. This work was supported
by Australian Government Research Training Scholarships (T.P.A, A.E.R), The
Emeritus Professor George Rodgers AO Supplementary Scholarship (T.P.A, A.E.R).
The Playford Memorial Trust Thyne Reid Foundation Scholarship (A.E.R). The George
Fraser Supplementary Scholarship (A.E.R), The University of Adelaide Biochemistry
Trust Fund (D.J.P. and M.L.W) and the Bill and Melinda Gates Foundation
Contraceptive Discovery Program [OPP1771844] (D.C.B, D.L.R).

Author contributions. Study was initially conceived by D.C.B and M.L.W. T.P.A,
D.C.B., A.E.R designed and performed experiments. T.P.A, D.C.B., M.L.W, M.L. and
R.J.Q. performed and analysed the bimodal screening campaign. M.R. and A.E.R.
derived FIH KO cell line. T.P.A, D.C.B and M.L.W wrote the manuscript with input
from all authors. Work was supervised by D.J.P, D.L.R. & M.L.W.

Source Data. Source data for figures is available with this manuscript.

Competing interests. The authors declare no competing interests.

Correspondence and requests for materials. Should be addressed to David C.
Bersten.

1180
1181

Supplementary Table 1: Synthetic toolkit for generation of reporter cell lines

Deposit Name:	Availability	Purpose
Dual fluorescent reporter constructs:		
pLV-REPORT(EF1a)	Addgene #172326	Reporter with mnucTomato and EF1a downstream promoter
pLV-REPORT(EF1a)-TTN	Addgene #172327	Reporter with mnucTomato-HSVtk-2A-NeoR and EF1a downstream promoter
pLV-REPORT(PGK)	Addgene #172328	Reporter with mnucTomato-HSVtk-2A-NeoR and PGK downstream promoter
pLV-REPORT(PGK/CMV)	Addgene #172330	Reporter with mnucTomato-HSVtk-2A-NeoR and PGK/CMV downstream promoter
12xHRE-pLV-Report-EF1a	Addgene: #172333	Reporter with HRE enhancer
12xHRE-pLV-REPORT(PGK)	Addgene #172334	Reporter with HRE enhancer
12xHRE-pLV-REPORT(PGK/CMV)	Addgene #172335	Reporter with HRE enhancer
PREcat-pLV-REPORT(PGK/CMV)	By Request	Reporter with a PR-responsive concatemer, with enhancers from 5 target genes, containing 6 PR response elements.
5xGRE-pLV-REPORT(PGK/CMV)	Addgene #172336	Reporter with GRE enhancer
12xHRE-pLV-REPORT(EF1a)	By Request	Reporter with HRE
12xHRE- pLV-REPORT(EF1a)-tdnucTomato	By Request	Reporter with tdnucTomato and EF1a downstream promoter
Protein expression constructs:		
pLV-TET2Puro	By Request	Doxycycline-inducible expression vector
pLV-TET2BlastR	By Request	Doxycycline-inducible expression vector
pLV-TET2nucTomato	By Request	Doxycycline-inducible expression vector
pLV-TET2Puro-gal4DBD-miniVPR-HA	Addgene #207171	Doxycycline-inducible expression vector for GAL4DBD-miniVPR
pLV-TET2Puro-gal4DBD-HIFCAD	Addgene #207173	Doxycycline-inducible expression vector for GAL4DBD-HIFCAD (727-826) with Myc tag
pEF-IRES-puro6 gal4DBD-HIFCAD myc tag	Addgene #207171	Constitutively expresses GAL4DBD-HIFCAD (727-826) with Myc tag
pEF-IRES-puro6 gal4DBD-HIFCAD pGalO linker	Addgene #207172	Constitutively expresses GAL4DBD-HIFCAD (727-826) with Myc tag
pENTR1a-CRISPRoffv2.1	Addgene #207174	Lentiviral expression vector for CRISPRoffv2.1 with BFP tag
pLV-EgI-NeoR	Addgene #207175	Gateway-compatible lentiviral expression plasmid with Neomycin resistance
pLV-EgI-BlasR	Addgene #207176	Gateway-compatible lentiviral expression plasmid with Blasticidin resistance
pLV-EgI-HygroR	Addgene #207177	Gateway-compatible lentiviral expression plasmid with Hygromycin resistance
pLV-EgI-ZeoR	Addgene #207178	Gateway-compatible lentiviral expression plasmid with Zeocin resistance

1182
1183

Supplementary Table 2: Sequences for enhancer cloning

PRECat (G-block)
gaattacaaaaacaaattacaaaaattcaaaattttatcgatTGCATGCCTGCTTACATAAGGAAGTACAGAGTGTACCAAAACAGCAGACCCAAAAAAAGCCTGAAATGTGAGAACCCCAAAACTGTACAGCTTGTATTTCAGGAAGCAAAACTGAGGACGCAAGCCGTCTTCATGGAATAATACATCCTGTTCCCACAAGTGACGTTAGCTTCCAGACTGTGCACAGAGTGCACACTTCACCCAGTGTGTGTCATCATGGTCACACAGTGTTCTTTCCGTGGTCACATCTGTGTCCACATTTCCTCCTTTTGATGGGAACAAAGCAGTCATGTTAGGAAGGGAAAGGACACGGTGTTTAATCACACAATCCATGGACAGCCGTGGGCATCCAGTAATGCCTGGAATGAGTCAAGAAGGCATTGCCCCAGTTTTCACTAAGAGCTGCGAGGACAGCCTGTCCTGTTACAACCCACCCACAGCCTCCGTTGAGGCGCGCCAGCTTTAGGCGTGTACGGTGGGCGCCTATAAAGC
5xGRE
GGTACCAGCTTGCATGCCTGCAGGTCGGAGTACTGTCCTCCGAGCGGAGTACTGTCCTCCGAGCGGAGTACTGTCCTCCGAGCGGAGTACTGTCCTCCGAGCGGAGTACTGTCCTCCGAGCGGAGAC

Supplementary Table 3: Index of all sgGuide oligos used

	Upper (5'-3')	Lower (5'-3')
VHL Knockdown sgGuide	CACCGCCGGGTGGTCTGGATCGCGG	AAACCCGCGATCCAGACCACCCGGC
hHIF-1α CTD sgRNA	CACCGTGAAGAATTACTCAGAGCTT	AAACAAGCTCTGAGTAATTCTTCA
hHIF-2α CTD sgRNA	CACCGCCTCCTCAGAGCCCTGGACC	AAACGGTCCAGGGCTCTGAGGAGGC

Supplementary Table 4: Primer sets for qPCR and PCR confirmation

	Forward (5'-3')	Reverse (5'-3')
qPCR HIF-1α	TATGAGCCAGAAGAACTTTTAGGC	CACCTCTTTTGGCAAGCATCCTG
qPCR PolR2a	GCACCATCAAGAGAGTGCAG	GGGTATTTGATACCACCCTCT
HIF-1α gDNA primers	GGCAATCAATGGATGAAAGTGGATT	GCTACTGCAATGCAATGGTTTAAAT
HIF-2α gDNA primers:	ACCAACCCTTCTTTCAGGCATGGC	GCTTGGTGACCTGGGCAAGTCTGC

1   Beitz, A. M., Oakes, C. G. & Galloway, K. E. Synthetic gene circuits as tools for drug
discovery. *Trends Biotechnol* **40**, 210-225 (2022).
https://doi.org/10.1016/j.tibtech.2021.06.007
2   Bock, C. *et al.* High-content CRISPR screening. *Nature Reviews Methods Primers* **2**
(2022). https://doi.org/10.1038/s43586-021-00093-4
3   Lee, T. I. & Young, R. A. Transcriptional regulation and its misregulation in disease.
Cell **152**, 1237-1251 (2013). https://doi.org/10.1016/j.cell.2013.02.014
4   Bersten, D. C., Sullivan, A. E., Peet, D. J. & Whitelaw, M. L. bHLH-PAS proteins in
cancer. *Nat Rev Cancer* **13**, 827-841 (2013). https://doi.org/10.1038/nrc3621
5   Darnell, J. E., Jr. Transcription factors as targets for cancer therapy. *Nat Rev Cancer* **2**,
740-749 (2002). https://doi.org/10.1038/nrc906
6   Sahu, B. *et al.* Sequence determinants of human gene regulatory elements. *Nat*
Genet **54**, 283-294 (2022). https://doi.org/10.1038/s41588-021-01009-4
7   Tycko, J. *et al.* High-Throughput Discovery and Characterization of Human
Transcriptional Effectors. *Cell* **183**, 2020-2035 e2016 (2020).
https://doi.org/10.1016/j.cell.2020.11.024
8   DelRosso, N. *et al.* Large-scale mapping and mutagenesis of human transcriptional
effector domains. *Nature* (2023). https://doi.org/10.1038/s41586-023-05906-y
9   Ortmann, B. M. *et al.* The HIF complex recruits the histone methyltransferase SET1B
to activate specific hypoxia-inducible genes. *Nature Genetics* **53**, 1022-1035 (2021).
https://doi.org/10.1038/s41588-021-00887-y
10   Tan, X., Letendre, J. H., Collins, J. J. & Wong, W. W. Synthetic biology in the clinic:
engineering vaccines, diagnostics, and therapeutics. *Cell* **184**, 881-898 (2021).
https://doi.org/10.1016/j.cell.2021.01.017
11   Choe, J. H. *et al.* SynNotch-CAR T cells overcome challenges of specificity,
heterogeneity, and persistence in treating glioblastoma. *Science Translational*
Medicine **13** (2021).
12   Allen, G. M. *et al.* Synthetic cytokine circuits that drive T cells into immune-excluded
tumors. *Science* **378**, 1186-+ (2022). https://doi.org/ARTN eaba1624
10.1126/science.aba1624
13   Hernandez-Lopez, R. A. *et al.* T cell circuits that sense antigen density with an
ultrasensitive threshold. *Science* **371**, 1166-+ (2021).
https://doi.org/10.1126/science.abc1855
14   Roybal, K. T. *et al.* Engineering T Cells with Customized Therapeutic Response
Programs Using Synthetic Notch Receptors. *Cell* **167**, 419-+ (2016).
https://doi.org/10.1016/j.cell.2016.09.011
15   Hasle, N. *et al.* High-throughput, microscope-based sorting to dissect cellular
heterogeneity. *Mol Syst Biol* **16**, e9442 (2020).
https://doi.org/10.15252/msb.20209442
16   Tchasovnikarova, I. A., Marr, S. K., Damle, M. & Kingston, R. E. TRACE generates
fluorescent human reporter cell lines to characterize epigenetic pathways. *Mol Cell*
(2021). https://doi.org/10.1016/j.molcel.2021.11.035
17   Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables
Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882 e1821
(2016). https://doi.org/10.1016/j.cell.2016.11.048

18   Singhal, R. & Shah, Y. M. Oxygen battle in the gut: Hypoxia and hypoxia-inducible
factors in metabolic and inflammatory responses in the intestine. *J Biol Chem* **295**,
10493-10505 (2020). https://doi.org/10.1074/jbc.REV120.011188

19   Wegiel, B., Vuerich, M., Daneshmandi, S. & Seth, P. Metabolic Switch in the Tumor
Microenvironment Determines Immune Responses to Anti-cancer Therapy. *Front*
Oncol **8**, 284 (2018). https://doi.org/10.3389/fonc.2018.00284

20   Triner, D. & Shah, Y. M. Hypoxia-inducible factors: a central link between
inflammation and cancer. *J Clin Invest* **126**, 3689-3698 (2016).
https://doi.org/10.1172/JCI84430

21   Epstien, A. *et al.* C. elegans EGL-9 and Mammalian Homologs Define a Family of
Dioxygenases that Regulate HIF by Prolyl Hydroxylation. *Cell* **107** (2001).

22   Lando, D. *et al.* FIH-1 is an asparaginyl hydroxylase enzyme that regulates the
transcriptional activity of hypoxia-inducible factor. *Genes Dev* **16**, 1466-1471 (2002).
https://doi.org/10.1101/gad.991402

23   Tian, Y.-M. *et al.* Differential Sensitivity of Hypoxia Inducible Factor Hydroxylation
Sites to Hypoxia and Hydroxylase Inhibitors. *Journal of Biological Chemistry* **286**,
13041-13051 (2011). https://doi.org/10.1074/jbc.m110.211110

24   Razorenova, O. V., Ivanov, A. V., Budanov, A. V. & Chumakov, P. M. Virus-based
reporter systems for monitoring transcriptional activity of hypoxia-inducible factor 1.
Gene **350**, 89-98 (2005). https://doi.org/10.1016/j.gene.2005.02.006

25   Villemure, J. F., Savard, N. & Belmaaza, A. Promoter suppression in cultured
mammalian cells can be blocked by the chicken beta-globin chromatin insulator
5'HS4 and matrix/scaffold attachment regions. *J Mol Biol* **312**, 963-974 (2001).
https://doi.org/10.1006/jmbi.2001.5015

26   Emerman, M. & Temin, H. Comparison of promoter suppression in avian and murine
retrovirus vectors. *Nucleic Acids Res* **14** (1986).

27   O'Connell, R. W. *et al. Ultra-high throughput mapping of genetic design space* (Cold
Spring Harbor Laboratory, 2023).

28   Lando, D., Peet, D. J., Dean A. Whelan, Jeffery J. Gorman & Whitelaw, M. L.
Asparagine Hydroxylation of the HIF Transactivation Domain: A Hypoxic Switch.
Science **295** (2002).

29   Vora, S. *et al.* Rational design of a compact CRISPR-Cas9 activator for AAV-mediated
delivery. *bioRxiv*, 298620 (2018). https://doi.org/10.1101/298620

30   Lydon, J. P. *et al.* Mice lacking progesterone receptor exhibit pleiotropic reproductive
abnormalities. *Genes & Development* **9**, 2266-2278 (1995).
https://doi.org/10.1101/gad.9.18.2266

31   Dinh, D. T. *et al.* Tissue-specific progesterone receptor-chromatin binding and the
regulation of progesterone-dependent gene expression. *Scientific Reports* **9** (2019).
https://doi.org/10.1038/s41598-019-48333-8

32   Grimm, S. L., Hartig, S. M. & Edwards, D. P. Progesterone Receptor Signaling
Mechanisms. *J Mol Biol* **428**, 3831-3849 (2016).
https://doi.org/10.1016/j.jmb.2016.06.020

33   Giannoukos, G., Szapary, D., Smith, C. L., Meeker, J. E. & Simons, S. S., Jr. New
antiprogestins with partial agonist activity: potential selective progesterone receptor
modulators (SPRMs) and probes for receptor- and coregulator-induced changes in
progesterone receptor induction properties. *Mol Endocrinol* **15**, 255-270 (2001).
https://doi.org/10.1210/mend.15.2.0596

34   Zhang, J.-H., Chung, T. & Oldenburg, K. A Simple Statistical Parameter for Use in
Evaluation and Validation of High Throughput Screening Assays. *Journal of*
Biomolecular Screening **4** (1999).
35   Kampmann, M. CRISPRi and CRISPRa Screens in Mammalian Cells for Precision
Biology and Medicine. *ACS Chem Biol* **13**, 406-416 (2018).
https://doi.org/10.1021/acschembio.7b00657
36   Jaakkola, P. *et al.* Targeting of HIF-a to the von Hippel–Lindau Ubiquitylation
Complex by O2-Regulated Prolyl Hydroxylation. *Science* **292** (2001).
37   Appelhoff, R. J. *et al.* Differential function of the prolyl hydroxylases PHD1, PHD2,
and PHD3 in the regulation of hypoxia-inducible factor. *J Biol Chem* **279**, 38458-
38465 (2004). https://doi.org/10.1074/jbc.M406026200
38   Chen, N. *et al.* Roxadustat Treatment for Anemia in Patients Undergoing Long-Term
Dialysis. *N Engl J Med* **381**, 1011-1022 (2019).
https://doi.org/10.1056/NEJMoa1901713
39   Cai, Z., Luo, W., Zhan, H. & Semenza, G. L. Hypoxia-inducible factor 1 is required for
remote ischemic preconditioning of the heart. *Proc Natl Acad Sci U S A* **110**, 17462-
17467 (2013). https://doi.org/10.1073/pnas.1317158110
40   Masoud, G. N. & Li, W. HIF-1alpha pathway: role, regulation and intervention for
cancer therapy. *Acta Pharm Sin B* **5**, 378-389 (2015).
https://doi.org/10.1016/j.apsb.2015.05.007
41   Semenza, G. L. HIF-1 mediates metabolic responses to intratumoral hypoxia and
oncogenic mutations. *J Clin Invest* **123**, 3664-3671 (2013).
https://doi.org/10.1172/JCI67230
42   Semenza, G. L. Pharmacologic Targeting of Hypoxia-Inducible Factors. *Annual Review*
of Pharmacology and Toxicology **59**, 379-403 (2019).
https://doi.org/10.1146/annurev-pharmtox-010818-021637
43   Keith, B., Johnson, R. S. & Simon, M. C. HIF1α and HIF2α: sibling rivalry in hypoxic
tumor growth and progression. *Nat Rev Cancer* **12**, 9-22 (2012).
44   Bracken, C. P. *et al.* Cell-specific regulation of hypoxia-inducible factor (HIF)-1alpha
and HIF-2alpha stabilization and transactivation in a graded oxygen environment. *J*
Biol Chem **281**, 22575-22585 (2006). https://doi.org/10.1074/jbc.M600288200
45   Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nature Protocols*
8, 2281-2308 (2013). https://doi.org/10.1038/nprot.2013.143
46   Huang, L. *et al.* Inhibitory action of Celastrol on hypoxia-mediated angiogenesis and
metastasis via the HIF-1α pathway. *International Journal of Molecular Medicine* **27**
(2011). https://doi.org/10.3892/ijmm.2011.600
47   Ma, J. *et al.* Celastrol inhibits the HIF-1α pathway by inhibition of
mTOR/p70S6K/eIF4E and ERK1/2 phosphorylation in human hepatoma cells.
Oncology Reports **32**, 235-242 (2014). https://doi.org/10.3892/or.2014.3211
48   Shang, F.-F. *et al.* Design, synthesis of novel celastrol derivatives and study on their
antitumor growth through HIF-1α pathway. *European Journal of Medicinal Chemistry*
220, 113474 (2021). https://doi.org/10.1016/j.ejmech.2021.113474
49   Srinivasan, B., Johnson, T. E. & Xing, C. Chalcone-based inhibitors against hypoxia-
inducible factor 1—Structure activity relationship studies. *Bioorganic &*
Medicinal Chemistry Letters **21**, 555-557 (2011).
https://doi.org/10.1016/j.bmcl.2010.10.063

50 Wan, C. *et al.* Genome-scale CRISPR-Cas9 screen of Wnt/β-catenin signaling
identifies therapeutic targets for colorectal cancer. *Science Advances* **7**, eabf2567
(2021). https://doi.org/10.1126/sciadv.abf2567

51 Semesta, K. M., Tian, R., Kampmann, M., Von Zastrow, M. & Tsvetanova, N. G. A
high-throughput CRISPR interference screen for dissecting functional regulators of
GPCR/cAMP signaling. *PLOS Genetics* **16**, e1009103 (2020).
https://doi.org/10.1371/journal.pgen.1009103

52 Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables
Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882.e1821
(2016). https://doi.org/10.1016/j.cell.2016.11.048

53 Potting, C. *et al.* Genome-wide CRISPR screen for PARKIN regulators reveals
transcriptional repression as a determinant of mitophagy. *Proc Natl Acad Sci U S A*
115, E180-E189 (2018). https://doi.org/10.1073/pnas.1711023115

54 Ilegems, E. *et al.* HIF-1a inhibitor PX-478 preserves pancreatic Beta cell function in
diabetes. *Science Translational Medicine* **14** (2022).

55 Koh, M. Y. *et al.* Molecular mechanisms for the activity of PX-478, an antitumor
inhibitor of the hypoxia-inducible factor-1α. *Molecular Cancer Therapeutics* **7**, 90-
100 (2008). https://doi.org/10.1158/1535-7163.mct-07-0463

56 Welsh, S., Williams, R., Kirkpatrick, L., Paine-Murrieta, G. & Powis, G. Antitumor
activity and pharmacodynamic properties of PX-478, an inhibitor of hypoxia-
inducible factor-1A. *Molecular Cancer Therapeutics* **3** (2004).
https://doi.org/https://doi.org/10.1158/1535-7163.233.3.3

57 Xia, M. *et al.* Identification of small molecule compounds that inhibit the HIF-1
signaling pathway. *Mol Cancer* **8**, 117 (2009). https://doi.org/10.1186/1476-4598-8-
117

58 Yin, J.-A. *et al.* Robust and Versatile Arrayed Libraries for Human Genome-Wide
CRISPR Activation, Deletion and Silencing. *bioRxiv*, 2022.2005.2025.493370 (2023).
https://doi.org/10.1101/2022.05.25.493370

59 Feldman, D. *et al.* Optical Pooled Screens in Human Cells. *Cell* **179**, 787-799.e717
(2019). https://doi.org/10.1016/j.cell.2019.09.016

60 Feldman, D. *et al.* Pooled genetic perturbation screens with image-based
phenotypes. *Nat Protoc* **17**, 476-512 (2022). https://doi.org/10.1038/s41596-021-
00653-8

61 Yan, X. *et al.* High-content imaging-based pooled CRISPR screens in mammalian cells.
Journal of Cell Biology **220** (2021). https://doi.org/10.1083/jcb.202008158

62 Nandagopal, N. *et al.* Dynamic Ligand Discrimination in the Notch Signaling Pathway.
Cell **172**, 869-880.e819 (2018). https://doi.org/10.1016/j.cell.2018.01.002

63 Agarwal, V. *et al.* Massively parallel characterization of transcriptional regulatory
elements in three diverse human cell types. *bioRxiv* (2023).
https://doi.org/10.1101/2023.03.05.531189

64 Gordon, M. G. *et al.* lentiMPRA and MPRAflow for high-throughput functional
characterization of gene regulatory elements. *Nat Protoc* **15**, 2387-2412 (2020).
https://doi.org/10.1038/s41596-020-0333-5

65 Bersten, D. C. *et al.* Inducible and reversible lentiviral and Recombination Mediated
Cassette Exchange (RMCE) systems for controlling gene expression. *PLoS One* **10**,
e0116373 (2015). https://doi.org/10.1371/journal.pone.0116373

66 Dinh, T. *et al.* OR08-1 Context-Specific Chromatin Binding Properties of Progesterone
Receptor and Consequential Effects on Gene Expression in Mouse Reproductive
Tissues. *J Endocr Soc* **3** (2019).
67 Singhal, H. *et al.* Genomic agonism and phenotypic antagonism between estrogen
and progesterone receptors in breast cancer. *Sci Adv* **2**, e1501924 (2016).
https://doi.org/10.1126/sciadv.1501924
68 Chen, D.-Y. *et al.* Ankyrin Repeat Proteins of Orf Virus Influence the Cellular Hypoxia
Response Pathway. *Journal of Virology* **91**, JVI.01430-01416 (2017).
https://doi.org/10.1128/jvi.01430-16
69 Dehairs, J., Talebi, A., Cherifi, Y. & Swinnen, J. V. CRISP-ID: decoding CRISPR mediated
indels by Sanger sequencing. *Sci Rep* **6**, 28973 (2016).
https://doi.org/10.1038/srep28973
70 Sanson, K. R. *et al.* Optimized libraries for CRISPR-Cas9 genetic screens with multiple
modalities. *Nat Commun* **9**, 5416 (2018). https://doi.org/10.1038/s41467-018-
07901-8
71 Bersten, D. *et al. Core and Flanking bHLH-PAS:DNA interactions mediate specificity
and drive obesity* (Cold Spring Harbor Laboratory, 2022).
72 Ritz, C., Baty, F., Streibig, J. C. & Gerhard, D. Dose-Response Analysis Using R. *PLOS
ONE* **10**, e0146021 (2016). https://doi.org/10.1371/journal.pone.0146021
73 Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series
B (Methodological)* **57**, 289-300 (1995).
https://doi.org/https://doi.org/10.1111/j.2517-6161.1995.tb02031.x
74 Becton, Dickenson & Company. (Ashland, OR, 2021).
75 Wong, F. C. *et al.* Antioxidant, Metal Chelating, Anti-glucosidase Activities and
Phytochemical Analysis of Selected Tropical Medicinal Plants. *Iran J Pharm Res* **13**,
1409-1415 (2014).
76 Wickham, H. in *Elegant Graphics for Data Analysis* VIII, 213 (Springer New York, NY,
2009).
1410