Generation and evaluation of protease inhibitor-resistant SARS-CoV-2 strains

Hawa Sophia Bouzidi[1,#], Jean-Sélim Driouich[1,#], Raphaëlle Klitting[1,#], Ornéllie Bernadin[1],

Géraldine Piorkowski[1], Rayane Amaral[1], Laurent Fraisse[2], Charles E. Mowbray[2], Ivan

Scandale[2], Fanny Escudié[2], Eric Chatelain[2], Xavier de Lamballerie[1,], Antoine Nougairède[1], and

Franck Touret[1,*].

[1]: Unité des Virus Émergents, UVE : Aix Marseille Univ, IRD 190, INSERM 1207, Marseille,

France

[2] : Drugs for Neglected Diseases initiative, Geneva, Switzerland

# These authors contributed equally to this work

*Corresponding author: franck.touret@univ-amu.fr

Abstract

11
Since the start of the SARS-CoV-2 pandemic, the search for antiviral therapies has been at the
forefront of medical research. To date, the 3CLpro inhibitor nirmatrelvir (Paxlovid®) has shown
the best results in clinical trials and the greatest robustness against variants. A second SARS-
CoV-2 protease inhibitor, ensitrelvir (Xocova®), has been developed. Ensitrelvir, currently in
Phase 3, was approved in Japan under the emergency regulatory approval procedure in
November 2022, and is available since March 31, 2023. One of the limitations for the use of
antiviral monotherapies is the emergence of resistance mutations. Here, we experimentally
generated mutants resistant to nirmatrelvir and ensitrelvir *in vitro* following repeating
passages of SARS-CoV-2 in the presence of both antivirals. For both molecules, we
demonstrated a loss of sensitivity for resistance mutants *in vitro*. Using a Syrian golden
hamster infection model, we showed that the ensitrelvir M49L mutation, in the multi-passage
strain, confers a high level of *in vivo* resistance. Finally, we identified a recent increase in the
prevalence of M49L-carrying sequences, which appears to be associated with multiple
repeated emergence events in Japan and may be related to the use of Xocova® in the country
since November 2022. These results highlight the strategic importance of genetic monitoring
of circulating SARS-CoV-2 strains to ensure that treatments administered retain their full
effectiveness.
29

Keywords

31   COVID-19, SARS-CoV-2, antiviral therapy, protease inhibitors, resistance

Introduction

Since the beginning of the SARS-CoV-2 pandemic in 2020, the search for antiviral therapeutics
has been a major focus for medical research. As previously observed with Influenza and HIV,
effective countermeasures to treat people can be decisive in managing a health crisis. At the
start of the pandemic, the repositioning of broad-spectrum antivirals targeting the replication
complex raised great expectations(Wang et al., 2020; Williamson et al., 2020), which were
dashed by disappointing results in clinical trials(Beigel et al., 2020). Therapeutic antibodies
have subsequently demonstrated their usefulness(Pinto et al., 2020; Starr et al., 2021), but

failed to overcome the emergence of variants escaping the humoral response(Touret et al.,
2023). Ultimately, the best effects in clinical trials(Hammond et al., 2022) and the greatest
robustness against variants(Vangeel et al., 2022) were achieved with the protease inhibitor
nirmatrelvir(Owen et al., 2021) (PF-07321332, Paxlovid®), targeting the 3CLpro. This inhibitor
of the coronavirus protease 3CLpro (PF-00835231) was originally developed against SARS-
CoV-1 (Boras et al., 2021). It is also effective against SARS-CoV-2 by preventing the cleavage
of the polyproteins PP1a and PP1ab – effectively blocking the generation of the non-structural
proteins essential for viral replication(Jin et al., 2020).
The pharmacokinetics (PK) of nirmatrelvir, whose metabolism includes a predominant role for
CYP3A4(Owen et al., 2021), have been improved by the combination with ritonavir used as a
exposure booster. However, this combination may complicate the use of nimatrelvir in certain
patients receiving other drugs whose metabolism may also be modified by ritonavir(Hoertel
et al., 2022).
54
In the meantime, a second SARS-CoV-2 3CLpro inhibitor named ensitrelvir (S-217622), was
developed by the Japanese company Shionogi. The molecule was identified through a
combination of virtual and biological screenings followed by optimization using a structure-
based drug design strategy(Unoh et al., 2022). It has been proposed for use without an
association with ritonavir(Unoh et al., 2022). Following encouraging results in Phase 2 clinical
trials, it is now in Phase 3(Shionogi, 2023; University of Minnesota, 2023). In Japan, however,
ensitrelvir (Xocova®), was approved under the emergency regulatory approval procedure in
November 2022 and has been available on prescription since March 31, 2023.
63
Resistance mutations can represent a major limitation to the use of antiviral mono-therapies.
Such mutations may naturally be observed in a proportion of circulating strains in the absence
of any apparent specific selective pressure(Bloom et al., 2010; Ip et al., 2023; Kawashima et
al., 2023) and/or appear in patients in the context of conditions of use associated with sub-
optimal efficiency (e.g., during prolonged use in a patient with long lasting viral replication).
For instance, the emergence of resistance against antiviral monotherapy has been observed
in patients treated with the monoclonal antibody sotrovimab during the Delta wave of SARS-
CoV-2(Rockett et al., 2022).
72
Here, we experimentally generated nirmatrelvir and ensitrelvir resistance mutants. For both
molecules, we demonstrated a loss in sensitivity for the resistance mutants *in vitro.* Using a
Syrian golden hamster model of SARS-CoV-2 infection, we showed that the ensitrelvir
resistance mutation M49L confers a high level of resistance *in vivo*. Finally, we identified a
recent increase in the prevalence of M49L-carrying sequences, that we link to multiple
repeated emergence events in Japan, and may be associated with the commercialization of
the molecule in the country since April 2023.
80

Materiel and methods

Experimental model

Cell line

84 VeroE6/TMPRSS2 cells (ID 100978) were obtained from CFAR and were grown in MEM
85 (Minimal Essential Medium-Life Technologies) with 7.5% heat-inactivated Fetal Calf Serum
86 (FCS; Life Technologies with 1% penicillin/streptomycin PS, 5000U.mL^{-1} and 5000µg.mL^{-1}
87 respectively (Life Technologies) and supplemented with 1% non-essential amino acids (Life
88 Technologies) and G-418 (Life Technologies), at 37°C with 5% CO_2.
89
90 ***Viral strain***
91 SARS-CoV-2 strain BavPat1 was obtained from Pr. C. Drosten through EVA GLOBAL
92 (https://www.european-virus-archive.com/) and contains the D614G mutation.
93
Organisms/strains
Three-week-old female Syrian hamsters (Mesocricetus auratus) were purchased from Janvier
labs. Three-week-old Syrian hamsters (Janvier labs) were maintained in ISOcage P -
Bioexclusion System (Techniplast) with unlimited access to water/food and 14h/10h
light/dark cycle. Animals were monitored and weighed daily throughout the duration of the
study to detect the appearance of any clinical signs of illness/suffering.
100
101 ***Antiviral Compounds***
102 Remdesivir was purchased from BLDpharm. ensitrelvir and nirmatrelvir were purchased
103 from MedChemexpress
104
105

Method details

RNA extraction and quantification
Viral RNA was extracted from 100 µL of cell supernatant from passages P8 and P16 using a
QIAamp Viral RNA kit on the automated QIAcube (Qiagen), following the manufacturer's
instructions. Relative quantification of viral RNA was performed using the GoTaq® 1-Step RT-
qPCR System kit (Promega). The mixture contained 5 µL of 2x Master Mix, 0.25 µL of each
primer (250 nM), 0.07 µL of probe (75 nM), 0.2 µL of GoScript RT Mix and 3.8 µL of extracted
nucleic acids. Assays were performed using the QuantStudio 12 K Flex real-time PCR machine
(Life technologies) under the following conditions : 50°C for 15 min, 95°C for 2 min, followed
by 40 cycles of 95°C for 3 s, 60°C for 30 s. Data collection took place during the 60°C step.
Synthetic RNA was used to calculate the amount of viral RNA from standard curves.
118

EC$_{50}$ determination
One day prior to infection, $5{\times}10^4$ VeroE6/TMPRSS2 cells per well were seeded in 100µL assay
medium (containing 2.5% FBS) in 96 well culture plates.The next day, antiviral compounds
were added using the D300e dispenser (TECAN) with eight ½ dilutions . Then, 50 µL/well of a
virus mix diluted in medium was added to the wells. Each well was inoculated with 100 TCID$_{50}$
of virus which correspond here to a MOI at 0.002 as classically used for SARS-CoV-2(Touret et
al., 2020a). Prior to the assay it was verified for each variant that with this MOI, viruses in the
cell culture supernatants were harvested during the logarithmic growth phase of viral
replication at 48 hours post infection (Touret et al., 2022, 2019) . Four virus control wells were
included within the plate. Quantification of the viral genome by real-time RT-qPCR as
previously described (Touret et al., 2020b).Nucleic acid from 100µL of cell supernatant were

extracted using QIAamp 96 DNA kit and Qiacube HT robot (both from Qiagen). Viral RNA was quantified by real-time RT-qPCR (GoTaq 1 step RT-qPCR kit, Promega). Quantification was provided by serial dilutions of an appropriate T7-generated synthetic RNA standard. RT-qPCR reactions were performed on QuantStudio 12K Flex Real-Time PCR System (Applied Biosystems) and analyzed using QuantStudio 12K Flex Applied Biosystems software v1.2.3. Primers and probe sequences, which target SARS-CoV-2 N gene, were: Fw: 5'-GGCCGCAAATTGCACAAT-3' ; Rev : 5'-CCAATGCGCGACATTCC-3'; Probe: 5'-FAM-CCCCCAGCGCTTCAGCGTTCT-BHQ1-3'. Viral inhibition was calculated as follow: 100* (quantity mean VC- sample quantity)/ quantity mean VC. The 50% effective concentrations (EC50 compound concentration required to inhibit viral RNA replication by 50%) were determined using logarithmic interpolation after performing a nonlinear regression (log(inhibitor) vs. response --Variable slope (four parameters)) as previously described (Touret et al., 2019, 2021a, 2022). All data obtained were analyzed using GraphPad Prism 9 software (Graphpad software).

Viral titration

One day prior to infection, 5×10^4 VeroE6 cells per well were seeded in 100µL assay medium (containing 2.5% FBS) in 96 well culture plates.Virus titration was performed using 96-well culture plates containing confluent VeroE6 cells inoculated with 150µL per well of four-fold dilutions of samples (dilution with medium supplemented with 2.5% FBS)Plates were incubate 6 days (37°C, 5% CO2) The absence or presence of cytopathic effect in each well was read.Infectious titers were estimated using the Reed & Muench method (REED and MUENCH, 1938).

Mutants Generation

One day prior to infection, 5×10^5 VeroE6/TMPRSS2 cells per well were seeded in 2mL assay medium (containing 2.5% FBS) in 12 well culture plates.The EC50s of Nirmatrelvir and Ensitrelvir were determined as previously described.Once the EC50s of these two molecules had been determined, successive passages of the virus at an MOI of 0.01 with increasing concentrations of these molecules were carried out in triplicate. Two passages at the EC50 concentration, two passages at 2x [EC50], two passages at 4 x [EC_{50}] and ten passages at 8 x [EC_{50}]. A no-molecule control was also run in triplicate. Between each run, the plates were incubated at 37°C/5% CO2 for 72 h. We performed extraction with Qiagen's EZ1 automated system and NGS sequencing (Ion Torrent) for passages P8 and P16. We then performed viral production for passage P16. Viral titration by $TCID_{50}$/mL and growth kinetics of all P16 viruses were performed.

Growth kinetics

One day prior to infection, 5×10^6 VeroE6/TMPRSS2 cells per well were seeded in 3mL assay medium (containing 2.5% FBS) in 6 well culture plates. The following day, the wells were infected with each virus with MOI 0.01, in 1mL of medium (containing 2.5% FBS), and incubated at 37°C/5% Co2 for 2h. Sampling at 0h took place before incubation. The experiment was performed in duplicate. After incubation, a wash with HBSS was performed and 3mL of medium (containing 2.5% FBS) was added to each well. Samples were taken at 6h pi, 24h pi, 30h pi, 48h pi, 54h pi and 72h pi. They were extracted by Qiacube as described above and qRT performed.

Animal experiments

In vivo experiments were approved by the local ethical committee (C2EA—14) and the
French "Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation"
(APAFIS#23975). All experiments were conducted in BSL 3 laboratory.
Three-week-old Syrian hamsters (Janvier labs) were maintained in ISOcage P - Bioexclusion
System (Techniplast) with unlimited access to water/food and 14h/10h light/dark cycle.
Animals were monitored and weighed daily throughout the duration of the study to detect
the appearance of any clinical signs of illness/suffering. Groups of animals were intranasally
infected under general anesthesia (isofluorane) with 50µL containing $2x10^4$ TCID50 of virus
diluted in 0.9% sodium chloride solution. Animal were treated via oral route two times a day
with 30 or 60 mg/kg of Ensitrelvir suspended in 0.5 % methylcellulose
Nasal washes were performed under general anesthesia (isoflurane). Blood and organs were
collected immediately after euthanasia (cervical dislocation; realized under general
anesthesia (isofluorane)).
Nasal washes were performed with 150µl 0.9% sodium chloride solution which was
transferred into 1.5mL tubes containing 0.5mL of 0.9% sodium chloride solution, then
centrifuged at 16,200g for 10 minutes and stored at -80°C. Left pulmonary lobes were washed
in 10mL of 0.9% sodium chloride solution, blotted with filter paper, weighed and then
transferred into 2mL tubes containing 1mL of 0.9% sodium chloride solution and 3mm glass
beads. They were then crushed using a Tissue Lyser machine (Retsch MM400) for 20min at 30
cycles/s and centrifuged 10 min at 16,200g. Supernatant media were transferred into 1.5mL
tubes, centrifuged 10 min at 16,200g and stored at -80°C.
Right apical lobes were collected into 2mL tubes containing 0.75mL of Qiazol lysis reagent
(Qiagen) and 3mm glass beads. They were then crushed using a Tissue Lyser machine (Retsch
MM400) for 10min at 30 cycles/s and stored at -80°C.
203

Sequence analysis of the full-length genome

200µL of lung of infectious cell supernatant (virus stock and HAE) was inactivated with an
equal volume of VXL lysis buffer (Qiagen) and viral RNA was extracted using an EZ1 Advanced
XL robot with the EZ1 mini virus 2.0 kit (both from Qiagen) and linear acrylamide
(ThermoFisher Scientific) in place of carrier RNA. The extracts were then subjected to
quantitative real-time RT-PCR in order to standardize the amounts of viral RNA used during
complete genome amplification (see below).
A specific set of primers (Table XXX) was used to generate thirteen amplicons covering the
entire genome with the Superscript IV one step RT-PCR System (ThermoFisher Scientific). PCR
mixes (final volume 25µL) contained 2.5µL containing a standard quantity of viral RNA from
the nucleic acid extract, 0.75µL of each primer (10µM), 12.5 µL of 2X Platinum SuperFi RT-PCR
Master Mix, 8.25µL of RNA free water and 0.25µL SuperScript IV RT Mix. Amplifications were
performed with the following conditions: 15 sec at 55°C, 2 min at 98°C, then 40 cycles of 10
sec at 98°C, 10 sec à 56°C and 1.5 min at 72°C. Size of PCR products was verified by gel
electrophoresis. For each sample, an equimolar pool of all amplicons was prepared and
purified using Monarch PCR & DNA Cleanup Kit (New England Biolabs). After Qubit
quantification using Qubit® dsDNA HS Assay Kit and Qubit 2.0 fluorometer (Thermo
Fisher) amplicons were sonicated (Bioruptor®, Diagenode, Liège, Belgium) into 250pb long
fragments. Libraries were built adding to fragmented DNA barcode for sample identification
and primers with Ion Plus Fragment Library Kit using AB Library Builder System (Thermo

Fisher). To pool equimolarly the barcoded samples, a real time PCR quantification step was
performed using Ion Library TaqMan™ Quantitation Kit (Thermo Fisher). Next steps included
an emulsion PCR of the pools and loading on 530 chips performed using the automated Ion
Chef instrument (Thermo Fisher), followed by sequencing using the S5 Ion torrent technology
(Thermo Fisher), following manufacturer's instructions. Consensus sequence was obtained
after trimming of reads (reads with quality score <0.99, and length <100pb were removed and
the 30 first and 30 last nucleotides were removed from the reads) and mapping of the reads
on a reference (MT594401.1) using CLC genomics workbench software v.21.0.5 (Qiagen).
Parameters for reference-based assembly consisted of match score = 1, mismatch cost = 2,
length fraction = 0.5, similarity fraction = 0.8, insertion cost = 3, and deletion cost = 3. A de
novo contig was also produced to ensure that the consensus sequence was not affected by
the reference sequence.
Mutation frequency for each position was calculated as the number of reads with a mutation
compared to the consensus sequence divided by the total number of reads at that site.
238
Public sequence data analysis
240
Datasets
We downloaded all three sets of SARS-CoV-2 sequences from the GISAID public database: (1)
all sequences (and attached metadata) containing an M to L change at position 49 in the NSP5
; (2) all EG.5.1.1 sequences (filtering out low coverage and incomplete sequences) from Japan
collected between July and October 2023 ; (3) all EG.5.1.1 sequences (filtering out low
coverage and incomplete sequences) from North America collected between July and October
2023 ; (4) all sequences (and attached metadata) containing an M to I change at position 49
in the NSP5 .
249
We first removed all sequences with >5% of ambiguous nucleotides from all datasets except
for the one corresponding to EG.5.1.1 M49L sequence. Then, to obtain a computationally
tractable dataset for phylogenetic inference, we randomly downsampled the EG.5.1.1 North
America dataset (filtering out duplicates) to 500 sequences, and combined it with all EG.5.1.1
sequences from Japan. We added all EG.5.1.1 sequences carrying the M49L mutation. After
the first maximum likelihood (ML) inference, we removed three sequences corresponding to
extremely long terminal branches (GISAID Accession IDs: 18434659, 18226914, 18432517),
yielding a final dataset of 2603 sequences. All GISAID accession numbers are available in Supp.
Table 2, the alignment and trees are available at https://github.com/rklitting/SARS-CoV-2-
resistance.
260
Phylogenetic analysis
262 For the M49L and M49I sequences lineage analysis, we assigned lineages to each of the M49L
263 sequences using the Pangolin lineage assigner (version 4.3)(O'Toole et al., 2021) .
264 Classification failed for 3 out of the 267 sequences submitted (See Supp. Table 1).
265
For phylogenetic inference based on M49L-sequences, we aligned the final EG.5.1.1 dataset
to the reference genome WIV04 (GISAID Accession: 402124) using MAFFTv7(Katoh and
Standley, 2013) . To provide some phylogenetic context to the EG.5.1.1 lineage, we added the
set of genomes corresponding to WHO reference strains for VOCs, VUIs and VUMs provided

by the Los Alamos national laboratory(Korber et al., 2020) .Finally, we performed the masking
of sites previously identified as potential sequencing errors or suspect homoplasies("Issues
with SARS-CoV-2 sequencing data - SARS-CoV-2 coronavirus / nCoV-2019 Genomic
Epidemiology," 2020), and removed the 5' and 3' UTRs. We inferred a first ML phylogeny for
this dataset using IQTREE2 under ModelFinder with ultrafast bootstrap approximation (1000
replicates)(Minh et al., 2020, 2013). Finally, we performed 10 ML inference replicates, to
control for potential instability in the topology, in particular within the EG.5.1.1 lineage.
277

Visualizations

279 The plots were created using matplotlibv3.6.3(Hunter, 2007)and the tree was visualized using
280 baltic (https://github.com/evogytis/baltic).

Results

Generation of SARS-CoV-2 strains resistant to protease inhibitors

To generate mutants resistant to SARS-CoV-2 protease inhibitors, we first determined the
experimental conditions suitable for the emergence of such resistance mutations in cell
culture. We evaluated, for both molecules, the half maximal effective concentration EC_{50} in
the VeroE6 TMPRSS2 cell line, using remdesivir as a reference compound and the BavPat1
SARS-CoV-2 strain (B.1 lineage carrying the D614G mutation). In line with the literature(Imai
et al., 2022; Sasaki et al., 2022; Takashita et al., 2022; Unoh et al., 2022), we obtained EC_{50}s of
0.14µM and 2.26µM for ensitrelvir and nirmatrelvir, respectively (Fig1a). The EC_{50} of
nirmatrelvir appears to be higher than that of ensitrelvir, as we chose to use similar conditions
for all our *in vitro* experiments and therefore did not add an efflux pump inhibitor with
nirmatrelvir, as other groups may have done (Boras et al., 2021; Owen et al., 2021).

Then, for each of the compounds, we started to grow the virus at the EC_{50}, doubling the drug
amount every two passages to increase selective pressure while allowing the virus to replicate
(Fig1b). In parallel, we performed virus passages in the absence of antiviral molecules to
identify cell line adaptation mutations. We performed the experiment with 3 replicates of
each condition. To assess the emergence of potential resistance mutations at passage 8, we
sequenced the NSP5 gene -which contains the 3CLpro region- but found no mutations. As it
was not possible to further increase the concentration of antiviral molecule in the
supernatant, we carried out 8 additional passages at the maximum concentration (8 times
higher than the EC_{50}). Throughout the experiment, we were able to maintain a stable level of
viral replication without any remarkable drop in viral load (Fig. 1c).

By sequencing full viral genomes from all experimental conditions at passage 16, we identified
mutations in the 3CLpro region, including some that were conserved across all replicates
(Fig1d).

All three replicates of the cultures performed in the presence of nirmatrelvir exhibited the T21I mutation. The other mutations -T304I, L50F, E166A- were observed in one out of three replicates. We observed T21I either alone (replicate A3), combined with T304I (A2), or combined with both L50F and E166A (A1). All four mutations have been previously described as conferring nimatrelvir resistance(Hu et al., 2023; Iketani et al., 2023; Kawashima et al., 2023).

In the case of ensitrelvir, we only observed a single mutation, M49L, conserved across the three replicates. This mutation has already been described both in clinical isolates(Ip et al., 2023; Kawashima et al., 2023) and following experiments conducted *in vitro* (Kiso et al., 2023b) and shown to reduce sensitivity to the drug.

In vitro characterization of resistant strains

To characterize our mutant strains *in vitro*, we first monitored viral replication in VeroE6 TMPRSS2 cells in the absence of compound (Fig.2a), using a low MOI (0.02), and a RT-qPCR assay.

Figure 1: Generation of nirmatrelvir and ensitrelvir resistant strains. a) Pilot experiment to determine the EC_{50s} of nirmatrelvir and ensitrelvir. b) Graphical representation of the strategy used to obtain resistant strains. c) Viral replication measured by the amount of viral RNA at each passage in each replicate. d) Table of NSP5 amino acid mutations obtained at passage 16 in the different replicates

Based on molecular viral load, we found no significant difference among the strains tested. These results are in line with the literature for T21I, T21I+T304I (Iketani et al., 2023) and M49L(Kiso et al., 2023b). They indicate that the mutations emerged in cell culture in the presence of nirmatrelvir and ensitrelvir do not alter significantly *in vitro* replication (Fig.2a)

Using a dose-response approach, we then characterized the sensitivity of these strains to the
two protease inhibitors, using remdesivir as a control. In addition to our wild-type reference
strain (BavPat), we tested one of the strains that had undergone the same number of passages
as the resistance mutants but without inhibitors (VC3). For the remdesivir control, we
obtained very similar EC_{50} values for the 8 viruses tested, with a maximum variation of 0.5µM
in their EC_{50} (Fig 2d, Table 1). This indicates that the 16 passages in culture did not alter the
sensitivity of the strains to remdesivir or their basic replication properties.

Figure 2: In vitro characterization of nirmatrelvir and ensitrelvir resistant strains. a) Kinetics of replication in VeroE6 TMPRSS2 cells Data presented are from two technical replicates and error bars show mean±s.d ; Dose response curves reporting the susceptibility of SARS-CoV-2 mutants and two wild type strain against b) ensitrelvir c) nirmatrelvir and d) remdesivir. Data presented are from three technical replicates in VeroE6-TMPRSS2 cells, and error bars show mean±s.d

In the ensitrelvir sensitivity analysis, the VC3 strain at 16 passages showed a sensitivity very
close to that of the BavPat strain, with an EC_{50} of 0.49µM versus 0.51µM. This rules out an
effect of the number of passages on sensitivity (Fig. 2b, Table 1). Strains selected in the
presence of nirmatrelvir (nirmatrelvirR) showed a slightly reduced sensitivity to ensitrelvir (3
to 9-fold). On the other hand, the three M49L ensitrelvir-resistant (ensitrelvirR) mutants
showed a great decrease in sensitivity (around 40-fold), indicating that this mutation does
confer resistance to ensitrelvir (Fig. 2b, Table 1).
In the nirmatrelvir sensitivity analysis, the VC3 strain showed a sensitivity close to that of the
BavPat strain, with an EC_{50} of 4.82µM versus 3.70µM (Fig. 2cTable 1). As before, two profiles
were observed for the mutant strains (Fig. 2cTable 1): ensitrelvirR strains showed a slightly

reduced sensitivity to nirmatrelvir (3 to 6-fold). On the other hand, the three nirmatrelvirR
mutants showed a greater decreased in sensitivity (6 to 17-fold) (Fig. 2c Table 1). Interestingly,
strains nirmatrelvirR-A2 and nirmatrelvirR-A3 showed the greatest resistance, despite having
only one and two mutations, respectively. These results confirm that the T21I mutation and
the combinations T21I-T304I and T21I-L50F-E166A do confer resistance to nirmatrelvir.

		Wild type		ensitrelvirR			nirmatrelvirR		
		Bavpat	VC3	B1	B2	B3	A1	A2	A3
nirmatrelvir	EC$_{50}$ (µM)	3.7	4.8	10.5	22.3	13.1	24.0	57.5	62.2
	EC$_{90}$ (µM)	8.4	7.6	17.3	69.5	14.7	>60	>60	>60
	Fold change	-	1.3	2.8	6.0	3.5	6.5	15.5	16.8
ensitrelvir	EC$_{50}$ (µM)	0.5	0.5	25.0	20.3	22.7	1.5	2.5	4.8
	EC$_{90}$ (µM)	0.9	2.0	39.6	36.1	37.0	4.2	3.5	6.6
	Fold change	-	1.0	49.0	39.9	44.5	3.0	5.0	9.3
remdesivir	EC$_{50}$ (µM)	4.8	4.4	3.5	3.1	2.8	2.5	4.5	3.2
	EC$_{90}$ (µM)	5.4	4.8	4.1	3.9	3.9	2.7	4.8	4.0
	Fold change	-	0.9	0.7	0.7	0.6	0.5	0.9	0.7

364 **Table 1 : Activity of ensitrelvir, nirmatrelvir and remdesivir against wild type and mutants strains**. Interpolated
365 EC$_{50}$ values are expressed in µM. Fold change reductions were calculated in comparison with the EC$_{50}$ of the
366 BavPat strain. F.c : fold change

f.c<5	5<f.c<10	10<f.c<100

367
368

Ensitrelvir is not efficacious against M49L ensitrelvir -resistant strain *in vivo*

370
Nirmatrelvir mutants have already been extensively studied(Heilmann et al., 2022; Hu et al.,
2023; Iketani et al., 2023; Kiso et al., 2023a) but the ensitrelvir resistance mutation M49L has
never been evaluated alone *in vivo* in an antiviral susceptibility assay. To assess the ability of
this mutation to confer resistance to ensitrelvir *in vivo*, we used a hamster model of SARS-
CoV-2 infection previously developed for antiviral evaluation(Driouich et al., 2021; Touret et
al., 2021b; Driouich et al., 2022). We infected groups of 6 hamsters intranasally with either
the wild type virus (BavPat), or M49L-R strain (B2). In addition to untreated groups, hamsters
were administered 30 or 60 mg/kg of ensitrelvir orally twice a day. This regimen has previously
been validated in two murine models(Kiso et al., 2023b; Unoh et al., 2022). It should be noted
that in our study, unlike the one exploring loss of sensitivity to ensitrelvir in vivo, we used the
strain from repeated passages and not generated by reverse genetics.
We observed that the M49L- ensitrlevirR strain and the wild type exhibited similar levels of
replication in lungs of untreated hamster (no statistical differences) both in terms of viral RNA
yield and infectious titers (Fig. 3 a and b). Regarding clinical follow-up, both the M49L-R strain
and the wild-type induced a similar weight loss in hamsters (Fig. 3 c and d). These results show
that, as observed *in vitro*, the selection process for resistance mutations has not deeply altered
the replication characteristics and the pathogenicity of this strain in our model.

Regarding the antiviral effect of ensitrelvir on the wild-type strain, we observed a significant
decrease in the amount of viral RNA and infectious titer in lungs, with a dose-response effect
(Fig 3 a and b). Furthermore, animals treated with 60 mg/kg had a similar clinical follow-up to
uninfected hamsters with no statistical differences in weight change, suggesting a beneficial
effect of treatment on clinical outcome (Fig 3 c and Supplemental Fig.1). Animals treated with
30 mg/kg had an intermediate clinical follow-up between untreated infected hamsters and
uninfected hamsters, with a similar trend to uninfected hamsters despite a decline between
day 1 and 2 (Fig. 3 c and Supplemental Fig.1). These results indicate significant antiviral activity
of ensitrelvir at both doses tested on the wild-type strain.
Regarding the antiviral effect of ensitrelvir on the resistant strain, we observed no significant
decrease in viral RNA levels or infectious titers in lungs compared to control, regardless of the
dosing regimen (Fig. 3 a and b). Furthermore, animals treated with 30 and 60 mg/kg had a

Figure 3: Antiviral activity of oral ensitrelvir treatment in a hamster model against wild type and resistant SARS-CoV-2 strains Groups of 6 hamsters were intranasally infected with 10^4 TCID50 of virus. Animals received ensitrelvir orally twice a day at 0 (untreated group), 30 or 60 mg/kg. a) Viral replication in lung based on viral RNA yields at day 3 post infection (measured using an RT-qPCR assay) expressed in viral genome copies/g of lung. b) Viral replication in lung based on infectious titers at day 3 post-infection (measured using a TCID50 assay) expressed in TCID50/g of lung. c-d) Clinical course of the disease (n=6 animals/group). Normalized weight at day n was calculated as follows: % of initial weight of the animal at day n. Data represent mean±SD. Two-sided statistical analysis was performed using Shapiro-Wilk normality test and Student t-test, ** and **** indicate and average significant value lower than that of the untreated group, with a p-value ranging between 0.001-0.01 and lower to 0.0001 respectively.

clinical follow-up similar to untreated infected hamsters with no statistical differences in weight change to untreated infected hamsters, suggesting no beneficial effect of treatment on clinical outcome (Fig. 3d and Supplemental Fig.1). These results indicate a lack of efficacy of ensitrelvir at both doses tested on the mutant strain containing the M49L mutation in NSP5. This result is in line with that obtained with the M49L/E166A double mutant in the same hamster model(Kiso et al., 2023b), and demonstrates the ability of M49L to confer resistance to ensitrelvir even if in the absence of E166A.

The M49L mutation increased in prevalence over the last 6 months

The emergence of resistance mutations can drastically limit the use of the antiviral therapy affected. Resistance may emerge during prolonged use of the molecule in a patient with ongoing viral replication but can also occur naturally – without any specific selective pressure due to the use of the molecule in question. Our experiments show that the M49L mutation alone can confer resistance to ensitrelvir and previous reports have described M49L in circulating SARS-CoV-2 variants(Ip et al., 2023; Kawashima et al., 2023). The prevalence of this mutation in recent months and in particular, since the commercialization of ensitrelvir in Japan, has, however, not been reported.

We therefore searched on GISAID for all SARS-CoV-2 sequences carrying the M49L mutation. We found that, as of 30th of October 2023, only 267 (0.00165%) of the 16,167,749 SARS-CoV-2 sequences available on GISAID present an M to L amino acid substitution at position 49 in the NSP5 gene. Using sequence collection dates, we evaluated the distribution over time of all M49L-carrying SARS-CoV-2 sequences. We found that M49L was first reported on the 13th August 2020, and has been observed almost monthly since then (Figure 4a), with an apparent global prevalence of 0.00168% (total number of sequences with collection date on or after 13th August 2020: 15,929,475). When looking at the most recent time span, we found that more M49L-carrying sequences have been reported between May and October 2023 (count: 154; prevalence: 0.036%) than over the previous 21 months (count: 113 M49L; prevalence: 0.0007%). These numbers indicate that while M49L corresponds to a very small proportion of the sequences observed until today, its prevalence has increased during the last 6 months.

Figure 4: M49L distribution in time, space, and across lineages. a) Global monthly counts (upper panel) and prevalences (lower panel) of M49L-sequences, based on all SARS-CoV-2 genomes publicly available on GISAID as of 2023-11-13. b) Number of M49L sequences observed by lineage, for each of the 60 Pango lineages observed most frequently among M49L sequences. c) Number of M49L sequences observed by country of collection.

The repeated observations of low numbers of M49L-carrying sequences since the first year of
the pandemic suggest that the mutation has emerged on multiple occasions and in several
SARS-CoV-2 lineages. To determine the Pango lineage of each of the 267 sequences with the
M49L mutation, we used the Pangolin online lineage assigner(O'Toole et al., 2021). We found
that M49L-carrying sequences are distributed across 90 different lineages, including Alpha,
Delta, and Omicron, with up to 31 sequences belonging to the same lineage (Figure 4b, Supp.
Table 1). The most prevalent lineages among M49L-carrying sequences are EG.5.1.1 (11.6%),
XBB.1.9.1 (5.6%), and EG.5.1 (4.9%). These three lineages are "active" (last observed less than
a year ago) and classified either as "variant of interest" (VOI) or as "variant under monitoring"
(VUM) by the WHO("Tracking SARS-CoV-2 variants," n.d.). Overall, these results indicate that
NSP5 M49L can emerge in multiple genomic backgrounds and is present in active, currently
prominent, lineages.
444
445   **The recent increase in the prevalence of M49L appears to be linked to a specific selective**
446   **pressure in Japan**
447

Figure 5: M49L distribution in time, space and, across lineages between May and October 2023 in Japan. a) Number of M49L sequences observed by country of collection for sequences collected between May and October 2023. b) Monthly counts (upper panel) and prevalences (lower panel) of M49L-sequences in Japan, based on all SARS-CoV-2 genomes publicly available on GISAID as of 2023-11-13. c) Number of recent M49L sequences (collection date between October and May 2023) observed by lineage, for each of the Pango lineages observed among recent M49L sequences.

Resistance against an antiviral therapy may occur naturally in circulating lineages but the use
of a drug may favor the emergence of resistance mutations. Using the collection dates of
M49L-carrying sequences, we showed that their proportion has been multiplied by more than
50 times in the last 6 months. Several scenarios may explain this increase in observation
frequency: (i) a specific selective pressure; (ii) a more favorable genomic background; (iii)
increased transmission.
454
The increased prevalence of the M49L resistance mutation observed in this study may be due
to the existence of specific selective pressure, *i.e.* the use of the drug to which the compound
confers resistance, ensitrelvir. The only country where this drug is used is Japan, where it first
received an emergency use approval in November 2022, and was made commercially available
in April 2023("Xocova to Be Commercially Available from March 31," n.d.). When we evaluated
the spatial distribution -in terms of country of origin- of M49L sequences, we found that Japan
is by far the most important country of origin for M49L sequences, with 59.9% of sequences
originating in Japan overall, a number that increases to 88.3% when considering only
collection dates between May and October 2023 (Figures 4c and 5a). These results indicate

that the vast majority of the M49L-carrying sequences collected over recent months originate from Japan.

When analyzing the time distribution of all M49L-sequences from Japan, we found that M49L has been observed only 4 times before December 2022 but almost monthly since then (Figure 5b), with especially high monthly prevalence values (above 0.2%) since June 2013. The timing of the increase in the prevalence of M49L and the origin of M49L sequences both align well with the use of Xocova® in Japan, suggesting that this selective pressure may favor the emergence of the mutation.

Another factor that may however, explain the recent increase in M49L sequences overall and in Japan more specifically, is the local circulation of lineages with a genomic background more favorable for the emergence of the mutation. To evaluate if recent M49L-sequences were associated with a specific genomic background, we analyzed their lineage distribution. We found that between May and October 2023, M49L occurred in 44 different lineages globally and in 40 different lineages in Japan including, EG.*, FL.*, GJ.*, HF.*, XBB.1.9.*, and XBB.1.16.* lineages (Figure 5c). These results indicate that multiple distinct lineages can accommodate the M49L mutation.

When considering M49L-sequences from the most prevalent lineage, EG.5.1.1, 93.5% of them appear to originate in Japan, and only 3.2% both in the US and in Canada. Based on GISAID data, EG.5.1.1 appears to have circulated to levels on the same order of magnitude (or greater) to that observed over the last 6 months in Japan (7.6% of 37550 total sequences), in the US (3.8% of 146709 total sequences), in Canada (6.8% of 46555 total sequences), and in China (18.2% of 34404 total sequences). If the increase in M49L prevalence was driven by the genomic background of EG.5.1.1 being more favorable, this mutation would have also emerged more frequently in other locations where the lineage has established sustained transmission. These observations indicate that the recent increase in M49L prevalence is not driven by a single -or a few- more favorable genomic background.

Repeated emergence, rather than increased transmission, contributed to the recent increase in M49L prevalence

Another phenomenon that may have participated in increasing the prevalence of M49L in recent month is the formation of large transmission chains by naturally-occurring M49L mutant. The distribution of recent M49L sequences across 44 different lineages suggests that increased emergence events, rather than increased transmission, fueled the rise in M49L detections. However, the presence of a large transmission chain within the main lineage EG.5.1.1 may have been an exacerbating factor.

To determine whether recent M49L-carrying sequences from the most prominent lineage, EG.5.1.1, correspond to a single or multiple emergence events, we performed a phylogenetic inference based on a set of publicly available SARS-CoV-2 genomes. Using a maximum-likelihood approach, we inferred the phylogenetic relationships between all M49L-carrying EG.5.1.1 sequences and a set of EG.5.1.1 sequences collected between July and October in Japan and North America (to match the spatio-temporal distribution of the M49L genomes). In the resulting phylogeny, we identified 12 different M49L emergence events, associated either with single tips, or with entire well-supported clades encompassing up to 10 sequences (Figure 6). Our findings were broadly similar across 10 inference replicates. These results show

that recent M49L-carrying sequences do not form a single clade -- the mutation emerged at least on 12 independent occasions within that lineage.

Figure 6: Maximum-likelihood phylogeny with a zoom on the EG.5.1.1 lineage. Branches and tips associated with M49L-carrying sequences are highlighted in orange. Orange arrows indicate M49L emergence events.

Discussion

In this study, we experimentally generated six viruses carrying resistance mutations to clinical stage SARS-CoV-2 protease inhibitors. We obtained three viruses with different sets of mutations for nirmatrelvir. For ensitrelvir, on the other hand, all three strains had the same mutation M49L. *In vitro* characterisation showed that nirmatrelvir-resistant strains had low resistance to ensitrelvir, but higher resistance to nirmatrelvir. Similarly, ensitrelvir-resistant strains showed low resistance to nirmatrelvir but much higher resistance to ensitrelvir. Finally, we investigated *in vivo* the resistance of an EnsitrelvirR M49L strain to two regimens of ensitrelvir and found that it completely escaped the inhibition induced by this antiviral compound.

We obtained resistance mutations after 16 passages with an increasing concentration of nirmatrelvir, which is similar to what has already been shown by Iketani and colleagues(Iketani et al., 2023). Indeed, these specific mutations have already been obtained using a similar approach: the T21I mutation was generated alone or in combination with E166V or T304I, exactly as in our study. This mutation, along with T304I, was predominantly acquired as initial mutation(Iketani et al., 2023). Interestingly, in the VeroE6 TMPRSS2 cell line, we obtained a mutation at residue 166, whereas in the study by Iketani and colleagues(Iketani et al., 2023) a mutation at this position only appeared in HUH.7 and not in VeroE6 , and the authors supposed that it was specific to this cell line. The mutation on residue 166 had already been described and was shown to reduce viral replicative fitness, but this fitness could be restored by the addition of the T21I mutation(Iketani et al., 2023). In our study, we confirmed this observation by showing that the replication capacity of our mutant (A1) carrying both T21I and E166A was the same as that of the wild type. Finally the mutation at residue 166 was the only one obtained in the substrate binding site; the others are situated at a distal position from nirmatrelvir predicted binding site (over 5 Å(Iketani et al., 2023)).

For ensitrelvir, we obtained three viruses with the same mutation, M49L. This mutation has already been obtained in a similar experiment, but with a shorter passage number(Kiso et al., 2023a). It is also present in circulating isolates of SARS-CoV-2 (Hu et al., 2023; Kawashima et al., 2023; Moghadasi et al., 2023b; Noske et al., 2023).
In a report identifying M49I as an ensitrelvir resistance mutation its visualization in complex with the 3CLpro shows that the greater hydrophobicity of isoleucine, compared to the methionine, caused the inhibitor to move to another site, which affects its antiviral activity(Noske et al., 2023). According to another study, the presence of a leucine instead of an isoleucine at this position could cause a greater resistance due the gamma-carbon branching of the leucine side chain, which is nearer to H41 one of the catalytic dyad in its active site (Moghadasi et al., 2023b; Unoh et al., 2022).
We found *in vitro* moderate cross resistance between NirmatrelvirR strains and ensitrelvir, and the same for EnsitrelvirR strains and Nirmatrelvir, in accordance with previous studies(Iketani et al., 2023; Moghadasi et al., 2023b). This suggests that replacing one of the two compounds with the other could compensate for the emergence of a resistance mutation during treatment.

We observed the greatest loss in activity with the couple ensitrelvir/M49L with a ~40-fold reduction in susceptibility for our three ensitrelvirR strains. This mutation had already been

tested, alone, in two distinct enzymatic assays(Moghadasi et al., 2023b, 2023a) and in a VSV based cleavage system(Moghadasi et al., 2023b), with observed fold change values of 21.4, 25.4 and 10 respectively. In the only previous study that explored the effect of this mutation with a replicative virus (a mutated Delta strain generated by reverse genetic(Kiso et al., 2023b)) M49L was associated with a 37.4 fold change in a focus reduction assay(Kiso et al., 2023a). This result is perfectly in line with our observation with the same range of reduction despite the use of the strain generated from repeated passages in a different variant backbone. The fact that we found the same effect for a single amino acid mutation but with a different backbone not only confirms but reinforces our result.

In the abovementioned study, the authors only studied the *in vivo* antiviral resistance of the M49L/166A mutation pair(Kiso et al., 2023b)*. Regarding the single mutation M49L, they only explored its impact on the *in vitro* replication fitness and the *in vivo* pathogenicity. We went a step further and explored the effect of this single mutation on ensitrelvir sensitivity, using both the ensitrelvir dose referred to in the study above (60mg/kg twice daily)(Kiso et al., 2023b) and a lower dose as in the original S-217622/ensitrelvir article (30mg/kg twice daily)(Unoh et al., 2022). When animals were infected with the ensitrelvirR strain, we observed no antiviral activity for either of ensitrelvir dose regimen and, in contrast to what was observed with the wild-type strain, no clinical improvement. This implies that the M49L mutation can be solely responsible for *in vivo* resistance. The same result at the highest regimen was observed *in vivo* with the double mutants M49L/E166A(Kiso et al., 2023b). The *in vitro* fold change of this combination is almost 200(Kiso et al., 2023b). Our result implies that a single mutation inducing a change in susceptibility of 40, *i.e.* five times lower, is enough to make ensitrelvir treatment ineffective in our animal model.

In the current study, we detailed a recent increase in the prevalence of M49L-carrying sequences, linked it to a specific region, Japan, and showed that this increase is concomitant with the use of ensitrelvir in the country (emergency approval and then commercialization) rather than with the emergence of a more favorable viral genomic background. These initial results are concerning and warrant further investigation, especially as several countries are currently considering the drug for emergency approval("Drugmaker Shionogi inks deal to market COVID pill in China," n.d.; "Xocova Filed for Conditional Approval in South Korea," n.d.). It is crucial to better characterize the conditions and frequency of emergence of the M49L mutation in patients treated with ensitrelvir. If an increased risk of resistance emergence during patient treatment is confirmed, genomic monitoring may allow to both optimize the medical efficacy of the treatment provided and prevent the spread of resistant strains. In case of resistance, it would still be possible to replace ensitrelvir with nirmatrelvir, given the low level of cross-resistance between the two molecules. However, this would be difficult for patients who had received ensitrelvir treatment in the first instance, due to a contraindication to ritonavir, which is present in Paxlovid®, the pharmaceutical formulation of nirmatrelvir.

To evaluate the potential duration of transmission for EG.5.1.1 M49L-mutants, we calculated minimum clade-durations for all M49L-clades by calculating the difference in time between the earliest and latest sampling dates among sequences of a given clade. We found that clade-duration could be up to 58 days for M49L clades belonging to the EG.5.1.1 lineage. This first assessment however, is based on a small number of clades (3) and on a limited sampling of the recent circulating diversity of SARS-CoV-2. For computational tractability reasons we focused our phylogenetic analysis on sequences from a single lineage (EG.5.1.1), specific countries (Japan, North America), and subsampled our dataset. We believe that it is of the

612 utmost importance to pursue these investigations and to continue to monitor the ability of
613 M49L-carrying lineages to establish sustained transmission and to assess the risk of fixation of
614 the mutation.
615
Finally, while we showed that the use of ensitrelvir is probably a driver of the increased
frequency of emergence of M49L in recent months in Japan, it is likely that the 107 early
sequences carrying M49L (identified between August 2020 and October 2023) emerged
spontaneously as the antiviral was not in use before November 2022. The natural occurrence
of strains carrying M49L may thus facilitate the emergence of resistance to ensitrelvir.
Other variations in the NSP5 protein provide resistance to ensitrelvir and have been observed
in circulating isolates. They may present a risk of increased emergence in case of wide use of
ensitrelvir, akin to what was showed for M49L in the present study. In particular, the mutation
M49I provides resistance *in vitro* (Moghadasi et al., 2023b) and is present in 2188 SARS-CoV-
2 sequences on GISAID (as of October 30th). For this mutation however, we found no apparent
increase in prevalence over recent months in Japan (Supplemental Figure 2a) with most M49I
sequences observed between June and December 2021. Along with the fact that M49I has
never been observed in experiments of generation of ensitrelvir-resistant mutants (current
study and in (Kiso et al., 2023b), this finding suggests that under selective pressure, the M49L
mutation is preferred, likely because it provides greater resistance by generating a branching
pattern more favorable to the disruption of a base stacking interaction with the inhibitor
(Moghadasi et al., 2023b; Noske et al., 2023). In addition, the lower prevalence of M49I over
recent months in Japan compared to M49L (Supplemental Figure 2b) may be due to less
favorable circulating genetic backgrounds, as most M49I sequences are associated with
lineages AY.4 (714/2188 sequences) and AY.44 (172/2188), and circulated during the pre-
Omicron period (Supplemental Figures 2a and 2c). These results suggest the risk of emergence
of resistance associated with M49I is currently lower than for M49L, but other mutations, in
the 45-49 and other regions of M^{pro}, require continuous monitoring in countries where
ensitrelvir is, or will soon be, approved for use.
640
These findings argues in favor of genetic monitoring of circulating strains to ensure that the
treatments administered retain their full effectiveness. In addition, the range of antiviral
treatments available against SARS-CoV-2 has become worryingly limited, with most
commercial monoclonal drugs losing their activity against the most recent variants. Obviously,
there is still a need to develop other antivirals targeting other viral proteins to broaden the
therapeutic armory, and possibly to develop dual therapies as has been done for other viral
diseases.
648

Acknowledgments

We thank Magali Gilles (UVE; Marseille), Gregory Moureau (UVE; Marseille) and Camille Placidi
(UVE; Marseille) for their valuable technical contribution. We thank Edyth Parker for her
recommendations on SARS-CoV-2 phylogenetic inference. We thank Pr. Drosten for providing
the SARSCoV-2 strain through the European Research infrastructure EVA GLOBAL.
This work was supported by the European Virus Archive Global (EVA GLOBAL) funded by the
European Union's Horizon 2020 research and innovation program under grant agreement No.
871029 and by DNDi under support by the Wellcome Trust Grant ref: 222489/Z/21/Z through
the COVID-19 Therapeutics Accelerator. This work was supported by the ARBOGEN project,

funded by the MSDAVENIR foundation. Part of the work was done on the Aix Marseille
University antivirals platform "MaSC".
660

Author Contributions
Conceptualization : AN, FE, IS, EC and FT
Methodology :, JSD, AN, and FT
Formal analysis : SHB, JSD, RK, OB,GP,RA and FT
Investigation : SHB, JSD,RK, GP ,RA, OBand FT
Resources : XdL, FT and AN
Writing original draft: SHB, JSD, AN and FT;
Writing review & editing : all co-authors
Visualization SHB, JSD, RK, OB, AN and FT
Funding acquisition LF, CM, EC, XdL, AN and FT.
671

Declaration of interests
673 The authors declare that they have no conflict of interest.
674

Supplemental Data
Supplemental figure 1: Clinical follow up after Wild-type (Bavpat) and EnsitrelvirR B2 SARS-
CoV-2 viral challenge. a-b : Weight change at day 2 and at day 3 post infection in percentage,
normalized to the body weight at day 0 of B.1 (G614) infection. C-D : Weight change at day 2
and at day 3 post infection in percentage, normalized to the body weight at day 0 of
EnsitelvirR B2 infection. Bars represent means ± SD. The data were analyzed with the
unpaired t test (fig.A-C) and Mann-Whitney test (fig b-d). *P < .05, **P < .01, ***P < .001.
Supplemental figure 2: M49I distribution in time, Japan, and across lineages. a) Global
monthly counts (upper panel) and prevalences (lower panel) of M49I-sequences, based on
all SARS-CoV-2 genomes publicly available on GISAID as of 2023-11-13. b) Monthly counts
(upper panel) and prevalences (lower panel) of M49I-sequences in Japan, based on all SARS-
CoV-2 genomes publicly available on GISAID as of 2023-11-13. c) Number of M49I sequences
observed by lineage
Supplementary Table 1: Accession IDs, collection dates, submission dates, collection
location, and Pango lineage (assigned using Pangolin lineage assigner) for all M49L
sequences analyzed in this study.
691
692 **Supplementary Table 2:** GISAID accession IDs for all SARS-CoV-2 genomes (and attached
693 metadata) in the M49L analysis.
694
695 **Supplementary Table 3**: Accession IDs, collection dates, submission dates, collection
696 location, and Pango lineage (assigned using Pangolin lineage assigner) for all M49I sequences
697 analyzed in this study.
698 **Supplementary Table 4**: GISAID accession IDs for all SARS-CoV-2 genomes (and attached
699 metadata) in the M49I analysis.
700
701

References
703

704 Beigel, J.H., Tomashek, K.M., Dodd, L.E., Mehta, A.K., Zingman, B.S., Kalil, A.C., Hohmann, E., Chu, H.Y.,
705   Luetkemeyer, A., Kline, S., Lopez de Castilla, D., Finberg, R.W., Dierberg, K., Tapson, V., Hsieh, L.,
706   Patterson, T.F., Paredes, R., Sweeney, D.A., Short, W.R., Touloumi, G., Lye, D.C., Ohmagari, N., Oh, M.,
707   Ruiz-Palacios, G.M., Benfield, T., Fätkenheuer, G., Kortepeter, M.G., Atmar, R.L., Creech, C.B.,
708   Lundgren, J., Babiker, A.G., Pett, S., Neaton, J.D., Burgess, T.H., Bonnett, T., Green, M., Makowski, M.,
709   Osinusi, A., Nayak, S., Lane, H.C., 2020. Remdesivir for the Treatment of Covid-19 — Final Report. New
710   England Journal of Medicine 383, 1813–1826. https://doi.org/10.1056/NEJMoa2007764
711 Bloom, J.D., Gong, L.I., Baltimore, D., 2010. Permissive Secondary Mutations Enable the Evolution of Influenza
712   Oseltamivir Resistance. Science 328, 1272–1275. https://doi.org/10.1126/science.1187816
713 Boras, B., Jones, R.M., Anson, B.J., Arenson, D., Aschenbrenner, L., Bakowski, M.A., Beutler, N., Binder, J., Chen,
714   E., Eng, H., Hammond, H., Hammond, J., Haupt, R.E., Hoffman, R., Kadar, E.P., Kania, R., Kimoto, E.,
715   Kirkpatrick, M.G., Lanyon, L., Lendy, E.K., Lillis, J.R., Logue, J., Luthra, S.A., Ma, C., Mason, S.W.,
716   McGrath, M.E., Noell, S., Obach, R.S., O' Brien, M.N., O'Connor, R., Ogilvie, K., Owen, D., Pettersson,
717   M., Reese, M.R., Rogers, T.F., Rosales, R., Rossulek, M.I., Sathish, J.G., Shirai, N., Steppan, C., Ticehurst,
718   M., Updyke, L.W., Weston, S., Zhu, Y., White, K.M., García-Sastre, A., Wang, J., Chatterjee, A.K.,
719   Mesecar, A.D., Frieman, M.B., Anderson, A.S., Allerton, C., 2021. Preclinical characterization of an
720   intravenous coronavirus 3CL protease inhibitor for the potential treatment of COVID19. Nat Commun
721   12, 6055. https://doi.org/10.1038/s41467-021-26239-2
722 Driouich, J.-S., Cochin, M., Lingas, G., Moureau, G., Touret, F., Petit, P.-R., Piorkowski, G., Barthélémy, K., Laprie,
723   C., Coutard, B., Guedj, J., de Lamballerie, X., Solas, C., Nougairède, A., 2021. Favipiravir antiviral
724   efficacy against SARS-CoV-2 in a hamster model. Nat Commun 12, 1735.
725   https://doi.org/10.1038/s41467-021-21992-w
726 Driouich, J.-S., Cochin, M., Touret, F., Petit, P.-R., Gilles, M., Moureau, G., Barthélémy, K., Laprie, C.,
727   Wattanakul, T., Chotsiri, P., Hoglund, R.M., Tarning, J., Fraisse, L., Sjö, P., Mowbray, C.E., Escudié, F.,
728   Scandale, I., Chatelain, E., de Lamballerie, X., Solas, C., Nougairède, A., 2022. Pre-clinical evaluation of
729   antiviral activity of nitazoxanide against SARS-CoV-2. eBioMedicine 82, 104148.
730   https://doi.org/10.1016/j.ebiom.2022.104148
731 Drugmaker Shionogi inks deal to market COVID pill in China [WWW Document], n.d. . Nikkei Asia. URL
732   https://asia.nikkei.com/Business/Pharmaceuticals/Drugmaker-Shionogi-inks-deal-to-market-COVID-
733   pill-in-China (accessed 11.14.23).
734 Hammond, J., Leister-Tebbe, H., Gardner, A., Abreu, P., Bao, W., Wisemandle, W., Baniecki, M., Hendrick, V.M.,
735   Damle, B., Simón-Campos, A., Pypstra, R., Rusnak, J.M., 2022. Oral Nirmatrelvir for High-Risk,
736   Nonhospitalized Adults with Covid-19. New England Journal of Medicine 386, 1397–1408.
737   https://doi.org/10.1056/NEJMoa2118542
738 Heilmann, E., Costacurta, F., Moghadasi, S.A., Ye, C., Pavan, M., Bassani, D., Volland, A., Ascher, C., Weiss,
739   A.K.H., Bante, D., Harris, R.S., Moro, S., Rupp, B., Martinez-Sobrido, L., von Laer, D., 2022. SARS-CoV-2
740   3CLpro mutations selected in a VSV-based system confer resistance to nirmatrelvir, ensitrelvir, and
741   GC376. Science Translational Medicine 15, eabq7360. https://doi.org/10.1126/scitranslmed.abq7360
742 Hoertel, N., Boulware, D.R., Sánchez-Rico, M., Burgun, A., Limosin, F., 2022. Prevalence of Contraindications to
743   Nirmatrelvir-Ritonavir Among Hospitalized Patients With COVID-19 at Risk for Progression to Severe
744   Disease. JAMA Network Open 5, e2242140. https://doi.org/10.1001/jamanetworkopen.2022.42140
745 Hu, Y., Lewandowski, E.M., Tan, H., Zhang, Xiaoming, Morgan, R.T., Zhang, Xiujun, Jacobs, L.M.C., Butler, S.G.,
746   Gongora, M.V., Choy, J., Deng, X., Chen, Y., Wang, J., 2023. Naturally Occurring Mutations of SARS-
747   CoV-2 Main Protease Confer Drug Resistance to Nirmatrelvir. ACS Cent. Sci. 9, 1658–1669.
748   https://doi.org/10.1021/acscentsci.3c00538
749 Hunter, J.D., 2007. Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering 9, 90–95.
750   https://doi.org/10.1109/MCSE.2007.55
751 Iketani, S., Mohri, H., Culbertson, B., Hong, S.J., Duan, Y., Luck, M.I., Annavajhala, M.K., Guo, Y., Sheng, Z.,
752   Uhlemann, A.-C., Goff, S.P., Sabo, Y., Yang, H., Chavez, A., Ho, D.D., 2023. Multiple pathways for SARS-
753   CoV-2 resistance to nirmatrelvir. Nature 613, 558–564. https://doi.org/10.1038/s41586-022-05514-2
754 Imai, M., Ito, M., Kiso, M., Yamayoshi, S., Uraki, R., Fukushi, S., Watanabe, S., Suzuki, T., Maeda, K., Sakai-
755   Tagawa, Y., Iwatsuki-Horimoto, K., Halfmann, P.J., Kawaoka, Y., 2022. Efficacy of Antiviral Agents
756   against Omicron Subvariants BQ.1.1 and XBB. New England Journal of Medicine 0, null.
757   https://doi.org/10.1056/NEJMc2214302
758 Ip, J.D., Wing-Ho Chu, A., Chan, W.-M., Cheuk-Ying Leung, R., Umer Abdullah, S.M., Sun, Y., To, K.K.-W., 2023.
759   Global prevalence of SARS-CoV-2 3CL protease mutations associated with nirmatrelvir or ensitrelvir
760   resistance. eBioMedicine 91, 104559. https://doi.org/10.1016/j.ebiom.2023.104559

Issues with SARS-CoV-2 sequencing data - SARS-CoV-2 coronavirus / nCoV-2019 Genomic Epidemiology [WWW
Document], 2020. . Virological. URL https://virological.org/t/issues-with-sars-cov-2-sequencing-
data/473/1 (accessed 11.14.23).
Jin, Z., Du, X., Xu, Y., Deng, Y., Liu, M., Zhao, Y., Zhang, B., Li, X., Zhang, L., Peng, C., Duan, Y., Yu, J., Wang, L.,
Yang, K., Liu, F., Jiang, R., Yang, Xinglou, You, T., Liu, Xiaoce, Yang, Xiuna, Bai, F., Liu, H., Liu, Xiang,
Guddat, L.W., Xu, W., Xiao, G., Qin, C., Shi, Z., Jiang, H., Rao, Z., Yang, H., 2020. Structure of Mpro from
SARS-CoV-2 and discovery of its inhibitors. Nature 582, 289–293. https://doi.org/10.1038/s41586-020-
2223-y
Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in
Performance and Usability. Molecular Biology and Evolution 30, 772–780.
https://doi.org/10.1093/molbev/mst010
Kawashima, S., Matsui, Y., Adachi, T., Morikawa, Y., Inoue, K., Takebayashi, S., Nobori, H., Rokushima, M.,
Tachibana, Y., Kato, T., 2023. Ensitrelvir is effective against SARS-CoV-2 3CL protease mutants
circulating globally. Biochem Biophys Res Commun 645, 132–136.
https://doi.org/10.1016/j.bbrc.2023.01.040
Kiso, M., Furusawa, Y., Uraki, R., Imai, M., Yamayoshi, S., Kawaoka, Y., 2023a. In vitro and in vivo
characterization of SARS-CoV-2 strains resistant to nirmatrelvir. Nat Commun 14, 3952.
https://doi.org/10.1038/s41467-023-39704-x
Kiso, M., Yamayoshi, S., Iida, S., Furusawa, Y., Hirata, Y., Uraki, R., Imai, M., Suzuki, T., Kawaoka, Y., 2023b. In
vitro and in vivo characterization of SARS-CoV-2 resistance to ensitrelvir. Nat Commun 14, 4231.
https://doi.org/10.1038/s41467-023-40018-1
Korber, B., Fischer, W.M., Gnanakaran, S., Yoon, H., Theiler, J., Abfalterer, W., Hengartner, N., Giorgi, E.E.,
Bhattacharya, T., Foley, B., Hastie, K.M., Parker, M.D., Partridge, D.G., Evans, C.M., Freeman, T.M., de
Silva, T.I., Angyal, A., Brown, R.L., Carrilero, L., Green, L.R., Groves, D.C., Johnson, K.J., Keeley, A.J.,
Lindsey, B.B., Parsons, P.J., Raza, M., Rowland-Jones, S., Smith, N., Tucker, R.M., Wang, D., Wyles,
M.D., McDanal, C., Perez, L.G., Tang, H., Moon-Walker, A., Whelan, S.P., LaBranche, C.C., Saphire, E.O.,
Montefiori, D.C., 2020. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases
Infectivity of the COVID-19 Virus. Cell 182, 812-827.e19. https://doi.org/10.1016/j.cell.2020.06.043
Minh, B.Q., Nguyen, M.A.T., von Haeseler, A., 2013. Ultrafast Approximation for Phylogenetic Bootstrap.
Molecular Biology and Evolution 30, 1188–1195. https://doi.org/10.1093/molbev/mst024
Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., Lanfear, R., 2020.
IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era.
Molecular Biology and Evolution 37, 1530–1534. https://doi.org/10.1093/molbev/msaa015
Moghadasi, S.A., Biswas, R.G., Harki, D.A., Harris, R.S., 2023a. Rapid resistance profiling of SARS-CoV-2 protease
inhibitors. npj Antimicrob Resist 1, 1–4. https://doi.org/10.1038/s44259-023-00009-0
Moghadasi, S.A., Heilmann, E., Khalil, A.M., Nnabuife, C., Kearns, F.L., Ye, C., Moraes, S.N., Costacurta, F., Esler,
M.A., Aihara, H., Harris, R.S., 2023b. Transmissible SARS-CoV-2 variants with resistance to clinical
protease inhibitors. SCIENCE ADVANCES.
Noske, G.D., Silva, E. de S., Godoy, M.O. de, Dolci, I., Fernandes, R.S., Guido, R.V.C., Sjö, P., Oliva, G., Godoy,
A.S., 2023. Structural basis of nirmatrelvir and ensitrelvir activity against naturally occurring
polymorphisms of the SARS-CoV-2 main protease. Journal of Biological Chemistry 299.
https://doi.org/10.1016/j.jbc.2023.103004
O'Toole, Á., Scher, E., Underwood, A., Jackson, B., Hill, V., McCrone, J.T., Colquhoun, R., Ruis, C., Abu-Dahab, K.,
Taylor, B., Yeats, C., du Plessis, L., Maloney, D., Medd, N., Attwood, S.W., Aanensen, D.M., Holmes,
E.C., Pybus, O.G., Rambaut, A., 2021. Assignment of epidemiological lineages in an emerging pandemic
using the pangolin tool. Virus Evol 7, veab064. https://doi.org/10.1093/ve/veab064
Owen, D.R., Allerton, C.M.N., Anderson, A.S., Aschenbrenner, L., Avery, M., Berritt, S., Boras, B., Cardin, R.D.,
Carlo, A., Coffman, K.J., Dantonio, A., Di, L., Eng, H., Ferre, R., Gajiwala, K.S., Gibson, S.A., Greasley,
S.E., Hurst, B.L., Kadar, E.P., Kalgutkar, A.S., Lee, J.C., Lee, J., Liu, W., Mason, S.W., Noell, S., Novak, J.J.,
Obach, R.S., Ogilvie, K., Patel, N.C., Pettersson, M., Rai, D.K., Reese, M.R., Sammons, M.F., Sathish, J.G.,
Singh, R.S.P., Steppan, C.M., Stewart, A.E., Tuttle, J.B., Updyke, L., Verhoest, P.R., Wei, L., Yang, Q.,
Zhu, Y., 2021. An oral SARS-CoV-2 Mpro inhibitor clinical candidate for the treatment of COVID-19.
Science 374, 1586–1593. https://doi.org/10.1126/science.abl4784
Pinto, D., Park, Y.-J., Beltramello, M., Walls, A.C., Tortorici, M.A., Bianchi, S., Jaconi, S., Culap, K., Zatta, F., De
Marco, A., Peter, A., Guarino, B., Spreafico, R., Cameroni, E., Case, J.B., Chen, R.E., Havenar-Daughton,
C., Snell, G., Telenti, A., Virgin, H.W., Lanzavecchia, A., Diamond, M.S., Fink, K., Veesler, D., Corti, D.,

2020. Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody. Nature 583,
290–295. https://doi.org/10.1038/s41586-020-2349-y

REED, L.J., MUENCH, H., 1938. A SIMPLE METHOD OF ESTIMATING FIFTY PER CENT ENDPOINTS12. American
Journal of Epidemiology 27, 493–497. https://doi.org/10.1093/oxfordjournals.aje.a118408

Rockett, R., Basile, K., Maddocks, S., Fong, W., Agius, J.E., Johnson-Mackinnon, J., Arnott, A., Chandra, S., Gall,
M., Draper, J., Martinez, E., Sim, E.M., Lee, C., Ngo, C., Ramsperger, M., Ginn, A.N., Wang, Q., Fennell,
M., Ko, D., Lim, H.L., Gilroy, N., O'Sullivan, M.V.N., Chen, S.C.-A., Kok, J., Dwyer, D.E., Sintchenko, V.,
2022. Resistance Mutations in SARS-CoV-2 Delta Variant after Sotrovimab Use. N Engl J Med 386,
1477–1479. https://doi.org/10.1056/NEJMc2120219

Sasaki, M., Tabata, K., Kishimoto, M., Itakura, Y., Kobayashi, H., Ariizumi, T., Uemura, K., Toba, S., Kusakabe, S.,
Maruyama, Y., Iida, S., Nakajima, N., Suzuki, T., Yoshida, S., Nobori, H., Sanaki, T., Kato, T., Shishido, T.,
Hall, W.W., Orba, Y., Sato, A., Sawa, H., 2022. S-217622, a SARS-CoV-2 main protease inhibitor,
decreases viral load and ameliorates COVID-19 severity in hamsters. Science Translational Medicine
15, eabq4064. https://doi.org/10.1126/scitranslmed.abq4064

Shionogi, 2023. A Phase 3, Multicenter, Randomized, Double-Blind, 24-Week Study of the Clinical and Antiviral
Effect of S-217622 Compared With Placebo in Non-Hospitalized Participants With COVID-19 (Clinical
trial registration No. NCT05305547). clinicaltrials.gov.

Starr, T.N., Czudnochowski, N., Liu, Z., Zatta, F., Park, Y.-J., Addetia, A., Pinto, D., Beltramello, M., Hernandez, P.,
Greaney, A.J., Marzi, R., Glass, W.G., Zhang, I., Dingens, A.S., Bowen, J.E., Tortorici, M.A., Walls, A.C.,
Wojcechowskyj, J.A., De Marco, A., Rosen, L.E., Zhou, J., Montiel-Ruiz, M., Kaiser, H., Dillen, J.R.,
Tucker, H., Bassi, J., Silacci-Fregni, C., Housley, M.P., di Iulio, J., Lombardo, G., Agostini, M., Sprugasci,
N., Culap, K., Jaconi, S., Meury, M., Dellota Jr, E., Abdelnabi, R., Foo, S.-Y.C., Cameroni, E., Stumpf, S.,
Croll, T.I., Nix, J.C., Havenar-Daughton, C., Piccoli, L., Benigni, F., Neyts, J., Telenti, A., Lempp, F.A.,
Pizzuto, M.S., Chodera, J.D., Hebner, C.M., Virgin, H.W., Whelan, S.P.J., Veesler, D., Corti, D., Bloom,
J.D., Snell, G., 2021. SARS-CoV-2 RBD antibodies that maximize breadth and resistance to escape.
Nature 597, 97–102. https://doi.org/10.1038/s41586-021-03807-6

Takashita, E., Kinoshita, N., Yamayoshi, S., Sakai-Tagawa, Y., Fujisaki, S., Ito, M., Iwatsuki-Horimoto, K.,
Halfmann, P., Watanabe, S., Maeda, K., Imai, M., Mitsuya, H., Ohmagari, N., Takeda, M., Hasegawa, H.,
Kawaoka, Y., 2022. Efficacy of Antiviral Agents against the SARS-CoV-2 Omicron Subvariant BA.2. N
Engl J Med 386, 1475–1477. https://doi.org/10.1056/NEJMc2201933

Touret, F., Baronti, C., Bouzidi, H.S., de Lamballerie, X., 2022. In vitro evaluation of therapeutic antibodies
against a SARS-CoV-2 Omicron B.1.1.529 isolate. Sci Rep 12, 4683. https://doi.org/10.1038/s41598-
022-08559-5

Touret, F., Baronti, C., Goethals, O., Van Loock, M., de Lamballerie, X., Querat, G., 2019. Phylogenetically based
establishment of a dengue virus panel, representing all available genotypes, as a tool in dengue drug
discovery. Antiviral Res. 168, 109–113. https://doi.org/10.1016/j.antiviral.2019.05.005

Touret, F., Driouich, J.-S., Cochin, M., Petit, P.R., Gilles, M., Barthélémy, K., Moureau, G., Mahon, F.-X., Malvy,
D., Solas, C., de Lamballerie, X., Nougairède, A., 2021a. Preclinical evaluation of Imatinib does not
support its use as an antiviral drug against SARS-CoV-2. Antiviral Research 193, 105137.
https://doi.org/10.1016/j.antiviral.2021.105137

Touret, F., Driouich, J.-S., Cochin, M., Petit, P.R., Gilles, M., Barthélémy, K., Moureau, G., Mahon, F.-X., Malvy,
D., Solas, C., de Lamballerie, X., Nougairède, A., 2021b. Preclinical evaluation of Imatinib does not
support its use as an antiviral drug against SARS-CoV-2. Antiviral Res 193, 105137.
https://doi.org/10.1016/j.antiviral.2021.105137

Touret, F., Gilles, M., Barral, K., Nougairède, A., van Helden, J., Decroly, E., de Lamballerie, X., Coutard, B.,
2020a. In vitro screening of a FDA approved chemical library reveals potential inhibitors of SARS-CoV-2
replication. Sci Rep 10, 13093. https://doi.org/10.1038/s41598-020-70143-6

Touret, F., Gilles, M., Barral, K., Nougairède, A., van Helden, J., Decroly, E., de Lamballerie, X., Coutard, B.,
2020b. In vitro screening of a FDA approved chemical library reveals potential inhibitors of SARS-CoV-2
replication. Scientific Reports 10, 13093. https://doi.org/10.1038/s41598-020-70143-6

Touret, F., Giraud, E., Bourret, J., Donati, F., Tran-Rajau, J., Chiaravalli, J., Lemoine, F., Agou, F., Simon-Lorière,
E., van der Werf, S., de Lamballerie, X., 2023. Enhanced neutralization escape to therapeutic
monoclonal antibodies by SARS-CoV-2 omicron sub-lineages. iScience 26, 106413.
https://doi.org/10.1016/j.isci.2023.106413

Tracking SARS-CoV-2 variants [WWW Document], n.d. URL https://www.who.int/activities/tracking-SARS-CoV-
2-variants (accessed 11.14.23).

University of Minnesota, 2023. Strategies and Treatments for Respiratory Infections &Amp; Viral Emergencies
(STRIVE): Shionogi Protease Inhibitor (Clinical trial registration No. NCT05605093). clinicaltrials.gov.
Unoh, Y., Uehara, S., Nakahara, K., Nobori, H., Yamatsu, Y., Yamamoto, S., Maruyama, Y., Taoda, Y., Kasamatsu,
K., Suto, T., Kouki, K., Nakahashi, A., Kawashima, S., Sanaki, T., Toba, S., Uemura, K., Mizutare, T.,
Ando, S., Sasaki, M., Orba, Y., Sawa, H., Sato, A., Sato, T., Kato, T., Tachibana, Y., 2022. Discovery of S-
217622, a Noncovalent Oral SARS-CoV-2 3CL Protease Inhibitor Clinical Candidate for Treating COVID-
19. J. Med. Chem. 65, 6499–6512. https://doi.org/10.1021/acs.jmedchem.2c00117
Vangeel, L., Chiu, W., De Jonghe, S., Maes, P., Slechten, B., Raymenants, J., André, E., Leyssen, P., Neyts, J.,
Jochmans, D., 2022. Remdesivir, Molnupiravir and Nirmatrelvir remain active against SARS-CoV-2
Omicron and other variants of concern. Antiviral Res 198, 105252.
https://doi.org/10.1016/j.antiviral.2022.105252
Wang, M., Cao, R., Zhang, L., Yang, X., Liu, J., Xu, M., Shi, Z., Hu, Z., Zhong, W., Xiao, G., 2020. Remdesivir and
chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. Cell
Research 1–3. https://doi.org/10.1038/s41422-020-0282-0
Williamson, B.N., Feldmann, F., Schwarz, B., Meade-White, K., Porter, D.P., Schulz, J., van Doremalen, N.,
Leighton, I., Yinda, C.K., Pérez-Pérez, L., Okumura, A., Lovaglio, J., Hanley, P.W., Saturday, G., Bosio,
C.M., Anzick, S., Barbian, K., Cihlar, T., Martens, C., Scott, D.P., Munster, V.J., de Wit, E., 2020. Clinical
benefit of remdesivir in rhesus macaques infected with SARS-CoV-2. Nature 585, 273–276.
https://doi.org/10.1038/s41586-020-2423-5
Xocova Filed for Conditional Approval in South Korea [WWW Document], n.d. . PHARMA JAPAN. URL
https://pj.jiho.jp/article/248085 (accessed 11.14.23).
Xocova to Be Commercially Available from March 31 [WWW Document], n.d. . PHARMA JAPAN. URL
https://pj.jiho.jp/article/248538 (accessed 10.31.23).

896
897