Adolescent Thalamocortical Inhibition Alters Prefrontal Excitation-Inhibition Balance

David Petersen[1,3], Christoph Kellendonk[1,2,3,#], Sarah Canetta[1,4,#],

[1]Department of Psychiatry, [2]Department of Molecular Pharmacology & Therapeutics, Vagelos

College of Physicians and Surgeons, Columbia University Irving Medical Center, New York, NY

10032, USA

[3]Division of Molecular Therapeutics, [4]Division of Developmental Neuroscience, , New York State

Psychiatric Institute, New York, NY, 10032

[#]Equal contribution

*Correspondence: ck491@cumc.columbia.edu, ses2119@cumc.columbia.edu

Abstract

Adolescent inhibition of thalamo-cortical projections leads to long lasting deficits in prefrontal

cortex function and cognition in the adult mouse. While this suggests a role of thalamic activity

in prefrontal cortex maturation, it is unclear how inhibition of these projections affects prefrontal

network function during adolescence. Here, we used chemogenetic tools to inhibit thalamo-

prefrontal projections in the mouse from postnatal day 20-35 and measured synaptic inputs to

prefrontal pyramidal neurons by layer (either II/III or V/VI) and projection target twenty-four

hours later using slice physiology. We found a decrease in the frequency of excitatory and

inhibitory currents in layer II/III nucleus accumbens projecting and layer V/VI mediodorsal

thalamus projecting neurons. In contrast, layer V/VI nucleus accumbens projecting neurons

showed an increase in the amplitude of excitatory and inhibitory currents. In addition, the

frequency of inhibitory currents in contralateral mPFC projecting neurons was also enhanced.

Notably, despite these complex changes in individual levels of excitation and inhibition, the

overall balance between excitation and inhibition was only changed in cortico-cortical

projections. This finding suggests homeostatic regulation occurs within subcortically- but not

cross-hemispheric intracortical projecting neurons, increased inhibition of intra-prefrontal

connectivity may therefore be particularly important for prefrontal cortex circuit maturation.

Keywords

Adolescence, Excitation-inhibition balance, Thalamocortical projections, Prefrontal cortex

maturation

Introduction

The prefrontal cortex (PFC) supports high level cognitive functioning including working memory and other executive functions (Fuster, 2001; Miller et al., 2018). An impairment in the development or maturation of the prefrontal cortex during adolescence has been postulated to contribute to the cognitive deficits observed in patients with schizophrenia (Feinberg, 1982; Weinberger, 1987; Paus et al., 2008; Gogtay et al., 2011; Rapoport et al., 2012; Sakurai and Gamo, 2019; Chini and Hanganu-Opatz, 2021; Dienel et al., 2022; Smucny et al., 2022). Thalamic inputs regulate sensory cortical maturation during sensitive time windows of development (Wiesel and Hubel, 1963; Nabel and Morishita, 2013; Benoit et al., 2022a). We recently established in the mouse that thalamic input activity during adolescence is necessary for adult prefrontal cortex function and behavior (Benoit et al., 2022b). More specifically, we found that decreasing activity in the mediodorsal thalamus from postnatal day 20-50 using pharmacogenetic tools led to deficits in the performance of two prefrontal dependent cognitive tasks addressing working memory and attentional set shifting. These deficits were measured in the adult mouse and associated with decreased anatomical thalamocortical projections and decreased functional excitatory post-synaptic currents measured in layer 2/3 prefrontal cortical pyramidal cells. Moreover, mice with thalamic inhibition during adolescence showed decreased correlated activity between neurons in the prefrontal cortex while performing an attentional set shifting task and the ability of the prefrontal cortex to decode task outcomes was impaired (Benoit et al., 2022b). Notably, decreasing the activity of thalamocortical projections for a comparable length period in adulthood, did not lead to the same long-lasting effects.

These findings established that thalamo-prefrontal activity is required during adolescence to

sustain normal thalamocortical connectivity, prefrontal network function, and cognitive behavior

in adulthood. However, it was not clear whether these effects on circuit connectivity were

apparent immediately following the adolescent manipulation or require maturation into

adulthood to manifest. Moreover, it was not clear whether the main impact of decreased

adolescent thalamo-prefrontal activity on prefrontal cortical circuitry is to reduce excitatory

thalamic afferents or whether this manipulation leads to more widespread changes in the intrinsic

connectivity of this brain region.

To address these questions, we undertook a detailed study of prefrontal cortical connectivity

using slice electrophysiology twenty-four hours following pharmacogenetic inhibition of

thalamo-prefrontal projections from postnatal day P20-35. We chose this reduced window of

inhibition because: 1) Prior work demonstrated changes in the density of thalamic projections to

the PFC following adolescent thalamocortical inhibition are already present at this time (Benoit

et al., 2022b) and 2) Unpublished work from our laboratory established that inhibiting thalamo-

prefrontal projections in this window is sufficient to induce lasting effects on cognitive behavior

in adulthood. We first recorded excitatory and inhibitory currents from layer II/III neurons as

previously done in the adult (Benoit et al., 2022b), to establish whether the previously observed

deficit arises during adolescence. We then restricted our recordings to specific mPFC projection

neurons identified by their projection target and cortical layer. We found excitatory post synaptic

currents are already reduced in layer II/III cells during adolescence, as in adult mice. When

analyzing specific projections, we found a reduction in frequency of excitatory and inhibitory

currents in layer V/VI cortico-thalamic cells and layer II/III cortico-accumbens projecting cells

while layer V/VI cortico-accumbens projecting neurons displayed increased amplitudes of

excitatory and inhibitory currents. While we observed changes in the absolute frequency of

excitatory or inhibitory inputs in all these subcortically projecting cells following adolescent

thalamic inhibition, notably the overall balance of excitatory and inhibitory currents experienced

by a given cell was not affected, suggesting homeostatic mechanisms may offset decreased

excitatory inputs. In contrast, in cross-hemispheric cortico-cortical projection neurons, we

observed a significant increase in the frequency of inhibitory inputs, and no change in excitatory

inputs, following adolescent thalamic inhibition, with a significant decrease in the overall ratio of

the frequency of excitatory to inhibitory currents. Together, these data show a complex

adaptation of cortical network activity in response to decreased thalamocortical activity during

adolescence and suggest that activity in cells projecting to contralateral PFC may be most

profoundly affected due to a lack of homeostatic adaptation.

Materials and Methods

Animal Husbandry

All procedures were performed in accordance with guidelines approved by the Institutional

Animal Care and Use Committees at Columbia University and the New York State

Psychiatric Institute (protocol NYSPI 1589). Animals were housed under a 12-h light/12-h

dark cycle in a temperature-controlled environment (22 °C, humidity of 30–70%) with food

and water available ad libitum, unless otherwise noted. C57/ BL6 females and males

(Jackson Laboratories, 000664) were used for all experiments. Littermates were randomly

assigned to each experimental group, with random distribution across males and females.

Mice were housed together with dams and littermates. Offspring were weaned at P28, and

group housed with same-sex littermates (no more than five mice per cage). For thalamic

inhibition, mice were given intraperitoneal (i.p.) injections of JHU37160 (J60, a CNO

analog) dissolved in 0.9% sterile saline at 0.1 mg/kg twice per day (Bonaventura et al., 2019).

J60 was used as an alternative to the widely used clozapine-N-oxide (CNO) because J60 is

known to have less off-target effects. New evidence has shown CNO might be reverse

metabolized into clozapine, a common antipsychotic medication with high affinity to

dopamine D2 and serotonin 2A receptors (Manvich et al., 2018). All mice were given J60

regardless of viral vector, except those given saline with hM4D expression. Surgeries were

conducted at P12-13, and all mice were injected from P20 to P35. 24 hours after the final

J60 injection, P35 (+/- 4 days), mice were anesthetized with isoflurane and decapitated for

slice electrophysiology. Throughout data collection and analysis, experimenters were

blinded to the group of the animal. For the dual virus approach, C57/BL6 males and females

were bred, and littermates were used for the experiment. We based sample sizes on previous

experiments, and no statistical methods were used to calculate sample sizes.

Surgical Procedures

For viral injections at P13, mice were anesthetized with isoflurane, and head-fixed in a

stereotactic apparatus (Kopf). Mice were injected bilaterally in the midline thalamus with

AAV5-hSyn-DIO-hM4D-mCherry (Addgene, 44362) or a control virus, AAV5-hSyn-DIO-

mCherry (Addgene, 50459), at a volume of 0.25 µl (0.1µl/min). Mice were also injected

bilaterally in the mPFC with retrograde AAV-hSyn-Cre-WPRE-hGH (Addgene, 105553) at a

volume of 0.2 µl (0.1 µl/min). For the tracer experiments mice were injected with retrograde

AAV-hSyn-EGFP (Addgene, 50465) at a volume of 0.25 µl for the MD, 0.2 µl for the mPFC, and

0.1µl for the NAc. The following were the P13 coordinates: MD thalamus, −0.85 anterior–

posterior (AP), ±0.20 medial–lateral (ML) and −3.25 dorsal–ventral (DV, zero at brain surface);

mPFC, 2.4 AP, ±0.2 ML and −1.9 DV (zero at brain surface); NAc, 1.7 AP, ±1.1 ML and −5.07

DV. All coordinates were zeroed at bregma unless otherwise noted.

Slice Electrophysiology

Whole-cell voltage clamp recordings were performed in layer II/III mPFC pyramidal cells

and layer V/VI mPFC pyramidal cells. Recordings were obtained with a Multiclamp 700B

amplifier (Molecular Devices) and digitized using a Digidata 1440B acquisition system

(Molecular Devices) with Clampex 10 (Molecular Devices) and analyzed with pClamp 10

(Molecular Devices). Following decapitation, 300-µm slices containing mPFC were

incubated in artificial cerebral spinal fluid containing 126mM NaCl, 2.5mM KCl, 2.0mM

$MgCl_2$, 1.25mM NaH_2PO_4, 2.0mM $CaCl_2$, 26.2mM $NaHCO_3$ and 10.0mM d-glucose (pH

7.45, 310 mOsm) bubbled with oxygen at 32 °C for 30min before being returned to room

temperature for at least 1 hour before use. During recording, slices were perfused in room

temperature artificial cerebral spinal fluid at a rate of 5 ml/min. Electrodes were pulled from

1.5-mm borosilicate glass pipettes on a P-97 puller (Sutter Instruments). Electrode

resistance was typically 1.5–3MΩ when filled with internal solution consisting of 130mM

cesium hydroxide monohydrate, 130mM D-gluconic acid, 10mM HEPES, 2.0mM MgCL2,

0.2mM EGTA, 2.5mM Mg-ATP, 0.3mM Na-GTP, and 5mM Lidocaine N-ethyl bromide (pH

7.3, 277mOsm).

mPFC recordings. Animals were sacrificed for recordings at P35 after adolescent

manipulation. mPFC pyramidal cells were visually identified based on their shape and

prominent apical dendrite at ×40 magnification under infrared and diffusion interference

contrast microscopy. Recordings were taken from either layer II/III of the mPFC (PrL) or

layer V/VI. Layer II/III was defined as any cell lying 150μm-350μm from the pial surface.

Layer V/VI was defined as cells within 400μm-600μm from the pial surface. Spontaneous

excitatory post synaptic currents (sEPSCs) were recorded in voltage clamp at a holding

potential of −65mV, and spontaneous inhibitory post-synaptic currents (sIPSCs) were

recorded in voltage clamp at a holding potential of +10mV; 60 s of the current recording for

each condition was analyzed. Recordings were filtered with a 2 kHz eight-pole low-pass

Bessel filter, and sEPSCs and sIPSCs were detected using MiniAnalysis (Synaptosoft).

Frequency, amplitude, and instantaneous frequency were calculated for all traces. Frequency

was calculated by dividing the total number of events by time (60s). Amplitude was the

distance from baseline to the tip of the event. Instantaneous frequency was the inverse of the

interevent-interval converted to a frequency. All event data were averaged by cell.

Statistics

Statistical analysis and graph preparations were done using Prism 9 software (GraphPad

Software), JASP (JASP Team (2023), JASP (Version 0.17.3) or custom scripts in R. Linear

mixed effects models with individual animal used as a random-effects grouping factor were

used to analyze slice physiology. Linear mixed effects models were used over a common

ANOVA or t-test to account for the possible non-independence of measurements from cells

from the same animal. Outliers were removed using the following equation Q1-1.5 x IQR or

Q3 + 1.5 x IQR, this corresponds to +/- 2.698σ from the mean. For the nonspecific L II/III

experiment, 2 outliers were removed from the EPSC hM4D+Saline amplitude group, 1

outlier was removed from the IPSC hM4D frequency group, and 1 outlier was removed from

the IPSC hM4D instantaneous frequency group. For the cortico-accumbens experiment, 2

outliers were removed from the IPSC hM4D group. For the cortico-cortical population

experiment, 2 outliers were removed from the IPSC mCherry frequency group. For the E/I

balance analysis, 2 outliers were removed from the hM4D cortico-cortical frequency group.

Results

Adolescent thalamocortical activity regulates excitatory transmission to

superficial mPFC.

We first performed patch clamp recordings in brain slices from prelimbic layer II/III neurons,

which is the mPFC population of projection neurons that obtain the most prominent input from

the MD (Anastasiades and Carter, 2021). To inhibit thalamic activity during adolescence on

postnatal day 13 (P13), we injected an adeno-associated virus (AAV) carrying a Cre-dependent

version of the inhibitory designer receptor hM4D into the thalamus of C57 mice and a retrograde

virus, rgAAV-CRE, into the mPFC. Retrograde transport will cause Cre recombinase expression

in the MD leading to hM4D expression selectively in medial prefrontal projecting neurons (Fig.

1a). We then chronically inhibited these cortically projecting MD cells with the hM4D ligand

JHU37160 (J60) (0.1 mg/kg) from postnatal day 20 until the day before they were recorded

(P35(-4days)) (Fig. 1b). To control for the possible non-ligand-dependent effects of the hM4D

receptor, we also included a group of mice expressing hM4D that were injected with saline

during the same developmental window. Twenty-four hours after the last J60 injection (at

P35±4days), we recorded layer II/III neurons and analyzed excitatory and inhibitory post-

synaptic currents.

Using linear mixed effects models, we found that inhibition of thalamocortical

projections from P20-35 decreased the frequency [$F(2, 5.93) = 6.854, p = 0.029$] and spatial

distribution (instantaneous frequency) [$F(1,7.02) = 8.129, p = 0.025$] of spontaneous excitatory

currents (sEPSCs), while there was no change in the amplitude [$F(2, 42) = 1.763, p = 0.184$] of

sEPSCs when compared to controls (Fig. 1f-i). Measuring spontaneous inhibitory currents

(sIPSCs) resulted in a trend of diminished frequency [$F(2, 9.78) = 2.906, p = 0.102$] while spatial

distribution (instantaneous frequency) [$F(2, 10.92) = 1.973, p = 0.185$] and amplitude [$F(2, 8.41)$

$= 0.458, p = 0.458$] remained unchanged when compared to controls (Fig. 1f-i). Furthermore,

when comparing the mCherry and Saline groups in a post-hoc contrast, no significant differences

were observed, signaling the lack of non-ligand dependent effects of the hM4D receptor (EPSC

frequency [$z(\infty) = 0.501, p_{bonferroni} = 1.000$]; EPSC median instantaneous frequency [$z(\infty) = 1.097$,

$p_{bonferroni} = 0.818$]; and no overall effect in EPSC amplitude, IPSC frequency, IPSC instantaneous

frequency, or IPSC amplitude. These results indicate that adolescent thalamocortical activity

causes significant alterations to excitatory transmission in the superficial regions of the mPFC,

consistent with what we previously found in adult mice. (Benoit et al., 2022b).

Adolescent thalamocortical activity regulates excitatory and inhibitory

transmission to superficial layer II/III cortico-accumbens projections.

We then determined whether the observed effects of adolescent thalamocortical inhibition

includes the layer II/III cortico-accumbens cell populations. To this end, we injected the rgGFP

virus into the nucleus accumbens (NAc) (Fig. 2a-b) and recorded from the superficial layers

(II/III) of the mPFC. We found that in superficial NAc-projecting neurons, excitatory and

inhibitory transmission were both dampened following adolescent thalamocortical inhibition. For

excitatory currents, the frequency $[F(1, 47) = 9.354, p = 0.004]$ was significantly decreased but

not the spatial distribution (instantaneous frequency) $[F(1, 6.77) = 1.926, p = 0.208]$ or the

amplitude $[F(1, 6.19) = 3.313, p = 0.117]$ (Fig. 2e-f). For inhibitory transmission in superficial

layers, sIPSC frequency was reduced $[F(1, 4.89) = 9.451, p = 0.028]$ but not the spatial

distribution (instantaneous frequency) $[F(1, 7.79) = 0.955, p = 0.358]$ or the amplitude $[F(1, 7.03)$

$= 3.402, p = 0.107]$ (Fig. 2g-h). These data show that excitatory and inhibitory inputs are

decreased onto layer II/III accumbens projecting neurons.

Adolescent thalamocortical activity regulates excitatory and inhibitory

transmission to cortico-thalamic projections.

We then determined the effects of chronic thalamocortical inhibition on a different subcortical

projection, cortical cells from deep layers V/VI projecting to the MD thalamus. A rgGFP virus

was injected into the MD and whole cell patch clamp recordings were taken in thalamic

projecting cortical cells (Fig. 3a-b). Like for non-specific layer II/III and cortico-accumbens

neurons, we observed significant decreases in excitatory synaptic transmission (Fig. 3e-h).

Significant decreases were observed in both the frequency $[F(1, 6.73) = 12.960, p = 0.009]$ and

spatial distribution (instantaneous frequency) $[F(1, 6.10) = 13.045, p = 0.011]$ of sEPSCs but the

amplitude $[F(1, 5.36) = 0.689, p = 0.442]$ of sEPSCs remained unchanged (Fig. 3e-f). In

addition, the frequency $[F(1, 6.39) = 7.829, p = 0.029]$ and spatial distribution (instantaneous

frequency) $[F(1, 5.96) = 6.501, p = 0.044]$ but not the amplitude $[F(1, 4.77) = 0.991, p = 0.367]$

of sIPSCs was reduced after P20-35 thalamic inhibition (Fig. 3g-h). These data show that like for

superficial layer II/III accumbens projections, adolescent thalamo-prefrontal inhibition also

decreases excitatory and inhibitory synaptic inputs to deep layer cortico-thalamic neurons.

Adolescent thalamocortical activity regulates the amplitude of excitatory and inhibitory currents in deep layer (V/VI) accumbens-projecting cortical cells.

Accumbens projecting prefrontal neurons originate from both superficial and deep cortical layers (Anastasiades and Carter, 2021). We therefore also recorded from accumbens-projecting cells in layer V/VI. We found significant increases to excitatory and inhibitory current amplitude, and no changes in frequency or spatial distribution, in deep layer accumbens-projecting cells (Fig 4e-h). When measuring excitatory currents in deep layers, amplitude was significantly increased [$F_{(1, 29)}$ = 8.976, p =0.006] however, frequency [$F_{(1, 5.49)}$ = 7.841x10^{-5}, p =0.993] and spatial distribution (instantaneous frequency) [$F_{(1, 6.24)}$ = 0.196, p =0.673] remained unchanged (Fig 4e-f). Similar to excitatory transmission, for inhibitory currents in deeper layers, amplitude was significantly increased [$F_{(1, 4.25)}$ = 14.809, p =0.013] however, frequency [$F_{(1, 6.82)}$ = 0.417, p =0.539] and spatial distribution (instantaneous frequency) [$F_{(1, 6.53)}$ = 0.910, p =0.374] remained unchanged (Fig. 4 g-h). These data show that unlike layer II/III accumbens projection neurons, excitatory and inhibitory drive onto layer V/VI accumbens neurons is not decreased and not enhanced except at the level of amplitude.

Adolescent thalamocortical activity regulates cross-hemispheric inhibition.

To determine how adolescent thalamocortical inhibition impacts cortico-cortical circuitry, a rGFP virus was injected into the dextral prelimbic mPFC and whole cell patch clamp recordings were taken in cortically-projecting cortical cells in the sinistral hemisphere (Fig. 5a-b). No significant differences were observed in the frequency [$F_{(1, 7.41)}$ = 0.012, p = 0.917], spatial distribution

(instantaneous frequency) [$F(1, 7.30) = 0.399, p = 0.547$] or amplitude [$F(1, 6.37) = 2.865, p =$

0.134] of excitatory events in cortical projecting cortical cells (Fig. 5e-f). Unlike excitatory

transmission, inhibitory transmission significantly increased in the distribution of events

(instantaneous frequency) [$F(1, 5.95) = 8.023, p = 0.030$] and trended in frequency [$F(1, 7.32) =$

$4.185, p = 0.078$] but not amplitude [$F(1, 62) = 0.112, p = 0.912$] after adolescent thalamocortical

inhibition (Fig. 5g-h). These findings were unexpected since callosal projecting prelimbic

neurons are enriched in layer II/III (Anastasiades et al., 2018) and the unspecific analysis of layer

II/III neurons displayed a decrease in the frequency of excitatory currents and no increase in

inhibition.

Excitation-Inhibition balance is only altered in callosally projecting mPFC

neurons.

Excitation and inhibition are well calibrated in prefrontal neurons (Yin et al., 2021) and in

agreement with this we found a strong correlation in the frequency of excitatory and inhibitory

currents for all control neurons across the cortex [$R^2=0.4935$, d.f. =1,121, P<0.0001] (Fig. 6a). In

many cell populations we observed changes in both excitation and inhibition, suggesting

homeostatic mechanisms may compensate for reduced excitatory drive from the thalamus. (Sup.

Fig. 1). We therefore calculated the excitation-inhibition balance in the different analyzed

populations of neurons. No changes in the excitation-inhibition balance were measured in layer

II/III and V/VI cortico-accumbens projecting neurons, and cortico-thalamic neurons (Fig. 6b-e).

In contrast, when analyzing the ratio of excitatory to inhibitory currents in callosally projecting

mPFC neurons, the frequency [$F(1, 11.37) = 5.697, p = 0.035$] and spatial distribution

(instantaneous frequency) [$F(1, 9.34) = 5.744, p = 0.039$] of excitatory to inhibitory currents was

significantly weighted towards inhibition (Fig. 6f).

<u>Discussion</u>

To determine the importance of thalamo-prefrontal activity for prefrontal circuit

maturation we inhibited thalamo-mPFC projections from P20-35 and recorded from prelimbic

neurons twenty-four hours later. We made several observations. First, our findings reveal that

excitatory transmission to layer II/III is already decreased during adolescence. This suggests that

the decrease in excitation measured in adult mice following this manipulation (Benoit et al.,

2022b) does not require full maturation of the circuitry to adulthood but arises early-on during

adolescence. Second, we measured decreased excitatory and inhibitory transmission to layer

II/III nucleus accumbens and layer V/VI thalamic-projecting neurons. These changes in

inhibition suggest that suppressing thalamic activity during adolescence leads to changes in

prefrontal connectivity beyond the loss of excitatory inputs from the thalamus that had

previously been observed at the anatomical level at P35 (Benoit et al., 2022b). Third, we

observed a surprising increase in excitatory and inhibitory transmission to layer V/VI NAc-

projecting neurons but at the level of sEPSC and sIPSC amplitude and not frequency. These

results suggest a post-synaptic adaptation in accumbens-projecting layer V/VI neurons. While

the reason for the specificity of this effect is unclear, cells projecting to the NAc located in deep

layers may have distinct molecular characteristics from those in superficial layers as molecular

characteristics differ between different layers (Bhattacherjee et al., 2019). Finally, we measured

an enhanced inhibition in cortico-cortical connectivity, which provides further evidence for a

complex prefrontal circuit adaptation in response to thalamic inhibition. Intriguingly, while

excitatory and inhibitory drive was altered onto subcortically projection neurons, a change in the

balance of excitation and inhibition was only measured in cross-hemispheric mPFC projections

and not in the three subcortical projections analyzed. As functional stability in neural circuits can

be sustained by homeostatic re-balancing of excitatory and inhibitory inputs, the largest

functional effect of adolescent thalamocortical inhibition on prefrontal circuitry may be a

decrease in the activity of contralaterally-projecting mPFC pyramidal cells. While it is difficult

to predict neuronal activity in vivo from slice physiological measures, enhanced inhibitory inputs

to cross-hemispheric projection neurons may lead to less activity of these neurons during

adolescence, which may have long lasting consequences for mPFC network function in the adult.

Future studies will be able to address whether disinhibition of cross hemispheric projections

during adolescence following thalamocortical inhibition will rescue the long-term negative

effects of the earlier thalamocortical inhibition.

Decreased excitatory transmission in superficial mPFC arises during

adolescence.

Previous work from our group demonstrated that adolescent thalamic inhibition during a

critical period (P20-P50) resulted in long term cognitive and physiological deficits in the

adult mouse, including a loss of excitatory transmission in non-projection-specific layer

II/III pyramidal cells. Here, we observe these deficits are already established during

adolescence, prior to being sustained into adulthood. This dampening of excitatory

transmission onto layer II/III neurons could result from the loss of primary MD innervation

to excitatory pyramidal cells or a loss of local excitation through local feed forward

excitation or both. The seemingly bimodal distribution of the effects in layer II/III neurons

led us to hypothesize that thalamocortical inhibition may be differentially affecting

subpopulations of neurons in layer II/III of the mPFC. To distinguish these effects on

distinct populations of cortical cells, we focused our first analysis on NAc projecting mPFC

neurons.

Adolescent thalamocortical activity regulates excitatory and inhibitory transmission to

superficial nucleus accumbens projecting pyramidal cells.

Excitatory and inhibitory currents were both decreased in NAc-projecting cortical cells

following adolescent thalamic inhibition. These data provide evidence that altering

thalamic input activity during adolescence may have widespread effects on cells projecting

to a variety of regions, including mPFC-NAc projections that are essential for reward driven

motivated behaviors (Kalivas et al., 2005; Mannella et al., 2013; McGlinchey et al., 2016)

Adolescent inhibited mice may therefore not only have deficits in attentional set shifting

and working memory task but also more broadly in reward-driven motivated behaviors.

The decrease in the frequency of inhibitory events could be the consequence of a reduction

in thalamic projections onto interneurons (Yang et al., 2021). For example, the MD directly

innervates layer II/III parvalbumin (PV+) and other interneurons, providing a source of

feedforward inhibition to cortical projection neurons (Delevich et al., 2015; Canetta et al., 2020).

Changes in interneuron-driven inhibition in NAc projecting cells measured in the slice could

reflect either reduced excitatory drive on PV+ cells, or anatomical and functional adaptations

within PV+ interneurons following thalamic inhibition. Future studies directly measuring

excitatory and inhibitory currents in PV+ cells or other cortical interneurons as well as using

optogenetic approaches to probe functional connectivity between those specific interneurons and

NAc-projecting neurons will be able to address this.

Moreover, the decrease in inhibitory transmission within this cell population is distinct from

other populations in the prelimbic cortex, namely layer II/III cortico-cortical cells, which show

increased levels of inhibition following adolescent thalamic inhibition. These findings

underscore the complex effects on excitatory and inhibitory connectivity elicited by adolescent

thalamocortical inhibition in layer II/III pyramidal cells in the prefrontal cortex. To determine

whether adolescent thalamic inhibition also alters excitatory and inhibitory connections in deeper

layers of the mPFC, we next recorded from cortico-thalamic neurons in layer V/VI.

Adolescent thalamic inhibition decreases excitatory and inhibitory drive to prefrontal

projections to the thalamus, further dysregulating the MD-mPFC circuit.

We found that adolescent thalamic inhibition resulted in a decrease in the frequency of

excitatory and inhibitory currents in thalamus projecting cells in layer V/VI, of the mPFC,

similar to the effects seen in NAc-projecting cells in layer II/III. The MD primarily projects

to layer II/III; however, slice physiology has demonstrated monosynaptic connection to the

deeper layers (V/VI) of the cortex, including direct reciprocal connection (Collins et al.,

2018). The origins for the decrease in excitatory and inhibitory drive may therefore be

similar to ones measured in layers II/III.

Deeper (V/VI) accumbens projecting cortical cells show post-synaptic adaptation to

MD inhibition.

Surprisingly, we found that following adolescent thalamic inhibition the amplitude of

excitatory and inhibitory currents increased in layer V/VI accumbens projecting neurons,

while the frequency and spatial distribution remained the same. While the thalamus makes

direct connections onto deep layer pyramidal cells (Collins et al., 2018) it is unknown

whether the MD directly innervates layer V/VI NAc projecting cells. One possibility is that

sEPSC frequency is unaffected in this cell population because it does not receive much

direct thalamic input. Adolescent thalamic inhibition further increased sEPSC amplitude in

these cells. As changes in amplitude are thought to reflect post-synaptic alterations, this

increase in the sEPSC amplitude may be a post-synaptic adaptation though it is currently

unclear what exactly the increased amplitude is compensating for. This increase in sEPSC

amplitude further emphasizes the diverse adaptive and complex changes that thalamocortical

inhibition can produce.

Callosally projecting cells display an increase in inhibitory inputs following adolescent

thalamic inhibition.

The frequency of excitatory inputs to contralaterally projecting mPFC neurons did not

change following adolescent thalamic inhibition. However, inhibitory inputs were increased.

The absence of an effect of adolescent thalamic inhibition on excitatory current frequency in

this population is surprising, given that they receive some of the strongest direct inputs from

the MD (Anastasiades et al., 2018). Why excitatory inputs to this population are spared

remains unknown, but it suggests that excitatory inputs from other brain regions may be

able to offset the reduction in excitatory inputs from the thalamus. This hypothesis could be

directly addressed in future studies using optogenetics to probe the strength of functional

connections made by distinct excitatory connections onto callosally projecting cells. The

increase in inhibition observed in this population is also surprising, given that inhibition was

decreased in NAc projecting cells found in the same cortical layer. Moreover, while changes

in excitation and inhibition driven by adolescent thalamic inhibition largely scaled in all

other cell populations analyzed, cortico-cortical projecting cells showed a significant

reduction in their E/I sPSC ratio.

Overall, the relative stability of the E/I sPSC ratio following adolescent thalamic inhibition

is consistent with the idea of a homeostatic regulation in subcortically projecting neurons

such that the decrease in inhibition may be an adaptive consequence to the decrease in

excitatory inputs they receive from the thalamus. However, in cortically projecting neurons,

thalamic inhibition resulted in an increase in inhibition with no effect on excitation, and a

significant reduction in E/I balance in these cells. One prediction of this finding would be

that cortically projecting neurons might experience the greatest overall change in their

activation profile following adolescent thalamic inhibition, with the net effect of being less

active in vivo. This change may in turn contribute to a lack of cross-hemispheric cortical

communication, and bilateral recruitment of the prefrontal cortex has been hypothesized to

be important for cognitive processes (Hasegawa et al., 1998; Holler-Wallscheid et al., 2017).

This change may also decrease cortical output to other neural circuits, including the nucleus

accumbens and mediodorsal thalamus, thereby having widespread effects on prefrontal

network function.

Alternatively, the increase in contralateral inhibition following adolescent thalamocortical

silencing may have resulted from disinhibition via either the release of inhibition on

somatostatin positive interneurons (SST+) by vasoactive intestinal polypeptide (VIP+)

interneurons or release of inhibition on PV+ interneurons by somatostatin positive

interneurons. VIP+ interneurons receive direct MD input and have strong connections with

SST+ neurons (Ahrlund-Richter et al., 2019; Sun et al., 2019; Canetta et al., 2020). SST+

neurons project onto PV+ cells and the apical dendrites of pyramidal cells. Like PV+ and

VIP+ interneurons, SST+ interneurons are also known to receive direct inputs from the MD

(Ahrlund-Richter et al., 2019; Sun et al., 2019). Therefore, the overall increase in inhibition

could be due to the complex disinhibitory circuit between VIP+, SOM+, PV+ and pyramidal cell

types. It is unclear why this disinhibition should be specific only to cross hemispheric,

intracortical connectivity. While no studies have directly addressed this question, PV

interneurons target callosally projecting pyramidal cells (Cho et al., 2023) while VIP+ and SST+

neurons preferentially target other interneuron populations in the cortex (Pi et al., 2013). Given

the specific increase in the density of inhibitory transmission, this could be the result of

decreased inhibition of PV+ interneurons by SST+ or VIP+ interneurons causing an increase in

the inhibition of callosal projections.

Possible mechanisms of mPFC plasticity induced by adolescent thalamic inhibition.

Recent studies also point to intra-cortical mechanisms in the maturation of cortical circuitry.

Anterior cingulate (ACC) neurons show greater excitatory synaptic inputs during adolescence

than adulthood and DREADD-mediated inhibition of ACC to visual cortex projection neurons

during adolescence has been shown to disrupt the maintenance of local connectivity within the

ACC (Nabel and Morishita, 2013). We also measured an increase in the frequency of excitatory

currents in adolescent mice (5.631+/- 0.466) when compared to adult mice in Benoit et al. 2022

(4.4+-0.4 Hz). By analogy, if inhibition of thalamic inputs decreases activity of mPFC-thalamic

or accumbens projection neurons during adolescence, this may lead to a disruption in the

maintenance of local excitation in the mPFC. A different mechanism has also been described

within the visual cortex. DREADD-mediated inhibition of layer II/III neurons during the critical

period of primary visual cortex development (P24-29) (Wen and Turrigiano, 2021) (Nabel et al.,

2020) led to excitatory synaptic scaling and increased intrinsic excitability suggesting

homeostatic plasticity as a mechanism affecting visual cortex maturation. Future studies will be

able to identify the exact plasticity mechanisms in mPFC circuitry that are induced by thalamic

inhibition during adolescence.

How these findings will scale to the adult mouse remain unknown.

Finally, how do the changes observed in adolescence (P35) relate to physiology and behavior in

the adult mouse? While we have observed similar effects on sEPSC frequency in non-specific

layer II/III cells at both P35 and in adulthood, it will be of interest to see whether the other

changes we have identified also scale into adulthood. If they did, this would be consistent with

the idea that P20-35 is an activity dependent sensitive period in which manipulating the activity

of thalamic inputs to the mPFC leads to changes in the wiring of that circuitry that are then

'hardwired' until adulthood.

Alternatively, changes identified in adolescence may represent a transient change in circuit

dynamics, that then sets in motion subsequent and different changes in circuit dynamics that

result in altered prefrontal network function in adulthood. In this scenario, changes in synaptic

connectivity in adolescent inhibited mice may be different immediately after the inhibition than

later during adulthood, a hypothesis that could be addressed by further recordings in adult

animals.

<u>Conclusions:</u>

In sum, our data demonstrate thalamic inhibition during a sensitive period in adolescence results

in projection specific changes in synaptic connectivity in the mPFC that become evident already

during adolescence. The dysregulation of multiple neural circuits demonstrates the widespread

effects of thalamo-prefrontal inhibition that impact more than just thalamocortical circuitry.

Further, the observed deficits cannot be explained just by a lack of anatomical input. Rather, they

generate widespread adaptative or possibly maladaptive physiological responses within

prefrontal circuitry. Future studies will examine which of these projection specific changes

persist into adulthood, whether new ones emerge and whether they represent appropriate targets

for intervention during adolescence.

Figure 1

Figure 1) **Adolescent thalamocortical activity regulates excitatory transmission to superficial mPFC**. **a**, Adolescent experimental timeline, and schematic. Whole-cell patch clamp recordings were made from pyramidal cells in layer II/III of the mPFC from hM4D and control mice. These pyramidal cells receive excitatory inputs from the thalamus as well as inhibitory inputs from local interneurons; created with https://biorender.com. **b**, Example image illustrating hM4D-mCherry expression in the mediodorsal thalamus (orange outline) in adolescent animals. Histology images were collected from each animal for each cohort, with at least 6 slices taken for each animal. Scale bar represents 300μm. **c**, Example image illustrating the recording location in the prelimbic (PrL) subregion of the medial prefrontal cortex. All recordings for this experiment were conducted in this region approximately 200-350μm from the pial surface (L II/III) denoted by the orange dotted line. Recording pipette outlined in white. Scale bar represents 500μm **d**, Representative traces showing sEPSCs of hM4D animals and mCherry controls. **e**, Representative traces showing sIPSCs of hM4D animals and mCherry controls. **f**, sEPSC frequency (right) and median instantaneous frequency (middle) is significantly reduced following adolescent thalamic inhibition relative to control mice, but sEPSC amplitude (left) is unchanged; mCherry control, $n = 26$ cells and 6 animals; hM4D, $n=14$ cells and 5 animals; hM4D+Saline $n=9$ cells and 2 animals; frequency: mCherry control 4.617(3.050) Hz, hM4D 2.833(3.858) Hz, and hM4D+Saline 5.733(2.700) Hz, contrasts of estimated marginal means revealed a significant difference between hM4D animals and both mCherry, $z = -3.311.129$, d.f. $= \infty$ *$Pbonferroni = 0.003$, and hM4D+Saline, $z = 3.035$, d.f. $= \infty$ *$Pbonferroni = 0.007$ but no significant difference between mCherry and hM4D+Saline, $z = 0.501$, d.f. $= \infty$ *$Pbonferroni = 1.000$; median instantaneous frequency: mCherry control 7.892(5.340) Hz, hM4D 5.126(4.882) Hz, hM4D+Saline 10.005(4.989) Hz; contrasts of estimated marginal means revealed a significant difference between hM4D animals and both mCherry, $z = -3.032$, d.f. $= \infty$ *$Pbonferroni = 0.007$, and hM4D+Saline, $z = 3.348$, d.f. $= \infty$ *$Pbonferroni = 0.002$ but no significant difference between mCherry and hM4D+Saline, $z = 1.097$, d.f. $= \infty$ *$Pbonferroni = 0.818$; amplitude: mCherry control 18.785(5.250) pA, hM4D 20.916(4.882) pA, and hM4D+Saline 28.786(5.782) pA. All reported values are Median(Interquartile Range). **g**, Graph of the cumulative firing distribution for sEPSCs in L 2/3 pyramidal cells in the mPFC, IEI represents the interevent-interval between spontaneous events. **h**, sIPSC frequency (right) may trend downwards but median instantaneous frequency (middle) and amplitude (left) are unchanged following adolescent thalamic inhibition relative to control mice; mCherry control, $n = 26$ cells and 6 animals; hM4D, $n = 14$ cells and 5 animals; frequency: mCherry control 2.500(3.267) Hz, hM4D 1.000(0.550) Hz and hM4D+Saline 2.717(1.833); median instantaneous frequency: mCherry control 3.826(5.039) Hz, hM4D 1.989(1.256) Hz, and hM4D+Saline 4.014(2.366; amplitude: mCherry control 14.264(5.638) pA, hM4D 14.3(6.075) pA, and hM4D+Saline 18.849(6.737). All reported values are Median(Interquartile Range). **i**, Graph of the cumulative firing distribution for sIPSCs in L 2/3 pyramidal cells in the mPFC, IEI represents the interevent-interval between spontaneous events. *P<0.05, #P~0.10.

Figure 2

Figure 2) Adolescent thalamocortical activity regulates excitatory and inhibitory transmission to superficial layer II/III cortico-accumbens projections. a, Adolescent experimental timeline and schematic. Whole-cell patch clamp recordings were made from accumbens projecting pyramidal cells in layer II/III of the mPFC from hM4D and control mice; Created with https://biorender.com. **b**, Example image illustrating rGFP positive, accumbens projecting cells in the prelimbic subregion of the medial prefrontal cortex. Scale bar represents 500μm. **c**, Representative traces showing sEPSCs of hM4D animals and mCherry controls. **d**, Representative traces showing sIPSCs of hM4D animals and mCherry controls**. e**, sEPSC frequency (right) is significantly reduced in accumbens projecting cortical cells following adolescent thalamic inhibition relative to control mice, but sEPSC amplitude (left) and median instantaneous frequency (middle) are unchanged; mCherry control, $n = 21$ cells and 5 animals; hM4D, $n = 28$ cells and 6 animals; frequency: mCherry control 5.367(2.717) Hz, hM4D 2.975(1.871) Hz.; median instantaneous frequency: mCherry control 8.514(4.842) Hz and hM4D 5.579(4.803) Hz; amplitude: mCherry control 21.051(7.992) pA and hM4D 23.648(12.495) pA. All reported values are Median(Interquartile Range). **f**, Graph of the cumulative firing distribution for sEPSCs in L II/III accumbens projecting pyramidal cells in the mPFC, IEI represents the interevent-interval between spontaneous events. **g**, sIPSC frequency (right) is significantly reduced in accumbens projecting cortical cells following adolescent thalamic inhibition relative to control mice, but sIPSC amplitude (left) and median instantaneous frequency (middle) are unchanged; mCherry control, $n = 21$ cells and 5 animals; hM4D, $n = 28$ cells and 6 animals; frequency: mCherry control 3.350(2.350) Hz, hM4D 1.417(1.762) Hz; median instantaneous frequency: mCherry control 5.195(4.817) Hz and hM4D 2.776(2.753) Hz; amplitude: mCherry control 14.618(3.768) pA and hM4D 17.597(7.919) pA. All reported values are Median(Interquartile Range). **h**, Graph of the cumulative firing distribution for sIPSCs in L II/III accumbens projecting pyramidal cells in the mPFC. *P<0.05.

Figure 3

Figure 3) Adolescent thalamocortical activity regulates excitatory and inhibitory transmission to cortico-thalamic projections. a, Adolescent experimental timeline, and schematic. Whole-cell patch clamp recordings were made from cortical projecting pyramidal cells in layer V/VI of the mPFC from hM4D and control mice. These pyramidal cells receive excitatory inputs from the thalamus and superficial layers of the cortex as well as inhibitory inputs from local interneurons; created with https://biorender.com. **b**, Example image illustrating rGFP positive cells in the prelimbic subregion of the medial prefrontal cortex. All recordings for this experiment were conduced in this region approximately 500-600μm from the pial surface (L V/VI). Scale bar represents 300μm **c**, Representative traces showing sEPSCs of hM4D animals and mCherry controls. **d**, Representative traces showing sIPSCs of hM4D animals and mCherry controls. **e**, sEPSC frequency (right) and median instantaneous frequency (middle) is significantly reduced in thalamic projecting cortical cells following adolescent thalamic inhibition relative to control mice, but sEPSC amplitude (left) is unchanged; mCherry control, n = 19 cells and 4 animals; hM4D, n = 26 cells and 5 animals; frequency: mCherry control 6.717(4.392) Hz and hM4D 2.833(1.408) Hz; median instantaneous frequency: mCherry control 11.173(7.187) Hz and hM4D 4.204(1.997) Hz; amplitude: mCherry control 21.764(9.897) pA and hM4D 24.043(13.508) pA. All reported values are Median(Interquartile Range). **f**, Graph of the cumulative firing distribution for sEPSCs in L V/VI thalamic projecting pyramidal cells in the mPFC, IEI represents the interevent-interval between spontaneous events. **g**, sIPSC frequency (right) and median instantaneous frequency (middle) is significantly reduced in thalamic projecting cortical cells following adolescent thalamic inhibition relative to control mice, but sIPSC amplitude (left) is unchanged; mCherry control, n = 19 cells and 4 animals; hM4D, n = 26 cells and 5 animals; frequency: mCherry control 4.917(1.825) Hz and hM4D 2.100(1.350) Hz; median instantaneous frequency: mCherry control 7.896(4.100) Hz and hM4D 2.755(2.081) Hz; amplitude: mCherry control 19.267(7.264) pA and hM4D 15.720(7.127) pA. All reported values are Median(Interquartile Range). **h**, Graph of the cumulative firing distribution for sIPSCs in L V/VI thalamic projecting pyramidal cells in the mPFC. *P<0.05.

Figure 4

Figure 4) Adolescent thalamocortical activity results in enhanced amplitude in deeper (V/VI) accumbens projecting cortical cells. a, Adolescent experimental timeline and schematic. Whole-cell patch clamp recordings were made from accumbens projecting pyramidal cells in layer V/VI of the mPFC from hM4D and control mice; Created with https://biorender.com. **b**, Example image illustrating rGFP positive, accumbens projecting cells in the prelimbic subregion of the medial prefrontal cortex. Scale bar represents 500µm. **c**, Representative traces showing sEPSCs of hM4D animals and mCherry controls. **d**, Representative traces showing sIPSCs of hM4D animals and mCherry controls**. e**, sEPSC amplitude (left) is significantly increased in accumbens projecting cortical cells following adolescent thalamic inhibition relative to control mice, but sEPSC frequency (right) and median instantaneous frequency (middle) are unchanged; mCherry control, $n = 22$ cells and 5 animals; hM4D, $n = 9$ cells and 4 animals; frequency: mCherry control 4.450(2.012) Hz, hM4D 3.167(4.217) Hz.; median instantaneous frequency: mCherry control 7.371(4.043) Hz and hM4D 5.290(12.190) Hz; amplitude: mCherry control 20.460(11.638) pA and hM4D 29.858(5.869). All reported values are Median(Interquartile Range). **f**, Graph of the cumulative firing distribution for sEPSCs in L II/III accumbens projecting pyramidal cells in the mPFC, IEI represents the interevent-interval between spontaneous events. **g**, sIPSC frequency (right) is significantly reduced in accumbens projecting cortical cells following adolescent thalamic inhibition relative to control mice, but sIPSC amplitude (left) and median instantaneous frequency (middle) are unchanged; mCherry control, $n = 21$ cells and 5 animals; hM4D, $n = 28$ cells and 6 animals; frequency: mCherry control 1.883(1.817) Hz, hM4D 2.50(2.350) Hz; median instantaneous frequency: mCherry control 2.997(2.405) Hz and hM4D 4.005(4.349) Hz; amplitude: mCherry control 13.362(2.306) pA and hM4D 17.495(3.331). All reported values are Median(Interquartile Range). **h**, Graph of the cumulative firing distribution for sIPSCs in L V/VI accumbens projecting pyramidal cells in the mPFC. *P<0.05.

Figure 5

Figure 5) Adolescent thalamocortical activity regulates cross-hemispheric inhibition. a, Adolescent experimental timeline and schematic. Whole-cell patch clamp recordings were made from cortical projecting pyramidal cells in layer II/III of the mPFC from hM4D and control mice; Created with https://biorender.com. **b**, Example image illustrating rGFP positive, cortical projecting cells in the prelimbic subregion of the medial prefrontal cortex. Virus was injected into the dextral (right) hemisphere and cells were recorded only in the sinistral (left) hemisphere. Scale bar represents 500µm. **c**, Representative traces showing sEPSCs of hM4D animals and mCherry controls. **d**, Representative traces showing sIPSCs of hM4D animals and mCherry controls. **e**, sEPSC frequency (right), sEPSC amplitude (left) and sEPSC median instantaneous frequency are unchanged in cortical projecting cortical cells following adolescent thalamic inhibition relative to control mice; mCherry control, $n = 38$ cells and 5 animals; hM4D, $n = 26$ cells and 5 animals; frequency: mCherry control 3.042(1.713) Hz and hM4D 2.95(2.933) Hz; median instantaneous frequency: mCherry control 4.807(3.944) Hz and hM4D 5.042(5.890) Hz; amplitude: mCherry control 17.876(7.549) pA and hM4D 23.951(8.397) pA. All reported values are Median(Interquartile Range). **f**, Graph of the cumulative firing distribution for sEPSCs in cortical projecting pyramidal cells in the mPFC, IEI represents the interevent-interval between spontaneous events. **g**, sIPSC median instantaneous frequency (middle) is significantly increased and frequency (right) is trending towards an increase in cortical projecting cortical cells following adolescent thalamic inhibition relative to control mice, while amplitude (left) is unchanged; mCherry control, $n = 38$ cells and 5 animals; hM4D, $n = 26$ cells and 5 animals; frequency: mCherry control 1.233(0.954) Hz and hM4D 2.208(1.887) Hz; median instantaneous frequency: mCherry control 1.937(1.676) Hz and hM4D 5.551(2.498); amplitude: mCherry control 15.735(5.268) pA and hM4D 15.099(5.667) pA. All reported values are Median(Interquartile Range). **h**, Graph of the cumulative firing distribution for sIPSCs in cortical projecting pyramidal cells in the mPFC. *P<0.05, #P<0.10.

Figure 6

a.

E/I Ratio in Control Cortical Cells

b.

| Amplitude | Med. I. Frequency | Frequency |

Layer 2/3

c.

| Amplitude | Med. I. Frequency | Frequency |

2/3 Cortico-acc.

d.

Cortico-thalamic

e.

5/6 Cortico-acc.

f.

Cortico-cortical

Figure 6) Adolescent thalamocortical inhibition disrupts E/I Ratio in cortical projecting cortical cells. a, Excitatory to inhibitory regression in control cortical cells. There was a significant correlation between EPSCs and IPSCs [R^2=0.4935, d.f. =1,121, P<0.0001]. **b,** Excitatory/Inhibitory ratio in nonspecific layer II/III cells for amplitude (left), median instantaneous frequency (middle), and frequency (right). There was no significant difference in E/I ratio between mCherry 1.188(0.400) pA/pA, hM4D 1.432(0.704) pA/pA or hM4D+Saline 2.005(1.247) pA/pA for amplitude [f = 2.567, d.f. = 2, 11.07 P = 0.121]. There was no significant difference in E/I ratio between mCherry 2.131(2.020) Hz/Hz, hM4D 1.736(2.409) Hz/Hz or hM4D+Saline 2.183(1.306) Hz/Hz for median instantaneous frequency [f = 0.713, d.f. = 2, 11.65 P = 0.511]. There was no significant difference in E/I ratio between mCherry 2.265(4.677) Hz/Hz, hM4D 2.186(3.134) Hz/Hz or hM4D+Saline 2.490(1.325) Hz/Hz for median instantaneous frequency [f = 1.209, d.f. = 2, 43 P = 0.308]. **c,** Excitatory/Inhibitory ratio in layer II/III accumbens projecting cortical cells for amplitude (left), median instantaneous frequency (middle), and frequency (right). There was no significant difference in E/I ratio between mCherry 1.316(0.587) pA/pA or hM4D 1.524(0.505) pA/pA for amplitude [f = 0.543, d.f. = 1, 6.59 P = 0.487]. There was no significant difference in E/I ratio between mCherry 1.686(1.368) Hz/Hz and hM4D 1.999(1.715) Hz/Hz for median instantaneous frequency [f = 0.836, d.f. = 1, 7.91 P = 0.388]. There was no significant difference in E/I ratio between mCherry 1.663(0.963) Hz/Hz and hM4D 1.865(1.406) Hz/Hz for frequency [f = 0.750, d.f. = 1, 8.10 P = 0.411]. **d,** Excitatory/Inhibitory ratio in layer V/VI thalamic projecting cortical cells for amplitude (left), median instantaneous frequency (middle), and frequency (right). There was no significant difference in E/I ratio between mCherry 1.195(0.500) pA/pA or hM4D 1.376(0.858) pA/pA for amplitude [f = 3.625, d.f. = 1, 42 P = 0.064]. There was no significant difference in E/I ratio between mCherry 1.414(0.485) Hz/Hz and hM4D 1.660(1.143) Hz/Hz for median instantaneous frequency [f = 0.975, d.f. = 1, 5.21 P = 0.367]. There was no significant difference in E/I ratio between mCherry 1.414(0.485) Hz/Hz and hM4D 1.503(1.136) Hz/Hz for frequency [f = 1.115, d.f. = 1, 5.31 P = 0.337]. **e,** Excitatory/Inhibitory ratio in layer V/VI accumbens projecting cortical cells for amplitude (left), median instantaneous frequency (middle), and frequency (right). There was no significant difference in E/I ratio between mCherry 1.467(0.624) pA/pA or hM4D 1.730(0.618) pA/pA for amplitude [f = 2.220, d.f. = 1, 28 P = 0.147]. There was no significant difference in E/I ratio between mCherry 2.060(1.840) Hz/Hz and hM4D 2.432(1.435) Hz/Hz for median instantaneous frequency [f = 0.412, d.f. = 1, 12.33 P = 0.533]. There was no significant difference in E/I ratio between mCherry 1.829(1.623) Hz/Hz and hM4D 2.658(1.765) Hz/Hz for frequency [f = 0.014, d.f. = 1, 12.10 P = 0.908]. **f,** Excitatory/Inhibitory ratio in cortical projecting cortical cells for amplitude (left), median instantaneous frequency (middle), and frequency (right). There was no significant difference in E/I ratio between mCherry 1.209(0.359) pA/pA or hM4D 1.473(0.403) pA/pA for amplitude [f = 1.099, d.f. = 1, 59 P = 0.299]. There was a significant decrease in E/I ratio between mCherry 2.513(1.915) Hz/Hz and hM4D 1.517(1.670) Hz/Hz for median instantaneous frequency. There was

also a significant decrease in E/I ratio between mCherry 2.594(1.607) Hz/Hz and hM4D 1.564(1.544) Hz/Hz for frequency. All reported values are Median(Interquartile Range). *P<0.05.

between mCherry 2.594(1.607) Hz/Hz and hM4D 1.564(1.544) Hz/Hz for frequency. All reported values are Median(Interquartile Range). *P<0.05.

Sup. Figure 1.

Amplitude   **Instantaneous Frequency**   **Frequency**

Supplementary Figure 1) Regression analyses for E/I Ratio across all experiments.

Sup. Figure 2.

Supplementary Figure 2) Total viral spread of hM4D-mCherry Receptor in Red. B is

distance from bregma.

References

Ahrlund-Richter S, Xuan Y, van Lunteren JA, Kim H, Ortiz C, Pollak Dorocic I, Meletis K,
Carlen M (2019) A whole-brain atlas of monosynaptic input targeting four different cell
types in the medial prefrontal cortex of the mouse. Nature neuroscience 22:657-668.
Anastasiades PG, Carter AG (2021) Circuit organization of the rodent medial prefrontal cortex.
Trends in neurosciences 44:550-563.
Anastasiades PG, Marlin JJ, Carter AG (2018) Cell-Type Specificity of Callosally Evoked
Excitation and Feedforward Inhibition in the Prefrontal Cortex. Cell reports 22:679-692.
Benoit LJ, Canetta S, Kellendonk C (2022a) Thalamocortical Development: A
Neurodevelopmental Framework for Schizophrenia. Biological psychiatry.
Benoit LJ, Holt ES, Posani L, Fusi S, Harris AZ, Canetta S, Kellendonk C (2022b) Adolescent
thalamic inhibition leads to long-lasting impairments in prefrontal cortex function. Nature
neuroscience 25:714-725.
Bhattacherjee A, Djekidel MN, Chen R, Chen W, Tuesta LM, Zhang Y (2019) Cell type-specific
transcriptional programs in mouse prefrontal cortex during adolescence and addiction.
Nature communications 10:4169.
Bonaventura J et al. (2019) High-potency ligands for DREADD imaging and activation in
rodents and monkeys. Nature communications 10:4627.
Canetta S, Teboul E, Holt E, Bolkan SS, Padilla-Coreano N, Gordon JA, Harrison NL,
Kellendonk C (2020) Differential Synaptic Dynamics and Circuit Connectivity of
Hippocampal and Thalamic Inputs to the Prefrontal Cortex. Cereb Cortex Commun
1:tgaa084.
Chini M, Hanganu-Opatz IL (2021) Prefrontal Cortex Development in Health and Disease:
Lessons from Rodents and Humans. Trends in neurosciences 44:227-240.
Cho KKA, Shi J, Phensy AJ, Turner ML, Sohal VS (2023) Long-range inhibition synchronizes
and updates prefrontal task activity. Nature 617:548-554.
Collins DP, Anastasiades PG, Marlin JJ, Carter AG (2018) Reciprocal Circuits Linking the
Prefrontal Cortex with Dorsal and Ventral Thalamic Nuclei. Neuron 98:366-379 e364.
Delevich K, Tucciarone J, Huang ZJ, Li B (2015) The Mediodorsal Thalamus Drives
Feedforward Inhibition in the Anterior Cingulate Cortex via Parvalbumin Interneurons.
The Journal of neuroscience : the official journal of the Society for Neuroscience
35:5743-5753.
Dienel SJ, Schoonover KE, Lewis DA (2022) Cognitive Dysfunction and Prefrontal Cortical
Circuit Alterations in Schizophrenia: Developmental Trajectories. Biological psychiatry
92:450-459.
Feinberg I (1982) Schizophrenia: caused by a fault in programmed synaptic elimination during
adolescence? Journal of psychiatric research 17:319-334.
Fuster JM (2001) The prefrontal cortex--an update: time is of the essence. Neuron 30:319-333.

Gogtay N, Vyas NS, Testa R, Wood SJ, Pantelis C (2011) Age of onset of schizophrenia: perspectives from structural neuroimaging studies. Schizophrenia bulletin 37:504-513.

Hasegawa I, Fukushima T, Ihara T, Miyashita Y (1998) Callosal window between prefrontal cortices: cognitive interaction to retrieve long-term memory. Science 281:814-818.

Holler-Wallscheid MS, Thier P, Pomper JK, Lindner A (2017) Bilateral recruitment of prefrontal cortex in working memory is associated with task demand but not with age. Proceedings of the National Academy of Sciences of the United States of America 114:E830-E839.

Kalivas PW, Volkow N, Seamans J (2005) Unmanageable motivation in addiction: a pathology in prefrontal-accumbens glutamate transmission. Neuron 45:647-650.

Mannella F, Gurney K, Baldassarre G (2013) The nucleus accumbens as a nexus between values and goals in goal-directed behavior: a review and a new hypothesis. Frontiers in behavioral neuroscience 7:135.

Manvich DF, Webster KA, Foster SL, Farrell MS, Ritchie JC, Porter JH, Weinshenker D (2018) The DREADD agonist clozapine N-oxide (CNO) is reverse-metabolized to clozapine and produces clozapine-like interoceptive stimulus effects in rats and mice. Scientific reports 8:3840.

McGlinchey EM, James MH, Mahler SV, Pantazis C, Aston-Jones G (2016) Prelimbic to Accumbens Core Pathway Is Recruited in a Dopamine-Dependent Manner to Drive Cued Reinstatement of Cocaine Seeking. The Journal of neuroscience : the official journal of the Society for Neuroscience 36:8700-8711.

Miller EK, Lundqvist M, Bastos AM (2018) Working Memory 2.0. Neuron 100:463-475.

Nabel EM, Morishita H (2013) Regulating critical period plasticity: insight from the visual system to fear circuitry for therapeutic interventions. Front Psychiatry 4:146.

Nabel EM, Garkun Y, Koike H, Sadahiro M, Liang A, Norman KJ, Taccheri G, Demars MP, Im S, Caro K, Lopez S, Bateh J, Hof PR, Clem RL, Morishita H (2020) Adolescent frontal top-down neurons receive heightened local drive to establish adult attentional behavior in mice. Nature communications 11:3983.

Paus T, Keshavan M, Giedd JN (2008) Why do many psychiatric disorders emerge during adolescence? Nature reviews Neuroscience 9:947-957.

Rapoport JL, Giedd JN, Gogtay N (2012) Neurodevelopmental model of schizophrenia: update 2012. Molecular psychiatry 17:1228-1238.

Sakurai T, Gamo NJ (2019) Cognitive functions associated with developing prefrontal cortex during adolescence and developmental neuropsychiatric disorders. Neurobiology of Disease 131:104322.

Smucny J, Dienel SJ, Lewis DA, Carter CS (2022) Mechanisms underlying dorsolateral prefrontal cortex contributions to cognitive dysfunction in schizophrenia. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 47:292-308.

Sun Q, Li X, Ren M, Zhao M, Zhong Q, Ren Y, Luo P, Ni H, Zhang X, Zhang C, Yuan J, Li A, Luo M, Gong H, Luo Q (2019) A whole-brain map of long-range inputs to GABAergic interneurons in the mouse medial prefrontal cortex. Nature neuroscience 22:1357-1370.

Weinberger DR (1987) Implications of normal brain development for the pathogenesis of schizophrenia. Archives of general psychiatry 44:660-669.

Wen W, Turrigiano GG (2021) Developmental Regulation of Homeostatic Plasticity in Mouse Primary Visual Cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience 41:9891-9905.

578   Wiesel TN, Hubel DH (1963) Single-Cell Responses in Striate Cortex of Kittens Deprived of
579   Vision in One Eye. Journal of neurophysiology 26:1003-1017.
580   Yang SS, Mack NR, Shu Y, Gao WJ (2021) Prefrontal GABAergic Interneurons Gate Long-
581   Range Afferents to Regulate Prefrontal Cortex-Associated Complex Behaviors. Frontiers
582   in neural circuits 15:716408.
583