Adherent reformed islets: a long-term primary cell-based platform for exploring mouse and human islet biology.

Haq N.[1*], K.W. Toczyska[1*], Wilson M.E.[1], Jacobs M.[1], Min Zhao[1], Lei Y.[1], Pearson J.[2], S.J. Persaud[1], Pullen T.J.[1], Bewick G.A.[1]

Affiliations

1. Department of Diabetes, School of Cardiovascular and Metabolic Medicine & Sciences, King's College London, Guy's Campus, London SE1 1UL, UK.
2. Diabetes Research Group, Division of Infection and Immunity, School of Medicine, Cardiff University, Cardiff, UK.

Abstract

Pancreatic islets are 3-dimensional micro-organs that maintain β-cell functionality via cell-cell and cell-matrix communication. Isolated primary islets are the gold standard for in vitro models. However, native islets present experimental challenges for long-term mechanistic studies owing to their short culture life (approximately 1 week). We developed a novel long-term protocol to study the function of primary islets. The protocol employed reformed islets following dispersion and a fine-tuned culture environment. Reformed islets are highly similar to their primary counterparts across various physiological characteristics. Long-term culture of reformed islets enables high-resolution imaging, repeated functional assessment, and the study of cell-cell communication. Unlike other platforms such as stem cell-derived organoids, reformed islets retain their resident immune populations, making them ideal for studying both resident and infiltrating immune cells and their interactions with hormone-producing islet cells.

Qualitative and quantitative analyses revealed that the composition and cytoarchitecture of the reformed islets mimicked those found in primary islets, including the presence of macrophages and $CD4^+$ and $CD8^+$ T cells, which are the key resident immune cell types. Reformed islets secrete insulin and are glucose-responsive, and their β-cells can be stimulated to proliferate using GLP-1 receptor agonism. Furthermore, a comparison of the transcriptomic landscape of isolated human islets and reformed islets generated from the same donor demonstrated a high degree of similarity.

Our reformed islets provide an ideal platform to study diabetes pathology. We recapitulated both the T1DM and T2DM disease milieu and validated our model for studying islet immune trafficking and invasion using activated macrophages and T cells.

Our data illustrates that reformed islets are an anatomical and functional alternative to native human and mouse islets. Moreover, reformed islets have an advantage over mouse and human β-cell lines, including MIN6 and EndoC-βH1cells, that lack the signalling input of non-β-endocrine cells and immune cell crosstalk. In this study, we showed that reformed islets are a durable paradigm (cell-based model) for islet-based exploration and a means of target discovery/validation for diabetes research.

Introduction

The prevalence of both type 1 and type 2 diabetes mellitus (T1DM, T2DM), which are among the leading causes of morbidity and mortality, is rapidly increasing, making it one of the largest global health concerns, affecting 537 million adults worldwide [1-3]. Although effective treatments exist for T2DM, intensive insulin therapy and islet transplantation for a limited number of people are the only approved option for T1DM despite improvements in care. For both types, treatment aims to manage symptoms and reduce complications; however, there is no cure.

Diabetes pathogenesis involves a complex interplay of multiple mechanisms, including genetics, islet/β-cell dysfunction, and immune dysregulation. As such, efforts to understand disease pathology have focused on comprehensively exploring and understanding all aspects of islet biology. Islets are micro-organs that regulate glucose homeostasis through the secretion of the key hormones insulin, glucagon, and somatostatin. They comprise interspersed β, α, δ, ε, and PP cells intermingled with resident immune cells, stellate cells, smooth muscles, and fibroblasts [4]. β-cell functionality is maintained via cell-cell and cell-matrix communication. Therefore, understanding islet physiology and pathological dysfunction requires exquisite knowledge of how all islet cells communicate to maintain normal function, and how this is dysregulated to drive disease progression.

Primary islets, often isolated from rodents, are the gold standard for diabetes research. They are readily available and cost-effective, and their function is comparable to those observed *in vivo* [5]. However, isolating pancreatic islets requires significant time and effort, and limited experiments can be performed because of their short culture life, usually no more than a week before significant reductions in functionality are evident [6]. This is equally true for primary human islets which are available for research purposes from islet transplantation programmes. These limitations have restricted investigations, such as high-resolution live imaging of islets over prolonged periods or repeated functional assessments, owing to the short culture life and insufficient techniques for culturing adherent primary islets. Other alternatives, such as β-cell lines, including MIN6 cells (mouse), EndoC-βH1 (human) or EndoC-βH5 (human), can be cultured on glass, but they are transformed β-cell lines and therefore lack the signalling input of non-β cells, including other hormone-producing cells and immune cells [7]. Although two-dimensional culture of dispersed islet cells is possible on plastic surfaces, it requires a coating with cell-derived extracellular matrices. This includes extracellular matrix secreted from human carcinoma cells or bovine corneal epithelial cell matrix, which have high batch-to-batch variability and are linked to β-cell dedifferentiation ([8-10].

In recent years, 3D culture of organ-like structures has paved the way for understanding the
pathology of various diseases, aided drug discovery, enabled the identification of novel
therapeutic targets, and enhanced personalised medicine. Therefore, developing novel 3D
models of pancreatic islets is of high importance to aid the study of the pathophysiology of
diabetes and to better understand the response to various compounds and therapeutics.
Dispersion of whole islets and their subsequent reaggregation into spheroids offers one such
method with the advantage of adherent long-term culture under suitable conditions and the
potential for more efficient transgene delivery. A previous study attempted to use
reaggregation and the subsequent formation and maintenance of islet organoids by using
novel expansion and proliferation culture media. These islet clusters harboured key islet
hormone-producing cell types and showed a higher degree of proliferation than native islets.
However, this paradigm has limitations such as: (i) islet-derived cell clusters could only be
generated from pregnant mice or wild-type rats, (ii) cluster numbers were small and
inadequate for transcriptomic studies, (iii) the islet clusters were not functionally evaluated,
and (iv) the presence of a resident immune population was not determined [11]. Stem cell-
derived pancreatic organoids are an alternative. These cells are generated by directed
differentiation of pluripotent stem cells and can be cultured over the long term. However,
they also suffer from limitations, including not exhibiting all hormone-producing cell types,
non-physiological insulin secretion, and a lack of a resident immune population.

Given the need for alternative methods for the long-term culture of primary islets, we aimed
to refine reaggregation techniques to generate a superior adherent 3D islet culture platform
using native murine or human islets as the source. These reaggregated islets are referred to
as reformed islets. They can be maintained in culture for at least 4-6 weeks, making them
ideal for long-term studies, including high-resolution imaging, suitable for repeated functional
assessment, and studying cell-cell communication. We found the reformed islets to be
architecturally, transcriptionally, and functionally highly similar to the native islets they are
generated from.

Methods

2.1 Materials

Hanks' Balanced Salt Solution and Hepes buffer were obtained from Gibco, B27 (Thermo
Fisher Scientific). Fluoxetine hydrochloride, collagenase type XI, histopaque-1077, culture
media, and supplements were purchased from Sigma-Aldrich. The primary antibodies used
in this study are summarised in Supplementary Tables 1 and 2. Recombinant murine tumour
necrosis factor α (TNF-α), interferon γ (IFNγ), and interleukin-1β (IL-1β) were purchased
from PeproTech EC, and the Caspase-Glo 3/7 Assay Kit was purchased from Promega.

CD8[+] cells were isolated from the spleens of four 8-week-old diabetic female NY8.3 NOD

mice using a MojoSort Mouse CD8 T-cell isolation kit (Cat# 480043, BioLegend, San Diego,

CA, USA) and MojoSort buffer (Cat# 480017, BioLegend). Cell Biolabs' CytoSelect™ cell

migration assay from Cell Biolabs and EZCellTM cell migration/chemotaxis assay kits were

purchased from BioVision.

2.2 Animal Husbandry

CD-1 male mice were purchased from Charles River and were housed under temperature-

(22±2ºC) and light-controlled conditions (12 h light:12 h dark cycle) with ad libitum access to

drinking water and a standard rodent chow diet. Spleens for CD8+ cell isolation were

obtained from four 8-week-old female diabetic NY8.3 non-obese diabetic (NOD) mice.

Animal experiments were performed in accordance with the British Home Office Animal

Scientific Procedures Act (1986).

2.3 Human Donors

Human islets were retrieved from non-diabetic, heart-beating, and brain-dead donors and

isolated by cold collagenase digestion ([12, 13] with appropriate ethical approval from the

King's College London Human Islet Research Tissue Bank (KCL HI-RTB; 20/SW/0074).

Islets from two donors were used in this study. Donor 1 was 41yr old female with a BMI of 24

and donor 2 was 36yr old male with a BMI of 23.

2.4 Islet Isolation

Mouse islets were isolated from male CD-1 mice (≥ 20 g) aged 8-12 weeks and separated

from exocrine tissue by digestion with collagenase. Isolated mouse islets were cultured

overnight (37°C, 95%air/5%CO2) in RPMI supplemented with 10% (v/v) foetal bovine serum

(FBS), 100U/ml penicillin and 100µg/ml streptomycin, and 2mM L-glutamine.

2.5 Preparation of matrix-coated coverslips

Round borosilicate glass coverslips (12 mm diameter and 0.17 mm thickness) were

transferred to 24-well plates and coated with either laminin from Engelbreth-Holm-Swarm

murine sarcoma basement membrane at 50µg/ml or Collagen IV stock solution 1 mg/ml in

Ca^{2+}/Mg^{2+}supplemented with Hanks Balanced Salt Solution (HBSS) overnight at 37°C. The

coverslips were washed in HBSS and left to dry for 10 min before seeding.

2.6 Preparation of reformed islets

2000-2500 mouse or human islets were hand-picked from suspension cultures 24h after

isolation and washed with PBS. Islets were then dissociated into a suspension of single cells

through trypsin digestion, which was stopped by adding neurobasal media consisting of

MEM supplemented with 5% (v/v) FBS, 1x B-27, 1%(v/v) penicillin and streptomycin, HEPES

10 mM, Glutamax 1x, Na-pyruvate 1mM, and 11 mM glucose. Islet cells were seeded at a

density of 35,000 cells/cm^2 on laminin-coated or collagenase-coated glass coverslips. They

required 3-4 days of culture to adhere and spread on the glass surfaces, and 10-14 days to

form reformed islets. These reformed islets were used for all the experiments described

here. The reformed islets required a change in the media every 2 days.

2.7 Immunofluorescence and confocal microscopy

Immunofluorescence was used to study the cytoarchitecture of the reformed islets, as well

as proliferation and cytokine-induced damage. Coverslips with 6-10 reformed islets were

fixed with 4% PFA and incubated in blocking buffer (1%BSA + 10%donkey serum in 0.1%

PBST) for 1h at room temperature to eliminate non-specific antibody binding. Thereafter, the

reformed islets were incubated with primary antibodies against insulin, glucagon,

somatostatin, Ki67, CD80, IBA1, CD8, CD3e, and CD4 overnight at 4°C and washed in PBS

(Supplementary Table 1). The slides were subsequently exposed to appropriate secondary

antibodies for 1h at room temperature and washed with PBS (Supplementary Table 2). DAPI

Fluoromount-G was used for the mounting. Images of the reformed islets were taken with an

Eclipse Ti-E Inverted A1 confocal microscope or Zeiss LSM700 Confocal Microscope and

analysed using CellProfiler and ImageJ software.

2.8 Insulin secretion from mouse and human reformed islets

Insulin secretion from reformed mouse and human islets was measured by static incubation

experiments. Reformed islets were pre-incubated for 1h in physiological salt solution (Gey &

Gey, 1936) supplemented with 2mM glucose. Coverslips with 6-10 reformed islets were

counted under a Nikon TMS phase-contrast microscope and incubated with appropriate

volumes (100µl of buffer per reformed islet) of Gey Gey buffer supplemented with either

2mM or 20mM glucose. The supernatants were collected and stored at -20°C prior to insulin

radioimmunoassay.

2.9 Apoptosis: Caspase-Glo 3/7 assay

Caspase-Glo 3/7 assay was used to detect apoptosis by measuring caspase 3/7 activity.

Coverslips having 4-5 reformed islets and primary islets were exposed to a mix of pro-

inflammatory cytokine cocktail (CK) (0.05U/µl IL-1β, 1U/µl TNF-α and 1U/µl IFN-γ) diluted in

RPMI supplemented with 2% FBS and 1% Pen/strep for 24h. Caspase 3/7 activity was
quantified by bioluminescence according to the manufacturer's protocol.

2.10 Islet destruction: Migration assay

The Transwell cell migration assay measures the chemotactic capability of cells toward a
chemo-attractant. First, we isolated islets from CD1 mice, BALB/c mice, and human
cadavers. BALB/c mice were used, as these mouse strains express the MHC class H2Kd
haplotype and are therefore matched to antigen-specific T cells.

The reformed islets were incubated in RPMI supplemented with 2% FBS and 1% Pen/strep
for 24h. Subsequent studies using the selective serotonin reuptake inhibitor fluoxetine were
chosen for migration and invasion studies, as there is evidence to support the anti-
inflammatory and β-cell-protective properties of this drug [11, 14].

Simultaneously, RAW 264.7 cells were activated with LPS (200 ng/ml) and IFN-_ (2.5 ng/ml)
for 24 h [15]. After washing with PBS to remove LPS and IFN- _, the RAW cells were
trypsinised, pelleted, seeded ($1X10^6$ cells) into inserts (upper chamber), and placed over
coverslips with at least 3-4 reformed islets for an additional 24h.

CD8+ cells were isolated from the spleens of four 8-week-old diabetic female NY8.3 NOD
mice. Briefly, a single-cell suspension was prepared with MojoSort buffer at 1×10^8 cells/ml.
10 µl of biotinylated antibody containing antibodies against CD4, CD11b, CD11c, CD19,
CD24, CD45R/B220, CD49b, CD105, I-A/I-E, TER-119/Erythroid, and TCRγδ were added.
The mixture was then incubated for 15 min on ice, and 10 µL streptavidin nanobeads were
added. The final volume of 120 µL of the cell and bead mixture was incubated on ice for an
additional 15 min. Finally, 120 µL was diluted in in 1x MojoSort buffer (2.5 mL) and placed in
a MojoSort Magnet (Cat# 480019/480020, BioLegend) for 5 min for magnetic separation.
The free cells in the solution were collected as enriched CD8$^+$ T cells.

The migration assay was measured using an 8µm pore transwell plate (BioVision, Inc., USA)
according to the manufacturer's protocol. Briefly, BALB/c reformed islets were incubated for
48h in 2% FBS RPMI complete medium in the lower chamber of a Transwell culture plate.
Thereafter, $2X10^5$ CD8$^+$ T cells were seeded into the upper chamber and placed over
coverslips with reformed islets of both groups: untreated reformed islets and cytokine
cocktail-exposed reformed islets. After 18 h of incubation, the cells in the lower chamber
were quantified by measuring the O.D values at Ex/Em = 540/590 nm using a PHERAstar
Microplate Reader.

To investigate reformed islets as a platform to perform migration studies, the effects of the drug fluoxetine on macrophage migration and reformed-islet destruction in the presence of pro-inflammatory cytokines were studied using established migration assay kits. Briefly, mouse or human reformed islets were incubated for 48h in 2% FBS RPMI complete media with or without a drug of interest for the first 24h. After the initial 24h, activated monocyte/macrophage-like RAW 264.7, were seeded into the upper chamber ($2X10^5$ cells) and placed over coverslips with reformed islets. Pro-inflammatory cytokines were added for the final 24h. After 24 h of incubation, the cells in the lower chamber were quantified by measuring the O.D values at Ex/Em = 540/590 nm using a PHERAstar Microplate Reader.

2.11 Islet destruction: Invasion assay

A Transwell cell invasion assay was developed to measure both cell chemotaxis and invasion of islet cells through the extracellular matrix. This method is often used for studying cancer metastasis and embryonic development [16]. To this end, we studied the effects of cytokine-induced destruction of mouse and human reformed islets in the presence of $CD8^+$ cells and activated monocyte/macrophage-like cells RAW 264.7 and assessed by confocal imaging [17].

For this study, we used coverslips that underwent migration assays (described above). The coverslips were fixed and immunolabelled with insulin, CD8, or CD80 antibodies and counterstained with DAPI for the identification of nuclei. Qualitative analyses using light and confocal microscopy were used to assess the appearance of the reformed islets.

2.12 Bulk RNA-sequencing

Total RNA was extracted from three replicates of native human islets and reformed islets from one donor, using the TRIzol method. Samples were uniquely barcoded and pooled into one sample. After sequencing, all the samples were distinguished based on their unique barcodes. The next step included In Vitro Transcription, is a linear amplification step which results in amplified RNA (αRNA). This α-RNA was fragmented and run on an Agilent TapeStation to assess RNA integrity.

Reverse transcription, in addition to a PCR reaction, amplified the α RNA, and appropriate sequencing adapters were added. During the PCR reaction, the material was again amplified, and appropriate adapters for sequencing were added to the sequences. This resulted in a DNA library (cDNA) that could be used for sequencing. The cleaned cDNA library was run on an Agilent TapeStation instrument.

Library preparation and RNA sequencing were performed by Single-Cell Discoveries using
the QuantSeq 3 kit (Lexogen) and sequenced on an Illumina NextSeq 500 with a
sequencing depth of 10 million reads per sample. Raw sequencing data were aligned
against the human reference genome using STARsolo (2.7.10a). Read quality checks were
carried out using FastQC, after which BBDuk from the BBMap (version 38.87) software was
used to trim Illumina adapters and polyA tails. Reads were mapped to the hg38 reference
human genome using Star Aligner.

2.13 Data processing

2.13.1 Bulk RNA-Seq analysis
290

The R package DESeq2 v1.36.0 [18, 19] was used to determine differential gene
expression, and DEGs with padj <0.1 were selected. Variance-stabilising transformations of
raw count data were used to visualise the sample clustering. All data were subjected to
global-scale normalisation and log-transformation. Linear dimensionality reduction was
performed using principal component analysis on the top 500 genes from the variable-
stabilising transformation dataset using the DESeq2 package. The data was also subject to
shrinkage estimation of the $\log_2$ fold changes using apeglm.

2.13.2 Bioinformatic pathway analysis
300

Gene Set Enrichment Analysis (GSEA) using the Hallmark, Gene Ontology biological
processes, and KEGG databases was conducted with the fgsea package v1.22.0 [19].
Heatmaps were generated using normalised counts and Z-scores scaled using the Complex
Heatmaps package v2.12.1 [20]. Volcano plots were generated using SRPLOT software,
using a $\log_2$ fold-change of specific genes in the dataset. A gene was considered
significantly differentially expressed if it demonstrated an adjusted p-value < 0.05 and $\log_2$
FC> 1.5. Sequence data are available from the GEO database under accession number
GSE248341.

2.14 Statistical analysis

Statistical analysis and graphical representation of the data were performed using GraphPad
Prism software (version 9.5.1; GraphPad Software, La Jolla, CA, USA). Statistical
comparisons were performed using one-way ANOVA, followed by Šídák's or Dunnett's
multiple comparison tests. All results are expressed as the mean ± SEM. Statistical
significance was defined as a p-value < 0.05 and is denoted as stars in the graphs (* p <
0.05; ** p< 0.01; *** p< 0.001; **** p< 0.0001) in all figures, unless otherwise stated.
316

Results

3.1 Reformed islets present organotypic cytoarchitecture akin to native primary islets

We optimised the reaggregation of islet cells by integrating a previously published protocol
[10] with several enhancements and a more extensive validation process. The critical
modification involved extending the maturation time after cell disaggregation and seeding
from 4 days to 14 (Fig. 1). This extended period enabled the cells to re-aggregate and
develop into 3D spherical structures, whereas they remained as a monolayer using the
previous protocol. Furthermore, once the reformed islets reached maturity, they were
successfully maintained on the platform for 42-days. This extended culture time allowed us
to observe the long-term viability and functionality of the reformed islets.

After seeding, single islet cells adhered to the coated coverslip, and within several days,
these cells formed a nonconfluent but connected network. After seven days of culture, focal
points within the network began to form 3-dimensional spheroid structures (Fig. S1A) that
harboured multi-hormonal cells producing insulin, glucagon, and somatostatin (Fig. S1B). At
14-16 days, these spheroids were fully formed and exhibited a native islet cytoarchitecture
with distinct hormone-producing populations. The size of the reformed mouse islets was
comparable to that of the native mouse islets, whereas human reformed islets tended to be
slightly smaller than their native counterparts (181.1±2.07 vs 126.3±18.71) (Table 1) (Fig.
2A, B).

We evaluated the cellular composition of mouse reformed islets, with β cells comprising
72.64%, α cells constituting 4.86%, and δ cells comprising 3.62% of the total cell population
(Fig. 2C). Notably, the distribution of glucagon-positive cells in the reformed islets was
similar to that observed in the native islets and was localised predominantly in the periphery.
Similarly, in human reformed islets, the cellular composition (β-61.70%, α-28.06%, δ-9.23%)
closely resembled that of native human islets which contain 50-60% β cells, 30-50% α cells
and 9-10% δ cells (Fig. 2D).

Native mouse islets consist of approximately 65–80% insulin-producing β-cells which are
primarily located in the inner core and, 10–20% of the cells are glucagon-producing α cells
found in the outer region of the islet. Next, we compared the cellular composition of the
major cell types between native and reformed islets in both mouse and humans. In mouse,
the percentages of β (*p-val*=0.48) and α cell types (*p-val*=0.41) were similar to those of
reformed islets. The percentage of δ cell type was slightly lower (p =0.1) (Fig. 2E, F and G).
In human native and reformed islets, the percentages of β-cells (p =0.98) and α-cells (p
=0.99) were similar. However, there was a lower percentage of δ cells in the reformed islets
(p =0.59) (Fig. 2H, I and J).

Similar to mouse reformed islets, the distribution of human glucagon-positive and
somatostatin-positive cells mirrored that of native human islets, where they were
interspersed with β-cells (Fig. 3 A, B). The hormonal cell composition and positional pattern
in both mouse and human reformed islets support the notion that our modified platform is an
excellent mimic of primary native islets. In addition to hormone-producing cell types, a
resident immune population is present in both murine and human islets [32]. This immune
population is key to normal function and accounts for approximately 2-3% of all islet cells.
Macrophages (MΦ) are the predominant fraction in both mouse and human islets, but other
immune cell types, such as CD4 and CD8 T cells are also present (Fig 3A, S2 C-E).
Increasing evidence suggests that these immune cells play crucial roles in maintaining islet
homeostasis, fine tuning islet development, and modulating insulin secretion [20, 33].
Moreover, the resident immune population has been shown to contribute to the initiation and
progression of diabetes. For instance, the resident macrophage population is believed to be
one of the key initiators of islet T cell infiltration, a hallmark of T1DM. The immune population
in the reformed islets was verified by quantitative PCR. The transcriptional level of CD45
(Fig. S2A), and macrophages (Fig. S2B) were comparable to those of the native islets.
Additionally, immunolabelling of the reformed islets revealed the presence of both resident
macrophages (identified by Iba1, ionised calcium-binding adaptor molecule 1, and positivity)
and resident T-cells (identified by CD8 and CD4 immunoreactivity) (Fig. 2C, and 3A-lower
panel). The presence of a resident immune population is an important advantage of our
platform because all other long-term culture paradigms lack these cell types.

3.2 Reformed islets are able to proliferate when exposed to a GLP-1 receptor agonist

Mature β-cells within primary islets typically remain in a quiescent state. However, one
potential approach to increase functional β-cell mass to treat insulin-dependent diabetes is
to induce proliferation of the residual β-cell population. We investigated whether reformed
islets retained the ability to proliferate when treated with the GLP-1 receptor agonist exendin-

4, which is known to enhance β-cell proliferation [21]. Exposure to exendin-4 significantly

increased islet Ki67 immunolabelling (p <0.05), which is a marker of cell proliferation (Fig 4A,

B). This indicates that reformed islets exhibit a capacity for proliferation, highlighting their

utility in elucidating ways to promote β-cell mass expansion.

3.3 Functional insulin secretion machinery in reformed islets

The dissociation of whole islets into single cells leads to the loss of connection which can

cause dysregulated insulin secretion [22]. Insulin secretion in response to glucose in the

islets of Langerhans is a tightly regulated process that involves multiple cell types and

signalling pathways. In response to glucose, mature primary islets display a biphasic insulin

secretory response [23].

Insulin secretion was quantified in static incubation experiments using primary and reformed

islets. Acute exposure (1h) to supramaximal glucose levels (20 mM) resulted in significantly

increased levels of secreted insulin compared to 2 mM glucose (Fig 4 C, D) in both native

and reformed mouse islets. Similarly, the concentrations of secreted insulin were

comparable between the native and reformed human islets (Fig 4 E, F). This suggests that

the reformed islets are glucose sensing and capable of nutrient-driven insulin secretion.

3.4 Transcriptional Analysis of reformed islets

Human pancreatic islets were obtained from a 36-year-old male cadaver with a BMI of 23 (>

85% purity and >60% viability). Native islets were dissociated and seeded onto collagen-

coated coverslips. After 15 days of culture, native islets and reformed islet samples were

processed for paired bulk RNA-Seq analysis (Single-cell Discoveries Ltd.). More than 70% of

raw reads were mapped to the reference genome. Globally, 1800 genes were differentially

expressed with an adjusted p value < 0.05 when a cut-off value of > ±2 was applied to the

log_2 fold change (FC) (Fig.5A). Hierarchical clustering of the top 30 genes demonstrated

significant differences between native and reformed islets (Fig. S3A). To further analyse

these data, Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis (Fig.5

B, C) was performed. Overexpression RNA-Seq analysis using significant log_2FC discards a

large proportion of the dataset that may be of biological relevance. Therefore, using the

Molecular Signature Database collection of hallmark genes, we performed Gene Set

Enrichment Analysis (GSEA)[24, 25] on our dataset as a complementary method. The

hallmark gene dataset is a compilation of various published well-defined biological

processes or states that display coherent expression. GSEA uses only p-values, thereby

reducing the variation and redundancy of gene expression across various pathways.

The enriched upregulated pathways included epithelial-mesenchymal transition (EMT) (*VACN, TGFB1, THBS2, LOXL2, ACTA2, ADAM12, SGCG, POSTN, COL1A1, CDH2* etc.) (Fig. S3B), focal adhesion (*COL5A, PDGFRA, THBS2, COL1A2, FLNC, COL5A1, ITGA11, COL1A1, COMP*, etc.) (Fig. S3 C), and skeletal system development (*PDGRFA, HAND2, HOXA1, HOXA3, HOXA4, HOXA5, SULF1, HOXC6, HOXC5, COL1A1*, etc.) (Fig. S3 D). As expected, there were overlaps between genes with enriched pathways. The upregulation of transcripts encoding EMT genes, focal adhesions, and skeletal muscle development suggests that the EMT process is ongoing in cultured reformed islets. EMT has been found to be present in human epithelial cell cultures [13], and a recent study established that EMT occurs when human islets are allowed to expand in culture. These mesenchymal cells are highly proliferative and maintain the ability to differentiate into hormone-producing cells [26].

The downregulated differentially expressed genes (DEGs) were enriched in KEGG pathways related to ribosome biogenesis (*RPL11, RPL14, RPS6, RPL12, RPL21, RPL13, RPS15, RPS9, RPL3, RPS4X* etc.) (Fig. S3E), and Maturity Onset of Diabetes in the Young (MODY) (*NR5A, NKX6.1, GCK, INS, IAPP, ONECUT1, FOXAA3, FOXA2, HNF4A*, etc.) (Fig. S3F). Downregulation of ribosomal RNA (rRNA) transcription often occurs in response to intra- and extracellular stimuli in order to maintain homeostasis. Downregulation of rRNA transcription has been associated with cellular processes such as cell cycle arrest and reduced cell growth, which may positively regulate cell differentiation [27]. Genes within the MODY pathway have been shown to be positively correlated with insulin secretion, and a decrease in gene expression has been associated with β-cells being unable to maintain their mature phenotype [28, 29]. The expression of islet hormones, such as INS, GCG, SST, PYY, GHRL, and NPY, was generally mildly downregulated in reformed islets compared to their native source (Fig. S3G) despite the proportions of cell numbers being similar when assessed by immunostaining. Additionally, transcription factors related to the development of endocrine cells, such as RFX3, NKX6.1, FOXA2, and GLIS3, were downregulated (Fig.S3H). Collectively, the transcriptional landscape of the reformed islets suggests that they have a high degree of similarity to the primary islets they are derived from but exhibit a partially dedifferentiated and relatively immature state. These features may aid long-term culture stability.

3.5 Reformed islets response to cytokine-induced damage mimics that of primary islets

Inflammatory responses and the presence of cytokines secreted by immune and non-immune cell types are central to both types of diabetes [30, 31]. While T2DM is associated with insulin resistance and β-cell dysfunction, recent studies in mouse models have linked

an increase in circulating inflammatory factors, such as chemokines and cytokines, to islet
inflammation and insulitis [32]. This is mainly due to the increased number of intra-islet
macrophages that secrete cytokines such as IL-1β. Three cytokines have been implicated in
this process: IFN-γ, produced by CD4+ and CD8+ T cells, and TNF-α and IL-1β, produced
by islet-infiltrating dendritic cells and macrophages [33]. The presence of this 'cytokine
storm' orchestrates inflammation of islets with subsequent β-cell death in T1DM, and
enhancement of insulin resistance with impairment of glucose homeostasis in T2DM.

We modelled this 'cytokine storm' by addition of a cocktail of pro-apoptotic cytokines (IL-1β,
TNF-α and IFN-γ), and quantified caspase 3/7 activation in our reformed islets. Our results
illustrated that cytokine-induced damage in mouse reformed islets (Fig. 6A) closely
resembled that observed in primary mouse islets (Fig. 6B). We further qualitatively evaluated
this cytotoxic damage by immunolabelling untreated and cytokine-treated reformed islets
stained with antibodies directed against insulin and cleaved caspase-3, a surrogate marker
of apoptosis. We observed that the untreated mouse reformed islets mostly maintained a
round shape with clearly defined margins, in contrast to the cytokine-treated reformed islets
that exhibited a loss of architecture with increased granularity, indicating islet destruction
(Fig. 6C).

3.6 Reformed islets are a qualitative and quantitative platform to study immune
invasion and migration

The development of insulitis is a common feature in both T1DM and T2DM. Although their
aetiologies differ, the invasion and accumulation of macrophages and T cells in the islets
display a pro-inflammatory phenotype [43, 44] that is common to both inflammatory
processes.

The progression of T1DM involves immune cell infiltration of islets, specifically CD8+ and
CD4+ T cells. The exact mechanism remains unclear; however, resident antigen-presenting
cells (APCs), such as dendritic cells and macrophages, detect β-cell proteins as 'non-self'
(MHC class I molecules on β-cells), causing the processing of β-cell-specific antigens and
their presentation to CD4+ T lymphocytes via the MHC class II complex [34, 35]. CD4+ T
helper cells become activated, allowing them to further activate B lymphocytes and trigger
autoantibody production [36]. Additionally, CD8+ cytotoxic T cells are activated through β cell
antigen presentation on MHC class I, triggering the release of cytokines and resulting in β
cell death [37].

The disease progression of T2DM is associated with the development of insulin resistance
and β-cell dysfunction. Interestingly, recent studies in mouse models have found that the

number of intra-islet macrophages increases significantly with an increase in circulating
inflammatory factors, such as chemokines and cytokines, leading to islet inflammation and
insulitis [32].

Using our reformed islets, we modelled immune migration and invasion in both types of
diabetes by co-culturing them with either T cells or macrophages in the presence or absence
of a cytokine cocktail to drive a pro-inflammatory environment (Fig. 7A).

3.6.1 Migration assay
487

In native islets with insulitis, $CD8^+$ cells are the most abundant invading cell type and are
reactive against β cell antigen-presenting cells in T1D. To mimic *in vivo* infiltration by
autoreactive CD8 cells in islets, a transwell migration assay was performed to detect the
migratory ability of $CD8^+$ cells. Reformed islets were derived from BALB/c mice, and CD8
cells (haplotype matched) were harvested from the spleens of non-obese diabetic (NOD)
mice, seeded onto the membrane of the transwell insert, and placed over coverslips with
adherent reformed islets. As expected, haplotyped matched T-cells migrated to reformed
islets, and this process was significantly increased during inflammation after 18h (Fig. 7B).

An increase in the number of islet-resident macrophages is prevalent in both types of
diabetes [38, 39]. To model this, we co-cultured reformed islets with RAW264.7 cells a
murine macrophage-like cell line. Under inflammatory conditions, RAW cell migration was
significantly increased (Fig. 7C).

3.6.2 Invasion assay
501

Next, we quantified cell migration using an invasion assay. Reformed islets were co-cultured
with T cells or activated RAW cells at baseline and during the inflammatory stimulus.
Invasion was imaged by immunostaining of immune cells (CD8 or CD80 antibodies) and β-
cells (insulin antibody) in combination with confocal microscopy. Reformed islets derived
from BALB/c mice were co-cultured with $CD8^+$ T cells and qualitatively assessed at 2 and 18
h. After 2 h, we observed that reformed islets exposed to cytokines showed destruction of
their architecture and infiltration of $CD8^+$ cells compared with reformed islets not exposed to
$CD8^+$ cells (Fig. 7E). After 18 h, both cytokine-treated and untreated reformed islets were
completely invaded by CD8 + cells, with complete destruction of the islets (Fig. 7F.

To study macrophage invasion, reformed islets were exposed to an inflammatory stimulus
and the invasive capability of LPS and IFN-γ activated RAW 264.7 cells was assessed. The

number of infiltrating RAW 264.7 cells were qualitatively higher in the presence of cytokines
than in untreated controls (Fig.7G).

3.6.3 Islet protective agent can cloak islets from immune cell infiltration
516

Finally, as a proof of concept, we sought to validate the use of our platform to identify
compounds that could prevent islet invasion and protect islets from damage. To this end, we
assessed the islet-protective effects of fluoxetine. Fluoxetine is a commercially available
anti-depressant drug that has been shown to significantly increase β-cell proliferation and
protect islet cells from cytokine-induced apoptosis [14].To replicate the protective effect of
fluoxetine, reformed islets were exposed to a cytokine cocktail in the presence or absence of
a therapeutic concentration of fluoxetine (1 µM) for 24h while activated macrophages (RAW
264.7) were added to the inserts of the Transwell plate and cultured for a further 24h.
Quantitative analysis indicated a significant reduction in the number of migrating activated
RAW cells when reformed islets were cultured under inflammatory conditions and treated
with fluoxetine (Fig 8A). Additional qualitative analysis by confocal microscopy illustrated the
reduced migration and islet protection (Fig 8 B, C).

Collectively, we show that reformed islets can be modelled to replicate the *in vivo*
pathogenesis common to both T1DM and T2DM. Moreover, interrogation of reformed islets
in a T2DM setting in the presence of an islet protective compound, fluoxetine, shows that
this platform is ideal for exploring the function of drugs designed to protect islets from
damage.

Discussion

This study reports the development of a long-term biomimetic method to model native
mouse and human islets *ex vivo*, while maintaining long-term islet functionality. We
performed transcriptome profiling to evaluate how closely our model recapitulates native
islets and glucose-stimulated insulin secretion, to assess functionality and evaluate islet
morphology in terms of islet cell-type distribution and size. We further investigated the
suitability of this platform for studying beta cell proliferation and immune cell-islet cell
interactions by performing invasion and migration assays.

One of the major results of this study highlighted that reformed islets mimic primary islets
and can survive for long periods in culture. In our protocol, disassociated primary islet cells
are reaggregated and, over time, form 3D structures which mimic their primary counterparts.

This is in contrast to previously reported protocols, in which reaggregated islet cells remain
in a monolayer [9]. Both mouse and human reformed islets harbour the major islet hormone
cell types arranged architecturally, and in proportions similar to primary islets [40]. The
functionality of the reformed islets was assessed using insulin secretion, beta-cell
proliferation, and cytokine-induced apoptosis assays. Our results indicate that glucose-
stimulated insulin secretion is comparable to that observed in primary mouse and human
islets [41]. In addition, we observed that reformed-islet cells could be stimulated to proliferate
in the presence of exendin-4 and that cytokine-induced damage could easily be measured
using a caspase 3/7 assay [42-44]. Our findings illustrate that reformed islets are an
anatomical and functional alternative to native human and mouse islets, but which can be
cultured for longer periods (4-6 weeks).

To understand how closely our reformed islets modelled native islets, we performed bulk
RNA sequencing and transcriptomic analysis of freshly isolated native human islets and the
reformed islets on day 16. Global expression profiling revealed a number of differentially
expressed genes between native and reformed islets. However, functional clustering and
pathway analyses of differentially expressed genes indicated an association with Epithelial-
mesenchymal Transition (EMT) hallmark genes in the reformed islets. These EMT-enriched
genes have stem cell-like properties and may generate endocrine islets via committed
precursor cells [45, 46]. We postulate that dissociation of native islets results in

dedifferentiation and EMT activation of cells, which then redifferentiation over-time following

aggregation giving rise to all endocrine cell types. It is plausible that dedifferentiation and

reaggregation enables islets to withstand long-term culture conditions.

One of the key advantages of our model is the presence of a resident immune population

similar to that of native islets. Various studies have highlighted the role of resident immune

cells, particularly macrophages, in islet development, beta cell homeostasis, and

regeneration, as well as in the pathology of diabetes [47, 48]. Indeed, the onset of diabetes,

specifically T1D, results from macrophage activation which drives the recruitment of CD8+

and CD4+ T-cells to the islet, ultimately resulting in the destruction of beta-cells [49]. The

lack of *in vitro* models where resident immune cells can be studied has hampered our

understanding of this disease process. To our knowledge, this model is the first in which

most immune cell types can be found.

Using reformed islets, we were able to recapitulate features of the immune infiltration

observed in T1D mechanisms by coculturing reformed islets generated from BALB/c mice

and splenic CD8$^+$ T cells from haplotype-matched NOD mice. We speculate that this

platform can be extended to human reformed islets by co-culturing haplotype-matched

reformed human islets with human T-cell clones. Furthermore, we could also model T2D.

T2D patients have an elevated number of macrophages infiltrating their islets [50, 51]. We

modelled this phenotype by coculturing mouse reformed islets with macrophage-like RAW

264.7 and observed complete infiltration and destruction of islets in the presence of

cytokines.

Fluoxetine, an approved antidepressant drug has recently been shown to increase insulin

secretion in response to glucose and to increase beta cell mass [14, 52, 53]. As a proof of

concept, we chose this drug to validate our human reformed islet model and qualitatively and

quantitatively assess Fluoxetine's ability to protect islets against cytotoxic insults. As

expected, the migration and invasion assays showed that fluoxetine reduced macrophage

migration and apoptosis. Thus, our study demonstrated that reformed islets can be used to

identify new targets or could be employed as a screening platform for drug repurposing or

repositioning once optimised in a multi-well format [54].

Long-term imaging of pancreatic islets is challenging, and the gold standard method for

measuring β-cell mass is to evaluate the pathological staining of fixed pancreatic tissue

sections. However, this does not encompass the dynamic changes that the beta-cell mass

undergoes during the different stages of disease progression. Reformed islets allow a non-

invasive, high-resolution system to quantitatively image at the single-islet level which can be

useful for the evaluation of islet viability, proliferation, and the effects of novel anti-diabetic

compounds. For example, reformed islets can be used to report macrophage infiltration by
coculturing with transgenic-labelled fluorescent macrophages.

In summary, we have refined a method for generating a static reformed islet platform which
can be applied to both mouse and human cells and cultured over the long-term. Reformed
islets are genotypically, phenotypically, and functionally like the primary islets they are
derived from and importantly harbour a resident immune population, which is missing in
other platforms. These properties make reformed islets highly amenable to exploring islet
cell-cell communication for example using multiple imaging modalities and repeated
functional assessment. The platform could easily be optimised for use in a multi-well format
for high content screening programmes and will likely prove to be a relatively easy platform
to genetically manipulate given the dispersion step in the methodology. Dispersed islets cells
are far easier to transfect whether using lipid-based transfection reagents, viral vectors, or
electroporation [55, 56]. We hope the platform will be a useful tool to further our
understanding of diabetes pathology.

Supplemental Table 1: Primary Antibodies

Target	Company	Product #	Concentration
Insulin	Abcam	ab7842	1:200
Glucagon	Merck Life Science UK Limited	G2654	1:1000
Somatostatin	Abcam	ab30788-125ul	1:50
Ki67	Abcam	ab15580	1:200
CD80	Bio-Techne	AF740-SP	1:100
IBA1	Alpha Laboratories Limited	019-19741	1:200
CD45	Abcam	ab40763	1:100
CD8α	Cell Signaling	98941S	1:100
CD4	BD Pharmingen™	557667	1:50

Supplemental Table 2: Secondary Antibodies

Target	Company	Product #	Concentration
Alexa Fluor® 488-AffiniPure Donkey Anti-Guinea Pig IgG	Stratech	706-545-148-JIR-0.5mg	1:200
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 568	Invitrogen	A10042	1:1000
Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 647	Invitrogen	A32787	1:1000
Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 647	Invitrogen	A48272	1:1000
Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 594	Invitrogen	A-11058	1:1000

Figure legends

653 **Figure 1. Overview of reformed protocol**. Native islets isolated from mouse or human
654 donors were allowed to recover overnight. On day 0 they were trypsinised and seeded at a
655 density of 35,000 cells/cm^2 on laminin-coated or collagenase-coated glass coverslips. They
656 required 3-4 days of culture to adhere and spread on the glass surfaces, and 10-14 days to
657 form reformed islets. Experiments were performed thereafter.

Figure 2. Comparative analyses of size and cell composition of native and reformed islets. The native and reformed islets were immunostained for insulin, glucagon somatostatin and IBA analysed using the ImageJ software. Sizes of native and reformed mouse (A) and human (B) islets were compared and presented in Table 1. Percentages of different cell types in mature mouse (C) and human (D) reformed islets were calculated and compared to native islets (mouse:E-G, human: H-J). Numerical data are presented as the mean ± SEM, n=4-13. Ns represents p>0.05, unpaired Student t-test or One-way ANOVA, Dunnett's multiple comparisons test.

Figure 3. Immunofluorescence detection of islet hormones in mature mouse (A) and human (B) reformed islets formed from single islet cells 14 days after seeding. The reformed islets were immunoprobed with antibodies against insulin (INS; green), glucagon (GCG; purple), and somatostatin (SST; red). DAPI nuclear staining is shown in blue. Scale bar is 20µm. Percentage calculation of cell types in mature mouse (C) and human (D) reformed islets was performed using the ImageJ software. Numerical data are presented as the mean ± SEM, n=6-13.

Figure 4. Effects of exendin-4 on proliferation of mouse reformed islets. (A)The reformed islets were exposed to exendin-4 for 30h and immunoprobed with antibodies against insulin (INS; green) and Ki67 (red). DAPI is shown in blue in merged images. Scale bar is 20µm. Percentage calculation of Ki67^{+ve} cells within the reformed islets was performed using the ImageJ software (B). Data are expressed as mean ± SEM, n=5 observations from two separate experiments. *p<0.05 relative to the control samples, One-way ANOVA, Dunnett's multiple comparisons test. Insulin secretory response of primary islets (mouse – C, human – E) and reformed islets (mouse – D, human –F) to supramaximal glucose concentration (20mM). In static incubation experiments, islets or reformed islets were pre-incubated for 2h in Gey and Gey buffer supplemented with 2mM glucose and then incubated for 1h in the presence of either 2mM or 20mM glucose. All data are shown as the mean ± SEM, n=4-8 observations. *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001 relative to the control samples at 2mM glucose, unpaired Student t-test.

Figure 5. Transcriptomic profiling of reformed human islets. (A) Volcano plot of all genes significantly upregulated or downregulated in reformed islets (log2-fold-change >2, benjamini-hochberg corrected p-value threshold = 0.1). (B) Gene set enrichment analysis (GSEA) of RNA-seq data performed on reformed islets illustrates enriched pathways. (F) KEGG pathway analysis was performed for functional enrichment of genes in reformed islets.

Figure 6. The responses of mice and human reformed islets to pro-inflammatory
cytokines. Reformed islets were incubated with a mix of pro-inflammatory cytokines (IL-1β,
TNF-α, IFN-γ) for 24h. Apoptosis of mouse (A) and primary islets (B) was measured by
quantifying caspase 3/7 activities using the Caspase-Glo 3/7 assay following a 24h exposure
to cytokines. In parallel, reformed islets were immunoprobed with antibodies against insulin
(INS; green) and cleaved caspase 3 (CAS-3; red) (A,). DAPI is shown in blue, and the scale
bar is 20µm. Data are shown as mean ± SEM; n=8 observations representative of three
experiments. ****p<0.0001 versus control; One-way ANOVA, Dunnett's multiple
comparisons test.

Figure 7. Illustration of migration and invasion assays. Schematic diagram of *ex vivo*
migration and invasion assays of CD8+ T cell and activated macrophages (RAW 264.7 cells)
(A). Bar graph shows mean total percentage of CD8+ T cells migrating when cocultured with
haplotype matched BALB/c reformed islets (B). Bar graph shows mean total percentage
activated RAW 264.7 macrophage-like cells co-cultured with reformed CD1 mouse islets and
number of migrating cells□±□SEM. for three independent migration experiments (C). One-
way ANOVA or unpaired Student t-test were performed to assess significance.
****p□<□0.0001. In invasion experiments, the reformed islets were immunolabelled with
antibodies against insulin (Ins; green), CD8 (red) and a nuclear stain DAPI or investigated
using light microscopy (7 D-G). Cytokine-induced destruction of reformed islets was
investigated at time points 2h (E) and 18h (F) in comparison to non-cytokine treated (control;
RPMI) reformed islets.

Figure 8.

Invasion assay illustrating the effects of 1µM fluoxetine, on reformed mouse islet infiltration
by RAW 264.7 macrophage cells. The reformed islets were incubated with 1µM fluoxetine
for 48h and cytokines and LPS-activated macrophages were added during the last 24h of
culture. Reformed islets were fixed in 4% PFA prior to imaging and fluorescence
measurements were obtained (A). The reformed islets were visualised using light
microscopy and immunoprobed with antibodies directed against CD80 (red), and DAPI was
stained blue in parallel. Scale bars represent 100µm for light microscopy and 20µm for
fluorescence microscopy. The migratory properties of RAW 264.7 cells were additionally
quantified by Cell Biolabs' CytoSelect™ cell migration and EZCellTM cell
migration/chemotaxis assays (Fig.8A). Data are expressed as mean ± SEM (n = 3
observations).   ***p<0.0006; ****p<0.001 versus cytokine control, One-way ANOVA,
Dunnett's multiple comparisons test.

Figure S1. Light microscopy images show that human islet cells formed 3-dimensional
spheroid structures after 7 days of seeding. Scale bar is 100µm (A). Immunofluorescence
detection by confocal microscopy of islet hormones in maturing mouse reformed at day 7
confirmed that developing reformed islets expressed insulin (Ins; green), glucagon (Gcg;
red) and somatostatin (Sst; purple). Scale bar is 20µm.

Figure S2. Immune population in reformed islets. (A, B) Delta Ct values distribution
graphs of Ptprc (CD45, all immune cell marker) and Adgre1(Macrophage marker) in native
and reformed mouse islets detected by quantitative PCR. Immunostaining of the reformed
islets with antibodies directed against insulin (Ins; purple) and immune markers CD4 (C),
CD8a (D) and CD3 (F) (green) confirmed the presence of both resident macrophages and
resident T-cells in reformed islets. Scale bar is 20µm.

Figure S3. Transcriptomic analyses of native and reformed human islets. (A) Heatmap
illustrating top 30 differentially expressed genes in reformed islets. (B, C, D) Heatmaps
illustrating genes differentially enriched in EMT, Focal adhesion and Skeletal development
pathway process. (E) KEGG enriched pathways related to ribosome biogenesis. (F)
Pathways enriched in Maturity Onset of the Diabetes of the Young. (G) Heatmap of mRNA
expression of islets hormones. (H) Heatmap of selected subset of islet-enriched transcription
factors between native and reformed islets.

AUTHOR CONTRIBUTIONS

NH and GAB designed and managed the studies. All authors contributed to the writing and
editing of the manuscript. NH, KT, MZ, YL carried out all the islet studies. JP isolated and
provided the T-cells. MW, MJ, GAB and TP analysed the sequencing data.

References

1. Khan, M.A.B., et al., *Epidemiology of Type 2 Diabetes - Global Burden of Disease and Forecasted Trends.* J Epidemiol Glob Health, 2020. **10**(1): p. 107-111.
2. Lin, X., et al., *Global, regional, and national burden and trend of diabetes in 195 countries and territories: an analysis from 1990 to 2025.* Sci Rep, 2020. **10**(1): p. 14790.
3. R Paul Robertson, M., *Type 2 diabetes mellitus: Prevalence and risk factors.* UpToDate, Feb 22, 2023.
4. Segerstolpe, A., et al., *Single-Cell Transcriptome Profiling of Human Pancreatic Islets in Health and Type 2 Diabetes.* Cell Metab, 2016. **24**(4): p. 593-607.
5. Carter, J.D., et al., *A practical guide to rodent islet isolation and assessment.* Biol Proced Online, 2009. **11**: p. 3-31.
6. Marchini, A., et al., *Long-term cultures of human pancreatic islets in self-assembling peptides hydrogels.* Front Bioeng Biotechnol, 2023. **11**: p. 1105157.
7. Skelin, M., M. Rupnik, and A. Cencic, *Pancreatic beta cell lines and their applications in diabetes mellitus research.* ALTEX, 2010. **27**(2): p. 105-13.
8. Kaido, T., et al., *Impact of defined matrix interactions on insulin production by cultured human beta-cells: effect on insulin content, secretion, and gene transcription.* Diabetes, 2006. **55**(10): p. 2723-9.
9. Phelps, E.A., et al., *Advances in pancreatic islet monolayer culture on glass surfaces enable super-resolution microscopy and insights into beta cell ciliogenesis and proliferation.* Sci Rep, 2017. **7**: p. 45961.
10. Weinberg, N., et al., *Lineage tracing evidence for in vitro dedifferentiation but rare proliferation of mouse pancreatic beta-cells.* Diabetes, 2007. **56**(5): p. 1299-304.
11. Lin, J.Y., et al., *In vitro expansion of pancreatic islet clusters facilitated by hormones and chemicals.* Cell Discov, 2020. **6**: p. 20.
12. Hering, B.J., et al., *Phase 3 Trial of Transplantation of Human Islets in Type 1 Diabetes Complicated by Severe Hypoglycemia.* Diabetes Care, 2016. **39**(7): p. 1230-40.
13. *Scientific Summary of the Collaborative Islet Transplant Registry (CITR)*, T.a. report, Editor. May 2019.
14. Liu, B., et al., *The selective serotonin reuptake inhibitor fluoxetine has direct effects on beta cells, promoting insulin secretion and increasing beta-cell mass.* Diabetes Obes Metab, 2022. **24**(10): p. 2038-2050.
15. Li, P., et al., *Comparative Proteomic Analysis of Polarized Human THP-1 and Mouse RAW264.7 Macrophages.* Front Immunol, 2021. **12**: p. 700009.
16. Justus, C.R., et al., *In vitro cell migration and invasion assays.* J Vis Exp, 2014(88).
17. Taciak, B., et al., *Evaluation of phenotypic and functional stability of RAW 264.7 cell line through serial passages.* PLoS One, 2018. **13**(6): p. e0198943.
18. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.
19. Korotkevich, G., *Fast gene set enrichment analysis.* BioRxiv, 2019.
20. Gu, Z., R. Eils, and M. Schlesner, *Complex heatmaps reveal patterns and correlations in multidimensional genomic data.* Bioinformatics, 2016. **32**(18): p. 2847-9.

21. Xu, G., et al., *Exendin-4 stimulates both beta-cell replication and neogenesis, resulting in*
increased beta-cell mass and improved glucose tolerance in diabetic rats. Diabetes, 1999.
48(12): p. 2270-6.
22. Benninger, R.K. and D.W. Piston, *Cellular communication and heterogeneity in pancreatic*
islet insulin secretion dynamics. Trends Endocrinol Metab, 2014. **25**(8): p. 399-406.
23. Henquin, J.C., et al., *Dynamics of glucose-induced insulin secretion in normal human islets.*
Am J Physiol Endocrinol Metab, 2015. **309**(7): p. E640-50.
24. Liberzon, A., et al., *The Molecular Signatures Database (MSigDB) hallmark gene set*
collection. Cell Syst, 2015. **1**(6): p. 417-425.
25. Liberzon, A., et al., *Molecular signatures database (MSigDB) 3.0.* Bioinformatics, 2011.
27(12): p. 1739-40.
26. Moreno-Amador, J.L., et al., *Epithelial to mesenchymal transition in human endocrine islet*
cells. PLoS One, 2018. **13**(1): p. e0191104.
27. Hayashi, Y., et al., *Downregulation of rRNA transcription triggers cell differentiation.* PLoS
One, 2014. **9**(5): p. e98586.
28. Taneera, J., P. Storm, and L. Groop, *Downregulation of type II diabetes mellitus and maturity*
onset diabetes of young pathways in human pancreatic islets from hyperglycemic donors. J
Diabetes Res, 2014. **2014**: p. 237535.
29. Nishimura, W., H. Iwasa, and M. Tumurkhuu, *Role of the Transcription Factor MAFA in the*
Maintenance of Pancreatic beta-Cells. Int J Mol Sci, 2022. **23**(9).
30. Randeria, S.N., et al., *Inflammatory cytokines in type 2 diabetes mellitus as facilitators of*
hypercoagulation and abnormal clot formation. Cardiovasc Diabetol, 2019. **18**(1): p. 72.
31. Tsalamandris, S., et al., *The Role of Inflammation in Diabetes: Current Concepts and Future*
Perspectives. Eur Cardiol, 2019. **14**(1): p. 50-59.
32. de Candia, P., et al., *Type 2 Diabetes: How Much of an Autoimmune Disease?* Front
Endocrinol (Lausanne), 2019. **10**: p. 451.
33. Feuerer, M., et al., *How punctual ablation of regulatory T cells unleashes an autoimmune*
lesion within the pancreatic islets. Immunity, 2009. **31**(4): p. 654-64.
34. Mathis, D., L. Vence, and C. Benoist, *beta-Cell death during progression to diabetes.* Nature,
2001. **414**(6865): p. 792-8.
35. Li, Y., et al., *Revisiting the Antigen-Presenting Function of beta Cells in T1D Pathogenesis.*
Front Immunol, 2021. **12**: p. 690783.
36. Roep, B.O., et al., *Type 1 diabetes mellitus as a disease of the beta-cell (do not blame the*
immune system?). Nat Rev Endocrinol, 2021. **17**(3): p. 150-161.
37. Lehuen, A., et al., *Immune cell crosstalk in type 1 diabetes.* Nat Rev Immunol, 2010. **10**(7): p.
501-13.
38. Atkinson, M.A., et al., *Current concepts on the pathogenesis of type 1 diabetes--*
considerations for attempts to prevent and reverse the disease. Diabetes Care, 2015. **38**(6):
p. 979-88.
39. Ehses, J.A., et al., *Increased number of islet-associated macrophages in type 2 diabetes.*
Diabetes, 2007. **56**(9): p. 2356-70.
40. van Gurp, L., et al., *Generation of human islet cell type-specific identity genesets.* Nat
Commun, 2022. **13**(1): p. 2020.
41. Lorza-Gil, E., et al., *FFA2-, but not FFA3-agonists inhibit GSIS of human pseudoislets: a*
comparative study with mouse islets and rat INS-1E cells. Scientific Reports, 2020. **10**(1): p.
16497.
42. Bowe, J.E., et al., *The permissive effects of glucose on receptor-operated potentiation of*
insulin secretion from mouse islets: a role for ERK1/2 activation and cytoskeletal remodelling.
Diabetologia, 2013. **56**(4): p. 783-91.
43. Fusco, J., et al., *GLP-1/Exendin-4 induces beta-cell proliferation via the epidermal growth*
factor receptor. Sci Rep, 2017. **7**(1): p. 9100.

44.   Wang, C., et al., *Exendin-4 Promotes Beta Cell Proliferation via PI3k/Akt Signalling Pathway.*
Cell Physiol Biochem, 2015. **35**(6): p. 2223-32.

45.   Rukstalis, J.M. and J.F. Habener, *Snail2, a mediator of epithelial-mesenchymal transitions,*
expressed in progenitor cells of the developing endocrine pancreas. Gene Expr Patterns,
2007. **7**(4): p. 471-9.

46.   Gershengorn, M.C., et al., *Epithelial-to-mesenchymal transition generates proliferative*
human islet precursor cells. Science, 2004. **306**(5705): p. 2261-4.

47.   Homo-Delarche, F. and H.A. Drexhage, *Immune cells, pancreas development, regeneration*
and type 1 diabetes. Trends Immunol, 2004. **25**(5): p. 222-9.

48.   Denroche, H.C., et al., *T cells accumulate in non-diabetic islets during ageing.* Immun Ageing,
2021. **18**(1): p. 8.

49.   Suk, K., et al., *IFN-gamma/TNF-alpha synergism as the final effector in autoimmune*
diabetes: a key role for STAT1/IFN regulatory factor-1 pathway in pancreatic beta cell death.
J Immunol, 2001. **166**(7): p. 4481-9.

50.   Boni-Schnetzler, M. and D.T. Meier, *Islet inflammation in type 2 diabetes.* Semin
Immunopathol, 2019. **41**(4): p. 501-513.

51.   Banu, S. and D. Sur, *Role of Macrophage in Type 2 Diabetes Mellitus: Macrophage*
Polarization a New Paradigm for Treatment of Type 2 Diabetes Mellitus. Endocr Metab
Immune Disord Drug Targets, 2023. **23**(1): p. 2-11.

52.   Ye, Z., et al., *Metabolic effects of fluoxetine in adults with type 2 diabetes mellitus: a meta-*
analysis of randomized placebo-controlled trials. PLoS One, 2011. **6**(7): p. e21551.

53.   Ghaeli, P., et al., *Comparing the effects of 8-week treatment with fluoxetine and imipramine*
on fasting blood glucose of patients with major depressive disorder. J Clin Psychopharmacol,
2004. **24**(4): p. 386-8.

54.   Zhu, S., et al., *Drug repositioning in drug discovery of T2DM and repositioning potential of*
antidiabetic agents. Comput Struct Biotechnol J, 2022. **20**: p. 2839-2847.

55.   Mali, P., et al., *RNA-guided human genome engineering via Cas9.* Science, 2013. **339**(6121):
p. 823-6.

56.   Cong, L., et al., *Multiplex genome engineering using CRISPR/Cas systems.* Science, 2013.
339(6121): p. 819-23.

Graphical Abstract

Methodology

Pancreas

① Islet isolation

② Trypsinisation of islets

③ Seeding of islet cells on laminin/collagen coated coverslips

④ Formation of reformed islet 14-16 days after seeding

Analyses

Cytoarchitecture Analysis

Functional Assays

Transcriptomic Analysis

Migration and Invasion assays

Fig. 1

Fig. 2

Fig. 2

Table 1

Charecteristics	Mouse Native Islets	Mouse Reformed Islets	Human Native Islets	Human Reformed Islets
Size	192.8±13.54	173.7±19.11	126.3±18.71	181.1±20.7
beta cells	76.53±2.96	72.64±2.54	65.79±9.16	66.38±6.39
alpha cells	15.53±3.34	8.951±1.57	26.56±2.21	26.73±1.52
delta cells	11.75±1.77	4.33±0.52	12.31±4.16	9.225±1.25

Fig. 3

Fig. 4

A

B

GSEA Pathways

C

KEGG Pathways

Fig. 5

A

B

C

| Mouse reformed islets |

Untreated — Cas-3 | Ins | Merged

Cytokines — Cas-3 | Ins | Merged

Fig. 6

A

B

Fig. 7

D

RPMI | Ins | CD8 | Merged

E

2 h

RPMI | Ins | CD8 | Merged

Cytokine cocktail | Ins | CD8 | Merged

F

18 h

RPMI | Ins | CD8 | Merged

Cytokine cocktail | Ins | CD8 | Merged

Fig. 7

G

A

B

C

Fig. 8

A

B

Islets grown from single cells-after 7 days of seeding

Fig. S1

A

Immune markers in reformed islets CD45(Ptprc)

B

Immune markers in reformed islets-F4/80 (Adgre1)

Fig. S2

A

Top 30 DEGs

Fig. S3

Fig. S3