An Environmental Restriction impairs HIV-1 virion fusion and triggers innate immune recognition

Samy Sid Ahmed[a], Liv Zimmerman[b], Andrea Imle[a], Katrin Wuebben [c], Nadine Tibroni[a], Lena Rauch-Wirth[d], Jan Münch[d], Petr Chlanda [b], Frederik Graw [c,f,g] and Oliver T. Fackler[a,e]#

[a] Department of Infectious Diseases, Integrative Virology, CIID, Heidelberg University, Medical Faculty Heidelberg, Heidelberg, Germany

[b] Schaller Research Groups, Department of Infectious Diseases, Virology, Heidelberg University, Medical Faculty Heidelberg, 69120 Heidelberg, Germany

[c] BioQuant-Center for Quantitative Biology, Heidelberg University, 69120 Heidelberg, Germany

[d] Institute of Molecular Virology, Ulm University Medical Center, 89081 Ulm, Germany

[e] German Centre for Infection Research (DZIF), Partner Site Heidelberg, Germany

[f] Interdisciplinary Center for Scientific Computing, Heidelberg University, 69120 Heidelberg, Germany

[g] Friedrich-Alexander-Universität Erlangen-Nürnberg, Department of Medicine 5, 91054 Erlangen, Germany

Running title: TLR sensing by extracellular matrix

Corresponding author at:

Department of Infectious Diseases, Integrative Virology, University Hospital Heidelberg, Im Neuenheimer Feld 344, 69120 Heidelberg, Germany Phone: ++49-(0)6221-561322; Fax: ++49-(0)6221-565003; Email: oliver.fackler@med.uni-heidelberg.de

Abstract

In vivo, HIV-1 replicates within 3D tissues, yet the impact of tissue-like environments on viral spread is largely unknown. Our previous research identified that synthetic 3D environments impose an *Environmental Restriction to cell-free Virus Infectivity* (ERVI) that diminishes HIV-1 particle infectivity. Here, mechanistic studies reveal that ERVI is implemented within minutes, saturable and induced by different adhesive tissue-like 3D matrices. ERVI reduces infectivity across a wide range of primary HIV-1 strains and virions bearing distinct viral glycoproteins but does not damage virion morphology or affect their binding to target cells. Rather, ERVI impairs virion fusion with target cells and infectivity enhancing peptide nanofibrils can restore efficient infection. In addition, ERVI sensitizes HIV-1 particles for recognition by monocyte-derived macrophages via toll-like receptors 4 and 8, triggering pronounced pro-inflammatory cytokine secretion. These results suggest that ERVI represents a broadly acting, tissue-intrinsic barrier to virus spread that reduces the fusogenicity of cell-free virions and sensitizes them for innate immune recognition.

Key words: HIV-1, tissue-like 3D environments, collagen, virion infectivity restriction, innate immune sensing

Introduction

Many viruses including Human Immunodeficiency Virus (HIV) can spread from infected donor cells either by the release of cell-free particles into the extracellular space that then diffuse to new target cells (cell-free transmission) or via close physical contacts between donor and target cell (cell-associated transmission) (Hubner *et al*, 2009; Jolly *et al*, 2004; Phillips, 1994). In the case of HIV, cell-associated transmission occurs at highly polarized cell-cell contacts referred to as virological synapse and in cell culture models, this transmission mode is significantly more efficient than infections by cell-free virus particles (Dimitrov *et al*, 1993; Iwami *et al*, 2015; Kolodkin-Gal *et al*, 2013; Sourisseau *et al*, 2007). Virological synapses have been documented in small animal models for HIV but also murine leukemia virus (Murooka *et al*, 2012; Sewald *et al*, 2015), but which transmission mode predominates in infected tissue remained unclear. To start addressing this problem, we previously established three dimensional (3D) collagen matrices as a tissue-like cell culture model in which parameters such as cell density and 3D organization can be controlled (Ahmed *et al*, 2020; Imle *et al*, 2019). At conditions of limited cell density where virions either need to diffuse or donor cells have to migrate to new target cells to sustain HIV spread, computational modelling and subsequent experimental validation identified cell-associated transmission as the predominant mode of virus spread in 3D collagen. Under such conditions, motility was found to be a relevant parameter for target cells' permissivity for infection (Lopez *et al*, 2022). In contrast, in 3D collagen with very high cell density or in classical suspension cultures, cell-free and cell-associated transmission modes supported HIV-1 spread with comparable efficacy. This shift towards cell-associated virus transmission in tissue-like environments reflected an enhanced efficiency of cell-associated spread, but also a significant reduction of the infectivity of HIV-1 particles. Imaging analysis revealed that HIV-1 particles freely diffuse in the tissue-like environment but undergo short (milliseconds) and reversible physical interactions with collagen fibers. These findings suggested that in tissue, the physical

contact with extracellular matrix limits the infectivity of cell-free HIV-1 particles. In analogy

to intracellular barriers that counteract HIV-1 spread, such as host cell restriction factors as

part of the cell intrinsic immune system, we refer to this tissue intrinsic phenomenon as

Extracellular Restriction of Virion Infectivity (ERVI).

As an enveloped virus particle, the infectious potential of HIV-1 virions is determined by a

large number of parameters. This comprises basic properties such as particle integrity,

packaging of the viral genome and essential enzymes, incorporation of the viral glycoprotein

Env, and liquid-order membrane microdomain-like lipid composition of the viral envelope

(Brügger *et al*, 2006; Nieto-Garai *et al*, 2021; Sáez-Cirión *et al*, 2002; Yi *et al*, 2006).

Reflecting this complex set of requirements for the generation of an infectious HIV particle, a

wide range of cellular restriction factors including tetherin, SERINC proteins, 90K, IFTIM

proteins, GBP 2 and 5 and PSGL-1 reduce HIV-1 infectivity via distinct mechanisms (Braun

et al, 2019; Krapp *et al*, 2016; Liu *et al*, 2019; Lodermeyer *et al*, 2013; Neil *et al*, 2008; Rosa

et al, 2015; Usami *et al*, 2015; Venkatesh & Bieniasz, 2013). In this study, we set out to gain

more insight into the mechanism and functional consequences of ERVI.

Results

ERVI is a rapidly induced, saturable and conserved restriction to HIV-1 infectivity

Our previous results had established that tissue-like 3D collagen environments pose the ERVI barrier that markedly reduces the infectivity of HIV-1 particles recovered from the supernatant of 3D collagen matrices in which virus producing cells or cell-free virus particles had been embedded (Imle *et al.*, 2019). Single particle tracking revealed that HIV virions diffuse freely in the 3D matrix but undergo physical contacts with collagen fibers. These interactions were transient in the millisecond range and did not result in coating of the fibers with virus particles (Imle *et al.*, 2019). Since physical stress can impair virus infectivity by inducing shedding of the viral glycoprotein from the virion, and the efficacy of glycoprotein incorporation is an important determinant of infectivity (Chertova *et al*, 2002; Day *et al*, 2004) we first tested if contact with 3D collagen alters the amounts of the HIV-1 glycoprotein Env in these particles. However, Env protein levels were comparable between virions cultured in suspension or in 3D collagen (EV 1A, B). To gain more insight into the nature of ERVI, we conducted a kinetic analysis and compared the impact of culturing HIV-1 NL4.3 particles in suspension or in 3D collagen with different densities (dense: 3 mg/ml; loose: 1.6 mg/ml) (Fig. 1A). Virion infectivity was assessed on Tzm-bl reporter cells, in which the luciferase gene is under control of the HIV-1 promoter and de novo expression of the viral transactivator Tat in productively infected cells triggers luciferase expression. The amount of luciferase expression relative to the amounts of virus used for infection, as determined by quantifying the activity of viral reverse transcriptase by the SG-PERT assay, yielded the relative infectivity of HIV particles. In line with our previous findings, single rounds of infection revealed that culturing HIV-1 in dense or loose 3D collagen for 16h reduced their infectivity to 10.6% or 14.3% of the particles from parallel suspension cultures (Fig. 1B). Kinetic analysis revealed that infectivity was reduced immediately or within 4h after contact with dense 3D collagen in

dense and loose collagen, respectively. The subsequent reduction of the remaining virion

infectivity over time followed comparable kinetics under all culture conditions (Fig. 1C).

To assess how conserved the sensitivity to ERVI is among primary lentiviruses, we

next analyzed the impact of 3D collagen on a panel of lab-adapted and primary HIV-1 strains

as well as one HIV-2 strain on Tzm-bl reporter cells (Fig. 1D). The results revealed that the

infectivity of all HIV strains tested was significantly reduced by ERVI, but to varying

magnitude ranging from strong (52.9 fold, HIV-1 ADA, dense collagen) to very mild

inhibition (1.6 fold, HIV-1 RHGA, loose collagen). The reduction of infectivity by ERVI was

independent of the HIV-1 entry co-receptor preference but patient-derived transmitted

founder and chronic HIV-1 variants tended to be less sensitive to ERVI than lab-adapted

HIV-1. Notably, also HIV-1 virions lacking Env but pseudotyped with the glycoprotein of

vesicular stomatitis virus (VSVG) were sensitive to ERVI. To analyze further the impact of

the viral glycoprotein for the sensitivity to ERVI, we pseudotyped GFP encoding lentiviral

particles with the unrelated glycoproteins of the HCV isolates Con1 or JFH1 and tested their

infectivity on Huh 7.5 cells by flow cytometry-based quantification of GFP expressing cells

(Fig. 1E). While the infectivity of these virions was also reduced, the effects were moderate

and only reached statistical significance for particles pseudotyped with JFH1 glycoprotein in

dense collagen. The analysis of lentiviruses pseudotyped with the Influenza HA or the Ebola

virus glycoproteins was not informative since already 16h culture in suspension reduced the

infectivity of these particles below the detection limit (EV 1C,D). Together, these results

suggest that the viral glycoprotein is a key determinant for the sensitivity to ERVI. Reduction

of virus infectivity is observed with glycoproteins of different receptor specificity and

topology but type I (HIV) and type III (VSVG) glycoproteins may be more sensitive to ERVI

than type II glycoproteins (HCV).

Since cellular restrictions to virus infection typically act as physical barriers that can

be overcome by an excess of virus particles, we analyzed if the amounts of virions added to a

constant culture volume affected the magnitude of reduction in virion infectivity (Fig. 1F).

While the relative infectivity of virus particles in suspension was unaffected by the

concentration of virus, the infectivity reduction by ERVI was significantly less efficient at

higher virus concentrations in dense or loose collagen (Fig. 1F; relative infectivity 7- or 2-fold

higher at 10^7 vs. 10^5 BCUs in dense or loose collagen; $p<0.09$ and $p<0.015$ respectively).

We next sought to define how the architecture and biophysical properties of the 3D

matrix affect its ability to reduce the infectivity of HIV-1 virions and compared several

matrices that (i) can be polymerized without harming *per se* the infectivity of HIV particles

(e.g. high temperature or UV exposure) and (ii) result in pore size that allows HIV-1 particles

to diffuse within and out of the 3D matrix (Fig. 1G, EV 1E). The dense and loose collagen

matrices are assembled into fibers polymerized from purified type I collagen proteins, the

most abundant collagen type in tissue. Confocal reflection microscopy analysis of these

matrices confirmed the different density of both types of type I collagens after polymerization

and revealed that the dense collagen was enriched in branched collagen bundles (EV 1E).

Similarly, the architecture of 3D matrices made of type III collagen, the second most

abundant fibrillary collagen in tissue that is synthesized by reticular cells and lines e.g.

vasculature, resembled that of dense collagen, albeit with shorter and thinner collagen

bundles, and type III collagen reduced HIV-1 infectivity with similar efficacy. We also tested

Matrigel, a complex extracellular environment of the basal membrane rich in collagen I as

model for type I collagen fibers in a complex tissue environment. Matrigel assembles into 3D

matrices with higher pore sizes and heterogeneity than purified collagen matrices (Anguiano

et al, 2020). Although matrigel displayed a punctate morphology, its antiviral activity was

comparable to that of dense collagen (24.1+/- 4.4% of suspension or 4-fold reduction). In

contrast, agarose hydrogels, which are made of bundles of linear filaments without cell

adhesion features, did not impair the infectivity of HIV-1 virions.

Together, these results reveal that a broad range of HIV-1 variants are sensitive to

ERVI. This extracellular restriction is exerted rapidly upon contact with the 3D environment,

can be saturated by excess of virus particles, and is exerted by a variety of adhesive

extracellular matrix components.

ERVI restricts HIV-1 infectivity without affecting structural integrity of virus particles

We next addressed in more detail how the physical contact of HIV particles with the 3D

collagen environment affects their infectivity. We considered the possibility that following

transient contacts of HIV-1 particles with collagen fibers, collagen material remains attached

to the virions and compromises their infectivity. Embedding of HIV-1 particles in

fluorescently labelled collagen (EV 2A), which exerted ERVI with similar efficacy than non-

labelled collagen (Fig. 2A), resulted in a small population of fluorescent HIV-1 particles

(8.5+/-4.7 %, Fig. 2B, C). However, these events were not frequent enough to explain the

over 10-fold reduction in infectivity observed. To assess potential physical damage of the

virions, we next analyzed their morphology by cryo-electron tomography. HIV-1 particles

kept in suspension or in dense or loose 3D collagen were placed on EM grids and processed

for cryo-ET analysis. Under all three conditions, enveloped HIV-1 particles with the typical

conical core and a diameter ranging from 104 to 154 nm (mean values: suspension: 134.5+/-

11.5 nm, dense: 135.2 +/- 9.6nm, loose: 132.0 +/- 11.6 nm) were observed (Fig. 2D, E). All

analyzed particles appeared intact without appreciable membrane rupture or deformation and

showed sparsely distributed Env spikes. We did not observe aberrations in HIV-1 particle

architecture resulting from the interaction with 3D collagen. As size comparison, we also

analyzed the morphology of dense and loose collagen fibers, which displayed the

characteristic structure of collagen fibrils, with a tight packing of D-periodic polyproline type

II helices corresponding to the spacing between individual tropocollagen monomers

(Shoulders & Raines, 2009) (Fig. 2F, EV. 2B). However, none of the images obtained for

HIV particles displayed virion-associated material reminiscent of collagen fibers (Fig. 2D),

suggesting that the detection of fluorescent HIV particles after contact with fluorescent 3D

matrices may reflect the transfer of fluorescent dye rather than collagen material. Of note,

incubating HIV-1 particles with the peptide EF-C or RM-8, which boost the infectivity of

HIV-1 particles by enhancing their interaction with target cells (Rauch-Wirth L *et al*, 2023;

Yolamanova *et al*, 2013), increased the infectivity of all particles and almost fully overcame

the inhibitory effect of ERVI (Figs. 2G, H). Together, these results reveal that ERVI does not

result from global disruption of HIV-1 particle architecture and suggest that ERVI does not

affect the intrinsic replication potential of HIV particles but rather the efficacy of their

interaction with target cells.

ERVI is manifest at the step of virus fusion to TZM-bl reporter cells without affecting

virion binding to target cells

The finding that infection enhancers boost the infectivity of HIV-1 particles subject to ERVI

suggested that ERVI acts at the early step of the viral life cycle. To test if ERVI impairs the

ability of HIV-1 particles to bind to target cells, we generated virions that incorporated

Vpr.mRuby2 during virus production for visualization (Gallucci *et al*, 2023) (Fig. 3A).

Incorporation of Vpr.mRuby2 did not affect their sensitivity to infectivity reduction upon

contact with 3D collagen (Fig. 3B). To visualize their interaction with the surface of Tzm-bl

target cells by spinning disk microscopy, cells were incubated with virus particles for 2h at

4°C to avoid particle internalization and to detect individual fluorescent HIV-1 particles

attached to the cell surface (Fig. 3C, EV 3). While incubating these particles for 16 h in

suspension slightly reduced the number of virions detected at the surface of target cells (Fig.

3D, 2 +/- 0.5 bound virus particles/cell for fresh virus vs. 1.1 +/- 0.4 bound virus particles/cell

for suspension), no additional reduction in binding efficacy was observed for virions that had

been embedded in dense or loose 3D collagen. Notably however, prior contact with dense 3D

collagen resulted in significantly larger aggregates of virus particles at the surface of target

cells. Such particle aggregation was not observed upon contact with loose collagen, indicating

that this effect is not essential for ERVI. The predominant action of ERVI therefore is not at

the level of virus binding to target cells.

To assess the ability of HIV particles to fuse with target cells membranes, Vpr.Blam

containing particles were produced and used to measure the conversion of the ß-lactamase

(BLAM) substrate CCF2 to the cytosol of target cells by flow cytometry as a measure for

fusion (Cavrois *et al*, 2002; Gallucci *et al.*, 2023). The incorporation of Vpr.Blam did not

affect the sensitivity of the virions to ERVI in dense collagen, while the infectivity reduction

by loose collagen was slightly less pronounced (Fig. 4B). Analyzing the fusion capacity of

these particles revealed efficient cytosolic delivery of Vpr.Blam by HIV-1 particles kept in

suspension (10.4 +/- 2.4%), which was dependent on the viral glycoprotein Env and could be

inhibited by the HIV fusion inhibitor T20 (Fig. 4C, D). This fusion capacity was strongly

impaired for particles that had prior contact with dense 3D collagen (1.5 +/- 0.7 %). Loose

collagen (5.4 +/- 2.5%) had a less pronounced effect on the fusion capacity of HIV-1 particles

and overall, infection rates and fusion capacity under the different conditions analyzed were

moderately correlated (Fig. 4E). We conclude that the interaction of HIV-1 particles with

tissue-like environments reduces their infectivity by impairing their ability to fuse with target

cells and that ERVI may also affect additional post entry steps.

ERVI moderately reduces infection of primary CD4+ T cells and MDMs

HeLa-derived Tzm-bl cells are a convenient and widely used reporter cell to quantify the

infectivity of HIV-1 particles but cannot reflect the differences in entry binding and receptor

densities, membrane lipid composition as well as uptake pathways between different primary

target cells (Choudhry *et al*, 2006).We therefore sought to analyze the relevance of ERVI for

HIV-1 infection of primary human CD4 T cells and primary human monocyte-derived

macrophages (MDMs). To be able to use this analysis with the same virus, we employed a

HIV-1 variant that uses CCR5 as entry coreceptor (HIV-1 NL4.3 R5) (Fig. 5A). This co-

receptor tropism did not affect the sensitivity of these particles to ERVI when assessed on

Tzm-bl cells (Fig. 5B), which was again more pronounced in dense than loose collagen.

Productive infection of primary target cells was assessed by quantifying the number of cells

with intracellular p24 capsid by flow cytometry (CD4 T cells, Fig. 5C, MDMs, Fig. 5D). The

reverse transcriptase inhibitor efavirenz (EFZ) was used to define background detection of

input virus. On both CD4 T cells and MDMs, productive infection by particles with prior

contact to dense 3D collagen was significantly reduced (3.6-fold reduction to 27.5% of

suspension on CD4 T cells, 2.9-fold reduction to 34.8% of suspension on MDMs) and loose

collagen only mediated a very mild reduction that did not reach statistical significance (1.8-

fold on CD4 T cells, 1.4-fold on MDMs). These results reveal that ERVI reduces the

infectivity of cell-free HIV-1 particles on primary target cells, albeit with lower efficacy than

on e.g. TZM-bl cells.

With 27.5% infectivity remaining, the impact of ERVI on infection of primary CD4 T

cells was less pronounced than on Tzm-bl cells, where ERVI reduces virion infectivity to 14%

of that of virions in suspension culture. Our computational model of HIV-1 replication in 3D

collagen cultures of primary human mononuclear cells, which predicted that cell-associated

virus transmission largely dominates over cell-free infection in 3D collagen matrices (Imle *et*

al., 2019), was based on the value of 14% of cell-free infectivity remaining in 3D. We

therefore asked if the fact that the infectivity of these virions is higher on the primary target

cells present in these 3D cultures affects this conclusion. To this end, we revisited our

previous analyses estimating the efficacy of cell-free and cell-to-cell transmission within

suspension and 3D collagen matrices by mathematical modelling (Imle *et al.*, 2019). Varying

the parameter defining the reduced infectivity of cell-free infection within collagen compared

to suspension, we estimate that the contribution of cell-free transmission to viral transmission

within collagen continuously increases with increasing infectivity preservation (Fig. 5E).

Differences between estimates for loose and dense collagen are partly affected by technical

compensations within the fitting procedure due to model constraints, which lead to reduced

estimates of cell-to-cell transmission rates within dense collagen and thereby potentially

underestimating the contribution of cell-to-cell transmission for this environment (EV 4C).

Under these conditions, the best estimates predict that an infectivity reduction to 27.5%

results in a contribution of cell-cell transmission to overall virus spread of 60% and 30% in

loose and dense collagen, respectively. We conclude that in our primary cell 3D cultures, cell-

associated transmission remains an important driver of HIV-1 spread in 3D collagen even if

the infectivity of cell-free particles is less reduced than previously thought.

ERVI sensitizes virus particles for TLR-mediated recognition by MDMs

We tested next if ERVI has functional implications in MDMs in addition to reducing virion

infectivity. Since MDMs can exhibit efficient innate immune reactions to challenge with HIV

particles (Pierini *et al*, 2021; Rasaiyaah *et al*, 2013; Yin *et al*, 2020), we quantified the

amounts of a panel of cytokines in the supernatant of MDMs 3 days after challenge with HIV-

1 particles kept in suspension or in 3D collagen. Interestingly, prior encounter of HIV-1

particles with tissue-like 3D environments markedly and broadly altered the cytokine

response of MDMs, resulting in increased release of important pro-inflammatory cytokines

such as IL-6, IL-8 and TNFα but also G-CSF, GM-CSF, GROα, IL-17, IP-10, MCP-3,

CXCL9 and MIP-1α/β (Fig. 6B, compare suspension vs. dense and loose collagen). To gain

insight into the HIV replication intermediate as well as the MDMs sensors that respond to

HIV-1 particles subjected to ERVI, we challenged MDMs with HIV-1 particles derived from

suspension or dense collagen cultures in the presence of selective inhibitors (Fig. 6C).

Interestingly, interfering with virus fusion by the entry inhibitor T20 or with reverse

transcription by efavirenz (EFZ) did not impair the ERVI-mediated increase in the secretion

of IL-6 or IL-8 (EV 5A). Entry and early post-entry steps of the HIV-1 life cycle are thus not

required for ERVI-mediated induction of pro-inflammatory cytokine production, suggesting

that innate recognition occurs in the context of non-productive uptake of virus particles. In

line with this scenario, inhibition of the cytoplasmic DNA sensor cGAS had no effect on the

ERVI-mediated induction of IL-6 or IL-8 secretion. Similarly, pharmacological inhibition

suggested that toll-like receptors (TLR) 1/2, 3 and 9 are not involved in the recognition of

ERVI-treated HIV particles. In contrast, inhibiting the endosomal TLR-8 that recognizes

single stranded RNA (Heil *et al*, 2004), or TLR-4, which recognizes bacterial

lipopolysaccharide (LPS) but also viral proteins (Del Cornò *et al*, 2016), or the TLR signaling

adaptor MyD88 (Medzhitov *et al*, 1998) fully abrogated the induction of IL-6 production by

ERVI-treated HIV-1 (Fig. 6C). Inhibition of TLR-8 or MyD88, but not of TLR-4, also

decreased the release of IL-8 by MDMs after challenge with collagen cultured virions (EV

5A). Collectively, these results reveal that ERVI sensitizes HIV-1 particles for recognition by

TLR-8 and -4 in the context of non-productive uptake of HIV-1 particles by MDMs.

Discussion

Tissue-like environments shape the mode of HIV-1 spread towards cell-associated transmission by optimizing duration and architecture of cell-cell contacts but also by suppressing the infectivity of cell-free virus particles. Investigating the mechanism and relevance of this intrinsic antiviral property of extracellular environments, ERVI, defined that this restriction affects the function of a broad range of viral glycoproteins and impairs fusion but not binding of cell-free virus particles with target cells. Glycoprotein incorporation and virion morphology are unaffected by ERVI and the restriction can be overcome by using infectivity enhancing peptide nanofibrils, suggesting that ERVI acts directly on the fusogenicity of the glycoprotein. This may involve modification of glycoprotein conformation and/or lipid microenvironment that result from the short physical interaction of virions with the 3D matrix and may affect the positioning of the glycoprotein in the viral membrane or their ability to cluster for efficient receptor interactions (Boesze-Battaglia *et al*, 1996; Chojnacki *et al*, 2012; Watala *et al*, 2002). Viral glycoproteins with differing sensitivity to ERVI will provide valuable tools for futures studies into the molecular mechanism of the impairment of Env function by ERVI.

Assessing the impact of ERVI on MDM target cells revealed a functional consequence of this extracellular restriction in addition to reducing virion infectivity: HIV particles that underwent prior contact with tissue-like environments triggered innate immune recognition resulting in the production of proinflammatory cytokines. Since infection rates of ERVI-treated HIV particles do not correlate with cytokine production and blocking infection does not abrogate sensing, ERVI-induced innate sensing of HIV particles does not occur during productive infection. Rather, the reduced ability to fuse with target cells is likely associated with increased recognition of particles in a non-productive uptake pathway by TLR-8 and at the cell surface by TLR-4. Since most HIV particles are subject to non-productive endocytic

uptake already in the absence of ERVI (Fackler & Peterlin, 2000; Maréchal *et al*, 1998; Maréchal *et al*, 2001; Schaeffer *et al*, 2004), we hypothesize that ERVI enhances innate immune recognition by sorting of particles into an uptake pathway particularly prone to TLR-8 recognition (Fig. 7). TLR8-mediated sensing of HIV-1 genomes has been reported in a variety of cell systems and is associated with improved control of HIV replication in patients, the regulation of latency, and can be exploited by HIV to allow the productive infection of plasmacytoid dendritic cells (Gringhuis *et al*, 2010; Li *et al*, 2023; Meås *et al*, 2020; Oh *et al*, 2008). The role of TLR-4, which recognizes LPS but also viral proteins at the cell surface (Chow *et al*, 1999), in HIV-1 infection is less established. However, R5-tropic HIV-1 Env (Env), Vpr and Tat proteins have been reported to trigger TLR-4 signaling (Bahraoui *et al*, 2020; Del Cornò *et al.*, 2016; Hoshino *et al*, 2010) and in particular for the recognition of Env, ERVI may potentiate this effect. Increased innate immune recognition is also associated with the reduction of virion infectivity by the host cell factor SERINC5 (Pierini *et al.*, 2021) and the capacity of dendritic cells to adapt to tissue-like environments (Gallucci *et al.*, 2023). Regulation by the extracellular environment thus emerges as an important parameter for innate immune responses to HIV-1 infection.

Our previous computational analysis had identified cell-associated infection as the predominant HIV-1 transmission mode in 3D cultures (Imle *et al.*, 2019). This conclusion was based on virion infectivity quantifications on model cell lines and was now questioned by our experimental finding that the effect of ERVI on infection of primary human CD4 T cells is less pronounced. Varying the magnitude of infectivity impairment in the model however revealed that even at these lower levels of cell-free infectivity reduction, cell-associated infection remains an important transmission mode. This relevant role of cell-associated transmission likely reflects that the long diffusion times required for virions to reach new target cells in 3D matrices are associated with significant reduction in their infectivity. This in

turn raises the question why the production of large amounts of cell free virus particles is

maintained in lentiviral evolution. The sensitization of non-infectious particles for the

induction of pro-inflammatory cytokine responses by extracellular matrix identified in this

study may be one reason since cytokines such as TNF-α, IL-2, IL-1 and IL-6 can increase the

permissiveness of primary target cells to HIV-1 infection (Foli *et al*, 1995; Maréchal *et al*,

1999; Poli *et al*, 1994; Vyakarnam *et al*, 1990). Although the infection rates of local target

cells are reduced by ERVI, the induced innate recognition may create a replication-prone

tissue microenvironment and thereby indirectly facilitate virus spread in tissue. ERVI is

exerted by 3D collagen from different species that form fibers of distinct architectures,

suggesting that this restriction can be relevant in different HIV target tissues. Notably, chronic

immune activation and inflammation observed in HIV patients even under therapy is

associated with lymph node fibrosis, which is thought to contribute to HIV pathogenesis

(Estes, 2013). Fibrosis reflects the deposition of large amounts of collagen with altered

structural and biophysical properties (Jones *et al*, 2018), which may also impact its antiviral

ERVI activity. Collectively, our study suggests that the reduction of virus particle infectivity

coupled to their enhanced innate immune recognition constitutes a tissue-intrinsic antiviral

immune mechanism. Future studies will focus on the role of ERVI in different fibrotic and

non-fibrotic target tissues as well as in the context of virus-host evolution.

Limitations:

Our study employed various 3D matrices to define the properties of tissue-like environments

to suppress HIV particle infectivity but this does not assess directly if HIV-1 particles

generated in human tissue are subjected to ERVI. This could in principle be tested with HIV-1

particles derived from organotypic explants models, e.g. tonsil or cervix. However, it is

impossible to ensure that particles produced in such cultures were really produced in densely

packed tissue areas and underwent physical contact with tissue fibers. While both dense and

loose collagen exert ERVI, the restriction is less pronounced for loose collagen and further

mechanistic studies will be required to define the molecular basis for this difference.

Materials and Methods:

Cells. 293T cells (ATCC, CRL-3216) and TZM-bl reporter cells (courtesy of NIH AIDS Reagent Program (ARP-8129)) were cultured in Dulbecco's Modified Eagle's Medium (DMEM, Gibco) supplemented with 10% heat-inactivated fetal calf serum (FCS, Capricorn) and 1% penicillin/streptomycin (Gibco). Huh 7.5 cells were maintained in complete DMEM medium supplemented with non-essential amino acids.

Primary CD4+ T cells and MDMs. Human peripheral blood of healthy, HIV-negative donors was obtained from the blood bank HD, according to regulation by local ethics committee (S-024/2022). CD4 T cells were isolated from human peripheral blood of healthy, HIV-negative donors using the RosetteSep Human CD4 T cell enrichment kit (StemCell Technologies) according to the manufacturer's protocol. The cells were then activated with Dynabead Human T-Activator CD3/CD28 (Gibco), or in a 3x3 activation as described previously (Imle *et al.*, 2019)rd

 for 72h and cultured in Roswell Park Memorial Institute Medium (RPMI, Gibco) supplemented with 10% heat-inactivated FCS, 1% penicillin–streptomycin, and 10 ng/ml interleukin 2 (IL-2, Biomol). To generate monocyte-derived macrophages (MDMs), peripheral blood mononuclear cells (PBMCs) were isolated from buffy coats by Biocoll (Merck Biochrom) density gradient centrifugation. $CD14^+$ monocytes were then isolated from PBMCs by positive selection using magnetic beads (CD14 MicroBeads; Miltenyi Biotech) and an AutoMACS Pro Separator (Miltenyi Biotech). $1x10^5$ monocytes per well were then seeded in glass-bottom 96 well plates that were previously coated with fibronectin (2 µg/cm²) and maintained at 37°C with 5% CO_2 in complete RPMI in the presence of 5% human AB serum (Sigma-Aldrich) for differentiation into MDMs for 10-14 day (Pierini *et al.*, 2021).

Viruses. Virus stocks of replication competent HIV-1 and HIV-2 strains (pNL4.3 WT, pNL4.3 Δnef, pNL4.3 R5, ADA, CH077, CH0 58.c, CH0198, RHGA, CH0167, CH0293, HIV-2 Rod9-GFP) were generated by transfecting 293T cells (sub-confluent 15cm² dishes) with 25 μg of proviral constructs alongside 75 μl of linear polyethyleneimine (PEI, Sigma Aldrich) in Optimem medium (Gibco). Viruses containing Vpr-BlaM and Vpr-mRuby2 were similarly produced by PEI co-transfection of 293T cells using 25 μg of pNL4.3 provirus with 7.5 μg of Vpr fusion constructs. For single round HIV-1 virus production (pNL4.3 Δenv VSVg, pNL4.3 Δenv HIV-1 Env), 22μg of pNL4.3 Δenv were used, complemented with 3 μg of the respective glycoprotein expression vectors (Gallucci *et al.*, 2023; Pierini *et al.*, 2021). Two to three days post transfection, supernatants were harvested, filtered (0.45 μm) and ultracentrifuged through a 20% (w/v) sucrose cushion. Virus pellets were then resuspended in sterile filtered PBS 0.1% BSA, aliquoted and stored at -80°C. All replication competent virions were handled in a BSL-3 containment laboratory.

Lentiviral pseudotyping. To generate lentiviral stocks, sub-confluent 293T cultures (15 cm² dishes) were transfected with PEI as described above. Briefly, 22.5 μg of pWXPL-GFP lentiviral backbone were co-transfected alongside 2.3 μg of pAdvantage, 15 μg packaging vector Pax 2 and 8 μg of envelope glycoprotein encoding plasmids (VSVg, pcDNA Con1, pcDNA JFH1). Lentiviral vectors carrying Vpx_{mac239} were produced in 293 T cells by co-transfection of pWPI, pcDNA.Vpxmac239, pΔR8.9 NSDP, and VSV-G at a molar ratio of 4:1:3:1 as previously described (Pierini *et al.*, 2021). Cell culture supernatants were harvested and filtered 3 days post transfection, and further concentrated through a 20% (w/v) sucrose cushion ultracentrifugation. Aliquots were stored at -80°C. Virus titers were determined by SG-PERT analysis (see below).

Plasmids. The proviral plasmids pNL4.3 WT and Δnef have been previously described

(Fackler *et al*, 2006). The pNL4.3-R5 WT & Δnef, containing 7 point mutations in the NL4.3

Env were previously described (Bozek *et al*, 2013; Pierini *et al.*, 2021). The pNL4.3 Δenv,

and HIV-2 Rod9-GFP proviruses were previously described (Baldauf *et al*, 2012), provirus

encoding the CCR5 tropic ADA provirus were via the NIH AIDS reagent program (ARP-

416)(Gendelman *et al*, 1988). All Transmitted/Founder strains were obtained via the NIH

AIDS reagent program (pCR-XL-TOPO CH077 (ARP-11742)(Ochsenbauer *et al*, 2012),

pCR-XL-TOPO CH058.c (ARP-11856)(Ochsenbauer *et al.*, 2012), pCR-XL-TOPO_HIV-1

M subtype C CH198 (Parrish *et al*, 2013). And chronic strains (pBR322 HIV-1 M subtype B

STCO (Mlcochova et al, 2015), pBR322 HIV-1 M subtype B RHGA(ARP-

12421)(Mlcochova *et al*, 2015), pUC57_HIV-1 M subtype C CH167(ARP-13544),

pUC57rev_HIV-1 M subtype C CH293(ARP-13539)(Parrish *et al.*, 2013). The following

expression plasmids were used: pMM311 (Vpr-BlaM) (Cavrois *et al.*, 2002), the

pmRuby2.Vpr vector was previously described (McDonald *et al*, 2002) and kindly provided

by Dr. Tom Hope. The pcDNA3.1 Vpx SIVmac239-Myc and the pΔ R8.9 packaging vector

harboring a Vpx binding motif were previously described (Baldauf *et al.*, 2012).

Reagents. The following antibodies were used: Zombie Violet dye (Biolegend), anti-p24 KC-

57 antibody (Beckman Coulter), 2G12 anti-gp120 antibody (NIH HIV Reagent program:

ARP-1476). The following dyes were used: Concanavalin A-488 (Sigma Aldrich) was used a

concentration of 50 μ/ml to stain TZM-bl cell membrane for 10 min at RT in the dark, Alexa

Fluor 488 NHS Ester (Invitrogen) was used to fluorescently label loose collagen gels as

described below, Fluoromount DAPI (Thermo Sci) was used to mount coverslips and stain

cell nuclei, the CCF2-AM dye (Thermo Scientific K1023) was used to perform HIV-1 entry

assays The following reagents were used PEI (Sigma Aldrich, Servapore dialysis membranes

(3.5 kDa MWCO, SERVA).

Virus titer determination. One step Syber Green based Product Enhanced Reverse Transcriptase assay (SG-PERT) was used to assess HIV-1 virus titers as described previously (Vermeire et al 2012). Briefly, concentrated virus stocks were first diluted in PBS, or culture supernatants were directly lysed in 2x lysis buffer (50 mM KCL, 100 mM Tris-HCl pH 7.4, 40% glycerol, 0.25% Triton X-100) supplemented with 40 mU/µl Rnase inhibitor for 10 minutes at room temperature. Lysed samples were then exported from BSL-3 and further diluted 1:10 in dilution buffer (5 mM $(NH_4)_2SO_4$, 20 mM KCL, 20 mM Tris-HCl pH 8). In parallel, 10 µl per well of a dilution series of a virus standard (pCHIV, 8,09x10^8 pURT/µl) were also lysed for 10 min. All lysed samples were then incubated with 10 µl of 2x reaction buffer (1xdilution buffer, 10 mM $MgCl_2$. 2x BSA, 400 µM each dATP, dTTP, dCTP, dGTP, 1pmol of each RT forward and reverse primers, 8 ng MS2 RNA, SYBR Green 1:10000) supplemented with 0.5U of GoTaq HotStart Polymerase. RT-PCR reactions were carried out and read in a real-time PCR detector (CFX 96, Biorad) using the following program: (1) 42 °C for .20 min, (2) 95°C for 2 min, (3) 95°C for 5s, (4) 60°C for 5 s, (5) 72°C for 15 s, 80°C for 7 s, repeat steps 3 to 6 for 40 cycles with a melting curve read out as a final step. Sequence of the used primers are as follows: fwd RT primer (TCCTGCTCAACTTCCTGTCGAG) and rev RT primer (CACAGGTCAAACCTCCTAGGAATG.)

Virus infectivity determination. To assess the infectivity of virus stocks, TZM-bl reporter cells were infected as previously described (Wei *et al*, 2002). In brief, TZM-bl cells stably expressing HIV-1 entry receptors and containing luciferase and β-galactosidase genes under the control of the HIV-1 LTR promoter were infected using dilution series of concentrated virus. 72h post-infection, cells were fixed in 3% PFA, and incubated with a substrate solution (β-Gal supplemented with 200 µg/ml X-Gal) for 3h at 37°C. Blue cells were counted to determine infectious virus titers in the form of Blue Cell Units (BCUs). For virus containing

cell culture supernatants, TZM-bl cells were infected in triplicates, and were lysed 3 days post-infection using 1x lysis buffer (Promega) for 10 min. Lysates were then incubated with Luciferase substrate (Promega), and luciferase activity was measured for 5s at a Tecan Infinity luminometer.

For saturation experiments, 10^5, 10^6, or 10^7 BCUs were used for seeding or embedding in collagen matrices of equivalent volumes. The culture supernatants were then processed by SG-PERT. TZM-bl cells were infected using equivalent amounts of RT units for all conditions. Virus infectivity after seeding or embedding at different concentrations was then determined by measuring the average luciferase activity from cell lysates 3 days post infection. To assess the relative infectivity of viruses after seeding or embedding, the infectivity and RT activity of virus containing supernatants were assessed as described above. The relative infectivity of the respective supernatants was calculated as the average luciferase activity divided by the average amount of RT units measured from the culture supernatant.

Virus embedding in 3D matrices. Type I collagen matrices of different densities were polymerized as previously described (Imle *et al.*, 2019). In brief, dense collagen gels (3 mg/ml) were generated by mixing 10X MEM medium with 7.5% NaHCO$_3$ (both Gibco) and highly concentrated rat tail collagen I (Corning) at a 1:1:8 ratio on ice. Concentrated virus was then added to the neutralized and chilled collagen solution at a 1:1 ratio. 100 μl of virus containing collagen solutions were then distributed in each well in a 96 well-plate and allowed to polymerize within 15 minutes at 37°C. Loose collagen gels (1.7 mg/ml) were generated by mixing 10X MEM medium with 7.5% NaHCO$_3$ (both Gibco) and bovine skin collagen I (PureCol, Advanced Biomatrix) at a 1:1:8 ratio on ice. Concentrated virus was then added to the neutralized and chilled collagen solution at a 1:1 ratio and, and the mixture was allowed to pre-polymerize for 10 min at 37°C. 100 μl per well were then transferred to a 96-well plate. Gels were allowed to polymerize within 45 minutes at 37°C.

Human lyophilized placental type III collagen (Advance Biomatrix) was reconstituted using a

chilled 2 mM CH$_3$COOH solution on ice to reach 6 mg/ml. Type III collagen matrices (3

mg/ml) were generated by mixing 10X MEM with 7.5% NaHCO$_3$ (both Gibco) with the

reconstituted type III collagen solution at a 1:1:8 ratio on ice. Concentrated virus was then

added to the neutralized and chilled collagen solution at a 1:1 ratio. Gels were allowed to

polymerize within 30 min at 37°C.

Similarly, Matrigel (Matrigel Growth Factor Reduced Basement Membrane, Corning) gels (5

mg/ml) were generated according to manufacturer protocol. In brief, concentrated Matrigel

aliquots (8.9 mg/ml) were thawed at 4°C overnight. Thawed Matrigel was then combined

with DMEM containing concentrated virus at a 3:2 ratio on ice. 100 µl of mixture was

transferred to a 96-well plate, gels were allowed to polymerize within 45 min at 37°C.

Agarose gels were prepared by preparing a 0.8% agarose solution in PBS. The dissolved

agarose solution was combined 1:1 with DMEM containing concentrated virus. 100 µl of the

mixture was transferred to a 96-well plate, gels were allowed to polymerize at 4°C within 15

minutes.

All polymerized gels were then overlaid with pre-warmed with 100 µl DMEM and incubated

at 37°C. Culture supernatants were harvested at the indicated time points and processed for

relative infectivity determination.

Fluorescent collagen gels. Fluorescently labelled loose collagen gels were generated as

described previously (Sixt & Lammermann, 2011). In brief, 5 mg of Alexa Fluor 488 NHS

Ester dye (Invitrogen) was dissolved in 0.5 ml DMSO. The dissolved dye was then combined

with PureCol bovine skin collagen (3 mg/ml) at a 1:100 ratio on ice and stirred overnight at

4°C. After labelling, the excess dye was removed by dialysis using a 3.5 kDa molecular

weight cut off Servapore tubing (Serva), placed in 1 L of acetic acid solution (0.02 N, pH 3.9)

and stirred for one week at 4°C. The resulting collagen solution was kept at 4°C until further

use. Fluorescently labelled collagen gels were polymerized as described above, by combining labelled and unlabelled collagen solutions at a 1:10 ratio.

Western blot. For detection of gp120 on virions from supernatants of suspension and collagen cultures, supernatants were concentrated by ultracentrifugation through a 20% sucrose cushion (44000 rpm, 45 min) using an Optima LE-80k ultracentrifuge (Beckman Coulter), and lysed in 2x SDS sample buffer (10% glycerol, 6% SDS, 130 mM Tris Hcl pH 6.8, 10% β-Mercaptoethanol) and boiled for 5 min. Protein samples and were resolved with SDS-PAGE electrophoresis and blotted onto nitrocellulose membranes. Membranes were blocked in 4% milk/TBST for 1 h and were incubated with primary antibodies overnight. We used secondary antibodies conjugated to IRDye700/800 (1:20000, Rockland) for fluorescent detection with Licor (Odyssey).

Confocal reflection microscopy. Collagen gels were imaged by confocal reflection microscopy as previously described (Imle *et al.*, 2019; Wolf *et al*, 2009). Briefly, different collagen gels were generated in 15 well angiogenesis μ-slides (Ibidi) as described above. Point laser scanning confocal microscopy was then performed on a Leica SP8 microscope using an HC PL APO CS2 63x/1.4 N.A. oil immersion objective. Images were acquired using PMT detectors in reflection mode with a laser excitation at 567 nm, and a spectral detection window set between 550 nm and 570 nm wavelengths. Fluorescently stained collagen matrices were additionally imaged using a 488 nm laser.

Widefield imaging. To assess the presence of fluorescently labelled collagen at the surface of NL4.3 Vpr-mRuby2 virions, viruses were embedded in stained or unstained loose collagen matrices as described above. The culture supernatants were then harvested and ultracentrifuged through a 20% (w/v) sucrose cushion. Virus pellets were resuspended in 3%

PFA for 1h30. The fixed virus particles were then seeded on 0.01% poly-L-lysine coated

coverslips for 30 min and mounted on microscopy slides with Fluoromount DAPI

(Invitrogen). Samples were then imaged using an epifluorescence microscope (Olympus IX81

S1F-3) under a 100X oil objective (PlanApo, N.a. 1.40). Quantification of double positive

Vpr-mRuby2 and Alexa Fluor 488 signals was performed using the Spot Detector plugin of

the Icy Imaging analysis software.

Harvesting of collagen fibers for cryo-ET analysis. To assess the morphology of single

collagen fibers by cryo ET, dense or loose collagen gels were prepared as previously in 1.5 ml

Eppendorf tubes. Polymerized gels were disrupted by sonication on a Sonorex super RK 102

H sonicator for 10 sec on ice. Disrupted individual fibers were resuspended in PBS, and

further processed for cyro-ET.

Plunge freezing. Collagen fibers and supernatant for plunge freezing were collected as

described above. Holey carbon grids (Cu 200 mesh, R2/1, Quantifoil®) were plasma-cleaned

for 10 s in a Gatan Solarus 950 (Gatan). Samples were mixed with 10x concentrated 10 nm

protein A gold (Aurion) prior plunge freezing. A total volume of 3 μl was used for plunge

freezing into liquid ethane using an automatic plunge freezer EM GP2 (Leica). The ethane

temperature was set to −183 °C and the chamber to 24 °C with 80% humidity. Grids were

blotted from the back with Whatman™ Type 1 paper for 3 s. Grids were clipped into

AutoGrids™ (Thermo Fisher Scientific).

Cryo-electron tomography and tomogram reconstruction. Cryo-electron tomography was

performed using a Krios cryo-TEM (Thermo Fisher Scientific) operated at 300 keV and

equipped with a post-column BioQuantum Gatan Imaging energy filter (Gatan) and K3 direct

electron detector (Gatan) with an energy slit set to 15 eV. As a first step, positions on the grid

were mapped at 8,700× (pixel spacing of 10.64 Å) using a defocus of approximately -65 μm

in SerialEM (Mastronarde, 2005) to localize collagen fibers or HIV particles. Tilt series were

acquired using a dose-symmetric tilting scheme (Hagen *et al*, 2017) with a nominal tilt range

of 60° to -60° with 3° increments with SerialEM. Tilt series were acquired at target focus −4

μm, with an electron dose per record of 3 e$^-$/Å^2 and a magnification of 33,000× (pixel spacing

of 2.671 Å). Beam-induced sample motion and drift were corrected using MotionCor2 (Zheng

et al, 2017). Tilt series were aligned using AreTomo (Zheng *et al*, 2022) and tomograms were

reconstructed using R-weighted back projection algorithm with dose-weighting filter and

SIRT-like filter 5 in IMOD (Kremer *et al*, 1996). Tomograms were used to measure diameter

of HIV particles in IMOD. For visualization, tilt series were aligned using protein A gold as

fiducials in IMOD. Tomograms in Fig. 2 were reconstructed using R-weighted back

projection algorithm with 3DCTF, dose-weighting filter and SIRT-like filter 10. In IMOD, 15

slices of the final tomogram were averaged and Fourier filtered. The diameter of HIV

particles was measured from the outer leaflet of the viral membrane in IMOD. The average

diameter of two measurements per particle was used in the graph.

Infectivity enhancement experiments. Culture supernatants from suspension seeded or

collagen embedded virions were incubated with previously described infectivity enhancers to

overcome ERVI. In brief, equivalent amounts of RT units of virus containing supernatants

were incubated with different Peptide Nanofibrils (PNFs). EF-C (15 μg/ml) (Yolamanova *et*

al., 2013) or RM-8 (15 μg/ml) (Rauch-Wirth L *et al.*, 2023) were then incubated for 20 min at

37°C prior to infection of TZM-bl cells. Relative infectivity of the treated virions was

assessed as described previously.

Binding assay. To assess the binding capacity of collagen and suspensions treated virions,

NL4.3 Vpr mRuby2 virions were seeded or embedded as previously described. Culture

supernatants were harvested 16h later, and viral titers were quantified by SG-PERT.

Equivalent amounts of RT units were then used to infect TZM-bl cells seeded on glass

coverslips in 24-well plates 24h prior to infection. Infection was carried out for 2h at 4°C to

prevent virion internalization. The cells were then washed with PBS, and the plasma

membrane was stained with Concanavalin A Alexa Fluor-488 (Invitrogen) according to

manufacturer protocol. The samples were then fixed using 3% PFA for 1h30, and mounted on

microscopy slides using Fluoromount-DAPI (Invitrogen). Spinning disk confocal microscopy

was performed on a PerkinElmer UltraVIEW VoX microscope equipped with Yokogawa

CSU-X1 spinning disk head and Nikon TiE microscope body. An Apo TIRF 60x/1.49 N.A.

oil immersion objective and a Hamamatsu C9100-23B EM-CCD camera were used. Images

were acquired using solid state lasers with excitation at 405nm, 488nm and 561nm with

matching emission filters. Z-stacks were acquired with a z-spacing of 0.5 μM steps. To assess

levels of virus binding between conditions, Z-stacks were 3D reconstructed using the Imaris

software (Oxford instruments). Cell membranes and virions were segmented as individual

surfaces, and statistic values (surface volumes, relative distance of surfaces) were exported by

the software.

Vpr-BlaM entry Assay. TZM-bl reporter cells were infected at MOI 0.1 using HIV-1

pNL4.3 viruses containing Vpr-BlaM, seeded in suspension or embedded in rat tail or bovine

collagen for 16 hours. Concentrated virus that was not seeded in medium prior to infection

was used as positive control, virus particles lacking Env were used as negative control. Fusion

of HIV-1 particles was allowed to proceed for 4 h at 37°C. Cells were then washed twice in

PBS and stained with 2mM CCF2-AM dye (Invitrogen) supplemented with 2.5 mM

Probenecid in Fluorobrite DMEM 2% FCS for 6 h at 11°C to prevent particle fusion during

the staining process according to manufacturer instructions. Where indicated, T20 was added

to block fusion. Cells were then trypsinized fixed in 3% PFA in PBS at 4°C overnight and

analyzed by FACS.

Primary CD4+ T cell infection. Activated primary CD4+ T cells were infected in 96-well

plated in triplicate with suspension and collagen culture supernatants containing NL4.3 R5 as

previously described (Imle *et al.*, 2019). Briefly, equivalent RT units were used between the

different conditions, as quantified by SG-PERT. The cells were spin-infected at 2000 rpm for

1h30 in presence or absence of 3 µg/ml reverse transcriptase inhibitor Efavirenz (EFZ), then

cultured at 37°C for 3 days. The samples were then stained with a fixable Zombie Violet dye

(Biolegend), fixed in 3% PFA for 1h30, then stained with an anti-p24 KC-57 FITC antibody

(Beckman Coulter) according to manufacturer protocol. Samples were then measured by flow

cytometry.

Primary MDM infection. Differentiated macrophages were spin transduced at 37°C in a pre-

heated centrifuge with lentiviral vectors containing Vpx$_{mac239}$ for 1h at 300 rpm. Within 16h

post-transduction, cells were infected in triplicates with equivalent amounts of RT units as

measured by SG-PERT from suspension and collagen culture supernatants in a 96-well plate

format, in presence or absence of EFZ. MDMs were also treated with 50 ng/ml LPS as a

positive control. Infected MDM culture supernatants were harvested 3 days post-infection and

further processed for cytokine analysis. 5 days post-infection, cells were harvested by

trypsinization, and stained with a fixable Zombie Violet dye (Biolegend), fixed in 3% PFA for

1h30, then stained with an anti-p24 KC-57 FITC antibody (Beckman Coulter) according to

manufacturer protocol. Infection rates were then determined by flow cytometry.

Modelling. The mathematical model that we used previously to estimate the contribution of

cell-free and cell-to-cell transmission given different environmental conditions has been

described in detail within (Imle *et al.*, 2019). In brief, the model describes the turnover and

dynamics of (un-)infected CD4+ T cells, CD8+ T cells and the viral load within the different

culture systems by ordinary differential equations. The complete set of mathematical

equations and detailed description, as well as pre-defined parameter values used within the

analyses, are given within (Imle *et al.*, 2019). In the original publication, the model was fitted

simultaneously to the data of a co-transfer experiment of infected and uninfected CD4+ T

cells into 2D suspension, and 3D loose and dense collagen environments, with environmental

restriction reducing the infectivity of cell-free virions within collagen, i.e., the transmission

parameter β_f, to only 14% of the effectivity considered within suspension, i.e., $\beta_{f,loose} =$

$\beta_{f,dense} = \eta\beta_{f,sus}$ with $\eta = 0.14$. Here, we re-performed the analysis done within (Imle *et*

al., 2019) by varying η between 0 and 1 within steps of 0.1, and also considering $\eta = 0.275$

as experimentally determined for primary target cells. Fitting was performed as described

within (Imle *et al.*, 2019) using the *optim*-function within the *R*-language of statistical

computing. Posterior distributions of parameter estimates were obtained by performing

ensemble fits for each value of η based on different starting values. Subsequent filtering steps

of fits ensured convergence of parameter estimates by excluding unreasonable dynamics of

CD4+ T cell counts and viral loads, with posterior distributions based on ~110-145 successful

fits for each value of η (see EV 4 A&B)).

Flow cytometry. Samples were measured by flow cytometry in BD FACS Celesta with BD

FACS Diva Software. Compensation controls were added for each experiment. Gating was

performed using FlowJo software 10.4.2 and data were processed in GraphPad Prism 8.4.3

software.

TLR inhibitor treatments. To determine the sensing pathway involved in collagen mediated

sensitization of virus particles for innate immune recognition, MDMs were pre-treated with

the different inhibitors prior to infection. 5 µM of cGAS inhibitor (G140, Invivogen), 8 µM of

TLR 1/2 inhibitor (Cu-CPT22, Selleckchem), 49 µM of TLR 4/6 inhibitor (GIT27, Tocris) or

10 µM of TLR 8 inhibitor (Cu-CPT9a, Invivogen) were incubated with MDMs 3h prior to

infection. The Myd-88 inhibitor (Pepinh-MYD, Invivogen) was used at 20 µM and incubated

with MDMs 4h prior to infection. Finally, MDMs were also pre-treated for 1h with a

TLR3/dsRNA complex inhibitor (Merck Millipore) 5 µM final concentration. MDMs were

also treated with 100 µM of T20 HIV-1 fusion inhibitor (Roche) or 3 µg/ml EFZ. The

different inhibitors were supplemented again during virus inoculation. The cells were then

cultured for 3 days at 37°C, and supernatants were harvested for cytokine analysis.

Cytokine quantification. The amounts of cytokines and chemokines present in cell culture

supernatants were determined by Eve Technologies Corporation using the Discovery Assay®:

Human Cytokine Array/Chemokine Array 48-Plex. Results are expressed in pg/ml of

cytokines/chemokines according to the company protein standard. Cell-free supernatants were

also analyzed for levels of Il-6, Il-8 and TNF-α by enzyme-linked immunosorbent assay

(ELISA; BD Biosciences) according to manufacturer's instructions.

Statistical analysis. Statistical analysis of datasets was carried out using Prism version 8.4.3

(GraphPad). Statistical significance was calculated using paired or unpaired one-way

ANOVA tests, as well as Wilcoxon matched paired test. Correction for multiple comparisons

are indicated in figure legends. n.s., not significant; *, $p < 0.05$; **, $p < 0.01$, ***$p<0,005$.

Acknowledgments

This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) Projektnummer 240245660 – SFB 1129 (project 8 to OTF) and 316249678 –

SFB 1279 (project A03 to JM). We are grateful to Katharina Morath, Swetha Ananth,

Christine Selhuber Unkel and Ada Cavalcanti-Adam for advice and discussion and to Volker

Lohmann, Frank Kirchhoff and Tom Hope for sharing reagents.

Author Contributions

726 Conceptualization, O.T.F.; Methodology, S.S.A., F.G., Investigation, S.S.A., L.Z., A.I., N.T.,

727 K.W.; Data analysis, S.S.A., L.Z.; Bioinformatic analysis, K.W., F.G.; Writing – Original

728 Draft, O.T.F., S.S.A., L.Z. and F.G.; Writing – Review & Editing, O.T.F, S.S.A., F.G., P.C.;

729 Funding Acquisition, O.T.F.; Resources, L.R., J.M.; Supervision, O.T.F, P.C., F.G.

Declaration of Interests

732 The authors declare no competing interests.

References

References

Ahmed SS, Bundgaard N, Graw F, Fackler OT (2020) Environmental Restrictions: A New Concept
Governing HIV-1 Spread Emerging from Integrated Experimental-Computational Analysis of Tissue-
Like 3D Cultures. *Cells* 9

Anguiano M, Morales X, Castilla C, Pena AR, Ederra C, Martínez M, Ariz M, Esparza M, Amaveda
H, Mora M *et al* (2020) The use of mixed collagen-Matrigel matrices of increasing complexity
recapitulates the biphasic role of cell adhesion in cancer cell migration: ECM sensing, remodeling and
forces at the leading edge of cancer invasion. *PLoS One* 15: e0220019

Bahraoui E, Serrero M, Planès R (2020) HIV-1 Tat - TLR4/MD2 interaction drives the expression of
IDO-1 in monocytes derived dendritic cells through NF-κB dependent pathway. *Sci Rep* 10: 8177

Baldauf HM, Pan X, Erikson E, Schmidt S, Daddacha W, Burggraf M, Schenkova K, Ambiel I,
Wabnitz G, Gramberg T *et al* (2012) SAMHD1 restricts HIV-1 infection in resting CD4(+) T cells.
Nat Med 18: 1682-1687

Boesze-Battaglia K, Clayton ST, Schimmel RJ (1996) Cholesterol redistribution within human platelet
plasma membrane: evidence for a stimulus-dependent event. *Biochemistry* 35: 6664-6673

Bozek K, Lengauer T, Sierra S, Kaiser R, Domingues FS (2013) Analysis of physicochemical and
structural properties determining HIV-1 coreceptor usage. *PLoS Comput Biol* 9: e1002977

Braun E, Hotter D, Koepke L, Zech F, Groß R, Sparrer KMJ, Müller JA, Pfaller CK, Heusinger E,
Wombacher R *et al* (2019) Guanylate-Binding Proteins 2 and 5 Exert Broad Antiviral Activity by
Inhibiting Furin-Mediated Processing of Viral Envelope Proteins. *Cell Rep* 27: 2092-2104.e2010

Brügger B, Glass B, Haberkant P, Leibrecht I, Wieland FT, Kräusslich HG (2006) The HIV lipidome:
a raft with an unusual composition. *Proc Natl Acad Sci U S A* 103: 2641-2646

Cavrois M, De Noronha C, Greene WC (2002) A sensitive and specific enzyme-based assay detecting
HIV-1 virion fusion in primary T lymphocytes. *Nat Biotechnol* 20: 1151-1154

Chertova E, Bess JW, Jr., Crise BJ, Sowder IR, Schaden TM, Hilburn JM, Hoxie JA, Benveniste RE,
Lifson JD, Henderson LE *et al* (2002) Envelope glycoprotein incorporation, not shedding of surface
envelope glycoprotein (gp120/SU), Is the primary determinant of SU content of purified human
immunodeficiency virus type 1 and simian immunodeficiency virus. *J Virol* 76: 5315-5325

Chojnacki J, Staudt T, Glass B, Bingen P, Engelhardt J, Anders M, Schneider J, Müller B, Hell SW,
Kräusslich HG (2012) Maturation-dependent HIV-1 surface protein redistribution revealed by
fluorescence nanoscopy. *Science* 338: 524-528

Choudhry V, Zhang MY, Harris I, Sidorov IA, Vu B, Dimitrov AS, Fouts T, Dimitrov DS (2006)
Increased efficacy of HIV-1 neutralization by antibodies at low CCR5 surface concentration. *Biochem
Biophys Res Commun* 348: 1107-1115

Chow JC, Young DW, Golenbock DT, Christ WJ, Gusovsky F (1999) Toll-like receptor-4 mediates
lipopolysaccharide-induced signal transduction. *J Biol Chem* 274: 10689-10692

Day JR, Münk C, Guatelli JC (2004) The membrane-proximal tyrosine-based sorting signal of human
immunodeficiency virus type 1 gp41 is required for optimal viral infectivity. *J Virol* 78: 1069-1079
785

Del Cornò M, Cappon A, Donninelli G, Varano B, Marra F, Gessani S (2016) HIV-1 gp120 signaling through TLR4 modulates innate immune activation in human macrophages and the biology of hepatic stellate cells. *J Leukoc Biol* 100: 599-606

Dimitrov DS, Willey RL, Sato H, Chang LJ, Blumenthal R, Martin MA (1993) Quantitation of human immunodeficiency virus type 1 infection kinetics. *J Virol* 67: 2182-2190

Estes JD (2013) Pathobiology of HIV/SIV-associated changes in secondary lymphoid tissues. *Immunol Rev* 254: 65-77

Fackler OT, Moris A, Tibroni N, Giese SI, Glass B, Schwartz O, Kräusslich HG (2006) Functional characterization of HIV-1 Nef mutants in the context of viral infection. *Virology* 351: 322-339

Fackler OT, Peterlin BM (2000) Endocytic entry of HIV-1. *Curr Biol* 10: 1005-1008

Foli A, Saville MW, Baseler MW, Yarchoan R (1995) Effects of the Th1 and Th2 stimulatory cytokines interleukin-12 and interleukin-4 on human immunodeficiency virus replication. *Blood* 85: 2114-2123

Gallucci L, Abele T, Fronza R, Stolp B, Laketa V, Sid Ahmed S, Flemming A, Müller B, Göpfrich K, Fackler OT (2023) Tissue-like environments shape functional interactions of HIV-1 with immature dendritic cells. *EMBO Rep* 24: e56818

Gendelman HE, Orenstein JM, Martin MA, Ferrua C, Mitra R, Phipps T, Wahl LA, Lane HC, Fauci AS, Burke DS *et al* (1988) Efficient isolation and propagation of human immunodeficiency virus on recombinant colony-stimulating factor 1-treated monocytes. *J Exp Med* 167: 1428-1441

Gringhuis SI, van der Vlist M, van den Berg LM, den Dunnen J, Litjens M, Geijtenbeek TB (2010) HIV-1 exploits innate signaling by TLR8 and DC-SIGN for productive infection of dendritic cells. *Nat Immunol* 11: 419-426

Hagen WJH, Wan W, Briggs JAG (2017) Implementation of a cryo-electron tomography tilt-scheme optimized for high resolution subtomogram averaging. *J Struct Biol* 197: 191-198

Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, Lipford G, Wagner H, Bauer S (2004) Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* 303: 1526-1529

Hoshino S, Konishi M, Mori M, Shimura M, Nishitani C, Kuroki Y, Koyanagi Y, Kano S, Itabe H, Ishizaka Y (2010) HIV-1 Vpr induces TLR4/MyD88-mediated IL-6 production and reactivates viral production from latency. *J Leukoc Biol* 87: 1133-1143

Hubner W, McNerney GP, Chen P, Dale BM, Gordon RE, Chuang FY, Li XD, Asmuth DM, Huser T, Chen BK (2009) Quantitative 3D video microscopy of HIV transfer across T cell virological synapses. *Science* 323: 1743-1747

Imle A, Kumberger P, Schnellbacher ND, Fehr J, Carrillo-Bustamante P, Ales J, Schmidt P, Ritter C, Godinez WJ, Muller B *et al* (2019) Experimental and computational analyses reveal that environmental restrictions shape HIV-1 spread in 3D cultures. *Nat Commun* 10: 2144

Iwami S, Takeuchi JS, Nakaoka S, Mammano F, Clavel F, Inaba H, Kobayashi T, Misawa N, Aihara K, Koyanagi Y *et al* (2015) Cell-to-cell infection by HIV contributes over half of virus infection. *Elife* 4

Jolly C, Kashefi K, Hollinshead M, Sattentau QJ (2004) HIV-1 cell to cell transfer across an Env-induced, actin-dependent synapse. *J Exp Med* 199: 283-293

Jones MG, Andriotis OG, Roberts JJ, Lunn K, Tear VJ, Cao L, Ask K, Smart DE, Bonfanti A, Johnson P *et al* (2018) Nanoscale dysregulation of collagen structure-function disrupts mechano-homeostasis and mediates pulmonary fibrosis. *Elife* 7

Kolodkin-Gal D, Hulot SL, Korioth-Schmitz B, Gombos RB, Zheng Y, Owuor J, Lifton MA, Ayeni C, Najarian RM, Yeh WW *et al* (2013) Efficiency of cell-free and cell-associated virus in mucosal transmission of human immunodeficiency virus type 1 and simian immunodeficiency virus. *J Virol* 87: 13589-13597

Krapp C, Hotter D, Gawanbacht A, McLaren PJ, Kluge SF, Stürzel CM, Mack K, Reith E, Engelhart S, Ciuffi A *et al* (2016) Guanylate Binding Protein (GBP) 5 Is an Interferon-Inducible Inhibitor of HIV-1 Infectivity. *Cell Host Microbe* 19: 504-514

Kremer JR, Mastronarde DN, McIntosh JR (1996) Computer visualization of three-dimensional image data using IMOD. *J Struct Biol* 116: 71-76

Li Y, Wang Z, Hou Y, Liu X, Hong J, Shi X, Huang X, Zhang T, Liao X, Zhang L (2023) Novel TLR7/8 agonists promote activation of HIV-1 latent reservoirs and human T and NK cells. *Front Microbiol* 14: 1033448

Liu Y, Fu Y, Wang Q, Li M, Zhou Z, Dabbagh D, Fu C, Zhang H, Li S, Zhang T *et al* (2019) Proteomic profiling of HIV-1 infection of human CD4(+) T cells identifies PSGL-1 as an HIV restriction factor. *Nat Microbiol* 4: 813-825

Lodermeyer V, Suhr K, Schrott N, Kolbe C, Stürzel CM, Krnavek D, Münch J, Dietz C, Waldmann T, Kirchhoff F *et al* (2013) 90K, an interferon-stimulated gene product, reduces the infectivity of HIV-1. *Retrovirology* 10: 111

Lopez P, Ajibola O, Pagliuzza A, Zayats R, Koh WH, Herschhorn A, Chomont N, Murooka TT (2022) T cell migration potentiates HIV infection by enhancing viral fusion and integration. *Cell Rep* 38: 110406

Maréchal V, Arenzana-Seisdedos F, Heard JM, Schwartz O (1999) Opposite effects of SDF-1 on human immunodeficiency virus type 1 replication. *J Virol* 73: 3608-3615

Maréchal V, Clavel F, Heard JM, Schwartz O (1998) Cytosolic Gag p24 as an index of productive entry of human immunodeficiency virus type 1. *J Virol* 72: 2208-2212

Maréchal V, Prevost MC, Petit C, Perret E, Heard JM, Schwartz O (2001) Human immunodeficiency virus type 1 entry into macrophages mediated by macropinocytosis. *J Virol* 75: 11166-11177

Mastronarde DN (2005) Automated electron microscope tomography using robust prediction of specimen movements. *J Struct Biol* 152: 36-51

McDonald D, Vodicka MA, Lucero G, Svitkina TM, Borisy GG, Emerman M, Hope TJ (2002) Visualization of the intracellular behavior of HIV in living cells. *J Cell Biol* 159: 441-452

Meås HZ, Haug M, Beckwith MS, Louet C, Ryan L, Hu Z, Landskron J, Nordbø SA, Taskén K, Yin H *et al* (2020) Sensing of HIV-1 by TLR8 activates human T cells and reverses latency. *Nat Commun* 11: 147

Medzhitov R, Preston-Hurlburt P, Kopp E, Stadlen A, Chen C, Ghosh S, Janeway CA, Jr. (1998) MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways. *Mol Cell* 2: 253-258

Mlcochova P, Apolonia L, Kluge SF, Sridharan A, Kirchhoff F, Malim MH, Sauter D, Gupta RK
(2015) Immune evasion activities of accessory proteins Vpu, Nef and Vif are conserved in acute and
chronic HIV-1 infection. *Virology* 482: 72-78

Murooka TT, Deruaz M, Marangoni F, Vrbanac VD, Seung E, von Andrian UH, Tager AM, Luster
AD, Mempel TR (2012) HIV-infected T cells are migratory vehicles for viral dissemination. *Nature*
490: 283-287

Neil SJ, Zang T, Bieniasz PD (2008) Tetherin inhibits retrovirus release and is antagonized by HIV-1
Vpu. *Nature* 451: 425-430

Nieto-Garai JA, Arboleya A, Otaegi S, Chojnacki J, Casas J, Fabriàs G, Contreras FX, Kräusslich HG,
Lorizate M (2021) Cholesterol in the Viral Membrane is a Molecular Switch Governing HIV-1 Env
Clustering. *Adv Sci (Weinh)* 8: 2003468

Ochsenbauer C, Edmonds TG, Ding H, Keele BF, Decker J, Salazar MG, Salazar-Gonzalez JF,
Shattock R, Haynes BF, Shaw GM *et al* (2012) Generation of transmitted/founder HIV-1 infectious
molecular clones and characterization of their replication capacity in CD4 T lymphocytes and
monocyte-derived macrophages. *J Virol* 86: 2715-2728

Oh DY, Taube S, Hamouda O, Kücherer C, Poggensee G, Jessen H, Eckert JK, Neumann K, Storek A,
Pouliot M *et al* (2008) A functional toll-like receptor 8 variant is associated with HIV disease
restriction. *J Infect Dis* 198: 701-709

Parrish NF, Gao F, Li H, Giorgi EE, Barbian HJ, Parrish EH, Zajic L, Iyer SS, Decker JM, Kumar A
et al (2013) Phenotypic properties of transmitted founder HIV-1. *Proc Natl Acad Sci U S A* 110: 6626-
6633

Phillips DM (1994) The role of cell-cell transmission in HIV-1 infection. *AIDS*

Pierini V, Gallucci L, Stürzel CM, Kirchhoff F, Fackler OT (2021) SERINC5 Can Enhance
Proinflammatory Cytokine Production by Primary Human Myeloid Cells in Response to Challenge
with HIV-1 Particles. *J Virol* 95

Poli G, Kinter AL, Fauci AS (1994) Interleukin 1 induces expression of the human immunodeficiency
virus alone and in synergy with interleukin 6 in chronically infected U1 cells: inhibition of inductive
effects by the interleukin 1 receptor antagonist. *Proc Natl Acad Sci U S A* 91: 108-112

Rasaiyaah J, Tan CP, Fletcher AJ, Price AJ, Blondeau C, Hilditch L, Jacques DA, Selwood DL, James
LC, Noursadeghi M *et al* (2013) HIV-1 evades innate immune recognition through specific cofactor
recruitment. *Nature* 503: 402-405

Rauch-Wirth L RA, Kaygisiz K WT, Zimmermann L,, Rodriguez-Alfonso AA SD, Wiese S SL, Weil
T, Schmiedel D, Münch J (2023) Optimized peptide nanofibrils as efficient transduction enhancers for
in vitro and ex vivo gene transfer. *Front Immunol*

Rosa A, Chande A, Ziglio S, De Sanctis V, Bertorelli R, Goh SL, McCauley SM, Nowosielska A,
Antonarakis SE, Luban J *et al* (2015) HIV-1 Nef promotes infection by excluding SERINC5 from
virion incorporation. *Nature* 526: 212-217

Sáez-Cirión A, Nir S, Lorizate M, Agirre A, Cruz A, Pérez-Gil J, Nieva JL (2002) Sphingomyelin and
cholesterol promote HIV-1 gp41 pretransmembrane sequence surface aggregation and membrane
restructuring. *J Biol Chem* 277: 21776-21785

Schaeffer E, Soros VB, Greene WC (2004) Compensatory link between fusion and endocytosis of
human immunodeficiency virus type 1 in human CD4 T lymphocytes. *J Virol* 78: 1375-1383

Sewald X, Ladinsky MS, Uchil PD, Beloor J, Pi R, Herrmann C, Motamedi N, Murooka TT, Brehm
MA, Greiner DL *et al* (2015) Retroviruses use CD169-mediated trans-infection of permissive
lymphocytes to establish infection. *Science* 350: 563-567

Shoulders MD, Raines RT (2009) Collagen structure and stability. *Annu Rev Biochem* 78: 929-958

Sixt M, Lammermann T (2011) In vitro analysis of chemotactic leukocyte migration in 3D
environments. *Methods Mol Biol* 769: 149-165

Sourisseau M, Sol-Foulon N, Porrot F, Blanchet F, Schwartz O (2007) Inefficient human
immunodeficiency virus replication in mobile lymphocytes. *J Virol* 81: 1000-1012

Usami Y, Wu Y, Göttlinger HG (2015) SERINC3 and SERINC5 restrict HIV-1 infectivity and are
counteracted by Nef. *Nature* 526: 218-223

Venkatesh S, Bieniasz PD (2013) Mechanism of HIV-1 virion entrapment by tetherin. *PLoS Pathog* 9:
e1003483

Vyakarnam A, McKeating J, Meager A, Beverley PC (1990) Tumour necrosis factors (alpha, beta)
induced by HIV-1 in peripheral blood mononuclear cells potentiate virus replication. *Aids* 4: 21-27

Watala C, Waczulikova I, Wieclawska B, Rozalski M, Gresner P, Gwozdzinski K, Mateasik A,
Sikurova L (2002) Merocyanine 540 as a fluorescent probe of altered membrane phospholipid
asymmetry in activated whole blood platelets. *Cytometry* 49: 119-133

Wei X, Decker JM, Liu H, Zhang Z, Arani RB, Kilby JM, Saag MS, Wu X, Shaw GM, Kappes JC
(2002) Emergence of resistant human immunodeficiency virus type 1 in patients receiving fusion
inhibitor (T-20) monotherapy. *Antimicrob Agents Chemother* 46: 1896-1905

Wolf K, Alexander S, Schacht V, Coussens LM, von Andrian UH, van Rheenen J, Deryugina E, Friedl
P (2009) Collagen-based cell migration models in vitro and in vivo. *Semin Cell Dev Biol* 20: 931-941
Yi L, Fang J, Isik N, Chim J, Jin T (2006) HIV gp120-induced interaction between CD4 and CCR5
requires cholesterol-rich microenvironments revealed by live cell fluorescence resonance energy
transfer imaging. *J Biol Chem* 281: 35446-35453

Yin X, Langer S, Zhang Z, Herbert KM, Yoh S, König R, Chanda SK (2020) Sensor Sensibility-HIV-
1 and the Innate Immune Response. *Cells* 9

Yolamanova M, Meier C, Shaytan AK, Vas V, Bertoncini CW, Arnold F, Zirafi O, Usmani SM,
Müller JA, Sauter D *et al* (2013) Peptide nanofibrils boost retroviral gene transfer and provide a rapid
means for concentrating viruses. *Nat Nanotechnol* 8: 130-136

Zheng S, Wolff G, Greenan G, Chen Z, Faas FGA, Bárcena M, Koster AJ, Cheng Y, Agard DA
(2022) AreTomo: An integrated software package for automated marker-free, motion-corrected cryo-
electron tomographic alignment and reconstruction. *J Struct Biol X* 6: 100068

Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA (2017) MotionCor2: anisotropic
correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* 14: 331-332

1001

Figure legends

Figure 1: ERVI is Rapidly Induced, Saturable, and conserved across viruses and

different types of matrices. A. Experimental workflow. HIV-1 NL4.3 viruses were seeded in

suspension or embedded in type I collagen matrices of different densities. Supernatants of the

cultures were harvested at different time points; RT activity was measured by SG-PERT and

TZM-bl reporter cells were infected with equivalent volumes of culture supernatant to

determine virion relative infectivity. B. Relative infectivity of HIV-1 NL4.3 particles after

seeding or embedding for 16h. Data is normalized to suspension seeded virions (grey dotted

line) Significance was calculated by one-way ANOVA test after Dunnett multiple comparison

correction. C. Kinetics of ERVI. Culture supernatants were harvested at 0, 4, 8 or 16h post

seeding/embedding and relative infectivity of virions was determined as in B. D.

Conservation of ERVI across HIV-1, HIV-2 and single round lentiviral pseudotypes. Lab

adapted HIV-1 strains (NL4.3, NL4.3 ΔNef, NL4.3 R5, ADA) as well as primary isolates

(Transmitted/Founder: CH077, CH058.c and CH0198; Chronic; RHGA, CH0167, CH0293),

but also HIV-2 Rod9 GFP and VSVg pseudotyped NL4.3 ΔEnv virions were seeded in

suspension or embedded in collagen cultures for 16h as described in Fig.1.A. The relative

infectivity of the virions was then assessed by SG-PERT and TZM-bl infection as previously

described. Data is normalized to suspension condition for each virus (grey dotted line).

Significance was calculated by one-way ANOVA test after Dunnett's multiple comparison

correction. E. HCV lentiviral pseudotypes are more resistant to ERVI than VSVg

pseudotypes. Lentiviral particles were seeded/embedded and Huh 7.5 cells were transduced as

described in C. Transduced cells were quantified by flow cytometry analysis of GFP

expressing cells 5 days post transduction. Data is normalized to suspension condition for each

virus (grey dotted line). Significance was calculated by one-way ANOVA test by matched

two-way ANOVA after Tukey correction. F. Saturation of ERVI. Increasing concentrations of

virus were seeded or embedded in the equivalent volume of matrix/medium for 16h.

Supernatants were then processed to determine the HIV-1 relative infectivity as in B. Significance was calculated by matched two-way ANOVA test after Tukey correction. G. Experimental workflow. HIV-1 NL4.3 viruses were seeded in suspension or embedded in different types of 3D matrices: dense or loose type I collagen matrices, human type III collagen matrices, Matrigel or agarose gels. Supernatants of the cultures were harvested 16h post seeding/embedding; RT activity was measured by SG PERT and TZM-bl reporter cells were infected with equivalent amounts of RT units for 3 days. The cells were then lysed, and infectivity was measured by assessing luciferase activity from cell lysates. H. Collagen containing matrices, but not agarose are able to restrict HIV-1 infectivity. Relative infectivity of HIV-1 NL4.3 particles after seeding or embedding in different matrices for 16h. Data is normalized to respective suspension seeded virions (grey dotted line). Significance was calculated by one-way ANOVA test after Dunnett's multiple comparison correction. Results represent the mean ± SD of 3 independent experiments. Symbols indicate data from individual experiments. *P<0.05; **P<0.001; ***P<0.005; n.s. not significant.

Figure 2: ERVI does not result from collagen deposition or structural damages of embedded virions. A. Relative infectivity of Vpr-mRuby2 containing HIV-1 NL4.3 virions, seeded in suspension or embedded in unstained, or AlexaFluor-488 stained loose collagen matrices for 16h. Data is normalized to suspension condition (Grey dotted line). Significance was calculated by one-way ANOVA test after Dunnett's multiple comparison correction. B. Representative micrographs of virions used in A. White arrows indicate Vpr-mRuby2 and AlexaFluor-488 positive virions. C. Quantification of the frequency of double positive events shown in panel C. Significance was calculated by unpaired one-way ANOVA test after Tukey's multiple comparison correction. D. Averaged slices of a tomogram showing HIV-1 NL4.3 virions after 16h of culture in suspension (left panel), dense collagen (middle panel), or loose collagen (right panel). Scale bar = 50 nm. E. Quantification of the diameter of virus

particles treated as indicated in D. Violin plot shows individual data points with corresponding median, 25% and 75% quartiles. Significance was calculated by unpaired t-tests. F. Averaged slices of a tomogram showing disrupted dense (left panel) or loose (right panel) collagen fibres. Scale bar = 100 nm. G. Experimental workflow. HIV-1 NL4.3 virions were seeded in suspension or embedded in collagen matrices as previously described. 16h post seeding/embedding, virions were harvested and equivalent amounts of RT units were incubated with infectivity enhancing peptides for 10 minutes prior to infection of TZM-bl reporter cells. H. Relative infectivity of PNF treated virions. Results represent the mean ± SD of 3 independent experiments. Symbols indicate individual experiments. *$P<0.05$; ***$P<0.005$; n.s. not significant.

Figure 3: Collagen matrices do not affect the ability of viral particles to bind to target cells but triggers aggregation at the surface of target cells. A. Experimental workflow. HIV-1 NL4.3 Vpr-mRuby2 virions were seeded in suspension or embedded in dense or loose collagen matrices for 16h. Culture supernatants were then collected, and equivalent amounts of RT units were incubated with TZM-bl cells for 2h at 4°C. Additionally virus particles that were not seeded in suspension for 16h (control, untreated virus) were used. Cell membranes were then stained using Concanavalin-A AF-488, prior to microscopy processing. B. Relative infectivity of HIV-1 NL4.3 Vpr-mRuby2 virions after seeding/embedding for 16h. Significance was calculated by one-way ANOVA test after Dunnett's multiple comparison correction. C. Representative micrographs of cells incubated with HIV-1 NL4.3 Vpr-mRuby2 virions, either from concentrated stock, or seeded/embedded for 16h. Yellow arrows indicate Vpr-mRuby2+ spots detected at the surface of the target cells. Scale bar= 15 μm. D. Quantification of the average binding frequency of HIV-1 NL4.3 Vpr-mRuby2 virions to target cells as shown in C. Significance was calculated by one-way ANOVA test after Tukey's multiple comparison correction.E. Quantification of the average volume of the Vpr-

1080 mRuby2 spots detected as shown in C. Significance was calculated by one-way ANOVA test

1081 after Tukey's multiple comparison correction. Results represent the mean ± SD from 3

1082 independent experiment. Symbols indicate individual experiments. *P<0.05, **P<0.001,

1083 ***P<0.005; n.s. not significant.

1084

Figure 4: Collagen matrices impair viral fusion with target cells. A. Experimental

workflow. HIV-1 NL4.3 VprBlaM virions were seeded in suspension or embedded in dense

or loose collagen matrices for 16h. Culture supernatants were then collected, and equivalent

amounts of RT units were incubated with TZM-bl cells for 4h at 37°C, in the presence or

absence of the T20 fusion inhibitor. Additionally, virus particles that were not seeded in

suspension for 16h (control, untreated virus) were used. Cells were then loaded with the

CCF2-AM dye for 10h at 11°C and processed by flow cytometry. B. Relative infectivity of

HIV-1 NL4.3 VprBlaM virions 16h post seeding/embedding. Significance was calculated by

unpaired one-way ANOVA test after Dunnets's multiple comparison correction. C.

Representative flow cytometry dot plots depicting the appearance of CCF2 cleavage product

by B-lactamase 4h post-infection. Gates indicate cells in which virus entry occurred. D.

Quantification of the percentage of CCF2-product positive cells measured by flow cytometry.

Significance was calculated by one-way ANOVA test after Tukey's multiple comparison

correction. E. Reduction of HIV-1 virus entry in TZM-bl target cells correlates with their

reduced relative infectivity after ERVI. Correlation between Relative Infectivity was assessed

by linear regression. Results represent the mean ± SD from 3 independent experiments.

Symbols indicate individual experiments. *P<0.05; **P<0.001; **P<0.005, n.s.: not

significant.

1103

Figure 5: ERVI does not affect infection of primary cells but sensitizes virus particles for innate immune recognition by MDMs. A. Experimental workflow. HIV-1 NL4.3 R5 virions were seeded in suspension or embedded in dense or loose collagen matrices for 16h. Culture supernatants were then collected, and equivalent amounts of RT units were incubated with MDMs or activated CD4+ T cells, in the presence or absence of the reverse transcription inhibitor Efavirenz. CD4+ T cells were harvested 3 days post infection for p24 quantification by flow cytometry. Supernatants from MDMs were harvested 3 days post infection for cytokine analysis, and cells were harvested 5 days post infection for p24 quantification by flow cytometry. B. Relative infectivity of HIV-1 NL4.3 R5 virions 16h post seeding/embedding. Significance was calculated by unpaired one-way ANOVA test after Tukey's multiple comparison correction C. Quantification of the percentage of p24+ CD4+ T cells by flow cytometry. CD4+ T cells were activated using Dynabeads for 3 days prior to infection and treated as described in A. Data is displayed for 4 independent donors. Significance was calculated by matched two-way ANOVA test after Tukey's multiple comparison correction. D. Quantification of the percentage of p24+ MDMs by flow cytometry. Monocytes were differentiated to MDMs for 10 days and transduced with Vpx-VLPs 16h prior to infection. Cells were then treated as described in A. Graphs depict mean values $\pm$ SD for 6 donors. Significance was calculated by matched two-way ANOVA test after Tukey's multiple comparison correction. E. Estimated fraction of cells infected by cell-free transmission after 21 days for the three environmental conditions given different values for the reduced efficacy of virion infectivity by ERVI, η. The plots show the posterior distributions of estimates over ~110-140 fits per value of η with dots indicating the estimate of the best model fit (see also Materials & Methods). Results for measured values of $\eta = 0.14$ (Imle et al. 2019) and $\eta = 0.275$ (here) are shown in light colours. Results represent the mean $\pm$ SD from 3independent experiments, or 4/5 5 independent donors. *P<0.05; **P<0.001; n.s.: not significant.

1130

Figure 6: ERVI sensitizes HIV-1 virions for TLR-8 and TLR-4 innate immune

recognition. A. Experimental workflow. Monocytes were differentiated into MDMs for 10

days as described previously. MDMs were then challenged with HIV-1 NL4.3 R5 virus that

was kept in suspension or collagen, in presence or absence of TLR inhibitors, Efavirenz or

T20 for 3 days. Culture supernatants were then harvested and processed for IL-6 and IL-8

ELISA. B. Cytokine profiling of the supernatants of infected MDMs. Supernatants of infected

MDMs were harvested 3 days post infection and processed for cytokine analysis. Heatmaps

indicate Log2 fold change of cytokines in infected conditions as compared to mock infected

conditions for 2 donors. Cytokines that were induced by dense collagen cultured virions >5-

fold as compared to suspension conditions for both donors are highlighted in red. C. IL-6

concentration determined by ELISA analysis of infected MDM supernatants. MDMs were

infected with HIV-1 NL4.3 R5 kept in suspension or dense collagen in presence or absence of

antiviral drugs or PRR inhibitors as described in A. Supernatants were harvested 3 days post

infection and processed by ELISA for IL-6 concentration determination. Data is normalized to

mock infected condition. Significance was calculated by matched two-way ANOVA test after

Tukey's multiple comparison correction. Results represent the mean ± SD from 3 independent

donors. Symbols indicate individual donors. $*P<0.05$; $**P<0.001$; $***P<0.005$; n.s.: not

significant.

1149

Figure 7: Schematic model of reduction of cell-free virion infectivity and sensitization

for innate immune recognition by ERVI. Physical contact with collagen fibers reduces the

ability of HIV-1 particles to fuse with target cells, resulting in the recognition of HIV-1 Env

by TLR-4 or their uptake via an alternative pathway in which TLR-8-mediated sensing of the

HIV-1 genome occurs. Triggering of TLR-4 and -8 results in increased production of

proinflammatory cytokines. See text for details.

1156

Extended View 1: Determinants of virus sensitivity to ERVI and assessment of different matrices. Collagen embedding does not lead to Env shedding. HIV-1 NL4.3 virions were seeded in suspension or dense collagen cultures for 24h. Viruses were then collected from culture supernatants and concentrated by ultracentrifugation, and analyzed by western blot using the Licor technology. The gp120/gp41 ratio was determined by incubation of membranes with a human anti-gp120 antibody (2G12) and subsequent secondary antibodies conjugated to IRDye700/800 for fluorescent detection with Licor. Membranes were also blotted using a rabbit anti-p24 antibody as loading control. One representative blot is shown. B. Quantification of the gp120:p24 ratio from Western blots as in A. Significance was calculated by Wilcoxon matched paired test. C. Lentiviral pseudotype transduce Huh 7.5 cells more efficiently than Tzm-bl cells. Lentiviral particles were pseudotyped with type I (HA, Ebola virus GP), type II (HCV strains Con1 & JFH1) or type III (VSVg) fusion glycoproteins. TZM-bl or Huh 7.5 cells were transduced with lentiviral vectors for 5 days, and the percentage of GFP+ cells was quantified by flow cytometry. Data from one experiment is shown. D. Culture of lentiviral pseudotypes has differential effects on their infectivity. Lentiviral pseudotypes as in C. were seeded in suspension, dense or loose collagen cultures for 16h. Huh 7.5 cells were then transduced with equivalent RT units for each virus for 5 days. The percentage of GFP+ cells was then quantified by flow cytometry. Data from one experiment is shown. E. Representative confocal auto-reflection micrographs of different 3D matrices. Different matrices were polymerized (see methods section) in 15-well Ibidi slides and imaged by confocal microscopy to assess fiber morphology and density. Scale bar: 10 μm. Data is representative of one or three independent experiments. Results represent the mean ± SD from 3 independent donors. Symbols indicate individual donors. n.s.: not significant.

1181

Extended View 2: Characterization of collagen fibers. A. Representative confocal micrographs of fluorescently stained loose collagen gels. Acid dissolved loose collagen was stained using AlexaFluor-488 NHS Ester dye (see Materials & methods section). Fluorescently stained collagen gels were obtained by mixing stained and unstained collagen at a 1:10 ratio. Samples were then imaged by confocal microscopy, illuminated using a 488 nm laser for fluorescence detection, and imaged by auto-reflection. Scale bar: 10 μm. B. Low magnification image of collagen fibers on EM grids. Scale bar: 1 μm.

Extended View 3: Establishment of a workflow for automated detection of surface bound HIV-1 particles. HIV-1 NL4.3 VprmRuby2 virions were seeded in suspension or collagen cultures for 16h. Supernatants were harvested, and equivalent amounts of RT units were used to infect TZM-bl cells for 2h at 4°C. After staining of plasma membrane using Concanavalin-A 488, cells were fixed and processed for spinning disk microscopy image acquisition. The resulting z-stacks were loaded in Imaris for 3D reconstruction (left panel: raw data). Virions were segmented using the 568 channel and the "surfaces" tool in Imaris, with appropriate intensity and background thresholds (middle panel: red surfaces indicate segmented virions). Next, cells were then segmented using the DAPI and 488 channels, respectively staining nuclei and cellular membranes, by using the "cells" segmentation tool in Imaris (right panel: colored surfaces indicate single segmented cells). After segmentation, quantification of bound virus was performed by the quantification of virus surfaces that were distant no more than 0.5 μm of the nearest segmented cell surface. Quantification of the volumes of segmented virions was also performed by Imaris.

Extended View 4: **Modelling the impact of ERVI on HIV-1 spread in collagen lymphocyte cultures.** A. Filtering process of individual model fits to ensure comparability of estimates for different reduced efficacies η, used within the mathematical model shown in

(Imle et al. 2019). A sequence of different filtering steps is applied to the results of the

ensemble fits given different starting conditions only considering fits with residual sum of

squares (RSS) $<10^4$ (filter 1), disregarding fits with predictions of >6000 cells after 12.5 days

or viral concentrations $> 10^{6.8}$ RT μl^{-1} after 21 days in the supernatant for loose collagen (filter

2), disregarding fits with $>7.5\times10^4$ CD4 T cells after 5 days in suspension (filter 3), and

excluding fits with >5000 CD4 T cells within the first 5 days or $>2\times10^4$ cells after 20 days for

dense collagen. The procedure is repeated to ensure comparable number of estimates for each

value of η after the filtering for the analysis. B. Number of fits per value of η considered

within the analyses. C. Distribution of estimates for the individual parameters describing cell

infection dynamics within suspension and 3D collagen environments. Results of the best fit

for each value of η are indicated by the white dot. For the meaning of the individual

parameters and specific model equations see Imle et al., 2019.

1220

1221 **Extended View 5: ERVI induces IL-8 production by MDMs via TLR-8 sensing.** A.

1222 Supernatants form infected MDMs were harvested 3 days post infection and processed by

1223 ELISA for IL-8 concentration determination. Significance was calculated by paired one-way

1224 ANOVA after Dunnett's multiple comparison correction. Results represent the mean $\pm$ SD for

1225 3 independent donors. Symbols indicate individual donors. p*<0.05.

1226

Fig. 1: Sid Ahmed et al, 2023

Fig. 2: Sid Ahmed et al, 2023

Fig. 3: Sid Ahmed et al, 2023

Fig. 4: Sid Ahmed et al, 2023

Fig. 5: Sid Ahmed et al, 2023

Fig. 6: Sid Ahmed et al, 2023

Suspension cultured HIV-1

Collagen cultured HIV-1

Suspension culture

Type I/III Collagen culture

CCR5

CD4

vRNA

TLR-4

TLR-8

MyD88

NF-κB

Pro-inflammatory Cytokines (IL-6/IL-8/TNF-α)

Fig. 7: Sid Ahmed et al, 2023

Extended View 1: Sid Ahmed et al, 2023

A

| 488-channel | Autoreflection | Overlay |

B

Dense collagen fibers     Loose collagen fibers

Extended View 2: Sid Ahmed et al, 2023

| Raw Data | Virus segmentation | Cell membrane segmentation |

Extended View 3: Sid Ahmed et al, 2023

A Workflow for filtering process of individual model fitts

B Number of fits per ERVI reduced efficacy η

C

Extended View 4: Sid Ahmed et al, 2023

A

Extended View 5: Sid Ahmed et al, 2023