Upstream open reading frames dynamically modulate CLOCK protein translation to

regulate circadian rhythm and sleep

Yuanqiang Sun[1,#], Ke Shui[2,3,#], Qinyu Li[3], Chenlu Liu[1], Wanting Jin[1], Jian-Quan Ni[4], Jian Lu[1,*],

Luoying Zhang[3,5,*]

[1] China State Key Laboratory of Protein and Plant Gene Research, Center for Bioinformatics,

School of Life Sciences, Peking University, Beijing 100871, China

[2] College of Biomedicine and Health, College of Life Science and Technology, Huazhong

Agricultural University, Wuhan, Hubei 430070, China

[3] Key Laboratory of Molecular Biophysics of Ministry of Education, College of Life Science

and Technology, Huazhong University of Science and Technology, Wuhan, Hubei, 430074,

China

[4] Gene Regulatory Lab, School of Medicine, Tsinghua University, Beijing, China.

[5] Hubei Province Key Laboratory of Oral and Maxillofacial Development and Regeneration,

Wuhan, Hubei 430022, China

#These authors contributed equally to this work.

*Corresponding author.

Email: luj@pku.edu.cn (Jian Lu); zhangluoying@hust.edu.cn (Luoying Zhang)

Author Contributions: J.L. and L.Z. initiated the project and oversaw all aspects of the project.

With the help of Q.L. and J.N., Y.S., K.S., C.L. and W.J. performed research and analyzed data;

J.L. and L.Z. wrote the manuscript with input from all the authors.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences, Neuroscience

Keywords: circadian rhythm, *Clock*, uORF, sleep, *Drosophila*,

Abstract

The circadian rhythm is an evolutionarily conserved mechanism with translational regulation

increasingly recognized as pivotal in its modulation. In this study, we found that upstream open

reading frames (uORFs) are enriched in circadian rhythm genes, with particularly conserved

uORFs present in core circadian clock genes. We demonstrate evidence that the uORFs of the

core clock gene, *Clock* (*Clk*), rhythmically and substantially attenuate CLK protein translation

in *Drosophila*, with pronounced suppression occurring during daylight hours. Eliminating *Clk*

uORFs results in elevated CLK protein levels during the day and a compressed circadian cycle,

along with a broad shift in clock gene expression rhythms. Interestingly, *Clk* uORF deletion

also augments morning sleep by reducing dopaminergic activity. Beyond daily circadian

adjustments, *Clk* uORFs play a role in modulating sleep patterns in response to the varying day

lengths of different seasons, inhibiting translation in a day-length contingent manner.

Furthermore, the *Clk* uORFs act as a master regulator to shape the rhythmic expression of a

vast array of genes and influence multifaceted physiological outcomes. Collectively, our

research sheds light on the intricate ways uORFs dynamically adjust downstream coding

sequences to acclimate to environmental shifts.

Introduction

Circadian rhythms orchestrate ~24-hour cycles in a multitude of biological processes, with the sleep-wake cycle being a particularly salient manifestation. The molecular underpinnings of the circadian rhythms consist of a transcription-translation feedback loop (TTFL), an evolutionarily conserved mechanism evident across diverse taxa, ranging from bacteria to fungi, plants, and animals [1-4]. The circadian systems drive gene expression rhythms of thousands of genes, exerting pervasive influences on daily fundamental biological activities from the cellular level to complex behaviors and diseases such as cell proliferation, cell signaling, metabolism, rest/activity, cancer and neurodegeneration [5-7]. Despite shared rhythmic principles in the tree of life, species-specific divergence in clock components and mechanisms is evident, likely a result of distinct environmental pressures and life histories shaping unique adaptations after a single origin in a common ancestor of all life forms [4,8]. Alternatively, the difference in genetic components across different kingdoms of life implies that these rhythmic systems probably emerged separately in various lineages at an early stage via convergent evolution [9,10]. While the key genetic components that drive the circadian rhythms have been deciphered [5-7], the intricate regulatory mechanisms fine-tuning these rhythms remain partially obscured, with significant gaps in our understanding persisting.

Drosophila, with its genetic tractability and conservation of key circadian components, serves as an unparalleled model providing profound insights that resonate across species, including humans [4,11-14]. In *Drosophila*, two transcriptional activators, CLOCK (CLK) and CYCLE (CYC), constitute the positive limb of the loops. CLK and CYC heterodimerize and activate the transcription of *period* (*per*) and *timeless* (*tim*) via E-box enhancer elements [15,16]. As PER and TIM proteins accumulate in the cytoplasm, they bind to each other and translocate into the nucleus to suppress the transcriptional activities of CLK and CYC, thus repressing their own transcription and forming a pivotal negative feedback loop [17]. Throughout this process, the stoichiometry of these core regulators is precisely controlled to ensure the accuracy of the circadian rhythms. For instance, the overexpression of CLK-CYC increases their activity at *per* and *tim* E-box enhancers, consequently shortening the circadian rhythms [18,19]. Recent studies have revealed that the precision of circadian rhythms is regulated by multi-layered mechanisms beyond transcriptional modulation, which include the regulation of protein synthesis and post-translational stability. For instance, the translation of *tim* and *per* mRNAs is facilitated by TWENTY-FOUR and its activator, ATAXIN2, through their promotion of an association with polyadenylate-binding protein [20-23]. Additionally, the synthesis of CLK and TIM proteins is inhibited by the microRNAs *bantam* [24] and *mir-276a* [25], respectively, with the overexpression of either miRNA leading to prolonged periods or arrhythmicity [24-26]. Post-translational modifications (PTMs) also affect the stability and subsequent degradation of PER and TIM [1].

While recent advances have significantly enhanced our understanding of the regulatory
complexity of the classical TTFL, the field has not yet completely elucidated the intricate
network of components and interactions that govern the fine-scale regulation of circadian
rhythms, even in *Drosophila*, the quintessential model for circadian biology.

Recent studies underscore the pivotal role of translational regulation, particularly by
upstream open reading frames (uORFs), in the control of circadian rhythms in organisms with
divergent circadian architectures. uORFs, the short ORFs with start codons located within the
5' untranslated regions (5'UTRs) of eukaryotic mRNAs, are widely distributed in eukaryotic
genomes [27]. uORFs play a key role in suppressing translation initiation of the downstream
coding sequence (CDS) in an mRNA by either sequestering ribosomes or competing for binding
to translation initiation complexes [28-32]. In *Neurospora*, the central circadian clock component,
frequency (*frq*), has two isoforms—large (l-FRQ) and small (s-FRQ)—whose balance is
determined by temperature-dependent splicing and is essential for the temperature
compensation of circadian rhythms. Thermosensitive trapping of scanning ribosomes at the
uORFs of l-FRQ modulates FRQ protein levels in response to ambient temperatures, thus
calibrating the circadian clock [33]. Similarly, TIMING OF CAB EXPRESSION 1 (TOC1), a core
component of the *Arabidopsis* circadian clock, is regulated by uORFs within its mRNA leader
sequence [34]. Although *Neurospora* and *Arabidopsis* share similar design principles in circadian
rhythms as humans and flies, the proteins involved differ significantly in both structure and
conservation [35-37]. Recent studies have shown that uORFs in the mouse core clock gene *Period2*
(*Per2*) play a role in temperature entrainment of the circadian clock [38] and in regulating sleep
[39]. However, no significant differences in the period and amplitude of circadian rhythms were
observed between wild-type and *Per2* uORF mutant mice [39]. In addition, extensive ribosomal
binding to uORFs in rhythmically expressed mRNAs has been observed in mammalian cells,
yet the functional impacts of these uORFs in circadian regulation have not been characterized
[40,41]. In summary, although research into the translational regulation by uORFs is burgeoning,
the complete spectrum of regulatory functions of uORFs in circadian rhythm control remains
to be elucidated. Moreover, the potential impacts of uORF-mediated regulation on
physiological and pathological processes linked to circadian rhythms are currently elusive,
while their delineations can help reveal how circadian rhythms adapt to environmental
pressures.

Here, we show that circadian rhythm-related genes are significantly enriched with ultra-
conserved uORFs in *Drosophila*. These uORFs, which demonstrate translational activity in our
previously published translatome data [42], thus may play a crucial role in modulating circadian
rhythms. We then conducted an in-depth exploration of conserved and potentially functional

uORFs in *Clk* of *Drosophila melanogaster* and presented evidence that these uORFs substantially inhibit protein translation *in vitro* and *in vivo*. Knocking out (KO) *Clk* uORFs amplifies daytime CLK protein levels and shortens the circadian period of locomotor rhythm. Furthermore, *Clk*-uORF-KO mutants manifest increased morning sleep, likely stemming from diminished dopaminergic activity. These mutants also exhibit defects in modulating sleep duration in response to photoperiodic shifts. Molecular investigations underscore the requirement of *Clk* uORFs for photoperiod-driven regulation of CLK protein, highlighting their central role in both circadian and circannual orchestrations. Intriguingly, the mutants also display decreased fecundity and diminished resilience to starvation, indicating that the ablation of *Clk* uORFs has extensive physiological consequences beyond the regulation of circadian rhythms and sleep.

Results

Circadian rhythm genes are more likely to be regulated by uORFs

We identified 36,565 putative canonical uORFs across all isoforms of 13,470 protein-coding genes in *D. melanogaster*. Of these, 1,137 uORFs are associated with 169 circadian rhythm-related genes (GO:0007623), demonstrating a significant enrichment of uORFs in this category of genes ($P < 10^{-16}$, Fisher's exact test; Fig. 1A). Since conserved uORFs are more likely to be functional, we then focused on uORFs highly conserved across *Drosophila* species. As the start codon (uATG) is the most pivotal and definitive feature of a canonical uORF [30,43], we investigated the conservation patterns of the uATGs based on the multiple sequence alignments (MSA) of 23 *Drosophila* species. This analysis revealed 388 uATGs (within 250 genes) that are identical across the *Drosophila* species examined (Table S1). These genes are significantly enriched for functions in neuronal axon genesis and transcriptional regulation, implicating the potential roles of uORFs in modulating these processes (Fig. S1). Among the 169 circadian rhythm-related genes, 29 uORFs display conservation of uATGs, a higher ratio than that of uORFs in the remaining genes (29/1,137 vs. 359/35,428; $P = 2.3 \times 10^{-5}$, Fisher's exact test; Fig. 1B). For the ten core clock genes (Table1), we found that 7 out of 82 uORFs possess uATGs that are identical across *Drosophila* species, suggesting higher uORF conservation in the core versus other circadian genes (7/82 vs. 22/1,055; $P = 0.005$, Fisher's exact test; Fig. 1B). These seven conserved uORFs are distributed among four genes, with *Pdp1* containing four, and *Clk*, *sgg*, and *cwo* each containing one (Table 1). These patterns suggest that circadian rhythm genes, especially core clock genes, are more inclined to be regulated by uORFs.

Besides conservation, the regulatory functions of uORFs are also implied by their expression and translation patterns. Analysis of our previously published transcriptome and translatome data [42] indicates that 78% of the 1,137 uORFs in the circadian rhythm-related genes

are expressed in adult heads (mRNA RPKM > 0.1, 888 in males and 879 in females; Table 1),

and about 80% of these expressed uORFs display evidence of translation [Translation efficiency

(TE) > 0.1, 692/888 in males and 714/879 in females; Table 1]. Notably, 25 of the 29 highly

conserved uORFs exhibit translation signals in the heads of both sexes. Of the 82 uORFs from

the core clock genes, 60% show mRNA expression in adult heads (67 in males and 63 in

females), with 78% of these expressed uORFs exhibiting translation signals (52/67 in males

and 49/63 in females; Table 1), suggesting their potential functional importance.

To assess the impact of uORFs on translation efficiency of the downstream CDS, we

categorized the circadian rhythm-related genes expressed in heads (149 in male and 147 in

female) into three groups: Class I (genes with conserved uORFs that are translated, 12 in male

and 12 in female), Class II (genes with translated uORFs that are not highly conserved, 98 in

male and 100 in female), and Class III (genes without translated uORFs, 39 in male and 35 in

female). We observed that genes with translated uORFs (Class I and II) overall display a lower

TE than those without (Class III) ($P < 0.0002$ for both male and female heads, respectively,

Wilcoxon signed-rank test; Fig. 1C). While not statistically significant, Class I genes show a

tendency of lower TE compared to that of Class II in both sexes. Each of the four core clock

genes (*Clk*, *cyc*, *per*, and *tim*) demonstrated reduced TE relative to the genomic average.

Notably, *Clk* belongs to Class I and exhibits a significantly lower TE, implying a substantial

regulatory impact of uORFs on its translation (Fig. 1C).

uORFs suppress CLK and CYC translation *in vitro* and *in vivo*

In *Drosophila melanogaster*, the 5'UTR of *Clk* mRNA contains five canonical uORFs, among

which the second (uORF2) and the third uORF (uORF3) are in-frame to each other and

overlapping (Fig. 2A). Among the five uORFs of *Clk*, the start codon of uORF2 (uATG2) is the

most conserved, identical across all 23 *Drosophila* species (Fig. 2A). As CYC is the binding

partner of CLK, we also analyzed *cyc* for comparison. The 5'UTR of *cyc* mRNA contains four

canonical uORFs (SI Appendix, Fig. S2A). The uATGs of *cyc* are less conserved than those of

Clk in general, and none of them is identical across the 23 *Drosophila* species (SI Appendix,

Fig. S2A). We detected prominent mRNA expression and ribosome occupancy for uORF2 and

uORF3 of *Clk* in the head and body of male and female flies (Fig. 2B). As for *cyc*, prominent

mRNA expression and ribosome occupancy was observed for uORF2, uORF3 and uORF4 (SI

Appendix, Fig. S2B).

To further explore the regulatory impact of these uORFs on CDS translation, we performed

a dual-luciferase reporter assay *in vitro*. We engineered dual-luciferase constructs with either

wild-type (WT) or mutated 5'UTRs of *Clk* and *cyc* mRNAs, where the start codons for *Clk*

uORF2 and uORF3, as well as *cyc* uORF2, uORF3, and uORF4, were mutated from ATGs to

ATTs (Fig. 2C). In *Drosophila* S2 cells, we observed that the luminescence intensity, indicative

of CDS translational activity, was more than 70% higher in constructs with mutated 5'UTRs

compared to WT controls (WT vs. mutant: 1 vs. 1.79 and 1 vs. 1.83 for *Clk* and *cyc* 5'UTR,

respectively; $P < 0.001$ for 5'UTRs of both *Clk* and *cyc*; Fig. 2C). This significant increase

confirms the repressive function of these uORFs on the translation of downstream CDSs.

To investigate the function of *Clk* and *cyc* uORFs *in vivo*, we knocked out *Clk* uORF2 and

uORF3 simultaneously by mutating their start codon ATGs to ATTs, and established a

homozygous knock-out line (*Clk*-uORF-KO) (Fig. 2D). Similarly, we mutated the ATG start

codons of uORF2-4 to ATTs in *cyc* and generated a homozygous knock-out line (*cyc*-uORF-

KO) (Fig. 2D). We carried out ribosome fractionation followed by quantitative PCR (qPCR) to

compare the mRNA abundance in the polysome fraction to that in the monosome fraction (P-

to-M ratio) for *Clk* and *cyc*, respectively (Fig. S3). A larger P-to-M ratio means more mRNAs

are enriched in the polysome fractions and bound by more ribosomes, thus indicative of higher

translation efficiency [44-46]. We conducted experiments on male heads from *Clk*-uORF-KO, *cyc*-

uORF-KO, and WT flies at various Zeitgeber times (ZT), starting from ZT0 (the onset of lights)

to ZT20 at 4-hour intervals. P-to-M ratios of *Clk* mRNA in *Clk*-uORF-KO flies significantly

exceeded WT at ZT0, ZT4, and ZT8, with no notable changes at ZT12 and ZT16, and a decrease

at ZT20 (Fig. 3A). Conversely, P-to-M ratios for *cyc* mRNA in *cyc*-uORF-KO flies were

slightly higher at all but one ZT time point (ZT12) (Fig. 3B). Notably, the elevation in *cyc*

mRNA was not statistically significant except at ZT16. Overall, these patterns indicate *Clk*

uORFs exert a substantial impact on translation, especially during the light phase, while *cyc*

uORFs show modest effects. Consequently, our research has pivoted to focus on *Clk* uORFs.

Next, we measured CLK protein abundances for male and female heads at different time

points using immunoblotting (Fig. 3C and 3D). We found that CLK protein levels were

significantly higher in the heads of male *Clk*-uORF-KO flies compared to WT during the

morning hours of ZT0, ZT4, and ZT8, and marginally lower, though not significantly, at ZT16

and ZT20 (Fig. 3E). Similar patterns were observed for female heads (Fig. 3F). These protein

abundance trends corroborate the enhanced translational activity in *Clk*-uORF-KO mutants

compared to WT during the light phase. Overall, the data demonstrate that *Clk* uORFs

downregulate CLK protein translation during the daytime, with pronounced effects observed in

the early hours.

Clk uORFs modulate the pace of the circadian clock

To quantitatively assess the effects of *Clk* uORF KO on circadian rhythms, we utilized a

mathematical model previously established for simulating the *Drosophila* molecular clock [47].

This model employs nonlinear ordinary differential equations to model the regulatory weights

among mRNA and protein molecules within the circadian network, making it applicable for

predicting the increase in *Clk* translation when uORFs are disrupted. By incrementing the

regulatory weight from *Clk* mRNA to CLK protein from the baseline to higher values, we

incorporated the anticipated enhancement in *Clk* translation due to the uORF KO into our model

(Fig. 4A).

The simulation indicated an immediate increase in CLK protein levels correlating with a

higher regulatory weight from *Clk* mRNA to protein, which mimics the effect of *Clk* uORF KO

(Fig. S4A). This, in turn, led to elevated levels of *per* and *tim* mRNA and protein (Fig. S4A and

S4B). In the simulations, we analyzed the intervals between peaks of *tim* mRNA, using these

as a measure of circadian period length. The mean ± SEM of period length is 23.926 ± 0.003

hours, 20.967 ± 0.078 hours, 18.956 ± 0.061 hours, and 17.494 ± 0.043 hours for regulatory

weights from *Clk* mRNA to CLK protein set at 0.026 (baseline), 0.027, 0.028, and 0.030,

respectively (Fig. 4B and 4C; SI Appendix, Table S2). These findings demonstrate a shortening

of the circadian period as the regulatory weight increases, suggesting that enhanced *Clk*

translation—resulting from the disruption of *Clk* uORFs—compresses the circadian period.

To corroborate the simulation outcomes, we measured the mRNA level of *per* and *tim*,

observing significant elevation in *Clk*-uORF-KO flies with the peak level occurring at an earlier

time in female heads (Fig. 4D). As our simulations predicted, disrupting *Clk* uORFs led to a

reduced period length by over one hour for the locomotor rhythm of *Clk*-uORF-KO flies.

Specifically, the period's mean ± SEM for WT males is 24.0 ± 0.25, WT females 24.2 ± 0.14,

Clk-uORF-KO males 22.8 ± 0.21, and *Clk*-uORF-KO females 22.7 ± 0.09 (Fig. 4E and 4F; SI

Appendix, Table S3). These empirical results, together with our simulation data, strongly

indicate that *Clk* uORFs modulate CLK translation, thereby regulating the timing of the

molecular clock through the expression of downstream genes like *per* and *tim*.

***Clk* uORFs influence the temporal transcriptomic landscape**

CLK modulates the cyclical expression of a significant portion of *Drosophila* transcriptome [48].

To explore the impact of *Clk* uORFs on the global gene expression rhythms, we harvested heads

from WT and *Clk*-uORF-KO flies at four-hour intervals across the day, using three biological

replicates, and performed RNA sequencing for transcriptome analysis. Using JTK-CYCLE [49]

to detect genes with oscillating mRNA levels (Table S4), we found 3,370 and 3,859 genes

exhibited rhythmic expression in WT and *Clk*-uORF-KO female heads, respectively, with 2,235

genes in common (Fig. 5A and 5B). In male heads, 3,570 genes showed rhythmic expression

in WT, whereas 3,890 were rhythmic in *Clk*-uORF-KO, with 2,124 genes in common (Fig. 5C;

SI Appendix, Fig S5). Overall, knocking out *Clk* uORFs disrupted the rhythmic expression of

about 34-41% of genes in WT, while approximately 1,700 genes displayed *de novo* cyclic

expression patterns in *Clk*-uORF-KO heads. Gene ontology (GO) analysis demonstrated that

genes which lost rhythms in *Clk*-uORF-KO flies were mainly involved in rRNA processing,

ribosome biogenesis, translation and metabolism (SI Appendix, Fig. S6A and S6C), while genes

that gained rhythms when *Clk* uORFs were knocked out were enriched in pathways such as

translation, mitochondrial processes and development (SI Appendix, Fig. S6B and S6D).

Remarkably, genes related to circadian rhythms were significantly overrepresented

among rhythmically expressed genes in *Clk*-uORF-KO female heads (86 out of 150) compared

to the rest of the expressed genes (3,773 out of 9,051, Fisher's exact test, P = 0.02). This

enrichment was more significant in the heads of *Clk*-uORF-KO females than in WT females,

WT males, or *Clk*-uORF-KO males. Knocking out *Clk* uORFs led to an increase in oscillation

amplitude which was more significant in females than in males (Wilcoxon rank sum test; P =

5.72×10^{-9} in females vs. P = 1.29×10^{-5} in males; Fig. 5D and 5E). Furthermore, in females,

knocking out *Clk* uORFs led to desynchronized oscillations with peak expression time

distributed throughout the day (Fig. 5F and 5G). In summary, disrupting *Clk* uORFs markedly

influences gene expression rhythms in fly heads, exerting more pronounced effects in females

than males. These alterations in gene expression could result in diverse impacts on

physiological functions, especially in females, aspects of which are investigated in the

following sections.

***Clk* uORFs promote morning wakefulness by tuning up the dopaminergic tone**

Building on evidence that CLK promotes sleep [50], we first examined the impact of knocking

out *Clk* uORFs on sleep. We found that knocking out *Clk* uORFs significantly lengthened sleep

duration, with more prominent alteration in females (9.5% increase; Fig. 6A; SI Appendix, Fig.

S7A) compared to males (4.0% increase; SI Appendix, Fig. S7B and S7C). A closer

examination showed that in female *Clk*-uORF-KO flies, the duration of sleep was notably

prolonged by 38.65% during the initial 8 hours of the light phase, indicating that *Clk* uORFs

facilitate morning wakefulness (Fig. 6B). The increase in sleep duration stems from a longer

average sleep bout, implicating enhanced sleep continuity upon *Clk* uORF deletion (Fig. 6C).

Additionally, waking activity levels were higher in female *Clk*-uORF-KO flies, indicating that

the prolonged sleep is not due to impaired motor function (Fig. 6C). While in males, the

alterations in sleep were not as prominent as that observed in females (SI Appendix, Fig. S7D).

Given that CLK suppresses expression of the gene encoding tyrosine hydroxylase (*ple*),

which in turn diminishes dopamine signaling and activity levels [51], we investigated whether

dopamine signaling mediates the influences of *Clk* uORFs on sleep. Our qPCR assays revealed

a notable decrease in *ple* mRNA in female *Clk*-uORF-KO heads compared to WT (Fig. 6D),

aligning with the hypothesis that elevated CLK protein levels—resulting from *Clk* uORF

disruption—downregulate *ple* expression. Next, we treated female flies with dopamine

precursor L-DOPA, which facilitates dopamine synthesis [52]. This treatment reversed the

prolonged sleep duration in female *Clk*-uORF-KO flies (Fig. 6E and 6F). Thus, *Clk* uORFs

appear to promote morning alertness by modulating dopaminergic signaling.

Clk uORFs limit starvation-induced wakefulness and are beneficial for survival

Starvation typically reduces sleep through a mechanism dependent on CLK, a response thought

to enhance survival chances [53]. To determine if *Clk* uORFs modulate the effects of starvation

on sleep/wakefulness, we analyzed sleep patterns in female *Clk*-uORF-KO flies during

starvation. We observed an exaggerated sleep loss in *Clk*-uORF-KO flies, suggesting that *Clk*

uORFs may function to restrict starvation-induced sleep loss (Fig. 6G-I). Moreover, the

increased wakefulness in *Clk*-uORF-KO flies correlated with significantly faster mortality rates

under starvation compared to WT flies for both sexes [$P = 8.8 \times 10^{-13}$ for females (Fig. 6J) and

$P = 2.4 \times 10^{-9}$ for males (SI Appendix, Fig. S8), log-rank test]. These results indicate that *Clk*

uORFs function to restrict sleep loss during starvation, which potentially affects survival.

Clk uORFs modulate sleep/wakefulness in adaptation to seasonal photoperiodic changes

by suppressing CLK protein translation in a photoperiod-dependent manner

Given the known significance of the circadian clock in adapting to seasonal environmental

variations [54], we hypothesized that *Clk* uORFs might similarly regulate translation on a

circannual basis. To test this hypothesis, we subjected female *Clk*-uORF-KO and WT flies to

conditions that mimic seasonal changes. Because the most important seasonal signal for the

circadian clock is the shortening/lengthening of photoperiod [54], we monitored locomotor

rhythm and sleep under photoperiods of different lengths, ranging from 4 hours of light and 20

hours of dark (4L20D) to 20 hours of light and 4 hours of dark (20L4D). Our observations

indicate that *Clk* uORFs did not influence the timing of locomotor activity rhythm across these

varied photoperiods, suggesting that *Clk* uORFs are not essential for adapting locomotor

rhythms to changes in seasonal day length (SI Appendix, Fig. S9).

Nevertheless, analysis of sleep duration under varying photoperiods revealed that female

Clk-uORF-KO flies exhibited a greater disparity from WT as the photoperiod decreased (Fig.

7A and 7B). Notably, sleep duration showed a significant positive correlation with photoperiod

length in WT (Pearson's $r = 0.323$, $P = 6.5 \times 10^{-8}$), while this correlation was much weaker in

Clk-uORF-KO (Pearson's r = 0.172, P = 0.006) (Fig. 7C). As the consequence, the difference

in sleep duration between *Clk*-uORF-KO and WT became larger as the photoperiod shortened

(Fig. 7B). Under 4L20D, the shortest photoperiod tested here, WT displayed on average 918 ±

26 minutes of sleep, while that of *Clk*-uORF-KO females was 1,124 ± 18 minutes, with an

increase of 206 minutes on average (22.44%). In contrast, under 20L4D, the longest

photoperiod tested, WT females slept for 1,082 ± 21 minutes compared to 1,195 ± 18 minutes

for *Clk*-uORF-KO females, yielding a smaller difference of 113 minutes on average (10.44%).

These findings indicate that *Clk*-uORF-KO flies showed a diminished response to changes in

photoperiod length, implying that *Clk* uORF-mediated translational regulation plays a role in

synchronizing sleep patterns with seasonal daylight variations.

Given these observed behavioral changes, we posited that *Clk* uORFs participate in

translational regulation in a photoperiod-dependent manner. To test this, we carried out

ribosome fractionation followed by qPCR under 4L20D and 12L12D light-dark cycles. In WT

heads, we observed a consistent decrease in P-to-M ratios throughout the day under 12L12D

vs. 4L20D (Fig. 8A). However, such a pattern was not observed in *Clk*-uORF-KO (Fig. 8B).

The area under the curve (AUC) of these ratios, indicative of overall daily CLK protein

synthesis, was significantly reduced by long photoperiods in WT fly heads (~30% reduction),

while the reduction was much less (~8% reduction) in *Clk*-uORF-KO (Fig. 8C and 8D). This

photoperiod dependency was corroborated at the protein level. In WT heads, CLK protein levels

dipped significantly during ZT4-12 under 4L20D compared to 12L12D (Fig. 8E and 8F). In

contrast, CLK protein level was only significantly decreased at ZT0 in *Clk*-uORF-KO under

4L20D relative to 12L12D (Fig. 8E and 8F). Moreover, the AUC analysis demonstrated that

short photoperiod significantly reduced total daily CLK level in WT but not in *Clk*-uORF-KO

heads (SI Appendix, Fig. S10). Collectively, these findings suggest that *Clk* uORFs are involved

in photoperiod-dependent translation suppression, which likely aids in adjusting wakefulness

to seasonal variations.

Knocking out *Clk* uORFs impairs fecundity

Recognizing the roles of *Clk* uORFs in facilitating the adaptation to environmental changes, we

further investigated whether they enhance evolutionary fitness by assessing fecundity.

Remarkably, mated *Clk*-uORF-KO females laid significantly fewer eggs than their WT

counterparts during an eight-day monitoring duration, with the total egg production of mutants

markedly less ($P < 0.01$) (SI Appendix, Fig. S11A). This pattern persisted in *Clk*-uORF-KO

virgin females, which also laid fewer eggs than WT in both segmented (2-days) and pooled

(10-days) measurements (SI Appendix, Fig. S11B). Accordingly, female *Clk*-uORF-KO flies

produced significantly fewer offspring than WT counterparts at 25°C ($P = 1.5 \times 10^{-5}$ Wilcoxon

rank sum test) (SI Appendix, Fig. S11C). Even at 29°C, when both mutants and WT flies exhibited reduced offspring production compared to 25°C, the offspring counts for *Clk*-uORF-KO females were still significantly fewer than that for WT ($P = 0.00029$, Wilcoxon rank sum test) (SI Appendix, Fig. S11C). Collectively, these results indicate that *Clk* uORFs profoundly affect various aspects of physiological and behavioral processes, with implications for fitness beyond their established roles in circadian rhythm and sleep regulation.

Discussion

The evolutionary conservation of the circadian mechanisms across species underscores their fundamental significance. Our understanding of circadian rhythms has grown substantially over the years, providing critical insights into a range of physiological processes. Earlier investigations, including our own, have noted a very low correlation between the temporal variation pattern of the transcriptome and that of the proteome, indicating that regulations at the translational and post-translational levels play dominant roles in determining the daily variations of protein abundance [55]. Within this complex regulatory network, our study here has highlighted a pivotal role for translational regulation, specifically mediated by uORFs, in fine-tuning these rhythms. Beyond circadian rhythm and sleep, *Clk*-uORF-KO flies show diminished fecundity and reduced starvation resilience, suggesting that *Clk* uORFs might act as a master regulator, shaping the rhythmic expression of a vast array of genes and influencing multifaceted physiological outcomes.

In this study, we observed over one-hour shortening of the circadian period when *Clk* uORFs were knocked out, which means these uORFs play a crucial role in the core circadian timing mechanism. Consistent with this idea, we found that *Clk* uORFs mediate an instructive signal to the molecular clock by suppressing CLK protein translation during the daytime. Besides setting the pace of the clock, *Clk* uORFs also modulate cyclic gene expression globally. The oscillation of over 1/3 of the rhythmic genes requires *Clk* uORFs, as these oscillations were dampened in *Clk*-uORF-KO flies. This reveals the delicacy of the circadian system: an increase of CLK translation and, thus, CLK protein level caused by knocking out *Clk* uORFs can exert detrimental effects on the rhythmicity of the transcriptome. It is surprising that genes cycle with a more robust amplitude in *Clk*-uORF-KO flies relative to WT. Among these genes, about half did not display oscillatory expression patterns in WT. In other words, *Clk* uORFs suppress the mRNA rhythms of these genes. We reason that knocking out *Clk* uORFs increases CLK protein level in a time-dependent manner, which results in enhanced transcription of CLK target genes at a certain time of the day, and thus, these genes may acquire a *de novo* rhythmic expression pattern. Considering that *Clk* uORFs suppress CLK translation more under longer photoperiods and less under shorter photoperiods, we speculate this photoperiod-dependent variation in the

translational activities of *Clk* uORFs can shape the rhythmic transcription landscape in
adaptation to seasonal photoperiod changes.

Clk uORFs promote morning wakefulness, consistent with the time-dependent modulation
of these uORFs on CLK protein translation and abundance. This regulation is not only time of
the day-dependent, but also time of the year-dependent. WT flies display increased wakefulness
and decreased sleep in adaptation to a shortening of photoperiod (occurring in fall and winter
seasons), while *Clk* uORFs participate in modulating this process by suppressing CLK
translation in a photoperiod-dependent manner. Intriguingly, human individuals with seasonal
affective disorder (SAD, commonly known as winter depression) report a winter increase in
sleep, ranging from 30 min to 2 h longer in duration compared to controls, while the underlying
mechanism is yet unknown [56]. The long sleep phenotype caused by knocking out *Clk* uORFs is
most striking under a short photoperiod, reminiscent of the winter hypersomnolence in SAD.
Notably, while we observed a stronger sleep phenotype in female flies, SAD is also more
predominant in women [57]. From a mechanistic perspective, our results strongly suggest that *Clk*
uORFs promote wakefulness by increasing tyrosine hydroxylase expression, thus up-regulating
the dopaminergic tone. Similarly, the mammalian CLOCK protein has been shown to repress
the transcription of *Tyrosine hydroxylase*, while *Clock* mutation leads to decreased sleep and
depression-like behaviors in mice due to elevated dopamine signaling [58,59]. Taken together, our
findings here may provide some insights for understanding the pathological mechanism of SAD.
439

It seems contradictory that CLK translation is down-regulated under 12L12D vs. 4L20D,
while CLK protein level is higher under 12L12D relative to 4L20D. We reason that CLK protein
abundance is determined by coordinated actions of transcription, translation, and post-
translational modifications. A longer photoperiod may promote transcription and/or inhibit
degradation of CLK protein, while *Clk* uORFs counteract these influences by suppressing CLK
synthesis. In other words, *Clk* uORFs may serve as a buffer against the impacts of transcription
and/or PTMs, adjusting CLK protein abundance to maintain an optimal level throughout the
year. Of course, further investigations are needed to elucidate how *Clk* uORFs mediate
photoperiodic signals.
449

uORFs have been shown to dynamically modulate the translation of CDSs in stress
responses [30], and here we also observed a role for *Clk* uORFs in responses to starvation stress.
Although under food *ad libitum* condition, *Clk* uORFs promote wakefulness, under starvation
they function to reduce wakefulness, implicating the complexity of uORF regulation and their
multifaceted roles in varying physiological states. Despite these complicated and sometimes
contradictory actions, in the end, the presence of *Clk* uORFs confers adaptive advantages and

is beneficial for the survival (KO flies die earlier when starved) and fecundity of the organism.

To summarize, here we observed an enrichment of uORFs in circadian rhythm genes and

further identified functional uORFs in *Clk* that repress the translation of CLK protein in both

circadian and circannual manners. These regulations slow down the pace of the clock, modulate

global rhythms of gene expression, and promote morning wakefulness in adaption to seasonal

photoperiod changes. These findings, situated within the broader evolutionary, clinical, and

interdisciplinary contexts, underscore the pivotal role of uORFs in shaping biological processes

and potential therapeutic avenues.

Acknowledgements

This work was supported by grants from the Natural Science Foundation of China (31930021

and 32341021 to L.Z., 32070597 to J. L, and 32200956 to K.S.), the Ministry of Science and

Technology of China (STI 2030-Major Projects 2021ZD0203200-02 to L.Z, and

2022YFE0132000 to J.L.), the Department of Science and Technology of Hubei Province

(2022CFA049 to L.Z.) and Natural Science Foundation of Beijing (grant 5212006 to J.L.). We

thank the National Center for Protein Sciences at Peking University for technical assistance.

Some of the analyses were performed on the High-Performance Computing Platform of the

Center for Life Sciences. We thank Dr. Joanna Chiu for CLK antibody used in this study.

Data and code availability

477•   The raw fastq files for the RNA-seq libraries were deposited to NCBI Sequence Read Archive

478   (SRA) with BioProject accession PRJNA1026336 and are publicly available as of the date of

479   publication.

480•   Original Western blot images have been deposited to Mendeley and are publicly available on

481   the date of publication.

482•   Any additional information required to reanalyze the data reported in this paper is available

483   from the lead contact upon request.

484•

485   **References**

486   1.   Li, S., and Zhang, L. (2015). Circadian Control of Global Transcription. Biomed Res

487   Int *2015*, 187809. 10.1155/2015/187809.

488   2.   Dunlap, J.C. (1999). Molecular bases for circadian clocks. Cell *96*, 271-290.

489   10.1016/s0092-8674(00)80566-8.

490   3.   Patke, A., Young, M.W., and Axelrod, S. (2020). Molecular mechanisms and

491   physiological importance of circadian rhythms. Nature Reviews Molecular Cell

492   Biology *21*, 67-84. 10.1038/s41580-019-0179-2.

4.   Panda, S., Hogenesch, J.B., and Kay, S.A. (2002). Circadian rhythms from flies to

human. Nature *417*, 329-335. 10.1038/417329a.

5.   Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock.
Nature Reviews Genetics *18*, 164-179. 10.1038/nrg.2016.150.

6.   Reinke, H., and Asher, G. (2019). Crosstalk between metabolism and circadian clocks.
Nature Reviews Molecular Cell Biology *20*, 227-241. 10.1038/s41580-018-0096-9.

7.   Masri, S., and Sassone-Corsi, P. (2013). The circadian clock: a framework linking
metabolism, epigenetics and neuronal function. Nat Rev Neurosci *14*, 69-75.
10.1038/nrn3393.

8.   Palm, D., Uzoni, A., Simon, F., Fischer, M., Coogan, A., Tucha, O., Thome, J., and
Faltraco, F. (2021). Evolutionary conservations, changes of circadian rhythms and their
effect on circadian disturbances and therapeutic approaches. Neurosci Biobehav Rev
128, 21-34. 10.1016/j.neubiorev.2021.06.007.

9.   Rosbash, M. (2009). The Implications of Multiple Circadian Clock Origins. PLOS
Biology *7*, e1000062. 10.1371/journal.pbio.1000062.

10.   Young, M.W., and Kay, S.A. (2001). Time zones: a comparative genetics of circadian
clocks. Nat Rev Genet *2*, 702-715. 10.1038/35088576.

11.   Tataroglu, O., and Emery, P. (2014). Studying circadian rhythms in Drosophila
melanogaster. Methods *68*, 140-150. https://doi.org/10.1016/j.ymeth.2014.01.001.

12.   Rosato, E., Tauber, E., and Kyriacou, C.P. (2006). Molecular genetics of the fruit-fly
circadian   clock.   European   Journal   of   Human   Genetics   *14*,   729-738.
10.1038/sj.ejhg.5201547.

13.   Dubowy, C., and Sehgal, A. (2017). Circadian Rhythms and Sleep in Drosophila
melanogaster. Genetics *205*, 1373-1397. 10.1534/genetics.115.185157.

14.   Zehring, W.A., Wheeler, D.A., Reddy, P., Konopka, R.J., Kyriacou, C.P., Rosbash, M.,
and Hall, J.C. (1984). P-element transformation with period locus DNA restores
rhythmicity to mutant, arrhythmic Drosophila melanogaster. Cell *39*, 369-376.
10.1016/0092-8674(84)90015-1.

15.   Allada, R., White, N.E., So, W.V., Hall, J.C., and Rosbash, M. (1998). A mutant
Drosophila homolog of mammalian Clock disrupts circadian rhythms and transcription
of period and timeless. Cell *93*.

16.   Rutila, J.E. (1998). CYCLE is a second bHLH–PAS clock protein essential for
circadian rhythmicity and transcription of Drosophila period and timeless. Cell *93*.

17.   Sehgal, A., Price, J.L., Man, B., and Young, M.W. (1994). Loss of circadian behavioral
rhythms and per RNA oscillations in the Drosophila mutant timeless. Science *263*.

18.   Kadener, S., Menet, J.S., Schoer, R., and Rosbash, M. (2008). Circadian transcription
contributes to core period determination in Drosophila. PLOS Biol. *6*.

19.   Zhao, J. (2003). Drosophila clock can generate ectopic circadian clocks. Cell *113*.

20.   Lee, J., Yoo, E., Lee, H., Park, K., Hur, J.H., and Lim, C. (2017). LSM12 and
ME31B/DDX6 Define Distinct Modes of Posttranscriptional Regulation by ATAXIN-
2 Protein Complex in Drosophila Circadian Pacemaker Neurons. Mol Cell *66*, 129-140
e127. 10.1016/j.molcel.2017.03.004.

21.   Lim, C., and Allada, R. (2013). ATAXIN-2 activates PERIOD translation to sustain
circadian rhythms in Drosophila. Science *340*, 875-879. 10.1126/science.1234785.

22.   Lim, C., Lee, J., Choi, C., Kilman, V.L., Kim, J., Park, S.M., Jang, S.K., Allada, R.,

and Choe, J. (2011). The novel gene twenty-four defines a critical translational step in
the Drosophila clock. Nature *470*, 399-403. 10.1038/nature09728.

23.   Zhang, Y., Ling, J., Yuan, C., Dubruille, R., and Emery, P. (2013). A role for Drosophila
ATX2 in activation of PER translation and circadian behavior. Science *340*, 879-882.
10.1126/science.1234746.

24.   Kadener, S. (2009). A role for microRNAs in the Drosophila circadian clock. Genes
Dev. *23*.

25.   Chen, X., and Rosbash, M. (2016). mir-276a strengthens Drosophila circadian rhythms
by regulating timeless expression. Proc. Natl Acad. Sci. USA *113*.

26.   Chen, W. (2014). Regulation of Drosophila circadian rhythms by miRNA let-7 is
mediated by a regulatory cycle. Nat. Commun. *5*.

27.   Zhang, H., Wang, Y., Wu, X., Tang, X., Wu, C., and Lu, J. (2021). Determinants of
genome-wide distribution and evolution of uORFs in eukaryotes. Nature
Communications *12*, 1076. 10.1038/s41467-021-21394-y.

28.   Hinnebusch, A.G., Ivanov, I.P., and Sonenberg, N. (2016). Translational control by 5'-
untranslated regions of eukaryotic mRNAs. Science *352*, 1413-1416.
10.1126/science.aad9868.

29.   Morris, D.R., and Geballe, A.P. (2000). Upstream open reading frames as regulators of
mRNA translation. Mol Cell Biol *20*, 8635-8642. 10.1128/MCB.20.23.8635-
8642.2000.

30.   Zhang, H., Wang, Y., and Lu, J. (2019). Function and Evolution of Upstream ORFs in
Eukaryotes. Trends Biochem Sci *44*, 782-794. 10.1016/j.tibs.2019.03.002.

31.   Barbosa, C., Peixeiro, I., and Romão, L. (2013). Gene expression regulation by
upstream open reading frames and human disease. PLoS Genet *9*, e1003529.
10.1371/journal.pgen.1003529.

32.   Ruiz-Orera, J., and Alba, M.M. (2019). Translation of Small Open Reading Frames:
Roles in Regulation and Evolutionary Innovation. Trends Genet *35*, 186-198.
10.1016/j.tig.2018.12.003.

33.   Diernfellner, A.C., Schafmeier, T., Merrow, M.W., and Brunner, M. (2005). Molecular
mechanism of temperature sensing by the circadian clock of Neurospora crassa. Genes
Dev *19*, 1968-1973. 10.1101/gad.345905.

34.   Wu, H.W., Fajiculay, E., Wu, J.F., Yan, C.S., Hsu, C.P., and Wu, S.H. (2022). Noise
reduction by upstream open reading frames. Nat Plants *8*, 474-480. 10.1038/s41477-
022-01136-8.

35.   Montenegro-Montero, A., Canessa, P., and Larrondo, L.F. (2015). Around the Fungal
Clock: Recent Advances in the Molecular Study of Circadian Clocks in Neurospora
and Other Fungi. Adv Genet *92*, 107-184. 10.1016/bs.adgen.2015.09.003.

36.   Salichos, L., and Rokas, A. (2010). The diversity and evolution of circadian clock
proteins in fungi. Mycologia *102*, 269-278. 10.3852/09-073.

37.   Jabbur, M.L., and Johnson, C.H. (2022). Spectres of Clock Evolution: Past, Present,
and Yet to Come. Frontiers in Physiology *12*. 10.3389/fphys.2021.815847.

38.   Miyake, T., Inoue, Y., Shao, X., Seta, T., Aoki, Y., Nguyen Pham, K.T., Shichino, Y.,
Sasaki, J., Sasaki, T., Ikawa, M., et al. (2023). Minimal upstream open reading frame
of Per2 mediates phase fitness of the circadian clock to day/night physiological body

582   temperature rhythm. Cell Rep *42*, 112157. 10.1016/j.celrep.2023.112157.

583   39.   Millius, A., Yamada, R.G., Fujishima, H., Maeda, K., Standley, D.M., Sumiyama, K.,
584   Perrin, D., and Ueda, H.R. (2023). Circadian ribosome profiling reveals a role for the
585   Period2 upstream open reading frame in sleep. Proc Natl Acad Sci U S A *120*,
586   e2214636120. 10.1073/pnas.2214636120.

587   40.   Jang, C., Lahens, N.F., Hogenesch, J.B., and Sehgal, A. (2015). Ribosome profiling
588   reveals an important role for translational control in circadian gene expression. Genome
589   Res *25*, 1836-1847. 10.1101/gr.191296.115.

590   41.   Janich, P., Arpat, A.B., Castelo-Szekely, V., Lopes, M., and Gatfield, D. (2015).
591   Ribosome profiling reveals the rhythmic liver translatome and circadian clock
592   regulation by upstream open reading frames. Genome Res *25*, 1848-1859.
593   10.1101/gr.195404.115.

594   42.   Zhang, H., Dou, S., He, F., Luo, J., Wei, L., and Lu, J. (2018). Genome-wide maps of
595   ribosomal occupancy provide insights into adaptive evolution and regulatory roles of
596   uORFs during Drosophila development. PLoS Biol *16*, e2003903.
597   10.1371/journal.pbio.2003903.

598   43.   Zhang, H., Wang, Y., Wu, X., Tang, X., Wu, C., and Lu, J. (2021). Determinants of
599   genome-wide distribution and evolution of uORFs in eukaryotes. Nat Commun *12*,
600   1076. 10.1038/s41467-021-21394-y.

601   44.   Abdelmohsen, K., Panda, A.C., Kang, M.J., Guo, R., Kim, J., Grammatikakis, I., Yoon,
602   J.H., Dudekula, D.B., Noh, J.H., Yang, X., et al. (2014). 7SL RNA represses p53
603   translation by competing with HuR. Nucleic Acids Res *42*, 10099-10111.
604   10.1093/nar/gku686.

605   45.   Panda, A.C., Abdelmohsen, K., Martindale, J.L., Di Germanio, C., Yang, X.,
606   Grammatikakis, I., Noh, J.H., Zhang, Y., Lehrmann, E., Dudekula, D.B., et al. (2016).
607   Novel RNA-binding activity of MYF5 enhances Ccnd1/Cyclin D1 mRNA translation
608   during myogenesis. Nucleic Acids Res *44*, 2393-2408. 10.1093/nar/gkw023.

609   46.   Panda, A.C., Martindale, J.L., and Gorospe, M. (2017). Polysome Fractionation to
610   Analyze mRNA Distribution Profiles. Bio Protoc *7*. 10.21769/BioProtoc.2126.

611   47.   Fathallah-Shaykh, H.M., Bona, J.L., and Kadener, S. (2009). Mathematical model of
612   the Drosophila circadian clock: loop regulation and transcriptional integration. Biophys
613   J *97*, 2399-2408. 10.1016/j.bpj.2009.08.018.

614   48.   Abruzzi, K.C., Rodriguez, J., Menet, J.S., Desrochers, J., Zadina, A., Luo, W., Tkachev,
615   S., and Rosbash, M. (2011). Drosophila CLOCK target gene characterization:
616   implications for circadian tissue-specific gene expression. Genes Dev *25*, 2374-2386.
617   10.1101/gad.178079.111.

618   49.   Hughes, M.E., Hogenesch, J.B., and Kornacker, K. (2010). JTK_CYCLE: an efficient
619   nonparametric algorithm for detecting rhythmic components in genome-scale data sets.
620   J Biol Rhythms *25*, 372-380. 10.1177/0748730410379711.

621   50.   Liu, S., Lamaze, A., Liu, Q., Tabuchi, M., Yang, Y., Fowler, M., Bharadwaj, R., Zhang,
622   J., Bedont, J., Blackshaw, S., et al. (2014). WIDE AWAKE mediates the circadian
623   timing of sleep onset. Neuron *82*, 151-166. 10.1016/j.neuron.2014.01.040.

624   51.   Kumar, S., Chen, D., and Sehgal, A. (2012). Dopamine acts through Cryptochrome to
625   promote acute arousal in Drosophila. Genes Dev *26*, 1224-1234.

626   10.1101/gad.186338.111.

52. Pendleton, R.G., Rasheed, A., Paluru, P., Joyner, J., Jerome, N., Meyers, R.D., and Hillman, R. (2005). A developmental role for catecholamines in Drosophila behavior. Pharmacol Biochem Behav *81*, 849-853. 10.1016/j.pbb.2005.06.008.

53. Keene, A.C., Duboue, E.R., McDonald, D.M., Dus, M., Suh, G.S., Waddell, S., and Blau, J. (2010). Clock and cycle limit starvation-induced sleep loss in Drosophila. Curr Biol *20*, 1209-1215. 10.1016/j.cub.2010.05.029.

54. Wood, S., and Loudon, A. (2014). Clocks for all seasons: unwinding the roles and mechanisms of circadian and interval timers in the hypothalamus and pituitary. J Endocrinol *222*, R39-59. 10.1530/JOE-14-0141.

55. Wang, C., Shui, K., Ma, S., Lin, S., Zhang, Y., Wen, B., Deng, W., Xu, H., Hu, H., Guo, A., et al. (2020). Integrated omics in Drosophila uncover a circadian kinome. Nat Commun *11*, 2710. 10.1038/s41467-020-16514-z.

56. Wescott, D.L., Soehner, A.M., and Roecklein, K.A. (2020). Sleep in seasonal affective disorder. Curr Opin Psychol *34*, 7-11. 10.1016/j.copsyc.2019.08.023.

57. Jepson, T.L., Ernst, M.E., and Kelly, M.W. (1999). Current Perspectives on the Management of Seasonal Affective Disorder. Journal of the American Pharmaceutical Association (1996) *39*, 822-829. 10.1016/s1086-5802(15)30373-9.

58. Roybal, K., Theobold, D., Graham, A., DiNieri, J.A., Russo, S.J., Krishnan, V., Chakravarty, S., Peevey, J., Oehrlein, N., Birnbaum, S., et al. (2007). Mania-like behavior induced by disruption of CLOCK. Proc Natl Acad Sci U S A *104*, 6406-6411. 10.1073/pnas.0609625104.

59. Sidor, M.M., Spencer, S.M., Dzirasa, K., Parekh, P.K., Tye, K.M., Warden, M.R., Arey, R.N., Enwright, J.F., Jacobsen, J.P., Kumar, S., et al. (2015). Daytime spikes in dopaminergic activity drive rapid mood-cycling in mice. Mol Psychiatry *20*, 1479-1480. 10.1038/mp.2015.8.

60. Sisobhan, S., Rosensweig, C., Lear, B.C., and Allada, R. (2022). SleepMat: a new behavioral analysis software program for sleep and circadian rhythms. Sleep *45*. 10.1093/sleep/zsac195.

61. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

62. Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics *31*, 166-169. 10.1093/bioinformatics/btu638.

63. Gene Ontology, C. (2021). The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Res *49*, D325-D334. 10.1093/nar/gkaa1113.

Materials and Methods

Fly strains and cell line

All flies were reared on standard cornmeal-yeast-sucrose medium and kept in light/dark (LD)

cycles at 25°C except for specific experimental design denoted in the text. Food was accessible

ad libitum. *{nos-Cas9}attP40* and *{nos-Cas9}attP2* fly stocks used for embryo injection were

from the TsingHua Fly Center. The tool fly stocks (*y sc v* and *Dr, e/TM3, Sb*) used for mutant

screening were from the TsingHua Fly Center. *Clk*-uORF-KO and *cyc*-uORF-KO mutants were

generated and backcrossed to the Canton-S strain for nine generations. Homozygous mutants

were used throughout this study and the Canton-S strain served as the WT control flies.

Drosophila S2 cells used in this study were purchased from Life Technologies Corp

(http://www.lifetechnologies.com).

The conservation assessment of uORFs

We retrieved the whole genome sequence alignment (maf) of *D. melanogaster* (dm6) with 22

other *Drosophila* species from the UCSC genome browser (genome.ucsc.edu). Leveraging the

genome annotation of *D. melanogaster* (FlyBase r6.04, https://flybase.org/), the 5'UTR

sequence of each annotated transcript of *D. melanogaster* and its corresponding sequences in

other 22 *Drosophila* were extracted from the maf. We scanned all the AUG triplets within the

5'UTRs of *D. melanogaster* and then determined their presence or absence in the corresponding

positions in the other species based on maf. AUG triplets that overlapped with any annotated

CDS region were removed. The transcriptome and translatome data was obtained from the

previous study [42].

Generation of uORF-KO strain

We searched for possible sgRNA target sites near the start codon of the uORFs of *Clk* and *cyc*

genes using the Benchling website (https://www.benchling.com/crispr/) to design optimal

single guide RNA (sgRNA) sequences with high specificity and low off-target effects. Two

pairs of sgRNAs were used for each gene. We then synthesized single-stranded complementary

DNAs (ssDNAs) and annealed them to obtain double-stranded DNA (dsDNA), which served

as the template for sgRNA expression. The dsDNA was then ligated into the BbsI-digested

pU6B vector. The template sequences of the sgRNA uATG-KO are listed in SI Appendix, Table

S5. The screening procedures of mutant flies are provided in the (SI Appendix, SI Methods).

uORF mutation sequencing validation

The mutant fly was homogenized with DNA sequencing extract buffer [10 mM Tris-HCl at pH

8.0, 1 mM EDTA at pH 8.0, 25 mM NaCl and 0.02 % proteinase K (CW2584, CWBIO)] and

incubated at 37°C for 30 min. PCR was performed with Taq Plus MasterMix (CW2849, CWbiotech). The PCR reaction was performed as follows: 94 °C for 2 min followed by 94 °C for 10 s, 57°C for 15 s, and 72°C for 1 min 20 s for 35 cycles. The primers used were listed in (SI Appendix, Table S5). PCR products were sequenced using commercial Sanger sequencing (AuGCT Co.).

Cell culture and transfection

S2 cells were cultured in Schneider's *Drosophila* Medium (Sigma) containing 10% heat-inactivated fetal bovine serum (FBS), 100 U/ml penicillin, and 100 μg/ml streptomycin (Thermo Fisher) at 25°C for 24 h to reach $2\text{-}4 \times 10^6$ cells/ml before further treatments. Plasmid transfection was conducted with Lipofectamine 3000 (L3000001, Thermo Fisher) according to the supplier's protocol.

Dual-luciferase assay

The wild-type (WT) 5'UTR of *Clk* and *cyc* was cloned from cDNA by PCR, and uATG mutations were introduced into 5'UTR by the amplification primers. The PCR primers used for this purpose are listed in (SI Appendix, Table S5). The WT and mutated 5'UTR sequences were ligated into a linearized reporter plasmid (psiCHECK-2 vector, Promega). The entire sequences of all the plasmids were validated by Sanger sequencing. The *Renilla* luciferase activity associated with WT or uORF-mutated 5'UTRs was measured according to the manual of the Dual-Luciferase Reporter Assay System (Promega) 32 h after transfection and was normalized to the activity of firefly luciferase.

Fly head collection

Flies were collected within 3-7 days of eclosion and entrained in LD for three days. After that, they were transferred into constant darkness (DD). Flies were collected during LD or on the first day of DD and frozen at -80°C. Frozen flies were vortexed for 10 s to separate the head from the body.

Ribosome fraction analysis

The heads obtained from WT and mutant flies were collected and homogenized with lysis buffer, the lysates were clarified by centrifugation at 4°C and 20,000×g for 10 min, and the supernatants were applied to sucrose gradient ultracentrifugation. For each sample, the ratio of *Clk* and *cyc* mRNA abundance in the polysome fraction to that in the monosome fraction was calculated as the P-to-M ratio. Six biological replicates were performed for each sample. Details of the procedures can be found in the (SI Appendix, SI Methods).

Western blot

Proteins were extracted from fly heads and cultured cells using SDS lysis buffer [50 mM Tris-HCl at pH 7.6, 150 mM NaCl, 1% Triton X-100, 0.5% SDS, 1 mM EDTA at pH8.0, 2 mM DTT, 1 × proteinase inhibitor and phosphatase inhibitor (4693116001 and 04906845001, Roche)]. After homogenization, protein lysates were centrifuged at 12,000 × *g* for 15 min at 4°C and incubated at 95°C in the loading buffer for 5 min. Equal amounts of protein were loaded into each well on 6% SDS-PAGE gels and then transferred to nitrocellulose membranes for 2 h at 90 V. Membranes were incubated with primary antibody at 4°C overnight, followed by secondary antibody at room temperature for 1 h. The primary antibodies used were as follows: guinea pig anti-CLK (1:1000, gift from Dr. Joanna Chiu) and rabbit anti-ACTIN 57B (1:5000, AC026, ABclonal, CN). Donkey secondary antibodies (1:10000 dilution) were conjugated either with IRDye 680 or IRDye 800 (926-68072 and 926-32213, LI-COR Biosciences, US). Bands were visualized and quantified with an Odyssey Infrared Imaging System (LI-COR Biosciences).

Model simulation for the outcome of enhanced *Clk* translation

We adopted a previously published mathematical model of the *Drosophila* circadian clock [47]. This system relies on a nonlinear, autonomous, first-order system of ordinary differential equations to simulate the regulatory weights among mRNAs or protein molecules in the molecular clock network. Details of the simulation can be found in the (SI Appendix, SI Methods).

Total RNA extraction and RT-qPCR

RNA was extracted from frozen fly heads maintained at -80°C. 50 fly heads per sample were homogenized in Total RNA Isolation (TRIzol) Reagent (15596026, Invitrogen) by using a homogenizer with plastic beads. After mixing with trichloromethane, homogenates were centrifuged at 12,000 × *g* and the suspension was precipitated with 75% ethanol. After air dry, total RNA was reverse-transcribed into cDNA and genomic DNA was removed with One-step cDNA Synthesis SuperMix (AT311, Transgen). Quantitative real-time PCR was performed with One-Step RT-PCR SuperMix (AT141, Transgen). The PCR reaction was performed as follows: 45°C for 5 min; 94°C for 2 min; 94°C for 5 s, 60°C for 15 s, 72°C for 20 s for 40 cycles (Applied Biosystems). The ΔΔCT method was used for quantification. *Beta-Actin* was used as an internal control. The primers used are listed in (SI Appendix, Table S5).

Locomotor activity monitoring and analysis

Locomotor activity levels of adult flies were monitored by *Drosophila* Activity Monitoring System (TriKinetics) for 7 days of LD followed by 7 days of DD. For DD rhythmicity, chi-

squared periodogram analyses were performed by Clocklab (Actimetrics). Rhythmic flies were defined as those in which the chi-squared power was ≥ 10 above the significance line. Period calculations considered only rhythmic flies. Dead flies were defined by 0 activity on DD7 and removed from the analysis.

Sleep measurement and analysis

Drosophila Activity Monitor system (Trikinetics) was used to analyze fly sleep. Virgin female and male flies of 3~5 days old were used for the experiments unless otherwise specified. Flies were entrained under LD at 25°C for 3 days, and then their activities in the next 3 days under LD conditions were analyzed.

For sleep analysis during starvation, 3~5 days old virgin females were used. After 3 days of entrainment in monitoring tubes with standard media (2% agar and 5% sucrose), baseline sleep was measured for 2 days. Standard food was then replaced at ZT0 by starvation media (2% agar) for 24 h. Flies were then re-fed with standard media for an additional 24-hour recovery period.

Sleep is defined as 5 min consecutive inactivity. Sleep was analyzed with SleepMat following previously published protocol [60].

RNA-seq and circadian analysis

We collected heads from *Clk*-uORF-KO mutants and WT flies raised under LD cycles at 25°C at ZT 0, 4, 8, 12, 16 and 20. Library construction and sequencing with PE150 were conducted by Annoroad on the Illumina Nova6000 platform. Three biological replicates were sequenced for each sample. The clean data were mapped to the reference genome of *D. melanogaster* (FlyBase, r6.04) using STAR [61]. Reads mapped to the exons of each gene were tabulated with htseq-count [62]. Rhythmically expressed genes were identified by JTK_CYCLE under the threshold of < 0.05 on the permutation-based *P* values of the Jonckheere-Terpstra test (ADJ.P) implemented in the JTK CYCLE.

GO-based enrichment

For GO-based enrichment analysis, GO annotation files were downloaded on 15 November 2023 from the Gene Ontology Resource (http://geneontology.org/)[63] and contained 14,736 fly genes with at least one GO term. For each GO term *t*, we defined the following:

N = the total number of genes in the background distribution.

n = the total number of genes for GO analysis.

M = the total number of genes annotated by GO term *t*.

m = the number of genes annotated by GO term *t* in *n*.

The enrichment ratio (E-ratio) of *t* was then computed, and the *P* was calculated with the

hypergeometric distribution as follows:

$$E - ratio = \frac{m}{M}\bigg/\frac{n}{N} \tag{1}$$

$$P = \sum_{m'=m}^{n} \frac{\binom{M}{m'}\binom{N-M}{n-m'}}{\binom{N}{n}} (E - ratio \geq 1), or \tag{2}$$

$$P = \sum_{m'=0}^{m} \frac{\binom{M}{m'}\binom{N-M}{n-m'}}{\binom{N}{n}} (E - ratio < 1) \tag{3}$$

Drug treatment

For pharmacological experiments, 5 mg/ml L-DOPA (BBI Life Sciences) was mixed in the fly food, and the flies were fed with L-Dopa food during sleep monitoring.

Egg numbers quantification

Newly hatched virgins were picked out and allowed to mature for two days in separate vials. They were then mated by placing one virgin female with three male flies for two days. Subsequently, ten female parents from each group were transferred to a fresh dish containing a grape juice-based medium overlaid with yeast. This procedure was replicated in five separate dishes for each strain.

After one day of egg laying, the number of eggs deposited was meticulously documented. Following this assessment, the female parents were relocated to a new dish. This egg quantification process was conducted over a total duration of eight days. For the virgin female, egg counts were performed every two days, extending over a span of ten days.

Quantification of offspring number per female fly

Newly hatched virgins were picked out and allowed to mature for two days in individual vials. They were then mated by placing one virgin female with three male flies for two days. After that, each female parent was transferred to a new vial and the number of offspring produced in 10 days was measured at 25°C and 29°C. All assays were performed with 20 females per genotype.

Measurement of starvation resistance in adult flies

We selected 3- to 5-day-old adult males and females and placed them in a starvation medium (1.5% agar), with 10 flies per vial and 10 vials for both males and females from each strain. We observed every 6 hours or 12 hours to count the number of deaths under starvation conditions until all flies starved to death. The survival curves were plotted by the ggsurvplot package in R.

Quantification and statistical analysis

Quantifications in all data graphs represent the mean of at least four biological replicates, and

error bars represent the standard error of the mean (SEM). Two-tailed Mann-Whitney U test,

Wilcoxon sign rank test, log-rank test, and two-tailed Student's t test were used to calculate

significance. All statistical analysis was carried out using *Statskingdom*. In each statistical

analysis, the number of biological replicates and significant P values were noted in

corresponding Fig. legends.

Materials availability

All fly lines generated in this study are available from the Lead Contact.

Table 1. uORFs in circadian rhythm genes.

Description	Circadian rhythm genes	Core rhythm genes	*Clk*	*sgg*	*cwo*	*Pdp1*	*tim*	*cyc*	*vri*	*cry*	*Bdbt*	*per*
Total uORFs	1137	82	7	18	9	24	8	4	12	0	0	0
Conserved uORFs	29	7	1	1	1	4	0	0	0	0	0	0
Male head												
uORFs with mRNA RPKM > 0.1	888	67	6	15	8	24	6	3	5	0	0	0
uORFs with TE > 0.1	692	52	5	13	6	15	5	3	5	0	0	0
Conserved uORFs with mRNA RPKM > 0.1	28	7	1	1	1	4	0	0	0	0	0	0
Conserved uORFs with TE > 0.1	25	7	1	1	1	4	0	0	0	0	0	0
Female head												
uORFs with mRNA RPKM > 0.1	879	63	6	15	8	24	5	3	2	0	0	0
uORFs with TE > 0.1	714	49	6	13	5	16	5	3	1	0	0	0
Conserved uORFs with mRNA RPKM > 0.1	28	6	1	1	0	4	0	0	0	0	0	0
Conserved uORFs with TE > 0.1	25	6	1	1	0	4	0	0	0	0	0	0

860 The number of total uORFs or conserved uORFs in 169 circadian rhythm genes and ten core clock genes, as well as
861 each core clock gene, is shown. The number of transcribed and translated uORFs in heads of male and female flies
862 is shown. Conserved uORFs refer to the uATGs of uORFs that are present in all 23 *Drosophila* species. TE,
863 translational efficiency; RPKM, reads per kilobase of transcript per million mapped reads.

Fig. 1. Circadian rhythm genes are more likely to be regulated by uORFs.

(A) Average number of uORFs per gene for circadian rhythm-related genes and the other genes in *D. melanogaster*. The ratio of uORFs to genes is denoted on each bar (Fisher's exact test. ***, $P < 0.001$).

(B) Proportion of conserved uORFs in core clock genes, other circadian rhythm genes, and the other genes. Ratios reflect the number of conserved uORFs relative to the total count of uORFs for each category, depicted on each bar (Fisher's exact test. **, $P < 0.01$; ***, $P < 0.001$).

(C) The translational efficiency (TE) for different classes of circadian rhythm-related genes in male and female heads of *D. melanogaster*. Class I includes genes with conserved and translated uORFs. Class II consists of genes with non-conserved yet translated uORFs. Class III comprises genes without translated uORFs or lacking uORFs altogether. 'Other' encompasses all remaining genes expressed in heads that are not related to circadian rhythm. The gene counts of each class are denoted at the bottom of each bar. The ten core clock genes are highlighted with red dots, with *Clk*, *cyc*, *tim*, and *per* specifically marked by red arrows (Wilcoxon signed-rank test. ***, $P < 0.001$).

Fig. 2. Regulation of CLK and CYC Translation by uORFs in *Drosophila*.

(A) Multiple sequence alignment (MSA) of *Clk* uORFs among 23 *Drosophila* species. The start

codons (uATGs) of uORFs are highlighted by red boxes. The position schemes of uORFs and

partial CDSs are denoted above the MSA, with red and blue colors, respectively. The start and

end positions of each uORF (separated by a "-") in the 5' UTR are given in the parenthesis

above each uATG.

(B) The mRNA reads coverage (left) and ribosome-protected footprints (RPF) coverage (right)

of uORFs of *Clk* mRNA from bodies and heads of both sexes. The position schemes of uORFs

and partial CDSs were denoted at the bottom.

(C) Dual-luciferase assays conducted on WT and uORF-mutated 5' UTRs of *Clk* and *cyc* genes

in *Drosophila* S2 cells, with reporter constructs depicted top left. For the uORF mutants, the

start codons of *Clk* uORF2 and uORF3, as well as *cyc* uORF2, uORF3, and uORF4, were

changed from ATGs to ATTs. Renilla luciferase activity was normalized against firefly

luciferase and the relative activity in WT was normalized to 1. Six replicates were conducted

for each experiment. Data represent mean ± SEM (Two-tailed Student's *t*-test; ***, $P < 0.001$).

(D) Diagrams depicting WT and two uORF knockout strains (*Clk*-uORF-KO and *cyc*-uORF-

KO), created using CRISPR-Cas9. For the uORF mutants, the start codons of *Clk* uORF2 and

uORF3, as well as *cyc* uORF2, uORF3, and uORF4, were simultaneously changed from ATGs

to ATTs. Below each sequence, uORFs (red boxes) and partial CDSs (blue boxes) are shown.

Fig. 3. *Clk* uORFs suppress CLK protein translation in fly heads *in vivo*.

(A) P-to-M ratios of *Clk* mRNA from whole heads of male *Clk*-uORF-KO compared to that in male WT flies, sampled at indicated Zeitgeber times at 4-hour intervals. Each time point includes six biological replicates. Data are expressed as mean ± SEM (Wilcoxon signed-rank test. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; n.s., $P > 0.05$).

(B) P-to-M ratios of *cyc* mRNA from whole heads of male *cyc*-uORF-KO compared to that in male WT flies, sampled at indicated Zeitgeber times at 4-hour intervals. Each time point includes six biological replicates. Data are expressed as mean ± SEM (Wilcoxon signed-rank test. **, $P < 0.01$; n.s., $P > 0.05$).

(C) Representative Western blots of CLK protein from whole-head protein extracts of male

flies, sampled at indicated Zeitgeber times at 4-hour intervals. ACTIN 57B served as the

loading control.

(D) Representative Western blots of CLK protein from whole-head protein extracts of female

flies, sampled at indicated Zeitgeber times at 4-hour intervals. ACTIN 57B served as the

loading control.

(E) Quantification of CLK abundance relative to ACTIN 57B from (C), with the average

intensity for WT normalized to 1 at each Zeitgeber Time (ZT) point. The analysis included four

replicates. Data are presented as mean $\pm$ SEM (Two-tailed Mann-Whitney U test. *, $P < 0.05$;

n.s., $P > 0.05$).

(F) Quantification of CLK abundance relative to ACTIN 57B from (D), with the average

intensity for WT normalized to 1 at each Zeitgeber Time (ZT) point. The analysis included four

replicates. Data are presented as mean $\pm$ SEM (Two-tailed Mann-Whitney U test. *, $P < 0.05$;

**, $P < 0.01$; n.s., $P > 0.05$).

Fig. 4. Computational and experimental analysis of *Clk* uORFs on circadian rhythms.

(A) Diagram of the *Drosophila* circadian clock molecular network, adapted from a previous

study [47]. mRNA molecules are indicated in italicized lowercase, while protein molecules are in

uppercase. Red arrows represent activation, while blue lines indicate suppression. The bold red

arrow from *Clk* mRNA to CLK protein signifies the enhanced translation activity due to *Clk*

uORF knockout. The green arrow with an "X" indicates that CRY protein promotes the
degradation of TIM.

(B) The distribution of period length (hr) under different *Clk* mRNA to CLK protein regulatory
weights across approximately 400 simulation cycles. The original regulatory weight setting of
0.026 in the original model [47] acts as the baseline, emulating the presence of WT uORF. After
200 cycles the regulatory weight is increased from 0.0026 to several arbitrary higher levels
(0.027, 0.028, and 0.029) which is denoted by the red arrow, and continued for another 200
simulation cycles.

(C) The statistics of circadian period length (hr) for simulations in (B). Data are measured in
200 simulation cycles and are reported as the mean ± SEM (two-tailed Student's *t* test. ***, *P*
< 0.001).

(D) qRT-PCR measured mRNA levels of *per* and t*im* from total RNA extracts of female heads,
sampled at specified Zeitgeber time points. WT intensity at ZT0 is normalized to 1. Each time
point includes four biological replicates. Data are reported as the mean ± SEM (two-tailed
Student's *t* test. **, *P* < 0.01; ***, *P* < 0.001).

(E) Period of locomotor rhythms of *Clk*-uORF-KO flies and WT flies under constant darkness
conditions (DD). The period lengths (hr) and the number of flies tested (*n*) are displayed above
and below each bar, respectively.

(F) Representative double-plotted actograms of the locomotor activity of the indicated flies,
monitored under LD followed by DD. White and black bars represent light and dark phases,
respectively.

Figure 5. *Clk* uORFs modulate the temporal pattern of transcriptome expression landscape in fly heads.

(A) The number of rhythmically expressed genes identified from whole-head RNA-seq data of female WT and *Clk*-uORF-KO flies.

(B) Hierarchical clustering of genes expressed rhythmically in whole heads of female flies

across various Zeitgeber time points, with gene expression levels normalized to Z-scores. The

color scale reflects expression intensity, ranging from low (green) to high (red) Z-scores.

(C) The number of rhythmically expressed genes identified from whole-head RNA-seq data of

male WT and *Clk*-uORF-KO flies.

(D) The distribution of expression amplitudes for genes rhythmically expressed in the heads of

female WT and *Clk*-uORF-KO flies ($n = 2,235$; Wilcoxon rank sum test; $P = 5.72 \times 10^{-9}$).

(E) The distribution of expression amplitudes for genes rhythmically expressed in the heads of

male WT and *Clk*-uORF-KO flies ($n = 2,124$; Wilcoxon rank sum test; $P = 1.29 \times 10^{-5}$).

(F-G) Phase distribution of rhythmically expressed genes in whole heads of female WT and

Clk-uORF-KO flies (F) and male WT and *Clk*-uORF-KO flies (G). The *y*-axis of the radar chart

indicates the number of rhythmic genes.

Fig. 6. *Clk* **uORFs regulate sleep/wakefulness.**

(A) Sleep profile of female *Clk*-uORF-KO and WT flies under LD conditions.

(B) Sleep duration during ZT0-8, ZT8-24, and ZT0-24 of female *Clk*-uORF-KO and WT flies under LD condition.

(C) Daily sleep bout number, average bout length, and average waking activity of female *Clk*-

uORF-KO and WT flies under LD condition.

(D) The relative mRNA abundance of *ple* determined by RT-qPCR using whole-head total RNA

extracts collected from female flies under LD conditions. The average value of WT is set to 1.

(E) Sleep profile of female *Clk*-uORF-KO and WT flies treated with L-Dopa under LD

condition.

(F) Daily sleep duration of female *Clk*-uORF-KO *vs.* WT flies treated with L-Dopa.

(G) Sleep profile of female uORF-KO and WT flies under food ad libitum and starvation (the 3rd

day) condition. The yellow shade indicates period of starvation.

(H) Daily sleep duration of female *Clk*-uORF-KO and WT flies under baseline and starvation

conditions.

(I) Histogram shows starvation-induced sleep change (Δ sleep) compared with normal

conditions during daytime (L), nighttime (D), and the entire day (LD).

(J) Survival curves of female flies under starvation conditions ($n = 200$; log-rank test; ***, $P <$

0.001).

The numbers of flies tested are shown in the brackets of legend or the bottom of histograms.

Data are presented as the mean ± SEM. Asterisks indicate statistical significance (two-tailed

Student's *t* test. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; n.s., $P > 0.05$). ZT, Zeitgeber Time.

Fig. 7. *Clk* uORFs promote wakefulness in adaptation to seasonal photoperiod changes.

(A) Sleep profile of female *Clk*-uORF-KO and WT flies under 4 h L : 20 h D (4L20D) (A), 8 h

L : 16 h D (8L16D) (B), 12 h L : 12 h D (12L12D) (C), and 20 h L : 4 h D (20L4D) (D) condition.

The numbers of tested flies (*n*) are shown in the bottom-right.

(B) Daily sleep duration (min) of female *Clk*-uORF-KO and WT flies under different

photoperiods. Data are presented as the mean ± SEM. Asterisks and dollar signs indicate

statistical significance (two-tailed Student's *t* test. ***, *P* < 0.001).

(C) The correlation between daily sleep duration (min) and photoperiod length in female *Clk*-

uORF-KO and WT flies (Pearson's correlation).

1007

Fig. 8. ***Clk*** **uORFs suppress CLK protein translation in a photoperiod-dependent manner.**

(A-B) P-to-M ratios of *Clk* mRNA from whole-head extracts collected from female WT (A)

and *Clk*-uORF-KO (B) flies at indicated time points under 4L20D and 12L12D conditions.

Each time point includes six biological replicates and data are presented as the mean ± SEM.

(C-D) AUC analysis of daily P-to-M ratio in (A) and (B) of female WT (C) and *Clk*-uORF-KO

(D) flies under indicated photoperiod.

(E) Representative Western blots for CLK protein from whole-head total protein extracts

collected from female WT (upper) and *Clk*-uORF-KO (lower) flies at indicated time points

under 4L20D (left) and 12L12D (right). ACTIN 57B is used as a loading control.

(F) The relative CLK abundance normalized to ACTIN 57B in (E). The average value of WT

under 12L12D is set to 1. Each time point includes four biological replicates and data are

presented as the mean ± SEM.

Asterisks indicate statistical significance (two-tailed Mann-Whitney *U* test for unpaired

comparisons. **, $P < 0.01$; ***, $P < 0.001$). ZT, Zeitgeber Time.

1022