1

Zinc finger homeobox-3 (ZFHX3) orchestrates genome-wide daily gene expression in the suprachiasmatic nucleus

Akanksha Bafna[1, 2*], Gareth Banks[1, 3], Vadim Vasilyev[4], Robert Dallmann[4, 5], Michael H. Hastings[6], Patrick M. Nolan[1*]

[1]Medical Research Council, Harwell Science Campus, Oxfordshire, OX11 0RD, United Kingdom

[2]Nuffield Department of Clinical Neurosciences, University of Oxford, OX1 3QU, United Kingdom

[3]Nottingham Trent University, Nottingham, NG1 4FQ, United Kingdom

[4]Division of Biomedical Sciences, Warwick Medical School, University of Warwick, Coventry, United Kingdom

[5]Zeeman Institute for Systems Biology and Infectious Disease Epidemiology Research, University of Warwick, Coventry CV4 7AL, United Kingdom

[6]MRC Laboratory of Molecular Biology, Cambridge, CB2 0QH, United Kingdom

*Corresponding author: akanksha.bafna@ndcn.ox.ac.uk , pmnolan10@gmail.com

ABSTRACT

The mammalian suprachiasmatic nucleus (SCN), situated in the ventral hypothalamus, directs daily cellular and physiological rhythms across the body. The SCN clockwork is a self-sustaining transcriptional-translational feedback loop (TTFL) that in turn co-ordinates the expression of clock-controlled genes (CCGs) directing circadian programmes of SCN cellular activity. In the mouse, the transcription factor, ZFHX3 (zinc finger homeobox-3), is necessary for the development of the SCN and influences circadian behaviour in the adult. The molecular mechanisms by which ZFHX3 affects the SCN at transcriptomic and genomic levels are, however, poorly defined. Here, we used chromatin immunoprecipitation sequencing (ChIP-seq) to map the genomic localization of ZFHX3 binding sites in SCN chromatin. To test for function, we then conducted comprehensive RNA sequencing at six distinct times-of-day to compare the SCN transcriptional profiles of control and ZFHX3-conditional null mutants. We show that the genome-wide occupancy of ZFHX3 occurs predominantly around gene transcription start sites (TSS), co-localizing with known histone modifications, and preferentially partnering with clock transcription factors (CLOCK, BMAL1) to regulate clock gene(s) transcription. Correspondingly, we show that the conditional loss of ZFHX3 in the adult has a dramatic effect on the SCN transcriptome, including changes in the levels of transcripts encoding elements of numerous neuropeptide neurotransmitter systems while attenuating the daily oscillation of the clock TF *Bmal1*. Furthermore, various TTFL genes and CCGs exhibited altered circadian expression profiles, consistent with an advanced in daily behavioural rhythms under 12h light- 12h dark conditions. Together, these findings reveal the extensive genome-wide regulation mediated by ZFHX3 in the central clock that orchestrates daily timekeeping in mammals.

 INTRODUCTION

Circadian (approximately one day) clocks are internal biological time-keepers co-ordinating coherent 24-h molecular, behavioural and physiological rhythms that are synchronized with changing environmental conditions across day and night. In mammals, the circadian clock mechanism consists of cell-autonomous transcription-translation feedback loops (TTFLs) that drive rhythmic, ~24-h expression patterns of canonical clock components (Takahashi et al., 2008). The primary TTFL involves rhythmic transcription of period (*Per1*, *Per2*, and *Per3*) and cryptochrome genes (*Cry1* and *Cry2*). Thereafter, PER and CRY proteins form heterodimers that act on the dimeric complex of CLOCK (circadian locomotor output cycles kaput) and BMAL1 (brain and muscle ARNT-Like 1) to repress their own transcription, which is mediated by E-box (*CACGTG*) cis-regulatory sequences at their gene promoters. A secondary feedback loop is driven by *Rev-erbα* (*Nr1d1*), which is also directly transcribed through binding of CLOCK:BMAL1 at E-boxes. REV-ERBA feeds back to repress *Bmal1* transcription by competing with a retinoic acid-related orphan receptor (ROR) activator to bind ROR response elements (RREs) in the *Bmal1* promoter. Along with this, CLOCK:BMAL1 heterodimers also trans-activate a series of clock-controlled gene (CCGs), many of which encode transcription factors (TFs) that regulate additional downstream programmes of gene expression (Herzog et al., 2017; Hogenesch et al., 2003).

At an organismal level, mammalian circadian rhythms are directed by the suprachiasmatic nucleus (SCN) of the hypothalamus, which synchronizes robust 24-h oscillations across peripheral tissues (Hastings et al., 2018, 2019; Schibler et al., 2015). In the SCN, the cell-autonomous TTFL molecular clocks in individual cells are synchronized through intercellular coupling to form a coherent oscillatory network (Herzog et al., 2017; Maywood et al., 2021).

This intercellular coupling is maintained through several neuropeptide neurotransmitter

systems, including vasoactive intestinal peptide (VIP), arginine vasopressin (AVP) and

gastrin-releasing peptide (GRP) systems (Brown et al., 2005), while Prokinetin2 (PROK2)

serves as a pace-making element in the SCN (Morris et al., 2021). Moreover, zinc finger

homeobox-3, ZFHX3, is a key transcription factor in the SCN. It is highly expressed in

discrete populations of adult SCN cells and exerts robust control over circadian behavioural

rhythms. Intriguingly, while a dominant missense mutation in *Zfhx3* (short circuit, *Zfhx3*$^{Sci/+}$)

results in a short circadian period in mice (Parsons et al., 2015), a conditional null *Zfhx3*

mutation results in more severe behaviour with short circadian period and arrhythmic mice

(Wilcox et al., 2017).

Given the crucial role of the SCN in orchestrating daily timekeeping in mammals, it is

imperative to understand the role of ZFHX3 in regulating the molecular circadian clock, but

although the effects of mutations in *Zfhx3* are known at the level of circadian behaviour, its

molecular functions that underpin daily timekeeping are not well-characterized. Therefore,

we set out to assess systematically the function of ZFHX3 by employing a multi-omics

approach in the SCN. First, we analysed the genomic binding pattern of ZFHX3 by

conducting ZFHX3 targeted chromatin immunoprecipitation (ChIP-seq) of SCN tissue to

identify the genomic scope of ZFHX3 and its potential mechanistic roles. Second, we

implemented a detailed transcriptional profiling of the SCN in control and *Zfhx3* conditional

null mutants, using vibratome based microdissection (Bafna, Lau, et al., 2023). This allowed

us to map the effect of ZFHX3 on tissue-specific expression of both rhythmic and non-

rhythmic genes.

Overall, we have defined genome-wide occupancy of ZFHX3 in the SCN, showing that it is

found predominantly around actively transcribed genes and is accompanied by modified

histones and other SCN TFs. We noted that conditional (adult) deletion of ZFHX3 drastically

modifies the SCN transcriptome by affecting genes important for circadian timekeeping,

neuropeptide signalling and synaptic processes. An in-depth investigation on time-of-day

dependent gene expression clearly highlighted loss of *Bmal1* rhythmic transcription along

with aberrant 24-h oscillations for TTFL and non-TTFL genes. This change in rhythmic clock

gene transcriptional profiles contributed towards a phase-advanced internal central clock

mirroring the advance seen in the daily behavioural activity patterns of the null mutant mice.

These findings reveal a pivotal and genome-wide physiological role for ZFHX3 in the

maintenance of robust circadian timekeeping in the SCN and identify molecular substrates

that underpin this role.

RESULTS

ZFHX3 *binds to active promoter sites in the SCN*

Genome-wide binding of ZFHX3 was profiled in the SCN using ChIP-seq. Briefly, genomic

DNA was extracted from C57BL/6J mice raised in standard 12-h/12-h light-dark conditions

at two distinct time-points; *zeitgeber* time (ZT) 03 (3 hours after lights on) and ZT15, and

subjected to ChIP-seq (Methods). The sequenced reads were aligned to the mouse genome

and significant sites of ZFHX3 occupancy were assessed in the context of previously

identified modified histone binding regions (Bafna, Banks, et al., 2023) in the SCN. We

found a total of 43,236 ZFHX3-bound sites, overlapping with approximately 22% of total

modified histone regions in the SCN genome. About 60% of the ZFHX3 bound sites were

found near promoters (<= 3 Kb of TSS) and so would be likely to identify direct

transcriptional targets of ZFHX3, whilst ca. 21% were in distal intergenic regions potentially

involved in other modes of gene regulation.

When assessed in relation to modified histone binding, ZFHX3 sites fell into four categories.

First, the majority of ZFHX3 peaks coincided with both H3K4me3 (trimethylation of Histone

H3 at Lysine 4) and H3K27ac (Histone 3 lysine 27 acetylation) occupancy (48.9% of total

ZFHX3 peaks at ZT3, Supplementary Table 1), and were focussed near transcription start

sites (TSS $\pm$ 3kb) (Fig. 1B). Since such sites are likely responsible for active transcription

(Beacon et al., 2021), the ZFHX3 occupancy in these sites highlighted probable

transcriptional regulation by ZFHX3 in the SCN. Next, a subset of ZFHX3 sites (20.7%)

were bound only by H3K4me3 (no H3K27ac) as illustrated at Fig. 1C. These are indicative of

poised transcriptional regulation (Calo & Wysocka, 2013) and were also enriched near the

TSS. About 8.2 % of ZFHX3 sites were bound by H3K27ac exclusively (no H3K4me3) (Fig.

1D). These sites were commonly found at intergenic and intronic regions and suggestive of

enhancer-mediated gene regulation (Won et al., 2008). Finally, 22% of ZFHX3 sites were

found to be devoid of modified histone binding and were also concentrated at distal

intergenic regions (39.04%) (Fig. 1E). Overall, the broad occurrence of ZFHX3 alongside

histone modifiers near the gene TSS and at distal intergenic regions in the SCN genome

suggests its crucial role in gene regulation potentially via promoter-enhancer interaction.

Next, we assigned each ZFHX3 site to its closest transcribed gene ($\pm$ 3Kb) and performed

functional annotation upon these genes. The functional annotation of genes which were

associated with ZFHX3 and both the H3K4me3 and H3K27ac histone modifications were

found to be associated with neurodegeneration and axonal guidance, synaptic processes

known to be important in the SCN (glutamatergic, GABAeric and dopaminergic systems) and

various signalling pathways. Importantly, they also included pathways involved with circadian entrainment and timekeeping (Supplementary Fig. 1A), highlighting the reciprocal intersection between ZFHX3 and the SCN cell-autonomous clock. In contrast, genomic regions co-operatively bound by ZFHX3 and H3K4me3 but with no H3K27ac, or sites with ZFHX3 alone, were found to be associated with genes involved in several fundamental biological processes such as DNA replication or MAPK signalling, but notably not circadian rhythm (Supplementary Fig. 1B, C). Therefore, it is plausible that ZFHX3 binds at the TSS and regulates actively transcribing genes, particularly those involved in tissue-specific functions such as daily timekeeping, while it's occupancy without histone modifiers and/or at poised state regulates genes responsible for other basal functions. Interestingly, it is worth noting that ZFHX3 occupancy did not differ appreciably between the two time-points tested (ZT03 vs. ZT15): ZFHX3 was seen to bind consistently at the same genomic loci (Supplementary Table 1). This indicates that other factors/ mechanisms confer temporal control to these sites whilst the presence of ZFHX3 is permissive to such cis- and trans-regulatory cues.

We next examined the signature DNA binding motifs/ cis-regulatory elements occupied by ZFHX3, either alone or with specific histone modifications (H3K4me3, H3K27ac). As anticipated, this revealed an over-representation of AT-rich motifs (Parsons et al., 2015). Notably, it also suggested co-occurrence of ZFHX3 with SCN developmental and maintenance TFs such as DLX (distal-less homeobox) and LHX (LIM homeobox) (VanDunk et al., 2011) across the tested categories (Supplementary Fig. 1D). Furthermore, ZFHX3-bound sites were also enriched for non-AT-rich motifs such as the RFX family of TFs, which have been implicated in transcriptional responses to light in the SCN (Araki et al., 2004). Interestingly, we noted enrichment of ARNT (bHLH) domains but only where ZFHX3 peaks

were seen adjacent to H3K27ac and H3K4me3 (i.e., active promoters). This clearly points

towards the possibility of shared binding of ZFHX3 with the circadian clock TFs

BMAL1:CLOCK at E-boxes and highlights a point of interaction between the core TTFL and

ZFHX3 in the regulation of circadian gene transcription in the central pacemaker.

Zfhx3 knock-out affects circadian behaviour and entrainment of mice

To explore the contribution of _Zfhx3_ to SCN function in the adult animal, we compared

circadian wheel-running in cohorts of _Zfhx3_$^{\text{Flox/Flox}}$;UBC- mice that had been dosed with

tamoxifen (gene deletion, +Tamoxifen group) with mice that had not been dosed with

tamoxifen (no deletion genetic control, -Tamoxifen group). Comparisons of gene expression

between the –Tamoxifen and +Tamoxifen groups by real time PCR confirmed loss of _Zfhx3_

expression in the +Tamoxifen group. (Supplementary Fig. 2A). Behavioural analysis was

performed in conditions of constant darkness and in a 12 h light/12 h dark cycle. Consistent

with previous studies, this analysis showed that the free-running circadian period was

significantly shortened by conditional deletion of _Zfhx3_ (Fig. 2A). Furthermore, the overall

activity rhythm was less well defined with significantly reduced amplitude, increased intra-

daily variability and lower inter-daily stability in the deleted mice (Supplementary Fig. 2B, C,

D). The full analysis is given in Supplementary Table 2. The loss of ZFHX3 therefore

compromised free-running circadian behaviour.

Visual inspection of the actograms whilst the animals were in light/dark cycles suggested

that animals with gene deletion showed poor entrainment to the light cycle, with animals

showing increased activity in the hours prior to the light to dark transition (Fig. 2A, right). To

quantify this, we analysed the distribution of wheel running activity across 24-h, by

expressing wheel running activity within each 30 minute segment of the day as a proportion

of the total daily wheel running activity for that animal (Fig.2C). Repeated Measures

ANOVA demonstrated a significant interaction between tamoxifen dosing and time of day

(tamoxifen dosing x time: $F_{(47, 1316)} = 9.013$; p= <0.0001). Pairwise analysis demonstrated

that, compared to genetic control animals, gene deleted animals showed a significant increase

in wheel running activity in the hours prior to lights off (p= <0.05 for ZT 10.5-12.5) and a

significant reduction in wheel running activity in the hours prior to lights on (p= <0.05 for ZT

20 -23). Consistent with this, analysis of the activity onset demonstrated that gene deleted

animals showed an earlier onset of activity than controls (Supplementary Fig. 2E). This phase

advance under 12 h L/12 h D is consistent with the shortening of circadian period seen under

DD. Thus, ZFHX3 contributes to circadian time-keeping and entrainment in the adult SCN,

independently of any developmental roles.

ZFHX3 knock-out (KO) results in extensive effects across the SCN transcriptome

Having mapped the genomic binding sites of ZFHX3 and validated the behavioural effects of

deletion of ZFHX3, we then employed total directional RNA sequencing to define the impact

of *Zfhx3* deletion on the SCN transcriptome. Adult SCN tissues were collected from

Zfhx3$^{Flox/Flox}$;UBC-Cre$^-$ (Cre-neg) and *Zfhx3*$^{Flox/Flox}$;UBC-Cre$^+$ mice that were not treated

with tamoxifen (-Tamoxifen, genetic control) , both serving as independent control groups,

and tamoxifen treated *Zfhx3*$^{Flox/Flox}$;UBC-Cre$^+$ (+Tamoxifen) mice i.e., *Zfhx3*-deleted, at six

distinct times of light/dark cycle (Methods). As an initial analysis we compared overall (i.e.,

time-independent) gene expression levels between the groups (Cre-neg vs. +Tamoxifen) to

examine the global effect of *Zfhx3* deletion on the SCN transcriptome, and found that almost

36% (n = 5,637) of total assessed genes were affected by the acute loss of *Zfhx3* in the SCN.

Of these, 2,725 were down-regulated and 2,912 up-regulated in +Tamoxifen animals

compared to Cre-neg controls. (Supplementary Table 3, Fig. 3A). Furthermore, as expected,

both the control groups (Cre-neg and -Tamoxifen) showed consistent differential gene

expression levels when compared to the mutant which mitigated any potential downstream

effect of tamoxifen dosing *per se* (Fig. 3B, C).

To better understand the relationship between ZFHX3 occupancy on the genome and changes

in gene expression, we compared the results of our initial ChIP-seq with the RNA-seq

described above. Of the total differentially expressed genes, 80% (n = 4,543) revealed

ZFHX3 occupancy at their promoter sites along with histone modifications (H3K4me3 and

H3K27ac), showing a direct interaction between ZFHX3 bound TSS ($\pm$ 3Kb) and gene

expression (p = 0, odds ratio = 6.3, Jaccard index = 0.4, signifying strong association). In

contrast, only 7% of differentially expressed genes (n = 381) were solely bound by ZFHX3 at

the TSS (p = 0.37, odds ratio = 1, Jaccard Index = 0.1, signifying weak association),

indicating the effect of ZFHX3 on gene transcription is predominantly observed when jointly

occupied by histone modification at gene TSS.

We then sub-selected highly differential genes ($\log_2 FC \geq 1$, FDR ≤ 0.05) and conducted

functional gene enrichment analysis. Consistent with previous studies (Hughes et al., 2021;

Parsons et al., 2015; Wilcox et al., 2021; Wilcox et al., 2017), genes that were downregulated

were seen to be associated with neuropeptide signalling pathways (*Vip, Grp*), circadian

rhythm, neuron differentiation etc., while those that were upregulated showed functional

enrichment for cell differentiation and extracellular matrix organization processes (Fig. 3D).

Interestingly, our comprehensive assessment also highlighted an effect of loss of ZFHX3 on

signal transduction by affecting cholinergic receptor (*Chrna3, Chrnb4*), melatonin receptor

(*Mtnr1a*), prokineticin receptor (*Prokr2*), GRP receptor (*Grpr*), and the SCN peptide *Nmu*.

Therefore, it is likely that loss of ZFHX3 leads to physiological impairment by affecting a

diversity of neuropeptide neurotransmitter systems in the SCN and more generally, circadian-

relevant transcripts.

.

Loss of Zfhx3 affects rhythmic transcription of TTFL and clock-controlled genes

The present and recent studies have demonstrated severe disturbances of daily and circadian

behaviour in mice carrying a missense mutation (*Zfhx3^{Sci}*) (Parsons et al., 2015) or knock-out

(*Zfhx3$^{Flox/Flox}$*;UBC-Cre$^+$) (Wilcox et al., 2017) of ZFHX3. In addition to assessing general

gene up- or down-regulation in mutants (above), we next wanted to assess how deletion of

ZFHX3 might alter transcriptional rhythms. For this, we investigated gene expression levels

in the SCN at six distinct times-of-day (Hughes et al., 2017), starting at ZT02, at 4-h intervals

(Methods). In total, we analysed the daily transcriptional profile of 20,521 genes in the SCN

and compared the rhythmicity index between the control (Cre-neg) and mutant (+Tamoxifen)

groups using statistical framework *dryR* (Weger et al., 2021). This enabled us to categorise

the genes into one of 5 distinct modules (Methods): Module 1 refers to genes with no

detected 24-h rhythm in their expression levels in either condition (n = 16832, Supplementary

Fig. 3A); Module 2 consists of genes that were rhythmic in control SCN but lost rhythmicity

in the ZFHX3-KO (n =1566, Fig. 4A); Module 3 consists of genes that gained daily

rhythmicity after ZFHX3-KO (n = 403, Supplementary Fig. 3B); Module 4 was composed of

rhythmic genes that exhibited no change between control and mutants (n = 1653, Fig. 4B);

Module 5 included genes that retained rhythmic expression but with a change in either

amplitude or phase (peak expression) in the mutant group (n = 67) (Supplementary Table 4,

Fig. 5).

Thus, we noted ~82% of compared genes did not show a daily rhythm (module 1).

Conversely, ~18% of gene expressed in the SCN exhibited daily rhythmicity under one or

other condition. Of these, ~45% were not affected by loss of ZFHX3, whereas ~55% of

rhythmic transcripts showed alteration in their daily pattern following loss of ZFHX3. Indeed,

~42% of rhythmic genes required ZFHX3 to maintain that rhythm, whereas ~2% of genes

(403/20521) gained rhythmicity in the absence of ZFHX3. Thus, ZFHX3 contributes, directly

or indirectly, to a significant part of the daily rhythmic transcriptome of the SCN.

For each module, we compiled the gene list and examined the functional enrichment

pathways. The genes that lost rhythmic transcription after the deletion of ZFHX3 (module 2)

were seen to be involved in intracellular protein transport, signal transduction such as WNT-

pathway linking circadian and cell-cycle processes (Matsu-Ura et al., 2018), oxidative stress

response etc. (Supplementary Fig. 4A). On the contrary, genes that gain rhythmicity (module

3) were associated with RNA processing, stress and metabolic pathways. The genes that did

not show any change in rhythmic expression (module 4) were found to be enriched for

functional processes such as protein phosphorylation and de-phosphorylation, cytoskeleton

organization, DNA repair etc. Interestingly, only a tiny proportion (~1.8%) of the rhythmic

genes showed substantial changes in their daily cyclic gene expression profiles (phase and

amplitude, module 5), but these were primarily involved in circadian rhythm generation,

entrainment and clock-controlled gene expression (Supplementary Fig. 4B). This suggests

that in the absence of ZFHX3 the core circadian mechanism of the SCN is disrupted but is

not ablated.

Indeed, almost all the genes driving the cell-autonomous TTFL were severely affected by

Zfhx3 perturbation with *Bmal1* demonstrating a complete loss of 24-h rhythm, and its

counterpart *Clock* mRNA showing overall reduced expression levels (Fig. 4A). PER family

genes (*Per1-3*) showed a significant advancement in peak expression with low amplitude

(Fig. 5C). Specifically, we observed a phase advance of ca. 2-h for *Per2* and *Per3,* with *Per1*

(Kuhlman et al., 2003) which is also known to mediate light-responsive pathways, advancing

by only ~0.3-h. Strikingly, there was no obvious change in the rhythmic levels or phase for

CRY family genes (*Cry1, 2*) (Fig. 4B), indicating that ZFHX3 acts only on select genes

driving the cell-autonomous clock in the SCN. Given that *Bmal1* transcription is under the

control of ROR (*Rora, b, c*) and REV-ERB (*Rev-erb a, b*) factors (Liu et al., 2008), we also

noted loss of the rhythm in *Rorb* expression*,* whilst *Rev-erba* showed a phase advance by

almost one hour (Table1).

Next, we examined the effect of loss of ZFHX3 on rhythmic expression of the clock-

controlled genes (CCGs). Clock-controlled genes are basically genes whose expression is

regulated by circadian clock factors (such as CLOCK:BMAL1) that are directly associated

with the TTFL. Several CCGs like DBP (D site albumin promoter binding protein) show

rhythmic transcription (Lopez-Molina et al., 1997), and in turn control expression of a

repertoire of genes, generating a cascade of 24-h cycling genes in the SCN (Herzog et al.,

2017). In our mutant group, *Dbp* mRNA levels also presented an advanced peak (~1.66 h)

with reduced amplitude (Fig. 5C). Another such TF- *Creb3l1* (cAMP responsive element

binding protein 3-like 1), that has been implicated in intercellular coupling (Bedont et al.,

2017), showed a distinctive change in phase and amplitude. Therefore, the alteration seen in

the rhythmicity index at the level of TTFL genes due to the loss of ZFHX3 was recapitulated

in the diurnal expression of certain CCGs in the SCN.

Finally, we studied the effect of ZFHX3 deletion on the rhythmic pattern of intercellular

coupling agents (neuropeptides), whose overall mean expression levels were significantly

reduced (Fig.3, *Vip, Grp*). Along with lowered baseline expression, we noted severe loss in

24-h rhythm for crucial SCN neuropeptides such as *Avp* (Fig. 3a), *Prok2* and *Npy* (Fig. 5c).

For *Avp*, there was no apparent increase in daytime mRNA expression (Arima et al., 2002),

and the levels remained intermediate throughout the day. Conversely, for both *Prok2* and

Npy, transcript levels were reduced to minimal levels, possibly resulting in compromised

intercellular-coupling in the SCN.

The effects of ZFHX3 on daily transcriptional profiles therefore varied between different

gene categories. Whereas rhythmicity of downstream, clock-controlled genes was either

independent of ZFHX3 or required ZFHX3 to be sustained, core circadian function was

sustained but with altered dynamics. Intriguingly, this systematic approach not only

highlighted the genes that lost 24-h rhythmic oscillations, such as *Bmal1* and the

neuropeptide *Avp* (module 2, Fig. 4A), but also identified genes that retained rhythmicity

with changes in their waveform (module 5). Given the fact that ZFHX3 deficient mice do not

lose rhythmic behaviour in a daily LD cycle (Wilcox et al., 2017) (Fig. 2), but do exhibit a

short circadian period or arrhythmicity in free-running (complete darkness: DD) conditions,

we believe our targeted SCN-specific assessment unravels the vital role of ZFHX3 in the

mechanistic regulation of the daily molecular clock.

Table 1: *Effect of loss of ZFHX3 on core-clock genes in the SCN*

Gene	Rhythm	Phase	Amplitude
Per1	Altered	-0.36	0.72 (low)
Per2	Altered	-2.0	0.69 (low)
Per3	Altered	-2.1	0.30 (low)
Cry1	No change	-	-
Cry2	No change	-	-
Bmal1	Lost	-	-
Clock	No change	-	-
Nr1d1	Altered	-0.99	0.67 (low)
Nr1d2	No change	-	-
Rorb	Lost	-	-

The state of the daily SCN clock in ZFHX3 deficient mice

In order to link the perturbations of the core molecular clock in the SCN of ZFHX3 deficient

mice with their altered behavioural profiles, we used TimeTeller, a recently established

method to predict circadian clock phase alongside the parameter 'Theta' which provides an

index of clock disruption from transcriptomic data (Vlachou et al., 2024). This approach

allows one to assess the clock globally as a system instead of gene by gene. For this, an initial

training model was built using the available Cre-neg and -Tamoxifen (n = 35, control) mice

data (Fig. 6A). Cross-validation for the training model suggested that the model is

informative and predicts the control SCN time well with a median prediction error of -0.08

hours (95% CI: -0.83;1.33) and 0.50 hours (95% CI: -1.67;3.00), i.e., the difference between

the TimeTeller predicted time and the known ZT of SCN collection (Supplementary Fig. 5A).

This was further assessed by predicting SCN timing in an independent dataset of SCN RNA-

Seq (Deota et al., 2023). Similar to the TimeTeller predictions for the control group, median

prediction error was 0.00 hours (95% CI: -2.00;1.33) for the *ad libitum* fed and -0.66 hours

(95% CI: -1.83;1.00) for the temporary night-time restricted feeding groups. These are not

distinguishable from the control SCN in the current study (Fig. 6B). Similarly, the clock

disruption parameter theta is low (Medians: 0.03, 0.03 and 0.04) as is expected for a fully

functional and well predictable circadian clock in all three control groups. Furthermore, TT

clock functionality parameter theta suggests that +Tamoxifen SCN have a similarly

functional daily clock compared to the control SCN (Median 0.04 vs. 0.03 in controls

(Supplementary Fig. 5B).

While TT's theta metric suggests the clock in the ZFHX3-deleted SCN is not disrupted as it

is indistinguishable from controls, the phase of the +Tamoxifen SCN was consistently

advanced by a median 1.83 hours (95.0% CI 0.585, 2.67), when compared to the current

controls and the published data (Deota et al., 2023) as illustrated in Fig.6B. Of note, detecting

this ~2-h phase difference from data sampled at 4-h intervals showcases the advantages in

robustness and sensitivity of assessing the clock as a system rather than by a gene-by-gene

approach, and TimeTeller as a tool of inquiry. In summary, according to TimeTeller, the SCN

daily clock is intact but advanced by approximately 2-h in comparison to the control groups

after loss of ZFHX3. This is consistent with the observed daily behavioural activity in mutant

mice as shown in Fig. 2.

<u>DISCUSSION</u>

The SCN is distinctive in its ability to sustain autonomous rhythmicity which is driven by

circadian cascades of transcription that direct metabolic, electrophysiological and signalling

rhythms. Our current findings highlight the instrumental role of the transcription factor,

ZFHX3, in driving the daily timekeeping mechanism. By comparing the daily behavioural

activity of ZFHX3-KO mice with the genetic matched control, we clearly noted an

advancement in LD activity and shortening of circadian period in mutants. This finding is

consistent with previous studies (Parsons et al., 2015; Wilcox et al., 2021; Wilcox et al.,

2017) and prompted a detailed molecular investigation in the SCN. Through targeted ChIP-

seq we successfully demonstrated the genome-wide occupancy of ZFHX3 in SCN chromatin

with 78% of ZFHX3 bound sites found to be co-localized with modified histones

(H3K4me3, H3K27ac), and a vast majority of these concentrated near gene TSS . This

provides strong evidence in support of the observed pervasive regulation conferred by

ZFHX3 on the SCN transcriptome. That said, a considerable proportion of ZFHX3 sites were

also seen devoid of joint histone modifications (H3K4me3 + H3K27ac) and enriched for

CTCF binding sites. CTCF is a ubiquitous DNA binding protein that has exceptional roles in

genomic organization, chromatin looping and gene regulation (Holwerda & de Laat, 2013;

Kim et al., 2015; Mach et al., 2022). This co-occurrence hints on the potential accessory role

of ZFHX3 in modifying genome topology and chromosomal looping to bring about changes

in target gene expression by regulating promoter-enhancer interactions (Pugacheva et al.,

2020). Interestingly, along with CTCF binding sites, ZFHX3 sites were also enriched for

other SCN enriched regulatory factors; RFX, LHX, ZIC (Araki et al., 2004; Hatori et al.,

2014) (Supplementary fig. 1D), suggesting well-coordinated tissue-specific gene regulation.

Given the crucial role of ZFHX3 in daily timekeeping, we speculated a dynamic change in its

genomic occupancy and compared the binding intensity of the transcription factor at two anti-

phasic (12-h apart, ZT3 vs. ZT15) time-points (Methods). Surprisingly, we did not detect any

differential binding site(s) between the tested time-points, but rather noticed a uniform and

coherent localization. This suggests that ZFHX3 belongs to a family of poised transcription

factors, whose occupancy at regulatory sites does not change during the day, but could

potentially act as a co-activator/mediator to modulate oscillating gene transcription. Indeed,

in a recent unpublished study (Del Rocío Pérez Baca et al., 2023), ZFHX3 was shown to be

associated with chromatin remodelling and mRNA processing factors, strengthening the viewpoint as a co-regulator. It is, therefore, highly plausible that ZFHX3 occupies the targeted TSS and partners with clock TFs such as BMAL1 *(CACGTG)*, as observed around active sites, to regulate rhythmic gene transcription.

Along with ChIP-seq, we executed detailed RNA sequencing to study the effect of loss of ZFHX3 specifically in the SCN. Crucially, ZFHX3 affected almost a third of the transcriptome in the SCN. An overall (time-independent) assessment of deletion of *Zfhx3* highlighted a significant impact on expression of genes involved in several biological processes ranging from cellular differentiation to behavioural responses. We then performed a comprehensive time-dependent assessment to delineate the functional role of ZFHX3 in regulating the circadian timekeeping mechanism. We observed about 55% of rhythmic genes showed alteration at distinct levels in their daily transcriptional pattern following loss of ZFHX3.

At the TTFL level, ZFHX3-KO resulted in the loss of rhythmic *Bmal1* gene expression. It remains to be determined whether this is associated with changes at the posttranslational level, given the findings by Abe et al where arrhythmic *Bmal1* transcript did possess posttranslational rhythm (Abe et al., 2022), and could explain the subsequent 24-h driven *Per (1-3)* and *Cry (1-2)* mRNA rhythms noted in our study. Likewise, in a separate study constitutive *Bmal1* expression has been shown to sustain rhythmic levels of transcriptional and translational products of *Per, Cry, Clock, RORc,* and *Rev-erbα* by means of mathematic modelling (Mirsky et al., 2009). Arguably, clear circadian rhythmicity in BMAL1, CLOCK, or REV-ERBα protein levels were not sustained in association with constitutive *Bmal1* levels

under *in-vitro* conditions (Padlom et al., 2022). Therefore, in future it will be interesting to

study the daily abundance of protein levels of the clock genes in *Zfhx3*-deficient SCN, with

its associated loss of *Bmal1* mRNA 24-h rhythm, to gain a complete understanding of the

clock molecular circuitry.

Broadly speaking, aside from *Bmal1*, the majority of clock genes did not lose 24-h

rhythmicity but showed an early peak in expression, ranging between ~0.4 – 2-h that

perfectly resonates with the advanced rhythmic behaviour seen in mutants. In this case, it is

difficult to delineate a straightforward cause-and-effect axis in terms of the direct targets of

the perturbed ZFHX3 activity. Based on the current data, ZFHX3 is possibly partnering with

CLOCK:BMAL1 to regulate the daily transcription of clock genes (*Per, Ror, Reverb)* and

CCGs. Two possible scenarios could explain the noted effect on the TTFL genes due to lack

of ZFHX3 (Fig. 5). First, ZFHX3 preferentially controls *Per* family gene transcription (and

not *Cry*) by binding at the promoter (Fig. 1). Loss of ZFHX3 affects the daily rhythm of *Per*

gene transcription, and potentially PER protein levels, which in turn disturbs the

CLOCK:BMAL1 driven transcription cascades. This leads to an aberrant daily transcriptional

profile of the *Ror* (activator)/ *Reverb* (repressor) genes as observed, which are known to

control the amplitude and oscillation of *Bmal1* (Liu et al., 2008), and further results in loss of

Bmal1 transcriptional rhythm. Therefore, the observed change in *Ror* and *Reverb* family gene

expression levels could be reminiscent of the effect of ZFHX3 on the TTFL. In a second

scenario, the constitutive expression of *Bmal1* mRNA could result from the primary effect of

ZFHX3 on CLOCK:BMAL1 driven *Ror/Reverb* genes (affecting ROR/REVERB

translational products), which does not disrupt the TTFL, but weakens the daily clock. The

direct /indirect effect is seen in the cyclic expression of *Per* family genes, with relatively

greater phase change in *Per2* and *Per3* than the *Per1* gene (Kuhlman et al., 2003)

(Supplementary fig. 6). It is, however, noteworthy that the CRY family of genes *(Cry1, Cry2)*

did not show any obvious change in the daily expression and is supposedly shielding the

molecular clock , as recently postulated by Abe et al (Abe et al., 2022).

As previously reported (Parsons et al., 2015) , lack of ZFHX3 also resulted in severe

reduction in the SCN neuropeptides such as *Avp, Vip, Grp, Prok2*, that potentially

compromises intercellular coupling and synchronization of circadian timekeeping. This

universal effect, where loss of ZFHX3 does not deter a particular but all the key SCN

neuropeptides, clearly points that 1) ZFHX3 is prevalent in almost all neuronal populations

within the SCN, that are largely clustered based on their neuropeptide expression (Morris et

al., 2021; Wen et al., 2020) 2) potentially results in loosely coupled intercellular clocks that

are still able to drive 24-h behavioural and molecular rhythms but with advanced phase. The

prevalence of ZFHX3 in neuronal populations is also supported by a complete independent

assessment in which almost 60% of the top 100 categorised neuronal genes in the SCN (as

per single-cell atlas) were seen to be affected by ZFHX3 deletion based on the current

dataset. In contrast, only 20% of the 100 most expressed genes in the astrocytic cell

populations were impacted.

Along with this, lack of *Zfhx3* also leads to aberrant daily transcriptional rhythms of several

CCGs, with the majority of them seen to lose daily oscillations. This could potentially result

in hindrance of clock-regulated vital biological processes such as regulation of cell-cycle,

metabolic pathways etc. Indeed, ZFHX3 is reported to be crucial for the development of a

functional SCN (Wilcox et al., 2021), and could act as an 'integrator' of afferent signals such

as light and feeding to generate timely and robust physiological outputs. Moreover, genetic

ablation of *Zfhx3* has been shown to result in a compromised light sensing mechanism in

retina (Hughes et al., 2021), while the missense mutation ($Zfxh3^{Sci/+}$) results in reduced

growth and altered metabolic gene expression in the arcuate nucleus of the hypothalamus

(Nolan et al., 2023) converging on its functional roles in other brain regions. The pleitropic

roles of ZFHX3 further extend in non-brain tissues with its recently described crucial

involvement in angiogenesis for tumour development (Fu et al., 2020), cardiac function

(Jameson et al., 2023) and spinocerebellar ataxia (Figueroa et al., 2024; Wallenius et al.,

2024). It is important to note that although in the present study we used adult-specific *Zfhx3*

null mutants resulting in global loss of ZFHX3, the effects observed both at molecular and

behavioural levels are independent of its functional role(s) in other tissues. In future,

however, it will be beneficial to carry out the investigation using an inducible, SCN specific

genetically ablated model to circumvent any confounding effects. Regardless, the current

research is of great importance as it not only maps the genomic binding of ZFHX3, but also

demonstrates its critical association with both lineage- and clock-controlling TFs that

perfectly resonates with the observed profound effects of ZFHX3 on the SCN transcriptome,

regulating a vast majority of genes ranging from circadian timing to synaptic transmission

and inflammatory responses. This highlights how ZFHX3 regulates the core-clock (TTFL) in

addition to its control over CCGs in order to drive sustained transcriptional rhythms in the

SCN. Given the extensive pleiotropy of ZFHX3, it provides a solid substrate to further

understand the functional interaction of ZFHX3 in distinct tissues indispensable for overall

health and fitness.

METHODS

Mice

All animal studies were performed under the guidance issued by Medical Research Council

in Responsibility in the Use of Animals for Medical Research (July1993) and Home Office

Project Licence 19/0004. WT C57BL/6J were maintained and provided in-house by MRC

Harwell. Animals were group housed (4-5 mice per age) in individually ventilated cages

under 12:12h light-dark conditions with food and water available ad libitum. Adult specific

knock out *Zfhx3* mice were generated using an inducible Cre line as described by Wilcox et al

(Wilcox et al., 2017). Briefly, tamoxifen inducible Cre mice (UBC-Cre) were crossed to

Zfhx3 floxed (*Zfhx3*^Flox^) mice to produce an initial stock of *Zfhx3*^Flox/+^;UBC-Cre^+^ mice. These

were subsequently crossed to *Zfhx3*^Flox/+^ mice to generate experimental cohorts (Sun et al.,

2012). Out of the 6 possible genotype combinations, 2 genotypes; *Zfhx3*^Flox/Flox^;UBC-Cre^−^

(mice homozygous for *Zfhx3*^Flox^ allele but not carrying the Cre allele) and *Zfhx3*^Flox/Flox^;UBC-

Cre^+^ (mice homozygous for *Zfhx3*^Flox^ allele and hemizygous for Cre allele) were assigned as

"control" (Cre-neg) and "experimental" (+Tamoxifen) groups, respectively, and dosed with

tamoxifen. In addition, we also included a cohort of *Zfhx3*^Flox/Flox^;UBC-Cre^+^ mice without

tamoxifen treatment as a secondary control (-Tamoxifen). Animals were group housed (4-5

mice per age) in individually ventilated cages under 12:12h light-dark conditions with food

and water available ad libitum.

Experimental Design

WT C57BL/6J mice (aged between 8 to 12 weeks) were used for SCN tissue punch collection

as described by Jagannath et al. (Jagannath et al., 2013) at ZT3 and ZT15, where lights on at

7 am (ZT0) and lights off at 7 pm (ZT12), for ZFHX3-ChiP-seq. For SCN RNA-Seq, mice

from control (Cre-neg and -Tamoxifen) and *Zfhx3* knock out (+Tamoxifen) cohorts were used

for SCN tissue collection (Bafna, Lau, et al., 2023) at 6 distinct time-points starting from ZT2

at every 4h. To achieve high resolution transcriptomic dataset, four biological replicates per

time-point were collected independently from the control (Cre-neg) and experimental

(+Tamoxifen) groups. For the second control group viz. -Tamoxifen; 2-4 biological replicates

were collected due to time constraints. Furthermore, each biological replicate for all the three

tested condition constituted 3 individual SCN samples.

<u>Animal behaviour</u>

Circadian wheel running was performed as previously described (Banks & Nolan, 2011).

Briefly, mice were singly housed in cages containing running wheels, placed in light

controlled chambers and wheel running activity monitored via ClockLab (Actimetrics).

Animals were given a five-day acclimatisation period in a 12-h light/dark cycle (100 lux light

intensity) before their activity was monitored for 32 days in constant darkness and 24 days

under a 12-h light/dark cycle (100 lux light intensity). Circadian activity analysis was

performed upon the entire of the period in constant darkness and upon the final seven days in

the light/dark cycle. Circadian analysis was performed using ClockLab Version 6

(https://actimetrics.com/products/clocklab/clocklab-analysis-version-6/). *Zfhx3*$^{\text{Flox/Flox}}$;UBC-

Cre$^+$ animals were used for this analysis, with animals being randomly assigned into

tamoxifen un-dosed (-Tamoxifen) or tamoxifen dosed (+Tamoxifen) cohorts. The numbers of

animals per cohort were: male -Tamoxifen = 8; male +Tamoxifen = 8; female -Tamoxifen =

8; female +Tamoxifen = 8. Statistical analysis and graphing of circadian data was performed

using GraphPad Prism 8.

<u>SCN tissue collection for RNA-seq</u>

Animals were sacrificed at the aforementioned time-points by cervical dislocation and brains

were removed and flash frozen on dry ice. The frozen brains were placed on dry ice along

with the brain matrix (Kent Scientific, Torrington CT, USA) and razor blades to chill. The

brain dissection was performed on a vibrating microtome (7000smz-2 Vibrotome, Campden

Instruments) under chilled conditions (Bafna, Lau, et al., 2023). First, the frozen brain was

placed on the chilled brain matrix and a ca. 3mm thick region was removed from the caudal

end using a razor blade to discard the cerebellum and attain flat surface. Then, the dissected

brain was super-glued to a chilled (2 –3°C) metal chuck and placed into the metal specimen

tray of the Vibratome. The metal specimen tray was filled with pre-chilled 1X RNase free

PBS (phosphate buffered saline, ThermoFisher Scientific, #AM9625). Coronal slices

measuring 250 uM in thickness were cut from rostral to caudal at slow speed (0.07mm/s) and

pre-tested specifications ; Frequency: 70Hz, Amplitude: 1.00mm. Once cut, the slice with

intact optic chiasm, third ventricle and bilateral SCN tissues (Fig. S1) was selected and

transferred on to pre-chilled glass slides (ThermoFisher Scientific SuperFrost Plus Adhesion

slides) with the help of paint brush. The glass slide was placed on top of cold block and SCN

tissues were collected by pre-chilled forceps and dissection needles under dissection

microscope (Nikon SMZ645 Stereo Microscope).

ZFHX3 ChIP and sequencing

ChIP was conducted using the Active Motif ChIP☐IT enzymatic kit

(https://www.activemotif.com/catalog/868/chip-it-high-sensitivity) according to the

manufacturer's instructions. Briefly, SCN punches were pooled from four mice at designated

times (ZT3, ZT15) and immediately fixed for 10 min at room temperature with gentle

shaking in 1% formaldehyde. The cross☐linking reaction was stopped by addition of glycine

to a final concentration of 0.25 M. Cross☐linked chromatin was sheared by enzymatic

digestion following the manufacturer's recommendations. Samples were incubated with 3μg

of antiserum against ZFHX3 (Parsons et al., 2015), together with magnetic beads, on a rotator

at 4°C overnight. This was followed by crosslinking reversal and proteinase☐K digestion as

per the manufacturer's protocol. The recovered DNA was purified using PCR purification

reagents and subjected to quantitative PCR for quality inspection. ZFHX3 immuno-

precipitated genomic DNA along with their corresponding input samples was sequenced on

Illumina sequencing (using NextSeq 500) platform by Active Motif Epigenomic Services.

ZFHX3 ChIP-seq peak calling and data analysis

Raw sequence data was quality assessed by removing low-quality bases (Phred < 20) and

trimmed using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and

Trimmomatic v0.36 (http://www.usadellab.org/cms/index.php?page=trimmomatic),

respectively. FASTQ files containing trimmed sequences were then aligned to mm10 genome

assembly to generate binary alignment map (BAM) files and used for peak calling by MACS

(Zhang et al., 2008) at a threshold of q < 0.01. Finally, peaks from ZT3 and ZT15 were

analysed for differential binding by a Bioconductor package Diffbind v2.10.0

(http://bioconductor.org/packages/release/bioc/vignettes/DiffBind/inst/doc/DiffBind.pdf) at

default settings with p < 0.05 and fold change > $\pm$ 2. No significant differential ZFHX3

bound peaks were detected between ZT3 and ZT15. In order to investigate the ZFHX3 peaks

overlapping with histone modifications (H3K4me3, H3K27ac) in the SCN, data was

downloaded from NCBI Gene Expression Omnibus (GEO;

https://www.ncbi.nlm.nih.gov/geo/) accession number GSE217943 and processed using

BEDTools suite (Quinlan & Hall, 2010). ChIPSeeker package (Yu et al., 2015) was used to

retrieve the nearest gene around resultant peaks and to annotate the genomic region of the

peak.

RNA extraction

Total RNA from each sample replicate was extracted using RNeasy Micro Kit (QIAGEN,

#74004) following the protocol for microdissected cryosections as per manufacturer's

guidelines and stored at -80°C. Quality and quantity of RNA were measured using an Agilent

Bioanalyzer (Pico chip) and a Nanodrop1000 (Thermo Fisher Scientific, Waltham, MA

USA), respectively.

RNA Sequencing and data analysis

PolyA RNA-Seq libraries were prepared by Oxford Genomics Centre, University of Oxford

using NEBNext Ultra II Directional RNA Library Prep Kit (NEB, #E7760) and sequenced on

NovaSeq6000_150PE (150 bp paired-end directional reads: ~40 million reads/sample)

platform. Paired-end FASTQ files were quality assessed (Phred<20 removed) with FastQC

and Illumina adapters were trimmed with TrimGalore (v0.4.3). Then, the reads were aligned

to mm10 genome assembly using STAR (v2.7.8a) with MAPQ value for unique mappers set

to 60. Binary alignment map (BAM) files were used to generate read counts per gene by

FeatureCounts via Samtools (v1.11). Finally, limma-voom method (Liu et al. 2015) from the

Bioconductor package-limma (v3.48.0) was adopted to quantify differential gene expression

and normalised logarithmic CPM (counts per million) values were generated for downstream

analysis. Genes without more than 0.5 CPM in at least 2 samples are insignificant and filtered

out. 8582 of 24421 (35.14%) genes were filtered out for low expression. TMM was the

method used to normalise library sizes.

To study the overlap between genes with ZFHX3 occupancy at TSS and those impacted by

Zfhx3 deletion, R-package GeneOverlap

(https://bioconductor.org/packages/release/bioc/html/GeneOverlap.html) was adopted where

overlap between two gene lists was tested sing Fisher's exact test with P-value = 0 meaning

the overlap is highly significant. The Fisher's exact test also gives an odds ratio which

represents the strength of association. If an odds ratio is equal to or less than 1, there is no

association between the two lists. If the odds ratio is much larger than 1, then the association

is strong. The class also calculates the Jaccard index which measures the similarity between

two lists. The Jaccard index varies between 0 and 1, with 0 meaning there is no similarity

between the two and 1 meaning the two are identical.

Differential rhythm analysis

The read counts per gene (n = 24,421) for each analysed sample was obtained by

FeatureCounts and arranged in sample matrix by using coreutils (v8.25). The matrix file was

used as an input for *dryR* (R package for Differential Rhythmicity Analysis) available at

https://github.com/neaf-lab/dryR . Normalized gene expression from "control" (Cre-neg) and

"experimental" (+Tamoxifen) groups were categorized into 5 modules with a threshold of

BICW (Bayes information criterion weight) > 0.4 based on their time-resolved expression

pattern.

RT-qPCR

Total RNA was extracted from the SCN tissue as described above from three/ four biological

replicates per genotype (-Tamoxifen vs. +Tamoxifen), where each biological replicate

constitute one independent SCN tissue. 10 ng RNA per sample replicate was tested by one-

step RT_qPCR reagent (PrecisionPLUS OneStep RT-qPCR Master Mix with ROX at a

reduced level premixed with SYBRgreen, PrimerDesign) using an Applied Biosystems 7500

realtime PCR instrument (10 min at 55°C, 8 min at 95°C , 40× 10 sec at 95°C, and 60 sec at

60°C, followed by meltcurve analysis 65°C–95°C in 0.5°C increments at 5 sec per step) in

triplicates for *Zfhx3* expression in control and mutant groups . 7500 Applied Biosystems

Software v2.3 (https://www.thermofisher.com/uk/en/home/technical-resources/software-

downloads/applied-biosystems- 7500-real-time-pcr-system.html) was used to obtain the

relative quantification values and examine melt curves. All RT-qPCR data was normalized to

the *Rpl32* reference control gene and analysed using the standard curve method (Larionov et

al., 2005). The primers used for *Zfhx3* (CTGTGCCAAGACATGCTCAAC [forward] and

CAGCAGGGAGGAACATGCTA [reverse]) and *Rpl32* (AGGCACCAGTCAGACCGATA

[forward] and TGTTGGGCATCAGGATCTGG [reverse]) were designed and screened for

target specificity with the National Center for Biotechnology Information's (NCBI) Basic

Local Alignment Search Tool (BLAST) (https://blast.ncbi.nlm.nih.gov/). The primer sets

were validated before use (requirements: no predicted off-targets, primer efficiency 85-120%,

no significant signal in nontemplate control, single peak in melt curve, and single band at the

predicted size when separated via agarose gel electrophoresis). ZFHX3 gene expression

differences from qPCR were plotted and analyzed with GraphPad Prism v9.5.1 for Windows,

GraphPad Software (www.graphpad.com). Unpaired-two tailed t-test between control and

mutant group across the tested time points was set at alpha= 0.05.

Gene annotation

The gene list derived from Bioconductor-based package ChIPseeker v1.28.3 (Yu et al., 2015)

was fed into the Database for Annotation, Visualization, and Integrated Discovery (DAVID)

tool (Dennis et al., 2003). The functional annotation chart based on KEGG pathway, Gene

Ontology (GO): biological processes (BP) was plotted with the help of the ggplot2 package in

Rv4.0.5 (https://ggplot2.tidyverse.org/).

HOMER motif analysis

The findMotifsGenome.pl function within the HOMER v4.11 package (Heinz et al., 2010)

was used to identify enriched motifs and their corresponding TFs with options size 200 –len

8, 10, 12 –mask –preparse –dumpfasta, with default background regions.

TimeTeller modelling of the SCN clock

TimeTeller (TT) analysis was conducted using all 35 available SCN time-course RNA-Seq

samples from control (Cre-neg and -Tamoxifen) mice to train the model (Vlachou et al.,

2024). Data from +Tamoxifen SCN were then projected into the model as test samples.

Based on the available training data, we selected the 10 most rhythmic and consistent genes

to build the TimeTeller model, in this case: *Bmal1, Per2, Cry1, Tef, Hlf, Dbp, Nr1d2, Npas2,*

Ciart, and *Bhlhe41.* As a first step to cross-validate the model, we used a "leave-one-out"

approach. Iteratively, 5-6 randomly chosen samples from all Cre-neg and -Tamoxifen SCN

were left out from the training data to build the TT model, and the left out samples were

treated as test samples to determine the TT prediction for all Cre-neg and -Tamoxifen SCN.

In addition, we used published SCN RNA-Seq data to independently validate the full model

with all 35 control SCN from this study (Deota et al., 2023). TT analysis predicts phase and

clock functionality (theta) based on the averaged likelihood for the projected test sample. For

the SCN model, we used time-course normalization with log threshold selected at -6. Further,

to compare TT phase predictions between genotypes, we used estimation statistics (Ho et al.,

2019) and p-values were generated using a permutation testing with 5000 bootstrap samples.

FUNDING

685 PMN, GB and AB were supported by Medical Research Council (MC_U142684173). MHH

686 was supported by Medical Research Council (MC_U105170643). RD was supported by

687 Cancer Research UK and EPSRC (C53561/A19933), and VV was funded by the UK Medical

688 Research Council Doctoral Training Partnership (MR/N014294/1).

AUTHOR CONTRIBUTION

AB, GB and PMN conceived the study. AB and GB were involved in sample collection and

processing. AB and PMN carried out the investigation with AB conducting the formal data

analysis. Data visualization was performed by AB, GB, MHH and PMN. RD and VV carried

the data modelling and TT prediction. Supervision of the research was carried out by MHH

and PMN. The original draft was written by AB. Manuscript was reviewed, edited and

finalised by AB, GB, RD, VV, MHH and PMN.

COMPETING INTERESTS

Authors declare that they have no competing interests.

ACKNOWLEDGEMENTS

We thank the staff of the Mary Lyon Centre and core services at MRC Harwell Institute for

assistance with mouse studies. We thank Dr Benjamin Weger, University of Queensland for

his valuable suggestions and timely inputs during data analysis using dryR. We also thank

Richard Reeves for data accessibility on UCSC genome browser.

DATA AVAILIBILITY

All raw and processed sequencing data generated in this study have been submitted to the

NCBI Gene Expression Omnibus

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE261795) under accession number

GSE261795 .

REFERENCES

Abe, Y. O., Yoshitane, H., Kim, D. W., Kawakami, S., Koebis, M., Nakao, K., . . . Fukada, Y. (2022).
Rhythmic transcription of Bmal1 stabilizes the circadian timekeeping system in mammals.
Nat Commun, 13(1), 4652. https://doi.org/10.1038/s41467-022-32326-9
Araki, R., Takahashi, H., Fukumura, R., Sun, F., Umeda, N., Sujino, M., . . . Abe, M. (2004). Restricted
expression and photic induction of a novel mouse regulatory factor X4 transcript in the
suprachiasmatic nucleus. *J Biol Chem, 279*(11), 10237-10242.
https://doi.org/10.1074/jbc.M312761200
Arima, H., House, S. B., Gainer, H., & Aguilera, G. (2002). Neuronal activity is required for the
circadian rhythm of vasopressin gene transcription in the suprachiasmatic nucleus in vitro.
Endocrinology, 143(11), 4165-4171. https://doi.org/10.1210/en.2002-220393

Bafna, A., Banks, G., Hastings, M. H., & Nolan, P. M. (2023). Dynamic modulation of genomic
enhancer elements in the suprachiasmatic nucleus, the site of the mammalian circadian
clock. *Genome Res, 33*(5), 673-688. https://doi.org/10.1101/gr.277581.122
Bafna, A., Lau, P., Banks, G., & Nolan, P. M. (2023). Harvesting mouse suprachiasmatic nucleus by
vibrating microtome for diurnal transcriptome analysis. *STAR Protoc, 4*(4), 102618.
https://doi.org/10.1016/j.xpro.2023.102618
Banks, G. T., & Nolan, P. M. (2011). Assessment of Circadian and Light-Entrainable Parameters in
Mice Using Wheel-Running Activity. *Curr Protoc Mouse Biol, 1*(3), 369-381.
https://doi.org/10.1002/9780470942390.mo110123
Beacon, T. H., Delcuve, G. P., López, C., Nardocci, G., Kovalchuk, I., van Wijnen, A. J., & Davie, J. R.
(2021). The dynamic broad epigenetic (H3K4me3, H3K27ac) domain as a mark of essential
genes. *Clin Epigenetics, 13*(1), 138. https://doi.org/10.1186/s13148-021-01126-1
Bedont, J. L., LeGates, T. A., Buhr, E., Bathini, A., Ling, J. P., Bell, B., . . . Blackshaw, S. (2017). An LHX1-
Regulated Transcriptional Network Controls Sleep/Wake Coupling and Thermal Resistance of
the Central Circadian Clockworks. *Curr Biol, 27*(1), 128-136.
https://doi.org/10.1016/j.cub.2016.11.008
Brown, T. M., Hughes, A. T., & Piggins, H. D. (2005). Gastrin-releasing peptide promotes
suprachiasmatic nuclei cellular rhythmicity in the absence of vasoactive intestinal
polypeptide-VPAC2 receptor signaling. *J Neurosci, 25*(48), 11155-11164.
https://doi.org/10.1523/JNEUROSCI.3821-05.2005
Calo, E., & Wysocka, J. (2013). Modification of enhancer chromatin: what, how, and why? *Mol Cell,*
49(5), 825-837. https://doi.org/10.1016/j.molcel.2013.01.038
Del Rocío Pérez Baca, M., Jacobs, E. Z., Vantomme, L., Leblanc, P., Bogaert, E., Dheedene, A., . . .
consortium, Z. (2023). A novel neurodevelopmental syndrome caused by loss-of-function of
the Zinc Finger Homeobox 3 (ZFHX3) gene. *medRxiv*.
https://doi.org/10.1101/2023.05.22.23289895
Dennis, G., Sherman, B. T., Hosack, D. A., Yang, J., Gao, W., Lane, H. C., & Lempicki, R. A. (2003).
DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol, 4*(5),
P3.
Deota, S., Lin, T., Chaix, A., Williams, A., Le, H., Calligaro, H., . . . Panda, S. (2023). Diurnal
transcriptome landscape of a multi-tissue response to time-restricted feeding in mammals.
Cell Metab, 35(1), 150-165.e154. https://doi.org/10.1016/j.cmet.2022.12.006
Figueroa, K. P., Gross, C., Buena-Atienza, E., Paul, S., Gandelman, M., Kakar, N., . . . Pulst, S. M.
(2024). A GGC-repeat expansion in ZFHX3 encoding polyglycine causes spinocerebellar ataxia
type 4 and impairs autophagy. *Nat Genet*. https://doi.org/10.1038/s41588-024-01719-5
Fu, C., An, N., Liu, J., A, J., Zhang, B., Liu, M., . . . Dong, J. T. (2020). The transcription factor ZFHX3 is
crucial for the angiogenic function of hypoxia-inducible factor 1α in liver cancer cells. *J Biol*
Chem, 295(20), 7060-7074. https://doi.org/10.1074/jbc.RA119.012131
Hastings, M. H., Maywood, E. S., & Brancaccio, M. (2018). Generation of circadian rhythms in the
suprachiasmatic nucleus. *Nat Rev Neurosci, 19*(8), 453-469. https://doi.org/10.1038/s41583-
018-0026-z
Hastings, M. H., Maywood, E. S., & Brancaccio, M. (2019). The Mammalian Circadian Timing System
and the Suprachiasmatic Nucleus as Its Pacemaker. *Biology (Basel), 8*(1).
https://doi.org/10.3390/biology8010013
Hatori, M., Gill, S., Mure, L. S., Goulding, M., O'Leary, D. D., & Panda, S. (2014). Lhx1 maintains
synchrony among circadian oscillator neurons of the SCN. *Elife, 3*, e03357.
https://doi.org/10.7554/eLife.03357
Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., . . . Glass, C. K. (2010). Simple
combinations of lineage-determining transcription factors prime cis-regulatory elements
required for macrophage and B cell identities. *Mol Cell, 38*(4), 576-589.
https://doi.org/10.1016/j.molcel.2010.05.004

Herzog, E. D., Hermanstyne, T., Smyllie, N. J., & Hastings, M. H. (2017). Regulating the Suprachiasmatic Nucleus (SCN) Circadian Clockwork: Interplay between Cell-Autonomous and Circuit-Level Mechanisms. *Cold Spring Harb Perspect Biol*, *9*(1). https://doi.org/10.1101/cshperspect.a027706

Ho, J., Tumkaya, T., Aryal, S., Choi, H., & Claridge-Chang, A. (2019). Moving beyond P values: data analysis with estimation graphics. *Nat Methods*, *16*(7), 565-566. https://doi.org/10.1038/s41592-019-0470-3

Hogenesch, J. B., Panda, S., Kay, S., & Takahashi, J. S. (2003). Circadian transcriptional output in the SCN and liver of the mouse. *Novartis Found Symp*, *253*, 171-180; discussion 152-175, 102-179, 180-173 passim.

Holwerda, S. J., & de Laat, W. (2013). CTCF: the protein, the binding partners, the binding sites and their chromatin loops. *Philos Trans R Soc Lond B Biol Sci*, *368*(1620), 20120369. https://doi.org/10.1098/rstb.2012.0369

Hughes, M. E., Abruzzi, K. C., Allada, R., Anafi, R., Arpat, A. B., Asher, G., . . . Hogenesch, J. B. (2017). Guidelines for Genome-Scale Analysis of Biological Rhythms. *J Biol Rhythms*, *32*(5), 380-393. https://doi.org/10.1177/0748730417728663

Hughes, S., Edwards, J. K., Wilcox, A. G., Pothecary, C. A., Barnard, A. R., Joynson, R., . . . Nolan, P. M. (2021). Zfhx3 modulates retinal sensitivity and circadian responses to light. *FASEB J*, *35*(9), e21802. https://doi.org/10.1096/fj.202100563R

Jagannath, A., Butler, R., Godinho, S. I. H., Couch, Y., Brown, L. A., Vasudevan, S. R., . . . Peirson, S. N. (2013). The CRTC1-SIK1 pathway regulates entrainment of the circadian clock. *Cell*, *154*(5), 1100-1111. https://doi.org/10.1016/j.cell.2013.08.004

Jameson, H. S., Hanley, A., Hill, M. C., Xiao, L., Ye, J., Bapat, A., . . . Ellinor, P. T. (2023). Loss of the Atrial Fibrillation-Related Gene, . *Circ Res*, *133*(4), 313-329. https://doi.org/10.1161/CIRCRESAHA.123.323029

Kim, S., Yu, N. K., & Kaang, B. K. (2015). CTCF as a multifunctional protein in genome regulation and gene expression. *Exp Mol Med*, *47*(6), e166. https://doi.org/10.1038/emm.2015.33

Kuhlman, S. J., Silver, R., Le Sauter, J., Bult-Ito, A., & McMahon, D. G. (2003). Phase resetting light pulses induce Per1 and persistent spike activity in a subpopulation of biological clock neurons. *J Neurosci*, *23*(4), 1441-1450. https://doi.org/10.1523/JNEUROSCI.23-04-01441.2003

Larionov, A., Krause, A., & Miller, W. (2005). A standard curve based method for relative real time PCR data processing. *BMC Bioinformatics*, *6*, 62. https://doi.org/10.1186/1471-2105-6-62

Liu, A. C., Tran, H. G., Zhang, E. E., Priest, A. A., Welsh, D. K., & Kay, S. A. (2008). Redundant function of REV-ERBalpha and beta and non-essential role for Bmal1 cycling in transcriptional regulation of intracellular circadian rhythms. *PLoS Genet*, *4*(2), e1000023. https://doi.org/10.1371/journal.pgen.1000023

Lopez-Molina, L., Conquet, F., Dubois-Dauphin, M., & Schibler, U. (1997). The DBP gene is expressed according to a circadian rhythm in the suprachiasmatic nucleus and influences circadian behavior. *EMBO J*, *16*(22), 6762-6771. https://doi.org/10.1093/emboj/16.22.6762

Mach, P., Kos, P. I., Zhan, Y., Cramard, J., Gaudin, S., Tünnermann, J., . . . Giorgetti, L. (2022). Cohesin and CTCF control the dynamics of chromosome folding. *Nat Genet*, *54*(12), 1907-1918. https://doi.org/10.1038/s41588-022-01232-7

Matsu-Ura, T., Moore, S. R., & Hong, C. I. (2018). WNT Takes Two to Tango: Molecular Links between the Circadian Clock and the Cell Cycle in Adult Stem Cells. *J Biol Rhythms*, *33*(1), 5-14. https://doi.org/10.1177/0748730417745913

Maywood, E. S., Chesham, J. E., Winsky-Sommerer, R., & Hastings, M. H. (2021). Restoring the Molecular Clockwork within the Suprachiasmatic Hypothalamus of an Otherwise Clockless Mouse Enables Circadian Phasing and Stabilization of Sleep-Wake Cycles and Reverses Memory Deficits. *J Neurosci*, *41*(41), 8562-8576. https://doi.org/10.1523/JNEUROSCI.3141-20.2021

823 Mirsky, H. P., Liu, A. C., Welsh, D. K., Kay, S. A., & Doyle, F. J. (2009). A model of the cell-autonomous
824   mammalian circadian clock. *Proc Natl Acad Sci U S A*, *106*(27), 11107-11112.
825   https://doi.org/10.1073/pnas.0904837106
826 Morris, E. L., Patton, A. P., Chesham, J. E., Crisp, A., Adamson, A., & Hastings, M. H. (2021). Single-cell
827   transcriptomics of suprachiasmatic nuclei reveal a Prokineticin-driven circadian network.
828   *EMBO J*, *40*(20), e108614. https://doi.org/10.15252/embj.2021108614
829 Nolan, P. M., Banks, G., Bourbia, N., Wilcox, A. G., Bentley, L., Moir, L., . . . Dumbell, R. (2023). A
830   missense mutation in zinc finger homeobox-3 (ZFHX3) impedes growth and alters
831   metabolism and hypothalamic gene expression in mice. *FASEB J*, *37*(10), e23189.
832   https://doi.org/10.1096/fj.202201829R
833 Padlom, A., Ono, D., Hamashima, R., Furukawa, Y., Yoshimura, T., & Nishiwaki-Ohkawa, T. (2022).
834   Level of constitutively expressed BMAL1 affects the robustness of circadian oscillations. *Sci*
835   *Rep*, *12*(1), 19519. https://doi.org/10.1038/s41598-022-24188-4
836 Parsons, M. J., Brancaccio, M., Sethi, S., Maywood, E. S., Satija, R., Edwards, J. K., . . . Nolan, P. M.
837   (2015). The Regulatory Factor ZFHX3 Modifies Circadian Function in SCN via an AT Motif-
838   Driven Axis. *Cell*, *162*(3), 607-621. https://doi.org/10.1016/j.cell.2015.06.060
839 Pugacheva, E. M., Kubo, N., Loukinov, D., Tajmul, M., Kang, S., Kovalchuk, A. L., . . . Lobanenkov, V. V.
840   (2020). CTCF mediates chromatin looping via N-terminal domain-dependent cohesin
841   retention. *Proc Natl Acad Sci U S A*, *117*(4), 2020-2031.
842   https://doi.org/10.1073/pnas.1911708117
843 Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
844   features. *Bioinformatics*, *26*(6), 841-842. https://doi.org/10.1093/bioinformatics/btq033
845 Schibler, U., Gotic, I., Saini, C., Gos, P., Curie, T., Emmenegger, Y., . . . Franken, P. (2015). Clock-Talk:
846   Interactions between Central and Peripheral Circadian Oscillators in Mammals. *Cold Spring*
847   *Harb Symp Quant Biol*, *80*, 223-232. https://doi.org/10.1101/sqb.2015.80.027490
848 Sun, X., Fu, X., Li, J., Xing, C., Martin, D. W., Zhang, H. H., . . . Dong, J. T. (2012). Heterozygous
849   deletion of Atbf1 by the Cre-loxP system in mice causes preweaning mortality. *Genesis*,
850   *50*(11), 819-827. https://doi.org/10.1002/dvg.22041
851 Takahashi, J. S., Hong, H. K., Ko, C. H., & McDearmon, E. L. (2008). The genetics of mammalian
852   circadian order and disorder: implications for physiology and disease. *Nat Rev Genet*, *9*(10),
853   764-775. https://doi.org/10.1038/nrg2430
854 VanDunk, C., Hunter, L. A., & Gray, P. A. (2011). Development, maturation, and necessity of
855   transcription factors in the mouse suprachiasmatic nucleus. *J Neurosci*, *31*(17), 6457-6467.
856   https://doi.org/10.1523/JNEUROSCI.5385-10.2011
857 Vlachou, D., Veretennikova, M., Usselmann, L., Vasilyev, V., Ott, S., Bjarnason, G. A., . . . Rand, D. A.
858   (2024). TimeTeller: A tool to probe the circadian clock as a multigene dynamical system.
859   *PLoS Comput Biol*, *20*(2), e1011779. https://doi.org/10.1371/journal.pcbi.1011779
860 Wallenius, J., Kafantari, E., Jhaveri, E., Gorcenco, S., Ameur, A., Karremo, C., . . . Puschmann, A.
861   (2024). Exonic trinucleotide repeat expansions in ZFHX3 cause spinocerebellar ataxia type 4:
862   A poly-glycine disease. *Am J Hum Genet*, *111*(1), 82-95.
863   https://doi.org/10.1016/j.ajhg.2023.11.008
864 Weger, B. D., Gobet, C., David, F. P. A., Atger, F., Martin, E., Phillips, N. E., . . . Gachon, F. (2021).
865   Systematic analysis of differential rhythmic liver gene expression mediated by the circadian
866   clock and feeding rhythms. *Proc Natl Acad Sci U S A*, *118*(3).
867   https://doi.org/10.1073/pnas.2015803118
868 Wen, S., Ma, D., Zhao, M., Xie, L., Wu, Q., Gou, L., . . . Yan, J. (2020). Spatiotemporal single-cell
869   analysis of gene expression in the mouse suprachiasmatic nucleus. *Nat Neurosci*, *23*(3), 456-
870   467. https://doi.org/10.1038/s41593-020-0586-x
871 Wilcox, A. G., Bains, R. S., Williams, D., Joynson, E., Vizor, L., Oliver, P. L., . . . Nolan, P. M. (2021).
872   Zfhx3-mediated genetic ablation of the SCN abolishes light entrainable circadian activity

while sparing food anticipatory activity. *iScience, 24*(10), 103142.

https://doi.org/10.1016/j.isci.2021.103142

Wilcox, A. G., Vizor, L., Parsons, M. J., Banks, G., & Nolan, P. M. (2017). Inducible Knockout of Mouse

Zfhx3 Emphasizes Its Key Role in Setting the Pace and Amplitude of the Adult Circadian

Clock. *J Biol Rhythms, 32*(5), 433-443. https://doi.org/10.1177/0748730417722631

Won, K. J., Chepelev, I., Ren, B., & Wang, W. (2008). Prediction of regulatory elements in mammalian

genomes using chromatin signatures. *BMC Bioinformatics, 9*, 547.

https://doi.org/10.1186/1471-2105-9-547

Yu, G., Wang, L. G., & He, Q. Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP peak

annotation, comparison and visualization. *Bioinformatics, 31*(14), 2382-2383.

https://doi.org/10.1093/bioinformatics/btv145

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., . . . Liu, X. S. (2008).

Model-based analysis of ChIP-Seq (MACS). *Genome Biol, 9*(9), R137.

https://doi.org/10.1186/gb-2008-9-9-r137

Figure 1

A

Overlap of ZFHX3 binding with histone modifications

Category	Number of sites
Only H3K27ac	47552
Only H3K4me3	35673
Only ZFHX3	9512
H3K4me3 + ZFHX3	8992
H3K27ac + ZFHX3	3570
H3K27ac + H3K4me3	36819
H3K27ac + H3K4me3 + ZFHX3	21162

B H3K27ac+H3K4me3+ZFHX3

- Promoter (<=1kb) (70.69%)
- Promoter (1−2kb) (5.6%)
- Promoter (2−3kb) (2.26%)
- 5' UTR (0.23%)
- 3' UTR (0.53%)
- 1st Exon (0.01%)
- Other Exon (1.58%)
- 1st Intron (2.62%)
- Other Intron (4.52%)
- Downstream (<=300) (0.08%)
- Distal Intergenic (11.88%)

chr1:91,405,146-91,470,160 20 kb mm10
Per2
H3K4me3
H3K27ac
ZT3 ZFHX3
ZT15 ZFHX3

C H3K4me3+ZFHX3

- Promoter (<=1kb) (59.84%)
- Promoter (1−2kb) (9.21%)
- Promoter (2−3kb) (3.49%)
- 5' UTR (0.13%)
- 3' UTR (1.03%)
- 1st Exon (0.03%)
- Other Exon (3.06%)
- 1st Intron (2.16%)
- Other Intron (4.05%)
- Downstream (<=300) (0.12%)
- Distal Intergenic (16.87%)

chr3:103,560,990-103,622,359 20 kb mm10
Syt16
H3K4me3
H3K27ac
ZT3 ZFHX3
ZT15 ZFHX3

D H3K27ac+ZFHX3

- Promoter (<=1kb) (7.2%)
- Promoter (1−2kb) (3.73%)
- Promoter (2−3kb) (2.49%)
- 5' UTR (0.14%)
- 3' UTR (2.02%)
- Other Exon (3.73%)
- 1st Intron (10.7%)
- Other Intron (32.84%)
- Downstream (<=300) (0.22%)
- Distal Intergenic (36.93%)

chr12:109,385,255-109,398,349 5 kb mm10
H3K4me3
H3K27ac
ZT3 ZFHX3
ZT15 ZFHX3

E ZFHX3 ONLY

- Promoter (<=1kb) (13.37%)
- Promoter (1−2kb) (7.19%)
- Promoter (2−3kb) (5.17%)
- 5' UTR (0.2%)
- 3' UTR (2.63%)
- 1st Exon (0.03%)
- Other Exon (6.34%)
- 1st Intron (6.18%)
- Other Intron (19.52%)
- Downstream (<=300) (0.32%)
- Distal Intergenic (39.04%)

chr2:103,547,126-103,547,732
200 bases mm10
H3K4me3
H3K27ac
ZT3 ZFHX3
ZT15 ZFHX3

Figure 1. Genomic occupancy of ZFHX3 in the SCN. (A) left- Venn-diagram illustrating overlap of ZFHX3 peaks with histone marks, H3K4me3 and H3K27ac, right- table showing number of sites co-occupied by ZFHX3 and histone marks. (B,C,D and E) left- Genomic feature distribution of ZFHX3 peaks with histone marks (ChIPSeeker), right- UCSC genome browser tracks showing histone modifications H3K4me3(pink), H3K27ac (blue) and ZFHX3 (ZT3 = dark grey and ZT15 = light grey) normalized ChIP-seq read coverage at representative examples for each category. The chromosome location and scale (mm10 genome) indicated at the top.

Figure 2. Circadian wheel running analysis of control and mutant animals. A. Double plotted actograms showing wheel running activity of -Tamoxifen and +Tamoxifen animals through 32 days of constant darkness and 24 days under a 12h light/dark cycle. Regions in grey denote periods in darkness. B. +Tamoxifen animals show a reduced circadian period C: Graph showing the distribution of activity in -Tamoxifen and +Tamoxifen animals over 24 hours in a 12h light/dark cycle. Data is taken from the final seven days the animals were in the light/dark cycle shown in A. Grey denotes the dark period of the cycle.

Figure 3

Figure 3. Effect of ZFHX3-KO on SCN transcriptome. (A) Mean-difference plot (MD-plot) showing downregulated (red) and upregulated (blue) gene expression after the loss of ZFHX3. (B) Heatmap showing normalized expression of top 25 (by adj pvalue) differential genes for the compared groups. (C) Stripcharts of differentially expressed genes (*Dpy19l13, Grp, Vip*). (D) Functional annotation of downregulated (red) and upregulated (blue) genes after ZFHX3-KO using the GO::BP (biological processes) terms by DAVID.

Figure 4

Figure 4. Effect of ZFHX3-KO on rhythmic gene expression . (A) Left- Heatmap showing loss of rhythmic gene expression after *Zfhx3-KO* (module 2) as computed by dryR statistical framework , Right- Illustrative examples of daily abundance of module 2 genes, *Bmal1 and Avp,* in Cre-neg and +Tamoxifen conditions. (B) Left- Heatmap showing no effect on rhythmic gene expression after *Zfhx3-KO* (module 4) , Right- Illustrative examples of module 4 genes, *Cry1 and Cry2,* in Cre-neg and +Tamoxifen conditions.

Figure 5

Figure 5. Effect of ZFHX3-KO on clock gene expression clustered in module 5 . (A) Heatmap showing change in rhythmic gene expression after *Zfhx3-KO* (module 5) as computed by dryR statistical framework , (B) 24-h radial plots showing effect on phase (peak expression) of rhythmic genes in Cre-neg (WT) vs. +Tamoxifen (ZFHX3-KO) conditions (C)Illustrative examples of daily abundance of module 5 genes, showing change in phase and amplitude in ZFHX3-KO condition (blue) as compared to control (pink). The y-axis values represents normalised expression in the SCN.

Figure 6

Figure 6. TimeTeller model SCN circadian clock in control and ZFHX3-KO mice. (A) TimeTeller (TT) model build with the data of all 35 available SCN from Creneg and -Tam mice. Only the knots of the model at the collected ZTs, and predictions for test data of +Tam SCN according to when they were collected are shown. (B) Quantitation of SCN phases as difference between ZT to TT prediction for all genotypes of this study and previously published wild-type SCNs from ad libitum or night fed mice (Panda_ALF and Panda_TRF). Only the +Tam group is different from Creng & -Tam control training data (p=0.0028).

Supplementary Figure 1

Supplementary figure 1. Gene ontology and motif analysis of ZFHX3 bound sites. (A) left- Functional annotation of nearest neighbouring gene (TSS) to active promoter sites occupied by ZFHX3 and histone marks (H3K4me3 and H3K27ac) using KEGG pathway (DAVID), right-ciradian rhythm pathway genes that showed co-occupancy of ZFHX3 and histone marks at TSS illustrated using KEGG pathway . (B ,C) Functional annotation of nearest neighbouring gene (TSS) to sites occupied by ZFHX3 + H3K4me3 (light green) and ZFHX3 alone (dark green) (D) Over-represented transcription factor binding motifs at ZFHX3+H3K4me3+H3K27ac sites (shaded blue), ZFHX3+H3K27ac sites (shaded green) and ZFHX3+ H3K4me3 (shaded red)by HOMER.

Supplementary Figure 2. +Tamoxifen animals show changes in circadian wheel running parameters in conditions of constant darkness.
Compared to -Tamoxifen animals, +Tamoxifen show a significantly (A) reduced *Zfhx3* gene expression (p<0.02, unpaired t-test) (B) reduced circadian amplitude, (C) increased Intradaily variability (IV) , (D) reduced Interdaily stability(IS) and (E) advanced activity onset in LD.

Supplementary Figure 3. Effect of *Zfhx3*-KO on SCN transcriptome. (A) Heatmap showing module 1 genes (n = 16832) that show no rhythmic expression before and after *Zfhx3*-KO as computed by dryR statistical framework (B)Heatmap showing module 3 genes (n = 403) that showed rhythmic expression after *Zfhx3*-KO.

Supplementary Figure 4

Supplementary Figure 4. Functional annotation of rhythmic genes in the SCN. (A) Top 20 enriched functional pathways from genes that lost rhythmic transcription after gene deletion (module 2) and (B) genes that showed change in phase and/or amplitude (module 5) using the GO::BP (biological processes) terms by DAVID.

Supplementary Figure 5

A

B

Supplementary Figure 5. SCN circadian clock using TimeTeller. (A) (Left) Choosing genes for the TimeTeller model. Looking for good rhythmicity p-val and high R squared in population cosinor analysis leads to selection of a limited panel of well-behaved clock genes (named), i.e., clock genes with adjusted pval < 0.10 and R squared > 0.80.(Right) Results of the cross-validation of the TT model using the indicated genes in the left panel. (B) TimeTeller theta parameter is indistinguishable between the genotypes. Theta is a measure of certainty for phase prediction in TimeTeller, i.e., how closely the test samples can be projected into the training model.

Supplementary Figure 6. Effect of ZFHX3-KO on circadian clock . The effect of loss of ZFHX3 on the primary and ancillary TTFL in the SCN. Blue and red arrow depicts decreased/increased average gene expression in *Zfhx3* deficient mice as compared to the control, respectively. The waveform symbol next to the gene name depicts detected 24-h rhythmic expression in our study. The curly arrows (right) of the gene illustrates the change in phase of the rhythmic expression.

Lost rhythm

Avp Syt17 Wdr1
Bmal1 Prok2

Intracellular transport
Protein import to nucleus
mTOR signalling pathway
Spinocerebellar ataxia
Rhythmic process

49% ZFHX3 sites localized with modified histones

Clock genes

CCGs

Gained rhythm

Hdac3 Sox4 Aldh1
Notch1 Galnt4

RNA processing
Notch Signalling
Angiogenesis
Metabolic pathways
Cell proliferation

Retained rhythm

Dlk1 Cry1
Dusp16 Hlf Rfx5

Cell cycle
DNA repair
ER stress
MAPK Signalling
Protein phosphorylation

36% SCN genes affected by loss of ZFHX3

Clock genes

CCGs

Altered rhythm

Per2 Dbp Creb3l1
Hif3a Rgs16

Circadian rhythm
Ppar signalling
Response to hypoxia
Butanoate metabolism
Circadian gene regulation