Screening the MMV Pathogen Box reveals the mitochondrial *bc*$_1$-
complex as a drug target in mature *Toxoplasma gondii* bradyzoites.

Deborah Maus[1], Elyzana Putrianti[1], Tobias Hoffmann[3], Michael Laue[3], Frank Seeber[2], Martin
Blume[1*]

[1]P 6: Metabolism of Microbial Pathogens, Robert Koch Institute, 13353, Berlin, Germany

[2]FG 16: Mycotic and Parasitic Agents and Mycobacteria, Robert Koch Institute, 13353, Berlin,
Germany

[3]ZBS 4: Advanced Light and Electron Microscopy, Centre for Biological Threats and Special
Pathogens 4, Robert Koch-Institute, 13353, Berlin, Germany.

*corresponding author

Martin Blume
Project Group 6
Robert Koch Institute
Seestr. 10
13353 Berlin
Germany
Email: blumem@rki.de

Abstract

The apicomplexan parasite *Toxoplasma gondii* infects 25-30% of the global human population
and can cause life-threatening diseases in immunocompromised patients. The chronically
infectious forms of the parasite, bradyzoites, persist within cysts in brain and muscle tissue
and are responsible for its transmission and remission of the disease. Currently available
treatment options are very limited and are only effective against the fast-replicating tachyzoites
but fail to eradicate the chronic stages of *T. gondii*. The cause of these treatment failures
remains unclear. Here, we utilized our recently developed human myotube-based culture
model to screen compounds from the MMV Pathogen Box against pan-resistant in vitro
bradyzoites and identified multiple compounds with simultaneous activity against tachyzoites
and bradyzoites. Stable isotope-resolved metabolic profiling on tachyzoites and bradyzoites
identified the mitochondrial *bc*$_1$-complex as a target of bradyzocidal compounds and defines
their metabolic impacts on both parasite forms. Our data suggest that mature bradyzoites rely
on mitochondrial ATP production.

Key Words

36 Apicomplexa, in vitro tissue cysts, *bc*$_1$-complex, metabolomics, mitochondrial metabolism

Introduction

Toxoplasma gondii is an ubiquitous apicomplexan parasite that infects virtually all warm-blooded animals, including 25-30 % of humans (Montazeri et al. 2020). Of all food-borne pathogens, *T. gondii* causes 20% of the total disease burden in Europe (Havelaar et al. 2015). Although most acute infections caused by the fast replicating tachyzoite stage of the parasite remain asymptomatic in immunocompetent individuals, the parasite persists life-long in the form of encysted semi-dormant bradyzoites (Weiss and Dubey 2009). Currently, no medical treatment exists to eradicate these tissue cysts with their rigid cyst wall from infected individuals. *T. gondii* is transmitted through undercooked meat products of infected livestock but also via feline-shed oocysts that can contaminate water sources and soil (Weiss and Dubey 2009). Severe toxoplasmosis occurs after reactivation of cysts or primary infection of immunosuppressed individuals or fetuses of seronegative mothers and can lead to systemic infections, uveitis, encephalitis as well as congenital transmission with potentially fatal outcomes (Montoya and Liesenfeld 2004). Hence, tissue cysts of *T. gondii* constitute a long-term source for transmission and risk of acute disease. Also, other related cyst-forming coccidian parasites such as *Neospora caninum* and *Besnoitia besnoiti* pose an immense veterinary burden by causing abortion mainly in sheep and severe chronic infections in cattle and horses, respectively. Current treatment options exist only against the tachyzoite form of these parasites and include antifolates, such as pyrimethamine or trimethoprim, that inhibit the bifunctional dihydrofolate reductase-thymidylate synthetase, in combination with sulfadiazine, which blocks the dihydropteroate synthetase. These enzymes are implicated in DNA synthesis in both the parasite and side-effects often require pre-mature termination of the treatment (Katlama, De Wit, et al. 1996; Alday and Doggett 2017). Another potent drug target in coccidians and other apicomplexan parasites such as *Plasmodium* spp. is the cytochrome bc_1-complex in the mitochondrial respiratory chain, which is inhibited by a number of approved drugs, such as the naphthoquinones atovaquone (ATQ) (Meneceur et al. 2008; Nixon et al. 2013) and buparvaquone (BPQ) (Sharifiyazdi et al. 2012). However, both coenzyme Q analogs including bioavailable nanosuspensions (Shubar et al. 2011) fail to eradicate encysted bradyzoites of *T. gondii* and *N. caninum* from the brains of infected mice after multiple weeks of treatment (Doggett et al. 2012; Müller et al. 2015; Doggett et al. 2020). Experimental inhibitors of the bc_1-complex includes 1-hydroxy-2-dodecyl-4(1H)quinolone (HDQ) that arrests the growth of asexual blood stages of the related malaria parasite *P. falciparum* (Vallières et al. 2012). However, in *T. gondii* HDQ appears to inhibit multiple mitochondrial election donors such as the alternative NAD-dehydrogenase and the dihydroorotate dehydrogenase in tachyzoites (S. S. Lin, Gross, and Bohne 2009). It does not affect mature bradyzoites in vitro (Christiansen et al. 2022), and HDQ derivatives do not clear *T. gondii* cysts from the brain of infected mice (Bajohr et al. 2010). In addition, endochin-like quinolones constitute a promising series of potent bc_1-complex inhibitors that are active against *T. gondii* (McConnell et al. 2018) and *Plasmodium* parasites (Song et al. 2018). However, these compounds also do not fully eradicate the cysts *in vivo* (Doggett et al. 2012, 2020). Similarly, bumped kinase inhibitors that selectively inhibit parasite calcium-dependent protein kinases in various apicomplexans fail to clear tissue cysts *in vitro* (Christiansen et al. 2022) and *in vivo* (Hulverson et al. 2019). Despite these efforts, it remains unclear whether the target proteins of these drugs are rendered dispensable for the survival of bradyzoites or whether the eradication failures have other reasons, such as poor permeability across the blood brain barrier and/or the cyst wall. To date, the genetic tools available to directly characterize the function of essential genes in bradyzoites are limited to promoter replacements (Smith et al. 2021), while no genetic methods exist to

directly interrogate the fitness contribution of mitochondrially encoded genes such as the
cytochrome b subunit of the cytochrome bc_1-complex (J.-L. Wang et al. 2016). Consequently,
reverse pharmacological approaches, that involve screening of inhibitors against target
proteins, remain difficult and the underlying reasons for limited *in vivo* efficacies remain
uncertain.

Here we aim to address these shortcomings and identify essential processes in bradyzoite by
a combination of a phenotypic screen and subsequent metabolomics-driven identification of
the modes of actions of hit compounds. We employed a recently developed human myotube-
based culture system to generate mature *T. gondii* drug-tolerant bradyzoites (Christiansen et
al. 2022; Maus et al. 2024) to screen the MMV Pathogen Box. These *in vitro* cysts exhibit
resistance against antifolates, bumped kinase inhibitors and HDQ. They also express
hallmarks of *in vivo* cysts including oral infectivity and a cyst wall (Christiansen et al. 2022).
The Medicines for Malaria Venture (MMV) Pathogen Box provides a curated set of bioactive
molecules with activity against a number of pathogens including *P. falciparum*,
Cryptosporidium parvum, *Schistosoma spp.* and diseases caused by kinetoplastids (Veale
2019). Untargeted and stable isotope-resolved metabolomic analyses of active drugs identified
established and unsuspected modes of action in tachyzoites and bradyzoites, respectively.
Our data directly show metabolic consequences of bc_1-complex inhibition in bradyzoites and
suggest its crucial role in ATP production.

Material and Methods

Tachyzoite Culture
Prugeniaud Δhxgprt tdTomato (John et al. 2009) (PruTom) and RH-Rep1/2-S9(33-159)-GFP
(DeRocher et al. 2000; Thomsen-Zieger, Schachtner, and Seeber 2003) (RH-S9) parasites
were maintained in human foreskin fibroblasts (HFF) cells (BJ-5ta human foreskin fibroblasts
ATCC, CRL-4001) with Dulbecco's modified Eagle's medium (DMEM) containing 25 mM
glucose, 4 mM L-glutamine, 1 mM sodium pyruvate, 100 U/mL penicillin, 100 µg/mL
streptomycin and 1 % heat-inactivated fetal bovine serum (FBS, Gibco) at 37°C with 10 % CO_2
(Christiansen et al. 2022).

In vitro T. gondii **tissue cyst culture**
The *T. gondii in vitro* cyst culture was performed as in (Christiansen et al. 2022; Maus et al.
2024). In short, KD3 myoblasts (Shiomi et al. 2011) were grown to 70 % confluency, followed
by the induction of myotube differentiation by changing the medium to DMEM as described
above but with 2% horse serum instead of FBS and additionally supplemented with 10 µg/mL
human insulin, 5 µg/mL human holo-transferrin and 1.7 ng/mL Na_2SeO_3. After five days, the
myotubes were infected with PruTom (MOI 0.1) in bicarbonate-free Roswell Park Memorial
Institute medium (RPMI) containing 5.5 mM glucose, 50 mM HEPES, 4 mM L-glutamine,
100 U/mL penicillin, 100 µg/mL streptomycin, 2 % horse serum, 10 µg/mL human Insulin,
5 µg/mL human holo-transferrin and 1.7 ng/mL Na_2SeO_3 at pH 7.2. Cultures were incubated
at 37 °C at ambient CO_2 for four weeks and the medium was changed every two to three days.
A comprehensive scheme of the procedure is shown in Fig. S1.

MMV Pathogen Box Screening Procedure
All Pathogen Box compounds were stored at -20°C (Veale 2019). DMSO-solved compounds were diluted in DMEM or RPMI media and stored at 4°C until use. Host cells were grown in transparent, flat-bottom 96-well plates and infected as described above to cultivate either tachyzoites or in vitro tissue cysts. One column per plate was left uninfected to estimate fluorescence background. Four hours after infection with tachyzoites or after four weeks of bradyzoite maturation the culture medium was replaced by medium containing either 0.1 % DMSO (negative control) or 10 μM of the respective Pathogen Box compound. During the seven-day treatment period the medium was renewed every two to three days. It was then changed to tachyzoite growth medium to induce re-differentiation of viable bradyzoites to tachyzoites, and then the culture was monitored over 28 days. Prior to each medium exchange, the cultures were monitored for compound crystallization. The fluorescence was measured in a fluorescence plate reader (Tecan Infinite M200 PRO) (excitation Λ = 554 nm, emission Λ = 589 nm). Fluorescence values passing a threshold of 2,000 after background subtraction indicated proliferating parasites and hence an ineffective treatment (Fig. 1).

Resazurin viability assay and cytotoxicity
After the growth assays were completed, the host cell monolayers of both HFF and KD3 myotubes were washed with PBS and incubated with 500 nM resazurin in Milli-Q water at 37 °C for 4 hours. Fluorescence was measured (excitation Λ = 540 nm, emission Λ = 580 nm) in a fluorescence plate reader and normalized to the average fluorescence values of the DMSO-treated controls, which were set at 100 %. During bradyzoite assays KD3 host cells were visually inspected after drug exposure for one week. After completion of the re-growth phase for four weeks the colorimetric resazurin was performed. These values were reported as lowest cytotoxicity values (Supplemental Fig 2D). During secondary tests of putatively dually active compounds resazurin assays were performed in a concentration dependent manner (Supplemental File S1, Supplemental Fig 2D).

The dose response of dually active compounds
For the measurement of LC50 and minimal lethal concentrations tachyzoite and bradyzoite cultures were treated with the MMV compounds which were pipetted in a serial dilution across the 96-well plate and treated for 7 days. Afterwards, the compound containing medium was removed and possible regrowth was monitored for another 7 days (tachyzoite assay) or 21 to 28 days (bradyzoite assay) in a fluorescence plate reader (Tecan Infinite M200 PRO) (excitation Λ = 554 nm, emission Λ = 589 nm). The relative fluorescence units were background-subtracted. Depending on the development of the DMSO control, a late timepoint when DMSO-treated cultures still continued exponential growth was chosen to calculate the half-maximal inhibitory concentration (IC_{50}) for tachyzoites. This time point was between day 4 and 7. Bradyzoite cultures were of lower MOI, and we calculated LC_{50} values between experiments from 7 to 28 days post treatment when exponential growth occurred in mock treated cultures (Fig. 2). The concentration data was log-transformed and a non-linear regression (curve fit with variable slope) was performed to determine the dose response (Prism 9, GraphPad).

In silico predictions

MMV provided the SMILES IDs for every compound, which were compiled with LightBBB (Shaker et al. 2020) to predict the blood-brain permeability. SwissADME (Daina, Michielin, and Zoete 2017) was used to predict the gastrointestinal absorption capability. The logP values (Fig. S2) were provided by MMV and remaining logP values were calculated on SwissADME (Daina, Michielin, and Zoete 2017) as consensus values.

Fluorescence stain of mitochondrial activity

This assay was performed on tachyzoites and bradyzoites. For the tachyzoite experiment, HFF cells were infected with RH-S9 parasites at a MOI 0.1 and treated for 24 h with 1 µM atovaquone, 25 nM buparvaquone or 5 µM MMV1028806, respectively, corresponding to five-fold of the respective IC_{50} values, or 0.1% DMSO as a solvent control. For the bradyzoites experiment, 4 weeks-old ME49 *in vitro* cysts were treated for 24h with 5 µM atovaquone, 500 nM buparvaquone or 5 µM MMV1028806, respectively, corresponding to the lowest lethal concentration, or 0.1 % DMSO as a solvent control.

The assay was repeated with additional control treatments (see Supplemental Fig S5). In this experiment, HFF cells were infected with RH-S9 tachyzoites and, 4h post-infection, incubated for 24h with 0.1% DMSO (solvent control), 1 µM atovaquone, 25 nM buparvaquone, 5µM MMV1028806 (MMV), 1 µM pyrimethamine (Pyri), 10 nM clindamycin, or 25 µM 6-diazo-5-oxonorleucine (DON). Concentrations reflected the five-fold IC_{50} values, except for DON, which does not inhibit growth.

Subsequent staining and imaging procedures were identical across all experiments. Cultures were then stained with 500 nM Mitotracker Deep Red FM (MT; Thermo Fisher) in DMEM for 40 min at 37 °C, followed by a 20 min de-staining with DMEM and final fixation with 1 % paraformaldehyde for 20 min at room temperature. The cyst wall was decorated with *Dolichos biflorus* Agglutinin (DBA) and stained with Streptavidin-Cy2. Mowiol containing DAPI (1:3,000) was used to mount the cells. To avoid a biased image acquisition, all parasite vacuoles or cysts identified in a specimen along a horizontal axis were imaged. Tachyzoite images were taken with a Zeiss Axio Imager Z1 Microscope equipped with the ApoTome 2 at identical exposure times (DAPI 200 ms, MT 500 ms, S9-GFP 200 ms, Brightfield 10 ms) using a Zeiss Plan APOCHROMAT 63x/1.4 oil objective lens. Bradyzoite images were acquired on a Leica Stellaris 8 laser scanning confocal microscope using a Leica HC PL APO CS2 63x/1.4 oil objective lens. The pinhole diameter was set to 1 AU. All detectors were operating in photon counting mode. The acquisition parameters were set as follows. DAPI: diode laser at $405nm_{exc}$ with 2.2%, $430\text{-}550nm_{em}$ with a HyD S-detector. Cy2: WLL at $468nm_{exc}$ with 8.5%, $473\text{-}591nm_{em}$ with a HyD S-detector. Mitotracker: WLL at $641nm_{exc}$ with 2.7%, $650\text{-}784nm_{em}$ with a HyD X-detector (4x line average). The zoom was set to 7x, images were acquired in frame-sequential mode with a unidirectional scan speed of 400 Hz, the pixel dwell time for all channels was 3.18µs. For z-Stacks, z-step size was set to 0.299 µm.

Using ImageJ (Schindelin et al. 2012), the intensity threshold of individual pixels of MT (190 to 255 for tachyzoites and 125 to 65535 for bradyzoites) and DAPI (140 to 255 for tachyzoites and 363 to 65535) channels was cut off and the signal intensity was calculated per parasitophorous vacuole or cyst.

Oxygen consumption assay

To assess mitochondrial oxygen consumption in *T. gondii* RHΔku80 tachyzoites, we employed the MitoXpress Xtra Oxygen Consumption Assay (Agilent Technologies, Cat# MX-200-4) following the manufacturer's high-sensitivity (HS) method with modifications. Parasites were freshly harvested upon egress and resuspended at a density of 1×10^7 tachyzoites per reaction in phenol red-free D1 medium lacking glucose but supplemented with glutamine to promote mitochondrial electron transport chain (mETC) activity via the GABA shunt (MacRae, Sheiner et al. 2012).

Parasites were incubated in medium with one of the following treatments immediately before measurement: 0.1% DMSO (vehicle control), 1 µM antimycin A (complex III inhibitor, positive control), 10 µM FCCP (uncoupler, commonly used to determine the maximum respiration capacity), 1 µM ATQ, 1 µM BPQ, or 1 µM MMV1028806. Untreated control wells contained 1×10^7 tachyzoites in 0.1% DMSO. Background fluorescence was determined from wells containing MitoXpress Xtra reagent but no parasites.

Each 100 µL reaction was supplemented with 10 µL of reconstituted MitoXpress Xtra reagent. To prevent oxygen diffusion from ambient air, 100 µL of prewarmed HS mineral oil was carefully layered over each well. Plates were immediately transferred to a prewarmed Tecan Infinite 200 Pro plate reader, maintained at 37 °C. Fluorescence was recorded every 2 minutes for 2 hours using the following settings: excitation at 380 nm, emission at 650 nm, gain 220, 25 flashes, 30 µs integration delay, and 100 µs integration time.

Data were analyzed by determining the slope of the fluorescence lifetime signal (indicative of oxygen consumption) after an equilibration period of 20 minutes. The linear portion of the curve (typically over a 60-minute window) was used for slope determination. Background signal from cell-free wells was subtracted from all conditions to obtain net oxygen consumption rates. Data was normalized to the untreated control.

Thin section electron microscopy

The *T. gondii in vitro* cyst culture was performed as described above. After four weeks, cells were treated with 5 µM atovaquone, 500 nM buparvaquone, 5 µM MMV1028806, corresponding to the lowest lethal concentration, or 0.1% DMSO, as a solvent control, for 24h. Cells were fixed by covering the monolayer with a fixation solution of 1% paraformaldehyde and 2.5% glutaraldehyde (Electron Microscopy Services) in 0.05 M HEPES buffer (pH 7.4). After incubation at RT for 3h, samples were sealed with parafilm and stored in the fridge until further processing. Fixed cells were scraped, sedimented by centrifugation (3000 *g*, 10 min) using a swing-out rotor and washed twice with 0.05 M Hepes buffer. The cell sediment was embedded in low melting-point agarose and stored in 2.5% glutaraldehyde in 0.05 M HEPES buffer. Small blocks of the embedded sediment were subjected to postfixation (osmium tetroxide, tannic acid), *en bloc* contrasting (uranyl acetate), dehydration, and embedding in epoxy resin according to a standard protocol (Laue 2010).

Ultrathin sections were prepared with an ultramicrotome (UC7, Leica Microsystems, Germany) using a diamond knife (45°, Diatome, Switzerland). The sections were collected on bare copper grids (300 mesh, hexagonal mesh shape) or copper slot grids, contrasted with 2% uranyl acetate and 0.1% lead citrate, and coated with a thin (2-3 nm) layer of carbon. The ultrathin sections with a thickness of ~65 nm were examined with a transmission electron microscope (Tecnai Spirit, Thermo Fisher Scientific) operated at 120 kV. Images were recorded with a side-mounted CCD camera (Phurona, EMSIS, Germany) at 4095 x 2984 pixels.

Metabolomic analysis of drug-treated intracellular tachyzoites and bradyzoites

HFF cells were infected with PruTom at MOI 1. Approximately 64 h later, the cultures were treated with 1.5 µM MMV019189, 7 µM MMV1028806, 30 µM MMV228911, 1.2 µM MMV595321, 5 µM MMV688762, 2 µM MMV688854, respectively, and 0.1 % DMSO as a control for 3h. These concentrations correspond to the five-fold respective IC_{50} value. Cultures were quenched in ice-cold PBS and parasites were isolated from their host cells by passaging scraped monolayers through a 23G needle, filtered through a 3 µm polycarbonate filter and after microscopic counting distributed at 1×10^8 parasites per sample. Parasites were washed with ice-cold PBS via 5 min centrifugation at 21,500 x g at 0 °C. Metabolites were then extracted by sonication for 2 min in 50 µL 80 % acetonitrile in water. The insoluble particles were pelleted for 5 min at 21,500 x g at 0 °C. 5 µL of each sample was collected to generate a pooled biological quality control (PBQC) which was injected regularly within the otherwise randomized run list. 5 µL of each sample was injected onto either an ACQUITY UPLC BEH amide column (Waters) equipped with a VanGuard amide pre-column (Waters) or onto a SeQuant ZIC-pHILIC (Merck) column with an OPTI-LYNX ZIC-pHILIC guard column cartridge (Optimize Technologies) operating on a Vanquish Flex (Thermo Fisher). Analytes were chromatographically separated starting with 100 % eluent A (90 % acetonitrile in water with 10 mM ammonium carbonate) and 0 % eluent B (10 mM ammonium carbonate in water), decreasing to 40% A and 60% B over 25 min. Metabolites were detected on a Q Exactive Plus (Thermo Fisher) equipped with an electrospray ionization source operated in rapid polarity-switching mode at an MS^1 resolution of 70k and an MS^2 resolution of 35k (Top3 MSX1). Peak extraction, retention time alignment, and peak annotation were computed with Compound Discoverer 3.1 (Thermo Fisher) using an in-house library of authentic standards. Blank subtraction, gap-filling as well as statistical analysis were performed in Excel (Microsoft), and the data was plotted with GraphPad Prism 8.4.1. Principal component analysis was performed with ClustVis (Metsalu and Vilo 2015).

Isotopic labeling

Intracellular parasites were treated as mentioned above and incubated for three hours with DMEM containing either 25 mM U-^{13}C-glucose or 4 mM ^{15}N-amide-glutamine instead of the respective unlabeled carbon sources. Quenching, parasite isolation, metabolite extraction and liquid chromatography were performed as described. To increase coverage of the isotopologues, metabolites were measured in positive and negative mode separately without MS^2 scans but with an increased MS^1 scan resolution of 140k. For peak identification purposes, the unlabeled DMSO-treated sample was analyzed with intermittent MS^2 scans as described above.

ATP luciferase assay

PruTom tachyzoites were cultivated in HFF cells without glucose, isolated 48 h post-infection and treated with 1 µM atovaquone or HDQ for 3h. PruTom bradyzoites were matured for three weeks, glucose-starved for an additional week and treated for 24 h with 1 µM atovaquone or HDQ before being released by needle passage 23G, pepsin digestion for 20 min at 37 °C. The digestion reaction was stopped by neutralizing in 1.2 % sodium bicarbonate (pH ≈ 8.3) with 0.0159 g/l phenol red as pH indicator dye (Christiansen et al. 2022). They were isolated by douncing on ice and filtered through a 3 µm polycarbonate filter. 10^5 parasites were then

aliquoted in 100 µL ice-cold PBS in a 96-well plate. ATP was quantified using BacTiter-Glo™
Microbial Cell Viability Assay (Promega) according to the manufacturer's protocol using a
standard curve. Data were plotted in GraphPad Prism 8.4.1..

Metabolomic analysis drug responses on *in vitro* cysts
Intracellular cysts were isolated from their myotube host cells and extracted as described in
(Christiansen et al. 2022; Maus et al. 2024). In short, cysts were released by needle passage
(23G) and captured with DBA-coated magnetic beads. PBS (with 2% BSA) washed cysts were
extracted in 80% acetonitrile and analyzed as described above. One dish yielded
approximately 10^6 cysts and was distributed into 3 samples. Prior to a statistical analysis, the
dataset was normalized by dividing the compound intensities through the summed intensities
of the respective samples.

Results

The Pathogen Box contains dually active compounds

To initially identify compounds that simultaneously target the tachyzoite and the bradyzoite stage of *T. gondii*, we tested 371 of 400 compounds from the Pathogen Box at a concentration of 10 µM against both stages. We excluded previously reported active compounds against *T. gondii* tachyzoites (Spalenka et al. 2018) from this initial screen. HFF cells were infected with tachyzoites of the Pru strain expressing tandem Tomato fluorescent protein in its cytosol (PruTom) at an MOI of 1 (Fig. 1A). 47 compounds completely inhibited growth during 7 days of exposure while the parasitemia of DMSO-treated controls already peaked after 4 days (Fig. 1B). To also identify bradyzocidal compounds, we differentiated KD3 myoblasts for one week into multinucleated myotubes and infected them under bicarbonate-deplete conditions and at a multiplicity of infection of 0.1 for four weeks to allow maturation of tissue cysts (Fig. S1). We exposed the cultures to 10 µM of the 371 MMV compounds for one week, removed the inhibitors and stimulated re-differentiation into tachyzoites by bicarbonate supplementation (Fig. 1C). Growth of DMSO-treated cultures became detectable after 10 to 14 days. Interestingly, 31 compounds prevented any tachyzoite regrowth (Fig. 1D), indicating their bradyzocidal activity. To exclude host cytotoxicity-dependent suppression of parasite growth, we measured cell viability via the reductive potential of myotubes and HFF cells using a resazurin assay. At 10 µM, 30, 12 and 13 compounds inflicted cytotoxicity onto HFFs, myotubes or both cell types, respectively (Supplemental File 1). This initial screen identified 14 compounds with activity against both tachyzoites and bradyzoites and no apparent cytotoxicity (Fig. 1E).

Validation of screening hits

To validate our results, we determined half-inhibitory concentrations (IC_{50}) in PruTom tachyzoites (Fig. 2A, B) and half-maximal lethal concentration (LC_{50}) in bradyzoites. We also included 29 compounds that were previously reported to target tachyzoites (Spalenka et al. 2018). We confirmed dual activity of 16 compounds that exhibited mean IC_{50} values ranging between 70 pM and 1.1 µM for tachyzoites (Fig. 2E, Fig. S2D) and LC_{50} values between 0.15 and 8.2 µM for bradyzoites (Fig. 2F, Fig. S2D). To allow a comparison between the tachyzoite and bradyzoite data, we also determined the respective minimal lethal concentrations (MLC) over a seven-day treatment duration of tachyzoites as well (Fig. 2G). Except for four not significantly different compounds (MMV021013, MMV022478, MMV658988, MMV65900), minimal lethal concentrations appear generally higher in bradyzoites, potentially reflecting their lower metabolic activity and/or the cyst wall as a diffusion barrier.

Dually active MMV Pathogen Box hits share high lipophilicity and are indicated to be active against other pathogens

To gain insight into general properties of active compounds from our initial screen (Fig. 1E), we tabulated indicated activities as provided by MMV (Fig. S2A, D). As expected, these compounds were mostly earmarked as active against *T. gondii* tachyzoites (67%), however, none of those compounds exhibited bradyzocidal activity. Instead, 27% of dually active compounds were considered active against *Cryptosporidium*. Also, despite the relative

phylogenetic proximity of *P. falciparum,* only 5% of the 125 potential antimalarials blocked growth of both *T. gondii* stages. To characterize physicochemical properties of these dually active hits, we approximated their lipophilicity by their logP values (Supplemental File 1) and compared them to compounds that were inactive against *T. gondii* (Fig. S2B). We found that tachyzocidal, bradyzocidal and dually active compounds possess a statistically significantly higher lipophilicity than non-active compounds and this trend appeared more accentuated for bradyzocidal and dually active compounds. However, when comparing tachyzocidal with bradyzocidal compounds, including dually active ones, we did not find a difference (p = 0.06). We estimated permeability through the blood brain barrier (bbb) using LightBBB (Shaker et al. 2020) and gastrointestinal absorbability using SwissADME (Daina, Michielin, and Zoete 2017) (Supplemental File 1). Most of the dually active drugs were predicted to be both permeable across the bbb and absorbed in the intestine (Fig. S2C). Together, these data support phylogenetic relationship as a weak predictor of activity of Pathogen Box compounds against *T. gondii* bradyzoites and highlight the role of lipophilicity among the compounds tested here.

Metabolic drug responses in tachyzoites indicate conserved and novel modes of action

We chose to focus on dually active compounds, because they are clinically more relevant and investigation of their targets allows comparisons between on parasite stages. To that end, we choose five compounds based on their availability and yet unknown mode of action (Fig. 3, Supplemental File 2, Fig S3C) to test metabolite responses of tachyzoites. Infected monolayers were treated for three hours with compound concentrations five times their respective IC_{50} values or the solvent DMSO. The inhibitor MMV688854 (BKI 1294) (Doggett et al. 2014) is considered to inhibit a non-metabolic target (calcium-dependent protein kinase (CDPK1)) at the employed concentration (Hayward et al. 2023) and served as an additional control to estimate the metabolic impact of growth inhibition. We filter-purified parasites from quenched monolayers and analyzed their acetonitrile extracts by LCMS using both, pHILIC and BEH-amide chromatography columns. We generated 9 samples per condition from three repeats of the experiment with triplicate samples. A principal component analysis illustrates that batch effects are the main contributors to biological variances within the dataset (Fig. S3A, B). Despite that, all compounds elicited a distinct metabolic phenotype on both LCMS systems (Fig. 3). The CDPK1 inhibitor increased levels of AMP, oxidized glutathione, pseudo uridine, citrulline, glycolate and glutamyl glutamine, while reducing the abundances of GABA, carbamoyl aspartate, phosphoenolpyruvate and a hexose phosphate. Shifts in these metabolites are shared among many treatments and indicate their implication in a general stress response and thus do not signify direct interference with parasite metabolism.

Inhibitor-specific responses include the accumulation of cGMP-AMP after treatment with MMV019189, accumulation of carbamoyl aspartate and dihydroorotate upon MMV1028806 treatment, and a general depletion of central carbon metabolites (succinate, citrate, glyceraldehyde 3-phosphate, ATP, MEP) in MMV228911-treated parasites. MMV595321 caused an accumulation of N-acetylated amino acids, and MMV688762 led to uracil deficiency and an accumulation of dihydroorotate and orotate. These data suggest that growth inhibitors cause a bipartite metabolic response composed of generic and compound-specific changes. While these findings highlight compound-specific metabolic signatures, some observed alterations may represent downstream or systemic effects of inhibitor action, reflecting the inherent complexity of parasite metabolism.

MMV1028806 and buparvaquone elicit metabolic responses similar to the *bc1*-complex inhibitor atovaquone.

The metabolic impact of most tested MMV compounds indicated a novel mode of action that did not resemble known or recognizable metabolic fingerprints. However, the metabolic fingerprint of MMV1028806 was consistent with inhibition of pyrimidine biosynthesis at the bc_1-complex-dependent oxidation reaction of dihydroorotate to orotate, as has been reported for *P. falciparum* (Fig. 4A) (Allman et al. 2016; Creek et al. 2016; Antonova-Koch et al. 2018). A similar response would be expected to buparvaquone, which we found to be dually active against *T. gondii* tachyzoites and bradyzoites and which targets the bc_1-complex in *Theileria annulata* (McHardy et al. 1985), *Neospora caninum* (Müller et al. 2015) and *T. gondii* tachyzoites (Hayward et al. 2023).

To test these hypotheses, we exposed intracellular parasites to MMV1028806 or BPQ while substituting glucose with $^{13}C6$-glucose for three hours. We then measured the incorporation of ^{13}C into the central carbon metabolites as well as the intermediates of the pyrimidine synthesis pathway. As controls, we utilized the known bc_1-complex inhibitor atovaquone and the solvent DMSO (Supplemental File 3). Treatment with all three inhibitors led to a dramatic accumulation of +3 labeled aspartic acid, carbamoyl aspartate and dihydroorotate which was accompanied by a depletion of uracil and uridine, in particular of the +5-labeled uridine isotopologue. These changes are consistent with impaired function of the bc_1-complex-dependent dihydroorotate dehydrogenase (Fig. 4 B, C). Another common effect was the accumulation of succinate and depletion of fumarate, suggesting reduced recycling of $FADH_2$ via the bc_1-complex.

To independently verify our results, we hypothesized that *T. gondii* also incorporates glutamine-derived nitrogen into pyrimidines. We replaced glutamine with its ^{15}N-amide-labeled isotopologue in the tachyzoite culture media for three hours while simultaneously exposing the parasites to ATQ, BPQ or MMV1028806. As expected from our previous experiments, we observed a marked decrease of uracil and uridine levels while aspartate, carbamoyl aspartate and dihydroorotate accumulated again (Fig. 4D, Supplementary File 5). ^{15}N was incorporated into many metabolites (Fig. S4). Consistently, labelling of pyrimidines was diminished while carbamoyl aspartate and dihydroorotate were heavily labeled as M+1 isotopologues (Fig. 4E).

Together, these data show that treatments with all three compounds lead to the accumulation of metabolites upstream of the bc_1-complex-dependent reactions during pyrimidine biosynthesis and the TCA cycle and to the depletion of downstream metabolites (Fig. 4A). These metabolic signatures suggest an inhibition of the *T. gondii* bc_1-complex in tachyzoites.

Atovaquone, buparvaquone and MMV1028806 decrease the mitochondrial membrane potential of tachyzoites and bradyzoites and block the electron transport chain

The inhibition of the bc_1-complex by ATQ leads to the collapse of the mitochondrial membrane potential in *T. gondii* tachyzoites (Vercesi et al. 1998) and we anticipated a similar phenotype in BPQ or MMV1028806-treated parasites. We treated intracellular tachyzoites and bradyzoites for 24 h with the three inhibitors or 0.1 % DMSO alone and estimated the mitochondrial membrane potential and morphology using a fluorescent Mitotracker dye. The mitochondria of control tachyzoites appeared as extended loops but largely collapsed after

treatments with all three compounds (Fig. 5A, B). Notably, also host mitochondria appear influenced. To evaluate the specificity of this assay we used pyrimethamine, clindamycin and 6-diazo-5-oxonorleucine (DON) that target dihydrofolate reductase, apicoplast translation and glutaminase as controls. Mitotracker and GFP signal intensities remained comparable to DMSO controls (Fig. S5). DAPI signal intensity appeared reduced in pyrimethamine-treated samples. In bradyzoites, mitochondria appeared mostly as spheres or were donut-shaped (Supplementary File 6) and their relative fluorescence decreased after 24 h treatment with ATQ, BPQ or MMV1028806 at 5 µM, 500 nM or 5 µM, respectively while their morphology appeared unchanged (Fig. 5C, D). Analysis by thin section electron microscopy revealed a largely unaffected sub-mitochondrial ultrastructure but the areas of mitochondrial profiles were changed in comparison to control after exposure with ATQ and MMV1028806 but not with BPQ (Fig. S6). This indicates that the latter decreases Mitotracker fluorescence intensity directly, while ATQ and MMV1028806 cause both, a drop in Mitotracker fluorescence and a reduced size of the mitochondria which leaves open whether a change of the membrane potential, of the size, or even of both parameters were responsible for the drop in Mitotracker fluorescence.

We next tested directly whether the mETC was blocked by ATQ, BPQ and MMV1028806 by measuring oxygen consumption of treated tachyzoites using a fluorescence-based assay (Fig. 5E). The solvent DMSO, the decoupling agent carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP), and the $bc1$-complex inhibitor antimycin A served as controls. Treatment with FCCP led to a marked increase in oxygen consumption compared to the DMSO control, indicating maximal respiratory capacity of the parasites. In contrast, treatments with ATQ, BPQ, MMV1028806, and antimycin A resulted in substantially reduced oxygen consumption levels relative to the DMSO control and suggest indeed a blockage of the mETC consistent with the inhibition of the $bc1$-complex.

Inhibitors of the mitochondrial electron transport chain alter catabolic processes in bradyzoites

Next, we sought to confirm this mode of action directly on bradyzoites and record the metabolic impact of ATQ, BPQ or MMV1028806. To that end, we exposed intracellular four-week-old cysts to 5 µM, 500 nM or 5 µM of the respective compounds and for three hours and used 0.2% DMSO as a mock treatment. These concentrations represent their LC_{50} against bradyzoites and were chosen to minimize off-target effects. The cysts were isolated for LCMS analysis using magnetic beads (Christiansen et al. 2022) (Fig. 6A). Common in all three distinct metabolic phenotypes (Fig. S7) is the accumulation of AMP and NADH, while varying TCA cycle intermediates, such as malate, succinate and α-ketoglutarate and mitochondria-independent energy sources such as glucose and phosphocreatine were depleted (Fig. 6B-D). These metabolic changes suggest an energy starvation scenario marked by rising AMP levels and reduced recycling of NADH via the bc_1-complex. Interestingly, in BPQ and MMV1028806 treated cysts, we also observed prominent accumulation of several acyl-carnitines species which are mobile intermediates of ß-oxidation (Fig. 6B-D). Since this pathway is thought to be absent in *T. gondii,* we attribute these changes to potential inhibitory effects on host mitochondria. Notably, we did not observe changes in the pyrimidine synthesis pathway, which may reflect low nucleic acid demand from largely dormant bradyzoites. These data demonstrate that bradyzoites, which were considered metabolically inactive, exhibit metabolic responses to inhibitory compounds. Although we characterized AMP accumulation as part of a general stress response in tachyzoites, the metabolic profiles of inhibited

bradyzoites are consistent with inhibition of their mitochondrial bc_1-complex and suggest its
role in ATP generation.

The differential efficacy of atovaquone and HDQ against bradyzoites correlates
with the inhibition of ATP levels in bradyzoites
We sought to test the importance of the mETC chain in ATP generation in bradyzoites and
exploited the differential targets of the coenzyme Q analogs ATQ and 1-hydroxy-2-dodecyl-
4(1)quinolone (HDQ). However, while ATQ inhibits the bc_1-complex, HDQ inhibits
dihydroorotate dehydrogenase (DHODH) (Hegewald, Gross, and Bohne 2013) and the
alternative NADH dehydrogenase (NDH2) (Saleh et al. 2007). Both compounds block
tachyzoite growth at low nanomolar concentrations in vitro with IC50s at 20 and 4 nM,
respectively (Fig. 7A). However, in contrast to ATQ, HDQ does not affect the viability of mature
T. gondii bradyzoites despite diminishing their mitochondrial membrane potential (Christiansen
et al. 2022). Up to 20 µM HDQ did not arrest re-differentiation into proliferating tachyzoites,
while bradyzoites that were treated with 5 µM ATQ (LC_{50} 2.5 µM) did not generate proliferating
tachyzoites (Fig. 7B). To investigate how these modes of actions affect the mETC in
bradyzoites, we tested their influence on mitochondrial ATP production by luciferase-based
ATP assay. This assay shows a linear response to parasite cell equivalents between 10^3 and
10^6 parasites per 100 µL, and in order to minimize variance we decided to work with 10^5
parasites (Fig. S8). To maximize mitochondrial ATP production, we starved three-week-old
bradyzoites for one additional week of glucose. As a comparison, we utilized tachyzoites that
were cultured for two days in glucose-deplete conditions. Untreated tachyzoite samples
contained in average 196 ± 21 fmol ATP per 10^5 parasites, whereas untreated bradyzoites
only contained 14.1 ± 1.1 fmol. (Fig. 7C). Three hour-long treatment with 5 µM ATQ decreased
ATP levels in both stages by 75 % (Fig. 7D). In contrast, HDQ reduced ATP levels by 60 % in
tachyzoites but only marginally affected bradyzoites (Fig. 7D). These data indicate that
bradyzoites can maintain very low ATP levels and suggest the presence of a diminished but
obligate production of ATP in the mitochondria that functions independently of exogenous
glucose and HDQ-target enzymes.

Discussion

New treatment options for chronic *T. gondii* infections are needed, as chronic infections are
currently incurable and available treatments for acute toxoplasmosis are associated with
severe side-effects, in particular in infants after congenital transmission (Konstantinovic et al.
2019). To simultaneously identify drug targets in bradyzoites and along with their inhibitory
compounds, we utilized a recently developed *in vitro* culture system of pan-drug tolerant
bradyzoites (Christiansen et al. 2022) to screen 400 compounds of the MMV Pathogen Box.
16 compounds exhibited confirmed cidal activity against bradyzoites while also inhibiting
growth of tachyzoites. Metabolic profiling and stable isotope tracing in treated tachyzoites
suggested the inhibition of the mitochondrial bc_1-complex by MMV1028806 and the reference
compound BPQ.

Interestingly, of the 15 compounds in the Toxoplasmosis disease set, previously earmarked
by MMV, only 10 were found to be active against tachyzoites and none against bradyzoites.
These differences may be attributed to strain-specific susceptibilities which are known for anti-
folates and atovaquone (Meneceur et al. 2008) and differences in the design of growth assays.
MMV assessed activities via plaque assays that prevent media change and rely on visual
inspection of growth of a low number of parasites and might overestimate inhibitory effects.

We found that bradyzocidal compounds exhibit higher predicted logP coefficients, suggesting
that the parasite entry might occur via diffusion across the cyst wall and less via solute
transporter proteins. A known example of this principle is the anti-malarial drug fosmidomycin,
where high polarity and low import rates caused minimal uptake by and efficacy against *T.*
gondii tachyzoites (Nair et al. 2012; Baumeister et al. 2011). A similar case might also be made
for the antifolates, pyrimethamine, sulfadiazine, trimethroprim and sulfamethoxazole which
exhibit low consensus logP values of 2.09, 0.4, 1.16 and 0.92 and which also fail to eradicate
tissue cysts (Alday and Doggett 2017; Enshaeieh et al. 2021). Interestingly, MMV675968 is a
dually active compound with a logP value of 2.46, that has been shown to be a potent inhibitor
of the dihydrofolate reductase (DHFR) in a range of pathogens (Songsungthong et al. 2019,
2021; Kim et al. 2021; Borba-Santos, Vila, and Rozental 2020; Nugraha et al. 2019; Rollin-
Pinheiro et al. 2021; Vila and Lopez-Ribot 2017; Cantillon et al. 2022). Its target in *T. gondii*,
however, remains to be identified.

Another important diffusion barrier that needs to be overcome to eradicate *T. gondii* tissue
cysts is the blood brain barrier (BBB). Pyrimethamine and trimethroprim penetrate the blood
brain barrier (Geils et al. 1971; Ineichen et al. 2020), they do not clear brain-resident cysts
(Montazeri et al. 2018; Enshaeieh et al. 2021; Christiansen et al. 2022) highlighting the cyst
wall as a potentially decisive obstacle. Consistent with this hypothesis is the tolerance of
mature in vitro tissue cysts against high dose of sulfadiazine and pyrimethamine (Christiansen
et al. 2022).

To characterize the modes of action of four additional inhibitors, we used metabolomics. In
general, metabolic phenotypes reflect direct consequences of target inhibition as well as
indirect effects resulting from attenuated growth and stress responses. We chose the bumped-
kinase inhibitor MMV688854 (BKI-1294) that targets the calcium-dependent protein kinase 1
(CDPK1) as a control compound to identify metabolites implicated in a general stress response
and adaptation to slow growth (Winzer et al. 2015). CDPK1 is needed for invasion (Johnson
et al. 2012) and also affects formation of daughter cells. An accumulation of AMP and oxidized
glutathione are the most conserved responses across the six treatments and may indicate
depleted energy levels and oxidative stress (Caro et al. 2012). Also, pseudo uridine is elevated
in many treatments. Enhanced pseudo uridylation of RNA is considered to be implicated in
stress responses (Cerneckis et al. 2022) and modulates RNA stability (Davis 1995). It remains
to be tested which of these conserved metabolic responses signify cellular decay and which
ones take part in a productive metabolic response in *T. gondii*. Nonetheless, metabolic profiling
data on antibiotics-exposed bacterial pathogens has been used to design combination
therapies, by either interfering with productive metabolic stress responses (Zampieri et al.
2017) or by simultaneously inhibiting otherwise non-essential pathways (Campos and Zampieri
2019). We found that the systematic identification of target structures and drug modes of action
from profiling data alone remains challenging due to inherent non-linear relationships within
the cellular metabolic network, the underlying regulatory mechanisms of enzyme functions. A

systematic mapping of genetic disruptions to metabolic phenotypes, as demonstrated in *E.*
coli, may alleviate these restrictions (Anglada-Girotto et al. 2022).

Apart from the general metabolic stress response to the treatment with MMV688854, three
compounds caused apparent inhibitor-specific signatures. MMV019189, which is known to
inhibit *P. falciparum* (Duffy et al. 2017), but also *C. parvum* (Hennessey et al. 2018) and *L.*
mexicana (Berry et al. 2018), lead to the accumulation of c-GMP-AMP in the parasite. This
dinucleotide is produced by the c-GMP-AMP synthase (cGAS) in mammalian cells in response
to cytosolic DNA and triggers a type-I interferon response (Sun et al. 2013). While cGas is
considered absent in *T. gondii*, host c-GMP-AMP induces STING-signaling as a response to
T. gondii infection (P. Wang et al. 2019). Hence, MMV019189 may implicate the host in its
antiparasitic activity.

MMV228911-treated tachyzoites showed the depletion of the DXR product 2-C-
methylerythritol 4-phosphate (MEP), a metabolite that occurs in the non-mevalonate pathway
in *T. gondii'*s apicoplast. Other metabolites upstream of DXR are either not detected or
depleted, such as glyceraldehyde 3-phosphate, but also partake in other pathways.
Interestingly, MMV228911 also caused a broader dysregulation of central carbon metabolism
marked by a decrease of many phosphorylated metabolites such as nucleotide triphosphates
and PEP but also citric acid and acetyl-CoA. This is consistent with metabolic signaling beyond
the apicoplast, which also has been implicated in *P. falciparum*, where a cytosolic sugar
phosphatase is responsible for resistance to the DXR inhibitor Fosmidomycin (Guggisberg et
al. 2014).

We identified MMV1028806 as a candidate bc_1-complex inhibitor by comparing its metabolic
impact with the one of BPQ with ATQ. All three compounds cause a sharp increase of
pyrimidine synthesis substrates, depletion of pyrimidines and dysregulation of TCA cycle
intermediates. This is consistent with previous mass spectrometry-based studies that
characterized metabolic fingerprint of atovaquone on *P. falciparum* intraerythrocytic blood-
stages (Creek et al. 2016; Allman et al. 2016; Antonova-Koch et al. 2018). In addition to
abundance data, the incorporation of ^{13}C and ^{15}N stable isotopes from glucose and glutamine,
respectively, into TCA cycle and pyrimidine biosynthesis intermediates suggest the *bc1-*
complex as a target. Both, the isotopologue distributions and metabolite abundances strongly
suggest attenuation of pyrimidine production and TCA cycle fluxes by all three inhibitors
(Hortua Triana et al. 2016). TCA cycle intermediates show characteristic isotopologue
distributions that are indicative of a ^{13}C-label incorporation via acetyl-CoA (MacRae et al.
2012). A relatively increased M+3 isotopologue species in dicarboxylic acids in ATQ- and BPQ-
treated parasites may indicate enhanced anaplerotic synthesis from glycolytic intermediates.
These distinctions from MMV1028806 metabotypes may reflect a combination of off-target
effects and differential timing of the metabolic inhibition.

Interestingly, six compounds from the MMV Pathogen Box have recently been shown to inhibit
O_2 consumption in tachyzoites through bc_1-complex inhibition (Hayward et al. 2023). We co-
identified Buparvaquone as a bc_1-complex inhibitor but we did not test the remaining five
compounds for their targets. However, in our screen at 10 µM we found MMV024397 and
MMV688853 to be dually active, while Trifloxystrobin (MMV688754) and Auranofin
(MMV688978) lacked inhibitory activity in bradyzoites, and Azoxystrobin (MMV021057) was
inactive. In addition, the previously described bc_1-complex inhibitor ELQ400 (MMV671636)
(McConnell et al. 2018) was dually active. Overall, these data are consistent with the fact that

the bc_1-complex is indeed needed for survival of bradyzoites, while not every tachyzocidal bc_1-
inhibitor also inhibits bradyzoites.

In contrast to tachyzoites, the metabolic phenotype in bradyzoites is not dominated by
dihydroorotate dehydrogenase (Fig. 6B, C, and D), as we observed neither the accumulation
of carbamoyl-aspartate, dihydroorotate, and aspartate, nor the depletion of pyrimidines.
Instead, AMP and NADH accumulate, which in eukaryotes signifies a low cellular energy status
(S.-C. Lin and Hardie 2018; X. Fu et al. 2019) that typically induces AMP-activated protein
kinase and a glycolytic response. AMPK is a conserved enzyme with a pivotal role in cellular
homeostasis: it responds to elevated AMP levels by repressing anabolic activity and by
stimulating glucose uptake and glycolysis, β-oxidation, and ketogenesis (Winder and Hardie
1999; S.-C. Lin and Hardie 2018). *T. gondii* tachyzoites express AMPK which serves an
important role in metabolic reprogramming during the lytic cycle of tachyzoites (Li et al. 2023).
Transcriptomic and proteomic data indicate its presence in bradyzoites (Garfoot et al. 2019)
but its function remains unknown. We observed decreased levels of free glucose during all
three treatments that are consistent with stimulated glycolysis. We also observe depleted
phosphocreatine levels in bradyzoites (Fig. 6) during treatments. In muscle cells,
phosphocreatine acts as an ATP buffer system and replenishes ATP pools from ADP via
creatine kinase (Kuby, Noda, and Lardy 1954; Perry 1954). It remains to be seen whether
bradyzoites possess this mechanism. We also detect elevated levels of acyl-carnitines in BPQ
and MMV1028806 treated bradyzoites. These molecules act as shuttles for the mitochondrial
import of fatty acids for β-oxidation. However, this pathway has not been shown active and is
deemed absent in *T. gondii* (Seeber, Limenitakis et al. 2008, Shunmugam, Arnold et al. 2022).
The presence of acyl-carnitines in bradyzoites might reflect import from the host. It is
conceivable that their elevation in response to buparvaquone and MMV1028806 indicates
compromised functionality of the host bc_1-complex and subsequently accumulating β-oxidation
substrates. Indeed, BPQ has a very broad activity across Apicomplexa (Hudson et al. 1985)
and kinetoplastids (Croft et al. 1992).

We investigated the functionality of the mETC of tachyzoites and bradyzoites in response to
ATQ, BPQ, and MMV1028806 by Mitotracker staining and thin section electron microscopy.
All three treatments decrease overall Mitotracker fluorescence, consistent with a specific
decrease mitochondrial electron potential (Supplementary Fig. 5). In contrast, pyrimethamine,
clindamycin, and DON did not reduce the Mitotracker signal. Pyrimethamine-treated samples
displayed an increased Mitotracker-to-DAPI ratio, which can be attributed to diminished DAPI
intensity, likely resulting from inhibition of folate-dependent DNA synthesis via DHFR blockade.
Clin-treated samples showed no change in mitochondrial morphology or ratio, in line with its
delayed-death phenotype. Similarly, DON, which targets glutamine metabolism, did not alter
mitochondrial signal or ratio at 24 hours, though morphological changes were observed.
Together, these findings confirm that disruption of mitochondrial membrane potential is specific
to compounds targeting the electron transport chain, while inhibitors acting on nuclear or
metabolic pathways do not induce rapid mitochondrial dysfunction under the conditions tested.
ATQ and MMV1028806 but not BPQ also shrink areas of mitochondrial cross-sections
indicating a potential contribution to the observed change in overall fluorescence. These
differences might reflect distinct mitochondrial morphologies. In agreement with previous
observations (Place et al. 2023), we found that bradyzoite mitochondria differ from those in
tachyzoites, predominantly occurring as "blobs" rather than tubes or loops, and our
observations align with these findings (Supplementary File 6).

The oxygen consumption assays highlight the functional sensitivity of the T. gondii RHΔku80
mETC to pharmacological perturbation (Fig. 5E). As expected, FCCP, an uncoupler of
oxidative phosphorylation, markedly increased oxygen consumption, confirming the parasite's
ability to upregulate electron flow when the proton gradient is dissipated. Antimycin A nearly
abolished respiration, demonstrating the essential role of complex III in maintaining
mitochondrial activity. Likewise, ATQ and BPQ, both *bc1*-complex inhibitors, strongly
suppressed oxygen consumption. Notably, MMV1028806 produced a comparable inhibitory
effect, suggesting that it also targets complex III or a functionally linked site within the
mitochondrial electron transport chain. This observation differs from the findings of (Hayward
et al. 2023), who did not report MMV1028806 as an mETC inhibitor in their Seahorse-based
screen of the MMV Pathogen Box. A key difference is that our assays were performed at
10 µM, whereas (Hayward et al. 2023) used 1 µM. These results support a mitochondrial mode
of action for MMV1028806 under our experimental conditions and underline the value of
oxygen consumption assays for detecting dose-dependent effects on parasite respiration.

Our findings strongly suggest that bradyzoite survival is critically dependent on mitochondrial
electron transport. We investigated this by comparing the effects of two inhibitors: HDQ, which
kills tachyzoites but not bradyzoites, and ATQ, which kills both. Although HDQ enters
bradyzoites and reduces their mitochondrial membrane potential, its inability to kill them
implies its molecular targets are non-essential in this stage. To pinpoint the vital process, we
measured ATP levels and found that only the bradyzocidal compound, ATQ, caused a collapse
in bradyzoite ATP. HDQ did not, despite both drugs depleting ATP in tachyzoites. This directly
links bradyzoite viability to mitochondrial ATP production. The different outcomes are
explained by their distinct targets: ATQ inhibits the *bc1*-complex (McFadden et al. 2000), which
is central to ATP synthesis, whereas HDQ blocks other pathways like pyrimidine synthesis and
mitochondrial NADH oxidation through DHODH and NDH2. These results also support the
notion that the canonical TCA cycle is not the primary source of electrons for bradyzoites
(Christiansen et al. 2022). A variety of electron sources supply the electron transport chain
(Maclean et al. 2021). The sources that are crucial for ATP production remain to be identified.

Bradyzoites may also utilize glycolysis to generate ATP. This has been suggested to be the
case in early, seven-day-old bradyzoites where gliding motility, an ATP dependent process,
can be inhibited by withdrawal of exogenous glucose and by the glycolytic inhibitor 2-
deoxyglucose. However, in these parasites, also Oligomycin A inhibited motility suggesting a
simultaneously essential contribution of the mitochondrial ATP (Fu, Brown et al. 2021). In
addition, hexokinase has been shown to be needed for optimal cyst production in chronically
infected mice, suggesting an important role of this protein either though its enzymatic activity
or other functions (Shukla et al. 2018). It remains to be investigated how mature bradyzoites
rely on glycolysis to produce ATP and how this is integrated with mitochondrial ATP production.

Interestingly, despite its profound effect on ATP levels and viability of bradyzoites, ATQ fails
to eradicate all *T. gondii* cysts from infected mice and does not reliably prevent toxoplasmosis
relapse in AIDS patients and mice (Araujo, Huskinson, and Remington 1991; Kovacs 1992;
Ferguson et al. 1994; Katlama, Mouthon, et al. 1996). Underlying reasons for this discrepancy
need to be established but may include insufficient bioavailability. Quinolones are known to be
poorly soluble in aqueous environments. In addition, the predicted permeability of the blood-
brain barriers to either ATQ, BPQ and HDQ is low. Indeed, improved bioavailability of ATQ in
nanosuspensions improved efficacies (Shubar et al. 2011). Also, BPQ treatment of mice that
were experimentally infected with closely related *Neospora caninum* parasites did target

parasites in muscle but not in brain tissue (Müller et al. 2015). A similar conjecture arose when
comparing endochin-like quinolones. Here, favorable pharmacokinetics of a variant with lower
intrinsic potency likely conferred higher *in vivo* efficacy against cysts (Doggett et al. 2012).

Together, our data illustrate the functional importance of the bc_1-complex for the viability of
bradyzoites and encourage the pursuit of bc_1-complex inhibitors as candidates for curative
toxoplasmosis treatment. Both, efforts to develop multi-site-directed inhibitors (Doggett et al.
2020; Alday, Nilsen, and Doggett 2022) and re-purposing of respective drugs are promising
avenues. However, our data also underline the importance of favorable pharmacokinetics to
ensure passage through the blood-brain barrier and the cyst wall.

Figures

Figure 1: An in vitro screen reveals tachy- and bradyzocidal properties of the MMV Pathogen Box compounds.

(A) Experimental scheme of the MMV compound screen on *Toxoplasma gondii* tachyzoites in human foreskin fibroblast cells (B) Growth curves of tachyzoites treated with 10µM of each MMV compound for 7 days (shades of grey), a solvent control (green), and an inhibition control (red) (C) Experimental scheme of the MMV compound screen on bradyzoites matured in myotubes (D) Growth curves of bradyzoites treated with 10µM of each compound (shades of grey) or a solvent control (green) for 7 days, followed by a tachyzoite regrowth phase (E) Pie chart displaying the number of Pathogen Box compounds and their effects on our in vitro cultures.

747
Figure 2: Confirmed Pathogen Box compounds that target both tachyzoites and
mature bradyzoites.

(A) shows the growth curve of MMV689480 (Buparvaquone), one of the 16 dually active
compounds, in a tachyzoite growth assay (n=8). The infected cultures were treated for 7 days
and regrowth in absence of the inhibitor was observed for another week. Depending on the
development of the DMSO control, the half-maximal inhibitory concentration (IC_{50}) was
determined at either day 4 or 7 (arrow). (B) shown is the respective data plotted as relative
fluorescence units depending on the concentration at day 7. (C) With 4 weeks old, mature
bradyzoite cultures (n=6), the half-maximal lethal concentration (LC_{50}) was determined at the
time point which ideally reflected the dose-response curve (in the case of Buparvaquone at
day 31; arrow). (D) The LC_{50} of Buparvaquone on encysted bradyzoites determined at day 31.
All IC_{50} and LC_{50} of dually active compounds are summarized in (E) for tachyzoite and (F) for
bradyzoite cultures. The grey dotted line indicates 1 µM in both charts. (G) Comparison of the
minimal lethal concentration of each compound on tachyzoites and bradyzoites. Mann-Whitney
test of eight replicative measurements on tachyzoites and six replicates of bradyzoites from
two independent experiments each *** $p < 0.001$.

**Figure 3: Untargeted metabolomics of tachyzoites treated with 6 screening hits effective
against both stages of *T. gondii*.**

Volcano plots showing the parasitic, intracellular metabolic phenotype caused by MMV
Pathogen Box compounds that inhibited tachyzoites and bradyzoites during the screen. The
metabolome was measured with two columns of complementary chemistries to enhance
metabolite coverage. Blue dots indicate metabolites which were significantly reduced
compared to DMSO treated parasites and red dots indicate significantly accumulated
metabolites (p < 0.05, log2 fold change <-1 or > 1, 3x biological replicates with 3x technical
replicates each, n = 9). Abbreviations: AcCoA - Acetyl-CoA, AcCys - Acetylcysteine, AcPhe -
Acetylphenylalanine, AMP - Adenosinemonophosphate, ATP - Adenosinetriphosphate,
CarbAsp - Carbamoylaspartate, cGMP-AMP - cyclic Guanosinemonophosphate-
adenosinemonophosphate, CMP - Cytosinemonophosphate, DHO - Dihydroorotate, GABA -
gamma-Aminobutyratic acid, GluCys - Glutamylcysteine, GluGln - Glutamylglutamine, Gro1P
- Glyceraldehyde 1-phosphate, Gro3P - Glyceraldehyde 3-phosphate, GSH - Glutathione
oxidized, GSSG - Glutathione reduced, GTP - Guanosinetriphosphate, HP4 - Hexose
phosphate No 4 (fourth chromatographic peak of 260.02972), HydKyn - Hydroxylkynurenine,
MEP - 2-C-Methyl-D-erythritol 4-phosphate, NAcAla - *N*-Acetylalanine, NAcCys - N-
Acetylcysteine, NAcMet - *N*-Acetylmethionine, NAcPhe - *N*-Acetylphenylalanine, OAcSer - *O*-
Acetylserine, PalmCarn - Palmitoylcarnitine, PEP - Phosphoenolpyruvate, PseudoU -
Pseudouridine, Trp - Tryptophan, UMP - Uridinemonophosphate, UTP - Uridinetriphosphate

A

B

^{13}C-Glucose — Log$_2$ fold changes

	ATQ			BPQ			MMV		
Succinate	2.6	2.2	2.4	2.5	2.5	2.5	1.5	1.5	2.4
Fumarate	-0.9	-1.0	-1.0	-1.2	-1.2	-1.4	-2.8	-2.9	-2.8
Aspartate	9.1	9.2	9.3	9.5	9.8	9.7	7.9	7.9	8.0
CarbAsp	10.0	10.3	9.8	10.4	10.7	10.3	8.2	8.5	8.1
DHO	9.4	9.3	9.3	9.4	9.3	9.3	7.8	7.7	7.7
Uracil	-1.6	-1.8	-1.8	-0.5	-0.6	-0.5	-0.9	-1.1	-1.0
Uridine	-0.8	-0.5	-0.7	-1.1	-1.0	-1.3	-0.9	-0.7	-1.0

C

Isotopologues

DMSO ATQ BPQ MMV

M+0, M+1, M+2, M+3, M+4, M+5, M+6, M+7, M+8, M+9

D

^{15}N-Glutamine — Log$_2$ fold changes

	ATQ			BPQ			MMV		
Aspartate	17.8	10.8	17.6	17.5	10.5	17.4	17.0	10.0	16.9
CarbAsp	6.7	6.3	6.7	6.6	6.4	6.6	5.9	5.7	3.0
DHO	10.0	10.2	9.6	10.7	10.9	10.4	10.1	10.3	9.9
Uracil	-2.7	-2.9	-2.9	-1.4	-1.4	-1.5	-1.6	-1.6	-1.7
Uridine	-0.4	-0.4	-0.4	-0.6	-0.5	-0.6	-0.8	-0.7	-0.7

E

Isotopologues

DMSO ATQ BPQ MMV

M+0, M+1, M+2

Figure 4: The metabolic response to electron transport chain inhibitors and MMV1028806.

(A) Scheme of the affected pathways upon *bc*$_1$-complex inhibition. Intracellular tachyzoites were grown in presence of U-^{13}C-glucose (B, C) or ^{15}N-amide-glutamine (D, E) instead of

unlabeled carbon sources and treated with atovaquone (ATQ), buparvaquone (BPQ) and
MMV1028806 (MMV) for 3h. (B, D) The treatment induced changes of mETC-related
metabolite abundances are shown as log_2-fold changes in comparison to DMSO treated
cultures. Shown are three replicate measurements. (C, E) Shown is the average
isotopologue distribution of mETC-related metabolites. The pie charts depict the number of
heavy carbon or nitrogen atoms incorporated into the respective metabolites (M+0 in light
grey represents the unlabeled metabolite, while M+1 in light blue indicates the integration of
one heavy atom into the molecule). Shown are the means of three replicate measurements.
Abbreviations: ATQ - atovaquone, BPQ - buparvaquone, CarbAsp - carbamoyl-aspartate,
DHO - dihydroorotate, DHODH - dihydroorotate dehydrogenase, GABA - gamma-
aminobutyrate, MMV - MMV1028806, PPP - pentose phosphate pathway, SSA - succinic
semialdehyde

Figure 5

Figure 5: Atovaquone, buparvaquone, and MMV1028806 reduce mitochondrial potential and respiration in *T. gondii*.

(A) HFF cells were infected with tachyzoites and treated with DMSO, atovaquone,
buparvaquone and MMV1028806 for 24 h, and mitochondria were stained with Mitotracker
(red) and DNA was stained with DAPI (blue). scale bar = 5 µm. (B) The ratios of fluorescence
intensities between Mitotracker (MT) and DAPI dyes were calculated per vacuole (DMSO n =
56, ATQ n = 45, BPQ n = 49, MMV1028806 n = 40; Blue lines represent the median, two-sided
Mann-Whitney test). (C) 4 weeks old, myotube derived ME49 in vitro cysts were treated with
DMSO, atovaquone, buparvaquone and MMV1028806 for 24 h, and mitochondria were
stained with Mitotracker (red), the cyst wall was stained with Dolichos Biflorus Agglutinin and
Streptavidin-Cy2 (green), and DNA was stained with DAPI (blue). scale bar = 20 µm. (D) The
ratios of fluorescence intensities between Mitotracker (MT) and DAPI dyes were calculated per
vacuole (DMSO n = 56, ATQ n = 42, BPQ n = 45, MMV1028806 n = 38; Blue lines represent
the median, two-sided Mann-Whitney test). (E) 10 million freshly egressed tachyzoites were
incubated with known bc_1 complex inhibitors ATQ and BPQ, specific control compounds
Antimycin and FCCP, and MMV1028806. Every treatment significantly impacts the oxygen
consumption rate (OCR) (blue line represents the median, n = 3) (Welch's test; ** p < 0.005).

Figure 6: Untargeted metabolomic analysis of bradyzoites treated with *bc*₁-complex inhibitors shows an energy imbalance.

(A) Four weeks old *in vitro* cysts were treated for three hours. The cell monolayer was scraped off and the cysts were syringe-released. With *Dolichos biflorus* agglutinin (DBA) coated magnetic beads, the isolated cysts were captured and washed with PBS. The metabolites were extracted with 80% acetonitrile (AcN) in water and sonication, and analyzed via hydrophilic-interaction ultra-high pressure liquid chromatography coupled mass spectrometry (HILIC-UHPLC-MS). The results are shown in three separate volcano plots: Metabolic phenotypes of bradyzoites treated with Atovaquone (B), Buparvaquone (C), and MMV1028806 (D). (n = 3, significance analyzed with a two-sided U-Mann-Whitney test) Abbreviations: ATQ - atovaquone, BPQ - buparvaquone, DMSO – dimethyl sulfoxide, LA-carnitine - linoleoylcarnitine, MMV - MMV1028806, NADH - reduced nicotinamide adenine dinucleotide, PA-carnitine - palmitoylcarnitine, SA-carnitine - stearoylcarnitine

Figure 7: Differential efficacy and ATP depletion by atovaquone and HDQ in tachyzoites and bradyzoites.

Atovaquone and HDQ were tested against *T. gondii* Pru parasites. (A) Showing the normalized half-maximal inhibitory concentration (IC_{50}) against tachyzoites in fibroblasts at day 4, n=8. (B) 4-week-old bradyzoites were treated for a week with Atovaquone or HDQ. Cultures were observed for tachyzoite re-growth for three weeks. The half-maximal lethal concentration (LC_{50}) was determined at the time point which ideally reflected the dose-response curve, in this case day 14, n=6. 10^5 tachyzoites and bradyzoites were lysed and the ATP concentration was determined with an enzymatic luciferase assay. (C) Showing the calculated concentration of ATP within an average parasite. Lines reflect the median, error bars represent SEM, significance test Mann-Whitney, n=12. (D) Parasites were treated with 1µM Atovaquone or HDQ. RFUs were normalized to the untreated control of either the tachyzoite or bradyzoite dataset. Shown are means, error bars represent SD, significance test is pairwise Mann-Whitney, n=6.

Acknowledgements / Funding

We thank Naohiro Hashimoto for sharing the KD3 cells and the Medicine for Malaria Venture for the Pathogen Box and Silvio Bürge for his imaging support.

MB, DM, EP are funded by the Federal Ministry of Education and Research (BMBF) under project number 01KI1715 as part of the "Research Network Zoonotic Infectious Diseases". MB, DM, FS, TH, ML are funded by the Robert Koch Institute.

Supplements

A

B

Figure S1: Longitudinal Screening Procedure

All 400 compounds of the MMV Pathogen Box were screened against (A) tachyzoites (n = 4) and (B) in vitro tissue cysts (n = 3). Plotted fluorescence values were background-subtracted and reflect the abundance of parasites per treatment over time (black line = solvent control, green line = inhibitory compound, red line = ineffective compound, dashed line = growth threshold).

Supplemental Figure 2

A

	tachyzoidal	bradyzoidal	dually active
Cryptosporidium (11)	36	36	27
Dengue (5)	0	20	0
Hookworm (1)	0	0	0
Kinetoplastids (70)	21	11	6
Lymphatic Filariasis (3)	0	0	0
Malaria (125)	13	8	5
Onchoceriasis (11)	9	18	9
Reference Compounds (26)	15	15	8
Schistosomiasis (13)	31	15	15
Toxoplasmamosis (15)	67	0	0
Trichuriasis (1)	0	0	0
Tuberculosis (116)	9	9	4
Wolbachia (3)	33	0	0

B

C

D

Compound	Tachyzoite IC$_{50}$ [µM] Mean ± SEM	HFF lowest cytotoxity [µM]	Bradyzoite LC$_{50}$ [µM] Mean ± SEM	KD3 myotube lowest cytotoxity [µM]
MMV011765	0.14 ± 0.05	5	0.8 ± 0.2	
MMV019189	0.28 ± 0.04		2.7 ± 0.8	
MMV021013	0.28 ± 0.05	10	1.4 ± 0.3	10
MMV022478	1 ± 0.2	5	2.79 ± 0.07	
MMV595321	0.28 ± 0.01		1.3 ± 0.2	
MMV658988	0.28 ± 0.01		3.4 ± 0.2	
MMV659004	0.25 ± 0.02		2.4 ± 0.2	10
MMV671636 (ELQ-400)	0.0059 ±0.0006		0.42 ± 0.04	
MMV675968	0.00105 ± 0.00007		0.9 ± 0.2	
MMV676477	0.034 ± 0.005	0.3	0.53 ± 0.07	5
MMV687807	0.01 ± 0.003	0.08	0.71 ± 0.01	2.5
MMV688762	1.1 ± 0.1		8.2 ± 0.9	
MMV688853	0.061 ± 0.009		1.8 ± 0.2	
MMV688854 (BKI-1294)	0.191 ± 0.007		3.6 ± 0.3	
MMV689480 (BPQ)	0.00007 ± 0.00006		0.15 ± 0.03	
MMV1028806	0.98 ± 0.07		4.4 ± 0.3	

Figure S2: Properties of dually active screening hits.

(A) Table showing the relative amount of tachyzocidal, bradyzocidal and dually-active
compounds per originally indicated target species. E.g. eleven compounds were indicated by
MMV to be active against *Cryptosporidium* and 36% and 27% of these are bradyzocidal and
dually active, respectively. In contrast, 67% of the 15 compounds indicated as active against
Toxoplasma possessed activity against tachyzoites but none were active against bradyzoites.
(B) Estimated lipophilicity approximated by predicted partition coefficient suggest that active
compounds generally exhibit an increased lipophilicity compared to inactive compounds. (C)
A Venn diagram showing intersections of tachyzocidal and bradyzocidal compounds as well
as their predicted gastrointestinal absorption capability (GIA) and blood brain barrier (bbb)
permeability. (D) Tabulation of compound IC_{50} and LC_{50} data of confirmed active compounds.
The calculated IC_{50}s and LC_{50}s of 16 dually active compounds from figure 2 are summarized
in this table. The tachyzoite IC_{50}values represent 8 replicates, the LC_{50} values where generated
from 6 replicate bradyzoite cultures. Lowest cytotoxicity values were recorded during the
primary screen as detailed in the methods section.

Supplemental Figure 3

C

Figure S3: Principal component analysis of LCMS metabolomes of treated
parasites using two chromatographic columns.

(A) Principal component analysis of metabolites from extracellular tachyzoites that were
treated with indicated MMV compounds. Metabolites were separated using BEH-Amide
column -HILIC chromatography and detected by mass spectrometry. (B) The same metabolite
extracts have been re-analyzed using pHILIC-column HILIC chromatography for separation
instead. (C) Chemical structures of dually active and tested compounds as provided by MMV.

	untreated			ATQ			BPQ			MMV			
Glutamine	87	84	87	91	91	91	93	94	93	93	92	92	amino acids & amino acid derivates
Glutamate	49	48	48	15	16	15	15	15	15	35	35	35	
GABA	36	38	39	14	13	13	17	17	15	29	32	29	
Alanine	40	37	38	10	9	9	8	9	8	23	28	22	
Aspartate	7	9	9	12	15	14	14	15	15	12	13	14	
Proline	10	11	11	4	4	5	4	4	5	6	5	5	
Oxoproline	17	12	17	2	2	2	2	1	2	10	11	10	
NAcAspGlu	53	42	44	4	3	7	1	7	6	36	33	34	
Leucine	1	1	0	2	2	1	1	1	1	2	1	1	
Isoleucine	2	1	2	3	3	4	2	2	2	2	2	3	
Glycine	3	2	2	0	0	1	1	0	0	1	0	1	
CarbAsp	2	1	25	93	92	93	95	96	96	94	94	94	pyrimidine metabolism
Dihydroorotate	47	49	42	94	93	93	96	96	96	95	95	95	
Uracil	66	67	65	33	33	33	33	34	33	24	24	24	
Uridine	69	69	68	37	38	38	40	38	39	33	33	33	
UDPGlucNAc	96	98	97	95	96	97	96	97	97	96	96	96	
Desoxyuridine	48	47	47	17	17	16	17	17	16	35	37	9	
Cytosine	78	75	80	76	77	77	76	76	75	78	78	78	
Cytidine	83	83	83	74	74	76	76	76	75	78	77	78	
Desoxycytidine	18	14	14	21	21	20	19	15	28	18	19	17	
Xanthosine	54	56	57	57	57	58	49	48	46	56	55	56	purine metabolism
Inosine	60	60	60	70	68	67	61	58	60	62	62	58	
Hypoxanthine	59	56	59	68	67	69	62	62	62	62	62	62	
Guanosine	85	86	86	70	70	70	66	66	66	76	76	75	
Guanine	89	84	86	60	61	62	64	65	65	77	76	77	
cAMP	59	59	60	69	68	69	60	58	60	63	62	64	
AMP	52	55	55	70	58	77	54	48	51	55	59	55	
Adenine	60	55	63	63	61	61	56	56	54	59	63	60	
Adenosine	65	65	65	69	69	69	62	62	61	65	65	64	
dInosine	54	52	47	32	34	32	43	42	42	40	40	39	
Nicotinamide	71	72	77	33	28	30	29	24	27	54	56	51	misc.
NAD+	96	95	95	70	72	69	67	68	69	79	80	80	
GlucNAcP	90	86	89	78	78	76	85	81	77	79	81	83	

0 %   50 %   100 %

Figure S4: Intracellular tachyzoites exhibit differential ^{15}N incorporation after
treatment with mETC inhibitors.
RHΔku80 were treated with atovaquone, buparvaquone and MMV1028806, labeled with ^{15}N-
amide-L-glutamine for 3h, followed by their isolation from the HFF cells. The heatmap shows
the overall isotopic labeling (100-"M+0") of each replicate.

Figure S5: Direct mitochondrial inhibitors reduce Mitotracker signal intensity
and MT/DAPI ratio in *T. gondii* tachyzoites.
HFF cells were infected with RH-S9 tachyzoites and treated for 24h with DMSO, atovaquone
(ATQ), buparvaquone (BPQ), MMV1028806 (MMV), pyrimethamine (Pyri), clindamycin (Clin),
and 6-Diazo-5-oxonorleucine (DON). * and **** denote $p < 0.05$ and $p < 0.000$ in Mann-Whitney
tests).
910

A

Figure S6: bc_1-complex inhibitors partially reduce mitochondrial profiles of bradyzoites.

Thin section electron microscopy of cysts that were treated with atovaquone (ATQ), buparvaquone (BPQ) or MMV1028806 (MMV) for 24 h (A). (B) Measured areas of

mitochondrial profiles from 21, 12, 15 and 26 images showing DMSO, ATQ, BPQ and

MMV1028806 treated parasites (* denotes $p < 0.05$ in Mann-Whitney tests).

Mitotracker signal appears red, DAPI stain is blue, and the mitochondrial signal originates from
GFP. Scale bars represent 10μm. The graph is showing the ratio of the Mitotracker to DAPI
signal intensity within each individual parasitophorous vacuole (DMSO n = 61, ATQ n = 32,
BPQ n = 35, MMV1028806 n = 36; Pyri n = 58, Clin n = 53, DON n = 68. Blue lines represent
the median, two-sided Mann-Whitney test).
924
925

Figure S7: Principal component analysis of bradyzoites treated with atovaquone (ATQ), Buparvaquone (BPQ) and MMV1028806. Four-week-old intracellular cysts were exposed for three hours, purified from their host cells using magnetic beads and analyzed by LCMS.

Figure S8: Luminescence-based ATP assay

(A) The linear response of the BacTiter-Glo™ assay has a certain range, depending on the ATP concentration of the sample. The assay is reliable between 100 fM and 1 nM ATP. (B) Increasing numbers of freshly isolated parasites were analyzed to determine a minimal parasite density necessary for steady readouts. 10^5 parasites per sample were both manageable and accurate.

References

Alday, P. Holland, and Joseph Stone Doggett. 2017. "Drugs in Development for
Toxoplasmosis: Advances, Challenges, and Current Status." *Drug Design,*
Development and Therapy 11 (January): 273–93.
Alday, P. Holland, Aaron Nilsen, and J. Stone Doggett. 2022. "Structure-Activity
Relationships of Toxoplasma Gondii Cytochrome Bc1 Inhibitors." *Expert Opinion on*
Drug Discovery 17 (9): 997–1011.
Allman, Erik L., Heather J. Painter, Jasmeet Samra, Manuela Carrasquilla, and Manuel
Llinás. 2016. "Metabolomic Profiling of the Malaria Box Reveals Antimalarial Target
Pathways." *Antimicrobial Agents and Chemotherapy* 60 (11): 6635–49.
Anglada-Girotto, Miquel, Gabriel Handschin, Karin Ortmayr, Adrian I. Campos, Ludovic Gillet,
Pablo Manfredi, Claire V. Mulholland, et al. 2022. "Combining CRISPRi and
Metabolomics for Functional Annotation of Compound Libraries." *Nature Chemical*
Biology 18 (5): 482–91.
Antonova-Koch, Yevgeniya, Stephan Meister, Matthew Abraham, Madeline R. Luth, Sabine
Ottilie, Amanda K. Lukens, Tomoyo Sakata-Kato, et al. 2018. "Open-Source
Discovery of Chemical Leads for next-Generation Chemoprotective Antimalarials."
Science 362 (6419). https://doi.org/10.1126/science.aat9446.
Araujo, F. G., J. Huskinson, and J. S. Remington. 1991. "Remarkable in Vitro and in Vivo
Activities of the Hydroxynaphthoquinone 566C80 against Tachyzoites and Tissue
Cysts of Toxoplasma Gondii." *Antimicrobial Agents and Chemotherapy* 35 (2): 293–
99.
Bajohr, Lara Liv, Ling Ma, Christian Platte, Oliver Liesenfeld, Lutz F. Tietze, Uwe Gross, and
Wolfgang Bohne. 2010. "In Vitro and in Vivo Activities of 1-Hydroxy-2-Alkyl-
4(1H)Quinolone Derivatives against Toxoplasma Gondii." *Antimicrobial Agents and*
Chemotherapy 54 (1): 517–21.
Baumeister, Stefan, Jochen Wiesner, Armin Reichenberg, Martin Hintz, Sven Bietz, Omar S.
Harb, David S. Roos, et al. 2011. "Fosmidomycin Uptake into Plasmodium and
Babesia-Infected Erythrocytes Is Facilitated by Parasite-Induced New Permeability
Pathways." *PloS One* 6 (5): e19334.
Berry, Sarah L., Hamza Hameed, Anna Thomason, Marissa L. Maciej-Hulme, Somaia Saif
Abou-Akkada, Paul Horrocks, and Helen P. Price. 2018. "Development of NanoLuc-
PEST Expressing Leishmania Mexicana as a New Drug Discovery Tool for Axenic-
and Intramacrophage-Based Assays." *PLoS Neglected Tropical Diseases* 12 (7):
e0006639.
Borba-Santos, Luana Pereira, Taissa Vila, and Sonia Rozental. 2020. "Identification of Two
Potential Inhibitors of Sporothrix Brasiliensis and Sporothrix Schenckii in the
Pathogen Box Collection." *PloS One* 15 (10): e0240658.
Campos, Adrian I., and Mattia Zampieri. 2019. "Metabolomics-Driven Exploration of the
Chemical Drug Space to Predict Combination Antimicrobial Therapies." *Molecular*
Cell 74 (6): 1291-1303.e6.
Cantillon, Daire, Aaron Goff, Stuart Taylor, Emad Salehi, Katy Fidler, Simon Stoneham, and
Simon J. Waddell. 2022. "Searching for New Therapeutic Options for the Uncommon
Pathogen Mycobacterium Chimaera: An Open Drug Discovery Approach." *The*
Lancet. Microbe 3 (5): e382–91.
Caro, Pilar, Amar U. Kishan, Erik Norberg, Illana A. Stanley, Bjoern Chapuy, Scott B. Ficarro,
Klaudia Polak, et al. 2012. "Metabolic Signatures Uncover Distinct Targets in
Molecular Subsets of Diffuse Large B Cell Lymphoma." *Cancer Cell* 22 (4): 547–60.
Cerneckis, Jonas, Qi Cui, Chuan He, Chengqi Yi, and Yanhong Shi. 2022. "Decoding
Pseudouridine: An Emerging Target for Therapeutic Development." *Trends in*
Pharmacological Sciences 43 (6): 522–35.
Christiansen, Céline, Deborah Maus, Ellen Hoppenz, Mateo Murillo-León, Tobias Hoffmann,
Jana Scholz, Florian Melerowicz, Tobias Steinfeldt, Frank Seeber, and Martin Blume.

2022. "In Vitro Maturation of Toxoplasma Gondii Bradyzoites in Human Myotubes and Their Metabolomic Characterization." *Nature Communications* 13 (1): 1168.

Creek, Darren J., Hwa H. Chua, Simon A. Cobbold, Brunda Nijagal, James I. MacRae, Benjamin K. Dickerman, Paul R. Gilson, Stuart A. Ralph, and Malcolm J. McConville. 2016. "Metabolomics-Based Screening of the Malaria Box Reveals Both Novel and Established Mechanisms of Action." *Antimicrobial Agents and Chemotherapy* 60 (11): 6650–63.

Croft, S. L., J. Hogg, W. E. Gutteridge, A. T. Hudson, and A. W. Randall. 1992. "The Activity of Hydroxynaphthoquinones against Leishmania Donovani." *The Journal of Antimicrobial Chemotherapy* 30 (6): 827–32.

Daina, Antoine, Olivier Michielin, and Vincent Zoete. 2017. "SwissADME: A Free Web Tool to Evaluate Pharmacokinetics, Drug-Likeness and Medicinal Chemistry Friendliness of Small Molecules." *Scientific Reports* 7 (March): 42717.

Davis, D. R. 1995. "Stabilization of RNA Stacking by Pseudouridine." *Nucleic Acids Research* 23 (24): 5020–26.

DeRocher, A., C. B. Hagen, J. E. Froehlich, J. E. Feagin, and M. Parsons. 2000. "Analysis of Targeting Sequences Demonstrates That Trafficking to the Toxoplasma Gondii Plastid Branches off the Secretory System." *Journal of Cell Science* 113 (Pt 22) (November): 3969–77.

Doggett, J. Stone, Aaron Nilsen, Isaac Forquer, Keith W. Wegmann, Lorraine Jones-Brando, Robert H. Yolken, Claudia Bordón, et al. 2012. "Endochin-like Quinolones Are Highly Efficacious against Acute and Latent Experimental Toxoplasmosis." *Proceedings of the National Academy of Sciences of the United States of America* 109 (39): 15936–41.

Doggett, J. Stone, Kayode K. Ojo, Erkang Fan, Dustin J. Maly, and Wesley C. Van Voorhis. 2014. "Bumped Kinase Inhibitor 1294 Treats Established Toxoplasma Gondii Infection." *Antimicrobial Agents and Chemotherapy* 58 (6): 3547–49.

Doggett, J. Stone, Tracey Schultz, Alyssa J. Miller, Igor Bruzual, Sovitj Pou, Rolf Winter, Rozalia Dodean, et al. 2020. "Orally Bioavailable Endochin-Like Quinolone Carbonate Ester Prodrug Reduces Toxoplasma Gondii Brain Cysts." *Antimicrobial Agents and Chemotherapy* 64 (9). https://doi.org/10.1128/AAC.00535-20.

Duffy, Sandra, Melissa L. Sykes, Amy J. Jones, Todd B. Shelper, Moana Simpson, Rebecca Lang, Sally-Ann Poulsen, Brad E. Sleebs, and Vicky M. Avery. 2017. "Screening the Medicines for Malaria Venture Pathogen Box across Multiple Pathogens Reclassifies Starting Points for Open-Source Drug Discovery." *Antimicrobial Agents and Chemotherapy* 61 (9). https://doi.org/10.1128/AAC.00379-17.

Enshaeieh, Marjan, Geita Saadatnia, Jalal Babaie, Majid Golkar, Samira Choopani, and Mohammad Sayyah. 2021. "Valproic Acid Inhibits Chronic Toxoplasma Infection and Associated Brain Inflammation in Mice." *Antimicrobial Agents and Chemotherapy* 65 (10): e0100321.

Ferguson, D. J., J. Huskinson-Mark, F. G. Araujo, and J. S. Remington. 1994. "An Ultrastructural Study of the Effect of Treatment with Atovaquone in Brains of Mice Chronically Infected with the ME49 Strain of Toxoplasma Gondii." *International Journal of Experimental Pathology* 75 (2): 111–16.

Fu, Xiaorong, Stanisław Deja, Blanka Kucejova, Joao A. G. Duarte, Jeffrey G. McDonald, and Shawn C. Burgess. 2019. "Targeted Determination of Tissue Energy Status by LC-MS/MS." *Analytical Chemistry* 91 (9): 5881–87.

Fu, Yong, Kevin M. Brown, Nathaniel G. Jones, Silvia Nj Moreno, and L. David Sibley. 2021. "Toxoplasma Bradyzoites Exhibit Physiological Plasticity of Calcium and Energy Stores Controlling Motility and Egress." *ELife* 10 (December). https://doi.org/10.7554/eLife.73011.

Garfoot, Andrew L., Gary M. Wilson, Joshua J. Coon, and Laura J. Knoll. 2019. "Proteomic and Transcriptomic Analyses of Early and Late-Chronic Toxoplasma Gondii Infection Shows Novel and Stage Specific Transcripts." *BMC Genomics* 20 (1): 859.

Geils, G. F., C. W. Scott Jr, C. M. Baugh, and Butterworth. 1971. "Treatment of Meningeal Leukemia with Pyrimethamine." *Blood* 38 (2): 131–37.

Guggisberg, Ann M., Jooyoung Park, Rachel L. Edwards, Megan L. Kelly, Dana M. Hodge, Niraj H. Tolia, and Audrey R. Odom. 2014. "A Sugar Phosphatase Regulates the Methylerythritol Phosphate (MEP) Pathway in Malaria Parasites." *Nature Communications* 5 (1): 4467.

Havelaar, Arie H., Martyn D. Kirk, Paul R. Torgerson, Herman J. Gibb, Tine Hald, Robin J. Lake, Nicolas Praet, et al. 2015. "World Health Organization Global Estimates and Regional Comparisons of the Burden of Foodborne Disease in 2010." *PLoS Medicine* 12 (12): e1001923.

Hayward, Jenni A., F. Victor Makota, Daniela Cihalova, Rachel A. Leonard, Esther Rajendran, Soraya M. Zwahlen, Laura Shuttleworth, et al. 2023. "A Screen of Drug-like Molecules Identifies Chemically Diverse Electron Transport Chain Inhibitors in Apicomplexan Parasites." *PLoS Pathogens* 19 (7): e1011517.

Hegewald, Jana, Uwe Gross, and Wolfgang Bohne. 2013. "Identification of Dihydroorotate Dehydrogenase as a Relevant Drug Target for 1-Hydroxyquinolones in Toxoplasma Gondii." *Molecular and Biochemical Parasitology* 190 (1): 6–15.

Hennessey, Kelly M., Ilse C. Rogiers, Han-Wei Shih, Matthew A. Hulverson, Ryan Choi, Molly C. McCloskey, Grant R. Whitman, et al. 2018. "Screening of the Pathogen Box for Inhibitors with Dual Efficacy against Giardia Lamblia and Cryptosporidium Parvum." *PLoS Neglected Tropical Diseases* 12 (8): e0006673.

Hortua Triana, Miryam Andrea, Daniela Cajiao Herrera, Barbara H. Zimmermann, Barbara A. Fox, and David J. Bzik. 2016. "Pyrimidine Pathway-Dependent and -Independent Functions of the Toxoplasma Gondii Mitochondrial Dihydroorotate Dehydrogenase." *Infection and Immunity* 84 (10): 2974–81.

Hudson, A. T., A. W. Randall, M. Fry, C. D. Ginger, B. Hill, V. S. Latter, N. McHardy, and R. B. Williams. 1985. "Novel Anti-Malarial Hydroxynaphthoquinones with Potent Broad Spectrum Anti-Protozoal Activity." *Parasitology* 90 (Pt 1) (February): 45–55.

Hulverson, Matthew A., Igor Bruzual, Erin V. McConnell, Wenlin Huang, Rama S. R. Vidadala, Ryan Choi, Samuel L. M. Arnold, et al. 2019. "Pharmacokinetics and In Vivo Efficacy of Pyrazolopyrimidine, Pyrrolopyrimidine, and 5-Aminopyrazole-4-Carboxamide Bumped Kinase Inhibitors against Toxoplasmosis." *The Journal of Infectious Diseases* 219 (9): 1464–73.

Ineichen, Benjamin V., Serena Di Palma, Endre Laczko, Shane A. Liddelow, Susanne Neumann, Martin E. Schwab, and Alice C. Mosberger. 2020. "Regional Differences in Penetration of the Protein Stabilizer Trimethoprim (TMP) in the Rat Central Nervous System." *Frontiers in Molecular Neuroscience* 13 (September): 167.

John, Beena, Tajie H. Harris, Elia D. Tait, Emma H. Wilson, Beth Gregg, Lai Guan Ng, Paulus Mrass, et al. 2009. "Dynamic Imaging of CD8+ T Cells and Dendritic Cells during Infection with Toxoplasma Gondii." *PLoS Pathogens* 5 (7): e1000505.

Johnson, Steven M., Ryan C. Murphy, Jennifer A. Geiger, Amy E. DeRocher, Zhongsheng Zhang, Kayode K. Ojo, Eric T. Larson, et al. 2012. "Development of Toxoplasma Gondii Calcium-Dependent Protein Kinase 1 (TgCDPK1) Inhibitors with Potent Anti-Toxoplasma Activity." *Journal of Medicinal Chemistry* 55 (5): 2416–26.

Katlama, C., S. De Wit, E. O'Doherty, M. Van Glabeke, and N. Clumeck. 1996. "Pyrimethamine-Clindamycin vs. Pyrimethamine-Sulfadiazine as Acute and Long-Term Therapy for Toxoplasmic Encephalitis in Patients with AIDS." *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* 22 (2): 268–75.

Katlama, C., B. Mouthon, D. Gourdon, D. Lapierre, and F. Rousseau. 1996. "Atovaquone as Long-Term Suppressive Therapy for Toxoplasmic Encephalitis in Patients with AIDS and Multiple Drug Intolerance. Atovaquone Expanded Access Group." *AIDS* 10 (10): 1107–12.

Kim, Haeun, Brianne J. Burkinshaw, Linh G. Lam, Kevin Manera, and Tao G. Dong. 2021. "Identification of Small Molecule Inhibitors of the Pathogen Box against Vibrio Cholerae." *Microbiology Spectrum* 9 (3): e0073921.

Konstantinovic, Neda, Hélène Guegan, Tijana Stäjner, Sorya Belaz, and Florence Robert-Gangneux. 2019. "Treatment of Toxoplasmosis: Current Options and Future

1107   Perspectives." *Food and Waterborne Parasitology* 15 (June): e00036.

1108 Kovacs, J. A. 1992. "Efficacy of Atovaquone in Treatment of Toxoplasmosis in Patients with
1109   AIDS." *The Lancet* 340 (8820): 637–38.

1110 Kuby, S. A., L. Noda, and H. A. Lardy. 1954. "Adenosinetriphosphate-Creatine
1111   Transphosphorylase. I. Isolation of the Crystalline Enzyme from Rabbit Muscle." *The*
1112   *Journal of Biological Chemistry* 209 (1): 191–201.

1113 Laue, Michael. 2010. "Electron Microscopy of Viruses." *Methods in Cell Biology* 96: 1–20.

1114 Li, Yaqiong, Zhipeng Niu, Jichao Yang, Xuke Yang, Yukun Chen, Yingying Li, Xiaohan Liang,
1115   et al. 2023. "Rapid Metabolic Reprogramming Mediated by the AMP-Activated Protein
1116   Kinase during the Lytic Cycle of Toxoplasma Gondii." *Nature Communications* 14 (1):
1117   422.

1118 Lin, Sheng-Cai, and D. Grahame Hardie. 2018. "AMPK: Sensing Glucose as Well as Cellular
1119   Energy Status." *Cell Metabolism* 27 (2): 299–313.

1120 Maclean, Andrew E., Hannah R. Bridges, Mariana F. Silva, Shujing Ding, Jana Ovciarikova,
1121   Judy Hirst, and Lilach Sheiner. 2021. "Complexome Profile of Toxoplasma Gondii
1122   Mitochondria Identifies Divergent Subunits of Respiratory Chain Complexes Including
1123   New Subunits of Cytochrome Bc1 Complex." *PLoS Pathogens* 17 (3): e1009301.

1124 MacRae, James I., Lilach Sheiner, Amsha Nahid, Christopher Tonkin, Boris Striepen, and
1125   Malcolm J. McConville. 2012. "Mitochondrial Metabolism of Glucose and Glutamine Is
1126   Required for Intracellular Growth of Toxoplasma Gondii." *Cell Host & Microbe* 12 (5):
1127   682–92.

1128 Maus, Deborah, Blake Curtis, David Warschkau, Estefanía Delgado Betancourt, Frank
1129   Seeber, and Martin Blume. 2024. "Generation of Mature Toxoplasma Gondii
1130   Bradyzoites in Human Immortalized Myogenic KD3 Cells." *Bio-Protocol* 14 (1): e4916.

1131 McConnell, Erin V., Igor Bruzual, Sovitj Pou, Rolf Winter, Rozalia A. Dodean, Martin J.
1132   Smilkstein, Alina Krollenbrock, et al. 2018. "Targeted Structure-Activity Analysis of
1133   Endochin-like Quinolones Reveals Potent Qi and Qo Site Inhibitors of Toxoplasma
1134   Gondii and Plasmodium Falciparum Cytochrome Bc1 and Identifies ELQ-400 as a
1135   Remarkably Effective Compound against Acute Experimental Toxoplasmosis." *ACS*
1136   *Infectious Diseases* 4 (11): 1574–84.

1137 McFadden, D. C., S. Tomavo, E. A. Berry, and J. C. Boothroyd. 2000. "Characterization of
1138   Cytochrome b from Toxoplasma Gondii and Q(o) Domain Mutations as a Mechanism
1139   of Atovaquone-Resistance." *Molecular and Biochemical Parasitology* 108 (1): 1–12.

1140 McHardy, N., L. S. Wekesa, A. T. Hudson, and A. W. Randall. 1985. "Antitheilerial Activity of
1141   BW720C (Buparvaquone): A Comparison with Parvaquone." *Research in Veterinary*
1142   *Science* 39 (1): 29–33.

1143 Meneceur, Pascale, Marie-Anne Bouldouyre, Dominique Aubert, Isabelle Villena, Jean
1144   Menotti, Virginie Sauvage, Jean-François Garin, and Francis Derouin. 2008. "In Vitro
1145   Susceptibility of Various Genotypic Strains of Toxoplasma Gondii to Pyrimethamine,
1146   Sulfadiazine, and Atovaquone." *Antimicrobial Agents and Chemotherapy* 52 (4):
1147   1269–77.

1148 Metsalu, Tauno, and Jaak Vilo. 2015. "ClustVis: A Web Tool for Visualizing Clustering of
1149   Multivariate Data Using Principal Component Analysis and Heatmap." *Nucleic Acids*
1150   *Research* 43 (W1): W566-70.

1151 Montazeri, Mahbobeh, Saeed Mehrzadi, Mehdi Sharif, Shahabeddin Sarvi, Shayesteh
1152   Shahdin, and Ahmad Daryani. 2018. "Activities of Anti-Toxoplasma Drugs and
1153   Compounds against Tissue Cysts in the Last Three Decades (1987 to 2017), a
1154   Systematic Review." *Parasitology Research* 117 (10): 3045–57.

1155 Montazeri, Mahbobeh, Tahereh Mikaeili Galeh, Mahmood Moosazadeh, Shahabeddin Sarvi,
1156   Samira Dodangeh, Javad Javidnia, Mehdi Sharif, and Ahmad Daryani. 2020. "The
1157   Global Serological Prevalence of Toxoplasma Gondii in Felids during the Last Five
1158   Decades (1967-2017): A Systematic Review and Meta-Analysis." *Parasites & Vectors*
1159   13 (1): 82.

1160 Montoya, J. G., and O. Liesenfeld. 2004. "Toxoplasmosis." *The Lancet* 363 (9425): 1965–76.

1161 Müller, Joachim, Adriana Aguado-Martinez, Vera Manser, Vreni Balmer, Pablo Winzer,
1162   Dominic Ritler, Isabel Hostettler, David Arranz-Solís, Luis Ortega-Mora, and Andrew

Hemphill. 2015. "Buparvaquone Is Active against Neospora Caninum in Vitro and in Experimentally Infected Mice." *International Journal for Parasitology, Drugs and Drug Resistance* 5 (1): 16–25.

Nair, Sethu C., Carrie F. Brooks, Christopher D. Goodman, Angelika Sturm, Geoffrey I. McFadden, Sandeep Sundriyal, Justin L. Anglin, Yongcheng Song, Silvia N. J. Moreno, and Boris Striepen. 2012. "Apicoplast Isoprenoid Precursor Synthesis and the Molecular Basis of Fosmidomycin Resistance in Toxoplasma Gondii." *The Journal of Experimental Medicine* 209 (5): 1051–1051.

Nixon, G. L., D. M. Moss, A. E. Shone, D. G. Lalloo, and N. Fisher. 2013. "O, Neill, PM; Ward, SA; Biagini, GA." *The Journal of Antimicrobial Chemotherapy*.

Nugraha, Arifin Budiman, Bumduuren Tuvshintulga, Azirwan Guswanto, Dickson Stuart Tayebwa, Mohamed Abdo Rizk, Sambuu Gantuya, Gaber El-Saber Batiha, et al. 2019. "Screening the Medicines for Malaria Venture Pathogen Box against Piroplasm Parasites." *International Journal for Parasitology, Drugs and Drug Resistance* 10 (August): 84–90.

Pan, Ming, Ceng-Ceng Ge, Yi-Min Fan, Qi-Wang Jin, Bang Shen, and Si-Yang Huang. 2022. "The Determinants Regulating Toxoplasma Gondii Bradyzoite Development." *Frontiers in Microbiology* 13 (November): 1027073.

Perry, S. V. 1954. "Creatine Phosphokinase and the Enzymic and Contractile Properties of the Isolated Myofibril." *Biochemical Journal* 57 (3): 427–34.

Rollin-Pinheiro, Rodrigo, Luana Pereira Borba-Santos, Mariana Ingrid Dutra da Silva Xisto, Yuri de Castro-Almeida, Victor Pereira Rochetti, Sonia Rozental, and Eliana Barreto-Bergter. 2021. "Identification of Promising Antifungal Drugs against Scedosporium and Lomentospora Species after Screening of Pathogen Box Library." *Journal of Fungi (Basel, Switzerland)* 7 (10). https://doi.org/10.3390/jof7100803.

Saleh, Ahmad, Johannes Friesen, Stefan Baumeister, Uwe Gross, and Wolfgang Bohne. 2007. "Growth Inhibition of Toxoplasma Gondii and Plasmodium Falciparum by Nanomolar Concentrations of 1-Hydroxy-2-Dodecyl-4(1H)Quinolone, a High-Affinity Inhibitor of Alternative (Type II) NADH Dehydrogenases." *Antimicrobial Agents and Chemotherapy* 51 (4): 1217–22.

Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair, Tobias Pietzsch, Stephan Preibisch, et al. 2012. "Fiji: An Open-Source Platform for Biological-Image Analysis." *Nature Methods* 9 (7): 676–82.

Shaker, Bilal, Myeong-Sang Yu, Jin Sook Song, Sunjoo Ahn, Jae Yong Ryu, Kwang-Seok Oh, and Dokyun Na. 2020. "LightBBB: Computational Prediction Model of Blood–Brain-Barrier Penetration Based on LightGBM." *Bioinformatics* 37 (8): 1135–39.

Sharifiyazdi, Hassan, Fatemeh Namazi, Ahmad Oryan, Reza Shahriari, and Mostafa Razavi. 2012. "Point Mutations in the Theileria Annulata Cytochrome b Gene Is Associated with Buparvaquone Treatment Failure." *Veterinary Parasitology* 187 (3–4): 431–35.

Shiomi, K., T. Kiyono, K. Okamura, M. Uezumi, Y. Goto, S. Yasumoto, S. Shimizu, and N. Hashimoto. 2011. "CDK4 and Cyclin D1 Allow Human Myogenic Cells to Recapture Growth Property without Compromising Differentiation Potential." *Gene Therapy* 18 (9): 857–66.

Shubar, Hend M., Sabrina Lachenmaier, Markus M. Heimesaat, Uwe Lohman, Rachmat Mauludin, Rainer H. Mueller, Rudolf Fitzner, Klaus Borner, and Oliver Liesenfeld. 2011. "SDS-Coated Atovaquone Nanosuspensions Show Improved Therapeutic Efficacy against Experimental Acquired and Reactivated Toxoplasmosis by Improving Passage of Gastrointestinal and Blood-Brain Barriers." *Journal of Drug Targeting* 19 (2): 114–24.

Shukla, Anurag, Kellen L. Olszewski, Manuel Llinás, Leah M. Rommereim, Barbara A. Fox, David J. Bzik, Dong Xia, et al. 2018. "Glycolysis Is Important for Optimal Asexual Growth and Formation of Mature Tissue Cysts by Toxoplasma Gondii." *International Journal for Parasitology* 48 (12): 955–68.

Smith, David, Geetha Kannan, Isabelle Coppens, Fengrong Wang, Hoa Mai Nguyen, Aude Cerutti, Einar B. Olafsson, et al. 2021. "Toxoplasma TgATG9 Is Critical for Autophagy and Long-Term Persistence in Tissue Cysts." *ELife* 10 (April).

https://doi.org/10.7554/eLife.59384.

Song, Zehua, Bogdan I. Iorga, Pierre Mounkoro, Nicholas Fisher, and Brigitte Meunier. 2018. "The Antimalarial Compound ELQ-400 Is an Unusual Inhibitor of the Bc1 Complex, Targeting Both Qo and Qi Sites." *FEBS Letters* 592 (8): 1346–56.

Songsungthong, Warangkhana, Sunisa Prasopporn, Louise Bohan, Potjanee Srimanote, Ubolsree Leartsakulpanich, and Suganya Yongkiettrakul. 2021. "A Novel Bicyclic 2,4-Diaminopyrimidine Inhibitor of Streptococcus Suis Dihydrofolate Reductase." *PeerJ* 9 (February): e10743.

Songsungthong, Warangkhana, Suganya Yongkiettrakul, Louise E. Bohan, Eric S. Nicholson, Sunisa Prasopporn, Pimchai Chaiyen, and Ubolsree Leartsakulpanich. 2019. "Diaminoquinazoline MMV675968 from Pathogen Box Inhibits Acinetobacter Baumannii Growth through Targeting of Dihydrofolate Reductase." *Scientific Reports* 9 (1): 15625.

Spalenka, Jérémy, Sandie Escotte-Binet, Ali Bakiri, Jane Hubert, Jean-Hugues Renault, Frédéric Velard, Simon Duchateau, Dominique Aubert, Antoine Huguenin, and Isabelle Villena. 2018. "Discovery of New Inhibitors of Toxoplasma Gondii via the Pathogen Box." *Antimicrobial Agents and Chemotherapy* 62 (2). https://doi.org/10.1128/AAC.01640-17.

Sun, Lijun, Jiaxi Wu, Fenghe Du, Xiang Chen, and Zhijian J. Chen. 2013. "Cyclic GMP-AMP Synthase Is a Cytosolic DNA Sensor That Activates the Type I Interferon Pathway." *Science* 339 (6121): 786–91.

Thomsen-Zieger, Nadine, Joachim Schachtner, and Frank Seeber. 2003. "Apicomplexan Parasites Contain a Single Lipoic Acid Synthase Located in the Plastid." *FEBS Letters* 547 (1–3): 80–86.

Vallières, Cindy, Nicholas Fisher, Thomas Antoine, Mohammed Al-Helal, Paul Stocks, Neil G. Berry, Alexandre S. Lawrenson, et al. 2012. "HDQ, a Potent Inhibitor of Plasmodium Falciparum Proliferation, Binds to the Quinone Reduction Site of the Cytochrome Bc1 Complex." *Antimicrobial Agents and Chemotherapy* 56 (7): 3739–47.

Veale, Clinton G. L. 2019. "Unpacking the Pathogen Box-an Open Source Tool for Fighting Neglected Tropical Disease." *ChemMedChem* 14 (4): 386–453.

Vercesi, A. E., C. O. Rodrigues, S. A. Uyemura, L. Zhong, and S. N. Moreno. 1998. "Respiration and Oxidative Phosphorylation in the Apicomplexan Parasite Toxoplasma Gondii." *The Journal of Biological Chemistry* 273 (47): 31040–47.

Vila, Taissa, and Jose L. Lopez-Ribot. 2017. "Screening the Pathogen Box for Identification of Candida Albicans Biofilm Inhibitors." *Antimicrobial Agents and Chemotherapy* 61 (1). https://doi.org/10.1128/AAC.02006-16.

Wang, Jin-Lei, Si-Yang Huang, Michael S. Behnke, Kai Chen, Bang Shen, and Xing-Quan Zhu. 2016. "The Past, Present, and Future of Genetic Manipulation in Toxoplasma Gondii." *Trends in Parasitology* 32 (7): 542–53.

Wang, Peiyan, Siji Li, Yingchi Zhao, Baohuan Zhang, Yunfei Li, Shengde Liu, Hongqiang Du, et al. 2019. "The GRA15 Protein from Toxoplasma Gondii Enhances Host Defense Responses by Activating the Interferon Stimulator STING." *The Journal of Biological Chemistry* 294 (45): 16494–508.

Weiss, Louis M., and Jitender P. Dubey. 2009. "Toxoplasmosis: A History of Clinical Observations." *International Journal for Parasitology* 39 (8): 895–901.

Winder, W. W., and D. G. Hardie. 1999. "AMP-Activated Protein Kinase, a Metabolic Master Switch: Possible Roles in Type 2 Diabetes." *The American Journal of Physiology* 277 (1): E1-10.

Winzer, Pablo, Joachim Müller, Adriana Aguado-Martínez, Mahbubur Rahman, Vreni Balmer, Vera Manser, Luis Miguel Ortega-Mora, et al. 2015. "In Vitro and In Vivo Effects of the Bumped Kinase Inhibitor 1294 in the Related Cyst-Forming Apicomplexans Toxoplasma Gondii and Neospora Caninum." *Antimicrobial Agents and Chemotherapy* 59 (10): 6361–74.

Zampieri, Mattia, Michael Zimmermann, Manfred Claassen, and Uwe Sauer. 2017. "Nontargeted Metabolomics Reveals the Multilevel Response to Antibiotic

1275   Perturbations." *Cell Reports* 19 (6): 1214–28.
1276

1277