Meteorins regulate the formation of the left-right organizer and the

establishment of vertebrate body asymmetry

Fanny Eggeler[1], Jonathan Boulanger-Weill[1#], Flavia De Santis[2,3#], Laura Belleri[1#], Karine

Duroure[1], Thomas O. Auer[2,4], Shahad Albadri[1*], Filippo Del Bene[1,5*]

[1] Sorbonne Université, INSERM, CNRS, Institut de la Vision, F-75012 Paris, France

[2] Institut Curie, PSL Research University, Inserm U934, CNRS UMR3215, Paris, France

[3] ZeClinics SL. Sant Feliu de Llobregat, Barcelona, Spain.

[4] Department of Biology, University of Fribourg, Fribourg, Switzerland

[5] Lead contact

[#] These authors contributed equally

* correspondence: shahad.albadri@inserm.fr; filippo.del-bene@inserm.fr

SUMMARY

While the exterior of vertebrate bodies appears bilaterally symmetrical, internal organ
positioning and morphology frequently exhibit left-right (L-R) asymmetries. In several
vertebrates, including human, mouse, frog and zebrafish, left-right symmetry-breaking during
embryonic development is initiated by a ciliated organ called the Node or left-right organizer.
Within the Node, a leftward flow of extraembryonic fluid named the Nodal flow mediates the
asymmetric expressions of Nodal factors. Although downstream Nodal pathway components
leading to the establishment of the embryonic left-right axis are well known, less is known
about the development and formation of the embryonic Node itself.

Here we reveal a novel role for the Meteorin protein family in the establishment of the left-right
axis and in the formation of the Kupffer's vesicle, the Node equivalent structure in zebrafish.
We show that the genetic inactivation of each or all three members of the zebrafish Meteorin
family (*metrn*, *metrn-like a* and *metrn-like b*) leads to defects in properties of the Kupffer's
vesicle, caused by impaired assembly and migration of the Kupffer's vesicle forming dorsal
forerunner cells. In addition, we demonstrate that Meteorins genetically interact with integrins
ItgαV and Itgβ1b regulating the dorsal forerunner cell clustering and that *meteorins* loss-of-
function results in disturbed Nodal factor expression and consequently in randomized or
symmetric heart looping and jogging.

These results identify a new role for the Meteorin protein family in the left-right asymmetry
patterning during embryonic vertebrate development.

INTRODUCTION

From the outside the vertebrate body plan appears bilaterally symmetric. However, internal organs positioning and morphology often displays left-right (L-R) asymmetries. For instance, in vertebrates the heart generally lies on the left side, the liver and the pancreas are positioned on the right and the gut presents asymmetric rotations.

One major regulator for the establishment of the L-R axis is Nodal, a ligand belonging to the TGFβ protein family [1–6]. Its signaling is activated in the left lateral plate mesoderm (LPM), whereas Nodal remains inactive in the right LPM, creating an embryo-scale left-right asymmetry [3]. In mice, this asymmetry is achieved by an oriented rotation of the cilia of a structure called the Node generating a leftward flow of extraembryonic fluid named the Nodal flow. This results in an asymmetric expression of several Nodal factors and Nodal signaling. Participating in this patterning, Leftys, soluble inhibitors belonging to a subclass of TGF factors, antagonize Nodal signaling [7,8].

The Node and the Nodal flow have been described in numerous vertebrates. For instance, in *Xenopus* the gastrocoel roof plate and in zebrafish the Kupffer's vesicle (KV) [9–14] are key structures of the L-R symmetry breaking during the embryonic development [11,13,15]. Similarly, in humans and other mammals like rabbits and mice, the systematic establishment of left-right body symmetry begins with asymmetric fluid flow driven by rotating cilia [16]. This occurs within the transient primitive node in humans or the embryonic node in mice and rabbits [13,15,17]. However, in other mammalian classes, although left-right symmetry breaking mechanisms are believed to be cilia-independent, they are still poorly understood at present [2,18]. In chick, the symmetry breaking structure known as the Hensen's node, serves as the ciliated organizer responsible for the left-right asymmetry establishment. In this species, the process of symmetry breaking is believed to occur independently of the non-motile cilia present within the Hensen's node. Instead, the breaking of symmetry is primarily attributed to the leftward movement of cells around the node [19–21].

Compared to the relatively flat-shaped mouse node, the zebrafish KV is a fluid-filled sphere with a ciliated epithelium and it is formed by dorsal forerunner cells (DFCs) [9]. DFCs first emerge as cells of the epithelium of the dorsal surface in direct contact with the yolk syncytial layer at 6 hours post-fertilization (hpf) [22]. Around 8 to 9 hpf, following the epibolic movement, 20 to 30 DFCs form a single cluster and migrate towards the vegetal pole in a non-involuting manner at the leading edge [22,23]. During this migration process, polarized DFCs organize into multiple focal points by retaining long-lasting apical contacts within the cluster and remaining delaminated DFCs maintain contact with the cluster by cell-cell contact mechanisms [22,24]. At the final phase of epiboly and after reaching the vegetal pole, the DFC cluster is separating from the dorsal marginal enveloping layer cells by losing the apical contacts [22,24].

The focal points of the cluster are then rearranged by integrating the unpolarized DFCs to form
a single focal point that will expand into a monolayer rosette structure containing a lumen to
form the KV by 12 hpf [9,22–24]. Along this process, non-motile and motile monocilia are
generated on the apical membrane facing the lumen that create a counterclockwise flow of
fluid by the KV, called Nodal flow [22,25]. It was shown that the leftward directed flow within
the KV results in asymmetric expression of zebrafish Nodal genes, like *spaw*, *lefty1* and *lefty2*
that is fundamental for proper L-R patterning [26,27]. However, the molecular mechanisms
involved in correct DFC migration and clustering that will form the KV remain largely elusive.
In this context, it has been reported that *nodal, lefty* as well as *p-smad2* expression are
downregulated in *Meteorin*-null ES-cells. As such, Meteorin (Metrn) was hypothesized to be a
novel important regulator of Nodal transcription [28].
Meteorin (Metrn), a secreted neurotrophic factor highly conserved among vertebrates, is
expressed during early mouse development. It is already detected at the blastocyst stage in
the inner cell mass and then expands through the extraembryonic ectoderm to the central
(CNS) and peripheral nervous system (PNS) during later developmental stages [28]. It was
first described to induce glial cell differentiation and to promote axonal extension in dorsal root
ganglion (DRG) explants *in vitro* [29]. The disruption of *Metrn* function in Mouse resulted in
early embryonic lethality [28] preventing the investigation of Metrn proteins function during
early embryonic development *in vivo* and in particular their role for L-R patterning. Also
conserved among vertebrates, its paralog Meteorin-like (Metrnl) has been first reported as a
downstream target of the Pax2/5/8 signaling pathway during otic vesicle development [30] and
was later shown to have neurotrophic properties comparable to the ones of Metrn [31].
Here, using the zebrafish larva as a model system, we generated knockout lines for all three-
existing zebrafish *metrn* genes (*metrn*, *metrnla* and *metrnlb*). While all three single and
compound mutants are viable, *metrns* mutant embryos displayed organ patterning defects
notably with randomized or symmetric heart looping at a significantly higher frequency. Like in
mice, *metrn* genes are expressed during early development already from the 2-cells stage and
transcripts are detected in DFCs as well as in the KV structure. We show that *metrns* loss of
function leads to DFC disorganization and Nodal flow formation defects within the KV.
Together our study reveals a critical role for Metrn proteins in the DFC clustering, KV formation
and in the establishment of the L-R axis.
111

RESULTS

Metrns loss-of-function affects proper heart looping and correct visceral organs
positioning

To study the functions of Meteorin proteins during early development, we generated knock out
zebrafish lines using the CRISPR/Cas9 technology to target the second exon of each gene
that encodes for the signal peptide sequence. The newly generated zebrafish mutant lines
carried out-of-frame deletions in the coding sequence of *metrn*, *metrnla* and *metrnlb* (Fig. S1A).
In contrast to published embryonic lethal *Metrn* mouse mutants [28], constitutive homozygote
zebrafish mutants for all three genes individually (*metrn*$^{-/-}$; *metrnla*$^{-/-}$; *metrnlb*$^{-/-}$), in double
(*metrn*$^{-/-}$, *metrnla*$^{-/-}$; *metrn*$^{-/-}$, *metrnlb*$^{-/-}$; *metrnla*$^{-/-}$, *metrnlb*$^{-/-}$) and triple (*metrn*$^{-/-}$, *metrnla*$^{-/-}$,
metrnlb$^{-/-}$ or triplMut), were all viable and did not show any obvious morphological defects (Fig.
1A).

We first validated that our mutagenic approach resulted in the generation of null alleles. To do
so, we performed *in situ* hybridization on *metrn*, *metrnla* and *metrnlb* from 2-cells stage to 1
day post-fertilization (dpf) on triplMut embryos (Fig. S1B). No expression could be detected for
neither of those two genes in the triplMut embryos, indicating the degradation of the transcripts
in the *metrn* mutant backgrounds (Fig. S1B). To confirm this observation, we compared the
expression level of all three *metrn* genes upon total mRNA extraction from 14 and 48 hpf WT
and triplMut embryos by qRT-PCR. *Metrn* and *metrnla* expression was detected in samples of
14 hpf WT embryos and *metrnlb* expression was detected in samples of 48 hpf WT embryos.
A significant reduction of the expression of all *metrn* genes could be measured using triplMut
samples of the same age. These results suggest that the mRNA produced by the loci that we
targeted is degraded by nonsense-mediated decay (Fig. S1C).

Despite the absence of gross morphological defects, using *myl7* antisense riboprobe *in situ*
hybridization in single mutant embryos, we observed a mis-positioning of the heart in triplMut
mutant embryos. Indeed while 88.43% of the wild type control embryos had D-looped shaped
hearts, only 38.17% of the triplMut embryos exhibited a WT-like heart looping phenotype
(Table 1 and Fig. 1B-C). Similarly, *metrn*$^{-/-}$ and *metrnla*$^{-/-}$ single mutant embryos displayed a
high proportion of randomized heart phenotypes, whereas *metrnlb*$^{-/-}$ single mutant embryos
showed a small percentage of heart looping phenotypes (Table 1 and Fig. 1B-C). The injection
of *metrn* and/or *metrnla* mRNA into one-cell stage triplMut zebrafish embryos could partially
rescue the observed phenotypes (Table 1, Fig. S1D). In addition to the heart, morphological
L–R asymmetry of the visceral organs was also perturbed in triplMut embryos (using *gata6*
expression as a marker). Indeed, 10.49% of the triplMut embryos exhibited symmetrical
placement of the gut and pancreas with respect to the midline and 9.88% heterotaxy
phenotypes compared to wild type embryos (WT: 93.84% L–R asymmetry, 4.79% L–R

symmetry and 1.37% heterotaxia) (Fig. 1D-E). These results indicate that while Metrns are not

implicated in the specification of neither the heart nor the visceral organs (*myl7* and *gata6*

expression being detected in triplMut embryos and in single mutants for *metrns*), their loss of

function reveal that they are implicated in the L-R asymmetry positioning of these organs. The

left-jogged and D-looped heart or the right-positioned liver are two of many examples of early

L-R asymmetry determination in zebrafish and are well-studied readouts for altered L-R

patterning processes. Therefore, our results suggest that Metrns, in particular Metrn and

Metrnla, may have a crucial function in establishing L-R asymmetry during the early stages of

development. To exclude potential genetic compensation effects among paralogs that were

reported in other studies [32], we focused all subsequent analyses (if not stated otherwise) on

the triple mutant line (triplMut).

Metrns are required for proper Nodal factor gene expression

Nodal signaling is one of the major regulatory pathways for the determination of the L-R axis

during vertebrate development. In zebrafish, null mutants for several Nodal factors like *spaw*

(*Nodal* zebrafish orthologue), *dand5* or *lefty1* (*lft1*) all display symmetric or randomized heart

looping and jogging similarly to *Nodal* mutant mice [33–36]. We therefore asked whether

Metrns loss of function could affect the expression of these factors. To do so, we assessed

dand5, *spaw*, *lft1* and *lft2* expression in our *metrns* mutant embryos. By qRT-PCR, we

measured a significant reduction of *dand5* expression at 14 hpf in triplMut embryos compared

to wild type controls (Fig. 2A). Similarly, the expression of *spaw*, *lft1* as well as *lft2* in the

absence of Metrns proteins was also downregulated in triplMut embryos compared to wild type

embryos (Fig. 2A).

In order to assess whether Metrns loss of function altered the expression patterns of *spaw*, *lft1*

and *lft2*, we performed *in situ* hybridization experiments for all three genes at 16 hpf (Fig. 2B).

At 16 hpf *spaw* is normally expressed adjacent to the KV and in the left LPM. In *metrn*[-/-] and

metrnla[-/-] single mutants as well as triplMut embryos, we could not detect any *spaw* expression

in the left LPM and only faintly around the KV (Fig. 2B). The spaw inhibitor *lft1* at this stage is

expressed along the midline. In all analyzed *metrn* mutants (single *metrn* and *metrnla* and

triplMut), its expression was severely disrupted at the midline (Fig. 2B, middle panel). The

same was observed for *lft2* which expression in the left heart field was either absent or

randomized in *metrn* mutants, single or triplMut embryos, compared to the wild type embryos

(Fig. 2B, lower panel).

We next characterized *dand5* specific R > L bias expression pattern in triplMut embryos by *in*

situ hybridization. In more than 61% triplMut embryos, we could detect particularly reduced

dand5 expression at 14 hpf and 23% displayed an abnormal *dand5* R > L expression bias (Fig.

2 C-D). In single *metrn*[-/-] and *metrnla*[-/-] mutant embryos, *dand5* expression pattern was also

altered (Fig. S2). These results are in line with previous findings reporting that *dand5*
expression in the KV is only altered if upstream symmetry breaking mechanisms are disrupted
[37–39]. These results indicate that Metrns are required for the proper expression of Nodal
factor genes that are determinants of proper L-R patterning.
190
***Metrns* are expressed during early zebrafish development**
To investigate how Metrns affect the expression of Nodal factors during early development,
we analyzed the pattern of expression of all three zebrafish *metrn* genes from 2 cell stage to
48 hpf. We first investigated the expression pattern of *metrn*, *metrnla* and *metrnlb* by *in situ*
hybridization and hybridization chain reaction (HCR) experiments. At the 2-cells stage, we
detected both *metrn* and *metrnla* maternally expressed transcripts, whereas *metrnlb*
expression was not detected (Fig. 3A, Fig. S3A). During gastrulation, the expression of *metrn*
and *metrnla* could be detected, whereas *metrnlb* expression was again not observed in our *in*
situ hybridization analysis. The earliest time point for zygotic expression of *metrn* and *metrnla*
was at 6 hpf (shield stage). At this stage, *metrnla* expression could be detected over the whole
animal cap whereas zygotic *metrn* expression onset was observed only around the yolk
syncytial layer (YSL) at this developmental stage (Fig. 3A). *Metrn* expression was then
restricted to the developing LPM and to the leading edge of the former shield area from 9 hpf
(Fig. 3A, Fig. S3B). In comparison, *metrnla* was found expressed throughout the whole
enveloping layer and developing midline (Fig. 3A, Fig. S3B). At 12 hpf, the expression of *metrn*
was restricted to the KV area while *metrnla* transcripts were mainly found in the midline (Fig.
3A-B). By 14 hpf, *metrn* expression was present in the KV area as well as in the developing
brain whereas *metrnla* transcripts were mainly found in the midline and the KV area (Fig. S3B).
In contrast, from the 2-cell stage to 14 hpf no *metrnlb* transcripts could be detected (Fig. 3A,
Fig. S3A-B). By HCR, we found *metrn* and *metrnla* expression colocalizing in the area of the
KV at 12 and 14 hpf (Fig. 3B). By 24 and 48 hpf, both genes were found strongly expressed in
the central nervous system (CNS) (Fig. S3C). Finally, *metrnlb* expression onset started from
24 hpf broadly in the embryonic brain (Fig. S3C). At 48 hpf, *metrnlb* transcripts were mostly
found in the otic vesicles and at the somite boundaries (Fig. S3C).
We next analyzed the relative expression of *metrn* genes to landmarks of the KV structure like
sox32 a well-studied DFC marker [40,41]. We observed a complete colocalization of
metrn/*sox32* and *metrnla*/*sox32* signals at 10 hpf, indicating that *metrns* genes are expressed
in DFCs, the cells that will later form the KV (Fig. 3C-D). To assess whether *metrn* and *metrnla*
expression in the KV structure is conserved among vertebrates, we performed *in situ*
hybridization for *metrn* and *metrnla* in the chick embryo at early stages. Around stage HH6, we
could observe *metrn* and *metrnla* expression at the Hensen's node and around the primitive
streak (marked by primitive streak landmark *fgf8*) (Fig. S3D). Remarkably, the chick *metrn* and

metrnla expression patterns were reminiscent of the ones that we observed in zebrafish,
suggesting that *metrn* genes expression at early developmental stages in embryonic Node and
midline structures is conserved throughout vertebrates.
226

***Metrns* loss-of-function leads to DFC disorganization and migration defects**
The specific expression of *metrn* and *metrnla* in DFCs and in the KV area at early
developmental stages led us to ask what role they may play for these cells and the KV
formation and function. To assess whether the loss of Metrn proteins function could affected
the migratory and clustering properties of DFCs, we labeled these cells in 9 hpf (80-90%
epiboly) gastrulating mutant embryos using the DFC landmarks *sox32* [40,41]*, tbxta* [42] or
sox17 [40]. Compared to wild type embryos displaying ovoid DFC clustering, about 65% of the
triplMut embryos displayed multiple segregated DFC clusters or abnormal linear cellular
distributions as shown using *sox32* marker in 9 and 10 hpf triplMut embryos (Fig. 4A-B, Fig.
S4B, Table 2). Similarly, *metrn*[-/-] as well as *metrnla*[-/-] single mutant embryos presented
clustering defects of *sox32*-expressing DFC compared to wild type embryos (Fig. 4B, Fig. S4A-
B, Table 2). In single or triplMut[+/-] heterozygous embryos, the DFC clustering was instead not
significantly or very mildly affected as revealed (Fig. S4B, Table 2). The observed defects were
also present when using *tbxta* and *sox17* as DFC markers whose expression was detected at
8 and 10 hpf respectively, indicating that although disorganized, the DFCs identity was not
altered by the absence of *metrns* expression (Fig. S4C). Together, these results indicate that
Metrns are required for the proper clustering of DFCs.
244

It was reported that proper adherent cell assembly is crucial for DFCs directional collective cell
migration during their animal to vegetal pole (AP to VP) journey [43] (Fig. 4C). We therefore
evaluated the migration capacity of DFCs during early development in the absence of Metrn
proteins. To do so, we first labeled *sox32*-expressing DFCs by *in situ* hybridization in wild type
and triplMut embryos and measured the position of *sox32*-expressing DFCs along the AP-VP
axis divided by the total embryo length at 6 hpf, 8 hpf and 10 hpf. Compared to wild type
embryos, DFCs in triplMut embryos showed migratory delays of about 8% to 13% at all
analyzed developmental stages (Fig. S5B-C, Table 3). By live imaging, we tracked the *in vivo*
migration of *sox17*:GFP-labeled DFCs both in triplMut and wild type embryos from 50% epiboly
to 90% epiboly (from 6 hpf) (Fig. 4D, Fig. S5D , Video S1-2). In line with the DFC clustering
abnormalities in fixed triplMut embryos that we observed, the tracking of these cells in triplMut
embryos *versus* wild type control embryos revealed a diminished convergence rate for triplMut
DFCs (Fig. 4E, upper panel). Furthermore, we measured a significant decrease of their
migration speed compared to the wild type DFCs (Fig. 4E, lower panel). Intriguingly,
individually tracked DFCs exhibited a consistent movement toward the vegetal pole, regardless

of the absence of Metrns proteins (Fig. 4D, Fig. S5D, Video S1-2). Taken together, these

findings indicate that the absence of Metrn proteins does not affect DFC migration directionality

but alters instead their migratory abilities including migration speed and convergence rate.

It was shown that during the process of epiboly, initial EVL cells destined to develop into DFCs

exhibit an accumulation of ZO-1 along their apical junctions. Subsequently, their apical surface

gradually diminishes in size, resulting in the formation of discrete apical domains enriched with

ZO-1 [22,24]. DFCs as such originate from dorsal EVL cells through a mechanism involving

the delamination of epithelial cells, facilitated by apical constriction. We asked whether the

observed DFC organization and migration disruptions in the absence of Meteorin proteins

might be attributed to perturbations in both ZO-1 accumulation and apical constriction

disruption. To explore this hypothesis, we examined the spatial arrangement of these regions

enriched with ZO-1 in triplMut embryos and WT controls during epiboly at 6 hpf and 8 hpf. As

expected, immunohistochemistry against ZO-1 on 6 and 8 hpf *Tg*(*sox17:GFP*) wild type control

embryos where DFCs are labeled with GFP were enriched in ZO-1 at their apical side and

apical junctions (Fig. 4F-G, Fig. S5E). At this same developmental stage, a reduced ZO-1

enrichment at the apical junctions of triplMut GFP-positive DFCs could be detected (Fig. 4F,

Fig. S5E left plot). Similarly at 8 hpf, GFP-positive triplMut DFCs exhibited diminished ZO-1 at

their apical region (Fig. 4G, Sy Fig. 5E right plot). Furthermore, compared to wild type DFCs,

triplMut GFP-positive DFCs appear to have lost their apical attachments to the EVL (Fig. 4F-

G, right panels). Indeed, the distance measured between triplMut DFCs and the EVL/YSL at 6

and 8 hpf was significantly higher compared to the one of wild type DFCs (6 hpf: *p-value:

0.039, N = 4 WT, 30 DFCs measured, triplMut N = 6, 46 DFCs measured; 8 hpf: *p-value:

0.021, N = 7 WT embryos, 47 DFCs measured, N = 7 triplMut, 54 DFCs measured; Fig. S5F).

Together, our results demonstrate that triplMut DFCs migrate significantly slower and

convergence less compared to wild type DFCs. Furthermore, in the absence of Metrns, DFCs

loose apical ZO1-enrichment and attachment to the EVL/YSL, indicating that Metrn proteins

are important for proper DFC clustering, migration along the AP-VP axis and their loss affect

ZO-1 accumulation for the connection to the EVL/YSL.

***Metrns* loss-of-function impairs the KV formation and function**

At the end of gastrulation, DFCs that migrated to deeper layers of the developing zebrafish

embryo form a lumen to shape the KV, a ciliated organ transiently present (from around 10.5

hpf to 30 hpf [44]) that is crucial for initiating L-R asymmetric patterning [9,23]. As we observed

at 14 hpf, it was reported that Dand5, a member of Cerberus/Dan family of secreted factors, is

expressed adjacent to the KV with a right – left (R > L) bias between 11 and 14 hpf in zebrafish

(Fig. 2) [45]. The right-biased expression of *dand5* was shown to be directly influenced by the

direction and the strength of the Nodal flow generated by the KV [46]. In return, Dand5

contributes to the left-biased expression of *spaw* in the LPM by antagonizing Spaw activity
[45]. The leftward-directed flow generated by the KV was also shown to be important for the
asymmetric expression of other Nodal genes, like *spaw*, *lefty1* (*lft1*) and *lefty2* (*lft2*) [26,27].
The observed disturbed expression of these Nodal factors in the absence of Metrns led us to
hypothesize that Metrn proteins might not only be critical for the proper clustering and migration
of DFCs but also for the proper formation and function of the KV. To assess this question, we
first labeled the cilia present in the KV of wild type and triplMut embryos at 14 hpf by anti-
acetylated tubulin immunostaining (Fig. 5A). We then measured the number of cells in the KV,
the KV diameter and the number of cilia, which were correlating with the number of cells, of
both wild type and triplMut embryos (Fig. 5B-D). Previous work has shown that the KV diameter
and the number of KV cilia cells as well as KV cilia can be highly variable from embryo to
embryo [47]. Nevertheless, despite the wide variability, we observed highly significant
reduction in all cases (Fig. 5B-D). In addition, compared to the wild type controls, triplMut
embryos also displayed a significant reduction in the KV cilia length (Fig. 5E).
Furthermore, we investigated both the KV lumen and the process of DFC-to-KV organization.
As described recently and observed in ZO-1 stained embryos, by 14 hpf DFCs clustering into
a rosette structure lead tight junctions to form the KV lumen [22] (Fig. S6A, left panel). In
contrast, ZO-1 immunostained triplMut embryos of the same developmental stage displayed
an irregular ZO-1 lattice pattern and disrupted KV lumen shape (Fig. S6A, right panel).
Additionally, unlike in control embryos, triplMut embryos displayed only partially polarized
DFCs indicated by absent or irregular aPKC-ζ immunolabeling (Fig. S6B). These results
indicate that the loss of Metrns function also impacts the proper DFC polarization and
subsequent KV formation.
320
We next evaluated whether the deficient formation of the lumen as well as the reduced number
and cilia length in triplMut altered the KV function. To do so, we compared the Nodal flow
generated by the KV of triplMut embryos to the one of wild type embryos. The movement
generated by the KV cilia was monitored by injecting fluorescent microspheres (0.5 μm) into
the KV at 12-14 hpf. To quantify the flow inside the KV, fluorescent microbeads movement was
tracked in WT control (n = 363 beads in 4 embryos) and in triplMut embryos (n = 318 beads in
4 embryos) (Fig. 5F). Beads movement analysis within the KV showed a counter-clockwise
rotation of the flow in wild type control (Fig. 5F, Video S3). In contrast, this movement was
severely impaired in triplMut embryos (Fig. 5F, Video S4). To quantify the properties of the
beads displacement, we computed the mean square displacement (MSD), which enables to
distingue between directed and confined motion. This analysis revealed a directed motion of
microbeads in the wild type KV (MSD quadratic dependence on Δt) and a confined motion
(MSD asymptotic behavior) in triplMut [48]. Accordingly, measurement of average bead

velocity showed an overall reduction in the beads speed in the triplMut mutant embryos
compared to the wild types (WT mean = 0.96212 µm/sec; TriplMut mean = 0,47996 µm/sec;
Fig. S6C). These results demonstrate that the observed defects in the KV formation in the
absence of Metrns also severely impacts its function and thereby the Nodal flow generation,
responsible for proper Nodal factors expression.
339

Genetic interaction of ItgαV/Itgβ1b and Metrns on the level of the DFCs
In order to further uncover the mechanisms through which Metrn proteins act in the
establishment of early L-R patterning, we finally assessed the possible relationship between
Metrns and mediators of the extracellular matrix like Integrin αV1 (ItgαV1) and Integrin β1b
(Itgβ1b). Integrins are membrane receptors involved in cell adhesion and recognition during
embryogenesis amongst others [49]. During gastrulation, *itgαV* as well as *itgβ1b* were shown
to be expressed in DFCs and *itgαV* and *itgβ1b* morpholino-injected embryos display DFC
phenotypic defects reminiscent to those observed in *metrn* mutants [50].
We therefore asked whether Metrns loss of function could affect the expression of both
integrins. To do so, we quantified *itgαV* and *itgβ1b* expression levels in wild type and triplMut
embryos at 6, 9 and 24 hpf by qRT-PCR. No significant change in *itgαV* and *itgβ1b* expression
levels at either tested developmental stage was measured in triplMut embryos compared to
wild type controls (Fig. S7A).
In order to test whether ItgaV1 and Itgβ1b and Metrns genetically interact, we injected
insufficient doses of morpholinos for *itgβ1b* (0.5 ng) or *itgαV1* (0.41 ng) into 1 to 4 -cell stage
in a heterozygous *metrns* mutant background. The injection of the morpholinos at these doses
and at sufficient doses (*itgβ1b:* 0.7 ng, *itgαV1:* 1.25 ng) reproduced the previously reported
defects on DFCs (Fig. 6A, Table 4) [50]. Instead, the injection of low doses of either *itgβ1b* or
itgαV1 morpholino did not affect DFC clustering, similarly to non-injected triplMut[+/-] embryos
(Fig. 6A, Table 4). However, a significantly higher percentage of embryos with DFC clustering
defects could be observed upon the injection of either morpholino at low dose into triplMut[+/-]
(Fig. 6A, Table 4). These results are consistent with the hypothesis that Metrn proteins act
together with both Integrins for the proper clustering and migration of DFCs during early
embryonic development without directly affecting their expression.
364

DISCUSSION

Proteins of the Meteorin family were first described as secreted neurotrophic factors highly conserved among vertebrates [28–31]. In mouse *Metrnla* mutants display phenotypes as impaired angiogenesis after ischemic injury [51] or dysregulated cytokine production [52]. The disruption of *Metrn* resulted in early embryonic lethality [28] preventing to investigate the combined role of Metrn proteins in early embryonic development *in vivo* and in particular its role in L-R patterning. Here, using the zebrafish larva as model system, we generated knockout lines for all three zebrafish *metrn* genes, *metrn*, *metrnla* and *metrnlb*, which are all viable and fertile allowing for the study of Meteorins functions during early embryonic development.

Our results demonstrate that Metrns are required for symmetry breaking and the establishment of L-R asymmetry, a novel role for Meteorin proteins during vertebrate embryonic development. All three paralogs analyzed, when genetically deleted, had an effect on heart looping formation, with the strongest effect for Metrnla and Metrn while Metrnlb had a smaller but still significant effect. In our *in situ* analysis, we failed to detect early *metrnlb* expression. We do not exclude that at early stages, *metrnlb* could be expressed at very low level or in a very small number of cells that we cannot detect. However, in the *metrnlb* mutant, the loss of this expression could still be sufficient to induce the low penetrant phenotype. Interestingly, when analyzing the expression of these genes in a single cell transcriptomic atlas at the earliest available time point (10 hpf), in contrast to the robust expression of the two first genes, *metrnlb* expression is indeed found in very few cells [53].

We show that Metrns play a crucial role in ensuring this function the genetic interaction with the integrins ItgαV1 and Itgβ1b known to facilitate the proper clustering and migration of DFCs based on the sufficient accumulation of ZO-1 and apical constriction, essential to form the KV [22,50]. Furthermore, our results reveal that Metrns are essential for several aspects of the KV formation and function including the cilia length, cilia number, KV size and lumen formation. Consequently, Metrns are important for the regulation of the proper expression of Nodal flow genes and that this regulation is critical for the proper asymmetric positioning and morphology of vital organs like the heart (Fig. 6B).

Previous work has shown that zebrafish midline mutants display altered *lft1* and *lft2* expression patterns as well as discordance among heart, gut and brain gene expression patterns [54]. From 9 hpf *metrn* expression was visible around the developing LPM close to the midline and the leading edge of the shield. *Metrnla* expression was found throughout the whole enveloping layer and the developing midline which plays an important role in vertebrate L-R development. Despite the expression of both paralog close to or at the midline, we hypothesize that the described body asymmetry defects in triplMut embryos do not originate from impaired midline

structures but rather impaired DFC clustering and migration. Indeed, triplMut[-/-] embryos display
intact *tbxta*-positive notochords (Fig. S4C), as well as randomized visceral organs positioning
(Fig. 1D-E), phenotypes that are not characteristic for defects of the anterior midline [54].
DFCs are precursors of the zebrafish KV, organ of laterality, essential for the L-R axis
establishment. DFCs develop at the beginning of gastrulation and form the KV by the end of
gastrulation [9,23]. Given the early onset of *metrn* and *metrnla* expression and their specific
expression within the DFCs, we investigated their role for the clustering and migration capacity
of DFCs during these early embryonic developmental steps. In the course of their migration,
DFC progenitors intercalate mediolaterally and results in an oval-rosette shape cluster of DFCs
by mid-gastrulation that will form the KV [22]. *Metrn[-/-], metrnla[-/-]* and triplMut zebrafish embryos
displayed DFC assembly defects and delayed migration of these cells during gastrulation.
Since DFCs in triplMut still express DFC specific markers (*sox17, sox32, tbxta*), we concluded
that Metrns are not required for the specification of DFCs *per se* but rather that Metrns are
important for their proper clustering and migration.
416
It is important to note that the observed DFC clustering and migration abnormalities in Metrn
mutants primarily affect the apico-ventral axis rather than the mediolateral axis. This is evident
from the disrupted connection between DFCs and the YSL/EVL in these mutants, while the
migration directionality remains intact. Since a proper connection between DFCs and the EVL
facilitates a subset of DFCs to establish apical contacts with the EVL and become pre-
polarized, one could speculate that Metrns are involved in the polarization and establishment
of direct apical contacts of DFCs with the EVL [22–24,55]. This disrupted connection in *metrns*
loss-of-function embryos, which usually links DFCs with the vegetal spread of extra-embryonic
tissues [24], likely results in the observed DFCs vegetal motion defects.
It has been demonstrated that these apical attachments function as tissue connectors, linking
DFCs with the spreading of extra-embryonic tissues towards the vegetal pole [24].
Therefore, the loss of apical attachments in the mutant embryos causes all DFCs to converge
less and migrate more slowly than in wild-type controls, as shown by our *in vivo* imaging
analysis.
This would imply that a number of DFCs was simply too slow to reach the vegetal pole in time
to form a fully functional KV. Given that KV cilia are monocilia, this subsequently results in a
diminished count of KV cells (Fig. S5A), along with the observed structural abnormalities in the
KV lumen and impairments in its overall functionality.
Additionally it was shown that the formation of the developing KV lumen was directly connected
to ciliogenesis as polarized DFCs initiate developing cilia at their membrane domain facing the
forming lumen [22]. This indicates that proper DFC polarization is essential for the KV
formation and for ciliogenesis. In *metrns* mutants the reduced cilia length and the lack of PKCζ

staining in the developing KV further highlights an impairment of DFC polarization in the
absence of Metrns consequently leading to KV formation and function defects. Further work
would be needed to explore whether Metrns directly impact the motility of KV cilia or if the
disrupted Nodal flow observed is solely caused by the disturbed KV lumen.

It has been demonstrated that an appropriate asymmetrical Nodal flow plays a crucial role in
the asymmetrical expression of Nodal genes such as *dand5, spaw, lefty1 (lft1),* and *lefty2 (lft2)*
[26,27,45]. As such, the disrupted formation and function of the KV in *metrns* mutants might
contribute to the evident perturbation in Nodal factor expression and may also induce
randomization of Nodal factor expression in L-R asymmetry [9,15,47,50]. We do not exclude
however that Metrns directly impact the expression of *spaw, leftys* and *dand5* in addition to
their role in DFC cell behaviors and KV formation. In this context, several studies have reported
that Nodal signaling was shown to regulate DFC specification [56,57].
In line with the impaired KV formation and function that we observed in *metrns* mutants, it was
shown that knockdown of Integrin αV as well as Integrin β1b in zebrafish results in L-R
asymmetry defects [50]. This was explained by DFC organization and migration defects and
disturbed KV formation. Integrin αVβ1 specific functions have not yet been completely resolved
in mammals, mostly due to their broad expression and redundant function with other integrins,
as well as the lack of antibodies and inhibitors specific for αVβ1. However, several *in vitro*
studies have already connected αVβ1 to developmental processes. For instance, αVβ1 was
shown to be implicated in embryonic astrocytes and oligodendrocyte precursors migration in
rodents [58] and neural cell adhesion molecule L1 cell binding [59]. Additionally, the fibronectin
receptor integrin α5β1 has been found to be crucial for regulating extracellular matrix assembly
along tissue boundaries, in coordination with Eph/Ephrin signaling [60].
Our results suggest that one mechanism by which Metrns act is through their genetic
interaction with Integrin αV/β1b to mediate the proper clustering of DFCs. These interactions
could influence Integrin's adhesive functions (or other properties) necessary for DFC
clustering. If so, the absence of Meteorins would lead to disturbed adhesion and/or Integrin
signaling, resulting in DFC migration defects and disturbed KV development. Interestingly, it
was shown that in αV and β1b morphants DFC-EVL connections remain intact [50], unlike in
our *meteorin* mutants. This contrast suggests that Meteorins alone are crucial for DFC
polarization and that their interaction with integrins αVβ1b may rather be important for the
subsequent steps leading to KV formation.

Despite the recent advancements in uncovering the multiple functions of Meteorins, their
receptors during early development are still unknown. It was recently shown that METRNL is

promoting heart repair through its binding to the KIT tyrosine kinase receptor, establishing
METRNL as a KIT receptor ligand in the context of ischemic tissue repair [51]. In another study,
it was reported that Metrn is a ligand for the 5-hydroxytryptamine receptor 2b (Htr2b) [61].
Additionally Metrn was associated with the regulation of reactive oxygen species levels in
hematopoietic stem/progenitor cells via directing phospholipase C signaling [61]. When we
analyzed the expression of *kita*, one of the two zebrafish orthologs to KIT tyrosine kinase
receptor, at 9 hpf we found that *kita* is solely expressed in the prechordal plate and later (from
11 hpf) in the lateral borders of the anterior neural plate (Fig. S7B, left panel). Its paralog *kitb*
is expressed from 11 hpf in the anterior ventral mesoderm (Fig. S7B, right panel), as previously
reported [62]. Additionally, the analysis of *htr2b* expression during zebrafish development
revealed that its expression only starts from around 2 dpf in the heart (Fig. S7C). The absence
of *kita/b* as well as *htr2b* expression at early embryonic stages in or at proximity of DFCs and
in the forming KV suggests that Metrns rather act through other receptor(s) to ensure their L-
R patterning functions.
489
Mechanisms underlying L-R patterning are known to be highly conserved within vertebrates.
Hence, we raised the question whether Meteorin protein family is conserved across different
species. Genes coding for *metrn* and *metrnl* exist in several vertebrates classes but could not
be found in the genomes of invertebrates [63]. *Metrnl* orthologs genes could instead be found
in amphioxus and tunicates. Therefore, we propose that Metrns function in the establishment
of the L-R axis that we have uncovered here emerged during Chordate evolution (Fig. S7D).
496

MATERIAL AND METHODS

Zebrafish husbandry

Zebrafish (*Danio rerio*) were maintained at 28°C overnight (O/N) at 14 hours light/10 hours dark cycle. Fish were housed in the animal facility of our laboratory which was built according to the respective local animal welfare standards. All animal procedures were performed in accordance with French and European Union animal welfare guidelines with protocols approved by the committee on ethics of animal experimentation of Sorbonne Université (APAFIS#21323- 2019062416186982). The *Tg*(*sox17:GFP*) lines were created via injection of a *pTol2-sox17:GFP* plasmid (kindly gifted from Didier Stanier and Stephanie Woo) into one-cell stage zebrafish embryos together with 150 ng/ml of *tol2* mRNA. Injected embryos were grown to adulthood and screened for GFP in their offspring.

Metrn, *metrnla* and *metrnlb* CRISPR-Cas9-mediated mutagenesis

SgRNA guide sequences were cloned into a BsaI-digested DR274 plasmid vector (Addgene 42250). The sgRNAs were synthesized by *in vitro* transcription using (using the Megascript T7 transcription kit, Ambion, AM1334). After transcription, sgRNAs were purified using a RNAeasy Mini Kit (Qiagen). The quality of purified sgRNAs was checked by electrophoresis on a 2 % agarose gel. The target sequences were the following:

	Sequence (5' - 3')
metrn	CCACCACACCCGGCCCAGCC
metrnla	GGTGTATCTCCGCTGCGCCC
metrnlb	CCAGAAACAGCGCCCCCTCC

Cas9 mRNA was generated as described in Hwang, Fu et al. 2013 [64]. To induce targeted mutagenesis at the *metrn, metrnla* and *metrnlb* loci, 200 ng/ml of sgRNA were injected into one-cell stage zebrafish embryos together with 150 ng/ml of *cas9* mRNA. Injected embryos were grown to adulthood and screened for mutation in their offspring.

DNA extraction and sequencing for analysis of CRISPR/Cas9- mediated mutagenesis

For genomic DNA extraction, pools of 5 dpf embryos were digested for 1 h at 55 °C in 0.5 mL lysis buffer (10 mM Tris, pH 8.0, 10 mM NaCl, 10 mM EDTA, and 2% SDS) with proteinase K (0.17 mg/mL, Roche Diagnostics) and inactivated 10 min at 95 °C. To check for frequency of indel mutations, target genomic loci were PCR amplified using a Phusion High-Fidelity DNA polymerase (Invitrogen). PCR amplicons were subsequently cloned into a *pCR-bluntII-TOPO* vector (Thermo Fisher). Plasmid DNA was isolated from single colonies and Sanger

530   sequencing was performed by Eurofins. Mutant alleles were identified by comparison with the
531   wild type sequence using Geneious software.
532
Mutants genotyping and generation of triple mutants
534   Genotyping of *metrns* mutants was performed as follows: after genomic DNA extraction PCR
535   amplification reactions were conducted in final volumes of 20 µl containing 1x PCR reaction
536   buffer, 1,5mM MgCl2, 70 ng of gDNA, Taq DNA Polymerase (5U/µl) and 0.5 µM of each primer.
537

	Sequence (5' - 3')
metrn-fw	CTGTGTTGACTGCTGGCTG
metrn-rv	GTGGTTTAGTGGTGTTCTTACAATGA
metrnla-fw	TCCCATGCCTGGACCTCATA
metrnla-rv	AGACGGAGAGAAGAGACGCT
metrnlb-fw	TGTTGATCAGCAGTGTGTGCGTAGC
metrnlb-rv	GTCCTCCGCTGATCTACGTG

538
The DNA amplification was performed with 35 cycles at the annealing temperature of 60°C.
Metrn and *metrnlb* amplicons were loaded on 2 % agarose gel to discriminate between wild
type and mutant alleles (for *metrn*, the expected size of wild type amplicon is 598 bp while the
mutant amplicon is 482 bp long using the same primer set; for *metrnlb*, the size of the expected
wild type amplicon is 242 bp while the mutant band is 156 bp long using the same primer set).
Metrnla PCR product was digested at 55°C O/N with the PasI restriction enzyme and the
resulting digestion was subsequently analyzed on a 2% agarose gel. The wild type amplicon
results in two fragments of 107 bp and 94 bp length, respectively. The mutant amplicon band
size is instead 201 bp using the same primer set. Triple mutants were obtained first by
incrossing homozygous *metrn* and *metrnla* mutants to generate double heterozygous mutants.
Double heterozygous mutants were then crossed with homozygous *metrnlb*[-/-] fish and
genotyped to identify triplMut[+/-] triple heterozygous mutants. triplMut[+/-] fish were incrossed and
genotyped to identify triplMut[-/-] triple homozygous mutants. In all described experiments of this
study, triplMut[+/-] were obtained by outcrossing female triplMut[-/-] fish with male wild type fish, if
not stated differently to prevent from *metrns* maternal contribution.
554
Hybridization chain reaction (HCR)
All HCR probes and solutions were purchased from Molecular Instruments®. Dechorionated
embryos at the developmental stage(s) of interest were fixed in 4% paraformaldehyde in PBS
(pH 7.4) for 2 hours at room temperature, followed by several washes with PBS to stop the
fixation. Embryos were then dehydrated with a series of methanol (MeOH) washes and stored

at -20°C. The HCR was performed following the manufacturer's instructions. HCR samples

were imaged on an inverted confocal microscope Olympus FV-1000, employing a 20x oil

immersion objective (NA 0.85). Z-volumes were acquired with a 5 μm resolution and images

were processed and analyzed using ImageJ.

***In situ* hybridization in zebrafish and chicken**

The respective cDNA fragments were amplified by PCR from total zebrafish or chicken cDNA

of stage 12 hpf – 48 hpf and HH6. *In vitro* transcription of Digoxigenin/Fluorescent-labeled

probes was performed using an RNA Labeling Kit (Roche Diagnostics Corporation) according

to manufacturer's instructions. Zebrafish and chicken whole-mount *in situ* hybridizations were

performed as previously described [65] [66]. Stained embryos were then imaged on a Leica

MZ10F stereomicroscope. Images were processed and analyzed using ImageJ software.

Color balance, brightness and contrast were applied uniformly. Fluorescently-labeled samples

were imaged using an inverted confocal microscope Olympus FV-1000, employing a 40x oil

immersion objective (NA 1.30). Z-volumes were acquired with a 0.5 μm resolution and images

were processed and analyzed using ImageJ.

Analysis of DFC phenotype and migration

For the DCF clustering analysis, Bayesian inference was used to recover the maximum-

likelihood value for the malformation probability. The outcome of an experiment was

considered to be a binary random variable (malformed or wild type phenotype) whose

probability depends solely on the experimental condition. The total number of malformed

phenotypes per experiment is therefore binomially distributed. The posterior distribution is

obtained by inverting this binomial distribution using Bayes theorem, under a uniform prior.

Under these hypotheses the maximum-likelihood estimate for the malformation probability is

simply given by the empirical malformation frequency. The standard deviation of the posterior

distribution was used to provide a confidence interval for the estimate (black error bars). *P*-

values were evaluated using Fisher exact test on the number of malformed and wild type

phenotypes. AP to VP DFC migration was calculated as the percentage of the total embryo

length.

For *in vivo* live DFC imaging, a *pEGFP-sox17* construct (Addgene #31400) and *H2B-RFP*

mRNA were co-injected either in wild type or triplMut embryos that were embedded at 50%

epiboly (6 hpf) in 1% low melting-point agarose diluted in E3 water. DFC were identified as

recently described in [24]. Imaging was performed in a temperature constant environment with

a 10x water dipping objective using an upright Yokogawa CSU-X1 spinning disk scan head,

mounted on a DM6000 upright Leica microscope and a CCD CoolSnap HQ2 camera at 4 Hz.

Volumetric image stacks were acquired every 6 minutes which was sufficient to manually track

individual DFC migration during 3 hours from 50% epiboly. No drift correction was required.

Convergence ratio was computed as the ratio between the maximal extent of the tracked cells

along the y axis measured at the onset and at the end of each recording.

601   qRT-PCR and analysis

For gene expression analysis, total RNA extraction was prepared from zebrafish embryos with

TRIzol reagent (Thermo Fisher) and TURBO DNA-free reagents (Invitrogen™ AM1907M).

Total RNA was cleaned up using RNeasy Mini Kit (Qiagen) following the manufacturer's

instructions and treated twice with DNase I (1 U/µg RNA, Qiagen). The RNA concentration

was quantified using the Nanodrop2000 (Thermo Fisher). RNA (1 µg) was retro-transcribed

using random primers and a SuperScript III First-Strand Synthesis system (Invitrogen

18080051). For qRT-PCR, the SYBR Green PCR Master Mix (Thermo Fisher) was used

according to the manufacturer's protocol. *Ef1a* was used as reference genes as previously

reported [67]. All assays were performed in triplicates and repeated in 3 independent

experiments. The mean values of triplicate experiments were calculated according to the delta

CT quantification method [68] and a student T-test was applied for the *p*-value calculation.

614   mRNA synthesis and injection

Zebrafish *metrn* and *metrnla* cDNA fragments were amplified by PCR from zebrafish cDNA of

1 dpf. The fragments were cloned into a *pCS2+* plasmid linearized with Xho1 and Xba1

restriction enzymes using a Quick Ligation Kit (NEB). The *pCS2:h2b-rfp* plasmid readily

available in the laboratory was linearized with the NotI restriction enzyme and the mRNA

synthesis was performed by *in vitro* transcription using the mMESSAGE mMACHINE Sp6 Ultra

kit (AM1340, Ambion) adding 1 µL of GTP as recommended by the manufacturer's instructions

and followed by lithium chloride precipitation. 150 ng.µL^{-1} of the synthesized mRNA was then

injected into 1- to 4-cell-stage zebrafish embryos.

624   Design and injection of antisense morpholino oligonucleotides

The published morpholinos (MO) used in this study for *itgαV* (5'-

AGTGTTTGCCCATGTTTTGAGTCTC-3'), *itgβ1b1* (5'-GGAGCAGCCTTACGTCCATCTT

AAC-3') and for control (ctrl MO) (5'-CCTCTTACCTCAGTTACAATTTATA-3') were injected

using the same reported concentration of 0.41 ng (*itgαV*) and 0.5 ng (*itgβ1b1*) for insufficient

doses, 1.25 ng (*itgαV*) and 0.7 ng (*itgβ1b1*) for full knock down effect and 2.5 ng for the ctrl

MO [50]. All MOs were obtained from Gene Tools (Philomath, OR, USA). MOs were injected

in 1 to 4 -cells stage zebrafish embryos.

Whole-mount immunohistochemistry

Embryos were fixed in 4% paraformaldehyde in PBS-Tween for 2h at room temperature (RT) and subsequently washed three times in PBS-1% Triton X-100 to promote their permeabilization. They were then incubated for 1 hour at RT in blocking solution (10% Normal Goat Serum, in PBS-Tween) followed by overnight incubation, at 4°C with primary antibodies diluted at 1:200 in 1% blocking solution. The following primary antibodies were used: anti-acetylated tubulin (Sigma T6793); anti-ZO1 (339100, Invitrogen, Carlsbad, CA, USA); anti-GFP (GTX13970, GeneTex); anti-aPKCζ (sc-216, Santa Cruz Biotechnology Inc., CA, USA). After five washes in 1x PBS-Tween, the embryos were then incubated O/N with respective Alexa Fluor IgG secondary antibody (Life Technologies) diluted 1:200 in 1% blocking solution. The following day embryos were washed five times in 1x PBS-Tween and mounted in 1% low-melting point agarose in glass-bottom cell tissue culture dish (Fluorodish, World Precision Instruments, USA). An inverted confocal microscope Olympus FV-1000 was used for imaging, employing a 20x objective (NA 0.75) or a 40x oil immersion objective (NA 1.30). Z-volumes were acquired with a 0.5 μm resolution and images were processed and analyzed using ImageJ and Imaris. The distances between the DFCs and the EVL at 6 and 8 hpf were measured in fixed whole-mount samples. Using ImageJ, the distances were analyzed from the center of the leading DFCs to the leading edge of the EVL, as marked by ZO-1. The area of ZO-1 enrichments was measured using Fiji in cropped images that included only DFCs and at 6 hpf also the apical part of the EVL.

Fluorescent microspheres injection and analysis

FluoSpheresH fluorescent microspheres injection was conducted as previously described [69]. Embryos were embedded in 1% low melting-point agarose diluted in egg water. Imaging was performed with a 40x water dipping objective using an upright Yokogawa CSU-X1 spinning disk scan head, mounted on a DM6000 upright Leica microscope and a CCD CoolSnap HQ2 camera at 4 Hz using the green channel at 0.25 μm resolution on a single Z-plane. Beads trajectory tracking was performed using an already developed toolbox [70]. The size detection parameter was set to 0.75 – 1.25 μm objects with a brightness superior to the 0.01 percentile of the whole image. Objects that traveled less than 2.5 μm between two consecutive frames were linked into the same trajectory and considered as a single bead. Objects traveling longer distances were analyzed as separate trajectories, even if they originated from the same bead. Analyses were then performed using a custom-made Matlab code. For better visualization, trajectory plots only show 20 tracked beads. MSD analysis was conducted using an already published code (https://tinevez.github.io/msdanalyzer/) [71]. Average mean square displacement (MSD) plots only show particles tracked between 1 and 8 seconds. Very short

669   tracking durations do not permit to correctly estimate displacement properties and less than
670   1% of the particles were tracked for more than 8 seconds.
671

Data analysis
Experiments for heart looping, *dand5* expression and visceral organ phenotypes have
multinomial outcomes, and the *p*-values for these distributions were evaluated accordingly.
For any given pair of conditions α (control) and β (mutated condition), the number of observed
experimental outcomes were counted. The number of outcomes for each condition was
denoted as (nα, dα, rα) and (nβ, dβ, rβ), with Nα and Nβ denoting the total number of
observations. The frequency of each outcome under a particular condition is simply the number
of observations for the given outcome divided by the total number of observations. The
dissimilarity between two conditions α and β was measured as the sum of absolute differences
between their respective frequencies δ = |fnα - fnβ| + |fdα - fdβ| + |frα - frβ|. Then the probability
of the null hypothesis was estimated, i.e., that the multinomial distributions of outcomes related
to the two conditions are the same, as the probability of obtaining a dissimilarity equal or higher
to the one observed by simply pooling all the observations together, and again randomly
subdivided in two sets of observations of size Nα and Nβ. To numerically estimate this
probability the procedure of pooling the observations together was repeated for K = 100'000
times, randomly separating them again in two sets, and evaluating the new dissimilarity δ'. The
probability of the null hypothesis is then given by the numerical frequency with which the
dissimilarity δ' ≥ δ is greater or equal to the one observed in the experiments. [72,73]
Python and Matlab codes used for beads and DFC tracking are available at
https://github.com/jboulanger91/meteorin. Codes for statistical analysis are available at
https://github.com/FannyEggeler/Meteorin_analysis.
693

Phylogenetic analysis
Protein sequences were obtained from the Ensembl Genome Browser or UniProt. The
following sequences were used: *D. rerio* Metrn (ENSDARP00000041804.5), *D. rerio* Metrnla
(ENSDARP00000131064.1), *D. rerio* Metrnlb (ENSDARP00000126894.1), *G. gallus* Metrn
(ENSGALP00010026416.1), *G. gallus* Metrnl (ENSGALP00010044844.1), *H. sapiens* Metrn
(ENSP00000455068.1), *H. sapiens* Metrnl (ENSP00000315731.6), *M. musculus* Metrn
(ENSMUSP00000127275.2), *M. musculus* Metrnl (ENSMUSP00000038126.8), *X. leavis*
Metrnl (ENSXETP00000107425.1), *L. oculatus* Metrn (ENSLOCP00000003703.1), *L. oculatus*
Metrnl (ENSLOCP00000017224.1), *B. lanceolatum* (A0A8K0F3K3), *S. clava*
(XP_039261436). *L. oculatus* was chosen as an outgroup, *B. lanceolatum* as an example for
Cephalochordata and *S. clava* as a representative for Tunicata. The phylogenetic tree was
reconstructed with the phylogeny analysis from www. phylogenie.fr [74–80].
706

FIGURE LEGENDS

Figure 1

Fig. 1. Metrns loss-of-function causes heart jogging/looping and visceral organ positioning defects. (**A**) 48 hpf triplMut zebrafish display no gross phenotypic defects compared to wild type (WT) embryos. (**B**) Examples of 48 hpf embryos showing mRNA expression of the heart marker *myl7* and the different heart looping phenotypes (ventral view). (**C**) Quantification in percentage of heart looping phenotypes at 48 hpf in WT, triplMut, *metrn*$^{-/-}$, *metrnla*$^{-/-}$, *metrnlb*$^{-/-}$, triplMut$^{+/-}$ embryos; displayed *p*-values compared to WT: ****p-value: <1.0e-5 for triplMut, ****p-value: <1.0e-5 for *metrn*$^{-/-}$, ****p-value: <1.0e-5 for *metrnla*$^{-/-}$, *p-value: 0.003 for *metrnlb*$^{-/-}$, ns *p*-value: 0.49 for triplMut$^{+/-}$, *p*-value compared to *metrnlb*$^{-/-}$: *p-value: 0.034 for triplMut$^{+/-}$ (not displayed). (**D**) Example of 56 hpf embryos showing *gata6* mRNA expression highlighting the different visceral organ positioning phenotypes (ventral view). (**E**) Quantification in percentage of visceral organ positioning phenotypes at 56 hpf in WT and triplMut embryos, ***p-value: 0.00034. In (**B**) L = left, R = right. In (**D**) L = liver, P = pancreas, I = intestine. Scale bars in (**B**): 100 µm, (**D**): 500 µm.

Fig. 2. Metrns are required for proper Nodal factor genes expression. (**A**) qRT-PCR analysis for *dand5*, *spaw, lft1* and *lft2* expression in 14 hpf wild type (WT) and triplMut embryos (Student t-tests, ****p*-value: 0.00019 for *dand5; *p*-value: 0.014 for *spaw*; ****p*-value: 0.0004 for *lft1*; ****p*-value: 0.00019 for *lft2*). (**B**) Dorsal view of 16 hpf triplMut, *metrn*$^{-/-}$ and *metrnla*$^{-/-}$ embryos with reduced expression of *spaw*, *lft1* and *lft2* as revealed by *in situ* hybridization. (**C**) *Dand5* expression at 14 hpf as revealed by *in situ* hybridization showing different transcripts distribution of *dand5* mRNA. (**D**) Quantification in percentage of *dand5* expression phenotypes at 14 hpf in WT and triplMut embryos, *****p*-value: <1.0e-5. Scale bar in (**B**): 250 µm, in (**C**): 100 µm; L= left, R=right.

Figure 3

Fig. 3. *Metrns* are expressed during early zebrafish development. (**A**) Expression patterns of
metrn, metrnla and *metrnlb* during early embryonic development from 2-cell stage to 12 hpf. (9hpf

and 12 hpf dorsal view) (**B**) Confocal cross-section of the midline region of 12 and 14 hpf zebrafish embryo showing mRNA expression of the *metrn* (magenta) and *metrnla* (cyan) and their co-expression in the area of the Kupffer's vesicle (KV, indicated by arrowheads) by HCR. (**C**) *Metrn* and *sox32* are co-expressed by dorsal forerunner cells (DFCs) as shown by double fluorescence *in situ* hybridization against *metrn* (magenta) and *sox32* (cyan) on 10 hpf wild type embryos. (**D**) *Metrnla* and *sox32* are co-expressed by DFCs as shown by double fluorescence *in situ* hybridization against *metrnla* (magenta) and *sox32* (cyan) on 10 hpf wild type embryos. Scale bars in (**A**): 500 µm, (**B**) 250 µm, (**C**) and (**D**): 50 µm. L = left, R = right.

Figure 4

Fig. 4. *Metrns* loss-of-function leads to DFC clustering and migration defects. (**A**) Dorsal views of s*ox32* expression in DFCs at 9 hpf and 10 hpf (cyan) in wild type (WT) and triplMut embryos reveals

DFC misclustering in triplMut embryos. (**B**) Quantification in fraction of embryos with DFC clustering defects at 9 hpf in WT, *metrn*[-/-], *metrnla*[-/-], triplMut and and triplMut[+/-] embryos (Fisher exact test, ***p*-value: 1.6e-07 for WT vs. *metrn*[-/-]; ***p*-value: 3.5e-40 for WT vs. *metrnla*[-/-]; ***p*-value: 1.5e-49 for WT vs. triplMut and ***p*-value: 7.9e-15 for triplMut vs. triplMut[+/-]). (**C**) Lateral view of a schematic representation of an 8 hpf zebrafish embryo visualizing animal pole (AP) to vegetal pole (VP) dorsal forerunner cell (DFC) migration from both poles, adapted from [44]. (**D**) Tracking plots of combined DFC movement in wild type (left) and triplMut embryos (right) (n = 3 embryos per conditions, single embryo traces see Fig. S5D) showing directed DFC movement in both conditions. (**E**) Plots for convergence (upper) and migration speed (lower) analyzed from DFC tracking data show significant decrease in these parameters in triplMut embryos compared to WT controls. (*p-value convergence ratio: 0.0295; ***p*-value for migration speed 1.217e-04) (**F-G**) GFP and ZO-1 immunostainings of the dorsal margin and confocal microscopy ZY-planes (right panels) of *Tg*(*sox17*:GFP) WT and triplMut embryos at (**F**) 6 hpf and (**G**) 8 hpf (shield stage and 75% epiboly) showing the apical domains of marginal DFCs with ZO-1 enriched junction points (arrowheads) and revealing the absence of apical ZO1-enrichment and detachment from the EVl/YSL in triplMut embryos. EVL, enveloping layer; DC, deep cells; YSL, yolk syncytial layer; D, dorsal; V, ventral. Scale bars in (**A**): 50 µm, (**F-G**): 50 µm. AP = animal pole, VP = vegetal pole.

Figure 5

Fig. 5. *Metrns* loss-of-function impairs the Kupffer's vesicle formation and function. (**A**) Confocal cross-section of 14 hpf wild type (WT) and triplMut counterstained with acetylated tubulin

(AcTub) labeling the cilia of the Kupffer's vesicle (KV). (**B**) The quantification of the cell number per KV in WT and triplMut embryos at 14 hpf shows a significant difference. (**C**) The KV diameter size measurement, (**D**) cilia number per KV quantification and (**E**) individual cilia length measurement in WT and triplMut embryos at 14 hpf are all significantly decreased in triplMut (N WT: 23, N triplMut: 26, Student t-tests, ***_p_-value: 0.00028 for KV cell number, ***_p_-value: 9.7e-06 for KV diameter size, **_p_-value: 0.0012 for cilia number. For cilia length: average per KV and ***_p_-value: 5.8e-12). (**F**) Single microbeads tracking in the KV of 12 hpf -14 hpf WT and triplMut embryos and the mean square displacement (MSD), revealing a directed trajectory in WT samples in contrast to triplMut, displaying short and undirected trajectories**.** Scale bar in (**A**): 20 µm.

Fig. 6. Metrns and ItgαV/Itgβ1b act together for the proper clustering and migration of DFCs.
(**A**) Quantification in percentage of embryos with DFC clustering defects at 9 hpf in non-injected wild
type (WT), WT + 2.5 ng control morpholino (WT + ctrl MO), WT + insufficient doses of *itgαV/itgβ1b*
MOs (WT + 0.41 ng *itgαV*; WT + 0.5 ng *itgβ1b*), WT + sufficient *itgαV/itgβ1b* MO doses (WT+ 1.25
ng *itgαV*; WT + 0.7 ng *itgβ1b*), triplMut[+/-] non-injected, triplMut[+/-] + 2.5ng ctrl MO and triplMut[+/-] with
insufficient doses of *itgαV/itgβ1b* MOs (triplMut[+/-] + 0.41 ng *itgαV*; triplMut[+/-] + 0.5 ng *itgβ1b*). Fisher

exact tests indicate *p-value < 0.01, **p-value < 0.001 and ***p-value < 0.0001 and ns = not significant. (**B**) Schematic representation modeling the novel role of Metrns for DFCs assembly and migration, for the proper KV formation and subsequent Nodal factors expression distribution at the base of the correct L-R axis establishment during early development (adapted from [44]).

REFERENCES

[1] Blum, M., Feistel, K., Thumberger, T., and Schweickert, A. (2014) The evolution and conservation of left-right patterning mechanisms. *Development*. 141 (8), 1603–1613.

[2] Blum, M. and Ott, T. (2018) Animal left–right asymmetry. *Current Biology*. 28 (7), R301–R304.

[3] Grimes, D.T. and Burdine, R.D. (2017) Left–Right Patterning: Breaking Symmetry to Asymmetric Morphogenesis. *Trends in Genetics*. 33 (9), 616–628.

[4] Kawasumi, A., Nakamura, T., Iwai, N., Yashiro, K., Saijoh, Y., Belo, J.A., et al. (2011) Left–right asymmetry in the level of active Nodal protein produced in the node is translated into left–right asymmetry in the lateral plate of mouse embryos. *Developmental Biology*. 353 (2), 321–330.

[5] Branford, W.W. and Yost, H.J. (2004) Nodal Signaling: CrypticLefty Mechanism of Antagonism Decoded. *Current Biology*. 14 (9), R341–R343.

[6] Zhou, X., Lowe, L., Sasaki, H., Hogan, B.L.M., and Kuehn, M.R. (1993) Nodal is a novel TGF-β-like gene expressed in the mouse node during gastrulation. *Nature*. 361 (6412), 543–547.

[7] Chen, Y. and Schier, A.F. (2002) Lefty Proteins Are Long-Range Inhibitors of Squint-Mediated Nodal Signaling. *Current Biology*. 12 (24), 2124–2128.

[8] Chen, C. and Shen, M.M. (2004) Two Modes by which Lefty Proteins Inhibit Nodal Signaling. *Current Biology*. 14 (7), 618–624.

[9] Essner, J.J., Amack, J.D., Nyholm, M.K., Harris, E.B., and Yost, H.J. (2005) Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left-right development of the brain, heart and gut. *Development*. 132 (6), 1247–1260.

[10] Vick, P., Schweickert, A., Weber, T., Eberhardt, M., Mencl, S., Shcherbakov, D., et al. (2009) Flow on the right side of the gastrocoel roof plate is dispensable for symmetry breakage in the frog Xenopus laevis. *Developmental Biology*. 331 (2), 281–291.

[11] Nonaka, S., Shiratori, H., Saijoh, Y., and Hamada, H. (2002) Determination of left–right patterning of the mouse embryo by artificial nodal flow. *Nature*. 418 (6893), 96–99.

[12] Schweickert, A., Weber, T., Beyer, T., Vick, P., Bogusch, S., Feistel, K., et al. (2007) Cilia-Driven Leftward Flow Determines Laterality in Xenopus. *Current Biology*. 17 (1), 60–66.

[13] Okada, Y., Takeda, S., Tanaka, Y., Belmonte, J.-C.I., and Hirokawa, N. (2005) Mechanism of Nodal Flow: A Conserved Symmetry Breaking Event in Left-Right Axis Determination. *Cell*. 121 (4), 633–644.

[14] Tanaka, Y., Okada, Y., and Hirokawa, N. (2005) FGF-induced vesicular release of Sonic hedgehog and retinoic acid in leftward nodal flow is critical for left–right determination. *Nature*. 435 (7039), 172–177.

[15] Nonaka, S., Tanaka, Y., Okada, Y., Takeda, S., Harada, A., Kanai, Y., et al. (1998) Randomization of Left–Right Asymmetry due to Loss of Nodal Cilia Generating Leftward Flow of Extraembryonic Fluid in Mice Lacking KIF3B Motor Protein. *Cell*. 95 (6), 829–837.

[16] Smith, D.J., Montenegro-Johnson, T.D., and Lopes, S.S. (2019) Symmetry-Breaking Cilia-Driven Flow in Embryogenesis. *Annual Review of Fluid Mechanics*. 51 (1), 105–128.

[17] Schoenwolf, G.C., Bleyl, S.B., Brauer, P.R., and Francis-West, P.H. (2014) Larsen's Human Embryology. Elsevier Health Sciences, .

[18] Hamada, H., Meno, C., Watanabe, D., and Saijoh, Y. (2002) Establishment of
vertebrate left–right asymmetry. *Nature Reviews Genetics*. 3 (2), 103–113.
[19] Gros, J., Feistel, K., Viebahn, C., Blum, M., and Tabin, C.J. (2009) Cell
Movements at Hensen's Node Establish Left/Right Asymmetric Gene
Expression in the Chick. *Science*. 324 (5929), 941–944.
[20] Essner, J.J., Vogan, K.J., Wagner, M.K., Tabin, C.J., Yost, H.J., and Brueckner,
M. (2002) Conserved function for embryonic nodal cilia. *Nature*. 418 (6893), 37–
38.
[21] Stephen, L.A., Johnson, E.J., Davis, G.M., McTeir, L., Pinkham, J., Jaberi, N.,
et al. (2014) The chicken left right organizer has nonmotile cilia which are lost in
a stage-dependent manner in the talpid3 ciliopathy. *Genesis (New York, N.y. :*
2000). 52 (6), 600–613.
[22] Oteíza, P., Köppen, M., Concha, M.L., and Heisenberg, C.-P. (2008) Origin and
shaping of the laterality organ in zebrafish. *Development*. 135 (16), 2807–2813.
[23] Cooper, M.S. and D'amico, L.A. (1996) A Cluster of Noninvoluting Endocytic
Cells at the Margin of the Zebrafish Blastoderm Marks the Site of Embryonic
Shield Formation. *Developmental Biology*. 180 (1), 184–198.
[24] Pulgar, E., Schwayer, C., Guerrero, N., López, L., Márquez, S., Härtel, S., et al.
(2021) Apical contacts stemming from incomplete delamination guide progenitor
cell allocation through a dragging mechanism. *eLife*. 10 e66483.
[25] Kramer-Zucker, A.G., Olale, F., Haycraft, C.J., Yoder, B.K., Schier, A.F., and
Drummond, I.A. (2005) Cilia-driven fluid flow in the zebrafish pronephros, brain
and Kupffer's vesicle is required for normal organogenesis. *Development*. 132
(8), 1907–1921.
[26] Hirokawa, N., Tanaka, Y., Okada, Y., and Takeda, S. (2006) Nodal Flow and
the Generation of Left-Right Asymmetry. *Cell*. 125 (1), 33–45.
[27] Harvey, R.P. (1998) Links in the Left/Right Axial Pathway. *Cell*. 94 (3), 273–276.
[28] Kim, Y.-Y., Moon, J.-S., Kwon, M., Shin, J., Im, S.-K., Kim, H.-A., et al. (2014)
Meteorin Regulates Mesendoderm Development by Enhancing Nodal
Expression. *PLoS ONE*. 9 (2), e88811.
[29] Nishino, J., Yamashita, K., Hashiguchi, H., Fujii, H., Shimazaki, T., and Hamada,
H. (2004) Meteorin: a secreted protein that regulates glial cell differentiation and
promotes axonal extension. *The EMBO Journal*. 23 (9), 1998–2008.
[30] Ramialison, M., Bajoghli, B., Aghaallaei, N., Ettwiller, L., Gaudan, S., Wittbrodt,
B., et al. (2008) Rapid identification of PAX2/5/8 direct downstream targets in the
otic vesicle by combinatorial use of bioinformatics tools. *Genome Biology*. 9 (10),
R145.
[31] Jørgensen, J.R., Fransson, A., Fjord-Larsen, L., Thompson, L.H., Houchins,
J.P., Andrade, N., et al. (2012) Cometin is a novel neurotrophic factor that
promotes neurite outgrowth and neuroblast migration in vitro and supports
survival of spiral ganglion neurons in vivo. *Experimental Neurology*. 233 (1),
172–181.
[32] El-Brolosy, M.A., Kontarakis, Z., Rossi, A., Kuenne, C., Günther, S., Fukuda, N.,
et al. (2019) Genetic compensation triggered by mutant mRNA degradation.
Nature. 568 (7751), 193–197.
[33] Montague, T.G., Gagnon, J.A., and Schier, A.F. (2018) Conserved regulation of
Nodal-mediated left-right patterning in zebrafish and mouse. *Development*. 145
(24), dev171090.

[34] Long, S., Ahmad, N., and Rebagliati, M. (2003) The zebrafish *nodal* -related
gene *southpaw* is required for visceral and diencephalic left-right asymmetry.
Development. 130 (11), 2303–2316.
[35] Brennan, J., Norris, D.P., and Robertson, E.J. (2002) Nodal activity in the node
governs left-right asymmetry. *Genes & Development*. 16 (18), 2339–2344.
[36] Saijoh, Y., Oki, S., Ohishi, S., and Hamada, H. (2003) Left–right patterning of
the mouse lateral plate requires Nodal produced in the node. *Developmental*
Biology. 13.
[37] Pelliccia, J.L., Jindal, G.A., and Burdine, R.D. (2017) Gdf3 is required for robust
Nodal signaling during germ layer formation and left-right patterning. *eLife*. 6
e28635.
[38] Hojo, M., Takashima, S., Kobayashi, D., Sumeragi, A., Shimada, A., Tsukahara,
T., et al. (2007) Right-elevated expression of charon is regulated by fluid flow in
medaka Kupffer's vesicle: Left-right asymmetric expression of charon.
Development, Growth & Differentiation. 49 (5), 395–405.
[39] Gourronc, F., Ahmad, N., Nedza, N., Eggleston, T., and Rebagliati, M. (2007)
Nodal activity around Kupffer's vesicle depends on the T-box transcription
factors notail and spadetail and on notch signaling. *Developmental Dynamics*.
236 (8), 2131–2146.
[40] Alexander, J., Rothenberg, M., Henry, G.L., and Stainier, D.Y.R. (1999)
casanova Plays an Early and Essential Role in Endoderm Formation in
Zebrafish. *Developmental Biology*. 215 (2), 343–357.
[41] Kikuchi, Y., Agathon, A., Alexander, J., Thisse, C., Waldron, S., Yelon, D., et al.
(2001) *casanova* encodes a novel Sox-related protein necessary and sufficient
for early endoderm formation in zebrafish. *Genes & Development*. 15 (12), 1493–
1505.
[42] Amack, J.D. and Yost, H.J. (2004) The T Box Transcription Factor No Tail in
Ciliated Cells Controls Zebrafish Left-Right Asymmetry. *Current Biology*. 14 (8),
685–690.
[43] Matsui, T., Hiroshi, I., and Yasumasa, B. (2015) Cell collectivity regulation within
migrating cell cluster during Kupffer's vesicle formation in zebrafish. *Frontiers in*
Cell and Developmental Biology. 3 8.
[44] Kimmel, C.B., Ballard, W.W., Kimmel, S.R., Ullmann, B., and Schilling, T.F.
(1995) Stages of embryonic development of the zebrafish. *Developmental*
Dynamics. 203 (3), 253–310.
[45] Hashimoto, H., Rebagliati, M., Ahmad, N., Muraoka, O., Kurokawa, T., Hibi, M.,
et al. (2004) The Cerberus/Dan-family protein Charon is a negative regulator of
Nodal signaling during left-right patterning in zebrafish. *Development*. 131 (8),
1741–1753.
[46] Sampaio, P., Ferreira, R.R., Guerrero, A., Pintado, P., Tavares, B., Amaro, J.,
et al. (2014) Left-Right Organizer Flow Dynamics: How Much Cilia Activity
Reliably Yields Laterality? *Developmental Cell*. 29 (6), 716–728.
[47] Gokey, J.J., Ji, Y., Tay, H.G., Litts, B., and Amack, J.D. (2016) Kupffer's vesicle
size threshold for robust left-right patterning of the zebrafish embryo: Size
Threshold For KV Function. *Developmental Dynamics*. 245 (1), 22–33.
[48] de Bruin, K., Ruthardt, N., von Gersdorff, K., Bausinger, R., Wagner, E., Ogris,
M., et al. (2007) Cellular Dynamics of EGF Receptor–Targeted Synthetic
Viruses. *Molecular Therapy*. 15 (7), 1297–1305.
[49] Hynes, R.O. (1992) Integrins: Versatility, modulation, and signaling in cell
adhesion. *Cell*. 69 (1), 11–25.

[50] Ablooglu, A.J., Tkachenko, E., Kang, J., and Shattil, S.J. (2010) Integrin αV is
necessary for gastrulation movements that regulate vertebrate body asymmetry.
Development. 137 (20), 3449–3458.
[51] Reboll, M.R., Klede, S., Taft, M.H., Cai, C.-L., Field, L.J., Lavine, K.J., et al.
(2022) Meteorin-like promotes heart repair through endothelial KIT receptor
tyrosine kinase. *Science*. 376 (6599), 1343–1347.
[52] Ushach, I., Arrevillaga-Boni, G., Heller, G.N., Pone, E., Hernandez-Ruiz, M.,
Catalan-Dibene, J., et al. (2018) Meteorin-like/Meteorin-β Is a Novel
Immunoregulatory Cytokine Associated with Inflammation. *The Journal of
Immunology*. 201 (12), 3669–3676.
[53] Kim, Y.J., VijayKumar, S., Iovino, B., Granados, A., Ancheta, S., Zhao, X., et al.
(2024) Zebrahub-Multiome: Uncovering Gene Regulatory Network Dynamics
During Zebrafish Embryogenesis.
[54] Bisgrove, B.W., Essner, J.J., and Yost, H.J. (2000) Multiple pathways in the
midline regulate concordant brain, heart and gut left-right asymmetry.
Development. 127 (16), 3567–3579.
[55] Solnica-Krezel, L., Stemple, D.L., Mountcastle-Shah, E., Rangini, Z., Neuhauss,
S.C., Malicki, J., et al. (1996) Mutations affecting cell fates and cellular
rearrangements during gastrulation in zebrafish. *Development*. 123 (1), 67–80.
[56] Oteiza, P., Köppen, M., Krieg, M., Pulgar, E., Farias, C., Melo, C., et al. (2010)
Planar cell polarity signalling regulates cell adhesion properties in progenitors of
the zebrafish laterality organ. *Development*. 137 (20), 3459–3468.
[57] Choi, W.-Y., Giraldez, A.J., and Schier, A.F. (2007) Target Protectors Reveal
Dampening and Balancing of Nodal Agonist and Antagonist by miR-430.
Science. 318 (5848), 271–274.
[58] Milner, R., Edwards, G., Streuli, C., and ffrench-Constant, C. (1996) A Role in
Migration for the αvβ1 Integrin Expressed on Oligodendrocyte Precursors. *The
Journal of Neuroscience*. 16 (22), 7240–7252.
[59] Felding-Habermann, B., Silletti, S., Mei, F., Siu, C.-H., Yip, P.M., Brooks, P.C.,
et al. (1997) A Single Immunoglobulin-like Domain of the Human Neural Cell
Adhesion Molecule L1 Supports Adhesion by Multiple Vascular and Platelet
Integrins. *Journal of Cell Biology*. 139 (6), 1567–1581.
[60] Jülich, D., Mould, A.P., Koper, E., and Holley, S.A. (2009) Control of extracellular
matrix assembly along tissue boundaries via Integrin and Eph/Ephrin signaling.
Development. 136 (17), 2913–2921.
[61] Dai, Y.-W., Ma, J.-K., Jiang, R., Zhan, X.-L., Chen, S.-Y., Feng, L.-L., et al.
(2022) Meteorin links the bone marrow hypoxic state to hematopoietic
stem/progenitor cell mobilization. *Cell Reports*. 40 (12), 111361.
[62] Mellgren, E.M. and Johnson, S.L. (2005) kitb, a second zebrafish ortholog of
mouse Kit. *Development Genes and Evolution*. 215 (9), 470–477.
[63] Zheng, S., Li, Z., Song, J., Liu, J., and Miao, C. (2016) Metrnl: a secreted protein
with new emerging functions. *Acta Pharmacologica Sinica*. 37 (5), 571–579.
[64] Hwang, W.Y., Fu, Y., Reyon, D., Maeder, M.L., Tsai, S.Q., Sander, J.D., et al.
(2013) Efficient genome editing in zebrafish using a CRISPR-Cas system.
Nature Biotechnology. 31 (3), 227–229.
[65] Thisse, C. and Thisse, B. (2008) High-resolution in situ hybridization to whole-
mount zebrafish embryos. *Nature Protocols*. 3 (1), 59–69.
[66] Streit, A. and Stern, C.D. (2001) Combined Whole-Mount in Situ Hybridization
and Immunohistochemistry in Avian Embryos. *Methods*. 23 (4), 339–344.

[67] Tang, R., Dodd, A., Lai, D., McNabb, W.C., and Love, D.R. (2007) Validation of
Zebrafish (Danio rerio) Reference Genes for Quantitative Real-time RT-PCR
Normalization. *Acta Biochimica et Biophysica Sinica*. 39 (5), 384–390.
[68] Valcu, M. and Valcu, C.-M. (2011) Data transformation practices in biomedical
sciences. *Nature Methods*. 8 (2), 104–105.
[69] Borovina, A., Superina, S., Voskas, D., and Ciruna, B. (2010) Vangl2 directs the
posterior tilting and asymmetric localization of motile primary cilia. *Nature Cell
Biology*. 12 (4), 407–412.
[70] Sbalzarini, I.F. and Koumoutsakos, P. (2005) Feature point tracking and
trajectory analysis for video imaging in cell biology. *Journal of Structural Biology*.
151 (2), 182–195.
[71] Tarantino, N., Tinevez, J.-Y., Crowell, E.F., Boisson, B., Henriques, R.,
Mhlanga, M., et al. (2014) TNF and IL-1 exhibit distinct ubiquitin requirements for
inducing NEMO–IKK supramolecular structures. *Journal of Cell Biology*. 204 (2),
231–245.
[72] Zwillinger, D. (2022) CRC Standard Mathematical Tables and Formulae, 31st
Edition (Advances in Applied Mathematics) 31st Edition. 31st ed. Chapman and
Hall/CRC, .
[73] Papoulis, A. (1984) Probability, random variables, and stochastic processes.
McGraw-Hill Inc.,US, .
[74] Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., et al.
(2008) Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic
Acids Research*. 36 (Web Server issue), W465–W469.
[75] Dereeper, A., Audic, S., Claverie, J.-M., and Blanc, G. (2010) BLAST-
EXPLORER helps you building datasets for phylogenetic analysis. *BMC
Evolutionary Biology*. 10 8.
[76] Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy
and high throughput. *Nucleic Acids Research*. 32 (5), 1792–1797.
[77] Castresana, J. (2000) Selection of Conserved Blocks from Multiple Alignments
for Their Use in Phylogenetic Analysis. *Molecular Biology and Evolution*. 17 (4),
540–552.
[78] Guindon, S. and Gascuel, O. (2003) A Simple, Fast, and Accurate Algorithm to
Estimate Large Phylogenies by Maximum Likelihood. *Systematic Biology*. 52 (5),
696–704.
[79] Anisimova, M. and Gascuel, O. (2006) Approximate Likelihood-Ratio Test for
Branches: A Fast, Accurate, and Powerful Alternative. *Systematic Biology*. 55
(4), 539–552.
[80] Chevenet, F., Brun, C., Bañuls, A.-L., Jacq, B., and Christen, R. (2006) TreeDyn:
towards dynamic graphics and annotations for analyses of trees. *BMC
Bioinformatics*. 7 439.
952

RESOURCE AVAILABILITY

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Filippo Del Bene (filippo.del-bene@inserm.fr).

Materials availability

All unique/stable reagents generated in this study are available from the lead contact with a completed materials transfer agreement.

Data and Code Availability

•All data reported in the paper are available from the lead contact upon request.

•This paper does not report original code.

•Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGEMENTS

We thank all members of the Del Bene lab for discussions, the Institut Curie and Institut de la Vision imaging and animal facilities. We acknowledge the lab of Jean Livet for their support with the chick analysis. We would like to thank Pierre Luc Bardet from IBPS (Sorbonne Université) for fruitful discussions. We thank Clémence Gentner from Gilles Tessier's lab at the Institut de la Vision for the microspheres sample. Work in the Del Bene laboratory was supported by ANR MetAxon [ANR-17-CE16-0007], the IHU FOReSIGHT [ANR-18-IAHU-0001], CNRS, INSERM, and Sorbonne Université core funding. F.E. was supported by the École des Neurosciences de Paris (ENP) Ile-de-France network and the Fondation pour la Recherche Médicale (FRM). F.D.S. was supported by a doctoral fellowship of the Curie International PhD program. T.O.A. was supported by a Boehringer Ingelheim Fonds PhD Fellowship.

AUTHORS CONTRIBUTION

F.E. did the experimental work and analyzed the results. L.B. contributed to the experimental work. F.D.S., T.A and K.D. generated the CRISPR/Cas9 *metrn* mutant lines. J.B.W performed the KV microbeads tracking analysis. F.E, S.A. and F.D.B. conceived the project and wrote the manuscript. S.A. and F.D.B. co-supervised the study. All the authors reviewed and edited the manuscript.

DECLARATION OF INTERESTS

985   The authors declare no competing interests.

SUPPLEMENTARY FIGURES

Supp Figure 1

Fig. S1. (A-C) *Metrn* **genes CRISPR/Cas9 mediated knockout generation and validation. (D)** *Metrns* **loss of function effect on heart morphology.** (**A**) Targeted *metrns* loci in the zebrafish genome for knockout generation. Red highlights the sgRNA homologous sequence and PAM sequences are highlighted in green. The generated mutation is indicated below by dashed lines and the number of deleted nucleotides is indicated after the Δ sign. (**B**) *Metrn, metrnla* and *metrnlb* expression in triplMut embryos from 2-cell stage to 1 day post fertilization (dpf) is highly reduced or undetectable by *in situ* hybridization (lateral views). (**C**) *Metrns* expression levels are reduced in triplMut embryos as shown by qRT-PCR analysis of *metrn*, *metrnla*, *metrnlb* expression level at 14 hpf for *metrn* and *metrnla* and at 48 hpf for *metrnlb* in wild type (WT) and triplMut embryos. (Student t-test, ***p-value: 3.1e-06 for *metrn*; *p-value: 0.014 for *metrnla*; *p-value: 0.04 for *metrnlb*). (**D**) Quantification in percentage of the number of embryos with S-looped, D-looped or mild/no looped heart phenotypes at 2 dpf upon injection of *metrnla* mRNA, *metrn* mRNA or both in triplMut embryos. Displayed p-values compared to WT: ****p-value: <1.0e-5 for *metrnla* mRNA, ****p-value: <1.0e-5 for *metrn* mRNA, ****p-value: <1.0e-5 for *metrn + metrnla* mRNA. Scale bar in (**B**) 250 μm.

Supp Figure 2

	WT-like	abnormal	reduced
triplMut⁺/⁻	60.63%	7.87%	31.50%
metrn⁻/⁻	39.50%	40.33%	28.57%
metrnla⁻/⁻	53.16%	20.89%	25.95%

Fig. S2. Nodal factors gene expression is altered in the absence of Metrns. Quantification of the number of embryos in percentage with reduced, inverted or wild type (WT) -like *dand5* expression pattern in 14 hpf triplMut$^{+/-}$, *metrn$^{-/-}$* and *metrnla$^{-/-}$* embryos, compared to WT, Permutation test, ****p-value <1.0e-5 for triplMut$^{+/-}$, ****p-value <1.0e-5 for *metrn$^{-/-}$*, ****p-value <1.0e-5 for *metrnla$^{-/-}$*.

Supp Figure 3

Fig. S3. *Metrns* expression during early zebrafish and chick development. (**A**) No expression can be detected for *metrn, metrnla* and *metrnlb* upon *in situ* hybridization using sense *metrn, metrnla*

and *metrnlb* riboprobes on 2-cell stage wild type embryos. (**B**) *Metrn* expression at 10 hpf can be found in the leading edge of the shield and in the forming Kupffer's vesicle as shown by *in situ* hybridization (dorsal view) and at 14 hpf *metrn* transcripts can be detected in the area of the KV and the developing brain (later view). While *metrnla* at 10 hpf is expressed in the whole enveloping layer and developing midline as revealed here by *in situ* hybridization (dorsal view). At 14 hpf *metrnla* transcripts can be found in the area of the forming KV and midline (later view). No expression can be detected for *metrnlb* at these stages. (**C**) From 24 hpf, all *metrn* genes are expressed in the developing central nervous system as shown by *in situ* hybridization (lateral views). (**D**) Dorsal view of *metrn*, *metrnl* and *fgf8 in situ* hybridization in HH6 chick embryos reveal that *metrn*, *metrnl* are expressed around the Hensen's node (*) and the primitive streak during chick early embryonic development. Scale bars: in (**A, B** and **D**) 250 µm and in (**C**) 100 µm.

Fig. S4. *Metrns* **loss-of-function leads to DFC disorganization and migration defects.** (**A**) *Sox32* antisense riboprobe *in situ* hybridization revealing DFC misclustering in 9 hpf *metrn^-/-* and *metrnla^-/-* embryos. (**B**) Quantification in percentage of the number of embryos with DFC clustering defects at 9 hpf in wild type (WT) and several *metrns* mutant backgrounds (Fisher exact test, ***p-value: 1.5e-

05 for *metrn*[-/-] vs. *metrn*[+/-]; ***p-value: 2.5e-26 for *metrnla*[-/-] vs. *metrnla*[+/-]; ***p-value: 1.2e-13 for *metrn*[-/-] x *metrnla*[-/-] vs. *metrn*[+/-] x *metrnla*[+/-]; ***p-value: 2.2e-18 for triplMut vs. ♀WT x ♂triplMut and ***p-value: 2.1e-13 for triplMut vs. ♀triplMut x ♂WT) (**C**) *tbxta* and *sox17* DFC specification markers are expressed in the absence of Metrn proteins as shown by *in situ* hybridization on WT and triplMut at 8 and 10 dpf respectively. Scale bars: in **A** and **C** 0.5 mm.

Supp Figure 5

Fig. S5. *Metrns* loss-of-function leads to DFC disorganization and migration defects. (**A**) The KV cilia number and cells per KV quantification in WT and triplMut embryos at 14 hpf show no significant difference between cilia number and KV cells (N WT: 23, N triplMut: 26, Student T-tests, n.s.: 0.67 for WT KV cells, 0.96 for triplMut KV cells). (**B**) DFC migration quantification calculated as the percentage of the total embryo length at 6, 8 and 10 hpf in wild type (WT) *vs*. triplMut embryos. Per condition and group 16 to 27 embryos were analyzed (Fisher exact test, ***p-value: 2.5e-06 for 6 hpf, ***p-value: 0.00038 for 8 hpf and ***p-value: 1.1e-08 for 10 hpf). (**C**) *Sox32* antisense riboprobe *in situ* hybridization highlighting DFC migration between 6 hpf and 10 hpf in WT and triplMut embryos (point line marks the margin of epiboly). All depicted images are dorsal views. (**D**) Single traces of *in vivo* tacked DFCs in WT control (upper panel) and triplMut (lower panel) showing the vegetal movement and convergence. (**E**) Quantification at 6hpf (left plot) and 8 hpf (right plot) of of DFC adjacent ZO-1 enrichments areas reveal a decreased in triplMut embryos in contrast to respective WT controls (T-test; 6hpf *p-value: 0.043, 8hpf *p-value: 0.028). (**F**) The distance of the leading DFCs to the EVL is increased in triplMut compared to WT controls visualized at 6 hpf and 8 hpf (nested T-test, *p-value: 0.039 N = 4 WT, 30 DFCs measured,

triplMut N = 6, 46 DFCs measured; for 6 hpf and *p*-value: 0.021 N = 7 WT embryos, 47 DFCs measured, N = 7 triplMut, 54 DFCs measured for 8 hpf). Scale bars: in (**C**) 0.25 mm.

Fig. S6. *Metrns* loss-of-function impairs the Kupffer's vesicle formation and function. (**A**) Dorsal views of 14 hpf wild type (WT) and triplMut embryos immunostained with ZO-1. Wild type embryos display a large KV lumen with uniform ZO-1 labeled tight junction lattice. TriplMut embryos show disturbed KV lumen and dysmorphic ZO-1 lattice. (**B**) WT KV cells of 14 hpf *Tg*(*sox17*:GFP) show proper polarization as visualized with anti-αPKCζ immunostaining (white arrowhead). 14 hpf TriplMut embryos in contrast exhibit a reduced polarization (white arrowhead). (**C**) Tracked microbeads velocity analysis in 12-14 hpf WT and triplMut embryos. The purple line marks the mean and the standard deviation (std) for each condition (WT mean= 0.96212; std=0.048143; TriplMut mean=0.47996; std=0.023682). Scale bars: in (**A-B**) 20 µm.

Supp Figure 7

Fig. S7. *Kita, kitb* and *htr2b* **expression patterns during early zebrafish development.** (**A**) qRT-PCR analysis for *itgaV* and *itgb1b* expression in 6 hpf, 9 hpf and 24 hpf wild type (WT) and triplMut embryos (Fisher exact tests, n.s. = non-significant, *p*-values: 0.95 for 24 hpf *itgaV* WT *vs. itgaV* triplMut, 0.37 for 9 hpf hpf *itgaV* WT *vs. itgaV* triplMut, 0.92 for 6 hpf *itgaV* WT *vs. itgaV* triplMut, 0.64 for 24 hpf *itgb1b* WT *vs. itgb1b* triplMut, 0.7 for 9 hpf *itgb1b* WT *vs. itgb1b* triplMut, 0.46 for 6 hpf *itgb1b* WT *vs. itgb1b* triplMut) (**B**) At 9 hpf *kita* is expressed in the prechordal plate (black arrowhead, dorsal view) and from 11 hpf in the lateral borders of the anterior neural plate (black arrowhead, lateral view). *Kitb* expression could be detected from 11 hpf in the anterior ventral mesoderm (black arrowhead, lateral view). (**C**) At 9 hpf, *htr2b* is not yet expressed. From 2 days post-fertilization (dpf), *htr2b* transcripts could be detected in the developing heart (black arrowhead, lateral views). (**D**) Phylogenetic tree of Metrn proteins. (Scale bar indicates number of amino acids substitutions per site). Scale bars: in (**B – C**) 250 µm.

SUPPLEMENTARY VIDEOS

Video S1: *In vivo* imaging of GFP-positive DFCs (single cells labelled with *H2B-mRFP* mRNA, full stack) in a WT (WT 2 from Fig. S5D) embryo from 50% epiboly for 3 hours showing directed DFC movement and clustering.

Video S2: *In vivo* imaging of GFP-positive DFCs (single cells labelled with *H2B-mRFP* mRNA, full stack) in a triplMut embryo (triplMut 2 from Fig. S5D) from 50% epiboly for 3 hours showing directed DFC movement, but clustering defects.

Video S3: Microbeads tracking the Kupffer's vesicle of 12-14 hpf wild type embryos showing a distinctive anti-clockwise rotation.

Video S4: Microbeads tracking the Kupffer's vesicle of 12-14 hpf triplMut embryos showing the loss of the distinctive anti-clockwise rotation.

SUPPLEMENTARY TABLES

	D-loop	mild/no-loop	S-loop
WT (N = 121)	88.43%	11.57%	0%
***metrn*^{-/-}** (N = 131)	61.19%	29.10%	8.21%
***metrnla*^{-/-}** (N = 104)	28.85%	51.92%	19.23%
***metrnlb*^{-/-}** (N = 104	74.21%	21.38%	4.41%
triplMut (N = 134)	38.17%	25.95%	35.88%
triplMut^{+/-} (N = 182)	85.16%	14.29%	0.55%
triplMut + *metrnla* mRNA (N = 160)	37.5%	53.13	9.37%
triplMut + *metrn* mRNA (N = 156)	45.51%	42.31%	12.18%
triplMut + *metrnla* + *metrn* mRNA (N = 162)	52.47%	27.16%	20.37%

Table 1: Quantification of heart looping phenotypes at 2 dpf using *myl7* riboprobe *in situ* hybridization on wild-type (WT), *metrn*s single and triple mutants (triplMut) and *metrn* and/or *metrnla* mRNAs -injected embryos. 'Mild/no loop' designates hearts with no particular L-R orientation. D-loop, dextral loop; mild/no-loop and S-loop, sinistral loop.

	WT-like (% of embryos)	DFC clustering defects (% of embryos)
WT (N = 202)	93.07	6.93
metrn^{-/-} (N = 140)	63.57	36.43
metrnla^{-/-} (N = 245)	46.94	53.06

triplMut (N = 182)	34.62	65.38
triplMut^{+/-} (N = 140)	79.51	20.49

1023

1024 **Table 2:** Quantification in percentage of embryos with DFC clustering defects at 9 hpf in WT, *metrn*^{-/-},

1025 *metrnla*^{-/-}, triplMut and and triplMut^{+/-} embryos.

1026

	6 hpf	8 hpf	10 hpf
WT	59.32% (+/- 7.65%, N = 27)	78.18% (+/- 7.94%, N = 19)	93.67% (+/- 2.68%, N = 16)
triplMut	46.26% (+/- 9.83%, N = 24)	69.94% (+/- 6.22%, N = 24)	80.67% (+/- 6.56%, N = 20)

1027

1028 **Table 3:** Quantification of the DFC migration calculated as the percentage of the total embryo length at

1029 6, 8 and 10 hpf in wild type (WT) *vs*. triplMut embryos.

1030

Experiment	Number of embryos
WT non injected	346
WT + ctrl MO	192
WT + 0.41 ng *itgαV* MO	208
WT + 0.5 ng *itgβ1b* MO	623
WT + 1.25 ng *itgαV* MO	412
WT + 0.7 ng *itgβ1b* MO	336
triplMut^{+/-} non injected	219
triplMut^{+/-} + ctrl MO	176
triplMut^{+/-} + 0.41ng *itgαV* MO	182
triplMut^{+/-} + 0.5 ng *itgβ1b* MO	559

1031 **Table 4:** Number of embryos with DFC clustering defects at 9 hpf upon *itgβ1b*, *itgαV* or control (ctrl)

1032 morpholino (MO) injection in wild type (WT) or triplMut^{+/-} embryos.

1033