A Unified Approach to Investigating 4 dpf Zebrafish Larval Behaviour through a Standardised

Light/Dark Assay

Courtney Hillman[1], James Kearn[2], and Matthew O. Parker[1,3]

[1]Surrey Sleep Research Centre, University of Surrey, Guildford, UK

[2]Defence Science and Technology Laboratory (DSTL), UK

[3]School of Pharmacy and Biomedical Sciences, University of Portsmouth, Portsmouth, UK

Declaration of interest: none

Author note

Correspondence concerning this article should be addressed to Matthew O. Parker, PhD.
Surrey Sleep Research Centre, University of Surrey, Guildford, GU2 7XH, UK. Email:
matthew.parker@surrey.ac.uk, phone number: +44 7986 205349

Email addresses of authors:

Courtney Hillman, BSc.: c.hillman@surrey.ac.uk

James Kearn, PhD.: jkearn@dstl.gov.uk

Matthew O. Parker, PhD.: matthew.parker@surrey.ac.uk

27   **Abstract**

Zebrafish have emerged as a dynamic research model in the domains of neuropsychopharmacology,

biological psychiatry and behaviour. Working with larvae $\leq$ 4 days post-fertilisation (dpf) offers an

avenue for high-throughput investigation whilst aligning with the 3Rs principles of animal research.

The light/dark assay, which is the most used behavioural assay for larval neuropharmacology

research, lacks experimental reliability and standardisation. This study aimed to formulate a robust,

reproducible and standardised light/dark behavioural assay using 4 dpf zebrafish larvae. Considerable

between-batch and inter-individual variability was found, which we rectified with a normalisation

approach to ensure a reliable foundation for analysis. We then identified that 5-minute light/dark

transition periods are optimal for locomotor activity. We also found that a 30-minute acclimation in

the light was found to produce significantly increased dark phase larval locomotion. Next, we

confirmed the pharmacological predictivity of the standardised assay using ethanol which, as

predicted, caused hyperlocomotion at low concentrations and hypolocomotion at high

concentrations. Finally, the assay was validated by assessing the behavioural phenotype of

hyperactive transgenic (*adgrl3.1$^{-/-}$*) larvae, which was rescued with psychostimulant medications. Our

standardised assay not only provides a clear experimental and analytical framework to work with 4

dpf larvae, but also facilitates between-laboratory collaboration using our normalisation approach.

Keywords: Zebrafish, light/dark transition test, standardized testing, locomotor activity, 3Rs

of animal research

Abbreviations[1]

____

[1] ATO: atomoxetine
dpf: days post-fertilization
LRT: likelihood ratio test
MPH: methylphenidate
SD: supplementary data
WT: wild-type

1. Introduction

Zebrafish (*Danio rerio,* Hamilton 1822) larvae are an increasingly popular research model
with a breadth of application and tractability for neuropsychopharmacology (Cassar et al., 2020). Due
to their rapid breeding and small size, working with larvae provides potential for high-throughput
approaches, with the added benefit that larvae up to 5 days post-fertilisation (dpf) are exempt from
protection by national legislation (e.g., the (Animals (Scientific Procedures) Act 1986) 1986) in the
United Kingdom (UK) and the European Union (EU) Directive 2010/63/EU of the European
Parliament. Hence, 4 dpf larval research has significant interest for meeting the criteria of the 3Rs
(replacement, refinement and reduction) of *in vivo* animal research (Crilly et al., 2018). Indeed, as a
first-tier model, 4 dpf zebrafish larvae can reduce the number of protected animals required at later
testing stages as well as minimising the suffering of experimental animals when testing toxic
substances. Therefore, from a 3Rs perspective, zebrafish larvae are a unique and appealing research
model.

The light/dark assay remains the most used for studying the behaviour of zebrafish larvae
owing to its cost and time-effectiveness, as well as its high face and pharmacological predictivity
(Haigis et al., 2022). The protocol manipulates the innate tendency for larvae to display hyperactivity
following a sudden change from light to dark conditions (Bachour et al., 2020). Some have suggested
that this locomotor response represents a stress or alarm response, and it has been used for
modelling anxiety-like phenotypes in larvae (Rock et al., 2022; Steenbergen et al., 2012). Its
popularity stems from its ease of manipulation for toxicity and drug analysis and the robust
behavioural response observed (Chatzimitakos et al., 2022; Haigis et al., 2022; Kirla et al., 2021;
Licitra et al., 2021; Nixon et al., 2021; Pesavento et al., 2022; Sales Cadena et al., 2021; Zaig et al.,
2021). As seen in Table 1, there is, however, a lack of consistency between light/dark assay
experimental protocols in 4 dpf larvae. The lack of experimental consistency can impact the
behavioural responses observed and hence the reproducibility of results between laboratories and
researchers (Pesavento et al., 2022). This lack of standardised testing and variability in protocols has
been highlighted previously by Legradi (2015). However, no standardised or reproducible assay has
been published to date.

Author and year of study	Age of larvae (dpf)	Acclimation period	Transition period
Y. Wang, 2023	4	15 min in light	10 min light to dark x 3 (60 min total)
Okeke, 2022	4	10 min in dark	10-min light to dark to light x1 (30 min total)

Scarlett, 2022	4	10 min in dark	10-min light to dark x 2 (40 min total)
Mundy, 2021	4	5 min in light	5-min light to dark to light followed by 15-min dark x1 (30 min in total)
Hasumi, 2020	4-5	N/A	15-min light to dark x 6 (180 min total)
Basnet, 2019	4-5	30 min in dark	10-min light to dark x 6 (120 min)
Walter, 2019	3 – 5	10 min in light	5-min dark to 5-min light to 15-minute dark (25 min total)
Best, 2017	4	Overnight acclimation in dark + 1 hour	7.5-min light to dark x8 (60 min total)
Fraser, 2017	4-5	20 min in light	10-min dark to light x1 (20 min total)
Liang, 2017	4	840 min in light	10-min dark to light x4.5 (50 min total)
Thit, 2017	4	N/A	5-min dark to light x 9 (90 min total)
Liu, 2015	3 – 9	210 min in dark	30-min light to dark x 3 (180 min total)
Zhao, 2014	4 – 6	20 min in dark	10-min in dark into 10-min light to dark x 3 (70 min total)
Beker van Woudenberg, 2013	4	15 min in light	15-min dark to light to dark (45 min total)
Padilla, 2011	4, 5 and 6	20 min in dark	10-min light to dark x 2
Jin, 2010	4	20 min in dark	10-min light to dark x 3.5 (70 min in total)

Table 1. List of previous publications using 4 dpf zebrafish larvae and the different light dark parameters
used. dpf: days post-fertilisation, min: minutes.

Here, we aimed to develop a standardised robust and reproducible light/dark behavioural
assay using 4 dpf zebrafish larvae and to validate this assay using pharmacological and genetic
manipulations. After assessing different parameters to determine the optimal experimental
conditions, we validated the assay pharmacologically in two-ways. Wild-type larvae were exposed to
ethanol and genetically modified (*adgrl3.1* knock-out *(adgrl3.1$^{-/-}$)*) larvae were used to study
behavioural interventions of two Attention-Deficit Hyperactivity Disorder (ADHD) drugs. Ethanol was
selected due to its well-known ability to induce low concentration hyperactivity and high
concentration hypoactivity, which we postulated would impact the dark phase responses (McLellan,
2017). Furthermore, it is widely available, is globally abused, and is commonly used in larval
light/dark assay research (Davis et al., 2021; Irons et al., 2010; MacPhail et al., 2009). The *adgrl3.1$^{-/-}$*
mutant line demonstrates phenotypes typical of ADHD and therefore we hypothesised that the
larvae would display hyperactivity in the dark phase (Sveinsdóttir et al., 2023) which would be
rescued/reversed with methylphenidate (MPH), a standard stimulant drug used to treat ADHD. We
also assessed atomoxetine (ATO), another common ADHD drug known to induce sedation in patients,

which we postulated would result in hypoactivity (Bastiaens, 2007). In addition to the validation, we
also aimed to provide a consolidated method of analysis and interpretation of the behavioural data,
including radar charts and heat maps of different swim patterns, which will be of use to the zebrafish
research community.

2. Materials and Methods
2.1. Animal husbandry

AB wild-type (WT) zebrafish were bred in-house from the University of Portsmouth colony.
Breeding adults in the colony were housed in ~50:50 male:female groups of 8-10 per 2.8 L tank, or up
to 20 per 6 L tank, on a re-circulating system (Aquaneering, Fairfield Controlec Ltd., Grimsby, UK).
Sample sizes (n = 16) were calculated based on power analyses from effect sizes observed in our pilot
studies (*see section 2.8. Power Analysis*). $adgrl3.1^{-/-}$ larvae were used for the genetic phenotype
experiment. The $adgrl3.1^{-/-}$ line was created by CRISPR-Cas9 genome engineering as previously
described (Sveinsdóttir et al., 2023) and maintained via homozygous in-cross at the University of
Portsmouth fish facility (with matched-age WT [AB] controls). Room and tank water temperatures
were maintained at 25-27°C, pH 8.4 (±0.4), on a 14:10-hour light/dark cycle with the lights switched
on at 08:30. Fish were fed on ZM-100 fry food (Zebrafish Management Ltd., Winchester, UK) from 5
dpf until adulthood when they were moved onto a diet of flake food and live brine shrimp 3
times/day (once/day on weekends). Larvae were bred for experimentation either by adding marbles
to the base of the tank overnight (to induce spawning) or by pair breeding. The larvae used in these
experiments originated from over 10 different batches from different breeders. Larvae were reared
in a translucent incubator (28.5°C) in the fish facility until the day of testing, where they were moved
into our behavioural testing room and placed in a small countertop incubator at 28°C. On completion
of behavioural testing, larvae were killed using Aqua-Sed (Aqua-Sed™, Vetark, Winchester, UK) in
accordance with manufacturer guidelines. All experiments were approved by the University of
Portsmouth Animal Welfare and Ethical Review Board. Genetically altered fish were used under
license from the UK Home Office (Animals (Scientific Procedures) Act, 1986, Licence number
PP8708123).

2.2. Behavioural apparatus

All behavioural testing was carried out between 09:00 and 18:00 (Mon-Fri) in Zantiks MWP
units (Zantiks Ltd, Cambridge, UK). Previous literature determined that 48-well plates are the optimal
format for consistent and robust behavioural responses and high-throughput analysis (Widrick et al.,
2023). However, our preliminary studies suggest any well-plate size is suitable *(see SD 1)*. Therefore,
with respect to previous literature and both throughput and time-efficiency, all behavioural testing

took place using 48 well plates. Filming was carried out directly from above the plates, which allowed

live monitoring within the behavioural system.

Figure 1. The experimental protocol for the standardised light/dark assay including drug exposure

procedures. (A) the fish are transferred from the incubator to a 48 well plate and placed in the

behavioural unit with the lights on for 30-minutes of acclimation prior to the recording. (B)

Breakdown of the L/D (light/dark) recording which incorporates 5-minute light phase to dark phase

transitions for 30-minutes. (C) Overview of drug exposure procedure from a 96-well dosing plate to

the fish. Behaviour was then monitored using a Zantiks MWP unit.

2.3. Standardisation of Light/Dark Assay

A breakdown of the experimental procedure is seen in Figure 1. At 08:30 the larvae Petri dishes
were transferred into an incubator in our behavioural testing room at 28°c to allow the fish to
acclimate to the new environment for 30 minutes, prior to experimentation. 225 μL of Petri dish
water and a randomly selected larva were transferred into each well of a 48-well plate using trimmed
pipette tips (~ 2 mm removed from the tip end of a 1000 μL pipette tip). The larvae were selected
from different Petri dishes at random for each experiment. The well plate was transferred to the
Zantiks MWP unit, and the overhead lights in the unit were switched on for 30-minutes for
acclimation (Fig. 1A). Following acclimation, the light/dark tracking script was run for a 30-minute
baseline recording. This comprised three sets of five-minute light phase (white light, 350 lux) to dark
phase transitions (Fig. 1B). The MWP unit was temperature controlled throughout experimentation
at 28.02°c ± 0.07.

2.4. The Light/Dark Assay for Chemical Exposures

The drug exposure protocol incorporated the standardised protocol (*Section 2.3 of this paper*).
Following baseline recording, the plate was removed from the MWP, and a multi-channel pipette was
used to administer 25 μL of different ethanol concentrations *(CAS number: 64-17-5, tested at 0.5 –*
4.0 %) (Merck Life Sciences UK Ltd., Gillingham, UK) from a 96-well plate to achieve the desired final
exposure concentration (96-well plate contained 5-40% concentrations) (Fig.1C). The 48-well plate
was returned to the unit and the larvae were tracked during the light/dark alternations (Fig 1B).

2.5. The Light/Dark Assay for Genomic Phenotype Evaluation

Assays with adgrl3.1$^{-/-}$ larvae incorporated the standardised protocol and both WT and adgrl3.1$^{-/-}$
larvae were used (*Section 2.3 of this paper*). Following the baseline recording, the plate was
removed, and a multi-channel pipette was used to administer 0.05 – 20 μM MPH *(CAS number: 298-*
59-9) (Merck Life Sciences UK Ltd., Gillingham, UK) *and* 2.93 - 10.2 μM ATO *(CAS number: 82248-59-*
7) (Merck Life Sciences UK Ltd., Gillingham, UK) (Fig. 1C). Both MPH and ATO were initially diluted in
100% dimethyl sulfoxide (DMSO) for a final concentration of 0.5%, which is a safe concentration in
larvae (Hoyberghs et al., 2021). The 48-well plate was returned to the unit and the larvae were
subjected to the light/dark protocol (Fig 1B).

2.6. Optimisation of the Light/Dark Assay

Alterations in a range of parameters were investigated as part of the standardisation of the
light/dark assay. This included assessing previously published protocols (Acevedo-Canabal et al.,
2019; Bachour et al., 2020; Haigis et al., 2022), for determining the effect of altering the length of

light/dark transition periods. The length of acclimation period and lighting conditions were assessed
as well as the effect of the time of day upon behavioural responses.

2.7. Statistics

The data files were imported into a custom R processing script (R Core Team, 2021) where
they were normalised to baseline to overcome larval batch variability. Normalisation was carried out
by determining the percentage of movement compared to the overall baseline mean response of all
the larvae. Outliers were identified and eliminated if they were greater than the absolute deviation
around the median, a method commonly used in neuropsyhcopharmacological research for outlier
identification and removal (Leys et al., 2013). All graphs and analyses were produced and performed
using the R script. Data was assessed for distribution using the Shapiro-Wilk test and Levine's test for
homogeneity. Statistically significant differences were assessed using a one-way analysis of variance
(ANOVA) and for chemical/ strain analyses, linear mixed models were implemented with movement
as the dependent variable, concentration or strain of fish, light phase and time as independent
variables and fish ID as the random effect. A second linear mixed model analysis was run on the dark
phase data only with movement as the dependent variable, concentration or strain of fish and time
as independent variables and fish ID as the random effect. Likelihood ratio tests (LRT) were
performed on all models (i.e., the fitted models compared to null models which included only
random effects) to test the null hypotheses that added parameters (independent variables) did not
have a significant effect on the outcomes. Post-hoc tests incorporated into the analysis included
Tukey's multiple comparison for comparing dark phase responses and linear regressions analysing
dark phase movement with time as the independent variable. The alpha levels are $p^{***} < 0.001$, p^{**}
< 0.01, $p^* < 0.05$.

Radar charts and heat maps were produced by adapting an approach described by Haigis (2022)
and Steele (2018). Briefly, within the custom R processing script, the percentage of baseline data was
analysed as either darting (>2000 %/s), steady swimming (690 – 2000 %/s), slow swimming (0.1 - 690
%/s) or zero (0 %/s) movements, the values of which were adapted from Haigis (2022), Steele (2018)
and Scarlett (2022) according to relative movement in our experiments. Each observation had an
addition of one to overcome the limitations of percentage determination using a zero response. The
endpoints evaluated were total movements, darting swimming movements, steady swimming
movements, slow swimming movements and the absence of movements (zero movements). The
mean response for each of these observations was calculated for controls. The percentage of
movement compared to vehicle-controls was calculated for each drug concentration and a base 10
logarithmic conversion was performed.

2.8. Power Analysis

Observed power and where appropriate, sample size, was calculated for all tested concentrations and strains of fish. When observed power was below 0.8 (80%), a sample size calculation was performed. If the observed power was $\geq$ 0.8, then the sample size was n = 16. Figure 2 provides an overview of this. All power analysis results, code, data and comprehensive statistical results can be found at the OSF linked to this paper: https://osf.io/k7y2c/?view_only=61e23ea4bb2d452dab76646931e3f332.

Figure. 2. Flow chart depicting the power analysis and sample size determination procedure for larval zebrafish drug exposures.

All figures were created with Biorender.com and data is represented as mean ± SEM. The minute analysis graphs have data points which represent averaged locomotor responses of the fish. Comparatively, the bar charts have data points which represent individual fish responses.

3.　Results

3.1. Five-minute light/dark protocol produces a robust response in 4 dpf zebrafish larvae allowing for high throughout behavioural analysis

Previous publications have lacked consistency with respect to the length and number of light/dark transition periods for assays using 4 dpf zebrafish larvae (Table 1). Therefore, we assessed the optimal length of light to dark transition periods by replicating the protocols employed by Acevedo-Canabal (2019), Bachour (2020) and Haigis (2022), as well as assessing our own protocol. There were reproducible and robust dark phase locomotor responses with all transition periods. In addition, the five-minute (one-way ANOVA: $F_{(2, 41)}$ = 0.656, p = 0.524) and ten-minute transition (one-way ANOVA: $F_{(3, 60)}$ = 0.21, p = 0.889) protocols, had no significant change in locomotion across successive dark phases (Fig. 3D and 3F). In both the two-minute (one-way ANOVA: $F_{(4, 75)}$ = 3.654, p = 0.008) and fifteen-minute transition (one-way ANOVA: $F_{(5, 86)}$ = 2.731, p = 0.024) protocols, however, we observed a significant reduction across successive dark phases (Fig. 3B and 3H). With the two-minute transition period, locomotion in dark phases four and five is significantly less than dark phase

one (Fig. 3B). Similarly, with the fifteen-minute transition period, dark phase six is significantly less

than dark phase one (Fig. 3H).

Figure 3. Effect of varying light and dark phase length on larval behavioural response. Light/dark
locomotion displayed as a percentage of baseline response are represented in minutes for (A) two-
minute transitions (C) five-minute transitions (E) ten-minute transitions and (G) fifteen-minute
transitions. The mean dark phase responses are numbered as D1-D6 for (B) two-minute transitions,
(D) five-minute transitions, (F) ten-minute transitions and (H) fifteen-minute transitions.

We next looked in detail at locomotion within the dark phases, and calculated the
locomotion within the first minute, half-way through, and during the final minute of the dark phase
(Fig. 4) (n.b., the two-minute period was only sufficient for a single dark phase analysis). There was a
significantly reduced initial dark phase locomotion during the fifteen-minute protocol compared to
the two-minute transition period (one-way ANOVA: $F_{(3, 60)}$ = 4.002,   = 0.0115) (Fig. 4A). No
significant difference in locomotion is seen half-way through the dark phase regardless of length of
transition period (one-way ANOVA: $F_{(2, 44)}$ = 2.543,   = 0.0901) (Fig. 4B). However, by the end of the
recording period, the two-minute, five-minute and ten-minute locomotion's are significantly greater
than the fifteen-minute protocol (one-way ANOVA: $F_{(3, 60)}$ = 7.021,   = 0.0004) (Fig. 4C).

Figure 4. Time-dependent effects of varying light and dark phase length on larval behavioural
responses. Locomotion displayed as a percentage of baseline response represented as mean dark
phase responses for (A) the first minute of each dark phase (B) the median minute of each dark
phase and (C) the final minute of each dark phase for the two-minute, five-minute, ten-minute and
fifteen-minute transition protocols.

In view of the sustained increased locomotion throughout the recording period with the five-
minute transition protocol, and relatively higher throughput potential, we proceeded with further
testing using the five-minute transition protocol.

3.2. The acclimation period affects locomotor behavioural responses in the larval light/dark
assay

The importance of the length and lighting conditions of the acclimation period has been
previously reported (Makaras et al., 2021). Larval light/dark research lacks standardisation of the
length and lighting conditions of the acclimation period (Table 1). The effects of the length and
lighting conditions of acclimation within the behavioural monitoring unit on light/dark assay
locomotor responses were addressed by comparing no acclimation, five-minute, ten-minute, 20-
minute or 30-minute acclimation periods (Fig. 5), as well as varying the lighting conditions (Fig. 6).
Our findings in Figure 5 show a significant effect of acclimation period length on locomotor
responses (one-way ANOVA: $F_{(4,\ 75)} = 7.249$, < 0.0001). 30-minute acclimation produced significantly
greater dark phase locomotion compared to the other tested acclimation periods (Fig. 5F).
Therefore, we chose a 30-minute acclimation for all further experimentation due to the increased
dark phase responsiveness.

273
Figure 5. The light/dark response is unaffected by acclimation period length. The light/dark mean
locomotor responses represented in minutes for (A) no acclimation, (B) five-minute acclimation in
the light, (C) ten-minute acclimation in the light, (D) 20-minute acclimation in the light and (E) 30-
minute acclimation in the light. (F) represents the mean dark phase locomotor responses for each
acclimation period length.

Varying the lighting conditions (light or dark) during the acclimation period revealed a
significant effect on light (one-way ANOVA: $F_{(1,\ 30)} = 4.611$, $= 0.04$) and dark phase (one-way ANOVA:
$F_{(1,\ 30)} = 18.09$, < 0.0001) locomotion (Fig. 6). Therefore, due to the increased overall locomotion,
light acclimation was selected for further experimentation.

283
284   **Figure 6. Lighting conditions in the acclimation period do not affect the subsequent light/dark**

285   **assay behavioural response.** Larvae were acclimated in the light (A) or dark (B) for 30 minutes and

286   then underwent the standardised light/dark assay (section 2.3 of this paper).

3.3. The light/dark assay is suitable for assessing the effects of chemicals on larval behaviour

We next validated the assay in the context of behavioural pharmacology by examining larval

behavioural responses to different concentrations of ethanol. Ethanol was selected due to its

widespread availability and common use in larval light/dark assay research (Davis et al., 2021; Irons

et al., 2010; MacPhail et al., 2009). We also sought to identify meaningful and effective

interpretations of the light/dark assay data (Fig. 7). By applying these straightforward analyses,

distinctive concentration-dependent altered locomotor responses in both light and dark phases were

revealed.

Figure 7. Concentration-dependent effects of ethanol on larval behaviour in the light/dark assay. The minute analysis (A) is represented as mean percentage of baseline movement over the 30-minute recording period. The heat map (B) and radar chart (D) show light phase responses and (C) and (E) show dark phase responses.

Figure 7A shows that ethanol exposure has a significant effect upon both light and dark phase locomotion (linear mixed model analysis (lights effect): β = 81.58, SE = 3.30, t = 24.76). To compare this model to the null model, an LRT was conducted. The LRT showed a significant difference between the fitted model and the null model, $\chi^2(1)$ = 543.11, p < 0.0001. Specifically, low concentrations of ethanol (1.0%) increase locomotion in the dark phase compared to control larvae. 3.5% and 4.0% exposed larvae had decreased locomotion in the dark phase compared to controls (Fig. 7A). The fish exposed to the 2.0% ethanol appeared to increase locomotion in the light phase

307 compared to control larvae with relatively unchanged dark phase locomotion. We hypothesised that
308 this altered locomotion was reduced responsiveness to the light/dark stimuli (Fig. 8). However,
309 further analysis disregarded our hypothesis with 2.0% ethanol exposed larvae having no significant
310 difference in light phase locomotion compared to controls following a student's t-test analysis (Paired
311 t-test, t(15) = -1.3924,   = 0.1847).

312
Figure 8. Increased light phase responsiveness with exposure to 2.0% ethanol in zebrafish larvae.
The minute analysis (A) and light vs dark average (B and C) are represented as mean percentage of
baseline movement over the 30-minute recording period.

To gain more intricate concentration-dependent behavioural responses, we next furthered
our analysis and interpretation of the data by adapting heat maps and radar charts (Haigis (2022) and
Steele (2018)) to the larval light/dark analysis. The type of swim movement that is increased
following addition of low concentrations of ethanol is 'total movements' (one-way ANOVA: $F_{(5, 90)}$ =
0.856,   = 0.514) (Fig. 7C and E). 4.0%-exposed larvae had equivalent darting responses to control
larvae; however, they had reduced total and steady movements (Fig. 7D and E). Furthermore, in the
light phase all concentrations had reduced total movement compared to control larvae (one-way
ANOVA: $F_{(5,90)}$ = 1.393,   = 0.235) (Fig. 7B and D). 2.0%-exposed larvae had increased darting
responses compared to controls and 4.0%-exposed fish had increased slow swim responses
compared to the other concentrations. These specific swim observations we have highlighted here
are not seen on the line graph and bar chart. Therefore, the radars and heat maps facilitate
interpretation of the data and provide insight to more advanced swim responses which cannot be
achieved with basic movement analyses.

Next, using the same data, we examined time-dependent effects of ethanol exposures for
the first 10-minutes, 10-20 minutes and 20-30 minutes post-exposure (Fig. 9). By the first dark phase,
there is a significant reduction in locomotion with 4% ethanol compared to controls (one-way
ANOVA: $F_{(5, 87)}$ = 5.55,   < 0.0001) (Fig. 9B). The significant increase in locomotion with 1.0% ethanol
is not observed until the second dark phase (one-way ANOVA: $F_{(5, 84)}$ = 13.02,   < 0.0001) (Fig. 9D).

4.0% ethanol exposed larvae had reduced dark phase locomotion as well (Fig. 9D). By the final 10-

minutes, the 2.0% exposed larvae showed increased light phase locomotion (one-way ANOVA: $F_{(5, 80)}$

= 14.45,   < 0.0001) (Fig. 9E). This demonstrates the potential of the assay to examine behavioural

expressions of pharmacokinetics/dynamics and highlights the importance of evaluating time-

dependence in addition to overall recording responses over the full 30-minutes.

Figure 9. Time-dependent effects of ethanol exposure on larval light/dark behaviour. Mean
locomotor responses represented as a percentage of baseline movement for (A and B) light and dark
phase one (0-10-minutes post-exposure), (C and D) light and dark phase two (10-20-minutes post-
exposure) and (E and F) light and dark phase three (20-30-minutes post-exposure).

3.4. The light/dark assay is suitable for assessing behavioural phenotypes of transgenic
zebrafish lines

We next validated the assay by examining the phenotypes of *adgrl3.1$^{-/-}$* larvae, which have
previously been shown to express hyperactivity (Sveinsdóttir et al., 2023). In previous work, the
noradrenergic reuptake inhibitor, MPH, a therapeutic drug that reduces hyperactivity in children with
ADHD, has been shown to rescue the hyperactive phenotype. We examined the effects of this drug
on the light/dark assay responses at concentrations determined from previous publications (Lange et
al., 2012).

Our findings in figure 10 show a significant increase in dark phase locomotion in the *adgrl3.1^{-}*
$^{/-}$ larvae compared to WT larvae, which is partially rescued following addition of 5 and 20 µM MPH
(linear mixed model analysis (lights effect): β = 140.24 (SE = 2.6, t = 53.97) (Fig. 10A). The LRT showed
a significant difference between the fitted model and the null model, $\chi2$ (7) = 2085.4, p < 0.0001. No
significant changes were observed for the WT larvae treated with 0.05 µM and 5 µM MPH compared
to untreated WT fish in either light condition, but 20 µM MPH significantly reduced dark phase
responses (Fig. 10A).

The *adgrl3.1$^{-/-}$* larvae express phenotypic hyperactivity with increased dark phase locomotion
compared to WT (dark phase linear mixed model analysis (strain concentration): β = 79.68 (SE =
31.55, t = 2.53) (Fig. 10). The LRT showed a significant difference between the fitted model and the
null model, $\chi2$ (7) = 61.64, p < 0.0001. 5 µM MPH exposed *adgrl3.1$^{-/-}$* larvae display similar
movement responses to that of WT fish, with reduced total movements compared to *adgrl3.1$^{-/-}$*
larvae. 20 µM MPH exposed *adgrl3.1$^{-/-}$* larvae display similar behavioural responses to 5 µM MPH
exposed; however, results in significant reductions in WT total movements. These findings suggest
that 5 µM MPH is optimal for reversing the hyperactive phenotype of *adgrl3.1$^{-/-}$* larvae without
impacting WT responses compared to the other tested concentrations. These observations highlight
the value of including these additional analyses and interpretations of the larval light/dark data.

Like MPH, ATO is a noradrenergic reuptake inhibitor which reduces hyperactivity in children
with ADHD and has previously been shown to reverse the hyperactive phenotype of *adgrl3.1$^{-/-}$* larvae
(Sveinsdóttir et al., 2023). However, studies have reported that a major side-effect of ATO in patients
is sedation (Bastiaens, 2007). Therefore, here we examined the effects of ATO on the light/dark assay

response at concentrations previously used with *adgrl3.1*$^{-/-}$ larvae with the objective to determine

the translational relevance of our assay for evaluating complex effects such as sedation (Sveinsdóttir

et al., 2023).

Our findings show that like 20 μM MPH, the WT larvae exposed to ATO experienced

significant reductions in locomotion compared to untreated WT larvae (linear mixed model analysis

(lights effect): *β* = 121.06 (SE = 3.52, *t* = 34.43) (Fig. 10F). The LRT showed a significant difference

between the fitted model and the null model, χ2(7) = 104.78, *p* < 0.0001. *adgrl3.1*$^{-/-}$ larvae exposed

to 2.93 μM and 10.2 μM ATO had reduced dark phase locomotion compared to untreated WT fish

(Fig. 10F). However, 5.87 μM ATO exposed *adgrl3.1*$^{-/-}$ larvae had unchanged dark phase locomotion

compared to untreated WT fish (Fig. 10F).

ATO exposure has a significant effect on both light and dark phase swim phenotypes (Fig.

10G-J). The radar charts and heat maps demonstrate that the major differences between the WT and

adgrl3.1$^{-/-}$ larvae exposed to ATO are in the total movement responses. The WT larvae exposed to all

three ATO concentrations experienced significantly reduced total movement compared to untreated

WT larvae in both light and dark phases (Fig. 10G-J). Comparatively, the *adgrl3.1*$^{-/-}$ 5.87 μM ATO

treated larvae had comparable total movements compared to untreated WT larvae. Our ATO

observations highlight the importance of including the additional analyses for depicting adverse

effects of therapeutic substances in larval zebrafish and promotes the translational relevance of our

assay.

Figure 10. The effects of AB wild type and transgenic adgrl3.1$^{-/-}$ larval phenotypes on light/dark locomotion as well as the potential for assessing therapeutic agents. The minute analyses (A and F) are represented as mean percentage of baseline movement over the 30-minute recording period. The heat maps (B and G) and radar charts (D and I) show light phase responses and (C and H) and (E and J) show dark phase responses.

4. Discussion

Here, we aimed to determine the optimal conditions for the light/dark behavioural assay in 4 dpf zebrafish larvae. Our primary objective was to develop a robust and reproducible light/dark assay for 4 dpf zebrafish larvae with an in-depth analysis plan, and to validate the assay both pharmacologically, and with genetically modified larvae. The premise of this study was for providing a standardised assay for future larval research in neuropsychopharmacology and biological psychiatry.

There were several key findings here that should be highlighted. First, high levels of between-batch and inter-individual variability were found, which were corrected with normalisation. Second, we identified that a period of five-minutes for each dark and light period was optimal in terms of locomotor activity in 4 dpf larvae. A 30-minute light phase acclimation period resulted in significantly increased dark phase larval locomotion. We confirmed the pharmacological predictivity of the assay using ethanol, which caused low concentration hyperlocomotion and high concentration hypolocomotion, as predicted. Finally, the behavioural phenotypes of transgenic ($adgrl3.1^{-/-}$) larvae were reported on, where hyperlocomotion behaviour was rescued with both MPH and ATO. We also demonstrate the translational relevance of our assay for observing potential adverse effects of therapeutic substances, with ATO exposure resulting in hypolocomotion in WT larvae. As well as validation of our assay with transgenic larvae and ethanol, we have provided a comprehensive overview of alternative methods of analysis and interpretation of the data.

In previous work, we had noted variations in larval responses between different laboratories -- for example, if larvae are transported to different sites, or when working with collaborators – even when running identical light/dark assay protocols with fish bred at the same facility (*see SD 2*). Larval zebrafish variability in behavioural responses is well established (Jacobs & Ryu, 2023) with both genetic and environmental factors impacting batch-batch variability (Xie et al., 2019). For instance, hatching time has a significant effect upon dark phase responsiveness in the light/dark assay; our time-of-day analysis identified no significant effects upon the behavioural responses (Rock et al., 2022). However, we did observe large variations in behavioural responses during our time-of-day analysis, which was rectified with normalisation (*see SD 3*). Therefore, normalisation of the data is essential (Xie et al., 2019). We established a normalisation technique where the dataset of interest is analysed as a percentage against the average baseline locomotor responses, which was able to overcome the batch variability, time-of-day variability and variability between laboratories *(see SD 2).*

We encourage normalisation of the light/dark data for facilitating statistical comparisons and
collaboration between research laboratories (Xie et al., 2019).

Development of the assay involved confirmation of the optimal transition period length and
number of repetitions. Researchers have previously used a wide range of different transition periods
with further variation in the number of repetitions in the assay (Table 1). Here, we evaluated the
protocols employed by other researchers (Acevedo-Canabal et al., 2019; Bachour et al., 2020; Haigis
et al., 2022), as well as assessing our own developed protocol. Light to dark transitions were used, as
our preliminary findings showed no significant difference between light to dark or dark to light
transitions for larval locomotion *(SD 4)*. We report that the ten-minute and five-minute protocols are
both suitable for assessing behavioural responses in 4 dpf zebrafish larvae. We also suggest that the
five-minute protocol is superior due to its time-effectiveness, enhanced throughput and sustained
increased dark phase locomotion compared to the ten-minute protocol. The larvae exposed to the
two-minute transition and fifteen-minute transition protocols had reduced dark phase locomotor
responses by the final dark phases compared to the first (Fig. 3B and H). This indicates that the
longer paradigms are unsuitable for 4 dpf light/dark assay studies due to reduced locomotion, which
hinders the potential use of the assay for assessing drug effects for example. These findings are
supported by Maeda (2021) who demonstrated that larvae exposed to 15-minute intervals had
gradual decreases in dark phase locomotion. They did not however observe this with the five-minute
protocol of the same length. A possible reason for the reduced locomotion observed could be
habituation to the dark (Tuz-Sasik et al., 2022). Habituation is non-associative learning defined here
as a decrease in dark phase locomotion following repeated exposure to the light/dark transitions
(Tuz-Sasik et al., 2022). In the wild, repeated exposure to harmless stimuli causes high energy
expenditure and motor fatigue. Therefore, habituation ensures the fish conserves energy for survival
(Tuz-Sasik et al., 2022). Another possible reason for the reduced dark phase locomotion may be
exhaustion brought-on by excessive energy use (Tuz-Sasik et al., 2022). However, confirmation of
these hypotheses requires further investigation.

Previous research has highlighted the importance of the length and lighting conditions of the
acclimation period used prior to behavioural evaluation with fish (Makaras et al., 2021). In these
studies, the effect of changing the acclimation period conditions on locomotor responses was also
observed. Using our reported protocol, we identified a significant increase in dark phase locomotion
with the 30-minute acclimation period (Fig. 5) as well as increased overall locomotion with light
acclimation compared to dark (Fig. 6). Our evidence therefore supports the use of a 30-minute light
phase acclimation period for increased locomotor responses in 4 dpf larvae.

It has been suggested that the zebrafish circadian rhythm has no influence upon light/dark
assay locomotion (Krylov et al., 2021). Our findings in *SD 3* confirms this with no difference in dark
phase locomotion regardless of the time of day of recording. However, observations by Fitzgerald
(2019), Kristofco (2016) and MacPhail (2009), disagreed with our findings. The discrepancy between
observations could be methodology-dependent, which highlights the importance of the
experimental parameters of the light/dark assay with zebrafish larvae (Fitzgerald et al., 2019). Slight
differences in dark phase locomotion at different times of the day was observed when using the
distance travelled data (*SD 2A*). However, this variation is removed through normalisation to baseline
(*SD 2B*). Therefore, we strongly encourage incorporation of normalisation to baseline findings for
consistent and comparable results throughout the day, alleviating the need to limit behavioural work
to a certain time of day.

We aimed to further validate the assay by administering ethanol, which we selected due to
its robust concentration-dependent behavioural effects (low concentration hyperactivity and high
concentration hypoactivity in the dark phase (Cohen et al., 1997)), its widespread availability and
common global abuse (McLellan, 2017) and its frequent use in larval light/dark assay research (Davis
et al., 2021; Irons et al., 2010; MacPhail et al., 2009). The diverse and well understood
neuropharmacology of ethanol (low concentration hyperactivity and high concentration hypoactivity)
allowed for suitable assay validation. Low concentration ethanol hyperactivity has previously been
reported in older larvae using the light/dark assay (MacPhail et al., 2009) and a similar finding was
reported here with 1.0 % ethanol exposure (Fig. 7). This low concentration hyperactivity is a result of
elevated dopamine levels and is commonly reported as the positive reinforcing effects of ethanol in
rodents (Cohen et al., 1997). Mid-range concentrations are known to result in behavioural
disinhibition in humans (Harrison et al., 2017) and similar dissociative-like swim behavioural
responses have previously been reported (MacPhail et al., 2009). This dissociative-like response can
be seen here in larvae exposed to 2.0 % ethanol (Fig. 7). High concentration hypoactivity stems from
ethanol's sedative-like properties in rodents and mammals, which are attributed to its GABAergic
effects (Clayman & Connaughton, 2022). Like previous publications, (Irons et al., 2010; MacPhail et
al., 2009), we report ethanol-induced sedative-like responses with exposure to high concentrations
(Fig. 7). These observations with ethanol in the light/dark assay indicate consistent
neuropharmacological mechanisms between larval zebrafish and previous findings and suggests that
the assay is suitable as a first-tier model for predicting drug and chemical effects.

We further validated the assay for neuropsychopharmacology evaluation using an
established transgenic line, *adgrl3.1$^{-/-}$*, which was employed because previous studies had shown

robust hyperactivity compared to WT (Sveinsdóttir et al., 2023). This hyperactivity was successfully

reduced with exposure to both ATO and MPH (Fig. 10), which has previously been reported in ATO

treated adult *adgrl3.1*[-/-] zebrafish (Fontana et al., 2023) and in 6 dpf larval *adgrl3.1*[-/-] zebrafish

treated with ATO and MPH (Lange et al., 2012). Here, we demonstrated the ability of our assay to

distinctly establish more specific drug effects by demonstrating the common adverse effect,

sedation, in the WT ATO exposed larvae (Bastiaens, 2007) (Fig.10). In contrast, MPH only reduced

movement in WT larvae at the top concentration, consistent with its stimulant mechanism of action

(Challman & Lipsky, 2000). These observations indicate this light/dark assay is appropriate for

differentiating and highlighting intrinsic and potentially complex effects of compounds.

There are several ways that the larval light/dark assay data can be presented, and typically

this takes the form of mean distance travelled (Haigis et al., 2022). Although this has advantages

(e.g., it is quick and uses the raw data), we believe this approach has its limitations. First, as

highlighted, when analysing the distance travelled data at face value there was major variation

between different laboratories, and even batches within the same laboratory, using fish bred in the

same facility and with identical experimental parameters. We postulate that this is associated with

environmental stressors, transport, or other generic inter-batch variability (Xie et al., 2019).

Secondly, different common swim phenotypes such as the freezing response cannot be accurately

identified with just distance travelled data (Kalueff et al., 2013). These limitations prompted us to

adopt a broader analytical approach (from Steele (2018)) and analyse responses as different swim

parameters. Haigis (2022) also analysed the data in this format to produce a radar chart; however,

they chose to normalise the data to control response and perform a $\log_2$ transformation. We similarly

normalised the data to baseline responses, followed by analysis of the different swim parameters.

Next, we normalised the individual swim responses to control larval response and performed a

logarithmic transformation to overcome any exponential increases in swimming behaviour which

was identified as impacting the readability of the radar charts and heat maps.

The swim parameters analysed here include total movement, darting movement, steady

swimming movement, slow swimming movement and zero movements. We believe these swim

phenotypes will facilitate more complex behavioural analysis with 4 dpf zebrafish larvae. We

demonstrate that the radar charts and heat maps can distinguish more precise movement patterns

and drug/genomic induced behavioural responses which cannot be ascertained solely from the line

graph or bar chart (Fig. 7 and 10). Darting swimming is an important swim factor to analyse because

it is an indicator of the initial dark escape/light-seeking response as well as a sign of anxiety-like

responses (Haigis et al., 2022). Slow swimming responses highlight more clearly any reduced swim

response whilst zero movements show the innate freezing common with zebrafish as well as
complete loss of movement because of sedation, paralysis or death (Kalueff et al., 2013). A steady
swim response was also included, which we postulate is continuous swimming which is not
commonly observed in control 4 dpf larvae. As highlighted by, Haigis (2022), including analyses such
as the radar chart permits identification of behavioural responses which may go unnoticed in
standard analyses. We believe our findings highlight the radar chart and heat map value in analysis
and strongly encourage the inclusion of both to facilitate the readability and interpretation of future
light/dark larval behavioural research.

5. Potential applications

Our validated assay promotes and encourages the use of 4 dpf zebrafish larvae for
neuropsychopharmacology experimentation, which is beneficial from a 3R's of animal research
perspective (Sneddon et al., 2017). The model can be used for studying potential therapeutic agents.
For example, we used adgrl3.1$^{-/-}$ transgenic larvae to observe the effects of MPH and ATO, known
anti-ADHD drugs, (Yang et al., 2018), as well as assessing the genetic behavioural phenotypes. There
is also the potential for the model to be used for toxicology and ecotoxicology research (Haigis et al.,
2022). Previous research has identified the use of the light/dark assay for assessing anxiogenic and
anxiolytic substances (Facciol et al., 2019). However, here we suggest the assay can be used to screen
for a range of pharmacologically induced behavioural parameters, including hyperlocomotion and
sedation. Additionally, we believe a potential application for the assay would be to assess
environmental effects on behaviour (e.g., water condition) (Rock et al., 2022). These potential
applications we believe highlight the unique flexibility of this larval assay.

6. Limitations

We acknowledge several limitations. First, although we highlight excellent translational potential
of the 4 dpf behavioural data, they remain a preliminary model only. Analysis of more complex
behaviours is likely not possible at this age due to their underdeveloped CNS (Blader, 2000). Second,
this method is reliant upon accurate behavioural tracking and therefore the tracking settings may
need adjustment prior to investigation. Third, our new analytical method is more complex than
simplistic averaging of distance travelled; however, the custom R script available on OSF overcomes
this and facilitates analysis.

7. Conclusions

Here we present a standardised light/dark behavioural assay for use in 4 dpf zebrafish larvae,
which was validated using exposure to ethanol with wild type AB and exposure to ADHD drugs with

adgrl3.1$^{-/-}$ transgenic larvae. We report that the five-minute transition period is optimal for consistent

and reproducible locomotor responses and that an acclimation period is not required for robust

responses to be seen, which improves the throughput of the assay. We present the influence of time

of day on 4 dpf larval responses whilst providing a comprehensive analysis of different methods of

interpretation and presentation of the behavioural data. Finally, this work has strong 3Rs

applications. Despite some limitations due to their reduced ability to determine complex behaviours,

we strongly encourage the incorporation of 4 dpf studies prior to adult or mammalian

neuropsychopharmacology studies to reduce the overall number of protected animals used in

studies.

8.   Declarations/Acknowledgments

Funding: CH was funded by DSTL (UK). JK and MP received funding from the NC3Rs (NC/W00092X/1)

Conflicts of interest/Competing interests: There is no conflict of interest to declare.

Ethics approval: All experiments were approved by the University of Portsmouth Animal Welfare and
Ethical Review Board, and under license from the UK Home Office (Animals [Scientific Procedures]
Act, 1986, license number PP8708123).

Consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: All data and analysis materials are available on the Open Science
Framework. https://osf.io/k7y2c/?view_only=61e23ea4bb2d452dab76646931e3f332

Code availability: Not applicable.

Authors' contributions:

Conceptualization: CH, JK, MP. Formal analysis: CH. Funding acquisition: JK, MP. Investigation: CH.
Methodology: CH, JK, MP. Project administration: MP. Resources: JK, MP. Supervision: JK, MP.
Visualization: CH. Writing – original draft preparation: CH. Writing – review & editing: CH, JK, MP.

9. References

Acevedo-Canabal, A., Colón-Cruz, L., Rodriguez-Morales, R., Varshney, G. K., Burgess, S., González-
Sepúlveda, L., Yudowski, G., & Behra, M. (2019). Altered swimming behaviors in zebrafish larvae
lacking cannabinoid receptor 2. *Cannabis and Cannabinoid Research*, *4*(2), 88.
https://doi.org/10.1089/CAN.2018.0025

Animals (Scientific Procedures) Act 1986, 14 (1986).
https://www.legislation.gov.uk/ukpga/1986/14/contents

Bachour, R. L., Golovko, O., Kellner, M., & Pohl, J. (2020). Behavioral effects of citalopram, tramadol,
and binary mixture in zebrafish (*Danio rerio*) larvae. *Chemosphere*, *238*, 124587.
https://doi.org/10.1016/J.CHEMOSPHERE.2019.124587

Basnet, R. M., Zizioli, D., Taweedet, S., Finazzi, D., & Memo, M. (2019). Zebrafish larvae as a
behavioral model in neuropharmacology. *Biomedicines*, *7*(1).
https://doi.org/10.3390/biomedicines7010023

Bastiaens, L. (2007). Effectiveness and tolerability of atomoxetine in a real-world ADHD population:
nonrandomized comparison with stimulants. *Psychiatry (Edgmont (Pa.:Township))*, *4*(12), 44–
48.

Beker van Woudenberg, A., Wolterbeek, A., te Brake, L., Snel, C., Menke, A., Rubingh, C., de Groot,
D., & Kroese, D. (2013). A category approach to predicting the developmental (neuro) toxicity of
organotin compounds: The value of the zebrafish (*Danio rerio*) embryotoxicity test (ZET).
Reproductive Toxicology, *41*, 35–44. https://doi.org/10.1016/j.reprotox.2013.06.067

Best, C., Kurrasch, D. M., & Vijayan, M. M. (2017). Maternal cortisol stimulates neurogenesis and
affects larval behaviour in zebrafish. *Scientific Reports*, *7*, 40905.
https://doi.org/10.1038/srep40905

Blader, P. (2000). Zebrafish developmental genetics and central nervous system development. *Human
Molecular Genetics*, *9*(6), 945–951. https://doi.org/10.1093/hmg/9.6.945

Cassar, S., Adatto, I., Freeman, J. L., Gamse, J. T., Iturria, I., Lawrence, C., Muriana, A., Peterson, R. T.,
Van Cruchten, S., & Zon, L. I. (2020). Use of zebrafish in drug discovery toxicology. *Chemical
Research in Toxicology*, *33*(1), 95–118.
https://doi.org/10.1021/ACS.CHEMRESTOX.9B00335/ASSET/IMAGES/LARGE/TX9B00335_0008.J
PEG

Challman, T. D., & Lipsky, J. J. (2000). Methylphenidate: Its pharmacology and uses. *Mayo Clinic
Proceedings*, *75*(7), 711–721. https://doi.org/10.4065/75.7.711

Chatzimitakos, T., Chousidis, I., Leonardos, D., Stalikas, C., & Leonardos, I. (2022). In the swim of
cannabis: developmental toxicity and metabolomic pathway alterations of zebrafish larvae
exposed to THC for the assessment of its potential environmental and human health impact.
Molecules, *27*(17). https://doi.org/10.3390/MOLECULES27175506/S1

Clayman, C. L., & Connaughton, V. P. (2022). Neurochemical and behavioral consequences of ethanol
and/or caffeine exposure: effects in zebrafish and rodents. *Current Neuropharmacology*, *20*(3),
560–578. https://doi.org/10.2174/1570159X19666211111142027

Cohen, C., Perrault, Gh., & Sanger, D. J. (1997). Evidence for the involvement of dopamine receptors
in ethanol-induced hyperactivity in mice. *Neuropharmacology*, *36*(8), 1099–1108.
https://doi.org/10.1016/S0028-3908(97)00100-7

Crilly, S., Njegic, A., Laurie, S. E., Fotiou, E., Hudson, G., Barrington, J., Webb, K., Young, H. L., Badrock,
A. P., Hurlstone, A., Rivers-Auty, J., Parry-Jones, A. R., Allan, S. M., & Kasher, P. R. (2018). Using
zebrafish larval models to study brain injury, locomotor and neuroinflammatory outcomes
following intracerebral haemorrhage. *F1000Research 2018 7:1617*, *7*, 1617.
https://doi.org/10.12688/f1000research.16473.2

Davis, R., Luchtenburg, F., Richardson, M., Schaaf, M., Tudorache, C., & Slabbekoorn, H. (2021). The
importance of individual variation for the interpretation of behavioural studies: ethanol effects
vary with basal activity level in zebrafish larvae. *Psychopharmacology*, *238*(11), 3155–3166.
https://doi.org/10.1007/s00213-021-05932-6

Facciol, A., Tran, S., & Gerlai, R. (2019). A standardized tank design for the light dark task in zebrafish.
BIO-PROTOCOL, *9*(14). https://doi.org/10.21769/BioProtoc.3306

Fitzgerald, J. A., Kirla, K. T., Zinner, C. P., & Vom Berg, C. M. (2019). Emergence of consistent intra-
individual locomotor patterns during zebrafish development. *Scientific Reports*, *9*(1), 13647.
https://doi.org/10.1038/s41598-019-49614-y

Fontana, B. D., Reichmann, F., Tilley, C. A., Lavlou, P., Shkumatava, A., Alnassar, N., Hillman, C.,
Karlsson, K. Æ., Norton, W. H. J., & Parker, M. O. (2023). adgrl3.1-deficient zebrafish show
noradrenaline-mediated externalizing behaviors, and altered expression of externalizing
disorder-candidate genes, suggesting functional targets for treatment. *Translational Psychiatry*,
13(1), 304. https://doi.org/10.1038/s41398-023-02601-4

Fraser, T. W. K., Khezri, A., Jusdado, J. G. H., Lewandowska-Sabat, A. M., Henry, T., & Ropstad, E.
(2017). Toxicant induced behavioural aberrations in larval zebrafish are dependent on minor
methodological alterations. *Toxicology Letters*, *276*, 62–68.
https://doi.org/10.1016/j.toxlet.2017.05.021

Haigis, A. C., Ottermanns, R., Schiwy, A., Hollert, H., & Legradi, J. (2022). Getting more out of the
zebrafish light dark transition test. *Chemosphere*, *295*, 133863.
https://doi.org/10.1016/J.CHEMOSPHERE.2022.133863

Harrison, N. L., Skelly, M. J., Grosserode, E. K., Lowes, D. C., Zeric, T., Phister, S., & Salling, M. C.
(2017). Effects of acute alcohol on excitability in the CNS. *Neuropharmacology*, *122*, 36–45.
https://doi.org/10.1016/j.neuropharm.2017.04.007

Hasumi, A., Maeda, H., & Yoshida, K. I. (2020). Analyzing cannabinoid-induced abnormal behavior in
a zebrafish model. *PLoS ONE*, *15*(10). https://doi.org/10.1371/JOURNAL.PONE.0236606

Hoyberghs, J., Bars, C., Ayuso, M., Van Ginneken, C., Foubert, K., & Van Cruchten, S. (2021). DMSO
concentrations up to 1% are safe to be used in the zebrafish embryo developmental toxicity
assay. *Frontiers in Toxicology*, *3*. https://doi.org/10.3389/ftox.2021.804033

Irons, T. D., MacPhail, R. C., Hunter, D. L., & Padilla, S. (2010). Acute neuroactive drug exposures alter
locomotor activity in larval zebrafish. *Neurotoxicology and Teratology*, *32*(1), 84–90.
https://doi.org/10.1016/J.NTT.2009.04.066

Jacobs, E. A. K., & Ryu, S. (2023). Larval zebrafish as a model for studying individual variability in
translational neuroscience research. *Frontiers in Behavioral Neuroscience*, *17*.
https://doi.org/10.3389/fnbeh.2023.1143391

Jin, M., Zhang, Y., Ye, J., Huang, C., Zhao, M., & Liu, W. (2010). Dual enantioselective effect of the
insecticide bifenthrin on locomotor behavior and development in embryonic-larval zebrafish.
Environmental Toxicology and Chemistry, *29*(7), 1561–1567. https://doi.org/10.1002/etc.190

Kalueff, A. V., Gebhardt, M., Stewart, A. M., Cachat, J. M., Brimmer, M., Chawla, J. S., Craddock, C.,
Kyzar, E. J., Roth, A., Landsman, S., Gaikwad, S., Robinson, K., Baatrup, E., Tierney, K., Shamchuk,
A., Norton, W., Miller, N., Nicolson, T., Braubach, O., … Schneider, and the Z. N. H. (2013).
Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. *Zebrafish*, *10*(1), 70–
86. https://doi.org/10.1089/zeb.2012.0861

Kirla, K. T., Erhart, C., Groh, K. J., Stadnicka-Michalak, J., Eggen, R., Schirmer, K., & Kraemer, T. (2021).
Zebrafish early life stages as alternative model to study 'designer drugs': Concordance with
mammals in response to opioids | Elsevier Enhanced Reader. *Toxicology and Applied*
Pharmacology, *419*, 115483. https://doi.org/10.1016/j.taap.2021.115483

Kristofco, L. A., Cruz, L. C., Haddad, S. P., Behra, M. L., Chambliss, C. K., & Brooks, B. W. (2016). Age
matters: Developmental stage of *Danio rerio* larvae influences photomotor response thresholds
to diazinion or diphenhydramine. *Aquatic Toxicology*, *170*, 344–354.
https://doi.org/10.1016/j.aquatox.2015.09.011

Krylov, V. V., Izvekov, E. I., Pavlova, V. V., Pankova, N. A., & Osipova, E. A. (2021). Circadian rhythms in
zebrafish (*Danio rerio*) behaviour and the sources of their variability. *Biological Reviews*, *96*(3),
785–797. https://doi.org/10.1111/brv.12678

Lange, M., Norton, W., Coolen, M., Chaminade, M., Merker, S., Proft, F., Schmitt, A., Vernier, P., Lesch,
K.-P., & Bally-Cuif, L. (2012). The ADHD-susceptibility gene lphn3.1 modulates dopaminergic
neuron formation and locomotor activity during zebrafish development. *Molecular Psychiatry*,
17(9), 946–954. https://doi.org/10.1038/mp.2012.29

Legradi, J., el Abdellaoui, N., van Pomeren, M., & Legler, J. (2015). Comparability of behavioural
assays using zebrafish larvae to assess neurotoxicity. *Environmental Science and Pollution*
Research, *22*(21), 16277–16289. https://doi.org/10.1007/s11356-014-3805-8

Leys, C., Ley, C., Klein, O., Bernard, P., & Licata, L. (2013). Detecting outliers: Do not use standard
deviation around the mean, use absolute deviation around the median. *Journal of Experimental*
Social Psychology, *49*(4), 764–766. https://doi.org/10.1016/j.jesp.2013.03.013

Liang, X., Souders, C. L., Zhang, J., & Martyniuk, C. J. (2017). Tributyltin induces premature hatching
and reduces locomotor activity in zebrafish (*Danio rerio*) embryos/larvae at environmentally
relevant levels. *Chemosphere*, *189*, 498–506.
https://doi.org/10.1016/j.chemosphere.2017.09.093

Licitra, R., Martinelli, M., Jasinski, L. P., Marchese, M., Kiferle, C., & Fronte, B. (2021). in vivo
Evaluation of Cannabis sativa Full Extract on Zebrafish Larvae Development, Locomotion

Behavior and Gene Expression. *Pharmaceuticals (Basel, Switzerland)*, *14*(12).
https://doi.org/10.3390/PH14121224

Liu, Y., Carmer, R., Zhang, G., Venkatraman, P., Brown, S. A., Pang, C.-P., Zhang, M., Ma, P., & Leung, Y.
F. (2015). Statistical Analysis of Zebrafish Locomotor Response. *PLoS One*, *10*(10), e0139521.
https://doi.org/10.1371/journal.pone.0139521

MacPhail, R. C., Brooks, J., Hunter, D. L., Padnos, B., Irons, T. D., & Padilla, S. (2009). Locomotion in
larval zebrafish: Influence of time of day, lighting and ethanol. *NeuroToxicology*, *30*(1), 52–58.
https://doi.org/10.1016/j.neuro.2008.09.011

Maeda, H., Fukushima, N., & Hasumi, A. (2021). Standardized Method for the Assessment of
Behavioral Responses of Zebrafish Larvae. *Biomedicines*, *9*(8), 884.
https://doi.org/10.3390/biomedicines9080884

Makaras, T., Stankevičiūtė, M., Šidagytė-Copilas, E., Virbickas, T., & Razumienė, J. (2021). Acclimation
effect on fish behavioural characteristics: determination of appropriate acclimation period for
different species. *Journal of Fish Biology*, *99*(2), 502–512. https://doi.org/10.1111/jfb.14740

McLellan, A. T. (2017). Substance Misuse and Substance use Disorders: Why do they Matter in
Healthcare? *Transactions of the American Clinical and Climatological Association*, *128*, 112.
/pmc/articles/PMC5525418/

Mundy, P. C., Pressly, B., Carty, D. R., Yaghoobi, B., Wulff, H., & Lein, P. J. (2021). The efficacy of γ-
aminobutyric acid type A receptor (GABA AR) subtype-selective positive allosteric modulators in
blocking tetramethylenedisulfotetramine (TETS)-induced seizure-like behavior in larval zebrafish
with minimal sedation. *Toxicology and Applied Pharmacology*, *426*, 115643.
https://doi.org/10.1016/J.TAAP.2021.115643

Nixon, J., Abramovici, H., Cabecinha, A., Martinez-Farina, C., Hui, J., & Ellis, L. (2021). Assessing the
bioactivity of cannabis extracts in larval zebrafish. *Journal of Cannabis Research*, *3*(1), 44.
https://doi.org/10.1186/S42238-021-00103-Y

Okeke, E. S., Qian, X., Che, J., Mao, G., Chen, Y., Xu, H., Ding, Y., Zeng, Z., Wu, X., & Feng, W. (2022).
Transcriptomic sequencing reveals the potential molecular mechanism by which
Tetrabromobisphenol A bis (2-hydroxyethyl ether) exposure exerts developmental neurotoxicity
in developing zebrafish (Danio rerio). *Comparative Biochemistry and Physiology Part C:*
Toxicology & Pharmacology, *262*, 109467. https://doi.org/10.1016/j.cbpc.2022.109467

Padilla, S., Hunter, D. L., Padnos, B., Frady, S., & MacPhail, R. C. (2011). Assessing locomotor activity
in larval zebrafish: Influence of extrinsic and intrinsic variables. *Neurotoxicology and Teratology*,
33(6), 624–630. https://doi.org/10.1016/j.ntt.2011.08.005

Pesavento, S., Bilel, S., Murari, M., Gottardo, R., Arfè, R., Tirri, M., Panato, A., Tagliaro, F., & Marti, M.
(2022). Zebrafish larvae: A new model to study behavioural effects and metabolism of fentanyl,
in comparison to a traditional mice model: *Https://Doi.Org/10.1177/00258024221074568*.
https://doi.org/10.1177/00258024221074568

Rock, S., Rodenburg, F., Schaaf, M. J. M., & Tudorache, C. (2022). Detailed Analysis of Zebrafish Larval
Behaviour in the Light Dark Challenge Assay Shows That Diel Hatching Time Determines
Individual Variation. *Frontiers in Physiology*, *13*.
https://doi.org/10.3389/FPHYS.2022.827282/FULL

Sales Cadena, M. R., Cadena, P. G., Watson, M. R., Sarmah, S., Boehm, S. L., & Marrs, J. A. (2021).
Zebrafish (Danio rerio) larvae show behavioral and embryonic development defects when
exposed to opioids at embryo stage. *Neurotoxicology and Teratology*, *85*, 106964.
https://doi.org/10.1016/J.NTT.2021.106964

Scarlett, K. R., Lovin, L. M., Steele, W. B., Kim, S., & Brooks, B. W. (2022). Identifying Behavioral
Response Profiles of Two Common Larval Fish Models to a Salinity Gradient. *Archives of*
Environmental Contamination and Toxicology, *83*(2), 180–192. https://doi.org/10.1007/s00244-
022-00951-6

Sneddon, L. U., Halsey, L. G., & Bury, N. R. (2017). Considering aspects of the 3Rs principles within
experimental animal biology. *Journal of Experimental Biology*, *220*(17), 3007–3016.
https://doi.org/10.1242/jeb.147058

Steele, W. B., Kristofco, L. A., Corrales, J., Saari, G. N., Haddad, S. P., Gallagher, E. P., Kavanagh, T. J.,
Kostal, J., Zimmerman, J. B., Voutchkova-Kostal, A., Anastas, P., & Brooks, B. W. (2018).
Comparative behavioral toxicology with two common larval fish models: Exploring relationships
among modes of action and locomotor responses. *Science of The Total Environment*, *640–641*,
1587–1600. https://doi.org/10.1016/j.scitotenv.2018.05.402

Steenbergen, P. J., Richardson, M. K., & Champagne, D. L. (2012). *The Light–Dark Preference Test for*
Larval Zebrafish (Vol. 66). Humana Press, Totowa, NJ. https://doi.org/10.1007/978-1-61779-
597-8_2

Sveinsdóttir, H. S., Christensen, C., Þorsteinsson, H., Lavalou, P., Parker, M. O., Shkumatava, A.,
Norton, W. H. J., Andriambeloson, E., Wagner, S., & Karlsson, K. Æ. (2023). Novel non-stimulants
rescue hyperactive phenotype in an adgrl3.1 mutant zebrafish model of ADHD.
Neuropsychopharmacology, *48*(8), 1155–1163. https://doi.org/10.1038/s41386-022-01505-z

Thit, A., Skjolding, L. M., Selck, H., & Sturve, J. (2017). Effects of copper oxide nanoparticles and
copper ions to zebrafish (Danio rerio) cells, embryos and fry. *Toxicology in Vitro*, *45*, 89–100.
https://doi.org/10.1016/j.tiv.2017.08.010

Tuz-Sasik, M. U., Boije, H., & Manuel, R. (2022). Characterization of locomotor phenotypes in
zebrafish larvae requires testing under both light and dark conditions. *PLOS ONE*, *17*(4),
e0266491. https://doi.org/10.1371/journal.pone.0266491

Walter, K. M., Miller, G. W., Chen, X., Harvey, D. J., Puschner, B., & Lein, P. J. (2019). Changes in
thyroid hormone activity disrupt photomotor behavior of larval zebrafish. *NeuroToxicology*, *74*,
47–57. https://doi.org/10.1016/j.neuro.2019.05.008

Wang, Y., Liu, C., Deng, J., Xu, Q., Lin, J., Li, H., Hu, M., Hu, C., Li, Q., & Xu, X. (2023). Behavioral and
Sensory Deficits Associated with Dysfunction of GABAergic System in a Novel shank2-Deficient
Zebrafish Model. *International Journal of Molecular Sciences*, *24*(3).
https://doi.org/10.3390/ijms24032208

Widrick, J. J., Lambert, M. R., Kunkel, L. M., & Beggs, A. H. (2023). Optimizing assays of zebrafish
larvae swimming performance for drug discovery. *Expert Opinion on Drug Discovery*, *18*(6),
629–641. https://doi.org/10.1080/17460441.2023.2211802

Xie, R., Zhang, M., Venkatraman, P., Zhang, X., Zhang, G., Carmer, R., Kantola, S. A., Pang, C. P., Ma, P.,
Zhang, M., Zhong, W., & Leung, Y. F. (2019). Normalization of large-scale behavioural data

collected from zebrafish. *PloS One*, *14*(2), e0212234.
https://doi.org/10.1371/journal.pone.0212234

Yang, L., Chang, S., Lu, Q., Zhang, Y., Wu, Z., Sun, X., Cao, Q., Qian, Y., Jia, T., Xu, B., Duan, Q., Li, Y.,
Zhang, K., Schumann, G., Liu, D., Wang, J., Wang, Y., & Lu, L. (2018). A new locus regulating
MICALL2 expression was identified for association with executive inhibition in children with
attention deficit hyperactivity disorder. *Molecular Psychiatry*, *23*(4), 1014–1020.
https://doi.org/10.1038/mp.2017.74

Zaig, S., Scarpellini, C., & Montandon, G. (2021). Respiratory depression and analgesia by opioid
drugs in freely-behaving larval zebrafish. *ELife*, *10*. https://doi.org/10.7554/ELIFE.63407

Zhao, J., Xu, T., & Yin, D.-Q. (2014). Locomotor activity changes on zebrafish larvae with different
2,2⬚,4,4⬚-tetrabromodiphenyl ether (PBDE-47) embryonic exposure modes. *Chemosphere*, *94*,
53–61. https://doi.org/10.1016/j.chemosphere.2013.09.010

10. Supplementary Data

SD 1: Previous studies have shown that the size of well plate used for larval behavioural research significantly impacts the behavioural response (Widrick et al., 2023). However, here we demonstrate that any well plate size is suitable for 4 dpf zebrafish larval light/dark behavioural research (SD 1). No significant difference was seen following normalisation in both light and dark phases for the well plates. However, the lack of movement in the light phase with the 96-well plate is extreme and these floor effects means it is hard to observe any movement suppression with pharmacological interventions (SD 1). Therefore, we selected the 48-well plate for further analysis due to its enhanced throughput compared to the 24-well plate and improved time-efficiency for plating compared to the 96-well plate as well as increased reliability. However, our evidence suggests that well plate size should not impact the behavioural results seen if normalisation is performed.

SD 1: The minute analysis comparing the average locomotion for 24-well plates, 48-well plates and 96-well plates.

SD 2: We noted substantial differences between our two laboratories running identical light/dark protocols with fish obtained from the same fish facility (University of Portsmouth). We highlight the importance of normalising the data before analysis rather than using the distance travelled data (*SD* 2A-F). Baseline dark phase locomotion was significantly different between the two laboratories using distance travelled data (one-way ANOVA: $F_{(1, 30)}$ = 14.3, p < 0.0001) (*SD* 2C). No difference was observed between the light phases (one-way ANOVA: $F_{(1, 30)}$ = 1.028, p = 0.319). Comparatively, following normalisation to baseline no significant difference in locomotion was observed for either dark phase or light phase.

SD 2: Comparison of baseline data obtained from our two separate laboratories using fish from the same fish facility.
The data is presented as either distance travelled data as a line graph (A) or normalised data as a line graph (B). The dark
and light phase responses for distance travelled (C and E) and normalised data (D and F) are also shown.

SD 3: It has previously been reported that the time of day affects larval zebrafish behavioural

responses using the light/dark assay (Fitzgerald et al., 2019; Kristofco et al., 2016; MacPhail et al.,

2009). We observed no significant effect of time of (working) day (morning (09:00 – 11:59),

afternoon (12:00 and 14:29) and evening (14:30– 17:00)) on larval light/dark assay (SD3 A and B).

There is slight variability in dark phase locomotion using the distance travelled data (one-way

ANOVA: $F_{(2, 45)}$ = 1.183, p = 0.316); however, this is fully removed following normalisation (one-way

ANOVA: $F_{(2, 45)}$ = 0, p = 1).

SD 3. Time of day of the light/dark assay recording affects behavioural responses observed in 4 dpf zebrafish larvae represented as (A) distance travelled data and (B) normalised to baseline data.

915
916

918 **SD4**: We observed no significant difference in locomotion starting with light to dark vs dark to light

919 transitions in the light phase or dark phase (*SD* 4C and D).

921   **SD 4**: The behavioural effects of running the assay light to dark (A) or dark to light (B) reported as percentage of baseline
922   recording in minutes. The mean light phase (C) and dark phase (D) comparison is reported.