Perceptual error based on Bayesian cue combination

drives implicit motor adaptation

Zhaoran Zhang[1,†], Huijun Wang[1,†], Tianyang Zhang[1], Zixuan Nie[1], Kunlin Wei[1,2,3,4 *]

[1] School of Psychological and Cognitive Sciences, Peking University, Beijing, China

[2] Beijing Key Laboratory of Behavior and Mental Health, Beijing, China

[3] Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China

[4] National Key Laboratory of General Artificial Intelligence, Beijing, China

[†]Zhaoran Zhang and Huijun Wang are co-first authors.

[*]Correspondence to: Kunlin Wei

        School of Psychological and Cognitive Sciences

        Peking University

        5 Yiheyuan Road

        Beijing, 100871, China

        wei.kunlin@pku.edu.cn

Abstract

The sensorimotor system can recalibrate itself without our conscious awareness, a type of procedural learning whose computational mechanism remains undefined. Recent findings on implicit motor adaptation, such as over-learning from small perturbations and fast saturation for increasing perturbation size, challenge existing theories based on sensory errors. We argue that perceptual error, arising from the optimal combination of movement-related cues, is the primary driver of implicit adaptation. Central to our theory is the increasing sensory uncertainty of visual cues with increasing perturbations, which was validated through perceptual psychophysics (Experiment 1). Our theory predicts the learning dynamics of implicit adaptation across a spectrum of perturbation sizes on a trial-by-trial basis (Experiment 2). It explains proprioception changes and their relation to visual perturbation (Experiment 3). By modulating visual uncertainty in perturbation, we induced unique adaptation responses in line with our model predictions (Experiment 4). Overall, our perceptual error framework outperforms existing models based on sensory errors, suggesting that perceptual error in locating one's effector, supported by Bayesian cue integration, underpins the sensorimotor system's implicit adaptation.

Introduction

To achieve and sustain effective motor performance, humans consistently recalibrate their sensorimotor systems to adapt to both internal and external environmental disturbances (Berniker & Kording, 2008; Shadmehr et al., 2010; Wolpert et al., 2011). For instance, transitioning to a high-sensitivity gaming mouse, which drives cursor movement at an accelerated rate compared to a standard computer mouse, may initially result in decreased performance in computer-related tasks. However, humans are capable of rapidly adapting to this new visuomotor mapping within a short period of time. While conscious corrections can facilitate this adaptation process, our sensorimotor system often times adapts itself implicitly without our conscious efforts (Albert et al., 2021; Krakauer et al., 2019).

While recent research has intensively examined the interplay between explicit and implicit learning systems (Albert et al., 2022; Miyamoto et al., 2020), several characteristics of implicit motor adaptation have emerged that challenge traditional theories. Conventionally, motor adaptation is conceptualized as error-based learning, in which learning accrues in proportion to the motor error experienced (Cheng & Sabes, 2006; Donchin et al., 2003; Thoroughman & Shadmehr, 2000). However, implicit adaptation exhibits an overcompensation phenomenon where the extent of adaptation surpasses the error induced by visual perturbations (Kim et al., 2018; Morehead et al., 2017). Additionally, implicit adaptation manifests a saturation effect; it increases with perturbations but plateaus across a broad range of larger perturbations (Bond & Taylor, 2015; Kim et al., 2018; Morehead et al., 2017; Neville & Cressman, 2018). These observations of overcompensation and saturation are incongruent with prevailing

state-space updating models, which presuppose that incremental learning constitutes

only a fraction of the motor error (McDougle et al., 2015; Smith et al., 2006). Another

aspect of implicit adaptation that remains mechanistically unexplained pertains to its

impact on proprioception. In traditional motor adaptation, proprioception is biased

towards visual perturbation, maintaining a stable bias throughout the adaptation process

(Ruttle et al., 2016, 2021). In contrast, implicit adaptation initially biases proprioceptive

localization of the hand towards the visual perturbation, but this bias gradually drifts in

the opposite direction over time (Tsay et al., 2020).

Causal inference of motor errors has been suggested to explain the discounting of large

perturbations (Wei & Körding, 2009). However, the causal inference account predicts a

decline in adaptation with increasing perturbation, diverging from the observed

ramp-like saturation effect. Tsay and colleagues recently synthesized existing evidence

to propose that implicit adaptation reaches an upper bound set by cerebellar error

correction mechanisms, reflected in a ramp-like influence of vision on proprioception

(Tsay, Kim, Saxena, et al., 2022). While this ramp function (instantiated as

proprioceptive re-alignment mode, PReMo) could explain the observed saturation, the

postulate of an upper bound on visual influence lacks empirical validation. Some

research supports the idea of saturation in proprioceptive recalibration (Modchalingam

et al., 2019), yet other studies suggest a linear increase with visual perturbations (Rossi

et al., 2021; Salomonczyk et al., 2011). Additionally, current models fall short of

quantitatively capturing the time-dependent shifts in proprioceptive bias associated with

implicit adaptation.

In this study, we put forth a unified model to account for the distinct features of implicit

adaptation based on the Bayesian combination of movement-related cues. Prior models

have overlooked the fact that visual uncertainty related to the perturbation increases

with the size of the perturbation as the cursor moves further from the point of fixation

and into the visual periphery (Klein & Levi, 1987; Levi et al., 1987). This is particularly

pertinent for implicit adaptation that is widely investigated by the so-called error-clamp

paradigm, in which participants are instructed to fixate on the target and disregard the

perturbing cursor. Moreover, conventional theories of motor adaptation define motor

error according to the sensory modality of the perturbation, i.e., visual errors for visual

perturbations (Tsay, Kim, Saxena, et al., 2022; Wei & Körding, 2009). We propose an

alternative: perceptual error drives implicit adaptation, as the sensory perturbation

influences the perception of the effector's position and, subsequently, motor adaptation.

Through a series of experiments, we aim to demonstrate that combining

eccentricity-induced visual uncertainty (Experiment 1) with a traditional motor adaptation

model (state-space model) and a classical perception model (Bayesian cue

combination) can explain both over-compensation and saturation effects (Experiment

2), as well as the time-dependent changes in hand localization(Tsay et al., 2020), 2020).

Finally, to provide causal evidence supporting our Perceptual Error Adaptation (PEA)

model, we manipulated visual uncertainty and observed that subsequent adaptation

was attenuated for large perturbations but not for small ones—a finding that contradicts

existing models but aligns well with the PEA model. Across the board, our model

outperforms those based on ramp error correction (Tsay, Kim, Saxena, et al., 2022) and

causal inference of errors (Wei & Körding, 2009), offering a more parsimonious

explanation for the salient features of implicit adaptation.

Results

The perceptual error adaptation model with varying visual uncertainty

We start by acknowledging that the perceptual estimation of effector position is

dynamically updated and influenced by sensory perturbations during motor adaptation.

For implicit adaptation studied via the error-clamp paradigm, participants are required to

bring their hand to the target while ignoring the direction-clamped cursor (Morehead et

al., 2017). Accordingly, the perceptual estimation of the hand movement direction relies

on three noisy sensory cues: the visual cue from the cursor, the proprioceptive cue from

the hand, and the sensory prediction of the reaching action (Figure 1A). Without loss of

generality, we posit that each cue is governed by an independent Gaussian distribution:

the visual cue x_v follows $N(\theta, \sigma_v^2)$, where θ is the cursor direction and σ_v^2 is visual

variance, the proprioceptive cue x_p follows $N(x_{hand}, \sigma_p^2)$, where x_{hand} is the hand

movement direction and σ_p^2 is proprioceptive variance, and the sensory prediction cue x_u

follows $N(T, \sigma_u^2)$, where T is the target direction and σ_u^2 is prediction variance.

Participants aim for the target, expecting their hand to reach it. Using the Bayesian cue

combination framework (Berniker & Kording, 2011), the perceived hand location $(\hat{x}_{Hand})$

on trial *n* can be derived:

127
$$\hat{x}_{Hand,n} = \sum_{i} W_i x_{i,n}, \quad with \quad W_i = \frac{1/\sigma_i^2}{\sum_{j} 1/\sigma_j^2}, \quad i,j = u,p,v \tag{1}$$

This estimated hand position is derived using maximum likelihood estimation from the

three noisy cues. Given that the clamped cursor deviates the target by θ, the visual cue

x_v biases the hand estimate $\hat{x}_{Hand}$ towards the cursor's direction. This deviation from the

target direction T constitutes the perceptual error, which drives adaptation on the

subsequent trial $n+1$ (Eq. 2). Consisting with existing models (Albert et al., 2022; Cheng

& Sabes, 2006; Herzfeld et al., 2014; McDougle et al., 2015), trial-to-trial adaptation is

modeled using a state-space equation:

135
$$x_{p,n+1} = Ax_{p,n} + B\left(T - \hat{x}_{Hand,n}\right), \tag{2}$$

136 where A is the retention rate capturing inter-movement forgetting and B is the learning

137 rate capturing the proportion of error corrected within a trial. The interplay between

138 forgetting and learning dictates the overall learning extent, i.e., the asymptote of x_p:

139
$$x_p^{asym} = -\frac{B/\sigma_v^2}{B/\sigma_p^2 + (1-A)\sum_{j} 1/\sigma_j^2}\theta, \quad j = v,p,u \tag{3}$$

140 Thus, the positive influence of perturbation size θ on the adaptation extent is

141 counterbalanced by the rise in visual uncertainty σ_v, since sensory uncertainty of

142 various visual stimuli increases linearly with eccentricity (Klein & Levi, 1987; Levi et al.,

143 1987). As participants are instructed to fixate on the target, an increase in θ leads to

144 increased eccentricity. Hence, we model this linear increase in visual uncertainty by

145
$$\sigma_v = a + b\theta, \tag{4}$$

where *a* and *b* are free parameters. We conducted simulations of implicit adaptation

with varying error clamp size (θ). The model simulation closely resembles the saturated

adaptation in three independent experiments (Kim et al., 2018; Morehead et al., 2017).

In fact, our PEA model predicts a concave adaptation pattern, contrasting with the ramp

pattern suggested by the PReMo model (Tsay, Kim, Saxena, et al., 2022). In

Experiment 1, we aim to validate the assumption of a linear increase in visual

uncertainty (Eq. 1); in Experiment 2, we seek to verify whether implicit adaptation

adheres to a concave pattern as prescribed by the PEA model. Subsequent

experiments, namely Experiments 3 and 4, will test the model's additional novel

predictions concerning changes in proprioception and the impact of experimentally

manipulated visual uncertainty on adaptation, respectively.

158   ---- insert Figure 1 here ----

Experiment 1: Visual Uncertainty Increases with Perturbation Size

To quantify visual uncertainty in a standard error-clamp adaptation setting, we employed

psychometric methods. Occluded from seeing their actual hand, participants (n=18)

made repetitive reaches to a target presented 10 cm straight head while an

error-clamped cursor moving concurrently with one of three perturbation sizes (i.e., 4°,

16°, and 64°), randomized trial-by-trial. In alignment with the error-clamp paradigm,

participants were instructed to fixate on the target and ignore the rotated cursor

feedback. Eye-tracking confirmed compliance with these instructions (Figure S1).

Perturbation directions were counterbalanced across trials, with equal probability of

clockwise (CW) and counterclockwise (CCW) rotation. Post-movement, participants

were required to judge the cursor's rotation direction (CW or CCW) relative to a briefly

displayed reference point (Figure 2A & Figure 6A). Employing this two-alternative

forced-choice (2AFC) task and the Parameter Estimation by Sequential Testing (PEST)

procedure (Lieberman & Pentland, 1982), we derived psychometric functions for visual

discrimination of cursor movement direction (Figures 6 and Figure S2). Our findings

reveal a significant increase in visual uncertainty (σ_v) with perturbation size for both CW

and CCW rotations (Friedman test, CW direction: $\chi^2(2) = 34.11$, p = 4e-8; CCW: $\chi^2(2) =$

26.47, p = 2e-6). Given the symmetry for the two directions, we collapsed data from

both directions, and confirmed the linear relationship between σ_v and θ by a generalized

linear model: $\sigma_v = a + b\theta$, with $a = 1.853$ and $b = 0.309$, $R^2 = 0.255$ (F = 51.6, p =

2.53e-9). The 95% confidence intervals (CI) for a and b are [0.440, 3.266] and [0.182,

0.435], respectively. The intercept was similar to the visual uncertainty estimated in a

previous study (Tsay, Avraham, et al., 2021). We thus observed a striking seven-fold

increase in visual uncertainty from a 4° perturbation to a 64° perturbation (22.641 ±

6.024° vs. 3.172 ± 0.453°). In addition, we observed significant correlations of σ_v

between different perturbation sizes (Spearman correlation, 4° and 16°: $\rho = 0.795$,

p<0.001; 16° and 64°: $\rho = 0.527$, p = 0.026). This finding further confirmed that the

relative magnitude of visual uncertainty among individuals was consistent across

perturbation sizes.

---- insert Figure 2 here ----

Experiment 2: Visual Uncertainty Modulated Perceptual Error Accounts for Overcompensation and Saturation in Implicit Adaptation

The critical test of the PEA model lies in its ability to employ the increase in visual uncertainty obtained from Experiment 1 to precisely explain key features of implicit adaptation. Earlier research mostly scrutinized smaller perturbation angles when reporting saturation effects (Bond & Taylor, 2015; Kim et al., 2018). In contrast, Experiment 2 involved seven participant groups (n = 84) in characterizing implicit adaptation across an extensive range of perturbation sizes (i.e., 2°, 4°, 8°, 16°, 32°, 64°, and 95°). After 30 baseline training cycles without perturbations, each group underwent 80 cycles of error-clamped reaching and 10 washout cycles without visual feedback (Figure 3A). We replicated key features of implicit adaptation: it incrementally reached a plateau and then declined during washout. Small perturbations led to overcompensation beyond visual errors: for 2°, 4°, 8°, and 16° clamp sizes, the adaptation was substantially larger than the perturbation itself. Across perturbation sizes, the faster the early adaptation, the larger the final adaptation level (Figure S4). We noticed that our adaptation phase might not be long enough for small clamp-size conditions to fully adapt (e.g., 2°, 4°, Figure 3A). Nevertheless, we operationally used the adaptation level achieved at the end of the adaptation phase as the measure of adaptation extent. Critically, the adaptation extent displayed a concave pattern: increasing steeply for smaller perturbations and tapering off for larger ones (Figure 3B). A one-way ANOVA revealed a significant group difference in adaptation extent ($F(6,83) = 12.108$, $p =$

1.543e-09). Planned contrasts indicated that 8°, 16°, and 32° perturbations did not differ

from each other (all p>0.417, with Tukey-Kramer correction), consistent with earlier

evidence of invariant implicit adaptation (Kim et al., 2018). However, 64° and 95°

perturbations led to significantly reduced adaptation extents compared to 8° (p =

3.194e-05 and 5.509e-06, respectively), supporting the concave pattern as a more

accurate portrayal of implicit adaptation across varying perturbation sizes.

Importantly, the PEA model, when augmented with visual uncertainty data from

Experiment 1, precisely predicts this size-dependent adaptation behavior (Figure 3B).

Beyond adaptation extent, the model also accurately predicts the trial-by-trial adaptation

across all seven participant groups, employing a single parameter set (R^2 = 0.975;

Figure 3A). The model had only four free parameters (A = 0.974, B = 0.208, σ_p =

11.119°, σ_u = 5.048°; Table S1). Remarkably, both the retention rate A and learning rate

B are consistent with previous studies focusing on visuomotor rotation adaptation

(Albert et al., 2022). We also quantified proprioceptive uncertainty (σ_p) in a subset of

participants (n=13) using a similar 2AFC procedure as in Experiment 1. We found that

σ_p was 9.737°±5.598° (Figure S6), which did not statistically differ from the σ_p value

obtained from the model fitting (two-tailed t-test, p = 0.391). To further verify the model,

a separate data fitting without assuming a linear function in the visual uncertainty was

conducted on the data of Experiment 2 (see details in Supplementary Text 2 and Figure

S7). The estimated visual uncertainty has a strong linear relationship with perturbation

size (R = 0.991, p<0.001). In addition, the slope of model-fitted visual uncertainty is very

close to the values we obtained in Experiment 1. In summary, the perceptual

parameters obtained in Experiment 1, when incorporated into the PEA model,

effectively explain the implicit adaptation behaviors observed in different participant

groups in Experiment 2.

239   ---- insert Figure 3 here ----

In comparative analysis, the PReMo model yields a substantially lower R^2 value of

0.749 (Figure S3B). It tends to underestimate adaptation for medium-size perturbations

and overestimate it for large ones (Figure 3C; see also Figure S3B for trial-by-trial

fitting). Another alternative is the causal inference model, previously shown to account

for nonlinearity in motor learning (Mikulasch et al., 2022; Wei & Körding, 2009).

Although this model has been suggested for implicit adaptation (Tsay, Avraham, et al.,

2021), it fails to reproduce the observed concave adaptation pattern (Figures S3C and

3D). The model aligns well with adaptations to medium-size perturbations (8°, 16°, and

32°) but falls short for small and large ones, yielding an R^2 value of 0.711 (see Figure

S3C for trial-by-trial fits). Model comparison metrics strongly favor the PEA model over

both the PReMo and causal inference models, as evidenced by AIC scores of 2255,

3543, and 3283 for the PEA, PReMo, and causal inference models, respectively (Table

S2). In summary, it is the eccentricity-induced visual uncertainty that most accurately

accounts for the implicit adaptation profile across a broad spectrum of perturbation

sizes, rather than saturated visual influence or causal inference of error.

Experiment 3: Cue Combination Accounts for Changes in Proprioception Measures During Implicit Adaptation

Motor adaptation not only recalibrates the motor system but also alters proprioception (Rossi et al., 2021) and even vision (Simani et al., 2007). In traditional motor adaptation involving both explicit and implicit components, the perceived hand location is initially biased towards the visual perturbation and subsequently stabilizes (Ruttle et al., 2016). However, in implicit adaptation, the perceived hand location initially aligns with but later drifts away from the visual feedback (Tsay et al., 2020). This perceived hand location was verbally reported after an active movement under the influence of the error clamp, with the hand staying near the target at the end of the reach. Thus, it is not the so-called proprioceptive recalibration that is typically probed using a passively located hand (Cressman & Henriques, 2009; Mostafa et al., 2019; 't Hart & Henriques, 2016; Tsay, Avraham, et al., 2021). Nevertheless, this intriguing gradual drift of active localization of the hand is informative of the underlying mechanism of implicit adaptation. The PReMo model proposes that this drift comprises two phases: initial proprioceptive recalibration and subsequent visual recalibration (Tsay, Kim, Saxena, et al., 2022). However, this assumption lacks empirical validation, and how visual recalibration relates to a proprioceptive measurement without visual cues is not laid out either. In contrast, we suggest that the perceived hand location is based on the same Bayesian cue combination principle we laid out in the PEA model. In this particular setting, the perceived hand location at the end of each reach is determined by the proprioceptive cue (x_p) and the estimated hand position under the influence of clamped feedback ($\hat{x}_{Hand}$, Eq. 1), the two relevant cues at the time of reporting relative to the target position.

During early adaptation, $\hat{x}_{Hand}$ is biased towards the clamped feedback (small negative

value, Figure 4A), while x_p remains near the target as the motor system has yet to adapt

(close to 0, Figure 4A). This results in an initial negative proprioceptive bias. As

adaptation progresses, although $\hat{x}_{Hand}$ remains biased with a small negative value, x_p

gradually shifts in the positive direction due to adaptation (value changed from near

zero to increasingly positive), resulting in an increasingly positive perceived hand

location. Remarkably, the PEA model can predict these temporal changes in the

localization of an actively located hand with high accuracy (R^2 = 0.982; Figure 4A).

We postulate that the same Bayesian cue combination governs not only the active

localization of the hand in Figure 4A but also the proprioceptive recalibration that is

probed by the passive localization of the hand. With the biased $\hat{x}_{Hand}$ during adaptation,

our PEA model can make specific quantitative predictions about the relationship

between proprioception changes, measured by a passively located hand, and visual

perturbation size. While traditional visuomotor paradigms suggest either invariant

(Modchalingam et al., 2019) or linear increases in proprioceptive recalibration with an

increase in visual-proprioceptive discrepancy (Salomonczyk et al., 2011), the PEA

model prescribes a concave function in relation to visual perturbation size simply due to

the fact that $\hat{x}_{Hand}$ follows a concave function in relation to the perturbation size (Figure

4B).

To empirically test this prediction, Experiment 3 (n=11) measured participants'

proprioceptive recalibration during implicit adaptation using a procedure similar to the

error-clamp perturbations in Experiment 2. After each block of six adaptation trials,

participants' right hands were passively moved by a robotic manipulandum, and they

indicated the perceived direction of their right hand using a visually represented "dial"

controlled by their left hand (Figure 7B). This method quantifies proprioceptive

recalibration during adaptation (Cressman & Henriques, 2009). Each adaptation block

was followed by three such proprioception test trials. The alternating design between

adaptation and proprioception test blocks allowed us to assess proprioceptive biases

across varying perturbation sizes, which consisted of ±10°, ±20°, ±40°, and ±80°, to

cover a wide range (Figure 4D).

Our findings confirmed a typical proprioceptive recalibration effect, as the perceived

hand direction was biased towards the visual perturbation (Figure 4E). Importantly, the

bias in the initial proprioception test trial exhibited a concave function of perturbation

size. A one-way repeated-measures ANOVA revealed a significant effect of perturbation

size ($F_{(3,30)}=3.603, p=0.036$), with the 20° and 40° conditions displaying significantly

greater proprioceptive bias compared to the 80° condition (pairwise comparisons: 20°

v.s. 80°, p = 0.034; 40° v.s. 80°, p = 0.003). The bias was significantly negative for 20°

and 40° conditions (p = 0.005 and p = 0.007, respectively, with one-tailed t-test) but not

for 10° and 80° conditions (p = 0.083 and p = 0.742, respectively). The concave pattern

aligns well with the PEA model's predictions (Figure 4B), further consolidating its

explanatory power.

---- insert Figure 4 here ----

This stands in contrast to the PReMo model, which assumes a saturation for the

influence of the visual cue on the hand estimate (Eq. 12-13). As a result, PReMo's

predicted proprioceptive bias follows a ramp function, deviating substantially from our

empirical findings (Figure 4C). The causal inference model, which mainly focuses on the

role of visual feedback in error correction, cannot directly predict changes in

proprioceptive recalibration.

Interestingly, we observed that the proprioceptive bias was reduced to insignificance by

the third trial in each proprioception test block (one-tailed t-test, all $p > 0.18$; Figure 4E,

yellow line). The observed proprioceptive bias is formally modeled as a result of the

biasing effect of the perceived hand estimate $\hat{x}_{Hand}$. In our mini-block of passive

localization, the participants neither actively moved nor received any cursor

perturbations for three trials in a row. Thus, the fact that the measured proprioceptive

bias is reduced to nearly zero at the third trial suggests that the effect of perceived hand

estimate $\hat{x}_{Hand}$ decays rather rapidly.

Experiment 4: Differential Impact of Upregulated Visual Uncertainty on Implicit

Adaptation Across Perturbation Sizes

Thus far, we have presented both empirical and computational evidence underscoring

the pivotal role of perceptual error and visual uncertainty in implicit adaptation. It is

crucial to note, however, that this evidence is arguably correlational, arising from natural

variations in visual uncertainty as a function of perturbation size. To transition from

correlation to causation, Experiment 4 (n = 19) sought to directly manipulate visual

uncertainty by blurring the cursor, thereby offering causal support for the role of

multimodal perceptual error in implicit adaptation.

By increasing visual uncertainty via cursor blurring, we hypothesized a corresponding

decrease in adaptation across all perturbation sizes. Notably, the PEA model predicts a

size-dependent attenuation in adaptation: the reduction is less marked for smaller

perturbations and more pronounced for larger ones (Figure 5A). This prediction

diverges significantly from those of competing models. The PReMo model, operating

under the assumption of a saturation effect for large visual perturbations, predicts that

cursor blurring will only influence adaptation to smaller perturbations, leaving adaptation

to larger perturbations unaffected (Figure 5B). The causal inference model makes an

even more nuanced prediction: it anticipates that the blurring will lead to a substantial

reduction in adaptation for small perturbations, a diminishing effect for medium

perturbations, and a potential reversal for large perturbations (Figure 5C). This

prediction results from the model's core concept that causal attribution of the cursor to

self-action—which directly dictates the magnitude of adaptation—decreases for small

perturbations but increases for large ones when overall visual uncertainty is elevated.

363   ---- insert Figure 5 here ----

Starting from the above predictions, Experiment 4 was designed to assess the impact of

elevated visual uncertainty across small (4°), medium (16°), and large (64°) perturbation

sizes. Visual uncertainty was augmented by superimposing a Gaussian blurring mask

on the cursor (Burge et al., 2008). Each participant performed reaching tasks with either

a standard or blurred clamped cursor for a single trial, bracketed by two null trials

devoid of cursor feedback (Figure 5D). These three-trial mini-blocks permitted the

quantification of one-trial learning as the directional difference of movements between

the two null trials. To preclude the cumulative effect of adaptation, perturbation sizes

and directions were randomized across mini-blocks.

Crucially, our findings corroborated the predictions of the PEA model: visual uncertainty

significantly diminished adaptation for medium and large perturbations (16° and 64°),

while leaving adaptation for small perturbations (4°) largely unaffected (Figure 5E). A

two-way repeated-measures ANOVA, with two levels of uncertainty and three levels of

perturbation size, revealed a significant main effect of increased visual uncertainty in

reducing implicit adaptation ($F(1,18) = 42.255$, $p = 4.112e-06$). Furthermore, this effect

interacted with perturbation size ($F(2,36) = 5.391$, $p = 0.012$). Post-hoc analyses

demonstrated that elevated visual uncertainty significantly attenuated adaptation for

large perturbations ($p = 2.877e-04$, $d = 0.804$ for 16°; $p = 1.810e-05$, $d = 1.442$ for 64°)

but exerted no such effect on small perturbations ($p = 0.108$, $d = 0.500$). These

empirical outcomes are not congruent with the predictions of either the PReMo or the

causal inference models (Figure 5B and 5C). This lends compelling empirical support to

the primacy of perceptual error in driving implicit adaptation, as posited by our PEA

model.

Discussion

In this study, we elucidate the central role of perceptual error, derived from multimodal sensorimotor cue integration, in governing implicit motor adaptation. Utilizing the classical error-clamp paradigm, we uncover that the overcompensation observed in response to small perturbations arises from a sustained perceptual error related to hand localization, and the saturation effect commonly reported in implicit adaptation is not an intrinsic characteristic of cerebellum-based learning but is attributable to increasing sensory uncertainty with increasing visual perturbation eccentricity—a factor hitherto neglected in existing models of sensorimotor adaptation. Contrary to conventional theories that describe implicit adaptation as either saturated or invariant (Kim et al., 2018; Tsay, Kim, Saxena, et al., 2022), our data reveal a concave dependency of implicit adaptation on visual perturbation size, characterized by diminishing adaptation in response to larger perturbations. Notably, our Perceptual Error Adaptation (PEA) model, calibrated using perceptual parameters from one set of participants, provides a robust account of implicit adaptation in separate groups subjected to varying perturbations. The model further successfully captures the perceptual consequences of implicit adaptation, including the continuous shifts in hand localization after active movements during adaptation process (Tsay et al., 2020) and the proprioceptive recalibration probed by localization of the passively moved hand. Lastly, we manipulated visual uncertainty independently of perturbation size and demonstrated that upregulating visual uncertainty selectively attenuated adaptation in the context of larger perturbations while leaving smaller perturbations unaffected. These empirical results, inconsistent with predictions from existing models, underscore the conceptual

and quantitative superiority of our PEA model. Our study advocates for a revised

understanding of implicit motor adaptation; that is, it is driven not by sensory prediction

errors but by perceptual errors in localizing one's effector, conforming to Bayesian cue

combination.

The importance of perceptual error was first implied in the PReMo model (Tsay, Kim,

Haith, et al., 2022), which also proposed that a misperceived hand location is the driving

signal for implicit adaptation. However, this previous work improperly called this

perceptual error a proprioceptive error, leading to the impression that it is a type of

sensory error (but see their new unpublished Tsay et al., 2024). The two models differ

fundamentally in their conceptualization of how different cues contribute to the error

signal. The PReMo model posits two intermediate perceptual variables with Bayesian

cue integration: a visual estimate of the cursor and a proprioceptive estimate of the

hand (Tsay, Kim, Saxena, et al., 2022). The final error signal for adaptation is presumed

to be a proprioceptive error, not from further Bayesian cue combination, but from a

visual-to-proprioceptive bias that is governed by a predetermined, ramp-like visual

influence that saturates around a 6–7° visual-proprioceptive discrepancy (Eq. 13).

These assumptions lack empirical validation. In fact, our findings in Experiment 3

indicate that proprioceptive recalibration follows a concave function with respect to

visual perturbation size, contradicting the ramp-like function assumed by PReMo.

Moreover, the presupposed ramp-like visual influence generates a rigid prediction for a

ramp-like adaptation extent profile, which is at odds with the concave adaptation pattern

observed in Experiment 2 and in a similar study involving trial-by-trial learning (Tsay,

Avraham, et al., 2021). Furthermore, the PReMo model predicts that upregulating visual

uncertainty will selectively reduce adaptation to small perturbations while sparing large

ones. This is inconsistent with our findings in Experiment 4, which demonstrated that

upregulating visual uncertainty substantially impacted adaptation more to larger

perturbations than small ones. Lastly, though PReMo has the potential to explain the

temporal shifts in perceived hand location during adaptation, the authors resorted to

separate mechanisms of proprioceptive and visual recalibration at different phases of

adaptation to explain these shifts (Tsay et al., 2020). In summary, the PReMo model's

assumptions introduce limitations that make it less consistent with empirical

observations, particularly concerning the nonlinearities observed in both motoric and

perceptual aspects of implicit adaptation.

The causal inference framework (Wei & Körding, 2009) fails to predict sensorimotor

changes in implicit adaptation adequately. For instance, it underestimates the

adaptation extent for large perturbations and incorrectly predicts that increasing visual

uncertainty would augment, rather than reduce, adaptation to large perturbations. The

casual inference model is also based on Bayesian principle, then why does it fail to

account for the implicit adaptation? We postulate that the failure of the causal inference

model is due to its neglect of visual uncertainty as a function of perturbation size, as we

revealed in Experiment 1. In fact, previous studies that advocating the Bayesian

principle in motor adaptation have largely focused on experimentally manipulating

sensory cue uncertainty to observe its effects on adaptation (Burge et al., 2008; He et

al., 2016; Körding & Wolpert, 2004; Wei & Körding, 2010), similar to our Experiment 4.

Our findings suggest that causal inference of perturbation alone, without incorporating

visual uncertainty, cannot fully account for the diverse findings in implicit adaptation.

The increase in visual uncertainty by perturbation size is substantial: our Experiment 1

yielded an approximate seven-fold increase from a 4° perturbation to a 64° perturbation.

We have attributed this to the fact that people fixate in the desired movement direction

during movements. Interestingly, even for conventional visuomotor rotation paradigm

where people are required to "control" the perturbed cursor, their fixation is also on the

desired direction, not on the cursor itself (de Brouwer, Albaghdadi, et al., 2018; de

Brouwer, Gallivan, et al., 2018). Thus, we postulate that a similar hike in visual

uncertainty in other "free-viewing" perturbation paradigms. Future studies are warranted

to extend our PEA model to account for implicit adaptation in other perturbation

paradigms.

Our research contributes to an ongoing debate concerning the driving forces behind

error-based motor learning, specifically addressing the question of whether implicit

adaptation is driven by target error or sensory prediction error (Albert et al., 2022; Izawa

& Shadmehr, 2011; Leow et al., 2020; Mazzoni & Krakauer, 2006; McDougle et al.,

2015; Miyamoto et al., 2020; Taylor & Ivry, 2011; Tseng et al., 2007). Most empirical

data fueling this debate stem from traditional motor adaptation paradigms where explicit

and implicit learning co-occur and interact. In these paradigms—visuomotor rotation

being a prime example—target error (TE) is defined as the disparity between the target

and the perturbed cursor, while sensory prediction error (SPE) is the disparity between

the predicted and actual cursor. These two types of error, sensory (specifically, visual) in

nature, differ simply due to the presence of explicit learning: the predicted (desired and

aiming) direction deviates from the target direction when explicit learning is

present(Taylor et al., 2014). Our study suggests that neither target error nor sensory

(visual) prediction error drives implicit adaptation in the error-clamp paradigm. The

error-clamp paradigm isolates implicit learning and thus eliminates potential confounds

from explicit learning. In this paradigm, the predicted and target directions are aligned,

and the target error and sensory prediction error effectively refer to the same visual

discrepancy. However, eitherTE or SPE, if plugged in the classical state-space models,

is able to account for the nuanced features of implicit adaptation (Tsay, Kim, Saxena, et

al., 2022). In contrast, our PEA model reframes the perturbing cursor as a visual cue

influencing the perceptual estimation of hand location rather than as a source of visual

error. The resultant bias in hand estimation from the desired target serves as the actual

error signal. This leads us to posit that the error signal driving implicit sensorimotor

adaptation is fundamentally perceptual rather than sensory. From a normative

standpoint, this perceptual error could be construed either as a predictive or

performance error (Albert et al., 2022), but importantly, it is not tied to a specific modality

(i.e., vision or proprioception). Instead, it directly pertains to the perceptual estimate that

is crucial for task execution, i.e., bringing the hand in the desired direction.

The concept of perceptual error-driven learning can be extrapolated to various motor

learning paradigms, including other motor adaptation tasks. For instance, in visuomotor

rotation tasks, explicit learning manifests as a deviation in the aiming direction from the

visual target, whereas implicit learning manifests as a further deviation the actual hand

position from this aiming direction (Taylor et al., 2014). With the presence of re-aiming,

the perturbed cursor still deviates from the re-aiming direction and thus produces the

perceptual bias of the hand from the desired direction, which subsequently drives

implicit adaptation. In this scenario, the perceptual error is defined as the difference

between the perceptual estimate of the hand and the "re-aiming" direction, which serves

as the new "target" when explicit learning is in play. Our PEA model would predict

similar saturation effects in implicit adaptation for this conventional adaptation paradigm,

comparable to for the error-clamp paradigm. Indeed, existing findings from the

conventional adaptation support this prediction: the implicit adaptation follows either a

saturation effect (Bond & Taylor, 2015; Neville & Cressman, 2018) or a concave pattern

(Tsay, Haith, et al., 2022) across a range of perturbation sizes. Furthermore, according

to the PEA framework, this perceptual error is anchored on the aiming target, thereby

naturally predicting that implicit and explicit adaptations should interact in a

complementary manner, a notion that aligns with recent theories on their interaction

(Albert et al., 2022; Miyamoto et al., 2020).

Besides visuomotor rotation, all the primary motor adaptation paradigms, including

prism adaptation (e.g., Petitet et al., 2018; von Helmholtz, 1867), visuomotor gain

adaptation (e.g., Pearson et al., 2010), and force field adaptation (e.g., Shadmehr &

Mussa-Ivaldi, 1994), involves locating one's effector under the influence of perturbed

sensory feedback. Thus, we postulate that perceptual error elicited by sensory

perturbation might play a similar significant role in driving implicit learning in these

diverse task paradigms. It is noteworthy that error-based learning also plays a major

role in motor skill learning, including de novo skill learning and motor acuity learning

(Krakauer et al., 2019). Unlike motor adaptation, motor skill learning currently lacks

computational models to account for its error-based learning component, owing to the

complexity and diversity of skill learning tasks. We believe that Bayesian cue

combination and perceptual error about locating one's effect might serve as a starting

point for theorization of the implicit processes in skill learning.

Our study also provides a new perspective on explaining proprioceptive changes during

motor adaptation, advocating for a Bayesian cue combination framework. Previously,

the change in proprioceptive hand localization during motor adaptation has been

ascribed to visual-proprioceptive discrepancy-induced recalibration (Ruttle et al., 2018;

Salomonczyk et al., 2013) and/or altered sensory prediction driven by the adapted

forward internal model (Mostafa et al., 2019; 't Hart & Henriques, 2016). To dissect

these components, researchers have often compared proprioceptive localization in

actively moved (Ruttle et al., 2021; 't Hart & Henriques, 2016) versus passively placed

(passive localization, e.g., Experiment 3) hands during adaptation, attributing the

smaller bias in passive localization to recalibration alone. The difference between the

two is then considered to reflect altered sensory prediction due to motor adaptation

(Mostafa et al., 2019; Rossi et al., 2021). But these conceptual divisions lack

computational models for validation. For instance, researchers have shown that

proprioceptive recalibration in visuomotor adaptation is either a fixed proportion (e.g.,

20%) of the visual-proprioceptive discrepancy (Henriques & Cressman, 2012; Ruttle et

al., 2021) or largely invariant (Modchalingam et al., 2019). However, a computational

model has yet to be proposed elucidate the underlying mechanism for these diverse

findings. In fact, cross-sensory calibration typically follows the Bayesian principle, as

shown in task paradigms other than motor adaptation (Stetson et al., 2006; Wozny &

Shams, 2011). We propose that proprioceptive changes elicited by motor adaptation, no

matter measured by a passive or active localization method, are governed by the same

Bayesian cue combination principle and the same critical cue, i.e., (mis)perceived hand

location ($\hat{x}_{Hand}$). Given that $\hat{x}_{Hand}$ has a concave dependency to visual perturbation size,

we would observe a corresponding concave proprioceptive recalibration profile.

Besides highlighting the importance of Bayesian cue combination, our study also

suggests that proprioceptive measurements should be understood by considering

available cues in the specific task setting. For both localization tests, i.e., the passive

(Experiment 3) and active localization (Tsay et al., 2020; Figure 4A), the hand stayed at

the trial end when the participants reported their perceived hand location. Thus, the

reported hand location is determined by the just-experienced $\hat{x}_{Hand}$ and the actual

proprioceptive cue. In the case of active localization, the proprioceptive cue becomes

increasingly large (biased to the positive direction in reference to the target), driven by

the adaptation process. In this sense, active localization indeed serves as a

multifaceted reflection of both the internal model (the adapted hand, supplying x_p) and

proprioceptive recalibration (the perceived hand after a movement, $\hat{x}_{Hand}$), as proposed

by previous researchers (Mostafa et al., 2019; Rossi et al., 2021). During the initial

stages of perturbation, the immediate negative bias in active localization is

predominantly attributable to rapid proprioceptive recalibration. This is evidenced by a

sudden shift in the estimated hand position ($\hat{x}_{Hand}$; Figure 4A), occurring before the

internal model has had sufficient time to adapt. Then, why does active localization in

traditional motor adaptation paradigms yield a largely stable bias (Ruttle et al., 2016,

2021)? We note that traditional visuomotor rotation paradigms invokes a rapid initial

explicit learning, driving the adaptation to its asymptote quickly. At the same time,

previous investigations have predominantly measured active localization when

adaptation has asymptoted (Henriques & Cressman, 2012; Modchalingam et al., 2019;

Mostafa et al., 2019; Salomonczyk et al., 2011, 2013; Tsay, Kim, et al., 2021).

Consequently, these studies overlook the evolving effect of the adaptation. In contrast,

the gradual nature of implicit adaptation, shown in the error-clamp paradigm here,

provides a unique opportunity to uncover the underlying mechanisms governing

changes in active localization during the adaptation process.

Notably, our model aligns with previous findings that show a positive correlation

between proprioceptive recalibration, measured by the passive localization method, and

motor adaptation based on individual differences (Ruttle et al., 2021; Salomonczyk et

al., 2013; Tsay, Kim, et al., 2021). Unlike existing theories that posit proprioceptive

recalibration either as a component (Modchalingam et al., 2019; Mostafa et al., 2019;

Ruttle et al., 2021) or a driver for implicit adaptation (Tsay, Kim, Saxena, et al., 2022),

our PEA model provides a mechanistic and empirically testable framework. It posits that

the misestimation of hand position ($\hat{x}_{Hand}$) —induced by the recent perturbation—serves

as the driving factor for both implicit adaptation and changes in proprioception. In other

words, this misestimation is the common cause for both implicit adaptation and

proprioceptive recalibration. This is why the concave dependency of $\hat{x}_{Hand}$ on

perturbation leads to similar concave perturbation dependencies in both implicit

adaptation and proprioceptive recalibration. Updated on a trial-by-trial basis, this

misestimation exerts immediate effects, manifesting as an abrupt negative bias (Figure

4A). Additionally, its influence decays rapidly, becoming negligible within three trials

(Figure 6C). These converging lines of evidence strongly suggest that perceptual

misestimation of hand position is central to the process of proprioceptive recalibration

during adaptation.

Our findings contribute nuanced perspectives to the modulation of implicit learning rate

by factors beyond visual perturbation size. Previous studies have shown that

environmental inconsistency -- defined as the inconsistency of visual errors -- reduced

the rate (Herzfeld et al., 2014; Hutter & Taylor, 2018) or extent (Albert et al., 2021) of

implicit adaptation. Baseline motor variance in unperturbed conditions has been shown

to increase implicit adaptation rate, proposed as a sign of better exploratory learning

(Wu et al., 2014). These studies interpret such phenomena as parametric changes in

the learning rate in relation to visual errors, conceptualized as alterations to the *B*

parameter in existing models. However, an apparent change in learning rate due to

visual errors does not necessarily signify parametric modification, but may attribute to

other factors that influence the use of visual cues (He et al., 2016), such as visual

uncertainty in our case. Previous research has also pointed to various alternative

factors for determining apparent learning rate, including error discounting based on

causal inference of error (Wei & Körding, 2009), proprioceptive uncertainty (Ruttle et al.,

2021; Tsay, Kim, et al., 2021), and state estimation uncertainty (He et al., 2016). In line

with these previous studies, our work suggests a shift in perspective: the driving error

signal for implicit learning should be considered as perceptual, rather than merely

visual. Hence, all the aforementioned factors can be considered as contributing to the

perceptual estimate. This paradigmatic shift could serve as a cornerstone for future

research aimed at understanding how learning rates change under varying conditions.

Our new framework opens avenues for exploring the memory characteristics of implicit

learning. Traditional motor adaptation often exhibits 'savings,' or accelerated relearning

upon re-exposure to a perturbation (Della-Maggiore & McIntosh, 2005; Huberdeau et

al., 2019; Krakauer et al., 2005; Landi et al., 2011). In contrast, implicit adaptation has

been found to exhibit a decreased learning rate during re-adaptation (Avraham et al.,

2021), a phenomenon attributed to conditioning (Avraham et al., 2021) or associative

learning mechanisms (Avraham et al., 2022). Investigating this 'anti-saving' effect will

yield insights into the unique memory properties of implicit learning. Although our

current PEA model is structured around single-epoch learning and does not directly

address this question, it does raise new, testable hypotheses. For example, is the

reduced adaptation rate during relearning attributable to a down-weighting of perturbed

visual feedback in cue combination, or does it reflect a parametric alteration in the

learning rate? Another noteworthy aspect of implicit learning is its remarkably slow

decay rate. It has been observed that the number of trials required to wash out the

implicit adaptation exceeds the number of trials needed to establish it (Avraham et al.,

2021; Tsay et al., 2020). In the context of our perceptual error framework, this raises the

possibility that washout phases might be governed by state updating involving a distinct

set of sensorimotor cues or an alternative updating mechanism, such as memory

formation and selection (Oh & Schweighofer, 2019).

Data availability

Data and codes presented in this work are available at:

https://doi.org/10.6084/m9.figshare.24503926.v2 and

https://github.com/ZhaoranZhang/PEAmodel.git

Acknowledgment

This work was supported by STI2030-Major Projects (2021ZD0202600) and the

National Natural Science Foundation of China (62061136001, 32071047, 31871102),

awarded to KW, and the National Natural Science Foundation of China (32300868)

awarded to ZZ.

649 Methods

650 Participants

We recruited 115 college students from Peking University (77 females, 38 males, 22.05

± 2.82 years, mean ± SD). Participants were all right-handed according to the

Edinburgh handedness inventory (Oldfield, 1971) and had normal or

corrected-to-normal vision. Participants were naïve to the purpose of the experiment

and provided written informed consent, which was approved by the Institutional Review

Board of the School of Psychological and Cognitive Sciences, Peking University.

Participants received monetary compensation upon completion of the experiment.

658 Apparatus

In Experiment 1, 2, and 4, participants were seated in front of a vertically-placed LCD

screen (29.6 x 52.7 cm, Dell, Round Rock, TX, US). They performed the movement task

with their right hand, holding a stylus and slide it on a horizontally placed digitizing tablet

(48.8 x 30.5 cm, Intuos 4 PTK-1240, Wacom, Saitama, Japan). In Experiment 1, a

keyboard was provided to the participants' left hand to enable them to report the

direction of visual stimuli in the discrimination task. A customized wooden shelter was

placed above the tablet to block the peripheral vision of the right arm. In Experiment 1

and 4, participants placed their chin on a chin rest attached to the wooden shelter to

stabilize their heads. Their eye movement was recorded by an eye tracker (Tobii pro

nano, Tobii, Danderyd Municipality, Sweden) affixed at the lower edge of the screen.

The sampling rate was 160-200 Hz for the tablet and 60 Hz for the eye tracker.

Experiment 3 was conducted using the KINARM planar robotic manipulandum with a
virtual-reality system (BKIN Technologies Ltd., Kingston, Canada). Participants, seated
in a chair, held the robot handles with their left and right hands (Figure 7). The
movement task was performed with the right handle and the left handle was used to
indicate the perceived direction of the right hand in the proprioception test. A
semi-silvered mirror was placed below the eye level to block the vision of the hands and
the robotic manipulandum; it also served as a display monitor.

Experiment 1: measuring visual uncertainty in error-clamp adaptation

Eighteen among twenty participants finished the reaching with clamped error feedback
and visual discrimination task in 3 consecutive days; two participants withdrew during
the experiment. Participants made reaching movement by sliding the stylus from a start
position at the center of the workspace to towards a target (Figure 6A). The start
position, the target, and the cursor were represented by a gray dot, a blue cross and a
white dot on the screen, respectively. All these elements had a diameter of 5mm. The
procedure of the motor and visual discrimination task is illustrated in Figure 2A. To
initiate a trial, participants moved the cursor into the start position. Following an 800ms
holding period, a target appeared 10 cm away in the twelve o'clock direction, and
participants were instructed to slide through the target rapidly while maintaining a
straight hand trajectory. The trial terminated when the distance between the hand and
the start position exceeded 10 cm, regardless of whether the target was hit. A warning
message, "too slow", would appear on the screen if participants failed to complete the
trial within 300 ms after initiating the movement. Each practice day began with 60
standard reaching trials, during which veridical feedback about hand location was

provided by the cursor. The target would change from blue to green if the cursor

successfully passed through it. In subsequent visual clamp trials, the cursor moved

along a predetermined direction set by the perturbation angle, while its position was

updated in real-time based on the hand's location. The cursor's distance from the start

position was equal to the distance between the hand and the start position until the end

of the trial.

Following each trial, the cursor remained frozen at its final position for an additional 800

ms before disappearing. The visual discrimination task commenced 1000 ms thereafter.

A yellow reference point, located 10 cm from the start position, was displayed for 150

ms near the cursor's final position (Figure 2A & Figure 6A). Subsequently, all visual

stimuli, except for the blue cross at the start position, were removed from the screen.

Participants were then required to judge whether the reference point was situated in a

clockwise (CW) or counterclockwise (CCW) direction relative to the cursor's final

position and to report their judgment by pressing a key on the keyboard. Participants

were informed that they no longer controlled the direction of cursor movement during

the task. They were instructed to fixate their gaze on either the start position or the blue

cross during the motor task, while actively ignoring the white cursor. During the

discrimination task, they were required to maintain their gaze on the blue cross. Eye

movements were monitored in real-time using an eye tracker. Participants received a

warning if their gaze was detected outside a 75-pixel-wide band-shaped region centered

on the line of gaze four consecutive times during the experiment (Figure S1).

715   ---- insert Figure 6 here ----

In each trial, the angular deviation between the error-clamped cursor and the reference point was determined using a PEST procedure (Lieberman & Pentland, 1982). Figure 6C-D illustrates the evolution of the deviation angle and step size for an exemplary participant experiencing a -16° perturbation. In each round, the deviation commenced at 30° (indicated by yellow points in Figure 6C-D) and was altered by one step size following each trial. The initial step size was set at 10° and was halved whenever the direction judgment changed (i.e., from "CW" to "CCW" or vice versa). For a specific perturbation angle, the initial deviation always started from the CW direction for the first round and flipped the direction at the beginning of the next round. A round terminated either when the step size fell below a predefined criterion (indicated by the red line in Figure 6D) or when the trial count exceeded 30. Six perturbation angles were randomly interleaved (Figure 6B), and the experiment concluded when four complete rounds of the PEST procedure had been completed for each perturbation angle. Consequently, the total number of trials varied among participants and across practice days (300-500 trials per day on average, with a maximum of 720 trials). Additionally, for some perturbation angles, more than four complete rounds could be conducted in a single day.

Experiment 2: Motor adaptation with different perturbation size

Eighty-four participants were randomly allocated into seven groups, each comprising 12 individuals. Each group performed a motor adaptation task featuring clamped visual feedback at different perturbation angles: 2°, 4°, 8°, 16°, 32°, 64°, and 95°. As in Experiment 1, participants were instructed to slide rapidly and directly through the

target, which was represented by a blue dot rather than a cross. In each trial, the target

appeared at one of four possible locations (45°, 135°, 225° or 315° counter-clockwise

from the positive x-axis). The sequence of target locations was randomized yet

constrained so that all four positions appeared in cycles of four trials. Each group

commenced with a baseline session that included 15 cycles of reaching trials with

veridical feedback, followed by 15 cycles without visual feedback. Subsequently, during

the perturbation session, participants completed 80 cycles of training trials featuring the

error-clamped cursor with one perturbation angle (i.e., clamp size), depending on their

group assignment. To assess the aftereffect, a session comprising 10 cycles of

movement without visual feedback was administered. In summary, each participant

practiced a total of 480 trials (120 cycles x 4 target directions) in Experiment 2.

Experiment 3: Proprioception test with different perturbation sizes

Eleven participants were recruited for testing their proprioceptive recalibration. This

experiment incorporated two types of trials: reaching trials and proprioception test trials.

During the reaching trials, participants were instructed to aim for a target, which could

appear at one of three possible locations (25°, 45°, or 65° counter-clockwise from the

positive x-axis, as represented by light blue dots in Figure 4C, right panel). The task

was similar to those in Experiments 1 and 2, with the key difference being that

participants performed the task using KINARM robots (as depicted in Figure 7A). The

dimensions and relative distances of the visual stimuli remained consistent with those

used in Experiments 1 and 2. As in previous experiments, three kinds of visual feedback

were provided during different sessions: no visual feedback, veridical feedback, and

feedback featuring an error-clamped cursor.

In the proprioception test, participants were instructed to hold the robot's right handle

and wait to be passively moved by the robot to one of six proprioception targets (small

red dots in Figure 4C, right panel). These targets were spaced at 10° intervals, ranging

from 20° to 70° counterclockwise from the positive x-axis, and flanked the three

reaching targets. The passive movement lasted for 1,000 ms and followed a

straight-line path at a speed consistent with a minimum jerk velocity profile. During this

movement, a ring with a 10 cm radius, centered at the start position, was displayed on

the screen (depicted as a red arc in Figure 7B). The cursor was also replaced by a ring,

its radius expanding as the hand moved toward the proprioception target.

After the right hand reached the proprioception target, participants were instructed to

maintain their right hand's position. Using the left handle, they were then asked to

indicate the perceived location of their right hand. The position of the left handle was

mapped to the rotation of a "dial," which was constrained to the target arc.

776   ---- insert Figure 7 here ----

The position of h_p was displayed on the target arc as a small red rectangle (a visual

"dial," as shown in Figure 7B). Participants were instructed to indicate the location of

their right hand by moving the red rectangle to the position they perceived as accurate.

The final position of h_p was recorded when its angular velocity remained below 1

degree/second for a duration exceeding 1000 ms. The proprioceptive bias was then

calculated as the angular deviation between the actual hand position (h_R) and the

perceived hand position (h_P).

Reaching trials and proprioception test trials were organized into blocks (Figure 4D).

Each reaching block consisted of 6 trials, targeting 3 different locations with 2

repetitions each. Each reaching block was followed by a proprioception test block

consisting of 3 trials. In these test trials, the robot moved the participant's right hand

toward a target position near one of the three reaching targets. These test targets were

randomly chosen from six possible locations (Figure 4C, right panel). The entire

experiment comprised 40 reaching blocks and 40 subsequent proprioception test

blocks. The first four reaching blocks provided veridical cursor feedback, the next four

offered no cursor feedback, and the remaining 32 featured one of eight possible

perturbation sizes (±10°, ±20°, ±40°, and ±80°). The size of the perturbation was

randomized between blocks.

Experiment 4: upregulating visual uncertainty affects implicit adaptation

Nineteen participants from Experiment 1 completed Experiment 4. The reaching task

employed the same setup as in Experiment 1. However, instead of performing

perceptual judgments of cursor motion direction, participants engaged in movements

with one of three types of cursor feedback: veridical feedback, no feedback, and

feedback with clamped perturbation. To assess the influence of visual uncertainty on

implicit learning, we modified the cursor to appear blurred in half of the clamped trials.

The blurring mask had a diameter of 6.8 mm, and the color intensity decreased from the

cursor's center following a two-dimensional Gaussian distribution with $\sigma_x = \sigma_y = 1.4$

mm. As depicted in Figure 5D, participants underwent the same procedures across

three consecutive days. Each day consisted of 60 baseline trials, followed by 15 training

blocks designed to assess single-trial learning. Within each training block, 12 trials

featured an error-clamped cursor, each flanked by a trial without feedback. The

difference between two adjacent no-feedback trials served as a measure of single-trial

learning at specific perturbation sizes. Each of the 12 perturbation trials was randomly

assigned one of 12 possible perturbations, comprising two cursor presentations (blurred

or clear) and six clamp sizes (±4°, ±16°, ±64°). Each participant practices 180 trials (15

blocks x 12 trials per block) with perturbation per day and 540 trials in total.

Data analysis

Processing of kinematic data

In Experiments 1, 2, and 4, hand kinematic data were collected online at a sampling

rate ranging between 160 and 200 Hz and subsequently resampled offline to 125 Hz.

The movement direction of the hand was determined by the vector connecting the start

position to the hand position at the point where it crossed 50% of the target distance,

i.e., 5 cm from the start position.

In Experiment 3, hand positions and velocities were directly acquired from the KINARM

robot at a fixed sampling rate of 1 kHz. The raw kinematic data were smoothed using a

fifth-order Savitzky-Golay filter with a window length of 50 ms. Owing to the high

temporal resolution and reliable velocity profiles provided by the KINARM system, the

heading direction in Experiment 3 was calculated as the vector connecting the start

position to the hand position at the point of peak velocity. For each participant, the

baseline hand direction and proprioceptive bias was subtracted from the data.

Psychometric curve

For the visual discrimination task, data of all three days were pooled together, the

probability of responding that "the reference point was in the counter-clockwise direction

of the cursor" was calculated as p for all angle differences (Figure S2). At each

perturbation size, a logistic function was used to fit the probability distribution for

individual participants:

834
$$p = \frac{1}{1+e^{-k\left(x-x_0\right)}},$$
(5)

835 where k is the slope, and x_0 is the origin of the logistic function. The visual uncertainty

836 was defined as the angle differences between 25% and 75% of the logistic function:

837
$$\sigma_v = \frac{\log\log\left(p_2/(1-p_2)\right) - \log\log\left(p_1/(1-p_1)\right)}{k},$$
(6)

where $p_1 = 25\%$ and $p_2 = 75\%$.

Statistical analysis

In Experiment 1, since the visual uncertainty σ_v follows a non-negative skewed

distribution among participants, it violated the assumption of the ANOVA test. We thus

applied Friedman's nonparametric test to determine whether σ_v changes with the

perturbation angle θ. Specifically, σ_v for both positive and negative θ were subjected to

Friedman's test separately, with θ serving as the factor. Spearman correlation analyses

were conducted between σ_v in different perturbation sizes. Given the symmetry between

positive and negative θ, we pool the data to quantify the linear dependency of σ_v on the

absolute θ (Eq. 4). Because σ_v is expected to be always positive and there are potential

positive outliers, we assume that it is generated from a gamma distribution rather than a

normal distribution, which is always positive and favors right-skewed distribution that

can well capture the positive outliers. Thus, the data was fitted by a generalized linear

regression model with the absolute value of θ as independent variable and σ_v as a

dependent variable.

In Experiment 2, the adaptation extent was defined as the mean hand angles in the last

10 cycles in the perturbation phase (cycle 101-110). A one-way ANOVA with

perturbation size serving as the factor to examine its influence on the adaptation extent.

Pairwise post-hoc comparisons were conducted using Tukey-Kramer correction.

In Experiment 3, proprioceptive biases were quantified as the angular difference

between the perceived and actual hand directions. A one-way repeated-measures

ANOVA was conducted on the data of the first trial, using perturbation size as the

within-subject factor. Greenhouse-Geisser corrections were applied when the

assumption of sphericity was violated (Kirk, 1968). Multiple pairwise comparisons were

conducted among different perturbation sizes for the first proprioception test. To

determine if the proprioceptive biases were significantly different from zero, one-tailed

(left) *t*-tests were conducted separately for the first and third proprioception test trials at

each perturbation size.

In Experiment 4, the single-trial learning data was subjected to a 2 (visual uncertainty) x

3 (perturbation size) repeated-measures ANOVA. Greenhouse-Geisser corrections were

applied as above, and the simple main effect of visual uncertainty was tested for each of

the three perturbation sizes.

Model fitting and simulations

Perceptual Error Adaptation (PEA) model

Model fitting for adaptation extent as a function of perturbation size. To fit the adaptation

extent data from three different experiments in previous studies in (Kim et al., 2018;

Morehead et al., 2017), Eq. 3 and Eq. 4 were modified for simplification. To avoid

overfitting of the small dataset, we reduced the number of model parameters by

assuming that $\hat{x}_{Hand}$ asymptote to the target direction in the final adaptation trials that

are used for computing adaptation extent, thus the retention rate $A \equiv 1$. Insert Eq. 4 to

Eq. 3, the asymptote hand angle with different perturbation size is:

879
$$x_p^{asym} = - \left(\frac{\sigma_p/a}{1+(b/a)\theta} \right)^2 \theta. \tag{7}$$

Two ratio parameters $R_{1,ext} = \sigma_p/a$ and $R_{2,ext} = b/a$ were used in data fitting. Three

datasets were fitted separately.

Model fitting for trial-by-trial adaptation and proprioception changes. The trial-by-trial

changes of adaptation (Figure 3A) and of proprioceptive changes (Figure 4A) were fitted

with Eq. 1, Eq. 2, and Eq. 4 based on the mean performance of all participants. The

PEA model only had four free parameters, $\Theta = [\sigma_u, \sigma_p, A, B]$. The slope *a* and intercept *b*

in Eq. 1 were obtained by psychometric tests from Experiment 1 (see statistical

analysis). The reported hand position (x_{report}, blue dots in Figure 4A) was based on the

proprioceptive cue x_p and the estimated hand $\hat{x}_{Hand}$ from the reaching trial. With the

Bayesian cue combination assumption, the reported hand position was biased by x_p

with a ratio determined by the variance of x_p and $\hat{x}_{Hand}$:

891
$$x_{report} = \hat{x}_{Hand} + \frac{\sigma^2_{Hand}}{\sigma^2_{Hand}+\sigma^2_p} (x_p - \hat{x}_{Hand}),$$
(8)

where σ^2_{Hand} and σ^2_p are the variance of $\hat{x}_{Hand}$ and x_p respectively (see Supplementary

Text 1 for further details of this fitting).

To verify if the slope b and intercept a obtained from Experiment 1 are consistent across

experiments, they were also estimated by fitting data from Experiment 2 (Figure 3). In

this case, the model fitting was performed with 6 free parameters, $\Theta = [\sigma_u, \sigma_p, a, b, A, B]$.

The fitted values of a and b are fallen into the 95% CI of estimated parameters in

Experiment 1 (purple line in Figure 2C, see details in Table S1).

The dependence of proprioceptive recalibration on perturbation size (Figure 4B) were

simulated by the PEA model with the parameter values estimated from Experiment 2.

We assumed that the proprioceptive bias results from the influence of a biased hand

estimate $(\hat{x}_{Hand})$ during adaptation and the influence is quantified as a percentage of its

deviation from the true hand location:

904
$$x_{bias} =- (0 - \hat{x}_{Hand})R_p,$$
(9)

where the actual hand location is 0, R_p is the percentage of influence, and $\hat{x}_{Hand}$ is

determined by Eq.1. In simulation, R_p varied from 0.05 to 0.8 to estimate the overall

dependence of proprioceptive recalibration on perturbation size.

Model fitting and simulation for single-trial learning. In the single-trial learning paradigm

(Figure S5), the average movement direction across trials aligns with the target direction

since the visual perturbations are evenly distributed in both directions. Thus, the

sensory cues x_u and x_p have the same mean value. For modeling single-trial learning,

instead of having two separate cues, we assume a combined cue of x_u and x_p to follow

$x_{int} \sim N(T, \sigma_{int}^2)$, where T is the target direction, $\sigma_{int}^2 = \frac{\sigma_u^2 \sigma_p^2}{\sigma_u^2 + \sigma_p^2}$ represents the variance of

integrated sensory signal of x_u and x_p. Single-trial learning was quantified as the

difference between the two null trials before and after the perturbation trial. As the

perturbation size in the triplet of trials varied randomly, we assume that the effects of

different perturbations are independent. Thus, single-trial learning was modeled as

learning from the current perturbation without history effect. It follows the equations

modified from Eq. 1 and 2:

920
$$x_{STL} = B(T - \hat{x}_{Hand}) \tag{10}$$

921
$$\hat{x}_{Hand} = W_{int}T + W_v x_v, \text{ with } W_{int} = \frac{1/\sigma_{int}^2}{\sum_j 1/\sigma_j^2}, \quad i, j = int, v, \tag{11}$$

where x_v is the visual perturbation, W_{int} and W_v are the weights of the cues, σ_v is the

standard deviation of the visual cue specified by Eq.4. Parameter set $\Theta = [\sigma_{int}, a, b, B]$

was fitted to the average data from all participants. Model simulations (Figure 5A) were

performed with the same single-trial learning equations. For the clear cursor condition,

we used the same parameter values estimated from Experiment 2 (see details in Table

S1). For the blurred cursor condition, the standard deviation of visual cue was changed

to:

$$\sigma_{v,blur} = R_v \sigma_v \tag{12}$$

930 for the simulation of the increase in visual uncertainty, the ratio R_v varied from 1.1 to 3.

931 **PReMo model**

932 We used the PReMo model to fit the average adaptation extent obtained from

933 Experiment 2 (Figure 3C & Figure S3B). Following the study by (Tsay, Kim, Saxena, et

934 al., 2022), the hand position at trial *n+1* is:

$$x_{p,n+1} = Ax_{p,n} + B\left(T - x_{per,n}\right), \tag{13}$$

936 where

$$x_{per,n} = \beta_p + \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2} x_{p,n}, \tag{14}$$

$$\beta_p = - min\left(\left|\beta_p^{sat}\right|, \left|\eta_p\left(\frac{\sigma_u^2}{\sigma_u^2 + \sigma_v^2} x_{v,n} - \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2} x_{p,n}\right)\right|\right). \tag{15}$$

In data fitting, we used two parameters to represent the ratio between sensory cues:

$R_1 = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_v^2}$ and $R_2 = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2}$. The data were fitted with the parameter set $\Theta = [R_1, R_2, \beta_p^{sat},$

η_p, *A, B*], where β_p^{sat} is the saturation angle, η_p is a scaling factor, *A* is the retention rate

and *B* is the learning rate. For simulating the proprioceptive localization of the hand

(Figure 4C), the parameter values estimated from Experiment 2 were used. The bias of

hand estimation in the proprioception trials is determined as: $x_{bias} = - (0 - x_{per})R_p,$

where ratio R_p varies from 0.05 to 0.8. Thus, similar to the PEA model simulation, the

proprioceptive bias is a fraction of the bias in the hand estimation from the adaptation

trials. Single-trial learning (Figure 5B) was simulated with:

$$x_{STL} = B(T - x_{per}),\qquad(16)$$

where x_{per} is determined by Eq. 12 and Eq. 13. For the clear condition, we used the

parameter values estimated from Experiment 2 with PReMo. For the blurred cursor

condition, the standard deviation of visual signal $\sigma_{v,\,blur}$ increases with a ratio R_v, as in

Eq. 12.

Causal inference model

The causal inference model by (Wei & Körding, 2009) was used to fit the data of

Experiment 2 (Figure 3D & Figure S3C). The hand position at trial n+1 is updated by

learning from visual error at trial n:

$$x_{p,n+1} = Ax_{p,n} + B\big(T - px_{v,n}\big),\qquad(17)$$

958 where *A* and *B* are the retention and learning rates, respectively; *T* is the target

959 direction. Specifically for this model, the learning from error is modulated by the

960 probability (*p*) of causal attribution of visual error to the action or proprioception:

$$p = S\frac{N(x_{v,n},0,\sigma^2)}{N\big(x_{v,n},0,\sigma^2\big)+C},\qquad(18)$$

962 where $x_{v,n}$ is the visual cue at trial n. *S* and *C* are the scaling factors, and σ is the

963 standard deviation of the integrated cue combining visual and proprioceptive cues,

964 following

965
$$\sigma^2 = \frac{\sigma_v^2 \sigma_p^2}{\sigma_v^2 + \sigma_p^2} \quad (19)$$

966 Thus, the data were fitted with five parameters $\Theta = [\sigma, S, C, A, B]$. For simulating

967 single-trial learning with cursor blurring (Figure 5C), the ratio between σ_v and σ_p is fixed

968 as ½. The single-trial learning was determined as:

969
$$x_{STL} = B(T - px_v), \quad (20)$$

where p is determined by Eq. 18. Put Eq. 12 and Eq. 19 into $\sigma_{blur}^2 = \frac{\sigma_{v,blur}^2 \sigma_p^2}{\sigma_{v,blur}^2 + \sigma_p^2}$, we can

calculate the standard deviation of the integrated sensory signal for the blurred cursor:

$\sigma_{blur} = \sigma\sqrt{\frac{5R^2}{R^2 + 4}}$. Simulation was performed with R ranging from 1.1 to 3.

Data fitting

All data were fitted using MATLAB 2022b (MathWorks, Natick, MA, US) built-in function

fmincon with 100 randomly sampled initial values of parameter sets. See Table S1 and

Table S2 for the fitted parameter values and comparisons between different models.

References

Albert, S. T., Jang, J., Modchalingam, S., Marius 't Hart, B., Henriques, D., Lerner, G., Della-Maggiore, V., Haith, A. M., Krakauer, J. W., & Shadmehr, R. (2022). Competition between parallel sensorimotor learning systems. *eLife*, *11*, e65361.

Albert, S. T., Jang, J., Sheahan, H. R., Teunissen, L., Vandevoorde, K., Herzfeld, D. J., & Shadmehr, R. (2021). An implicit memory of errors limits human sensorimotor adaptation. *Nature Human Behaviour*, 1–15.

Avraham, G., Morehead, J. R., Kim, H. E., & Ivry, R. B. (2021). Reexposure to a sensorimotor perturbation produces opposite effects on explicit and implicit learning processes. *PLoS Biology*, *19*(3), e3001147.

Avraham, G., Taylor, J. A., Breska, A., Ivry, R. B., & McDougle, S. D. (2022). Contextual effects in sensorimotor adaptation adhere to associative learning rules. *eLife*, *11*, e75801.

Berniker, M., & Kording, K. (2008). Estimating the sources of motor errors for adaptation and generalization. *Nature Neuroscience*, *11*(12), 1454–1461.

Berniker, M., & Kording, K. P. (2011). Estimating the relevance of world disturbances to explain savings, interference and long-term motor adaptation effects. *PLoS Computational Biology*, *7*(10), e1002210.

Bond, K. M., & Taylor, J. A. (2015). Flexible explicit but rigid implicit learning in a visuomotor adaptation task. *Journal of Neurophysiology*, *113*(10), 3836–3849.

Burge, J., Ernst, M. O., & Banks, M. S. (2008). The statistical determinants of adaptation rate in human reaching. *Journal of Vision*, *8*(4), 1–19.

Cheng, S., & Sabes, P. N. (2006). Modeling sensorimotor learning with linear dynamical systems. *Neural Computation*, *18*(4), 760–793.

Cressman, E. K., & Henriques, D. Y. P. (2009). Sensory recalibration of hand position following visuomotor adaptation. *Journal of Neurophysiology*, *102*(6), 3505–3518.

de Brouwer, A. J., Albaghdadi, M., Flanagan, J. R., & Gallivan, J. P. (2018). Using gaze behavior to parcellate the explicit and implicit contributions to visuomotor learning. *Journal of Neurophysiology*, *120*(4), 1602–1615.

de Brouwer, A. J., Gallivan, J. P., & Flanagan, J. R. (2018). Visuomotor feedback gains are modulated by gaze position. *Journal of Neurophysiology*, *120*(5), 2522–2531.

Della-Maggiore, V., & McIntosh, A. R. (2005). Time course of changes in brain activity and functional connectivity associated with long-term adaptation to a rotational transformation. *Journal of Neurophysiology*, *93*(4), 2254–2262.

Donchin, O., Francis, J. T., & Shadmehr, R. (2003). Quantifying generalization from trial-by-trial behavior of adaptive systems that learn with basis functions: theory and experiments in human motor control. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *23*(27), 9032–9045.

He, K., Liang, Y., Abdollahi, F., Bittmann, M. F., Körding, K., & Wei, K. (2016). The statistical determinants of the speed of motor learning. *PLoS Computational Biology*, *12*(9), e1005023.

Henriques, D. Y. P., & Cressman, E. K. (2012). Visuomotor adaptation and proprioceptive recalibration. *Journal of Motor Behavior*, *44*(6), 435–444.

Herzfeld, D. J., Vaswani, P. A., Marko, M. K., & Shadmehr, R. (2014). A memory of errors in sensorimotor learning. *Science*, *345*(6202), 1349–1353.

Huberdeau, D. M., Krakauer, J. W., & Haith, A. M. (2019). Practice induces a qualitative change in the memory representation for visuomotor learning. *Journal of Neurophysiology*, *122*(3), 1050–1059.

Hutter, S. A., & Taylor, J. A. (2018). Relative sensitivity of explicit reaiming and implicit motor adaptation. *Journal of Neurophysiology*, *120*(5), 2640–2648.

Izawa, J., & Shadmehr, R. (2011). Learning from sensory and reward prediction errors during motor adaptation. *PLoS Computational Biology*, *7*(3), e1002012.

Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R., & Ivry, R. B. (2018). Invariant errors reveal limitations in motor correction rather than constraints on error sensitivity. *Communications Biology*, *1*(1), 19.

Kirk, R. (1968). *Experimental design: Procedures for the behavioral sciences* (Vol. 1–f). Brooks/Cole.

Klein, S. A., & Levi, D. M. (1987). Position sense of the peripheral retina. *JOSA A*, *4*(8), 1543–1553.

Körding, K., & Wolpert, D. M. (2004). Bayesian integration in sensorimotor learning. *Nature*, *427*(6971), 244–247.

Krakauer, J. W., Ghez, C., & Ghilardi, M. F. (2005). Adaptation to visuomotor transformations: consolidation, interference, and forgetting. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *25*(2), 473–478.

Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L., & Haith, A. M. (2019). Motor learning. *Comprehensive Physiology*, *9*(2), 613–663.

Landi, S. M., Baguear, F., & Della-Maggiore, V. (2011). One week of motor adaptation induces structural changes in primary motor cortex that predict long-term memory one year later. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *31*(33), 11808–11813.

Leow, L.-A., Marinovic, W., de Rugy, A., & Carroll, T. J. (2020). Task errors drive memories that improve sensorimotor adaptation. *Journal of Neuroscience*, *40*(15), 3075–3088.

Levi, D. M., Klein, S. A., & Yap, Y. L. (1987). Positional uncertainty in peripheral and amblyopic vision. *Vision Research*, *27*(4), 581–597.

Lieberman, H. R., & Pentland, A. P. (1982). Microcomputer-based estimation of psychophysical thresholds: the best PEST. *Behavior Research Methods & Instrumentation*, *14*(1), 21–25.

Mazzoni, P., & Krakauer, J. W. (2006). An implicit plan overrides an explicit strategy during visuomotor adaptation. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *26*(14), 3642–3645.

McDougle, S. D., Bond, K. M., & Taylor, J. A. (2015). Explicit and Implicit Processes Constitute the Fast and Slow Processes of Sensorimotor Learning. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *35*(26), 9568–9579.

Mikulasch, F. A., Rudelt, L., & Priesemann, V. (2022). Visuomotor mismatch responses as a hallmark of explaining away in causal inference. *Neural Computation*, *35*(1), 27–37.

Miyamoto, Y. R., Wang, S., & Smith, M. A. (2020). Implicit adaptation compensates for erratic explicit strategy in human motor learning. *Nature Neuroscience*, *23*(3), 443–455.

Modchalingam, S., Vachon, C. M., 't Hart, B. M., & Henriques, D. Y. P. (2019). The effects of awareness of the perturbation during motor adaptation on hand localization. *PloS One*, *14*(8), e0220884.

Morehead, J. R., Taylor, J. A., Parvin, D. E., & Ivry, R. B. (2017). Characteristics of implicit sensorimotor adaptation revealed by task-irrelevant clamped feedback. *Journal of Cognitive Neuroscience*, *29*(6), 1061–1074.

Mostafa, A. A., 't Hart, B. M., & Henriques, D. Y. P. (2019). Motor learning without moving: Proprioceptive and predictive hand localization after passive visuoproprioceptive discrepancy training. *PloS One*, *14*(8), e0221861.

Neville, K. M., & Cressman, E. K. (2018). The influence of awareness on explicit and implicit contributions to visuomotor adaptation over time. *Experimental Brain Research. Experimentelle Hirnforschung. Expérimentation Cerebrale*, *236*(7), 2047–2059.

Oh, Y., & Schweighofer, N. (2019). Minimizing Precision-Weighted Sensory Prediction Errors via Memory Formation and Switching in Motor Adaptation. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *39*(46), 9237–9250.

Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia*, *9*(1), 97–113.

Pearson, T. S., Krakauer, J. W., & Mazzoni, P. (2010). Learning not to generalize: modular

adaptation of visuomotor gain. *Journal of Neurophysiology*, *103*(6), 2938–2952.

Petitet, P., O'Reilly, J. X., & O'Shea, J. (2018). Towards a neuro-computational account of prism

adaptation. *Neuropsychologia*, *115*, 188–203.

Rossi, C., Bastian, A. J., & Therrien, A. S. (2021). Mechanisms of proprioceptive realignment in

human motor learning. *Current Opinion in Physiology*, *20*, 186–197.

Ruttle, J. E., Cressman, E. K., 't Hart, B. M., & Henriques, D. Y. P. (2016). Time course of reach

adaptation and proprioceptive recalibration during visuomotor learning. *PloS One*, *11*(10),

e0163695.

Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. P. (2018). The fast contribution of

visual-proprioceptive discrepancy to reach aftereffects and proprioceptive recalibration.

PloS One, *13*(7), e0200621.

Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. P. (2021). Implicit motor learning within three

trials. *Scientific Reports*, *11*(1), 1627.

Salomonczyk, D., Cressman, E. K., & Henriques, D. Y. P. (2011). Proprioceptive recalibration

following prolonged training and increasing distortions in visuomotor adaptation.

Neuropsychologia, *49*(11), 3053–3062.

Salomonczyk, D., Cressman, E. K., & Henriques, D. Y. P. (2013). The role of the cross-sensory

error signal in visuomotor adaptation. *Experimental Brain Research. Experimentelle*

Hirnforschung. Experimentation Cerebrale, *228*(3), 313–325.

Shadmehr, R., & Mussa-Ivaldi, F. A. (1994). Adaptive representation of dynamics during

learning of a motor task. *Journal of Neuroscience*, *14*(5), 3208–3224.

Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and

adaptation in motor control. *Annual Review of Neuroscience*, *33*, 89–108.

Simani, M. C., McGuire, L. M. M., & Sabes, P. N. (2007). Visual-shift adaptation is composed of

separable sensory and task-dependent effects. *Journal of Neurophysiology*, *98*(5),

2827–2841.

Smith, M. A., Ghazizadeh, A., & Shadmehr, R. (2006). Interacting adaptive processes with

different timescales underlie short-term motor learning. *PLoS Biology*, *4*(6), e179.

Stetson, C., Cui, X., Montague, P. R., & Eagleman, D. M. (2006). Motor-sensory recalibration

leads to an illusory reversal of action and sensation. *Neuron*, *51*(5), 651–659.

Taylor, J. A., & Ivry, R. B. (2011). Flexible cognitive strategies during motor learning. *PLoS*

Computational Biology, *7*(3), e1001096.

Taylor, J. A., Krakauer, J. W., & Ivry, R. B. (2014). Explicit and implicit contributions to learning in

a sensorimotor adaptation task. *The Journal of Neuroscience: The Official Journal of the*

Society for Neuroscience, *34*(8), 3023–3032.

't Hart, B. M., & Henriques, D. Y. P. (2016). Separating predicted and perceived sensory

consequences of motor learning. *PloS One*, *11*(9), e0163556.

Thoroughman, K. A., & Shadmehr, R. (2000). Learning of action through adaptive combination

of motor primitives. *Nature*, *407*(6805), 742.

Tsay, J. S., Avraham, G., Kim, H. E., Parvin, D. E., Wang, Z., & Ivry, R. B. (2021). The effect of

visual uncertainty on implicit motor adaptation. *Journal of Neurophysiology*, *125*(1), 12–22.

Tsay, J. S., Chandy, A. M., Chua, R., Miall, R. C., Cole, J., Farnè, A., Ivry, R. B., & Sarlegna, F.

R. (2024). Minimal impact of proprioceptive loss on implicit sensorimotor adaptation and

perceived movement outcome. *bioRxiv : The Preprint Server for Biology*.

https://doi.org/10.1101/2023.01.19.524726

Tsay, J. S., Haith, A. M., Ivry, R. B., & Kim, H. E. (2022). Interactions between sensory prediction

error and task error during implicit motor learning. *PLoS Computational Biology*, *18*(3),

e1010005.

Tsay, J. S., Kim, H. E., Parvin, D. E., Stover, A. R., & Ivry, R. B. (2021). Individual differences in

proprioception predict the extent of implicit sensorimotor adaptation. *Journal of*

Neurophysiology, *125*(4), 1307–1321.

Tsay, J. S., Kim, H. E., Saxena, A., Parvin, D. E., Verstynen, T., & Ivry, R. B. (2022). Dissociable

use-dependent processes for volitional goal-directed reaching. *Proceedings. Biological*

Sciences / The Royal Society, *289*(1973), 20220415.

Tsay, J. S., Kim, H., Haith, A. M., & Ivry, R. B. (2022). Understanding implicit sensorimotor

adaptation as a process of proprioceptive re-alignment. *eLife*, *11*, e76639.

Tsay, J. S., Parvin, D. E., & Ivry, R. B. (2020). Continuous reports of sensed hand position

during sensorimotor adaptation. *Journal of Neurophysiology*, *124*(4), 1122–1130.

Tseng, Y., Diedrichsen, J., Krakauer, J. W., Shadmehr, R., & Bastian, A. J. (2007). Sensory

prediction errors drive cerebellum-dependent adaptation of reaching. *Journal of*

Neurophysiology, *98*(1), 54.

von Helmholtz, H. (1867). *Handbuch der physiologischen Optik*. Voss.

Wei, K., & Körding, K. (2009). Relevance of error: what drives motor adaptation? *Journal of*

Neurophysiology, *101*(2), 655–664.

Wei, K., & Körding, K. (2010). Uncertainty of feedback and state estimation determines the

speed of motor adaptation. *Frontiers in Computational Neuroscience*, *4*, 11.

Wolpert, D. M., Diedrichsen, J., & Flanagan, J. R. (2011). Principles of sensorimotor learning.

Nature Reviews. Neuroscience, *12*(12), 739.

Wozny, D. R., & Shams, L. (2011). Recalibration of auditory space following milliseconds of

cross-modal discrepancy. *Journal of Neuroscience*, *31*(12), 4607–4612.

Wu, H. G., Miyamoto, Y. R., Castro, L. N. G., Ölveczky, B. P., & Smith, M. A. (2014). Temporal

structure of motor variability is dynamically regulated and predicts motor learning ability.

Nature Neuroscience, *17*(2), 312.

1153

1154

1155

Figure 1. The Perceptual Error Adaptation (PEA) model for error-clamp adaptation. **(A)** Illustration of involved sensorimotor cues for estimating hand direction $\hat{x}_{Hand}$. The clamped cursor, the hand, and the sensory prediction of the reaching action provide the visual (x_v), proprioceptive (x_p), and the sensory prediction cue (x_u) of movement direction, respectively. The hand direction estimate is assumed to be based on maximum likelihood cue combination. **(B)** Assuming a linear dependency of visual uncertainty on eccentricity, the PEA model predicts that implicit adaptation extent is a concave function of perturbation size θ, a pattern qualitatively aligning with previous findings (Kim et al., 2018; Morehead et al., 2017).

1165

Figure 2. Experiment 1 measuring the dependency of visual uncertainty on perturbation size. **(A)** Illustration of all possible cursor endpoints during the experiment and the procedure of the 2AFC task for judging the cursor motion direction. In an exemplary trial, the participant reaches to a target while a direction-clamped cursor moves concurrently, serving as an error-clamp perturbation. Following a 1000-ms blank masking period, a reference point appears for 150ms, either clockwise or counterclockwise from the clamped cursor. The participant is then asked to make a binary judgment regarding the direction of the clamped cursor relative to the reference point. **(B)** The visual uncertainty, obtained from psychometrical estimation based on the 2AFC, is plotted as a function of perturbation size. Both individual estimates (red dots)

and group-level statistics (boxplots) are shown. Positive angles correspond to CW

rotations, negative angles to CCW rotations. **(C)** Collapsing data from both rotation

directions, we observe that visual uncertainty closely follows a linear function of

perturbation size. The dark gray line and its shaded region denote the regression line

and its 95% confidence interval, respectively. The purple line is generated with the

values of *a* and *b* fitted from data in Experiment 2 with *a* and *b* treated as free

parameters (See Methods for details).

1183

Figure 3. Results and model fitting of Experiment 2. **(A)** Implicit adaptation to error

clamps of varying sizes is depicted; colored dot-lines and colored shading area

represent the mean and standard error for each participant group. The light gray area

indicates trials with error-clamp perturbations. Adaptation starts after baseline, gradually

asymptotes to its final extent, and then decays with null feedback during washout.

Different perturbation sizes result in distinct adaptation rates and extents. Group

averages and standard error across participants are shown, along with predictions

(colored solid lines) from the PEA model. **(B)** The adaptation extent (cycle 100-110)

exhibits a nonlinear dependency on perturbation size, conforming to a concave function

as prescribed by PEA (purple line). Color dots and error bars denote the mean and

1194 standard error across participants in each group. **(C)-(D)** The same data fitted with the

1195 PReMo model and the causal inference model. For more details, refer to Figure S3.

1196

1197

Figure 4. Proprioceptive data fitting and results from Experiment 3. **(A)** The data from

(Tsay et al., 2020) are presented alongside the fitting of the PEA model. Participants

adapting to a 30° error-clamp perturbation were required to report the location of their

adapted hand using visual aids of numbers. The report was provided when the hand

stayed at the end of movement. Initially, the proprioceptive estimate of the hand is

biased toward the visual cursor (negative in the plot) and then gradually shifts toward

the hand (positive in the plot). This trend is accurately captured by the PEA model: lines

represent model fitting results, with the adapted hand direction in indigo and the

reported hand direction in blue. The hand direction estimate ($\hat{x}_{Hand}$, Eq.1) following a

reach movement is shown in red. **(B)-(C)** Model simulations for proprioceptive bias from

the PEA and PReMo models. Color gradients denote the simulations with varying ratio

between the weights of $\hat{x}_{Hand}$ and x_p, the two cues available for estimating the hand

direction. Note that the two models prescribe distinct profiles for the dependency of

proprioception bias on perturbation size. **(D)** Experimental design. A reaching block,

either with or without visual perturbations, is followed by a proprioception test block. The

size and direction of the visual perturbation vary across blocks. The proprioception test

is conducted when the hand is passively moved to a target (red dots) situated near the

reaching target (blue dots). **(E)** The observed proprioceptive bias as a function of

perturbation size. Data from the three proprioception test trials are separately plotted.

The first trial reveals proprioception biases that form a concave function of perturbation

size.

1219

Figure 5. Results of Experiment 4. **(A)-(C)** Model simulations for single-trial learning under different visual uncertainty levels, shown separately for the PEA, PReMo and causal inference models. Blue curves represent simulated learning based on model parameters estimated from Experiment 2. Curves with red gradient indicate simulations with increasing levels of visual uncertainty, color coded by the ratio of visual uncertainty for the blurred cursor to that of the clear cursor. **(D)** Experimental design. Following 60 baseline trials without perturbations, participants completed 15 mini-blocks of error-clamp adaptation over three successive days. Each mini-block features 12 different types of error-clamp perturbations, distinguished by two cursor presentations (blurred or clear cursor) and six clamp sizes. Each perturbation trial, varied randomly in perturbation type, is flanked by two no-feedback trials. The change in hand direction

1231 between these two no-feedback trials serves to quantify singe-trial learning. **(E)** The

1232 single-trial learning with the blurred cursor is less than that with the clear cursor, but the

1233 difference is non-monotonic across perturbation size (*** denote p < 0.001).

1234

Figure 6. Design of Experiment 1. **(A)** Top-down view of the setup in visual discrimination task. The reference point (yellow) was presented either CW or CCW relative to the clamped cursor (dashed circle), which has a perturbation size θ. **(B)** Trial structure of the visual discrimination task. Purple rectangles represent error-clamped trials with varying perturbation size, rectangles with yellow edges represent the ensuing visual discrimination test for each perturbation size. **(C)-(D)** Exemplary sequences of the reference point: These sequences illustrate the deviation of the reference point from the cursor (C) and the changing step size across trials (D), following the PEST algorithm. Individual trials are represented by blue dots. Yellow and red dots mark the initiation and termination of each round of trials, respectively. In each round, the reference point starts on either the CW or CCW side of the cursor; In the subsequent round, it starts on the opposite side.

1247

Figure 7. Setup for measuring proprioceptive recalibration in Experiment 3. **(A)** Reaching movement with error-clamped cursor, performed by the right hand holding a robot handle. **(B)** Passive movement in the proprioception test. The right hand was passively moved to the unseen target (h_R), depicted here as a small black dot. A red hollow circle with an expanding radius appears on the screen during passive movement, signaling the increasing distance between the hand and the start position. Subsequently, participants used their left hand to report the right-hand location (h_p) by aligning a red rectangle on the red circle, which is displayed at the target distance.

1256 **Supplementary Materials**

1257

1258

Figure S1. Heat map of eye fixations during the 2AFC task in Experiment 1. The screen

is partitioned into 10x10 pixel grids, and the cumulative number of gaze samples in

each grid is recorded. Data from all participants, aggregated across each day of

practice, are presented. The color map signifies the normalized count of gaze samples

in each grid. Data are separately displayed for the three distinct phases of a trial, as

delineated by the columns on the left, middle, and right. These correspond to periods

during hand movement, the appearance of the visual mask and reference point, and the

time allotted for manual response. On average, 95.06%, 89.93%, and 86.55% of gaze

samples fall within the ±50-pixel range of the central line during these three phases,

respectively. These results corroborate that participant adhered to the instructions and

refrained from looking at the cursor during the visual discrimination task.

1270

1271

Figure S2. Performance of an exemplary participant in Experiment 1. Six panels display the psychometric curves corresponding to different error-clamp sizes. The x-axis denotes the angular deviation between the clamped cursor and the reference point (as depicted in Figure 6A). A negative value implies that the reference point appears on the counterclockwise (CCW) side of the clamped cursor. The blue dots represent the proportion of trials in which the participant reported that "the yellow point is on the clockwise (CW) side of the clamped cursor" for various angular deviations between these two. Data were aggregated from all trials across three days of the experiment. The gray-shaded region represents the interquartile range (25th to 75th percentile) of the psychometric curve, and the width of this shaded region serves as an indicator of the amplitude of visual uncertainty.

1283

Figure S3. Model fitting for observed implicit adaptation in Experiment 2. This supplementary figure provides a comprehensive evaluation of the three competing models: the PEA model, the PReMo model, and the causal inference model. **(A)** Results of PEA Model Fitting: The layout of these plots mirrors that of Figures 3A and

3B, serving as a direct comparison between the empirical data and the predictions made by the PEA model. **(B)** Results of PReMo Model Fitting: The left panel is a duplicate of Figure 3C, while the right panel presents the trial-by-trial data fitting. This juxtaposition allows for a nuanced evaluation of the PReMo model's performance at both the aggregate and individual trial levels. **(C)** Results of Causal Inference Model: The arrangement of these plots is consistent with panels (A) and (B), facilitating a straightforward comparison of all three models. For a detailed assessment of the quality of model fitting and subsequent model comparisons, please refer to Table S1 and Table S2.

1298

Figure S4. Correlation between initial learning rate and adaptation extent in Experiment 2. For each participant, the initial learning rate is calculated as the change in hand angle between the 1st and 10th cycle, divided by 10. The adaptation extent is defined as the average hand angle across the last 10 adaptation cycles. When pooling data across all perturbation sizes, a significant correlation is observed between the initial learning rate and the adaptation extent.

1305

1306 **Figure S5.** Model fitting of single-trial learning from Experiment 2 of (Tsay, Avraham, et

1307 al., 2021). Blue dots represent the mean single-trial learning across varying perturbation

1308 size, with error bars denoting standard errors across participants. The left, middle and

1309 right panel present the fitting results for the PEA, PReMo, and causal inference models,

1310 respectively. For additional details, refer to the Methods, Results, and Table S1 & S2.

1311

Figure S6. Proprioception uncertainty estimation results. Thirteen participants from Experiment 1 participated in a proprioception discrimination task to measure their proprioceptive uncertainty in the setting of the error-clamp adaptation. The setup paralleled that used for estimating visual uncertainty in Experiment 1. In each trial, participants initially held their hand at the starting position. They were instructed to relax their arm while the experimenter, seated on the other side of the monitor, pulled their hand to a proprioceptive target near the straight-ahead target. The arms of the experimenter and the participant were blocked from the view of the participant. After 0.8 seconds, a yellow reference point appeared. The angular deviation between the participant's hand and this reference was determined using the same PEST procedure employed in Experiment 1. Participants indicated, by pressing left or right arrow keys by their left hand, whether the reference point appeared on the CW or CCW side of their actual right-hand position. The maximum deviation allowed was 30°, with an initial step

1325 size of 10° and a stop threshold of 0.5°. This task was conducted over six runs across

1326 three consecutive days. Similar to Figure S2, panel (A) to (M) show the psychometric

1327 curves for each participant with data from the three days pooled together. (N) and (O)

1328 present the measured proprioceptive uncertainty and bias for all participants (gray dots)

1329 and their mean ± standard deviation (red error bars).

Figure S7. Alternative Model fitting with PEA for Experiment 2; see details in Supplementary Text 2. **(A)** Results of PEA Model Fitting. The layout is similar to Figure 3A and Figure S3A. **(B)** Fitted ratio between uncertainties of different cues. **(C)** Estimated actual value of uncertainties of different cues.

1338 **Table S1.** Model fitting and simulation parameters with the PEA model.

	Data set	Parameters						Goodness-of-fit	
		σ_u (deg)	σ_p (deg)	a	b	A	B	R^2	RMSE (deg)
Adaptation extent fitting (Figure 1B)	Kim 2018, Exp1	--	σ_p/a = 3.406; b/a = 0.138			--	--	0.773	1.898
	Kim 2018, Exp2	--	σ_p/a = 4.758; b/a = 0.168			--	--	<0	2.163
	Morehead, 2017	--	σ_p/a = 1.639; b/a = 0.044			--	--	<0	2.937
Trial-by-trial adaptation fitting	Exp 2, Figure 3	5.048	11.119	*1.853	*0.309	0.970	0.208	0.975	1.222
		5.468	12.128	1.663	0.331	0.971	0.194	0.975	1.217
	Exp 2, Figure S7	--	--	--	--	0.991	0.364	0.989	0.815
	Tsay 2019, Figure 4A	1.896	7.959	*1.853	*0.309	0.978	0.525	0.991	0.716
Single-trial learning fitting	Tsay 2021, Figure S5	σ_{int} = 7.364		1.179	0.384	--	0.057	0.974	0.020
Proprioceptive recalibration simulation	Exp 3, Figure 4B	5.048	11.119	1.853	0.309	0.970	0.208	--	--
Adaptation affected by visual uncertainty simulation	Exp 4, Figure 5	5.048	11.119	1.853	0.309	--	0.208	--	--

1339 * Asterisks represent fixed parameters in specific data fitting. The fixed values equal to

1340 the slope and intercept estimated from Experiment 1.

1341 **Table S2.** Model comparisons.

Data set		PEA	PReMo	Causal Inference
Block-design learning fitting *Exp 2, Figure 3 & S3*	AIC	2255	3543	3283
	R^2	0.975	0.749	0.711
	$RMSE$ (deg)	1.222	3.896	4.151
Single-trial learning fitting *Tsay 2021, Figure S5*	AIC	-36.90	-15.98	-11.28
	R^2	0.974	0.584	0.305
	$RMSE$ (deg)	0.020	0.103	0.080

1342

1343 **Supplementary Text 1** Data fitting of Figure 4A

1344 We use u, v, p and h to denote the reciprocal of the variances of motor command cue,

1345 visual cue, proprioceptive cue and combined cue (hand estimation $\hat{x}_{Hand}$) respectively:

1346
$$u = 1/\sigma_u^2,\ v = 1/\sigma_v^2,\ p = 1/\sigma_p^2,$$

1347
$$h = 1/\sigma_{hand}^2 = 1/\sigma_u^2 + 1/\sigma_v^2 + 1/\sigma_p^2 = u + v + p.$$

1348 According to Eq1 in the main text, the estimated hand can be denoted by:

1349
$$\hat{x}_{Hand,n} = \frac{v}{h}x_v + \frac{p}{h}x_{p,n} + \frac{u}{h}x_{u,n}, \tag{S1}$$

1350 where $x_{u,n} = 0$ indicates the target position (T). We further define the weight parameters

1351 W_v and W_p as:

1352
$$W_v = \frac{v}{h},$$

1353
$$W_p = \frac{p}{h},$$

1354 and Eq S1 can be wrote as:

1355
$$\hat{x}_{Hand,n} = W_v x_v + W_p x_{p,n}, \tag{S2}$$

1356 insert Eq S2 into Eq 2 from the main text, the hand position at trial n is:

1357
$$x_{p,n+1} = Ax_{p,n} + B\left(0 - W_v x_v + W_p x_{p,n}\right). \tag{S3}$$

1358 Next, with the same Bayesian cue combination assumption, the reported hand position

1359 was combined with x_p and $\hat{x}_{Hand}$:

1360
$$x_{Report,n} = W_{\overline{Hand}}\hat{x}_{Hand,n} + W_{p^*}x_{p,n}, \tag{S4}$$

1361 where $W_{\overline{Hand}}$ and W_{p^*} are the weight parameters:

1362
$$W_{\widehat{Hand}} = \frac{h}{h+p} = \frac{1}{1+W_p},$$

1363
$$W_{p^*} = \frac{p}{h+p} = \frac{W_p}{1+W_p},$$

1364 put the weight parameters and Eq S2 into Eq S4, we can easily get:

1365
$$x_{Report,n} = \frac{W_v}{1+W_p}x_v + \frac{2W_p}{1+W_p}x_{p,n}, \tag{S5}$$

1366 The adaptation epoch of the actual and reported hand position in Figure 4A was fitted

1367 by Eq S3 and Eq S5 respectively with the parameter set $\Theta = [W_v, W_p, A, B]$.

1368 We simulated the wash out epoch based on the parameters estimated from the

1369 adaptation epoch with the method below (dashed line in Figure 4A).

1370 Since the visual cue was no longer available, the weight of cues changed to:

1371
$$W_{v[wo]} = 0,$$

1372
$$W_{p[wo]} = \frac{p}{u+p} = \frac{p}{h-v} = \frac{W_p}{1-W_v}.$$

1373 Since $x_{u,n} = 0$, the estimated hand in wash out epoch is:

1374
$$\hat{x}_{Hand,n} = W_{v[wo]}x_v + W_{p[wo]}x_{p,n}, \tag{S6}$$

1375 put $W_{v[wo]}$ $W_{p[wo]}$ and Eq S6 into Eq 2 in the main text, the hand position at trial n is:

1376
$$x_{p,n+1} = Ax_{p,n} + B\left(0 - \frac{W_p}{1-W_v}x_{p,n}\right). \tag{S7}$$

1377 Similar to S4, the reported hand position is:

1378
$$x_{Report,n} = W_{\widehat{Hand}[wo]}\hat{x}_{Hand,n} + W_{p^*[wo]}x_{p,n}, \tag{S8}$$

1379 where $W_{\widehat{Hand}[wo]}$ and $W_{p^*[wo]}$ denote the corresponding weight of each cue:

1380
$$W_{\widehat{Hand}[wo]} = \frac{h_{[wo]}}{h_{[wo]}+p} = \frac{1-W_v}{1-W_v+W_p},$$

1381
$$W_{p^*[wo]} = \frac{p}{h_{[wo]}+p} = \frac{W_p}{1-W_v+W_p},$$

1382 where $h_{[wo]}$ denotes the reciprocal of the variances of estimated hand during washout

1383 epoch:

1384
$$h_{[wo]} = u + p = h - v.$$

1385 Put the weight parameters and S6 into S8, the reported hand position can be easily

1386 derived:

1387
$$x_{Report,n} = (\frac{W_p(1-W_v)}{(1-W_v+W_p)(1-W_v)} + \frac{W_p}{1-W_v+W_p})x_{p,n} \tag{S9}$$

1388 Eq S7 and Eq S9 were used to simulate the hand position and report position during

1389 washout epoch with $\Theta = [W_v, W_p, A, B]$ estimated from the adaptation epoch.

1390 **Supplementary Text 2** Fitting block designed adaptation data (Experiment 2) without

1391 the assumption of linearity in visual uncertainty

1392 For each perturbation size i, the weight of visual and proprioceptive cues are:

$$1393 \qquad W_v^{[i]} = \frac{1/\sigma_{v[i]}^2}{1/\sigma_{u[i]}^2 + 1/\sigma_{v[i]}^2 + 1/\sigma_{p[i]}^2}, \tag{S10}$$

$$1394 \qquad W_p^{[i]} = \frac{1/\sigma_{p[i]}^2}{1/\sigma_{u[i]}^2 + 1/\sigma_{v[i]}^2 + 1/\sigma_{p[i]}^2}, i = 1 \cdots 7, \tag{S11}$$

1395 where $\sigma_{u[i]}$, $\sigma_{v[i]}$ and $\sigma_{p[i]}$ are the uncertainties of sensory cues under different

1396 perturbation sizes (2 to 95 degree). The estimated hand can be calculated by putting Eq

1397 S10 and Eq S11 into Eq 1 from the main text:

$$1398 \qquad \hat{x}_{Hand,n} = W_v^{[i]} x_v + W_p^{[i]} x_{p,n} + \frac{1/\sigma_{u[i]}^2}{1/\sigma_{u[i]}^2 + 1/\sigma_{v[i]}^2 + 1/\sigma_{p[i]}^2} x_{u,n},$$

1399 where $x_{u,n} = T = 0$ denotes the target location, thus:

$$1400 \qquad \hat{x}_{Hand,n} = W_v^{[i]} x_v + W_p^{[i]} x_{p,n}, \tag{S12}$$

1401 Insert Eq S12 into Eq 2 from the main text, the trial-by-trail hand position is:

$$1402 \qquad x_{p,n+1} = -BW_v^{[i]} x_v + (A - BW_p^{[i]}) x_{p,n} \tag{S13}$$

1403 The parameter set $\Theta = [W_v^{[1]} \cdots W_v^{[7]}, W_p^{[1]} \cdots W_p^{[7]}, A, B]$ were fitted to the data in

1404 Experiment 2. Data fitting was done with a Bayesian optimization algorithm (Acerbi and

1405 Ma, 2017) with 100 runs of different initial values between a set of predetermined

1406 preferred boundaries.

1407 The ratio between uncertainties of different sensory cues are:

1408
$$\sigma_{v[i]}/\sigma_{u[i]} = \sqrt{(1 - W_v^{[i]} - W_p^{[i]})/W_v^{[i]}}$$
(S14)

1409
$$\sigma_{p[i]}/\sigma_{u[i]} = \sqrt{(1 - W_v^{[i]} - W_p^{[i]})/W_p^{[i]}}$$
(S15)

The fitting results are presented in Figure S7A. Assuming σ_u remains constant across

different perturbation sizes, we convert σ_p and σ_v into ratios, Figure S7B illustrates the

ratio between estimated uncertainties at different perturbation sizes.

To determine the values shown in Figure S7C, we proceeded as follows: First, we found

no significant correlation between perturbation sizes and the estimated values of σ_p (R =

-0.476, p = 0.281). The mean of σ_p is thus set to the mean value derived from the

proprioception uncertainty experiment (Figure S6), with σ_p = 9.737° for all perturbation

sizes, and σ_u = 3.681° was also calculated as a ratio of σ_p. Those results are indicated

by the two horizontal dashed lines in Figure S7C. Subsequently, we estimated σ_v for

different perturbation sizes, represented by dark brown dots in Figure S7C. To compare

the visual uncertainty estimated from Experiment 2 with the results from Experiment 1,

we performed a linear fit of the estimated σ_v values and perturbation sizes using

first-order polynomials. The fitting yielded an intercept a = 3.617 and a slope b = 0.261

(R^2 = 0.982, p<0.001). The slope and intercept are in close agreement with those from

Experiment 1 (the gray line in Figure S7C), and the fitting results also confirmed a

strong linear relationship between σ_v and perturbation sizes.

Reference

Acerbi, L., & Ma, W. J. (2017). Practical Bayesian optimization for model fitting with

Bayesian adaptive direct search. Advances in Neural Information Processing Systems.