Perceptual error based on Bayesian cue combination

drives implicit motor adaptation

Zhaoran Zhang[1, †], Huijun Wang[1, †], Tianyang Zhang[1], Zixuan Nie[1], Kunlin Wei[1,2,3,4 *]

[1]School of Psychological and Cognitive Sciences, Peking University, Beijing, China

[2]Beijing Key Laboratory of Behavior and Mental Health, Beijing, China

[3]Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China

[4]National Key Laboratory of General Artificial Intelligence, Beijing, China

[†]Zhaoran Zhang and Huijun Wang are co-first authors.

[*]Correspondence to: Kunlin Wei
18   School of Psychological and Cognitive Sciences
19   Peking University
20   5 Yiheyuan Road
21   Beijing, 100871, China
22   wei.kunlin@pku.edu.cn

Abstract

The sensorimotor system can recalibrate itself without our conscious awareness, a type

of procedural learning whose computational mechanism remains undefined. Recent

findings on implicit motor adaptation, such as over-learning from minor perturbations and

swift saturation for increasing perturbation size, challenge existing theories based on

sensory errors. We argue that perceptual error, arising from the optimal combination of

movement-related cues, is the primary driver of implicit adaptation. Central to our theory

is the linear relationship between the sensory uncertainty of visual cues and perturbation,

validated through perceptual psychophysics (Experiment 1). Our theory predicts diverse

features of implicit adaptation across a spectrum of perturbation conditions on trial-by-

trial basis (Experiment 2) and explains proprioception changes and their relation to visual

perturbation (Experiment 3). By altering visual uncertainty in perturbation, we induced

unique adaptation responses (Experiment 4). Overall, our perceptual error framework

outperforms existing models, suggesting that Bayesian cue integration underpins the

sensorimotor system's implicit adaptation.

Introduction

To achieve and sustain effective motor performance, humans consistently recalibrate

their sensorimotor systems to adapt to both internal and external environmental

disturbances (Berniker & Kording, 2008; Shadmehr et al., 2010; Wolpert et al., 2011). For

instance, transitioning to a high-sensitivity gaming mouse, which drives cursor movement

at an accelerated rate compared to a standard computer mouse, may initially result in

decreased performance in computer-related tasks. However, humans are capable of

rapidly adapting to this new visuomotor mapping within a short period of time. While

conscious corrections can facilitate this adaptation process, our sensorimotor system

often times adapts itself implicitly without our conscious efforts (Albert et al., 2021;

Krakauer et al., 2019).

While recent research has intensively examined the interplay between explicit and implicit

learning systems (Albert et al., 2022; Miyamoto et al., 2020), several characteristics of

implicit motor adaptation have emerged that challenge traditional theories. Conventionally,

motor adaptation is conceptualized as error-based learning, in which learning accrues in

proportion to the motor error experienced (Cheng & Sabes, 2006; Donchin et al., 2003;

Thoroughman & Shadmehr, 2000). However, implicit adaptation exhibits an

overcompensation phenomenon where the extent of adaptation surpasses the error

induced by visual perturbations (Kim et al., 2018; Morehead et al., 2017). Additionally,

implicit adaptation manifests a saturation effect; it increases with perturbations but

plateaus across a broad range of larger perturbations (Bond & Taylor, 2015; Kim et al.,

2018; Morehead et al., 2017; Neville & Cressman, 2018). These observations of

overcompensation and saturation are incongruent with prevailing state-space updating

models, which presuppose that incremental learning constitutes only a fraction of the

motor error (McDougle et al., 2015; Smith et al., 2006). Another aspect of implicit

adaptation that remains mechanistically unexplained pertains to its impact on

proprioception. In traditional motor adaptation, proprioception is biased towards the visual

perturbation, maintaining a stable bias throughout the adaptation process (Ruttle et al.,

2016, 2021). In contrast, implicit adaptation initially biases proprioceptive localization of

the hand towards the visual perturbation, but this bias gradually drifts in the opposite

direction over time (Tsay et al., 2020).

Causal inference of motor errors has been suggested to explain the discounting of large

perturbations (Wei & Körding, 2009). However, the causal inference account predicts a

decline in adaptation with increasing perturbation, diverging from the observed ramp-like

saturation effect. (Tsay, Kim, et al., 2022) recently synthesized existing evidence to

propose that implicit adaptation reaches an upper bound set by cerebellar error correction

mechanisms, reflected in a ramp-like influence of vision on proprioception (Tsay, Kim, et

al., 2022). While this ramp function could explain the observed saturation, the postulate

of an upper bound on visual influence lacks empirical validation. Some research supports

the idea of saturation in proprioceptive recalibration (Modchalingam et al., 2019), yet other

studies suggest a linear increase with visual perturbations (Rossi et al., 2021;

Salomonczyk et al., 2011). Additionally, current models fall short of quantitatively

capturing the time-dependent shifts in proprioceptive bias associated with implicit

adaptation.

In this study, we put forth a unified model that aims to account for the distinct features of

implicit adaptation, based on the Bayesian combination of movement-related cues. Prior

models have overlooked the fact that visual uncertainty related to the perturbation

increases with the size of the perturbation as the cursor moves further from the point of

fixation and into the visual periphery (Klein & Levi, 1987; Levi et al., 1987). This is

particularly pertinent for implicit adaptation that is widely investigated by the so-called

error-clamp paradigm, in which participants are instructed to fixate on the target and

disregard the perturbing cursor. Moreover, conventional theories of motor adaptation

attribute motor error to the sensory modality of the perturbation, i.e., visual errors for visual

perturbations (Tsay, Kim, et al., 2022; Wei & Körding, 2009). We propose an alternative:

perceptual error drives implicit adaptation, as the perturbed sensory feedback influences

the perception of the effector and, subsequently, motor adaptation. Through a series of

experiments, we aim to demonstrate that combining eccentricity-induced visual

uncertainty (Experiment 1) with a traditional motor adaptation model (state-space model)

and a classical perception model (Bayesian cue combination) can explain both over-

compensation and saturation effects (Experiment 2), as well as the time-dependent

changes in proprioceptive bias (Tsay et al., 2020). Finally, to provide causal evidence

supporting our Perceptual Error Adaptation (PEA) model, we manipulated visual

uncertainty and observed that subsequent adaptation was attenuated for large

perturbations but not for small ones—a finding that contradicts existing models but aligns

well with the PEA model. Across the board, our model outperforms those based on ramp

error-correction (Tsay, Kim, et al., 2022) and causal inference of errors (Wei & Körding,

2009), offering a more parsimonious explanation for the salient features of implicit

adaptation.

Results

The perceptual error adaptation model with varying visual uncertainty

We start by acknowledging that the perceptual estimation of effector position is

dynamically updated and influenced by sensory perturbations during motor adaptation.

For implicit adaptation studied via the error-clamp paradigm, participants are required to

bring their hand to the target while ignoring the direction-clamped cursor (Morehead et

al., 2017). Accordingly, the perceptual estimation of the hand movement direction relies

on three noisy sensory cues: the visual cue from the cursor, the proprioceptive cue from

the hand, and the sensory prediction of the reaching action (Figure 1A). Without loss of

generality, we posit that each cue is governed by an independent Gaussian distribution:

the visual cue x_v follows $N(\theta, \sigma_v^2)$, where θ is the cursor direction and σ_v^2 is visual

variance, the proprioceptive cue x_p follows $N(x_{hand}, \sigma_p^2)$, where x_{hand} is the hand

movement direction and σ_p^2 is proprioceptive variance, and the sensory prediction cue x_u

follows $N(T, \sigma_u^2)$, where T is the target direction and σ_u^2 is prediction variance.

Participants aim for the target, expecting their hand to reach it. Using the Bayesian cue

combination framework (Berniker & Kording, 2011), the perceived hand location ($\hat{x}_{Hand}$)

on trial *n* can be derived:

$$\hat{x}_{Hand,n} = \sum_i W_i x_{i,n}, \quad with \quad W_i = \frac{1/\sigma_i^2}{\sum_j 1/\sigma_j^2}, \quad i,j = u, p, v \qquad (1)$$

This estimated hand position is derived using maximum likelihood estimation from the

three noisy cues. Given that the clamped cursor deviates the target by θ, the visual cue

x_v biases the hand estimate $\hat{x}_{Hand}$ towards the cursor's direction. This deviation from the

target direction T constitutes the perceptual error, which drives adaptation on the

subsequent trial *n+1* (Eq. 2). Consisting with existing models (Albert et al., 2022; Cheng

& Sabes, 2006; Herzfeld et al., 2014; McDougle et al., 2015), trial-to-trial adaptation is

modeled using a state-space equation:

$$x_{p,n+1} = Ax_{p,n} + B\left(T - \hat{x}_{Hand,n}\right), \tag{2}$$

134   where A is the retention rate capturing inter-movement forgetting and B is the learning

135   rate capturing the proportion of error corrected within a trial. The interplay between

136   forgetting and learning dictates the overall learning extent, i.e., the asymptote of x_p:

$$x_p^{asym} = -\frac{B/\sigma_v^2}{B/\sigma_p^2 + (1-A)\sum_j 1/\sigma_j^2}\theta, \ j = v, p, u \tag{3}$$

138   Thus, the positive influence of perturbation size θ on the adaptation extent is

139   counterbalanced by the rise in visual uncertainty σ_v, since sensory uncertainty of various

140   visual stimuli increases linearly with eccentricity (Klein & Levi, 1987; Levi et al., 1987). As

141   participants are instructed to fixate on the target, an increase in θ lead to increased

142   eccentricity. Hence, we model this linear increase in visual uncertainty by

$$\sigma_v = a + b\theta, \tag{4}$$

where *a* and *b* are free parameters. We conducted simulations of implicit adaptation with

varying error clamp size (θ). The model simulation closely resembles the saturated

adaptation in three independent experiments (Kim et al., 2018; Morehead et al., 2017). In

fact, our PEA model predicts a concave adaptation pattern, contrasting with the ramp

pattern suggested by the PReMo model (Tsay, Kim, et al., 2022). In Experiment 1, we

aim to validate the assumption of a linear increase in visual uncertainty (Eq. 1); in

Experiment 2, we seek to verify whether implicit adaptation adheres to a concave pattern

as prescribed by the PEA model. Subsequent experiments, namely Experiments 3 and 4,

will test the model's additional novel predictions concerning changes in proprioception

and the impact of experimentally manipulated visual uncertainty on adaptation,

respectively.

156   ---- insert Figure 1 here ----

Experiment 1: Visual Uncertainty Increases Linearly with Perturbation Size

To quantify visual uncertainty in a standard error-clamp adaptation setting, we employed

psychometric methods. Occluded from seeing their actual hand, participants (n=18) made

repetitive reaches to a target presented 10 cm straight head while an error-clamped

cursor moving concurrently with one of three perturbation sizes (i.e., 4°, 16° and 64°),

randomized trial-by-trial. In alignment with the error-clamp paradigm, participants were

instructed to fixate on the target and to ignore the rotated cursor feedback. Eye-tracking

confirmed compliance with these instructions (Figure S1). Perturbation directions were

counter-balanced across trials, with equal probability of clockwise (CW) and

counterclockwise (CCW) rotation. Post-movement, participants were required to judge

the cursor's rotation direction (CW or CCW) relative to a briefly displayed reference point

(Figure 2A & Figure 6A). Employing this two-alternative forced-choice (2AFC) task and

the Parameter Estimation by Sequential Testing (PEST) procedure (Lieberman &

Pentland, 1982), we derived psychometric functions for visual discrimination (Figures 6

and Figure S2). Our findings reveal a significant increase in visual uncertainty (σ_v) with

perturbation size, for both CW and CCW rotations (Friedman test, CW direction: $\chi^2(2) =$

34.11, p = 4e-8; CCW: $\chi^2(2)$ = 26.47, p = 2e-6). Given the symmetry for the two directions,

we collapsed data from both directions, and confirmed the linear relationship between σ_v

and θ by a generalized linear model: $\sigma_v = a + b\theta$, with a = 1.853 and b = 0.309, R^2 =

0.255 (F = 51.6, p = 2.53e-9). The 95% confidence intervals (CI) for a and b are [0.440,

3.266] and [0.182, 0.435], respectively. The intercept was similar to the visual uncertainty

estimated in a previous study (Tsay, Avraham, et al., 2021). The linear dependency

indicates a striking seven-fold increase in visual uncertainty from a 4° perturbation to a

64° perturbation (22.641 ± 6.024° vs. 3.172 ± 0.453°).

183   ---- insert Figure 2 here ----

Experiment 2: Visual Uncertainty Modulated Perceptual Error Accounts for

Overcompensation and Saturation in Implicit Adaptation

The critical test of the PEA model lies in its ability to employ the linear function of visual

uncertainty obtained from Experiment 1 to precisely explain key features of implicit

adaptation. Earlier research mostly scrutinized smaller perturbation angles when

reporting saturation effects (Bond & Taylor, 2015; Kim et al., 2018; Morehead et al., 2017).

In contrast, Experiment 2 involved seven participant groups (n = 84) to characterize

implicit adaptation across an extensive range of perturbation sizes (i.e., 2°, 4°, 8°, 16°,

32°, 64°, and 95°). After 30 baseline training cycles without perturbations, each group

underwent 80 cycles of error-clamped reaching and 10 washout cycles without visual

feedback (Figure 3A). We replicated key features of implicit adaptation: it incrementally

reached a plateau, and then declined during washout. Small perturbations led to

overcompensation beyond visual errors (for 2°, 4°, 8°, 16° clamp sizes). Across

perturbation sizes, the faster the early adaptation the larger the adaptation extent (Figure

S4). Critically, the adaptation extent displayed a concave pattern: increasing steeply for

smaller perturbations and tapering off for larger ones (Figure 3B). A one-way ANOVA

revealed a significant group difference in adaptation extent ($F(6,83) = 12.108$, $p = 1.543e$-

09). Planned contrasts indicated that 8°, 16°, and 32° perturbations did not differ from

each other (all $p>0.417$, with Tukey-Kramer correction), consistent with earlier evidence

of invariant implicit adaptation (Kim et al., 2018). However, 64° and 95° perturbations led

to significantly reduced adaptation extents compared to 8° ($p = 3.194e-05$ and $5.509e-06$,

respectively), supporting the concave pattern as a more accurate portrayal of implicit

adaptation across varying perturbation size.

Importantly, the PEA model, when augmented with visual uncertainty data from

Experiment 1, precisely predicts this size-dependent adaptation behavior (Figure 3B).

Beyond adaptation extent, the model also accurately predicts the trial-by-trial adaptation

across all seven participant groups, employing a single parameter set ($R^2 = 0.975$; Figure

3A). The model had only four free parameters ($A = 0.974$, $B = 0.208$, $\sigma_p = 11.119°$, $\sigma_u =$

$5.048°$; Table S1). Remarkably, both the retention rate A and learning rate B are

consistent with previous studies focusing on visuomotor rotation adaptation (Albert et al.,

2022). We also quantified proprioceptive uncertainty (σ_p) in a subset of participants (n=13)

using a similar 2AFC procedure as in Experiment 1. We found that σ_p was $9.737°±5.598°$

(Figure S6), which did not statistically differ from the σ_p value obtained from the model

fitting (two-tailed t-test, p = 0.391). In summary, the perceptual parameters obtained in

Experiment 1, when incorporated into the PEA model, effectively explain the implicit

adaptation behaviors observed in different participant groups in Experiment 2.

---- insert Figure 3 here ----

In comparative analysis, the PReMo model yields a substantially lower R^2 value of 0.749

(Figure S3B). It tends to underestimate adaptation for medium-size perturbations and

overestimate it for large ones (Figure 3C; see also Figure S3B for trial-by-trial fitting).

Another alternative is the causal inference model, previously shown to account for

nonlinearity in motor learning (Mikulasch et al., 2022; Wei & Körding, 2009). Although this

model has been suggested for implicit adaptation (Tsay, Avraham, et al., 2021), it fails to

reproduce the observed concave adaptation pattern (Figures S3C and 3D). The model

aligns well with adaptations to medium-size perturbations (8°, 16°, and 32°) but falls short

for small and large ones, yielding an R^2 value of 0.711 (see Figure S3C for trial-by-trial

fits). Model comparison metrics strongly favor the PEA model over both the PReMo and

causal inference models, as evidenced by AIC scores of 2255, 3543, and 3283 for the

PEA, PReMo, and causal inference models, respectively (Table S2). In summary, it is the

eccentricity-induced visual uncertainty that most accurately accounts for the implicit

adaptation profile across a broad spectrum of perturbation sizes, rather than saturated

visual influence or causal inference of error.

Experiment 3: Cue Combination Accounts for Changes in Proprioception During Implicit

Adaptation

Motor adaptation not only recalibrates the motor system but also alters proprioception

(Rossi et al., 2021) and even vision (Simani et al., 2007). In traditional motor adaptation

involving both explicit and implicit components, the perceived hand location is initially

biased towards the visual perturbation and subsequently stabilizes (Ruttle et al., 2016).

However, in implicit adaptation, the perceived hand location initially aligns with but later

drifts away from the visual feedback (Tsay et al., 2020). The PReMo model proposes that

this drift comprises two phases: initial proprioceptive recalibration and subsequent visual

recalibration (Tsay, Kim, et al., 2022), however, this assumption is lack of empirical

validation. In contrast, we suggest that the perceived hand location is based on the same

Bayesian cue combination principle. In this framework, the perceived hand location at the

end of each reach is influenced by both the proprioceptive cue (x_p) and the estimated

hand position under the influence of clamped feedback $(\hat{x}_{Hand}$, Eq. 1).

During early adaptation, $\hat{x}_{Hand}$ is biased towards the clamped feedback, while x_p remains

near the target as the motor system has yet to adapt (Figure 4A). This results in an initial

negative proprioceptive bias. As adaptation progresses, although $\hat{x}_{Hand}$ remains biased,

x_p gradually shifts in the positive direction due to adaptation, resulting in an increasingly

positive proprioceptive bias. Remarkably, the PEA model can predict these temporal

changes in proprioception with high accuracy (R^2 = 0.982; Figure 4A).

If the hand estimate $\hat{x}_{Hand}$ indeed influences proprioceptive recalibration during

adaptation, our PEA model can make specific quantitative predictions about the

relationship between proprioception changes and visual perturbation size. While

traditional visuomotor paradigms suggest either invariant (Modchalingam et al., 2019) or

linear increases in proprioceptive recalibration with visual-proprioceptive discrepancy

(Salomonczyk et al., 2011), the PEA model prescribes a concave function in relation to

visual perturbation size (Figure 4B).

To empirically test this prediction, Experiment 3 (n=11) measured participants'

proprioceptive recalibration during implicit adaptation, using a procedure similar to the

error-clamp perturbations in Experiment 2. After each block of six adaptation trials,

participants' right hands were passively moved by a robotic manipulandum, and they

indicated the perceived direction of their right hand using a visually represented "dial"

controlled by their left hand (Figure 7B). This method quantifies proprioceptive

recalibration during adaptation (Cressman & Henriques, 2009). Each adaptation block

was followed by three such proprioception test trials. The alternating design between

adaptation and proprioception test blocks allowed us to assess proprioceptive biases

across varying perturbation sizes, which consisted of $\pm 10°$, $\pm 20°$, $\pm 40°$, and $\pm 80°$, to

covering a wide range (Figure 4D).

Our findings confirmed a typical proprioceptive recalibration effect, as the perceived hand

direction was biased towards the visual perturbation (Figure 4E). Importantly, the bias in

the initial proprioception test trial exhibited a concave function of perturbation size. A one-

way repeated-measures ANOVA revealed a significant effect of perturbation size

$(F(3,30)=3.603, p=0.036)$, with the 20° and 40° conditions displaying significantly greater

proprioceptive bias compared to the 80° condition (pairwise comparisons: 20° v.s. 80°, p

= 0.034; 40° v.s. 80°, p = 0.003). The bias was significantly negative for 20° and 40°

conditions (p = 0.005 and p = 0.007, respectively with one-tailed t-test), but not for 10°

and 80° condition (p = 0.083 and p = 0.742, respectively). This concave pattern aligns

well with the PEA model's predictions (Figure 4B), further consolidating its explanatory

power.

290   ---- insert Figure 4 here ----

This stands in contrast to the PReMo model, which assumes a saturation for the influence

of the visual cue on the hand estimate (Eq. 12-13). As a result, PReMo's predicted

proprioceptive bias follows a ramp function, deviating substantially from our empirical

findings (Figure 4C). The causal inference model, which mainly focuses on the role of

visual feedback in error correction, lacks the capability to directly predict changes in

proprioceptive recalibration.

Interestingly, we observed that the proprioceptive bias reduced to insignificance by the

third trial in each proprioception test block (one-tailed t-test, all $p > 0.18$; Figure 4E, yellow

line). This suggests that the influence from implicit adaptation – manifested here as trial-

by-trial updates of the perceived hand estimate $\hat{x}_{Hand}$ – decays rapidly over time.

Experiment 4: Differential Impact of Upregulated Visual Uncertainty on Implicit Adaptation

Across Perturbation Sizes

Thus far, we have presented both empirical and computational evidence underscoring

the pivotal role of perceptual error and visual uncertainty in implicit adaptation. It is crucial

to note, however, that this evidence is arguably correlational, arising from natural

variations in visual uncertainty as a function of perturbation size. To transition from

correlation to causation, Experiment 4 (n = 19) sought to directly manipulate visual

uncertainty by blurring the cursor, thereby offering causal support for the role of

multimodal perceptual error in implicit adaptation.

By increasing visual uncertainty via cursor blurring, we hypothesized a corresponding

decrease in adaptation across all perturbation sizes. Notably, the PEA model predicts a

size-dependent attenuation in adaptation: the reduction is less marked for smaller

perturbations and more pronounced for larger ones (Figure 5A). This prediction diverges

significantly from those of competing models. The PReMo model, operating under the

assumption of a saturation effect for large visual perturbations, predicts that cursor

blurring will only influence adaptation to smaller perturbations, leaving adaptation to larger

perturbations unaffected (Figure 5B). The causal inference model makes an even more

nuanced prediction: it anticipates that the blurring will lead to a substantial reduction in

adaptation for small perturbations, a diminishing effect for medium perturbations, and a

potential reversal for large perturbations (Figure 5C). This prediction results from the

model's core concept that causal attribution of the cursor to self-action—which directly

dictates the magnitude of adaptation—decreases for small perturbations but increases

for large ones when overall visual uncertainty is elevated.

---- insert Figure 5 here ----

Starting from the above predictions, Experiment 4 was designed to assess the impact of

elevated visual uncertainty across small (4°), medium (16°), and large (64°) perturbation

sizes. Visual uncertainty was augmented by superimposing a Gaussian blurring mask on

the cursor (Burge et al., 2008). Each participant performed reaching tasks with either a

standard or blurred clamped cursor for a single trial, bracketed by two null trials devoid of

cursor feedback (Figure 5D). These three-trial mini-blocks permitted the quantification of

one-trial learning as the directional difference of movements between the two null trials.

To preclude the cumulative effect of adaptation, perturbation sizes and directions were

randomized across mini-blocks.

Crucially, our findings corroborated the predictions of the PEA model: visual uncertainty

significantly diminished adaptation for medium and large perturbations (16° and 64°),

while leaving adaptation for small perturbations (4°) largely unaffected (Figure 5E). A two-

way repeated-measures ANOVA, with two levels of uncertainty and three levels of

perturbation size, revealed a significant main effect of increased visual uncertainty in

reducing implicit adaptation ($F(1,18) = 42.255$, $p = 4.112e-06$). Furthermore, this effect

interacted with perturbation size ($F(2,36) = 5.391$, $p = 0.012$). Post-hoc analyses

demonstrated that elevated visual uncertainty significantly attenuated adaptation for large

perturbations ($p = 2.877e-04$, $d = 0.804$ for 16°; $p = 1.810e-05$, $d = 1.442$ for 64°) but

exerted no such effect on small perturbations ($p = 0.108$, $d = 0.500$). These empirical

outcomes are not congruent with the predictions of either the PReMo or the causal

inference models (Figure 5B and 5C). This lends compelling empirical support to the

primacy of perceptual error in driving implicit adaptation, as posited by our PEA model.

351
352

Discussion

In this study, we elucidate the central role of perceptual error, derived from multimodal sensorimotor cue integration, in governing implicit motor adaptation. Utilizing the classical error-clamp paradigm, we uncover that the overcompensation observed in response to small perturbations arises from a sustained perceptual error related to hand localization, and the saturation effect commonly reported in implicit adaptation is not an intrinsic characteristic but is attributable to increasing sensory uncertainty with increasing visual perturbation eccentricity—a factor hitherto neglected in existing models of sensorimotor adaptation. Contrary to conventional theories that describe implicit adaptation as either saturated or invariant (Kim et al., 2018; Tsay, Kim, et al., 2022), our data reveal a concave dependency of implicit adaptation on visual perturbation size, characterized by diminishing adaptation in response to larger perturbations. Notably, our Perceptual Error Adaptation (PEA) model, calibrated using perceptual parameters from one set of participants, provides a robust account of implicit adaptation in separate groups subjected to varying perturbations. The model further successfully captures the perceptual consequences of implicit adaptation, such as the continuous shifts in proprioceptive localization during the adaptation process (Tsay et al., 2020) and its correlation with perturbation size. Lastly, we manipulated visual uncertainty independently of perturbation size and demonstrated that this selectively attenuated adaptation in the context of larger perturbations while leaving smaller perturbations unaffected. These empirical results, inconsistent with predictions from existing models, underscore the conceptual and quantitative superiority of our PEA model. In summary, our findings advocate for a revised

understanding of implicit motor adaptation, suggesting that it is governed by Bayesian

cue combination-based perceptual estimation of effector localization.

Bayesian cue combination has been established as a foundational principle in various

perceptual phenomena, both intra- and inter-modally (Seilheimer et al., 2014). It has also

been implicated in motor adaptation (Burge et al., 2008; He et al., 2016; Körding &

Wolpert, 2004; Wei & Körding, 2010). However, previous studies have largely focused on

experimentally manipulating sensory cue uncertainty to observe its effects on adaptation

(Burge et al., 2008; Wei & Körding, 2010), similar to our Experiment 4. What has been

largely overlooked is the natural covariance between visual uncertainty and perturbation

size, which, when incorporated into classical state-space models, provides a compelling

explanation for implicit adaptation.

The causal inference framework (Wei & Körding, 2009) fails to adequately predict

sensorimotor changes in implicit adaptation. For instance, it underestimates the

adaptation extent for large perturbations and incorrectly predicts that increasing visual

uncertainty would augment, rather than reduce, adaptation to large perturbations. We

postulate that causal inference is more relevant to motor learning dominated by explicit

processes, such as traditional visuomotor rotations, rather than in implicit adaptations

where cue combination is obligatory.

Similar to our PEA model, the PReMo model also incorporates the integration of multiple

sensory cues. But two models differ fundamentally in their conceptualization of how these

cues contribute to the error signal. The PReMo model posits two intermediate perceptual

variables with Bayesian cue integration: a visual estimate of the cursor and a

proprioceptive estimate of the hand (Tsay, Kim, et al., 2022). The final error signal in

PReMo is presumed to be a proprioceptive error, not from further Bayesian cue

combination, but from a visual-to-proprioceptive bias that is governed by a predetermined,

ramp-like visual influence that saturates around a 6–7° visual-proprioceptive discrepancy

(Eq. 13). These assumptions lack empirical validation. Our findings in Experiment 3

indicate that proprioceptive recalibration follows a concave function with respect to visual

perturbation size, contradicting the ramp-like function assumed by PReMo. Moreover, the

presupposed ramp-like visual influence generates a rigid prediction for a ramp-like

adaptation extent profile, which is at odds with the concave adaptation pattern we

observed in Experiment 2 and in a similar study involving trial-by-trial learning (Tsay,

Avraham, et al., 2021). Furthermore, PReMo predicts that increasing visual uncertainty

will selectively reduce adaptation to small perturbations while sparing large ones. This is

inconsistent with our findings in Experiment 4, which demonstrated that increased visual

uncertainty substantially impacted adaptation more to larger perturbations than to small

ones. Lastly, PReMo's reliance on a proprioceptive bias constrains its ability to account

for the temporal shifts in perceived hand location during adaptation (Tsay et al., 2020). In

contrast to PEA's unified approach, PReMo must resort to separate mechanisms of

proprioceptive and visual recalibration at different phases of adaptation to explain these

shifts. In summary, the PReMo model's assumptions introduce limitations that make it

less consistent with empirical observations, particularly concerning the nonlinearities

observed in both motoric and perceptual aspects of implicit adaptation.

Our research contributes to an ongoing debate concerning the driving forces behind error-

based motor learning, specifically addressing the question of whether implicit adaptation

is driven by target error or sensory prediction error (Albert et al., 2022; Izawa & Shadmehr,

2011; Leow et al., 2020; Mazzoni & Krakauer, 2006; McDougle et al., 2015; Miyamoto et

al., 2020; Taylor & Ivry, 2011; Tseng et al., 2007). Most empirical data fueling this debate

stem from traditional motor adaptation paradigms where explicit and implicit learning co-

occur and interact. In these paradigms—visuomotor rotation being a prime example—

target error is defined as the disparity between the target and the perturbed cursor, while

sensory prediction error is the disparity between the predicted and actual cursor. Both

types of error are sensory (specifically, visual) in nature, yet they differ due to the

misalignment between the predicted or desired cursor direction and the target direction,

which is induced by explicit learning (Taylor et al., 2014).

By employing the error-clamp paradigm, our study was able to isolate implicit learning,

thereby eliminating potential confounds from explicit learning. Interestingly, in this

paradigm, the target error and sensory prediction error effectively refer to the same visual

discrepancy, as both the predicted and target directions are aligned. Despite this,

classical state-space models, which utilize this visual error, fail to account for the nuanced

features of implicit adaptation (Tsay, Kim, et al., 2022). In contrast, our PEA model

reframes the perturbing cursor as a visual cue influencing the perceptual estimation of

hand location, rather than as a source of visual error. The resultant bias in hand estimation

from the desired target serves as the actual error signal. This leads us to posit that the

error signal driving implicit sensorimotor adaptation is fundamentally perceptual, rather

than sensory. From a normative standpoint, this perceptual error could be construed

either as a predictive or performance error (Albert et al., 2022), but importantly, it is not

tied to a specific modality (i.e., vision or proprioception). Instead, it directly pertains to the

perceptual estimate that is crucial for task execution, i.e., bringing the hand to the target.

The concept of perceptual error-driven learning can be extrapolated to various motor

adaptation paradigms, including those involving explicit learning. For instance, in

visuomotor rotation tasks, explicit learning manifests as a deviation in the aiming direction

from the visual target, whereas implicit learning manifests as a further deviation the actual

hand position from this aiming direction (Taylor et al., 2014). Even in the presence of

explicit learning, the perturbed cursor continues to bias the perceptual estimate of the

hand, thereby potentially driving implicit adaptation. In this scenario, the perceptual error

is defined as the difference between the perceptual estimate of the hand and the altered

aiming direction, which serves as the new "target" when explicit learning is in play. Our

PEA model would predict similar saturation effects in implicit adaptation for this

conventional adaptation paradigm, similar to for the error-clamp paradigm. Indeed,

evidence from the conventional adaptation paradigm suggests that its implicit adaptation

follows either a saturation effect (Bond & Taylor, 2015; Neville & Cressman, 2018) or a

concave pattern (Tsay, Haith, et al., 2022) across a range of perturbation sizes.

Furthermore, according to the PEA framework, this perceptual error is anchored on the

aiming target, thereby naturally predicting that implicit and explicit adaptations should

interact in a complementary manner, a notion that aligns with recent theories on their

interaction (Albert et al., 2022; Miyamoto et al., 2020). Future research is warranted to

further investigate the role of perceptual error in driving implicit learning across diverse

motor learning paradigms.

Our study provides a new angle on explaining proprioceptive changes during motor

adaptation, advocating for a Bayesian cue combination framework. Previously, the

change in proprioceptive hand localization during motor adaptation has been ascribed to

visual-proprioceptive discrepancy-induced recalibration (Ruttle et al., 2018; Salomonczyk

et al., 2013) and/or altered sensory prediction driven by the adapted forward internal

model (Mostafa et al., 2019; 't Hart & Henriques, 2016). To dissect these components,

researchers have often compared proprioceptive localization in actively moved (Tsay et

al., 2020) versus passively placed (passive localization, e.g., Experiment 3) hands during

adaptation, attributing the smaller bias in passive localization to recalibration alone. The

difference between the two is then considered to reflect altered sensory prediction due to

motor adaptation (Mostafa et al., 2019; Rossi et al., 2021). But these conceptual divisions

lack computational models for validation. For instance, researchers have proposed that

proprioceptive recalibration in visuomotor adaptation is either a fixed proportion (e.g.,

20%) of the visual-proprioceptive discrepancy (Henriques & Cressman, 2012; Ruttle et

al., 2021) or largely invariant (Modchalingam et al., 2019). In fact, cross-sensory

calibration typically follows the Bayesian principle, as shown in other task paradigms other

than motor adaptation (Stetson et al., 2006; Wozny & Shams, 2011). Our Experiment 3

shows that proprioceptive recalibration exhibits a concave, instead of invariant or

proportional, dependency to visual perturbation size, a finding follows the Bayesian

principles of cue combination. Our results also confirm that the critical cue for passive

localization is the biased perceived hand position ($\hat{x}_{Hand}$) fueled by adaptation.

The same Bayesian framework applies to active localization, though this time $\hat{x}_{Hand}$ is to

be combined with the proprioceptive cue from the adapted hand. In this sense, active

localization indeed serves as a multifaceted reflection of both the internal model and

proprioceptive recalibration (Mostafa et al., 2019; Rossi et al., 2021). Specifically, the

proprioceptive cue continuously drifts by the adapted internal model, while the perceived

hand position encapsulates the effects of proprioceptive recalibration. During the initial

stages of perturbation, the immediate negative bias in active localization is predominantly

attributable to rapid proprioceptive recalibration. This is evidenced by a sudden shift in

the estimated hand position ($\hat{x}_{Hand}$; Figure 4A), occurring before the internal model has

had sufficient time to adapt.

Then, why does active localization in traditional motor adaptation paradigms yield a

largely stable bias (Ruttle et al., 2016, 2021)? We postulate that the rapid explicit learning

leads to a quick asymptotic adaptation, while previous investigations have predominantly

measured active localization after adaptation has plateaued (Henriques & Cressman,

2012; Modchalingam et al., 2019; Mostafa et al., 2019; Salomonczyk et al., 2011, 2013;

Tsay, Kim, et al., 2021). Consequently, these studies may overlook the evolving effect of

the adaptation. In contrast, the gradual nature of implicit adaptation provides a unique

opportunity to uncover the underlying mechanisms governing changes in proprioception

during the adaptation process.

Notably, our model aligns with previous findings that show a positive correlation between

proprioceptive recalibration and motor adaptation based on individual differences (Ruttle

et al., 2021; Salomonczyk et al., 2013; Tsay, Kim, et al., 2021). Unlike existing theories

that posit proprioceptive recalibration either as a component of (Modchalingam et al.,

2019; Mostafa et al., 2019; Ruttle et al., 2021) or a driver for implicit adaptation (Tsay,

Kim, et al., 2022), our PEA model provides a mechanistic and empirically testable

framework. It posits that the misestimation of hand position ($\hat{x}_{Hand}$) —induced by the

recent perturbation—serves as the driving factor for both implicit adaptation and changes

in proprioception. This misestimation is perturbation-dependent, resulting in both implicit

adaptation and proprioceptive recalibration exhibiting a concave profile relative to

perturbation size. Updated on a trial-by-trial basis, this misestimation exerts immediate

effects, manifesting as an abrupt negative bias (Figure 4A). Additionally, its influence

decays rapidly, becoming negligible within three trials (Figure 6C). These converging lines

of evidence strongly suggest that perceptual misestimation of hand position is central to

the process of proprioceptive recalibration during adaptation.

Our findings contribute nuanced perspectives to the modulation of implicit learning rate

by factors beyond visual perturbation size. Previous studies have shown that

environmental inconsistency -- defined as the inconsistency of visual errors -- reduced

the rate (Herzfeld et al., 2014; Hutter & Taylor, 2018) or asymptote (Albert et al., 2021) of

implicit adaptation. Baseline motor variance in unperturbed conditions has been shown

to increase implicit adaptation rate, proposed as a sign of better exploratory learning (Wu

et al., 2014). These studies interpret such phenomena as parametric changes in the

learning rate in relation to visual errors, conceptualized as alterations to the B parameter

in existing models. However, apparent change in learning rate to visual errors does not

necessarily signify parametric modification, but may attribute to other factors that

influence the use of visual cues (He et al., 2016), such as visual uncertainty in our case.

Previous research has also pointed to alternative factors like error discounting based on

causal inference of error (Wei & Körding, 2009), proprioceptive uncertainty (Ruttle et al.,

2021; Tsay, Kim, et al., 2021), and state estimation uncertainty (He et al., 2016; Wei &

Körding, 2010). Our work suggests a shift in perspective: the driving error signal for

implicit learning should be considered as perceptual, rather than merely visual. This

paradigmatic shift could serve as a cornerstone for future research aimed at

understanding how learning rates adapt under varying conditions.

Our new framework opens avenues for exploring the memory characteristics of implicit

learning. Traditional motor adaptation often exhibits 'savings,' or accelerated relearning

upon re-exposure to a perturbation (Della-Maggiore & McIntosh, 2005; Huberdeau et al.,

2019; Krakauer et al., 2005; Landi et al., 2011). In contrast, implicit adaptation has been

found to exhibit a decreased learning rate during re-adaptation (Avraham et al., 2021), a

phenomenon attributed to conditioning (Avraham et al., 2021) or associative learning

mechanisms (Avraham et al., 2022). Investigating this 'anti-saving' effect will yield insights

into the unique memory properties of implicit learning. Although our current PEA model

is structured around single-epoch learning and does not directly address this question, it

does raise new, testable hypotheses. For example, is the reduced adaptation rate during

relearning attributable to a down-weighting of perturbed visual feedback in cue

combination, or does it reflect a parametric alteration in the learning rate? Another

noteworthy aspect of implicit learning is its remarkably slow decay rate. It has been

observed that the number of trials required to washout the implicit adaptation exceeds the

number of trials needed to establish it (Avraham et al., 2021; Tsay et al., 2020). In the

context of our perceptual error framework, this raises the possibility that washout phases

might be governed by state updating involving a distinct set of sensorimotor cues or an

alternative updating mechanism, such as memory formation and selection (Oh &

Schweighofer, 2019).

Methods

Participants

We recruited 115 college students from Peking University (77 females, 38 males, 22.05

$\pm$ 2.82 years, mean $\pm$ SD). Participants were all right-handed according to the Edinburgh

handedness inventory (Oldfield, 1971) and had normal or corrected-to-normal vision.

Participants were naïve to the purpose of the experiment and provided written informed

consent, which was approved by the Institutional Review Board of the School of

Psychological and Cognitive Sciences, Peking University. Participants received monetary

compensation upon completion of the experiment.

Apparatus

In Experiment 1, 2 and 4, participants were seated in front of a vertically-placed LCD

screen (29.6 x 52.7 cm, Dell, Round Rock, TX, US). They performed the movement task

with their right hand, holding a stylus and slide it on a horizontally placed digitizing tablet

(48.8 x 30.5 cm, Intuos 4 PTK-1240, Wacom, Saitama, Japan). In Experiment 1, a

keyboard was provided to the participants' left hand to enable them to report the direction

of visual stimuli in the discrimination task. A customized wooden shelter was placed

above the tablet to block the peripheral vision of the right arm. In Experiment 1 and 4,

participants placed their chin on a chin rest attached on the wooden shelter to stabilize

their head. Their eye movement was recorded by an eye tracker (Tobii pro nano, Tobii,

Danderyd Municipality, Sweden) affixed at the lower edge of the screen. The sampling

rate was 160-200 Hz for the tablet and 60 Hz for the eye tracker.

Experiment 3 was conducted using the KINARM planar robotic manipulandum with a

virtual-reality system (BKIN Technologies Ltd., Kingston, Canada). Participants seated in

a chair and held the robot handles with their left and right hands (Figure 7). The movement

task was performed with the right handle and the left handle was used to indicate the

perceived direction of right hand in the proprioception test. A semi-silvered mirror was

placed below the eye level to block the vision of the hands and the robotic manipulandum;

it also served as a display monitor.

Experiment 1: measuring visual uncertainty in error-clamp adaptation

Eighteen among twenty participants finished the reaching with clamped error feedback

and visual discrimination task in 3 consecutive days, two participants withdrew during the

experiment. Participants made reaching movement by sliding the stylus from a start

position at the center of the workspace to towards a target (Figure 6A). The start position,

the target, and the cursor were represented by a gray dot, a blue cross and a white dot

on the screen, respectively. All these elements had a diameter of 5mm. The procedure of

the motor and visual discrimination task is illustrated in Figure 2A. To initiate a trial,

participates moved the cursor into the start position. Following an 800ms holding period,

a target appeared 10 cm away in twelve o'clock direction and participants were instructed

to slide through the target rapidly while maintaining a straight hand trajectory. The trial

terminated when the distance between the hand and the start position exceeded 10 cm,

regardless of whether the target was hit. A warning message, "too slow", would appear

on the screen if participants failed to complete the trial within 300 ms after initiating the

movement. Each practice day began with 60 standard reaching trials, during which

veridical feedback about hand location was provided by the cursor. The target would

change from blue to green if the cursor successfully passed through it. In subsequent

visual clamp trials, the cursor moved along a predetermined direction set by the

perturbation angle, while its position was updated in real-time based on the hand's

location. The cursor's distance from the start position was equal to the distance between

the hand and the start position until the end of the trial.

Following each trial, the cursor remained frozen at its final position for an additional 800

ms before disappearing. The visual discrimination task commenced 1000 ms thereafter.

A yellow reference point, located 10 cm from the start position, was displayed for 150 ms

near the cursor's final position (Figure 2A & Figure 6A). Subsequently, all visual stimuli,

except for the blue cross at the start position, were removed from the screen. Participants

were then required to judge whether the reference point was situated in a clockwise (CW)

or counterclockwise (CCW) direction relative to the cursor's final position and to report

their judgment by pressing a key on the keyboard. Participants were informed that they

no longer controlled the direction of cursor movement during the task. They were

instructed to fixate their gaze on either the start position or the blue cross during the motor

task, while actively ignoring the white cursor. During the discrimination task, they were

required to maintain their gaze on the blue cross. Eye movements were monitored in real-

time using an eye tracker. Participants received a warning if their gaze was detected

outside a 75-pixel-wide band-shaped region centered on the line of gaze four consecutive

times during the experiment (Figure S1).

622   ---- insert Figure 6 here ----

In each trial, the angular deviation between the error-clamped cursor and the reference

point was determined using a PEST procedure (Lieberman & Pentland, 1982). Figure 6C-

D illustrates the evolution of the deviation angle and step size for an exemplary participant

experiencing a -16° perturbation. In each round, the deviation commenced at 30°

(indicated by yellow points in Figure 6C-D) and was altered by one step size following

each trial. The initial step size was set at 10° and was halved whenever the direction

judgment changed (i.e., from "CW" to "CCW" or vice versa). For a specific perturbation

angle, the initial deviation always started from the CW direction for the first round and

flipped the direction at the beginning of the next round. A round terminated either when

the step size fell below a predefined criterion (indicated by the red line in Figure 6D) or

when the trial count exceeded 30. Six perturbation angles were randomly interleaved

(Figure 6B), and the experiment concluded when four complete rounds of the PEST

procedure had been completed for each perturbation angle. Consequently, the total

number of trials varied among participants and across practice days. Additionally, for

some perturbation angles, more than four complete rounds could be conducted in a single

day.

Experiment 2: Motor adaptation with different perturbation size

Eighty-four participants were randomly allocated into seven groups, each comprising 12

individuals. Each group performed a motor adaptation task featuring clamped visual

feedback at different perturbation angles: 2°, 4°, 8°, 16°, 32°, 64°, and 95°. As in

Experiment 1, participants were instructed to slide rapidly and directly through the target,

which was represented by a blue dot rather than a cross. In each trial, the target appeared

at one of four possible locations (45°, 135°, 225° or 315° counter-clockwise from the

positive x-axis). The sequence of target locations was randomized, yet constrained so

that all four positions appeared in cycles of four trials. Each group commenced with a

baseline session that included 15 cycles of reaching trials with veridical feedback,

followed by 15 cycles without visual feedback. Subsequently, during the perturbation

session, participants completed 80 cycles of training trials featuring the error-clamped

cursor with one perturbation angle (i.e., clamp size), depending on their group assignment.

To assess the aftereffect, a session comprising 10 cycles of movement without visual

feedback was administered.

Experiment 3: Proprioception test with different perturbation sizes

Eleven participants were recruited for testing their proprioception recalibration. This

experiment incorporated two types of trials: reaching trials and proprioception test trials.

During the reaching trials, participants were instructed to aim for a target, which could

appear at one of three possible locations (25°, 45°, or 65° counter-clockwise from the

positive x-axis, as represented by light blue dots in Figure 4C, right panel). The task was

similar to those in Experiments 1 and 2, with the key difference being that participants

performed the task using KINARM robots (as depicted in Figure 7A). The dimensions and

relative distances of the visual stimuli remained consistent with those used in Experiments

1 and 2. As in previous experiments, three kinds of visual feedback were provided during

different sessions: no visual feedback, veridical feedback, and feedback featuring an

error-clamped cursor.

In the proprioception test, participants were instructed to hold the robot's right handle and

wait for passive movement by the robot to one of six proprioception targets (small red

dots in Figure 4C, right panel). These targets were spaced at 10° intervals, ranging from

20° to 70° counter-clockwise from the positive x-axis, and flanked the three reaching

targets. The passive movement lasted for 1,000 ms and followed a straight-line path at a

speed consistent with a minimum jerk velocity profile. During this movement, a ring with

a 10 cm radius, centered at the start position, was displayed on the screen (depicted as

a red arc in Figure 7B). The cursor was also replaced by a ring, its radius expanding as

the hand moved toward the proprioception target.

After the right hand reached the proprioception target, participants were instructed to

maintain their right hand's position. Using the left handle, they were then asked to indicate

the perceived location of their right hand. The position of the left handle was mapped to

the rotation of a "dial," which was constrained to the target arc.

681   ---- insert Figure 7 here ----

The position of h_p was displayed on the target arc as a small red rectangle (a visual "dial,"

as shown in Figure 7B). Participants were instructed to indicate the location of their right

hand by moving the red rectangle to the position they perceived as accurate. The final

position of h_p was recorded when its angular velocity remained below 1 degree/second

for a duration exceeding 1000 ms. The proprioceptive bias was then calculated as the

angular deviation between the actual hand position (h_R) and the perceived hand position

(h_p).

Reaching trials and proprioception test trials were organized into blocks (Figure 4D). Each

reaching block consisted of 6 trials, targeting 3 different locations with 2 repetitions each.

Each reaching block was followed by a proprioception test block consisting of 3 trials. In

these test trials, the robot moved the participant's right hand toward a target position near

one of the three reaching targets. These test targets were randomly chosen from six

possible locations (Figure 4C, right panel). The entire experiment comprised 40 reaching

blocks and 40 subsequent proprioception test blocks. The first four reaching blocks

provided veridical cursor feedback, the next four offered no cursor feedback, and the

remaining 32 featured one of eight possible perturbation sizes ($\pm10°$, $\pm20°$, $\pm40°$, and

$\pm80°$). The size of the perturbation was randomized between blocks.

Experiment 4: upregulating visual uncertainty affects implicit adaptation

Nineteen participants from Experiment 1 completed Experiment 4. The reaching task

employed the same setup as in Experiment 1. However, instead of performing perceptual

judgments of cursor motion direction, participants engaged in movements with one of

three types of cursor feedback: veridical feedback, no feedback, and feedback with

clamped perturbation. To assess the influence of visual uncertainty on implicit learning,

we modified the cursor to appear blurred in half of the clamped trials. The blurring mask

had a diameter of 6.8 mm, and the color intensity decreased from the cursor's center

following a two-dimensional Gaussian distribution with $\sigma_x = \sigma_y = 1.4$ mm. As depicted in

Figure 5D, participants underwent the same procedures across three consecutive days.

Each day consisted of 60 baseline trials, followed by 15 training blocks designed to

assess single-trial learning. Within each training block, 12 trials featured an error-clamped

cursor, each flanked by a trial without feedback. The difference between two adjacent no-

feedback trials served as a measure of single-trial learning at specific perturbation sizes.

Each of the 12 perturbation trials was randomly assigned one of 12 possible perturbations,

comprising two cursor presentations (blurred or clear) and six clamp sizes ($\pm 4°$, $\pm 16°$,

$\pm 64°$).

Data analysis

Processing of kinematic data

In Experiments 1, 2, and 4, hand kinematic data were collected online at a sampling rate

ranging between 160 and 200 Hz and subsequently resampled offline to 125 Hz. The

movement direction of the hand was determined by the vector connecting the start

position to the hand position at the point where it crossed 50% of the target distance, i.e.,

5 cm from the start position.

In Experiment 3, hand positions and velocities were directly acquired from the KINARM

robot at a fixed sampling rate of 1 kHz. The raw kinematic data were smoothed using a

fifth-order Savitzky-Golay filter with a window length of 50 ms. Owing to the high temporal

resolution and reliable velocity profiles provided by the KINARM system, the heading

direction in Experiment 3 was calculated as the vector connecting the start position to the

hand position at the point of peak velocity.

Psychometric curve

731 For the visual discrimination task, data of all three days were pooled together, the

732 probability of responding that "the reference point was in the counter-clockwise direction

733 of the cursor" was calculate as p for all angle differences (Figure S2). At each perturbation

734 size, a logistic function was used to fit the probability distribution for individual participants:

$$p = 1/(1 + e^{-k(x-x_0)}), \tag{5}$$

736   where k is the slope and x_0 is the origin of the logistic function. The visual uncertainty was

737   defined as the angle differences between 25% and 75% of the logistic function:

738    $$\sigma_v = \frac{\log(p_2/(1-p_2)) - \log(p_1/(1-p_1))}{k},$$ (6)

739   where $p_1 = 25\%$ and $p_2 = 75\%$.

Statistical analysis

In Experiment 1, since the visual uncertainty σ_v follows a non-negative skewed

distribution among participants, it violated the assumption of the ANOVA test. We thus

applied Friedman's nonparametric test to determine whether σ_v changes with the

perturbation angle θ. Specifically, σ_v for both positive and negative θ were subjected to

Friedman's test separately, with θ serving as the factor. Given the symmetry between

positive and negative θ, we pool the data to quantify the linear dependency of σ_v on the

absolute θ (Eq. 4). Because σ_v is expected to be always positive, we assume that it is

generated from a gamma distribution rather than a normal distribution. Thus, the data

was fitted by a generalized linear regression model with the absolute value of θ as

independent variable and σ_v as dependent variable.

In Experiment 2, the adaptation extent was defined as the mean hand angles in the last

10 cycles in the perturbation phase (cycle 101-110). A one-way ANOVA with perturbation

size serving as the factor to examine its influence on the adaptation extent. Pairwise post-

hoc comparisons were conducted using Tukey-Kramer correction.

In Experiment 3, proprioceptive recalibration was quantified as the angular difference

between the perceived and actual hand directions. A one-way repeated-measures

ANOVA was conducted on the data of first trial, using perturbation size as the within-

subject factor. Greenhouse-Geisser corrections were applied when the assumption of

sphericity was violated (Kirk, 1968). Multiple pairwise comparisons were conducted

among different perturbation sizes for the first proprioception test. To determine if the

proprioceptive biases were significantly different from zero, one-tailed (left) *t*-tests were

conducted separately for the first and third proprioception test trials at each perturbation

size.

In Experiment 4, the single-trial learning data was subjected to a 2 (visual uncertainty) x

3 (perturbation size) repeated-measures ANOVA. Greenhouse-Geisser corrections were

applied as above, and the simple main effect of visual uncertainty was tested for each of

the three perturbation sizes.

Model fitting and simulations

Perceptual Error Adaptation (PEA) model

Model fitting for adaptation extent as a function of perturbation size. To fit the adaptation

extent data from three different experiments in previous studies in (Kim et al., 2018;

Morehead et al., 2017), Eq. 3 and Eq. 4 were modified for simplification. To avoid

overfitting of the small dataset, we reduced the number of model parameters by assuming

that $\hat{x}_{Hand}$ asymptote to the target direction in the final adaptation trials that are used for

computing adaptation extent, thus the retention rate $A \equiv 1$. Insert Eq. 4 to Eq. 3, the

asymptote hand angle with different perturbation size is:

777
$$x_p^{asym} = -(\frac{\sigma_p/a}{1+(b/a)\theta})^2\theta. \tag{7}$$

Two ratio parameters $R_{1,ext} = \sigma_p/a$ and $R_{2,ext} = b/a$ were used in data fitting. Three

datasets were fitted separately.

Model fitting for trial-by-trial adaptation and proprioception changes. The trial-by-trial

changes of adaptation (Figure 3A) and of proprioceptive localization (Figure 4A) was fitted

with Eq. 1, Eq. 2, and Eq. 4 based on the mean performance of all participants. The PEA

model only had four free parameters, $\Theta = [\sigma_u, \sigma_p, A, B]$. The slope *a* and intercept *b* in Eq.

1 were obtained by psychometric tests from Experiment 1 (see statistical analysis). The

reported hand position (x_{report}, blue dots in Figure 4A) was based on the proprioceptive

cue x_p and the estimated hand $\hat{x}_{Hand}$ from the reaching trial. With the Bayesian cue

combination assumption, the reported hand position was biased by x_p with a ratio

determined by the variance of x_p and $\hat{x}_{Hand}$:

789
$$x_{report} = \hat{x}_{Hand} + \frac{\sigma_{Hand}^2}{\sigma_{Hand}^2 + \sigma_p^2}(x_p - \hat{x}_{Hand}), \tag{8}$$

where σ_{Hand}^2 and σ_p^2 are the variance of $\hat{x}_{Hand}$ and x_p respectively. To verify if the slope *b*

and intercept *a* obtained from Experiment 1 are consistent across experiments, they were

also estimated by fitting data from Experiment 2 (Figure 3). In this case, the model fitting

was performed with 6 free parameters, $\Theta = [\sigma_u, \sigma_p, a, b, A, B]$. The fitted values of *a* and *b*

are fallen into the 95% CI of estimated parameters in Experiment 1 (purple line in Figure

2C, see details in Table S1).

The dependence of proprioceptive recalibration on perturbation size (Figure 4B) were

simulated by the PEA model with the parameter values estimated from Experiment 2. We

assumed that the proprioceptive bias results from the influence of a biased hand estimate

($\hat{x}_{Hand}$) during adaptation and the influence is quantified as a percentage of its deviation

from the true hand location:

801
$$x_{bias} = -(0 - \hat{x}_{Hand})R_p, \tag{9}$$

where the actual hand location is 0, R_p is the percentage of influence, and $\hat{x}_{Hand}$ is

determined by Eq.1. In simulation, R_p varied from 0.05 to 0.8 to estimate the overall

dependence of proprioceptive recalibration on perturbation size.

Model fitting and simulation for single-trial learning. In the single-trial learning paradigm

(Figure S5), the average movement direction across trials aligns with the target direction

since the visual perturbations are evenly distributed in both directions. Thus, the sensory

cue x_u and x_p have the same mean. For modeling single-trial learning, instead of having

two separate cues, we assume a combined cue of x_u and x_p to follow $x_{int} \sim N(T, \sigma_{int}^2)$,

where T is the target direction, $\sigma_{int}^2 = \frac{\sigma_u^2 \sigma_p^2}{\sigma_u^2 + \sigma_p^2}$ represents the variance of integrated sensory

signal of x_u and x_p. Single-trial learning was quantified as the difference between the two

null trials before and after the perturbation trial. As the perturbation size in the triplet of

trials varied randomly, we assume that the effects of different perturbations are

independent. Thus, single-trial learning was modeled as learning from the current

perturbation without history effect. It follows the equations modified from Eq. 1 and 2:

$$x_{STL} = B(T - \hat{x}_{Hand}) \tag{10}$$

$$\hat{x}_{Hand} = W_{int}T + W_v x_v, \text{ with } W_{int} = \frac{1/\sigma_{int}^2}{\sum_j 1/\sigma_j^2}, \ i, j = int, v, \tag{11}$$

where x_v is the visual perturbation, W_{int} and W_v are the weights of the cues, σ_v is the

standard deviation of the visual cue specified by Eq.4. Parameter set $\Theta = [\sigma_{int}, a, b, B]$ was

fitted to the average data from all participants. Model simulations (Figure 5A) were

performed with the same single-trial learning equations. For the clear cursor condition,

we used the same parameter values estimated from Experiment 2 (see details in Table

S1). For the blurred cursor condition, the standard deviation of visual cue was changed

to:

$$\sigma_{v,blur} = R_v \sigma_v \tag{12}$$

826   for the simulation of the increase in visual uncertainty, the ratio R_v varied from 1.1 to 3.

827   ***PReMo model***

828   We used the PReMo model to fit the average adaptation extent obtained from Experiment

829   2 (Figure 3C & Figure S3B). Following the study by (Tsay, Kim, et al., 2022), the hand

830   position at trial *n+1* is:

$$x_{p,n+1} = Ax_{p,n} + B(T - x_{per,n}), \tag{13}$$

832   where

$$x_{per,n} = \beta_p + \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2} x_{p,n}, \tag{14}$$

$$\beta_p = -min\left(\left|\beta_p^{sat}\right|, \left|\eta_p\left(\frac{\sigma_u^2}{\sigma_u^2 + \sigma_v^2} x_{v,n} - \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2} x_{p,n}\right)\right|\right). \tag{15}$$

In data fitting, we used two parameters to represent the ratio between sensory cues: $R_1 =$

$\sigma_u^2/(\sigma_u^2 + \sigma_v^2)$ and $R_2 = \sigma_u^2/(\sigma_u^2 + \sigma_p^2)$. The data were fitted with the parameter set $\Theta = [R_1,$

$R_2, \beta_p^{sat}, \eta_p, A, B]$, where β_p^{sat} is the saturation angle, η_p is a scaling factor, *A* is the

retention rate and *B* is the learning rate. For simulating the proprioceptive localization of

the hand (Figure 4C), the parameter values estimated from Experiment 2 were used. The

bias of hand estimation in the proprioception trials is determined as: $x_{bias} = -(0 -$

$x_{per})R_p$, where ratio R_p varies from 0.05 to 0.8. Thus, similar to the PEA model simulation,

the proprioceptive bias is a fraction of the bias in the hand estimation from the adaptation

trials. Single-trial learning (Figure 5B) was simulated with:

$$x_{STL} = B(T - x_{per}),\tag{16}$$

845   where x_{per} is determined by Eq. 12 and Eq. 13. For the clear condition, we used the

846   parameter values estimated from Experiment 2 with PReMo. For the blurred cursor

847   condition, the standard deviation of visual signal $\sigma_{v,blur}$ increases with a ratio R_v, as in Eq.

848   12.

Causal inference model

850   The causal inference model by (Wei & Körding, 2009) was used to fit the data of

851   Experiment 2 (Figure 3D & Figure S3C). The hand position at trial n+1 is updated by

852   learning from visual error at trial n:

$$x_{p,n+1} = Ax_{p,n} + B(T - px_{v,n}),\tag{17}$$

854   where *A* and *B* are the retention and learning rates, respectively; *T* is the target direction.

855   Specifically for this model, the learning from error is modulated by the probability (*p*) of

856   causal attribution of visual error to the action or proprioception:

$$p = S \frac{N(x_{v,n},0,\sigma^2)}{N(x_{v,n},0,\sigma^2)+C},\tag{18}$$

858   where $x_{v,n}$ is the visual cue at trial n. *S* and *C* are the scaling factors, and σ is the standard

859   deviation of the integrated cue combining visual and proprioceptive cues, following

$$\sigma^2 = \frac{\sigma_v^2 \sigma_p^2}{\sigma_v^2 + \sigma_p^2}\tag{19}$$

861   Thus, the data were fitted with five parameters $\Theta = [\sigma, S, C, A, B]$. For simulating single-trial

862   learning with cursor blurring (Figure 5C), the ratio between σ_v and σ_p is fixed as ½. The

863   single-trial leaning was determined as:

864
$$x_{STL} = B(T - px_v), \tag{20}$$

where p is determined by Eq. 18. Put Eq. 12 and Eq. 19 into $\sigma_{blur}^2 = \frac{\sigma_{v,blur}^2 \sigma_p^2}{\sigma_{v,blur}^2 + \sigma_p^2}$, we can

calculate the standard deviation of the integrated sensory signal for the blurred cursor:

$\sigma_{blur} = \sigma \sqrt{\frac{5R^2}{R^2+4}}$. Simulation was performed with R ranging from 1.1 to 3.

Data fitting

All data were fitted using MATLAB (2022b, MathWorks, Natick, MA, US) build-in function

fmincon with 100 randomly sampled initial values of parameter sets. See Table S1 and

Table S2 for the fitted parameter values and comparisons between different models.

Data availability

Data presented in this work are available at:

https://doi.org/10.6084/m9.figshare.24503926.v1.

References

Albert, S. T., Jang, J., Modchalingam, S., Marius't Hart, B., Henriques, D., Lerner, G.,

Della-Maggiore, V., Haith, A. M., Krakauer, J. W., & Shadmehr, R. (2022).

Competition between parallel sensorimotor learning systems. *ELife*, *11*, e65361.

Albert, S. T., Jang, J., Sheahan, H. R., Teunissen, L., Vandevoorde, K., Herzfeld, D. J.,

& Shadmehr, R. (2021). An implicit memory of errors limits human sensorimotor

adaptation. *Nature Human Behaviour*, 1–15.

Avraham, G., Morehead, J. R., Kim, H. E., & Ivry, R. B. (2021). Reexposure to a

sensorimotor perturbation produces opposite effects on explicit and implicit

learning processes. *PLoS Biology*, *19*(3), e3001147.

Avraham, G., Taylor, J. A., Breska, A., Ivry, R. B., & McDougle, S. D. (2022). Contextual

effects in sensorimotor adaptation adhere to associative learning rules. *ELife*, *11*,

e75801.

Berniker, M., & Kording, K. (2008). Estimating the sources of motor errors for adaptation

and generalization. *Nature Neuroscience*, *11*(12), 1454–1461.

Berniker, M., & Kording, K. (2011). Bayesian approaches to sensory integration for motor

control. *Wiley Interdisciplinary Reviews. Cognitive Science*, *2*(4), 419–428.

Bond, K. M., & Taylor, J. A. (2015). Flexible explicit but rigid implicit learning in a

visuomotor adaptation task. *Journal of Neurophysiology*, *113*(10), 3836–3849.

Burge, J., Ernst, M. O., & Banks, M. S. (2008). The statistical determinants of adaptation

rate in human reaching. *Journal of Vision*, *8*(4), 1–19.

Cheng, S., & Sabes, P. N. (2006). Modeling sensorimotor learning with linear dynamical

systems. *Neural Computation*, *18*(4), 760–793.

Cressman, E. K., & Henriques, D. Y. P. (2009). Sensory recalibration of hand position

following visuomotor adaptation. *Journal of Neurophysiology*, *102*(6), 3505–3518.

Della-Maggiore, V., & McIntosh, A. R. (2005). Time course of changes in brain activity

and functional connectivity associated with long-term adaptation to a rotational

transformation. *Journal of Neurophysiology*, *93*(4), 2254–2262.

Donchin, O., Francis, J. T., & Shadmehr, R. (2003). Quantifying generalization from trial-

by-trial behavior of adaptive systems that learn with basis functions: theory and

experiments in human motor control. *The Journal of Neuroscience: The Official*

Journal of the Society for Neuroscience, *23*(27), 9032–9045.

He, K., Liang, Y., Abdollahi, F., Bittmann, M. F., Körding, K., & Wei, K. (2016). The

statistical determinants of the speed of motor learning. *PLoS Computational*

Biology, *12*(9), e1005023.

Henriques, D. Y. P., & Cressman, E. K. (2012). Visuomotor adaptation and proprioceptive

recalibration. *Journal of Motor Behavior*, *44*(6), 435–444.

Herzfeld, D. J., Vaswani, P. A., Marko, M. K., & Shadmehr, R. (2014). A memory of errors

in sensorimotor learning. *Science*, *345*(6202), 1349–1353.

Huberdeau, D. M., Krakauer, J. W., & Haith, A. M. (2019). Practice induces a qualitative

change in the memory representation for visuomotor learning. *Journal of*

Neurophysiology, *122*(3), 1050–1059.

Hutter, S. A., & Taylor, J. A. (2018). Relative sensitivity of explicit reaiming and implicit

motor adaptation. *Journal of Neurophysiology*, *120*(5), 2640–2648.

Izawa, J., & Shadmehr, R. (2011). Learning from sensory and reward prediction errors

during motor adaptation. *PLoS Computational Biology*, *7*(3), e1002012.

Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R., & Ivry, R. B. (2018). Invariant

errors reveal limitations in motor correction rather than constraints on error

sensitivity. *Communications Biology*, *1*(1), 19.

Kirk, R. (1968). *Experimental design: Procedures for the behavioral sciences* (Vol. 1–f).

Brooks/Cole.

Klein, S. A., & Levi, D. M. (1987). Position sense of the peripheral retina. *JOSA A*, *4*(8),

1543–1553.

Körding, K., & Wolpert, D. M. (2004). Bayesian integration in sensorimotor learning.

Nature, *427*(6971), 244–247.

Krakauer, J. W., Ghez, C., & Ghilardi, M. F. (2005). Adaptation to visuomotor

transformations: consolidation, interference, and forgetting. *The Journal of

Neuroscience: The Official Journal of the Society for Neuroscience*, *25*(2), 473–

478.

Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L., & Haith, A. M. (2019). Motor learning.

Comprehensive Physiology, *9*(2), 613–663.

Landi, S. M., Baguear, F., & Della-Maggiore, V. (2011). One week of motor adaptation

induces structural changes in primary motor cortex that predict long-term memory

one year later. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *31*(33), 11808–11813.

Leow, L.-A., Marinovic, W., de Rugy, A., & Carroll, T. J. (2020). Task errors drive

memories that improve sensorimotor adaptation. *Journal of Neuroscience*, *40*(15),

3075–3088.

Levi, D. M., Klein, S. A., & Yap, Y. L. (1987). Positional uncertainty in peripheral and

amblyopic vision. *Vision Research*, *27*(4), 581–597.

Lieberman, H. R., & Pentland, A. P. (1982). Microcomputer-based estimation of

psychophysical thresholds: the best PEST. *Behavior Research Methods &*

Instrumentation, *14*(1), 21–25.

Mazzoni, P., & Krakauer, J. W. (2006). An implicit plan overrides an explicit strategy

during visuomotor adaptation. *The Journal of Neuroscience: The Official Journal*

of the Society for Neuroscience, *26*(14), 3642–3645.

McDougle, S. D., Bond, K. M., & Taylor, J. A. (2015). Explicit and Implicit Processes

Constitute the Fast and Slow Processes of Sensorimotor Learning. *The Journal of*

Neuroscience: The Official Journal of the Society for Neuroscience, *35*(26).

Mikulasch, F. A., Rudelt, L., & Priesemann, V. (2022). Visuomotor mismatch responses

as a hallmark of explaining away in causal inference. *Neural Computation*, *35*(1),

27–37.

Miyamoto, Y. R., Wang, S., & Smith, M. A. (2020). Implicit adaptation compensates for

erratic explicit strategy in human motor learning. *Nature Neuroscience*, *23*(3), 443–

455.

Modchalingam, S., Vachon, C. M., 't Hart, B. M., & Henriques, D. Y. P. (2019). The effects

of awareness of the perturbation during motor adaptation on hand localization.

PloS One, *14*(8), e0220884.

Morehead, J. R., Taylor, J. A., Parvin, D. E., & Ivry, R. B. (2017). Characteristics of implicit

sensorimotor adaptation revealed by task-irrelevant clamped feedback. *Journal of*

Cognitive Neuroscience, *29*(6), 1061–1074.

Mostafa, A. A., 't Hart, B. M., & Henriques, D. Y. P. (2019). Motor learning without moving:

proprioceptive and predictive hand localization after passive visuoproprioceptive

discrepancy training. *PloS One*, *14*(8), e0221861.

Neville, K. M., & Cressman, E. K. (2018). The influence of awareness on explicit and

implicit contributions to visuomotor adaptation over time. *Experimental Brain*

Research. Experimentelle Hirnforschung. Experimentation Cerebrale, *236*(7),

2047–2059.

Oh, Y., & Schweighofer, N. (2019). Minimizing Precision-Weighted Sensory Prediction

Errors via Memory Formation and Switching in Motor Adaptation. *The Journal of*

Neuroscience: The Official Journal of the Society for Neuroscience, *39*(46), 9237–

9250.

Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh

inventory. *Neuropsychologia*, *9*(1), 97–113.

Rossi, C., Bastian, A. J., & Therrien, A. S. (2021). Mechanisms of proprioceptive

realignment in human motor learning. *Current Opinion in Physiology*, *20*, 186–197.

Ruttle, J. E., Cressman, E. K., 't Hart, B. M., & Henriques, D. Y. P. (2016). Time course

of reach adaptation and proprioceptive recalibration during visuomotor learning.

PloS One, *11*(10), e0163695.

Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. P. (2018). The fast contribution of visual-

proprioceptive discrepancy to reach aftereffects and proprioceptive recalibration.

PloS One, *13*(7), e0200621.

Ruttle, J. E., 't Hart, B. M., & Henriques, D. Y. P. (2021). Implicit motor learning within

three trials. *Scientific Reports*, *11*(1), 1627.

Salomonczyk, D., Cressman, E. K., & Henriques, D. Y. P. (2011). Proprioceptive

recalibration following prolonged training and increasing distortions in visuomotor

adaptation. *Neuropsychologia, 49*(11), 3053–3062.

Salomonczyk, D., Cressman, E. K., & Henriques, D. Y. P. (2013). The role of the cross-

sensory error signal in visuomotor adaptation. *Experimental Brain Research.*

Experimentelle Hirnforschung. Experimentation Cerebrale, *228*(3), 313–325.

Seilheimer, R. L., Rosenberg, A., & Angelaki, D. E. (2014). Models and processes of

multisensory cue combination. *Current Opinion in Neurobiology*, *25*, 38–46.

Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction,

and adaptation in motor control. *Annual Review of Neuroscience*, *33*, 89–108.

Simani, M. C., McGuire, L. M. M., & Sabes, P. N. (2007). Visual-shift adaptation is

composed of separable sensory and task-dependent effects. *Journal of*

Neurophysiology, 98(5), 2827–2841.

Smith, M. A., Ghazizadeh, A., & Shadmehr, R. (2006). Interacting adaptive processes
with different timescales underlie short-term motor learning. *PLoS Biology*, *4*(6),
e179.

Stetson, C., Cui, X., Montague, P. R., & Eagleman, D. M. (2006). Motor-sensory
recalibration leads to an illusory reversal of action and sensation. *Neuron*, *51*(5),
651–659.

't Hart, B. M., & Henriques, D. Y. P. (2016). Separating predicted and perceived sensory
consequences of motor learning. *PloS One*, *11*(9), e0163556.

Taylor, J. A., & Ivry, R. B. (2011). Flexible cognitive strategies during motor learning.
PLoS Computational Biology, *7*(3), e1001096.

Taylor, J. A., Krakauer, J. W., & Ivry, R. B. (2014). Explicit and implicit contributions to
learning in a sensorimotor adaptation task. *The Journal of Neuroscience: The
Official Journal of the Society for Neuroscience*, *34*(8), 3023–3032.

Thoroughman, K. A., & Shadmehr, R. (2000). Learning of action through adaptive
combination of motor primitives. *Nature*, *407*(6805), 742.

Tsay, J. S., Avraham, G., Kim, H. E., Parvin, D. E., Wang, Z., & Ivry, R. B. (2021). The
effect of visual uncertainty on implicit motor adaptation. *Journal of
Neurophysiology*, *125*(1), 12–22.

Tsay, J. S., Haith, A. M., Ivry, R. B., & Kim, H. E. (2022). Interactions between sensory
prediction error and task error during implicit motor learning. *PLoS Computational
Biology*, *18*(3), e1010005.

Tsay, J. S., Kim, H. E., Parvin, D. E., Stover, A. R., & Ivry, R. B. (2021). Individual

differences in proprioception predict the extent of implicit sensorimotor adaptation.

Journal of Neurophysiology, *125*(4), 1307–1321.

Tsay, J. S., Kim, H., Haith, A. M., & Ivry, R. B. (2022). Understanding implicit sensorimotor

adaptation as a process of proprioceptive re-alignment. *ELife*, *11*, e76639.

Tsay, J. S., Parvin, D. E., & Ivry, R. B. (2020). Continuous reports of sensed hand position

during sensorimotor adaptation. *Journal of Neurophysiology*, *124*(4), 1122–1130.

Tseng, Y., Diedrichsen, J., Krakauer, J. W., Shadmehr, R., & Bastian, A. J. (2007).

Sensory prediction errors drive cerebellum-dependent adaptation of reaching.

Journal of Neurophysiology, *98*(1), 54.

Wei, K., & Körding, K. (2009). Relevance of error: what drives motor adaptation? *Journal*

of Neurophysiology, *101*(2), 655–664.

Wei, K., & Körding, K. (2010). Uncertainty of feedback and state estimation determines

the speed of motor adaptation. *Frontiers in Computational Neuroscience*, *4*, 11.

Wolpert, D. M., Diedrichsen, J., & Flanagan, J. R. (2011). Principles of sensorimotor

learning. *Nature Reviews. Neuroscience*, *12*(12), 739.

Wozny, D. R., & Shams, L. (2011). Recalibration of auditory space following milliseconds

of cross-modal discrepancy. *Journal of Neuroscience*, *31*(12), 4607–4612.

Wu, H. G., Miyamoto, Y. R., Castro, L. N. G., Ölveczky, B. P., & Smith, M. A. (2014).

Temporal structure of motor variability is dynamically regulated and predicts motor

learning ability. *Nature Neuroscience*, *17*(2), 312.

1045

Figure 1. The Perceptual Error Adaptation (PEA) model for error-clamp adaptation. **(A)**

Illustration of involved sensorimotor cues for estimating hand direction $\hat{x}_{Hand}$. The

clamped cursor, the hand, and the sensory prediction of the reaching action provide the

visual (x_v), proprioceptive (x_p), and the sensory prediction cue (x_u) of movement direction,

respectively. The hand direction estimate is assumed to be based on maximum likelihood

cue combination. **(B)** Assuming a linear dependency of visual uncertainty on eccentricity,

the PEA model predicts that implicit adaptation extent is a concave function of

perturbation size θ, a pattern qualitatively aligning with previous findings (Kim et al., 2018;

Morehead et al., 2017).

1055

Figure 2. Experiment 1 measuring the dependency of visual uncertainty on perturbation

size. **(A)** The 2AFC task for judging the cursor motion direction. In an exemplary trial, the

participant reaches to a target while a direction-clamped cursor moves concurrently,

serving as an error-clamp perturbation. Following a 1000-ms blank masking period, a

reference point appears for 150ms, either clockwise or counterclockwise from the

clamped cursor. The participant is then asked to making a binary judgement regarding

the direction of the clamped cursor relative to the reference point. **(B)** The visual

uncertainty, obtained from psychometrical estimation based on the 2AFC, is plotted as a

function of perturbation size. Both individual estimates (red dots) and group-level statistics

(boxplots) are shown. Positive angles correspond to CW rotations, negative angles to

CCW rotations. **(C)** Collapsing data from both rotation directions, we observe that visual

uncertainty closely follows a linear function of perturbation size. The dark gray line and

its shaded region denote the regression line and its 95% confidence interval, respectively.

The purple line is generated with the values of a and b fitted from data in Experiment 2

with a and b treated as free parameters (See Methods for details).

1071

Figure 3. Results and model fitting of Experiment 2. **(A)** Implicit adaptation to error clamps

of varying sizes is depicted; colored dot-lines and colored shading area represent the

mean and standard error for each participant group. The light gray area indicates trials

with error-clamp perturbations. Adaptation starts after baseline, gradually asymptotes to

its final extent, and then decays with null feedback during washout. Different perturbation

sizes result in distinct adaptation rates and extents. Group averages and standard error

across participants are shown, along with predictions (colored solid lines) from the PEA

model. **(B)** The adaptation extent (cycle 100-110) exhibits a nonlinear dependency on

perturbation size, conforming to a concave function as prescribed by PEA (purple line).

Color dots and error bars denote the mean and standard error across participants in each

1082   group. **(C)-(D)** The same data fitted with the PReMo model and the causal inference

1083   model. See more details, refer to Figure S3.

1084

1085

Figure 4. Proprioceptive data fitting and results from Experiment 3. **(A)** The data from

(Tsay et al., 2020) are presented alongside the fitting of the PEA model. Participants

adapting to a 30° error-clamp perturbation were required to report the location of their

adapted hand using visual aids of numbers. The report was provided when the hand

stayed at the end of movement. Initially, the proprioceptive estimate of the hand is biased

toward the visual cursor (negative in the plot) and then gradually shifts toward the hand

(positive in the plot). This trend is accurately captured by the PEA model: lines represent

model fitting results, with the adapted hand direction in indigo and the reported hand

direction in blue. The hand direction estimate ($\hat{x}_{Hand}$, Eq.1) following a reach movement

is shown in red. **(B)-(C)** Model simulations for proprioceptive bias from the PEA and

PReMo models. Color gradients denote the simulations with varying ratio between the

weights of $\hat{x}_{Hand}$ and x_p, the two cues available for estimating the hand direction. Note

that the two models prescribe distinct profiles for the dependency of proprioception bias

on perturbation size. **(D)** Experimental design. A reaching block, either with or without

visual perturbations, is followed by a proprioception test block. The size and direction of

the visual perturbation vary across blocks. The proprioception test is conducted when the

hand is passively moved to a target (red dots) situated near the reaching target (blue

dots). **(E)** The observed proprioceptive bias as a function of perturbation size. Data from

the three proprioception test trials are separately plotted. The first trial reveals

proprioception biases that form a concave function of perturbation size.

1106

Figure 5. Results of Experiment 4. **(A)-(C)** Model simulations for single-trial learning under different visual uncertainty levels, shown separately for the PEA, PReMo and causal inference models. Blue curves represent simulated learning based on model parameters estimated from Experiment 2. Curves with red gradient indicate simulations with increasing levels of visual uncertainty, color coded by the ratio of visual uncertainty for the blurred cursor to that of the clear cursor. **(D)** Experimental design. Following 60 baseline trials without perturbations, participants completed 15 mini-blocks of error-clamp adaptation over three successive days. Each mini-block features 12 different types of error-clamp perturbations, distinguished by two cursor presentations (blurred or clear cursor) and six clamp sizes. Each perturbation trial, varied randomly in perturbation type, is flanked by two no-feedback trials. The change in hand direction between these two no-

1118   feedback trials serves to quantify singe-trial learning. **(E)** The single-trial learning with the

1119   blurred cursor is less than that with the clear cursor, but the difference is non-monotonic

1120   across perturbation size (*** denote $p < 0.001$).

1121

Figure 6. Design of Experiment 1. **(A)** Top-down view of the setup in visual discrimination task. The reference point (yellow) was presented either CW or CCW relative to the clamped cursor (dashed circle), which has a perturbation size θ. **(B)** Trial structure of the visual discrimination task. Purple rectangles represent error-clamped trials with varying perturbation size, rectangles with yellow edges represent the ensuing visual discrimination test for each perturbation size. **(C)-(D)** Exemplary sequences of the reference point: These sequences illustrate the deviation of the reference point from the cursor (C) and the changing step size across trials (D), following the PEST algorithm. Individual trials are represented by blue dots. Yellow and red dots mark the initiation and termination of each round of trials, respectively. In each round, the reference point starts on either the CW or CCW side of the cursor; In the subsequent round, it starts on the opposite side.

1134

Figure 7. Setup for measuring proprioceptive recalibration in Experiment 3. **(A)** Reaching movement with error-clamped cursor, performed by the right hand holding a robot handle. **(B)** Passive movement in the proprioception test. The right hand was passively moved to the unseen target (h_R), depicted here as a small black dot. A red hollow circle with an expanding radius appears on the screen during passive movement, signaling the increasing distance between the hand and the start position. Subsequently, participants used their left hand to report the right-hand location (h_p) by aligning a red rectangle on the red circle, which is displayed at the target distance.

1143 **Supplementary Materials**

1144

1145

Figure S1. Heat map of eye fixations during the 2AFC task in Experiment 1. The screen

is partitioned into 10x10 pixel grids, and the cumulative number of gaze samples in each

grid is recorded. Data from all participants, aggregated across each day of practice, are

presented. The color map signifies the normalized count of gaze samples in each grid.

Data are separately displayed for the three distinct phases of a trial, as delineated by the

columns on the left, middle, and right. These correspond to periods during hand

movement, the appearance of the visual mask and reference point, and the time allotted

for manual response. On average, 95.06%, 89.93%, and 86.55% of gaze samples fall

within the ±50-pixel range of the central line during these three phases, respectively.

These results corroborate that participant adhered to the instructions and refrained from

looking at the cursor during the visual discrimination task.

1157

1158

Figure S2. Performance of an exemplary participant in Experiment 1. Six panels display

the psychometric curves corresponding to different error-clamp sizes. The x-axis denotes

the angular deviation between the clamped cursor and the reference point (as depicted

in Figure 6A). A negative value implies that the reference point appears on the

counterclockwise (CCW) side of the clamped cursor. The blue dots represent the

proportion of trials in which the participant reported that "the yellow point is on the

clockwise (CW) side of the clamped cursor" for various angular deviations between these

two. Data were aggregated from all trials across three days of the experiment. The gray-

shaded region represents the interquartile range (25th to 75th percentile) of the

psychometric curve, and the width of this shaded region serves as an indicator of the

amplitude of visual uncertainty.

1170

Figure S3. Model fitting for observed implicit adaptation in Experiment 2. This
supplementary figure provides a comprehensive evaluation of the three competing
models: the PEA model, the PReMo model, and the causal inference model. **(A)** Results
of PEA Model Fitting: The layout of these plots mirrors that of Figures 3A and 3B, serving
as a direct comparison between the empirical data and the predictions made by the PEA

model. **(B)** Results of PReMo Model Fitting: The left panel is a duplicate of Figure 3C,

while the right panel presents the trial-by-trial data fitting. This juxtaposition allows for a

nuanced evaluation of the PReMo model's performance at both the aggregate and

individual trial levels. **(C)** Results of Causal Inference Model: The arrangement of these

plots is consistent with panels (A) and (B), facilitating a straightforward comparison of all

three models. For a detailed assessment of the quality of model fitting and subsequent

model comparisons, please refer to Table S1 and Table S2.

1183

1184

Figure S4. Correlation between initial learning rate and adaptation extent in Experiment

2. For each participant, the initial learning rate is calculated as the change in hand angle

between the 1st and 10th cycle, divided by 10. The adaptation extent is defined as the

average hand angle across the last 10 adaptation cycles. When pooling data across all

perturbation sizes, a significant correlation is observed between the initial learning rate

and the adaptation extent.

1191

Figure S5. Model fitting of single-trial learning from Experiment 2 of (Tsay, Avraham, et

al., 2021). Blue dots represent the mean single-trial learning across varying perturbation

size, with error bars represent denoting standard errors across participants. The left,

middle and right panel present the fitting results for the PEA, PReMo, and causal

inference models, respectively. For additional details, refer to the Methods, Results, and

Table S1 & S2.

1198

Figure S6. Proprioception uncertainty estimation results. Thirteen participants from Experiment 1 participated in a proprioception discrimination task to measure their proprioceptive uncertainty in the setting of the error-clamp adaptation. The setup paralleled that used for estimating visual uncertainty in Experiment 1. In each trial, participants initially held their hand at the starting position. They were instructed to relax their arm while the experimenter, seated on the other side of the monitor, pulled their hand to a proprioceptive target near the straight-ahead target. The arms of the experimenter and the participant were blocked from the view of the participant. After 0.8 seconds, a yellow reference point appeared. The angular deviation between the participant's hand and this reference was determined using the same PEST procedure employed in Experiment 1. Participants indicated, by pressing left or right arrow keys by their left hand, whether the reference point appeared on the CW or CCW side of their actual right-hand position. The maximum deviation allowed was 30°, with an initial step

1212   size of 10° and a stop threshold of 0.5°. This task was conducted over six runs across

1213   three consecutive days. Similar to Figure S2, panel (A) to (M) show the psychometric

1214   curves for each participant with data from the three days pooled together. (N) and (O)

1215   present the measured proprioceptive uncertainty and bias for all participants (gray dots)

1216   and their mean ± standard deviation (red error bars).

1217

1218   **Table S1.** Model fitting and simulation parameters with the PEA model.

	Data set	Parameters						Goodness-of-fit	
		σ_u (deg)	σ_p (deg)	a	b	A	B	R^2	RMSE (deg)
Adaptation extent fitting (Figure 1B)	Kim 2018, Exp1	--	$\sigma_p/a = 3.406$; $b/a = 0.138$			--	--	0.773	1.898
	Kim 2018, Exp2	--	$\sigma_p/a = 4.758$; $b/a = 0.168$			--	--	<0	2.163
	Morehead, 2017	--	$\sigma_p/a = 1.639$; $b/a = 0.044$			--	--	<0	2.937
Trial-by-trial adaptation fitting	Exp 2, Figure 3	5.048	11.119	*1.853	*0.309	0.970	0.208	0.975	1.222
		5.468	12.128	1.663	0.331	0.971	0.194	0.975	1.217
	Tsay 2019, Figure 4A	1.896	7.959	*1.853	*0.309	0.978	0.525	0.991	0.716
Single-trial learning fitting	Tsay 2021, Figure S6	$\sigma_{int} = 7.364$		1.179	0.384	--	0.057	0.974	0.020
Proprioceptive recalibration simulation	Exp 3, Figure 4B	5.048	11.119	1.853	0.309	0.970	0.208	--	--
Adaptation affected by visual uncertainty simulation	Exp 4, Figure 5	5.048	11.119	1.853	0.309	--	0.208	--	--

1219   * Asterisks represent fixed parameters in specific data fitting. The fixed values equal to

1220   the slope and intercept estimated from Experiment 1.

1221 **Table S2.** Model comparisons.

Data set		PEA	PReMo	Causal Inference
Block-design learning fitting Exp 2, Figure 3 & S3	AIC	2255	3543	3283
	R^2	0.975	0.749	0.711
	$RMSE$ (deg)	1.222	3.896	4.151
Single-trial learning fitting Tsay 2021, Figure S5	AIC	-36.90	-15.98	-11.28
	R^2	0.974	0.584	0.305
	$RMSE$ (deg)	0.020	0.103	0.080

1222