Isogenic iPSC-derived proximal and distal lung-on-chip models: Tissue- and virus-specific immune responses in human lungs

Sachin Yadav[1,2], Kazuya Fujimoto[1,2], Toru Takenaga[2,3,4], Senye Takahashi[2,5], Yukiko Muramoto[2,3,4], Ryuta Mikawa[2,5,6], Takeshi Noda[2,3,4], Shimpei Gotoh[2,5,6], Ryuji Yokokawa[1,2]

1. Department of Micro Engineering, Kyoto University, Kyoto 615-8540, Japan
2. CREST, Japan Science and Technology Agency, Saitama 332-0012, Japan
3. Institute of Frontier Life and Medical Science, Kyoto University, Kyoto 606-8507, Japan
4. Graduate School of Biostudies, Kyoto University, Kyoto 606-8507, Japan
5. Center for iPS Cell Research and Application, Kyoto University, Kyoto 606-8507, Japan
6. Department of Drug Discovery for Lung Diseases, Kyoto University, Kyoto 606-8501, Japan

*Corresponding author email: yokokawa.ryuji.8c@kyoto-u.ac.jp (R.Y)

Abstract

Micro-physiological systems (MPS) are set to play a vital role in preclinical studies, particularly in the context of future viral pandemics. Nonetheless, the development of MPS is often impeded by the scarcity of reliable cell sources, especially when seeking various organs or tissues from a single patient for comparative analysis of the host immune response. Herein, we developed human airway-on-chip and alveolus-on-chip models using induced pluripotent stem cell (iPSC)-derived isogenic lung progenitor cells. Both models demonstrated the replication of two different respiratory viruses, namely SARS-CoV-2 and Influenza, as well as related cellular damage and innate immune responses-on-chip. Our findings reveal distinct immune responses to SARS-CoV-2 in the proximal and distal lung-on-chip models. The airway chips exhibited a robust interferon (IFN)-dependent immune response, whereas the alveolus chips exhibited dysregulated IFN activation but a significantly upregulated chemokine pathway. In contrast, Influenza virus infection induced a more pronounced immune response and cellular damage in both chip models compared to SARS-CoV-2. Thus, iPSC-derived lung-on-chip models may aid in quickly gaining insights into viral pathology and screening potential drugs for future pandemics.

Human lungs are prone to infections caused by respiratory viruses, including SARS-CoV-2 and Influenza A virus (IAV)[1,2]. These respiratory infections have been responsible for numerous pandemics and have placed a substantial burden on healthcare systems and management, as exemplified by the COVID-19 pandemic [3–7]. These infections affect various regions of the human lungs, from the proximal airway tract to the distal alveolar sac. Typically, a virus gains entry through cells that express virus receptors, such as multiciliated cells in the lung airway. Eventually, it reaches alveolar type I (AT1) or alveolar type 2 (AT2) cells in the alveoli, leading to alveolar diffusion and other pathological effects[8–12]. In severe cases, the infection can also spread to other organs, resulting in multiorgan damage[13–16].

Host cells detect pathogens through "pattern recognition receptors" which, upon activation, trigger innate immune responses and generate multiple interferons (IFN) and proinflammatory cytokines[17–19]. Additionally, chemokines, which are chemotactic cytokines controlling the migration and positioning of immune cells in tissues, play a critical role in the function of the innate immune system[20,21].

Viruses such as SARS-CoV-2 and IAV have developed strategies to evade the innate immune response, for example, by inhibiting the type I IFN pathway[22,23]. An attenuated type I IFN response can result in uncontrolled viral replication and the accumulation of immune cells at the infection site, ultimately leading to an excessive release of IFNs and the onset of a cytokine storm[24].

Recent research findings also indicate distinct tissue-specific innate immune responses in the upper and lower respiratory tracts, especially in the context of coronaviruses such as SARS-CoV-2. These findings suggest a role for the upper airway in active viral transmission and immune defense. The diminished innate immune response in the distal lung may potentially underlie the unique uncontrolled late-phase lung damage observed in advanced COVID-19[25].

The complications associated with the host immune response, especially following viral infections, highlight the need for urgently developing human-relevant models that can closely mimic the human lung microenvironment. Such models provide a robust platform for comprehending infection mechanisms and drug screening. While traditionally employed

animal models have been widely used for drug screening, they often encounter ethical and

technical challenges, including genetic differences[26].

In vitro models, such as dish and insert cultures fail to accurately replicate the *in vivo*

interface of human lungs. Since the onset of the recent COVID-19 pandemic, researchers

have increasingly turned to micro-physiological systems (MPS) to emulate viral infections

and the associated damage[27–29]. Si et al. have demonstrated the potential of MPS for drug

screening and repurposing[30]. Nevertheless, these models exhibit significant irregularities,

such as inadequate viral replication, atypical immune responses, and an inability to depict the

evasion of immune responses by viruses such as SARS-CoV-2. Overall, these models display

variability among cell types and struggle to accurately reproduce the diverse immune

signatures of viral infections.

One of the primary reasons for these irregularities is the lack of a suitable cell source, which

is a common limitation of lung MPS. Immortalized cells lack similarity to *in vivo* phenotypes

and genotypes, rendering them unsuitable for emulating the human lung tissue interface[31]. In

contrast, primary cells have limited proliferative capacity and instability under *in vitro*

conditions; for example, AT2 cells tend to differentiate into AT1-like cells when cultured *in*

vitro[32]. These limitations result in irregular viral replication and immune responses under *in*

vitro conditions, including MPS[27,28]. Moreover, SARS-CoV-2 exhibits varying infectivity in

different regions of the human lung owing to the spatial arrangement of receptor proteins[33]

and evokes different immune responses[25,34]. This underscores the critical need for a system

capable of emulating different regions of the human lung to compare host immune responses

and screen relevant drugs.

The integration of induced pluripotent stem cells (iPSCs) with MPS effectively overcomes

the limitations of traditional cell sources and *in vitro* models, such as their failure to mimic

the *in vivo* interface and their shortcomings in terms of cell availability and stability. The

utilization of iPSCs offers additional advantages, including the ability to develop different

regions of the human lung from a single-cell source and the potential for personalized models.

In this study, we developed airway-on-chip and alveolus-on-chip models using iPSC-derived

isogenic lung progenitor cells to replicate viral infections and the associated innate immune

responses on-chip.

Results

Differentiation of iPSCs into the airway and alveolar cells on-chip

We utilized iPSC-derived isogenic carboxypeptidase M (CPM+) lung progenitor cells to

establish airway-on-chip and alveolus-on-chip models (Fig. 1a). Briefly, iPSCs underwent a

21-day differentiation process on a culture plate, and magnetic-activated cell sorting (MACS)

was employed to isolate CPM+ lung progenitor cells. These CPM+ cells were then seeded

into the airway and alveolus chips with the respective culture medium (Fig. S1).

Immunofluorescence staining of ZO-1 and VE-Cadherin revealed the formation of tight

junctions in both the epithelium and endothelium tissues of the airway chip respectively (Fig.

1b, c). The optimal differentiation of (CPM+) cells into multiciliated cells was confirmed by

immunofluorescent staining of the airway multiciliated cell marker acetylated tubulin (Ac-

Tub). Additionally, other airway epithelial cell such as mucus-secreting cells (MUC5AC) and

basal cells (Cytokeratin 5) were also observed (Fig. 1d), consistent with prior reports[35]. Gene

expression analysis of key multiciliated cell markers, including *SNTN* and *FOXJ1*, indicated

progressive differentiation of CPM+ lung progenitor cells into multiciliated cells by day 28

(Fig. 1e). Furthermore, the presence of mucus-secreting club cells was verified by the

upregulation of *SCGB1A1 (CCSP)* and *SPDEF* genes (Fig. S2a).

Similarly, immunofluorescent staining of ZO-1 and VE-Cadherin showed the formation of

tight junctions in both the epithelium and endothelium tissues of the alveolus chip

respectively (Fig. 1f, g). Given that an SFTPC-GFP knock-in iPSC line[36] was utilized in the

experiments, alveolar type 2 (AT2) cells expressed GFP and were detected using an anti-GFP

antibody (Fig. 1h). The differentiation of CPM+ cells into AT2 cells was further confirmed

through immunofluorescent staining of SLC34A2 (Fig. 1i), a key marker for AT2 cells[37].

Additionally, some alveolar type 1 (AT1)-like cells were observed in the alveolar epithelium

by staining HT1-56 (Fig. 1j). Gene expression analysis of key AT2 cell markers, including

SFTPC, SFTPB, and *SLC34A2*, demonstrated the progressive differentiation of CPM+ cells

into AT2 cells by day 7 (Fig. 1k). Gene expression of AT1 cell marker (*AGER* and *PDPN*)

also increased from day zero to day 10 (Fig. S2b). Given that multiciliated cells and AT2

cells are the primary targets of SARS-CoV-2 in the proximal airway and alveoli, respectively,

our airway chip and alveolus chip offer an excellent means to replicate viral pathogenesis and

host immune responses.

Both chips exhibit high susceptibility to SARS-CoV-2 with minimal cellular damage

Transmembrane proteins, namely angiotensin-converting enzyme 2 (ACE2), transmembrane

serine protease 2 (TMPRSS-2), and neuropilin-1 (NRP-1), play a pivotal role in facilitating

the entry of SARS-CoV-2 into human host cells[38,39]. The gene expression of all three genes,

ACE2, *TMPRSS-2*, and *NRP-1* increased progressively as CPM+ lung progenitor cells

differentiated from day zero to day 28 in airway epithelium and from day zero to day 10 in

alveolus epithelium (Fig. 2a, b).

SARS-CoV-2 was introduced on day 32 in the top channel of the airway chips and on day 8

in the alveolus chips, respectively. The viral titer in the epithelium channel of the respective

chips increased from day 0 to day 3 post-infection, while no viral replication was detected in

the endothelial channel, consistent with previous reports[40,41] (Fig. 2c). The presence of

intracellular SARS-CoV-2 viral transcripts was also observed at 3dpi (Fig. 2d). SARS-CoV-2

infection resulted in the disruption of tight junctions between infected cells with minimal

decrease in number of ciliated cells (Fig. 2d).

Similarly, SARS-CoV-2 infection in the alveolus epithelium was confirmed by the presence

of intracellular SARS-CoV-2 viral transcripts at 3dpi (Fig. 2e). Interestingly, SARS-CoV-2

infection did not disrupt the tight junctions of the alveolus epithelium, and there was only a

minimal decrease in the GFP expression of AT2 cells (Fig. 2e). The cell number did not

change significantly after SARS-CoV-2 infection in either airway or alveolus epithelium (Fig.

2f, g). Comparatively, the cilia area percentage in mock and SARS-CoV-2 infected airway

chips showed no significant difference (Fig. 2f). The GFP area percentage quantification

revealed an insignificant change in the number of GFP-positive cells (Fig. 2g).

While SARS-CoV-2 infection led to a decrease in the expression of both *SNTN* and *FOXJ1*,

the changes were not statistically significant (Fig. 2h). In alveolus chips, SARS-CoV-2

infection resulted in a significant decrease in the expression of *SFTPC*, whereas the

expression of *SLC34A2* increased after infection (Fig. 2h).

RNA sequencing unveils a dysregulated type I IFN activation and strong chemokine response in alveolus epithelium

In the airway epithelial cells, RNA sequencing at 3 days post-infection revealed the

upregulation of 165 genes and the downregulation of 40 genes. The volcano plot depicted the

activation of innate immune response-related genes, including *IRF*s, *ISG*s, and *IFN*s (Fig. 3a).

Gene ontology Gene ontology (GO) analysis unveiled the upregulation of key biological

processes related to the immune response, particularly the type I IFN pathway. Among the

upregulated pathways, the response to interferon beta, interferon alpha, positive regulation of

interferon beta production, and interferon-gamma (type II IFN pathway) were the most

significant. Additionally, SARS-CoV-2 infection upregulated the MHC Class 1 related

pathway, indicating an overall robust innate immune response in airway epithelium (Fig. 3b).

A robust type I IFN activation, crucial for the host cell's antiviral response, was indicated by

the upregulation of negative regulation of viral genome replication (Fig. 3b). Further pathway

analysis using Reactome indicated the activation of interferon signaling through multiple

processes, confirming SARS-CoV-2 infection (Fig. S3).

In the alveolus epithelium, RNA sequencing following SARS-CoV-2 infection resulted in the

upregulation of 65 genes and the downregulation of 98 genes. The volcano plot indicated

limited upregulation of innate immune response genes compared to the airway epithelium

(Fig. 3c). GO analysis revealed limited activation of the immune response, particularly type I

IFN pathways (Fig. 3d). Analysis of biological processes unveiled a significant activation of

pathways related to immune cell recruitment, such as neutrophil chemotaxis and chemokine-

mediated signaling pathways, which played a major role in the inflammatory response

elicited by alveolus epithelium (Fig. 3d). Reactome analysis did not show a significant

increase in any immunity-related molecular pathways, further indicating limited IFN

activation in alveolus epithelium (Fig. S3). The absence of an innate immune response,

particularly type I IFN activation, resulted in a muted increase in antiviral activities by host

cells.

SARS-CoV-2 infection significantly increased the expression of genes related to innate

immune responses, including *ISG15*, *ISG56*, *IFNL1*, and *IFNB1* in airway epithelium.

Conversely, alveolus epithelium did not exhibit a significant increase in the gene expressions

of *ISG15, ISG56, IFNL1, and IFNB1* after SARS-CoV-2 infection, further confirming the

lack of innate immune activation in alveolus chips (Fig. 3e).

The heatmap analysis revealed the upregulation of key type I IFN genes in airway chips,

while no upregulation was observed in the alveolus epithelium after SARS-CoV-2 infection

(Fig. 3f). Interestingly, the investigation of chemokine-related genes in the alveolus

epithelium after infection showed a robust upregulation of key chemokine-related genes in

alveolus chips (Fig. 3g), possibly responsible for activating chemotaxis pathways, as

indicated in the GO analysis. A similar observation was previously reported in distal *in vivo*

lung samples with SARS-CoV-2 infection[34].

IAV causes heightened cellular damage in both airway and alveolus chips

In the airway chips, the titer of IAV increased from 1 dpi to 3 dpi in the top channel, while no

viral replication was observed in the lower channel (Fig. 4a). Immunofluorescent staining of

the airway chips confirmed IAV infection in epithelial cells. IAV infection resulted in

significant damage to the tight junctions of the airway epithelium and a decrease in

multiciliated cells (Ac-Tub) (Fig. 4b). Basal cells, which are adult stem cells located at the

base of the airway epithelium near the basement membrane, can increase in number and

differentiate into multiciliated cells in the event of an injury[42]. Immunofluorescent staining of

Cytokeratin 5, a marker for basal cells, revealed an increase in the number of basal cells after

infection (Fig. 4b). We also found a co-localization of IAV nucleoprotein antibody with

Cytokeratin-5. Additionally, IAV infection caused substantial damage to the stratified layer

in the airway chips, exposing the basal cells to the apical side of the cell layer (Fig. 4c),

which is a common feature of IAV infection[37]. Quantitative analysis of the immunostaining

images confirmed a decrease in the cell number, layer thickness, and cilia area percentage on

the apical side of the airway epithelium (Fig. 4d).

Like the airway chips, the titer of IAV increased from day 1 to day 3 post-infection in the top

channel of the alveolus chips, with no viral replication observed in the bottom channel (Fig.

4e). Immunofluorescent staining of the alveolus chips revealed IAV infection in epithelial

cells (Fig. 4f). IAV infection resulted in an increase in cell size but left tight junctions intact

(Fig. 4f). IAV caused substantial damage to AT2 cells, resulting in a decrease in GFP-

positive cells. Quantitative analysis of the immunostaining images confirmed a decrease in

cell number and GFP area percentage in the alveolus epithelium (Fig. 4g).

Gene expression of key multiciliated cells markers, *SNTN* and *FOXJ1*, was significantly

reduced after IAV infection in airway chips. While in alveolus chips, IAV infection led to a

significant decrease in key markers for AT2 cells, such as *SFTPC*. AT1 cell markers *AGER*

and *PDPN* were also significantly reduced after IAV infection (Fig. 4h). Interestingly,

SFTPB gene expression did not show any significant changes, while *SLC34A2* exhibited a

significant increase.

IAV indiscriminately activates type I pathway in both airway and alveolus chips

IAV infection resulted in the upregulation of 2271 genes and the downregulation of 3125

genes in airway epithelium, while in alveolus epithelium, IAV infection upregulated 1432

genes and downregulated 930 genes (Fig. 5a, b). Notably, IFNs were upregulated in both

chips. A heatmap comparison further confirmed the upregulation of crucial type I IFN genes,

including *IFNL1, IFNL2, IFNB1, IFNL3, OASL, ISGs, IFITMs, RIGI, IRF7,* and *STAT1* in

both airway and alveolus epithelium (Fig. 5c). Gene ontology analysis revealed the activation

of pathways such as negative regulation of virus replication, defense response against the

virus, and innate immune response, among others, in both chips (Fig. 5e). In line with our

previous observations using immunofluorescence assay (IFA), the GO of downregulated

genes indicated a decrease in cell-cell adhesion and extracellular matrix (ECM) production in

alveolus chips, while significant damage was observed to multiciliated cells and the

cytoskeleton in airway chips after IAV infection (Fig. 5d). These findings were further

validated using RT-PCR to demonstrate the activation of *ISG15*, *IFNL1*, and *IFNB1* in both

chips (Fig. 5e, f). The PCA plot displayed extreme coordinates in both airway and alveolus

epithelium after IAV infection compared with SARS-CoV-2 infection (Fig. 5g).

IAV induces an immune response in endothelial cells

Endothelial cells in both chips were not infected with SARS-CoV-2 or IAV (Fig. 2c, 3a, 3c).

In previous reports, endothelial cell infection has been rarely observed[28,30,40]. Subsequently,

we explored the impact of SARS-CoV-2 infection and IAV on both airway and alveolus

endothelium using RNA sequencing analysis. Interestingly, no significant gene expression

changes were observed in either the airway or alveolus chip endothelium after SARS-CoV-2

infection (Fig. S4a, b). However, after IAV infection, endothelial cells triggered a robust

innate immune response with the upregulation of innate genes, such as *OASs, ISGs,* and

IFITMs, in both airway and alveolus chips (Fig. 6a). GO analysis indicated the activation of

type I interferon and other innate immune response pathways in the endothelium of both the

airway and alveolus chips following IAV infection (Fig. S4c).

IFNB1 inhibits viral replication in alveolus chips

To investigate the impact of stunted IFN activation in alveolus epithelium, SARS-CoV-2

infected chips were cultured for 7 dpi, revealing consistent viral production in the top channel

(Fig. 6b). Notably, IFNB1 is considered one of the therapeutic options and has been shown to

inhibit SARS-CoV-2 replication[43]. Alveolus chips pretreated with IFNB1 showed a

significant inhibition of SARS-CoV-2 replication as compared to the untreated alveolus chips

(Fig. 6c).

Discussion

We developed human airway and alveolus chip models utilizing isogenic iPSCs for the first

time. Using iPSCs, we overcame common shortcomings of immortalized and primary cell

sources including genetic irrelevance and limited supply[31,32]. The usage of an isogenic iPSC

cell line allowed us to investigate and compare the host response between different regions of

human lungs. Our results reveal a differential immune response between the human airway

and alveolus against SARS-CoV-2. We also show that IAV causes multifold cellular damage

and immune response as compared to SARS-CoV-2. Our models unveil the complexities

associated with the immune response between different viruses and regions of human lungs.

By culturing iPSC-derived CPM+ lung progenitor cells on microfluidic devices, we

effectively obtained airway and alveolus on-chip models populated with different types of

airways and alveolus cells respectively. The airway epithelium consisted of multiciliated,

mucus-secreting, and basal cells. While alveolus epithelium consisted of AT2 and AT1-like

cells (Fig. 1b-k). Such cellular composition provides a close resemblance to the *in vivo*

cellular diversity of human lungs[44,45]. Further co-culturing endothelial cells with respective

tissues resulted in a close relatability to the *in vivo* interface of human airways and alveoli as

reported previous MPS studies[27–30]. Multiciliated cells and AT2 cells are the primary targets

of most respiratory viruses including SARS-CoV-2 and IAV infection[39,46,47]. Consequently,

dominant presence of multiciliated and AT2 cells in respective chips underscore its utility to

mimic respiratory viral infections *in vivo*.

iPSC-derived epithelial cells in both chips showed high permissibility to viral infections,

SARS-CoV-2, and IAV (Fig. 2c, d, f and 3a, e). Some previous MPS reports using primary

cells report lack of productive SARS-CoV-2 infection on-chip possibly owing to a lack of

genotypic similarities to the in vivo characteristics of lung epithelial cells[27]. The associated

cellular damage such as damage to ciliated cells and AT2 cells in respective chips was

significantly limited to post-SARS-CoV-2 infection as compared to IAV infection (Fig. 2f, g

and Fig. 4d, g). IAV infection leads to a significant decrease in the cilia area covered and the

GFP area covered infected chips. To our knowledge, no reports with MPS have shown such

comprehensive differences between cytopathies related to two viral infections. It shows the

capability of our models to accurately mimic cytopathies related to different viral infections.

IAV infection inflicted damage to the stratified layer of the airway epithelium, accompanied

by an increased population of basal cells (Fig. 4d), potentially indicating a regenerative phase

following the substantial loss of multiciliated cells[48–50]. Co-localization of IAV NP and

cytokeratin may also indicate the dedifferentiation of infected cells as reported previously[51].

In IAV-infected alveolus epithelium, the size of individual cells increased, while tight

junctions remained intact (Fig. 4f). AT2 cells play a crucial role in regenerating the alveolar

space after lung injury. During this process, AT2 cells may enter a transitional stage

characterized by an increase in cell size, also known as hypertrophy[52–54]. Increased cellular

size may indicate a regenerative phase in the alveolus epithelium post IAV-induced acute

injury. Recapitulation of the post-infection regenerative phase in both airways and alveoli is

reported for the first time in MPS models.

Leveraging the advantages of our unique isogenic iPSCs, we compared immune responses

between different regions of human lungs post-SARS-CoV-2 and IAV infections. Our results

show that SARS-CoV-2 can elicit a differential innate immune response between the human

airways and alveolar tissue (Fig. 3b-f). In airway chips, SARS-CoV-2 upregulates key genes

and pathways related to type I interferon, whereas in alveolus chips, type I interferon was

significantly suppressed. Dysregulated IFN activation after SARS-CoV-2 infection is well

reported[55]. Our system provides platform to understand differences of immune response

between different regions of the human lung, that is not well undestood[25]. The use of iPSCs

overcomes the irregularities related to SARS-CoV-2-related immune responses (low viral

proliferation, uncharacteristically high immune response) reported in previous MPS

reports[27,28].

Interestingly, despite a suppressed type I response, chemokine pathways were still

significantly upregulated in SARS-CoV-2 infected alveolus chip epithelium (Fig. 3g). A

delayed/suppressed IFN1 activation with upregulated chemokine pathways may lead to

uncontrolled viral replication and recruitment of immune cells[34,56,57].

The immune response elicited by both airway and alveolar epithelium to IAV infection was

multifold higher than SARS-CoV-2. IAV-infected airway and alveolus chips showed

upregulation of interferon pathways in both chips indiscriminately with widespread activation

of antiviral and apoptotic pathways (Fig. 5c-g). The lower immune response against SARS-

CoV-2 was presumably caused by a more effective suppression of the innate immune

response, specifically type I interferon response by SARS-CoV-2. Overall, activation of an
overwhelming immune response post-IAV also leads to a greater degree of cellular damage
compared to SARS-CoV-2. The differential immune response against different viruses shows
capability to emulate virus-specific responses isogenic iPSC-derived lung models.
360
We observe no upregulation of antiviral pathways in SARS-CoV-2 infected alveolus chips,
possibly owing to a lack of IFN activation (Fig. 3d). The viral titer of alveolus chips post-
infection until day 7 revealed a maintained peak of viral proliferation (Fig. 6b). Previous
reports provide hypothesis that a dysregulated IFN activation can lead to uncontrolled viral
growth in patients resulting in severe COVID-19[24]. Additionally, IFNB1 pretreatment of
infected alveolus chips inhibited SARS-CoV-2 infection (Fig. 6c). IFNB1 as a therapeutics
option was popular in the peak of COVID-19 pandemic owing to its ability to restrict viral
replication[58]. Inhibition of viral infection by IFNB1 pretreatment not only substantiates the
important role of type I interferon in viral inhibition but also provides a proof of concept of
our model as a potential drug screening tool.
371
SARS-CoV-2 and IAV infection in endothelium were not observed in either the airway or
alveolus chips at 3 dpi (Fig. 2c and Fig. 3a, e). RNA sequencing analysis of the endothelium
in the SARS-CoV-2 infected both chips, revealed lack of immune response related genes
upregulation (Fig. S4). While endothelial cells are rarely shown to be permissive to SARS-
CoV-2 infection[59], it was reported that infected epithelial cells can release factors that can
elicit immune response in the endothelial cells after crossing the porous membrane
mimicking the *in vivo* basement membrane[27,40]. The absence of immune response could be
explained by the limited migration of cytokines from the top to the bottom channel, owing to
less eminent immune response in epithelium. Conversely, RNA sequencing analysis of IAV-
infected airway and alveolus chip demonstrated upregulation of genes and pathways related
to type I interferon and other immune responses. (Fig. 6a). This aligns with prior reports of
immune system activation in endothelial cells post-IAV infection of epithelial cells is
reported previously[30].
385
386
387 **Conclusion**
388

Our iPSC-derived airway and alveolus chips closely replicate the *in vivo* interface of the

human upper airway and alveolar sac. We achieved optimal differentiation of isogenic lung

progenitor cells into airway and alveolar cells in respective chips. Our study reveals a tissue-

specific response to IAV and SARS-CoV-2 infection.

We demonstrate a differential innate immune response triggered by SARS-CoV-2 in our

iPSC-derived airway and alveolus chips, an aspect that cannot be captured using traditional

immortalized or primary cell sources. Furthermore, we provide insights into the conservative

nature of the immune response elicited by SARS-CoV-2 compared to IAV. With the ongoing

advancements in vaccine and drug administration methods, such as nasal delivery, our chips,

derived from identical cell sources, hold great potential for assessing the tissue-specific

effectiveness in future, alongside their utility in therapeutic drug screening.

Materials and Methods

Microfluidic device fabrication

The microfluidic device consists of two Polydimethylsiloxane (PDMS) layers with

microchannels measuring 1 mm × 1 mm, separated by a porous PET membrane (Corning

Falcon® Permeable 0.4 μm Transparent PET Membrane, 353090). The PDMS layers with 1

mm × 1 mm channels were fabricated using a standard soft lithographic technique. A PDMS

prepolymer (Sylgard 184; Dow Corning) at a ratio of 10:1 base to curing agent was cast

against a mold composed of SU-8 2150 (MicroChem) patterns formed on a silicon wafer.

To bond the two PDMS layers to the semipermeable PET membrane, liquid PDMS

prepolymer was applied through spin-coating at 4000 rpm for 60 s directly on a glass slide.

The thin layer of PDMS prepolymer was transferred to the PDMS layers with microchannels

by placing them directly on the glass slide before sandwiching the membrane between the

microchannels. The assembled PDMS layer with the PET membrane was then kept at 4 □ for

48 h to remove air bubbles, after which it was placed in an oven at 50 °C overnight to cure

the PDMS prepolymer.

iPSCs differentiation into NKX1+ lung progenitor cells

SFTPC-GFP knockin reporter iPSCs (B2-3)[36] were differentiated into definitive endodermal

cells in a stepwise manner, followed by anteriorization and ventralization to generate CPM+

lung progenitor cells in the specified medium in each step (Table S3), as previously

described[60]. In brief, iPSCs were initially seeded on Geltrex-coated (Thermo Fisher

Scientific) culture plates in RPMI medium with 2% B-27 supplement (Thermo Fisher

Scientific), containing 100 ng/ml Activin A (API), 1 μM CHIR99021 (Axon Medchem), 50

U/mL penicillin–streptomycin (Thermo Fisher Scientific), and 10 μM Y27632 (LC

Laboratories). On the following day, sodium butyrate (Fujifilm Wako) was added to the

medium to reach a final concentration of 0.25 mM. From Day 2 to Day 6, the cells were

cultured in medium supplemented with 0.125 mM sodium butyrate and without Y27632.

From Day 6 to Day 10 (Step 2), the cells were cultured in DMEM/F12 (Thermo Fisher

Scientific) basal medium with 2% B-27 supplement, 1x GlutaMax (Thermo Fisher Scientific),

50 U/ml penicillin–streptomycin, 0.05 mg/mL L-ascorbic acid (Fujifilm Wako), and 0.4 mM

monothioglycerol (Fujifilm Wako). During this period, the medium was supplemented with

100 ng/mL Noggin (R&D Systems) and 10 μM SB431542 (Fujifilm Wako) to induce

anteriorization. From Day 10 to Day 14 (Step 3), the cells were cultured in the basal medium

supplemented with 3 μM CHIR99021, 20 ng/mL BMP4 (Proteintech), and 0.05 μM all-trans

retinoic acid (Sigma-Aldrich) to induce ventralization.

From Day 14 to Day 21 (Step 4), the cells were cultured in the basal medium supplemented

with 3 μM CHIR99021, 10 ng/mL KGF (PeproTech), 10 ng/mL FGF10 (PeproTech), and 20

μM DAPT (Fujifilm Wako). Subsequently, the cells were labeled with a rat anti-CPM

antibody (in-house) and microbead-conjugated anti-rat immunoglobulin kappa light chain

antibody (#130-047-401, Miltenyi Biotec). This was followed by the isolation of CPM+ lung

progenitor cells using an autoMACS Pro Separator (Miltenyi Biotec) for the subsequent

experiments.

Differentiation on-chip

Airway-on-chip

Both channels of the devices were washed with 70% ethanol and dried at 50 ℃. Following

UV treatment, both channels were rinsed with 200 μl of phosphate-buffered saline (PBS).

Subsequently, the top side of the PET membrane was coated with 100 μl of iMatrix-511 (32

μg/ml) and incubated at 37 □ overnight. The following day, the coating material was

removed by introducing 100 μl of PneumaCult™-ALI Medium (#05001, Stemcell

Technologies) without DAPT into the top channel, along with the addition of 300 μl in the

bottom channel. The excess medium was removed from the top channel outlet reservoir using

an aspirator.

Fresh CPM+ cells were suspended in PneumaCult™-ALI Medium without DAPT at a

density of 2×10^7 cells/ml and introduced into the top channel of each device. After 4 h, an

additional 200 μl of PneumaCult™-ALI Medium without DAPT was directly added to the

top channel. Once confluence was achieved, all the medium from the top channel was

removed, and approximately 30 μl of PneumaCult™-ALI Medium with DAPT was

introduced, along with 300 μl of PneumaCult™-ALI Medium with DAPT in the lower

channel. The devices were then placed in an incubator at 37 ℃ for 28 days, with medium

changes every other day.

The devices were maintained for at least 28 days before the lower channel was washed, and

fibronectin (F4759, Sigma-Aldrich) was introduced as a coating material for endothelial cells.

HUVECs were suspended in EGM2 medium at a density of $1–2 \times 10^7$ cells/ml. After 4 h, the

PneumaCult™-ALI Medium with DAPT and EGM2 medium was added to their respective

channels.

Alveolus-on-chip

The top side of the PET membrane was coated with 100 μl of Geltrex (Thermo Fisher

Scientific) (0.12–0.18 mg/ml) and left to incubate overnight. The following day, the coating

material was removed by introducing 100 μl of alveolarization medium into the top channel,

along with the addition of 300 μl in the bottom channel. Any excess medium was removed

from the top channel outlet reservoir using an aspirator.

The alveolarization medium was prepared as follows: Ham's F12 (Fujifilm Wako)

supplemented with dexamethasone (50 nM), 3-Isobutyl-1-methylxanthine (IBMX) (100 μM),

B27 supplement (1%), BSA (0.25%), HEPES (15 mM), CaCl2 (0.8 mM), ITS premix (0.1%),

8-BrcAMP (100 μM), human KGF (10 ng/ml), CHIR99021 (3 μM), SB431542 (10 μM),

Y27632 (10 μM), and penicillin/streptomycin (50 U/ml) (Table S3). Frozen CPM+ cells,

suspended in the alveolarization medium at a density of 2×10^7 cells/ml, were introduced into

the top channel of each device. After 4 h, alveolarization medium was added directly to the

top channel. Once confluence was achieved, all the medium was removed from the top

channel, and approximately 30 μl of fresh alveolarization medium was introduced. In the

lower channel, 300 μl of alveolarization medium was replenished every alternate day. The

devices were incubated at 37 ℃ for 5 days before the lower channel was washed, and

fibronectin was introduced as a coating material for endothelial cells. HUVECs were

suspended in an EGM2 medium at a density of $1-2 \times 10^7$ cells/ml. After 4 h, alveolarization

medium and EGM2 medium were added to their respective channels.

SARS-CoV-2 and influenza virus

The SARS-CoV-2 isolate (SARS-CoV-2/Hu/DP/Kng/19-027), which was supplied by the
Kanagawa Prefectural Institute of Public Health, was cultured in VeroE6/TMPRSS2 cells.
The viral titers were determined using a 50% tissue culture infectious dose ($TCID_{50}$) method.
The influenza A virus (A/California/04/09 [H1N1]) was propagated and titrated in MDCK
cells through a plaque assay. All experiments involving SARS-CoV-2 were conducted within
a biosafety level 3 containment laboratory at the Institute for Life and Medical Sciences at
Kyoto University.

Viral infection-on-chip

For virus infection on the chip, the top channel was carefully rinsed with PBS, and each
device received 10^3 $TCID_{50}$ of SARS-CoV-2 or 10^3 PFU of IAV in 30 µl of medium.
Following a 1-h incubation at 37 °C, the virus solution was aspirated, and the top channel
was washed twice with PBS. Subsequently, 200 µl of medium was introduced into the upper
channel to sustain the epithelium. The medium in both the top and bottom channels was
changed daily until the devices were either fixed for IFA or the cells were lysed for RNA
examination.

Viral titer experiment

Virus titers were examined using a TCID50 assay for SARS-CoV-2 and a plaque assay for
IAV. Briefly, VeroE6/TMPRSS2 cells were prepared in a 96-well plate one day before
conducting the assay. The virus samples were serially diluted and introduced into the cells,
followed by an incubation at 37 °C for 4 days. The cells were monitored under a microscope
to detect any cytopathic effects. $TCID_{50}$/ml was calculated using the Reed–Muench method.

RNA extraction from chip

For RNA extraction from the single-layer airway or alveolus chips, the top channel was
washed with 200 µl DPBS. After washing, 350 µl cell lysis buffer (Qiagen kits) was
introduced in the top channel in two installments, and cell lysate solution was collected in a

1.5 ml tube. For bilayer chips, the top and bottom channels were washed with 200 µl each.

After washing, 350 µl cell lysis buffer (Qiagen RLT Buffer) was introduced in the bottom

channel in two installments, and cell lysate solution was collected in a 1.5 ml tube. Thereafter,

a similar procedure was performed for the top channel and cell lysate was collected for RNA

extraction and stored at -80 degree Celsius for further analysis.

RT-PCR analysis

Total mRNA was isolated from the cell lysates using a Qiagen RNeasy Micro Kit (Cat. No.

74004). The amount and quality of total RNA were assessed using a Thermo Fisher

NanoDrop spectrophotometer. Subsequently, cDNA was generated from each sample

employing the Prime Script RT Master Mix (Perfect Real Time) (Takara Bio Inc., Japan).

Quantitative RT-PCR was performed using TB Green Premix Ex Taq II (Takara Bio, Inc.,

Japan) in a Applied biosystems Quantstudio 5, and β-actin (ACTB) was used as the

housekeeping gene. The relative expression (fold change) of target genes was quantified

using the double delta C method. The primer sequences used are provided in Table 2.

Immunofluorescence microscopy

The top and bottom channels were washed with PBS before fixation with 4% PFA for 15

minutes. Following fixation, the cells were washed with PBS and permeabilized using 0.2%

Triton X-100 for 15 min. Alternatively, devices were stored after fixation at 4 ℃ for a

maximum of 7 days. A 5% goat serum blocking buffer was added to the permeabilized cells

at room temperature after washing, and primary antibodies in the blocking buffer were

introduced. The samples were stored overnight at 4 ℃or for 2 h at room temperature. A

secondary antibody in blocking buffer was applied to the cells for 40–50 min after washing at

room temperature. DAPI was used to counterstain the cellular nuclei after the secondary

antibody step, and imaging was performed using confocal microscope (Olympus FV-3000,

Tokyo, Japan). Image processing was conducted using ImageJ. A list of antibodies used is

provided in Table 1.

RNA sequencing

The iPSC-derived airway and alveolus chips, with mock and virus infections (both SARS-

CoV-2 and IAV), were utilized for RNA-seq analysis. Cell lysates were collected from both

the top and bottom channels, as described earlier, using Buffer RLT (Qiagen RNeasy Micro

Kit (Cat. No. 74004). RNA extraction was performed from each cell lysate employing a

Qiagen miniprep kit. The RNA quality was analyzed using the 2100 Bioanalyzer by Agilent

Technologies. Subsequently, each library was prepared following the manufacturer's

guidelines with the Illumina TruSeq Stranded mRNA Sample Prep Kit. Sequencing was

conducted using Illumina NextSeq2000, and FASTQ files were generated via bcl2fastq-2.20.

To prepare the data for analysis, adapter sequences and low-quality bases were trimmed from

the raw reads utilizing Cutadapt ver. 4.1. The trimmed reads were then mapped to the human

reference genome sequences (hg38) using STAR version 2.7.10a and a GENCODE (release

32, GRCh38.p13) GTF file.

Raw counts were calculated with htseq-count ver. 0.13.5, employing the GENCODE GTF

file. Subsequently, a differential expression analysis was carried out using DESeq2 in R

studio (R version 4.3.0). Differentially expressed SARS-CoV-2 and IAV genes were

excluded in further analysis. Gene Ontology analysis was conducted using the Database for

Annotation, Visualization, and Integrated Discovery (DAVID 2021, knowledgebase

(v2023q2))[61]. The generation of GO plots was achieved using the ggplot2 R package (version

3.4.2). Log$_2$ fold change values for each gene were utilized to create a heatmap using

GraphPad Prism (GraphPad software).

Image analysis and quantification

Cell number quantification: To quantify nuclei, we utilized 20× magnification DAPI channel

images (636.396 μm × 636.396 μm) from at least 3 regions of interest (ROI) in 3 different

devices. Z-stack images above the membrane level were employed for epithelial nucleus

quantification with ImageJ. Initially, the z-stack images were imported into ImageJ and

stacked. Subsequently, the image type was converted to 8-bit, and threshold was set.

Watershed segmentation was performed to separate boundaries, and then categorized regions

were counted using "Analyze Particles" function in ImageJ. The results were then plotted

using GraphPad Prism (GraphPad software).

Cilia area percentage quantification: The area covered by multiciliated cells were measured

by 60× magnification Alexa Fluor 488 channel images (Ac-Tub) (212.132 μm × 212.132 μm)

from at least 3 ROI in 3 different devices. Cilia are specifically localized on the apical cell

surface. Therefore, for analysis, the first 10 steps of z-stack images were considered. Initially,

the files were imported into ImageJ and z-stacked. Subsequently, the file type was converted

to 8-bit. The threshold was set, and the area was measured using built-in tools in ImageJ.

GFP area percentage quantification: To quantify GFP, we captured 60× magnification EGFP

channel images (212.132 μm × 212.132 μm) from at least 3 ROI in 3 different devices. All

images were adjusted for brightness using the same lookup table (LUT). Z-stack images were

imported into ImageJ and stacked before being converted to 8-bit. The threshold was set, and

the same procedure as for Ac-Tub was followed.

Layer thickness quantification: To determine the height of the cellular layer, 60x

magnification DAPI channel images (212.132 μm × 212.132 μm) from at least 3 ROI in 3

different devices were used. The top z-stack image with DAPI fluorescence intensity and the

last image before the intensity decline were manually selected. Images were acquired with a

z-stack gap of 1 μm, so the total number of z-stacks was calculated to determine the layer

height. The data was plotted on a graph using GraphPad Prism (GraphPad software).

IFNB1 experiment

IFN-beta (300-02BC, Peprotech) was introduced into the top channel at a concentration of 10

ng/ml, 24 h prior to the infection experiment. Subsequent infection experiments were

conducted following a similar protocol as described earlier.

Statistics

A Student's t-test was conducted using GraphPad Prism (GraphPad software) for RT-PCR

data and image analysis (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). For RNA-

seq data, further downstream analysis was carried out using criteria of $P_{adj} < 0.05$ and $\log_2$

fold change >1. Unless otherwise specified in the figure legends and figures, a minimum of 3

devices were used for each statistical comparison.

References

1. Khalbuss, W. E., Laucirica, R. & Pantanowitz, L. Pulmonary Infections. in 121–159 (2011). doi:10.1007/978-1-4614-0242-8_6.

2. Gorbalenya, A. E. *et al.* The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* **5**, 536–544 (2020).

3. Wang, H. *et al.* Modelling the economic burden of SARS-CoV-2 infection in health care workers in four countries. *Nat Commun* **14**, 2791 (2023).

4. Serrano-Ripoll, M. J. *et al.* Impact of viral epidemic outbreaks on mental health of healthcare workers: a rapid systematic review and meta-analysis. *J Affect Disord* **277**, 347–357 (2020).

5. Torres Munguía, J. A., Badarau, F. C., Díaz Pavez, L. R., Martínez-Zarzoso, I. & Wacker, K. M. A global dataset of pandemic- and epidemic-prone disease outbreaks. *Sci Data* **9**, 683 (2022).

6. Sampath, S. *et al.* Pandemics Throughout the History. *Cureus* (2021) doi:10.7759/cureus.18136.

7. Miller, I. F., Becker, A. D., Grenfell, B. T. & Metcalf, C. J. E. Disease and healthcare burden of COVID-19 in the United States. *Nat Med* **26**, 1212–1217 (2020).

8. Short, K. R., Kroeze, E. J. B. V., Fouchier, R. A. M. & Kuiken, T. Pathogenesis of influenza-induced acute respiratory distress syndrome. *Lancet Infect Dis* **14**, 57–69 (2014).

9. Torres Acosta, M. A. & Singer, B. D. Pathogenesis of COVID-19-induced ARDS: implications for an ageing population. *European Respiratory Journal* **56**, 2002049 (2020).

10. Luyt, C.-É., Combes, A., Trouillet, J.-L., Nieszkowska, A. & Chastre, J. Virus-induced acute respiratory distress syndrome: Epidemiology, management and outcome. *Presse Med* **40**, e561–e568 (2011).

11. Krynytska, I., Marushchak, M., Birchenko, I., Dovgalyuk, A. & Tokarskyy, O. COVID-19-associated acute respiratory distress syndrome versus classical acute respiratory distress syndrome (a narrative review). *Iran J Microbiol* (2021) doi:10.18502/ijm.v13i6.8072.

12. Xia, B. *et al.* SARS-CoV-2 envelope protein causes acute respiratory distress syndrome (ARDS)-like pathological damages and constitutes an antiviral target. *Cell Res* **31**, 847–860 (2021).

13. Liu, J. *et al.* SARS-CoV-2 cell tropism and multiorgan infection. *Cell Discov* **7**, 17 (2021).

14. Mokhtari, T. *et al.* COVID-19 and multiorgan failure: A narrative review on potential mechanisms. *J Mol Histol* **51**, 613–628 (2020).

15. Novak, T. *et al.* Transcriptomic profiles of multiple organ dysfunction syndrome phenotypes in pediatric critical influenza. *Front Immunol* **14**, (2023).

16. Tang, B. M., Cootes, T. & McLean, A. S. From Influenza-Induced Acute Lung Injury to Multiorgan Failure. in 449–458 (2019). doi:10.1007/978-3-030-06067-1_35.

17. Diamond, M. S. & Kanneganti, T.-D. Innate immunity: the first line of defense against SARS-CoV-2. *Nat Immunol* **23**, 165–176 (2022).

18. Takeuchi, O. & Akira, S. Pattern Recognition Receptors and Inflammation. *Cell* **140**, 805–820 (2010).

19. Sun, L., Wu, J., Du, F., Chen, X. & Chen, Z. J. Cyclic GMP-AMP Synthase Is a Cytosolic DNA Sensor That Activates the Type I Interferon Pathway. *Science (1979)* **339**, 786–791 (2013).

20. Melchjorsen, J., Sørensen, L. N. & Paludan, S. R. Expression and function of
chemokines during viral infections: from molecular mechanisms to in vivo function. *J*
Leukoc Biol **74**, 331–343 (2003).
21. Khalil, B. A., Elemam, N. M. & Maghazachi, A. A. Chemokines and chemokine
receptors during COVID-19 infection. *Comput Struct Biotechnol J* **19**, 976–988 (2021).
22. Kasuga, Y., Zhu, B., Jang, K.-J. & Yoo, J.-S. Innate immune sensing of coronavirus
and viral evasion strategies. *Exp Mol Med* **53**, 723–736 (2021).
23. Chen, X. *et al.* Host Immune Response to Influenza A Virus Infection. *Front Immunol*
9, (2018).
24. Kim, Y.-M. & Shin, E.-C. Type I and III interferon responses in SARS-CoV-2
infection. *Exp Mol Med* **53**, 750–760 (2021).
25. Alfi, O. *et al.* Human Nasal and Lung Tissues Infected *Ex Vivo* with SARS-CoV-2
Provide Insights into Differential Tissue-Specific and Virus-Specific Innate Immune
Responses in the Upper and Lower Respiratory Tract. *J Virol* **95**, (2021).
26. Gluck, J. P. & Bell, J. Ethical issues in the use of animals in biomedical and
psychopharmocological research. *Psychopharmacology (Berl)* **171**, 6–12 (2003).
27. Thacker, V. V *et al.* Rapid endotheliitis and vascular damage characterize
SARS□CoV□2 infection in a human lung□on□chip model. *EMBO Rep* **22**, (2021).
28. Zhang, M. *et al.* Biomimetic Human Disease Model of SARS□CoV□2□Induced
Lung Injury and Immune Responses on Organ Chip System. *Advanced Science* **8**,
2002928 (2021).
29. Bai, H. *et al.* Mechanical control of innate immune responses against viral infection
revealed in a human lung alveolus chip. *Nat Commun* **13**, 1928 (2022).
30. Si, L. *et al.* A human-airway-on-a-chip for the rapid identification of candidate
antiviral therapeutics and prophylactics. *Nat Biomed Eng* **5**, 815–829 (2021).
31. Marx, V. Cell-line authentication demystified. *Nat Methods* **11**, 483–488 (2014).
32. Beers, M. F. & Moodley, Y. When Is an Alveolar Type 2 Cell an Alveolar Type 2
Cell? A Conundrum for Lung Stem Cell Biology and Regenerative Medicine. *Am J*
Respir Cell Mol Biol **57**, 18–27 (2017).
33. Ortiz, M. E. *et al.* Heterogeneous expression of the SARS-Coronavirus-2 receptor
ACE2 in the human respiratory tract. *EBioMedicine* **60**, 102976 (2020).
34. Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives
Development of COVID-19. *Cell* **181**, 1036-1045.e9 (2020).
35. Konishi, S. *et al.* Directed Induction of Functional Multi-ciliated Cells in Proximal
Airway Epithelial Spheroids from Human Pluripotent Stem Cells. *Stem Cell Reports* **6**,
18–25 (2016).
36. Gotoh, S. *et al.* Generation of Alveolar Epithelial Spheroids via Isolated Progenitor
Cells from Human Pluripotent Stem Cells. *Stem Cell Reports* **3**, 394–403 (2014).
37. Kumar, P. A. *et al.* Distal Airway Stem Cells Yield Alveoli In Vitro and during Lung
Regeneration following H1N1 Influenza Infection. *Cell* **147**, 525–538 (2011).
38. Daly, J. L. *et al.* Neuropilin-1 is a host factor for SARS-CoV-2 infection. *Science*
(1979) **370**, 861–865 (2020).
39. Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e8 (2020).
40. Hashimoto, R. *et al.* SARS-CoV-2 disrupts respiratory vascular barriers by
suppressing Claudin-5 expression. *Sci Adv* **8**, (2022).
41. Nascimento Conde, J., Schutt, W. R., Gorbunova, E. E. & Mackow, E. R.
Recombinant ACE2 Expression Is Required for SARS-CoV-2 To Infect Primary
Human Endothelial Cells and Induce Inflammatory and Procoagulative Responses.
mBio **11**, (2020).

42.   Wu, M., Zhang, X., Lin, Y. & Zeng, Y. Roles of airway basal stem cells in lung
732   homeostasis and regenerative medicine. *Respir Res* **23**, 122 (2022).

733   43.   Rebendenne, A. *et al.* SARS-CoV-2 Triggers an MDA-5-Dependent Interferon
734   Response Which Is Unable To Control Replication in Lung Epithelial Cells. *J Virol* **95**,
735   (2021).

736   44.   Deprez, M. *et al.* A Single-Cell Atlas of the Human Healthy Airways. *Am J Respir*
737   *Crit Care Med* **202**, 1636–1645 (2020).

738   45.   Crystal, R. G., Randell, S. H., Engelhardt, J. F., Voynow, J. & Sunday, M. E. Airway
739   Epithelial Cells: Current Concepts and Challenges. *Proc Am Thorac Soc* **5**, 772–777
740   (2008).

741   46.   Ahn, J. H. *et al.* Nasal ciliated cells are primary targets for SARS-CoV-2 replication in
742   the early stage of COVID-19. *Journal of Clinical Investigation* **131**, (2021).

743   47.   Hou, Y. J. *et al.* SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient
744   in the Respiratory Tract. *Cell* **182**, 429-446.e14 (2020).

745   48.   Kiyokawa, H. *et al.* Airway basal stem cells reutilize the embryonic proliferation
746   regulator, Tgfβ-Id2 axis, for tissue regeneration. *Dev Cell* **56**, 1917-1929.e9 (2021).

747   49.   Tadokoro, T. *et al.* IL-6/STAT3 promotes regeneration of airway ciliated cells from
748   basal stem cells. *Proceedings of the National Academy of Sciences* **111**, (2014).

749   50.   Wu, M., Zhang, X., Lin, Y. & Zeng, Y. Roles of airway basal stem cells in lung
750   homeostasis and regenerative medicine. *Respir Res* **23**, 122 (2022).

751   51.   Robinot, R. *et al.* SARS-CoV-2 infection induces the dedifferentiation of multiciliated
752   cells and impairs mucociliary clearance. *Nat Commun* **12**, 4354 (2021).

753   52.   Bauer, L. *et al.* The pro-inflammatory response to influenza A virus infection is fueled
754   by endothelial cells. *Life Sci Alliance* **6**, e202201837 (2023).

755   53.   Weng, A. *et al.* Lung Injury Induces Alveolar Type 2 Cell Hypertrophy and Polyploidy
756   with Implications for Repair and Regeneration. *Am J Respir Cell Mol Biol* **66**, 564–
757   576 (2022).

758   54.   Ting, C. *et al.* Fatal COVID-19 and Non–COVID-19 Acute Respiratory Distress
759   Syndrome Is Associated with Incomplete Alveolar Type 1 Epithelial Cell
760   Differentiation from the Transitional State without Fibrosis. *Am J Pathol* **192**, 454–467
761   (2022).

762   55.   Acharya, D., Liu, G. & Gack, M. U. Dysregulation of type I interferon responses in
763   COVID-19. *Nat Rev Immunol* **20**, 397–398 (2020).

764   56.   Channappanavar, R. *et al.* Dysregulated Type I Interferon and Inflammatory
765   Monocyte-Macrophage Responses Cause Lethal Pneumonia in SARS-CoV-Infected
766   Mice. *Cell Host Microbe* **19**, 181–193 (2016).

767   57.   Acharya, D., Liu, G. & Gack, M. U. Dysregulation of type I interferon responses in
768   COVID-19. *Nat Rev Immunol* **20**, 397–398 (2020).

769   58.   Monk, P. D. *et al.* Safety and efficacy of inhaled nebulised interferon beta-1a
770   (SNG001) for treatment of SARS-CoV-2 infection: a randomised, double-blind,
771   placebo-controlled, phase 2 trial. *Lancet Respir Med* **9**, 196–206 (2021).

772   59.   Schimmel, L. *et al.* Endothelial cells are not productively infected by SARS☐CoV☐2.
773   *Clin Transl Immunology* **10**, (2021).

774   60.   Yamamoto, Y. *et al.* Long-term expansion of alveolar stem cells derived from human
775   iPS cells in organoids. *Nat Methods* **14**, 1097–1106 (2017).

776   61.   Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and
777   functional annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216–W221
778   (2022).

Acknowledgment

We thank our technical staff (Aki Kubo, Mayumi Morikawa, Megumi Kida, and S A

Nyoman Putri Triantini) for fabricating the devices, cell cultures, and RTPCR experiments.

We also thank Chiho Onishi for performing the infection experiments at the BSL3 facility. Dr.

Takuya Yamamoto, Kazusa Ohkita, and Satoko Sakurai for RNA-seq library preparation and

sequencing, and Dr. Koichi Igura and Koyori Ushitora for supporting the cell preparation.

Funding

This work was supported by JST-CREST (Grant Number JPMJCR20HA), Japan, JSPS Core-

to-Core Program A, Advanced Research Networks (JPJSCCA20190008), Grant for Joint

Research Project of the Institute of Medical Science at the University of Tokyo, and the Joint

Usage/Research Center Program of the Institute for Life and Medical Sciences at Kyoto

University.

Author Contributions

Conceptualization: SY, KF, TN, SG, RY

Funding acquisition: TN, SG, RY

Methodology: SY, YM, TN, SG, RY

Resources: SY, TN, SG, RY, RM

Software: SY, KF

Investigation: SY, TT, YM, ST

Visualization: SY,KF

Supervision: TN, SG, RY

Writing- original draft: SY

Review and editing: SY, KF, TN, SG, RY

Competing interests

All authors have agreed upon the publication of this manuscript. S.G. is one of founders and

shareholders of HiLung Inc. S.G. is one of the inventors of the patents related to the methods

of generating iPSC-derived airway and alveolar epithelial progenitor cells:

WO2014168264A1, WO2016143803A1, and WO2016148307A1. R.Y. is one of the

founders and shareholders of Physios Biotech, Inc.

Data and Material availability

Data supporting the findings of this study are available from the corresponding author upon

reasonable request.

Figures

Figure 1: Development of human iPSC-derived airway chips and alveolus chips

Human iPSC-derived CPM+ lung progenitor cells were seeded in the top channel for further

differentiation into airway and alveolar cells in the respective chips before seeding the

HUVECs on the opposite side of the porous membrane. (a) Schematic illustration of the

airway and alveolus chips (created with BioRender.com). (b) 3D volume view of tight

junctions immunoassayed for ZO-1 and VE-cadherin in the airway epithelium and

endothelium, respectively. (c) Maximum intensity z-stack immunofluorescent staining for

ZO-1 and VE-Cadherin in airway epithelium and endothelium respectively. (d)

Immunofluorescent staining for Ac-Tub, MUC5AC, Cyt 5 in airway epithelium. (e) Gene

expression levels of *SNTN* and *FOXJ1* in the airway epithelium (n = 3, except day 7, n = 2).

(f) 3D volume view of tight junctions immunoassayed for ZO-1 and VE-cadherin in the

alveolus epithelium and endothelium, respectively. (g) Maximum intensity z-stack

immunofluorescent staining for ZO-1 and VE-Cadherin in alveolus epithelium and

endothelium respectively. (h) GFP expression and Immunofluorescence staining of GFP in

the alveolus epithelium. (i) Immunofluorescence staining of SLC34A2 in alveolus epithelium.

(j) Immunofluorescence staining of HT1-56 in alveolus epithelium. (k) Gene expression

levels of *SFTPC*, *SFTPB*, *SLC34A2* in the alveolus epithelium (n = 3). (Immunofluorescence

images were taken on day 32 for airway chips and day 7-9 for alveolus chips).

Figure 2: SARS-CoV-2 infection of airway chips and alveolus chips

SARS-CoV-2 was introduced into the top channel of both chips at a concentration of 10^3

$TCID_{50}$/device, and the devices were cultured for three days. (a) Gene expression of the

SARS-CoV-2 receptors *ACE-2*, *TMPRSS-2*, and *NRP-1* in the airway epithelium measured

on day 0, 7, 21, and 28 (n = 3 chips, except day 7, n = 2). (b) Gene expression of the SARS-

CoV-2 receptors *ACE-2*, *TMPRSS-2*, and *NRP-1* in the alveolus epithelium measured on day

0, 4, 7, and 10 (n = 3 chips). (c) Viral titers in the airway and alveolus chips were measured at

3dpi using a 50% tissue culture infectious dose ($TCID_{50}$) method (n of at least 3 chips). (d)

Intracellular SARS-CoV-2 gene expression in airway chip epithelial cells normalized to

housekeeping gene *ACTB* and immunofluorescent staining for ZO-1, and Ac-Tub in mock

and infected airway epithelium at 3 dpi. (e) Intracellular SARS-CoV-2 gene expression in

alveolus chip epithelial cells normalized to housekeeping gene *ACTB* and immunofluorescent

staining for ZO-1, and GFP in mock and infected alveolus epithelium at 3 dpi. (f) Image

analysis of immunofluorescence staining in mock and infected airway chip samples to

determine the cell number and cilia area percentage (Ac-Tub) (n = at least three ROI from

independent 3 chips). (g) Image analysis of immunofluorescence staining in mock-and

infected alveolus chip samples for cell number and GFP area percentage (n = at least three

ROI from independent 3 chips). (h) Gene expression of key cellular markers *SNTN* and

FOXJ1 in the airway epithelium and *SFTPC* and *SLC34A2* in the alveolus epithelium in

mock and infected conditions at 3 dpi (n = 3).

Figure 3: RNA seq reveals a dysregulated IFN activation and robust chemokine

response in the alveolus epithelium.

(a) Volcano plot of differentially expressed genes between mock and infected airway

epithelium ($\log_2$ fold change > 1 and (P_{adj}) < 0.05, n = 3 chips). (b) GO analysis showing the

most significantly upregulated biological pathways in airway chips (P value shows $-\log_{10}$(P

value). (c) Volcano plot of differentially expressed genes between mock and infected

alveolus epithelium ($\log_2$ fold change > 1 and (P_{adj}) < 0.05, n = 3 chips). (d) GO analysis

showing the most significantly upregulated biological process pathways in the alveolus chips

(P value shows $-\log_{10}$(P value). (e) Expression of IFN-related genes in the airway alveolus

epithelium in mock and infection conditions at 3 dpi (n = 3). (f) $\log_2$ fold change heatmap

showing the expression of important innate immune response genes in the airway and

alveolus epithelium of mock and infected samples (n = 3). (g) Normalized count heatmap of

chemokine gene expression in mock and infected alveolus epithelium on alveolus chips.

876
877
878 **Figure 4: IAV inflicts multifold cellular damage and immune response in both airway**
879 **and alveolus epithelium in respective chips**
880
IAV was introduced into the top channel of both chips at a concentration of 10^3 PFU/device,
and the devices were cultured for three days. (a) The influenza viral titer in the airway chip
was measured at 3 dpi using a plaque-forming assay (n = 3). (b) Immunofluorescence

staining of the IAV N protein, Ac-Tub, ZO-1, and Cytokeratin 5 in mock and infected airway

epithelium in airway chips. (c) Orthogonal view of immunofluorescence staining of Ac-Tub

and Cytokeratin 5 in mock and infected airway epithelium in airway chips. (d) Image analysis

of immunofluorescence staining of mock and infected samples to determine the cell number,

layer thickness, and cilia area percentage (n = at least three ROI from three independent

chips). (e) Influenza viral titer in the alveolus chip measured at 3 dpi using a plaque forming

assay (n = 3). (f) Immunofluorescence staining of Influenza N protein, ZO-1, and GFP in the

mock and infected alveolus epithelium of alveolus chips. (g) Image analysis of

immunofluorescent staining in mock and infected samples for the number of nuclei and GFP

area percentage (n = at least three ROI from independent three chips). (h) Expression of

SNTN, FOXJ1, SFTPC, SFTPB, SLC34A2, AGER, and *PDPN* in respective chips of mock-

and infected samples, (n = 3).

Figure 5: IAV elicits a robust immune response in both airway chip and alveolus chip

unlike SARS-CoV-2

(a) Volcano plot of differentially expressed genes between mock and infected alveolus
epithelium ($\log_2$ fold change > 1 and (P_{adj}) < 0.05, n $= 3$ chips). (b) Volcano plot of
differentially expressed genes between the mock and infected airway epithelium ($\log_2$ fold
change > 1 and (P_{adj}) < 0.05, n $= 3$ chips). (c) $\log_2$ fold change heatmap showing the
expression of important innate immune response genes in the airway and alveolus epithelium
of mock and infected samples (n $= 3$). (d) GO analysis showing significantly upregulated and
downregulated pathways in the respective chips between the mock and infected samples (P
value shows $-\log_{10}$(P value)). (e) Expression of the key IFN genes *ISG15*, *IFNL1*, *IFNB1* in
mock and infected airway epithelium in airway chips (n $= 3$). (f) Expression of the key IFN
genes *ISG15*, *IFNL1*, and *IFNB1* in mock and infected alveolus epithelium in alveolus chips
(n $= 3$). (g) Principal component analysis (PCA) plot of mock and infected airway and
alveolus chips (n $= 3$).

914
915
916

Figure 6: IAV induces innate immune response in endothelial cells

(a) Volcano plot of differentially expressed genes between mock and IAV-infected airway

and alveolus endothelium ($\log_2$ fold change > 1 and (P_{adj}) < 0.05, n = 3 chips). (b) SARS-

CoV-2 viral titer in the alveolus chip was measured at 7 dpi using a plaque-forming assay (n

= 4). (c) SARS-CoV-2 viral titer in the alveolus chips was measured with and without IFNB1

pretreatment after 3 dpi (n = 3 chips; one chip without IFNB1 pretreatment infected with

SARS-CoV-2 was excluded owing failure to replicate the virus).

925
926 **Figure 7: Schematic of the host immune response against SARS-CoV-2 and IAV in**
927 **proximal and distal lung.**

928
In airway chips, SARS-CoV-2 infection induces an upregulation of type I IFN and
chemokine pathways. Conversely, in alveolus chips, the type I IFN response is suppressed,
and only the chemokine pathways are upregulated. In the case of Influenza infection, there is
a significant upregulation of type I interferon pathways in both airway and alveolus chips,
resulting in overwhelming cellular damage in both proximal and distal lung (Created with
BioRender.com).