Genomic-guided conservation actions to restore the most endangered conifer in the Mediterranean Basin

José Carlos del Valle[1]*, Montserrat Arista[1]*, Carmen Benítez-Benítez[1], Pedro Luis Ortiz[1], Francisco J. Jiménez-López[2], Anass Terrab[1], Francisco Balao[1]

[1] *Departamento de Biología Vegetal y Ecología, Universidad de Sevilla, Sevilla, Spain*

[2] *Departamento Biología y Geología, Física y Química Inorgánica, Universidad Rey Juan*

Carlos, Móstoles, Spain

The co-first authors are denoted by an asterisk.

Correspondence: Francisco Balao (fbalao@us.es)

Keywords: Abies nebrodensis, genetic conservation, paternity, SNP-Array, RADseq,

reforestation, selfing , assisted-gene flow.

Abstract

Species with extremely small population sizes are critically endangered due to reduced genetic diversity, increased inbreeding, and the added threat of hybridization. Genomic tools significantly advance conservation by revealing genetic insights into endangered species, notably in monitoring frameworks. Sicilian fir is the most endangered conifer in Europe with only 30 adult trees spread across an 84-hectare area. Using 20,824 SNPs from RAD-seq employing the silver fir genome assembly and a custom 120 SNP-array, we evaluated genetic diversity, mating patterns, and effective population size in adult trees, 118 natural seedlings, and 2,064 nursery seedlings from past conservation actions. We assessed introgression from neighboring non-native fir plantations and established an intra-population assisted gene flow program selecting the most genetically dissimilar individuals and investigating the outcome through simulations. Genomic analysis unveiled significant genetic diversity among adult Sicilian firs, comparable to non-endangered Mediterranean firs with larger populations. However, the genetic diversity of the forthcoming generation declined due to high self-fertilization, leading to marked inbreeding ($F_{is} = 0.38$) and an alarmingly low effective population size ($N_e = 6$). Nursery seedling monitoring revealed similar selfing rates but significant introgression (~50%) from non-native firs. Although intra-population assisted gene flow could help to mitigate genetic loss, it may not alleviate the species vulnerability to imminent environmental challenges, perpetuating the risk of an extinction vortex. Hence, investigating the impact of Sicilian fir population decline and selfing on inbreeding depression, along with exploring the potential of hybrids for genetic load alleviation and future adaptation, is crucial for effective conservation strategies. This study stands as a compelling model for guiding conservation strategies in similarly imperiled species characterized by extremely small populations.

Introduction

Over the past few centuries, many species of animals and plants have become extinct, while many others teeter on the brink of extinction due to habitat loss, fragmentation, and the reduction of population size to inviable levels (Palomares et al. 2012; Benazzo et al. 2017; Humphreys et al. 2019). Species with extremely small populations present a pressing challenge in contemporary biodiversity conservation, referred to as the 'small population paradigm' (Norris 2004). These species face an increased risk of extinction due to a confluence of factors, including reduced genetic diversity, increased inbreeding, vulnerability to stochastic events, and susceptibility to environmental pressures (Frankham 2015; Ralls et al. 2018). Hybridization presents an additional peril, endangering the species integrity and exacerbating the risks for its already impoverished populations (Balao et al. 2015). To persist at short-term without facing these detrimental effects, it is imperative to maintain an effective population size above 100 individuals (i.e., the minimum viable population sensu Frankham et al. 2014). Consequently, genetic management has become an essential tool to identify strategies for preserving their genetic integrity, mitigating inbreeding depression, and fostering the adaptive potential needed for their survival and long-term persistence in their respective ecosystems (Hogg et al. 2022). Furthermore, genomic tools facilitate the monitoring of current conservation status and the assessment of outcomes from conservation actions (Flanagan et al. 2018; Humble et al. 2023).

Assisted gene flow (AGF) replacing natural gene flow by guided-outcrossing can thus be a powerful conservation tool to preserve biodiversity, particularly for populations that are experiencing genetic erosion (Pregler et al. 2023). Restoring gene flow into small, isolated populations can increase genetic diversity and fitness (i.e., genetic rescue) reducing the risk of inbreeding depression and ultimately population extinction and, at the same time, accelerate the adaptive potential of species to environmental modifications such as climate change

(Aitken & Whitlock 2013; Castilla et al. 2019). Despite the benefits and risks of this practice

have been extensively discussed (Frankham 2016; Ralls et al. 2020; Grummer et al. 2022), the

practical use of the AGF in the tree conservation remains largely unexplored (Aitken et al.

2015; Browne et al. 2019).

Conifers, such as pines and spruces, have specific conservation needs due to their slow

growth, long reproductive cycles, low mutation rates, and specific breeding, pollination, and

dispersal systems, requiring a meticulous and sustained approach to population restoration

(Givnish 1980; Savolainen & Pyhäjärvi 2007; De La Torre et al. 2017). *Abies nebrodensis*

(Lojac.) Mattei (Sicilian fir) is the most threatened conifer in the Mediterranean Basin, and

likely one of the most endangered species in the world. It is categorized as "critically

endangered" by the IUCN Red List (Thomas 2017), listed in Appendix I of the Bern

Convention and considered a priority species in Annexes II and IV of the 92/43 EC Habitats

Directive (Code 9220). This species constitutes an example of a highly vulnerable species due

to its very low population size, with only 30-adult trees and 118 seedlings (over 5-yr old)

growing in a sole population over approximately 84 hectares within the integral reserve of

Parco delle Madonie in Sicily (Venturella et al. 1997). Previous studies suggested that

reproductive trees maintain high genetic diversity but typically exhibit reduced seed viability

(Vicario et al. 1995; Vendramin et al. 1996; Parducci et al. 2001; Scialabba 2019). This low

crop viability strongly indicates the existence of high rates of inbreeding. Moreover, fir

species show a pronounced tendency for hybridization (Klaehn & Winieski 1962; Balao et al.

2020). The presumed hybridization between the Sicilian fir and the non-native silver and

Greek firs (*A. alba* and *A. cephalonica*, respectively) from the surrounding areas of the natural

population may adversely impact seed viability and species integrity (Conte & Cristofolini

2003; Scialabba et al. 2005).

Many *in-situ* and *ex-situ* conservation efforts have been implemented over the last two

decades, encompassing breeding programs generating thousands of seedlings via natural

pollination and the removal of non-native fir species from the vicinity of the population

(Venturella et al. 1997; Frascella et al. 2022; Rogatis et al. 2022). However, there is scarce

genetic information available on both the natural occurring and the nursery-propagated

seedlings, hindering the assessment of a viable long-term conservation plan. To address this

limitation, we conducted a genome-wide analysis of Sicilian fir genetic variation and placed

our results in the context of genetic diversity observed in other Mediterranean fir species. We

then developed an SNP-array for genetic monitoring of the Sicilian firs. We evaluated the

genetic diversity and relationships among the 30 adult trees in the natural population. Through

simulations, we designed an AGF program that prioritizes crosses using individuals with

greater genetic dissimilarity, with the goal of ensuring the short-term survival. We determined

the parentage origins of 118 naturally regenerated seedlings, along with 2,064 seedlings

cultivated in the forest nursery 'Piano Noce,' aiming to ascertain outcrossing rates, and

occurrences of self-fertilization.   Finally, we assessed the degree of introgression with

neighboring plantations of non-native firs.

Materials and methods

Abies species sampling and RAD-seq libraries

We used restriction site associated DNA sequencing (RAD-Seq) to identify high-

quality and information-rich SNPs for genotyping of Sicilian firs and detect hybridization

with alien silver and Greek firs. We sampled five individuals from the unique population of

Sicilian fir, 15 individuals from three Italian silver fir populations and 15 individuals from

three populations of Greek fir (Supplementary Material; Table S1). We extracted DNA and

prepared RAD libraries following Balao et al. (2020). Genomic DNA was digested by using

high-fidelity SbfI (New England Biolabs) and the resulting fragments were double barcoded.

The library was sequenced in a separate lane of an Illumina flowcell HiSeq 2500 at the VBCF

NGS Unit (www.vbcf.ac.at/ngs) as 100 bp single-end (SE) reads.

SNP identification and population genomics

Quality filtering and demultiplexing of the RAD-seq library were performed with

deML (Renaud et al. 2015) and STACKS ver. 2.53 (Catchen et al. 2011, 2013, Rochette et al.,

2019). Following this, we aligned each raw read fastq file to the silver fir genome (Mosca et

al 2019), using Bowtie2 (Langmead & Salzberg 201) with --sensitive and --no-unal settings.

Single-best alignments were then sorted and converted from SAM to BAM format using

Samtools v1.10 (Li et al., 2009). To assemble RAD loci, we used the ref_map.pl pipeline

implemented through Stacks v.2.53. Then, we utilized the populations script to export SNP

data into various formats for subsequent analyses. We investigated the nucleotide diversity

(π), heterozygosity (H_s) and inbreeding (F_{IS}) of the Sicilian fir and compared to silver and

Greek firs populations.

OpenArray development for genotyping and hybrid detection

We customized a panel of 120 SNPs by PCR-based OpenArray Technology (Thermo

Fisher Scientific, USA) for genotyping the Sicilian firsamples. Firstly, we selected 100 highly

informative SNPs to assess the genetic structure of the natural population as well as the origin

of the seedling from the forest nursery 'Piano Noce'. These unlinked high-quality SNPs with

a maximum information for genotyping and paternity exclusion were selected using a

combination of Stacks, Plink2 v. 2.00a3.7 (Chang et al., 2015), PopLDdecay 3.40 (Zhang et

al., 2019) based on the following criteria: 'Biallelic', 'No other SNPs within 4 kb', '$r^2 < 0.01$',

'Minor allele frequency [MAF] > 0.05', 'Minimum 1 individual heterozygote', 'Minimum 1

individual homozygote for alternative alleles', '100 bp flanking sequence available on both

sides of the variable position'. To minimize the probability of SNPs being affected by natural

selection, we kept the SNPs in intergenic regions (avoiding 1kb downstream and 2 kb

upstream regions of genes) using SnpEff and SnpSift v4.3.1t (Cingolani et al., 2012).

Secondly, we selected 20 high-quality SNPs (unlinked markers with high MAF) to detect

hybridization between the Sicilian fir and the other firs inhabiting the Madonie Natural Park.

These selected makers followed the same selection criteria previously described and showed

the highest species-specific diagnostic power following the highest loadings in the first two

axes of a Discriminant Analysis of Principal Components (DAPC) performed with the

adegenet package v. 2.1.10 (Jombart, 2008) in R v.4.0.3 (R Core Team 2020). A Principal

Component Analysis (PCA) using the R package *dartR* 2.7.2 (Mijangos et al., 2022) was used

to corroborate the discriminant power of the 20 selected SNPs. We customized a panel of 120

SNPs by PCR-based OpenArray Technology (Thermo Fisher Scientific, USA) for genotyping

Sicilian fir samples. Firstly, we selected 100 highly informative SNPs to assess the genetic

structure of the natural population as well as the origin of the seedling from the forest nursery

'Piano Noce'. These unlinked high-quality SNPs with a maximum information for genotyping

and paternity exclusion were selected using a combination of Stacks, Plink2 v. 2.00a3.7

(Chang et al. 2015), PopLDdecay 3.40 (Zhang et al. 2019) based on the following criteria:

'Biallelic', 'No other SNPs within 4 kb', '$r^2 < 0.01$', 'Minor allele frequency [MAF] > 0.05',

'Minimum 1 individual heterozygote', 'Minimum 1 individual homozygote for alternative

alleles', '100 bp flanking sequence available on both sides of the variable position'. To

decrease the probability of selecting SNPs under selection, we kept the SNPs in intergenic

regions (avoiding 1kb downstream and 2 kb upstream regions of genes) using SnpEff and

SnpSift v4.3.1t (Cingolani et al. 2012). Secondly, we selected 20 high-quality SNPs (unlinked

markers with high MAF) to detect hybridization between the Sicilian fir and the other firs

occurring at the Madonie Regional Natural Park. These selected makers followed the same

selection criteria previously described and showed the highest species-specific diagnostic

power following the highest loadings in the first two axes of a Discriminant Analysis of

Principal Components (DAPC) performed with the *adegenet* package v. 2.1.10 (Jombart &

Ahmed 2011) in R v.4.0.3 (R Core Team 2020). A Principal Component Analysis (PCA)

using the R package *dartR* 2.7.2 (Gruber et al. 2018) was used to corroborate the discriminant

power of the 20 selected SNPs.

SNP-Array validation and Genotyping of plants from the natural population

To investigate the genetic diversity and relatedness of Sicilian fir individuals, we first

collected leaf material for genetic analysis from the 30 adult trees and the 118 seedlings

inhabiting the population. DNA was isolated using the NucleoMag Plant kit (Macherey-

Nagel, Germany) according to the manufacturer's protocol. Before starting the genotyping of

samples, we initially validated the specially developed 120-SNPs arrays for this species. To

do so, we genotyped 24 samples (12 in duplicate) and we assessed replicability and genotype

accuracy, as well as we calculated the genotyping error rate and allele drop-out. Then, we

carried out a PCA to verify that replicates clustered together.

We used the 120 SNPs panel to genotype the individuals in the natural population. We

explored the population genetic structuring of the natural population using Structure v. 2.3.4

(Pritchard et al. 2000). We estimated the number of genetic clusters (K) by assigning

individuals in undefined mixture clusters under a Bayesian framework. We conducted 10

independent runs of 1,000,000 iterations each one, with a burn-in period of 100,000 for each

value of K from 1 to 5. Then, the best K that fit the data was calculated using the Δk method

(Evanno et al. 2005) in Structure Harvester (Earl & VonHoldt 2011). Finally, the effective

population size (N_e) and inbreeding level (F_{IS}) were estimated using the software Colony

v.2.0.6.6 (Jones & Wang 2010).

We inferred the pedigree relationships in the natural population of *A. nebrodensis*,

using Colony2 with the full-likelihood approach. Our analysis incorporated prior information

on parental (all reproductive adults) and maternal genotypes (the closest adult to the seedling).

We assumed monoecy, diploidy, both male and female polygamy, and potential inbreeding.

For the full likelihood calculation, we employed very high precision setting, very long run

length, sibship scaling, updated allele frequencies, and weak sibship priors. The genotyping

error rate (estimated from replicates) was set to 0.01 for all loci. We performed three runs

with different random number seeds to check the reliability of the results. Additionally, the

seedlings with a putative hybrid origin (i.e., with an uncertain genetic origin) were examined

with a PCA using the 20 SNPs designed for the discrimination of fir species.

For the adult trees, we used the *adegenet* package to estimate the genetic diversity at

the individual level by calculating the following coefficients: *HsExp*, standardized

heterozygosity based on the mean expected heterozygosity; *HL*, homozygosity by loci; and

INBR, inbreeding coefficient. Additionally, we used Colony2 to estimate the effective

population size (N_e) and inbreeding level (F_{IS}). To assess whether genetic information could

be used in management strategies to mitigate the alleged low population diversity/inbreeding,

we studied the population genetic structure and relatedness of the 30-adult-trees to design a

genome-informed assisted gene flow (AGF) program. To do so, we used the *adegenet*

package to perform a DAPC using a *K*-mean clustering (*K* = 1-30, the total number of adult

trees). Additionally, we used the *related* package (Pew et al. 2015) to calculate the Ritland

estimator (RIT) of pairwise co-ancestry (Ritland 1996) to infer the genetic relatedness of all

adult trees. Finally, considering the absence of other Sicilian fir populations for outcrossing,

we selected the 12 pairs of adults with the lowest RIT values (i.e., those individuals

genetically more dissimilar) to simulate an artificial population in the AGF conservation

program. We used the function *hybridize* from the *adegenet* package to generate 10 progenies

from each of the 12 proposed crosses. Then, we compared the genetic diversity, inbreeding

and effective size of adults, natural seedlings, the simulated AGF population (n = 120) and the

AGF next generation population, which is composed by the 120 seedlings from the AGF

population, the 118 natural seedlings and 19 seedlings from the nursey originate by

outcrossing (see Results section).

Pedigree analysis of seedlings from the forest nursery

213   We investigated the genetic diversity and pedigree from 2,064 seedlings at the forest

nursery from eight maternal batches from the nursery. To assess a feasible parentage with any

of the 30-adult trees and detect putative hybrids with other alien firs, we genotyped the

samples with the 120 SNP panel and then we conducted paternity tests and determined the

rate of outcrossing, inbreeding and self-fertilization using Colony2, as described above.

Results

Genomic diversity of the Sicilian fir and relatives

220   The RAD-seq of the 35 individuals from populations of Sicilian, silver and Greek firs

produced a final STACKS catalogue a total of 365,582 RAD loci (composed of 34,325,726

genomic sites) with an average coverage per sample ($\pm$ s.d.) of 51.4 $\pm$ 38.5 reads per locus.

After filtering for polymorphic RAD loci with MAF > 0.05, we obtained a final data set with

20,824 SNPs. The raw data was deposited in the NCBI Short Reads Archive (BioProject ID

PRJNA563575).

Estimates of genetic diversity in the populations were notably consistent across the

three fir species (Figure 1). Although the Sicilian fir exhibited the lowest values for π and H_s,

these differences were not statistically significant when compared to the populations of the

other two firs (Kruskal-Wallis tests with $p > 0.05$). The Sicilian fir population showed similar

F_{IS} to those of the silver (0.07 ± 0.014) and Greek (0.06 ± 0.007) firs. Furthermore, it is

noteworthy that the Sicilian fir displayed the highest number of private alleles, with 1029,

surpassing the populations of the other species which ranged from 483 to 728 (Figure 1). The

pairwise F'_{ST} values varied between 0.07 (silver fir pop1- silver fir pop2) and 0.26 (Sicilian fir

– Greek fir pop 2). Finally, the Sicilian fir showed the lowest F'_{ST} pairwise with the silver fir

pop1 (0.14; Table S2).

Development of SNP-Array for conservation of Sicilian fir

We selected 120 high-quality SNPs from the 3,536 SNPs called with MAF > 0.05

present in the five individuals of the Sicilian fir, remaining only 459 SNPs after the quality

filtering. Functional annotations showed that 351 SNPs were in intergenic regions, and 268

SNPs remained after excluding promotors genes regions (46 downstream and 59 upstream).

The 20 SNPs selected to discriminate putative hybrids maximized the separation of the three

fir species (the first two components of the PCA accounted for 42.5% and 20.8% of the total

variation, respectively; Figure S1). The validation of the designed 120 SNP-array revealed a

replicability higher than 99% and the PCA successfully separated samples and clustered

duplicates together. Call rates (i.e., proportion of samples that were assigned a genotype call

compared to the total number of samples) of the analyzed samples ranged from 64.6% to

77.1%. High genotyping failures (> 50%) were detected at 15 loci and were discarded for

downstream analyses.

 Gene diversity and population structure in the natural population

The STRUCTURE analysis showed that the most likely number of genetic groupings

was $K = 2$ (Figure 2) with no clear clustering pattern. The first cluster comprised three adult

trees and 41 out of the 42 seedlings related to one of these trees. The second cluster included

the remaining adult trees and seedlings from the population. The effective population size was

very low (N_e= 6; 3–21; lower and upper confidence interval at 95%) and the level of

inbreeding was moderate (F_{IS}= 0.373).

The results of the paternity assignments disclosed the potential hybrid origin in nine

out of 118 seedlings from the natural population. The genetic origin of three of them could

not be assigned to any of the 30-adult trees occurring in the population, and for the remaining

six seedlings only one parental was assigned. 109 seedlings (92.4%) were identified as

purebred Sicilian firs. Of those, 103 seedlings originated by self-fertilization and only six of

them resulted from outcrossing (only five out of the 30 adult-trees were involved in

outcrossing). We analyzed in the PCA the nine seedlings with a putative hybrid origin,

finding that only one of them was genetically closer to the cloud point of the silver fir and,

therefore, it was confirmed as a hybrid (Figure 3).

For adult trees, the estimations of parental similarity showed mild genetic diversity at

the individual level. The mean value of the standardized observed heterozygosity (*HsExp*)

was 0.74 ± 0.19, varying from 0.32 to 1.13. The individual homozygosity by loci (*HL*) ranged

from 0.50 to 0.86, with an average value of 0.67 ± 0.09. Finally, the inbreeding coefficient

(*INBR*) showed values that ranged from -0.20 to 0.79, with an average value of 0.21 ± 0.23.

DAPC analysis conclusively found three genetic clusters with almost lacking genetic

admixture that were comprised of 12, 11 and seven individuals, respectively (Figure S2A).

Co-ancestry analysis using *RIT* pairwise estimations revealed overt differences of the genetic

relatedness of the 30-adult trees, ranging from -0.3837 to 1.0587 (a matrix of pairwise co-

ancestry among all adult trees is depicted in Figure S2B). Finally, the Mantel test showed

absence of any spatial correlation between geographical distance and genetic diversity from

adult trees (r = -0.06; p > 0.05).

Gene diversity and population structure in seedlings from the forest nursery

We were able to genetically characterize 1,776 of the 2,064 selected seedlings from

the nursery. Paternity tests confidently assigned 1,525 seedlings (85.8%) to one or more

parents of the Sicilian fir. We identified 897 purebred seedlings from this species with a very

high autogamy rate (97.9%), only 19 of them being the result of outcrossing. Additionally, we

identified 879 seedlings as putative hybrids. Finally, the effective population size of the

seedlings in the nursery was very low (N_e= 12; 6–26; lower and upper confidence interval at

95%) and its level of inbreeding was moderately high (F_{IS}= 0.354).

Simulating the effects of the Assisted Gene Flow

RIT values of 12 pairs of adult trees selected to simulate an artificial AGF population

ranged from –0.3837 to –0.2843. The simulated AGF population from the selected crosses

showed higher H_o (Kruskal-Wallis's test, χ^2 = 91.882, df = 3, $p < 0.001$) and lower

inbreeding coefficient (χ^2 = 179.82, df = 3, $p < 0.001$) than the natural population (i.e.,

adults and seedlings) and the next generation population (including natural seedlings,

seedlings from the simulated AGF event, and the 19 outbreed seedlings from the nursery).

This latter population maintained similar values than current adults in the population whereas

natural seedlings showed significatively the lowest H_o and the highest F_{IS} (Figure 4). The

nucleotide diversity was similar for all the groupings (χ^2 = 1.984, df = 3, $p > 0.05$). Natural

seedlings showed the lowest N_e (95% IC 2.6-2.9) and the adults the highest one (18-25.2).

AGP seedlings and the Next-Gen population displayed intermediate N_e values (Figure 4).

Discussion

Differential genetic diversity across age cohorts of the Sicilian fir

The genome-wide levels of genetic diversity among adult individuals in the Sicilian fir

population were comparable to those of the silver and Greek firs populations along their

native ranges. The significant genetic variation observed in adult Sicilian fir samples,

highlighted in prior studies (Vicario et al. 1995; Ducci et al. 1999; Conte et al. 2004),

correlated with the recent demographic dynamics of the Sicilian fir, as evidenced by the

increased private alleles, slower heterozygosity response, and stable inbreeding coefficients

(Llorens et al. 2018). Palaeobotanical and humanistic studies revealed the massive presence of

firs along Sicily from Paleogene to Holocene (Tinner et al. 2016), along with a significant

demographic decline since the Middle Ages (Pasta et al. 2020). Anthropogenic pressures

nearly decimated the Sicilian fir forests, driving the species to the verge of extinction, with

only a few surviving individuals comprising the current population (Venturella et al. 1997).

The extended lifespan of firs (> 150 years) and adaptive selection against homozygous

individuals could counteract the effects of demographic decline on genetic diversity (Yao et

al. 2007; Fraser et al. 2014; Su et al. 2018).

Despite the high genetic diversity observed in adult trees, genetic monitoring of

seedlings in the natural population using a developed SNP-array revealed a concerning

scenario for species conservation. The seedlings exhibited a substantial decrease in

heterozygosity and an increased inbreeding due to exceptionally high selfing rates (> 94%).

Only six outcrossed seedlings were found, which were traced back to just five parental

genotypes. This contrasts sharply with the typically high levels of outcrossing observed in

Mediterranean firs and other gymnosperms (Restoux et al. 2008), and it is likely influenced

by several factors such as the sparsely tree dispersion, with an average distance of 621.7

meters between them, and a limited pollen dispersal as occurs in other endangered

Mediterranean firs (Arista & Talavera 1994; Sánchez-Robles et al. 2014). The high selfing

rate and subsequent decline in genetic diversity across young cohorts of the Sicilian fir likely

account for the previously reported reduced seed and seedling viability (Conte et al. 2004;

Scialabba 2019; Mirabile et al. 2023). This reduced genetic diversity may also increase the

species' vulnerability to emerging environmental challenges, particularly those associated to

climate change (Frankham 2015; Ralls et al. 2020).

Remnants of recent hybridization in Sicilian fir population and implications in the

conservation

It is remarkable that, despite a high level of selfing, at least one seedling in the Sicilian

fir population likely derived from hybridization with an alien firs species. Previous

conservation efforts identified more than 5,000 such firs in reforested areas located within a

few hundred meters of the Sicilian fir population (Ducci 2014). While many of these non-

native firs were removed, some still persist, presenting a real risk of genetic exchange,

particularly the silver fir, the ancestral species that gave rise to the Sicilian fir (Balao et al.

2020). Hybridization is a widespread phenomenon present in evolutionary history of

Mediterranean firs (Krajmerová et al. 2016; Balao et al. 2020). While hybridization can

potentially harm endangered species (Balao et al. 2015), understanding the consequences of

hybridization requires detailed study (Arnold 2015; Gompert & Buerkle 2016). Notably, two

adult Sicilian firs exhibited indications of coancestry with silver firs in the PCA, suggesting

historical introgression of anthropical origin due to plantations from at least one century as

occurred in other tree species (Vanden Broeck et al. 2005; Meirmans et al. 2014; Scotti-

Saintagne et al. 2023). Considering that such introgressed trees could potentially enhance the

adaptability towards prevailing ecological stress (Kormutak et al. 2013; Stejskal et al. 2016),

it is cautious to prioritize their conservation as part of the Sicilian fir recovery program

(Jackiw et al. 2015; VonHoldt et al. 2022; Brauer et al. 2023).

Monitoring ex-situ regenerated seedlings with SNP-array confirms the prevalence of

hybridization and selfing

349   Similarly, SNP-array analyses of nursery-grown seedlings revealed a 50% prevalence

of hybridization, contrasting starkly with the minimal 1-in-118 ratio in natural seedlings. This

raises questions about the viability of hybrids in the wild (Campbell & Waser 2001). Clearly,

this prevalence of hybrids in the nursery may compromise the implementation of reforestation

actions. However, it seems worthwhile to delve deeper into investigating the fitness and

adaptability of these hybrids in their natural habitat, considering various environmental

factors.

356   Additionally, approximately 98% of the purebred Sicilian fir seedlings in the forest

nursery originated from self-pollination, mirroring the selfing rates observed in the wild.

While the effective population size (N_e) in the nursery population, established from cones

collected from adult trees, slightly surpassed that of the wild population, this discrepancy

likely stemmed from the inclusion of hybrid seedlings. However, the strikingly low ratio of

effective population size to census size (N_e/N=0.0058) highlights an alarming genetic

bottleneck. This compromises the genetic robustness of the forest restoration stock, akin to

endangered conifers such as the Boise araucaria or the Yellow Box (Kettle et al. 2008;

Broadhurst 2013).

Implementing within-population AGF as the last chance to avoid extinction

In a similar vein, the Sicilian fir population exhibited a notably low effective

population size (N_e=3-21), falling significantly below the recommended threshold (> 100),

which poses considerable challenges for its short-term conservation efforts (Frankham et al.

2014). The high selfing rate, coupled with the low mutation rate in firs (De La Torre et al.

2017), indicates that relying solely on spontaneous mutations will not sufficiently enhance the

population genetic diversity in the short- or medium- term, emphasizing the urgent need for

proactive genetic management strategies (Ralls et al. 2018). Consequently, genetic rescue,

through upcoming outcrossing events with genetically diverse individuals, stands as the final

opportunity to avert extinction by mitigating inbreeding and reducing levels of homozygosity

(Ingvarsson 2001; Fady et al. 2020). Given the single remaining wild Sicilian fir population,

we propose a within-population AGF approach, selecting crosses among trees with more

distant co-ancestry in the population. This method appears to be effective in enhancing

genetic diversity and *N*e, reducing inbreeding, and simultaneously mitigating potential

negative effects, such as outbreeding and the introduction of maladaptive alleles associated

with AGF (Aitken & Whitlock 2013; Grummer et al. 2022). However, the gain in genetic

diversity through AGF restoration appears insufficient to recover past diversity, and the long-

term effectiveness of this conservation action to avoid the extinction vortex remains

uncertain. Hence, there is a necessity to investigate the impact of Sicilian fir population

decline and selfing on inbreeding depression (Pečnerová et al. 2023), along with exploring the

potential of hybrids to alleviate genetic load and facilitate future adaptation (vonHoldt et al.

2018).

Conclusions

Genomic data revealed that adult Sicilian firs harbor high genetic diversity despite their small population size, comparable even to that observed in other Mediterranean firs spread across large populations. Despite this high genetic richness found in adult trees, the genetic variability of the population will dramatically decrease in the upcoming generations, even if human-mediated actions are taken to increase, or at least maintain, the genetic diversity of the population. Intra-population assisted gene flow can contribute to preventing the loss of genetic variation in the population; however, this does not seem to be sufficient to break free from the extinction spiral because of the predictable species' vulnerability in upcoming generations to emerging environmental challenges, such as the climate change. Interspecific crosses with other intersectional fir species, namely the silver fir, could represent a potential alternative strategy for the future conservation of the Sicilian fir that merits exploration. Our results emphasize the relevance of genomic-guided conservation actions, which can assist in identifying suitable individuals for reforestation and provide a solid foundation for conservation management. We expect that this approach could serve as a model to be replicated in the conservation strategies for other endangered firs, or even for other conifers, around the Mediterranean region.

Acknowledgments

The authors acknowledge the Ente Parco delle Madonie and the forest nursery of Piano Noce (Polizzi Generosa) for hosting the project area and maintaining *A. nebrodensis* seedlings. We also thank University of Seville Biology and Herbarium General Research Services (CITIUS) for providing the facilities. This research was funded by the European Union through the LIFE4FIR project, grant number LIFE18/NAT/IT/000164.

Credit authorship contribution statement

JC del Valle: Methodology, Investigation, Writing – original draft. **M Arista**: Conceptualization, Methodology, Investigation, Funding acquisition; **C Benítez-Benítez**: Methodology, Investigation; **PL Ortiz**: Conceptualization, Methodology, Investigation; **FJ Jiménez-López**: Methodology, Investigation; **A Terrab**: Conceptualization, Methodology, Investigation; **Francisco Balao**: Conceptualization, Methodology, Investigation, Writing – original draft. All authors have read and agreed to the published version of the manuscript.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Aitken SN, Bemmels JB, Sally Aitken CN. 2015. Time to get moving: assisted gene flow of forest trees. Evolutionary Applications **9**:271-290

Aitken SN, Whitlock MC. 2013. Assisted gene flow to facilitate local adaptation to climate change. Annual Review of Ecology, Evolution, and Systematics **44**:367–388.

Arista M, Talavera S. 1994. Phenology and anatomy of the reproductive phase of *Abies*

pinsapo Boiss (Pinaceae). Botanical Journal of the Linnean Society **116**:223–234.

Arnold ML. 2015. Divergence with Genetic Exchange. Page Divergence with Genetic

Exchange. Oxford University Press, Oxford, UK.

Balao F, Casimiro-Soriguer R, García-Castaño JL, Terrab A, Talavera S. 2015. Big thistle

eats the little thistle: Does unidirectional introgressive hybridization endanger the

conservation of *Onopordum hinojense*? New Phytologist **206**:448–458. Blackwell

Publishing Ltd.

Balao F, Lorenzo MT, Sánchez-Robles JM, Paun O, García-Castaño JL, Terrab A. 2020.

Early diversification and permeable species boundaries in the Mediterranean firs. Annals

of Botany **125**:495–507.

Benazzo A et al. 2017. Survival and divergence in a small group: The extraordinary genomic

history of the endangered Apennine brown bear stragglers. Proceedings of the National

Academy of Sciences of the United States of America **114**:E9589–E9597.

Brauer CJ, Sandoval-Castillo J, Gates K, Hammer MP, Unmack PJ, Bernatchez L,

Beheregaray LB. 2023. Natural hybridization reduces vulnerability to climate change.

Nature Climate Change **13**:282–289

Broadhurst LM. 2013. A genetic analysis of scattered Yellow Box trees (Eucalyptus

melliodora A.Cunn. ex Schauer, Myrtaceae) and their restored cohorts. Biological

Conservation **161**:48–57.

Browne L, Wright JW, Fitz-Gibbon S, Gugger PF, Sork VL. 2019. Adaptational lag to

temperature in valley oak (Quercus lobata) can be mitigated by genome-informed

assisted gene flow. Proceedings of the National Academy of Sciences of the United

States of America **116**:25179–25185.

Campbell DR, Waser NM. 2001. Genotype-by-environment interaction and the fitness of

plant hybrids in the wild. Evolution **55**:669–676.

Castilla AR, Garrote PJ, Żywiec M, Calvo G, Suárez-Esteban A, Delibes M, Godoy JA, Picó |

F Xavier, Fedriani JM. 2019. Genetic rescue by distant trees mitigates qualitative pollen

limitation imposed by fine-scale spatial genetic structure. Molecular Ecology **28**:4363.

Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-

generation PLINK: Rising to the challenge of larger and richer datasets. GigaScience

4:1–16.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM.

2012. A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;

iso-2; iso-3. Fly **6**:80–92.

Conte L, Cotti C, Schicchi R, Raimondo PM, Cristofolini G. 2004. Detection of ephemeral

genetic sub-structure in the narrow endemic *Abies nebrodensis* (Lojac.) Mattei

(Pinaceae) using RAPD markers. Plant Biosystems **138**:279–289.

Conte L, Cristofolini G. 2003. Assessment of RAPD variation in *Abies nebrodensis* (Lojac.)

Mattei (Pinaceae) using haploid tissue analysis. Israel Journal of Plant Sciences **51**:199–

206.

De La Torre AR, Li Z, Van de Peer Y, Ingvarsson PK. 2017. Contrasting Rates of Molecular

Evolution and Patterns of Selection among Gymnosperms and Flowering Plants.

Molecular Biology and Evolution **24**:1586–1591.

Ducci F. 2014. Species restoration through dynamic ex situ conservation: *Abies nebrodensis*

as a model in Genetic considerations in ecosystem restoration using native tree species.

State of the World's Forest Genetic Resources – Thematic Study. Bozzano M., Jalonen

R., Thomas E., Boshier D., Gallo L., Cavers S., Bordács S., Smith P., Loo J. eds,, p. 225-

233.

Ducci F, Proietti R, Favre JM. 1999. Allozyme assessment of genetic diversity within the relic

476   Sicilian fir Abies nebrodensis (Lojac.) Mattei. Annals of Forest Science **56**:345–355.

477   Earl DA, VonHoldt BM. 2011. STRUCTURE HARVESTER: a website and program for

478   visualizing STRUCTURE output and implementing the Evanno method. Conservation

479   Genetics Resources **4**:359–361.

480   Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals using

481   the software structure: a simulation study. Molecular Ecology **14**:2611–2620.

482   Fady B, Aravanopoulos F, Benavides R, González-Martínez S, Grivet D, Lascoux M, Lindner

483   M, Rellstab C, Valladares F, Vinceti B. 2020. Genetics to the rescue: managing forests

484   sustainably in a changing world. Tree Genetics and Genomes **16**.

485   Flanagan SP, Forester BR, Latch EK, Aitken SN, Hoban S. 2018. Guidelines for planning

486   genomic assessment and monitoring of locally adaptive variation to inform species

487   conservation. Evolutionary Applications **11**:1035–1052.

488   Frankham R. 2015. Genetic rescue of small inbred populations: meta-analysis reveals large

489   and consistent benefits of gene flow. Molecular Ecology **24**:2610–2618.

490   Frankham R. 2016. Genetic rescue benefits persist to at least the F3 generation, based on a

491   meta-analysis. Biological Conservation **195**:33–36.

492   Frankham R, Bradshaw CJA, Brook BW. 2014. Genetics in conservation management:

493   Revised recommendations for the 50/500 rules, Red List criteria and population viability

494   analyses. Biological Conservation **170**:56–63.

495   Frascella A et al. 2022. Innovative In Situ and Ex Situ Conservation Strategies of the

496   Madonie Fir Abies nebrodensis. Sustainability **14**:12643.

Fraser DJ, Debes P V, Bernatchez L, Hutchings JA. 2014. Population size, habitat

fragmentation, and the nature of adaptive variation in a stream fish. Proceedings of the

Royal Society B: Biological Sciences **281**.

Givnish T. 1980. Ecological constraints on the evolution of breeding systems in seed plants:

dioecy and dispersal in gymnosperms. Evolution **34**:959–972.

Gompert Z, Buerkle CA. 2016. What, if anything, are hybrids: enduring truths and challenges

associated with population structure and gene flow. Evolutionary Applications **9**:909–

923.

Gruber B, Unmack PJ, Berry OF, Georges A. 2018. dartr: An r package to facilitate analysis

of SNP data generated from reduced representation genome sequencing. Molecular

Ecology Resources **18**:691–699.

Grummer JA, Booker TR, Matthey-Doret R, Nietlisbach P, Thomaz AT, Whitlock MC. 2022.

The immediate costs and long-term benefits of assisted gene flow in large populations.

Conservation Biology **36**:e13911.

Hogg CJ, Ottewell K, Latch P, Rossetto M, Biggs J, Gilbert A, Richmond S, Belov K. 2022.

Threatened Species Initiative: Empowering conservation action using genomic

resources. Proceedings of the National Academy of Sciences of the United States of

America **119**.

Humble E et al. 2023. Conservation management strategy impacts inbreeding and mutation

load in scimitar-horned oryx. Proceedings of the National Academy of Sciences of the

United States of America **120**.

Humphreys AM, Govaerts R, Ficinski SZ, Nic Lughadha E, Vorontsova MS. 2019. Global

dataset shows geography and life form predict modern plant extinction and rediscovery.

Nature Ecology and Evolution **3**:1043–1047.

Ingvarsson PK. 2001. Restoration of genetic variation lost - The genetic rescue hypothesis.

522   Trends in Ecology and Evolution **16**:62–63.

523   Jackiw RN, Mandil G, Hager HA. 2015. A framework to guide the conservation of species

524   hybrids based on ethical and ecological considerations. Conservation Biology **29**:1040–

525   1051.

526   Jombart T, Ahmed I. 2011. adegenet 1.3-1: new tools for the analysis of genome-wide SNP

527   data. Bioinformatics **27**:3070–1.

528   Jones OR, Wang J. 2010. COLONY: A program for parentage and sibship inference from

529   multilocus genotype data. Molecular Ecology Resources **10**:551–555.

530   Kettle CJ, Ennos RA, Jaffré T, Gardner M, Hollingsworth PM. 2008. Cryptic genetic

531   bottlenecks during restoration of an endangered tropical conifer. Biological Conservation

532   **141**:1953–1961.

533   Klaehn FU, Winieski JA. 1962. Interspecific hybridization in the genus Abies.

534   Kormutak A et al. 2013. Artificial hybridization of some *Abies* species. Plant Systematics and

535   Evolution **299**:1175–1184.

536   Krajmerová D, Paule L, Zhelev P, Voleková M, Evtimov I, Gagov V, Gömöry D. 2016.

537   Natural hybridization in eastern-Mediterranean firs: The case of *Abies borisii-regis*.

538   Plant Biosystems **150**:1189–1199.

539   Llorens TM, Ayre DJ, Whelan RJ. 2018. Anthropogenic fragmentation may not alter pre-

540   existing patterns of genetic diversity and differentiation in perennial shrubs. Molecular

541   Ecology **27**:1541–1555.

542   Meirmans PG, Gros-Louis MC, Lamothe M, Perron M, Bousquet J, Isabel N. 2014. Rates of

543   spontaneous hybridization and hybrid recruitment in co-existing exotic and native mature

544   larch populations. Tree Genetics and Genomes **10**:965–975.

Mirabile G, Cirlincione F, Venturella G, Torta L. 2023. Seed vitality and fungal

contamination in Abies nebrodensis. Plant Biosystems **157**:112–118.

Norris K. 2004. Managing threatened species: the ecological toolbox, evolutionary theory and

declining-population paradigm. Journal of Applied Ecology **41**:413–426.

Palomares F, Godoy JA, López-Bao JV, Rodríguez A, Roques S, Casas-Marce M, Revilla E,

Delibes M. 2012. Possible Extinction Vortex for a Population of Iberian Lynx on the

Verge of Extirpation. Conservation Biology **26**:689–697.

Parducci L, Szmidt AE, Madaghiele A, Anzidei M, Vendramin GG. 2001. Genetic variation

at chloroplast microsatellites (cpSSRs) in *Abies nebrodensis* (Lojac.) Mattei and three

neighboring Abies species. Theoretical and Applied Genetics **102**:733–740.

Pasta S, Sala G, La Mantia T, Bondì C, Tinner W. 2020. The past distribution of *Abies*

nebrodensis (Lojac.) Mattei: results of a multidisciplinary study. Vegetation History and

Archaeobotany **29**:357–371.

Pew J, Muir PH, Wang J, Frasier TR. 2015. related: An R package for analysing pairwise

relatedness from codominant molecular markers. Molecular Ecology Resources **15**:557–

561.

Pregler KC, Obedzinski M, Gilbert-Horvath EA, White B, Carlson SM, Garza JC. 2023.

Assisted gene flow from outcrossing shows the potential for genetic rescue in an

endangered salmon population. Conservation Letters **16**.

Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using

multilocus genotype data. Genetics **155**:945–959.

Ralls K, Ballou JD, Dudash MR, Eldridge MDB, Fenster CB, Lacy RC, Sunnucks P,

Frankham R. 2018. Call for a Paradigm Shift in the Genetic Management of Fragmented

Populations. Conservation Letters **11**.

Ralls K, Sunnucks P, Lacy RC, Frankham R. 2020. Genetic rescue: A critique of the evidence

supports maximizing genetic diversity rather than minimizing the introduction of

putatively harmful genetic variation. Biological Conservation **251**:108784.

Restoux G, Silva DE, Sagnard F, Torre F, Klein E, Fady B. 2008. Life at the margin: The

mating system of Mediterranean conifers. Web Ecology **8**:94–102.

Ritland K. 1996. Estimators for pairwise relatedness and individual inbreeding coefficients.

Genetical Research **67**:175–185.

Rogatis A De, Ducci F, Guerri S, De Rogatis A, Ducci F, Guerri S, Teani A, Proietti R. 2022.

Genotyping ex situ trees of *Abies nebrodensis* translocated from the original Sicilian

population to enrich the gene pool. Journal of Forestry Research **34**:1095–1106.

Sánchez-Robles JMJM, García-Castaño JLJL, Balao F, Terrab A, Navarro-Sampedro L,

Tremetsberger K, Talavera S. 2014. Effects of tree architecture on pollen dispersal and

mating patterns in *Abies pinsapo* Boiss. (Pinaceae). Molecular Ecology **23**:6165–6178.

Savolainen O, Pyhäjärvi T. 2007. Genomic diversity in forest trees. Current Opinion in Plant

Biology **10**:162–167.

Scialabba A. 2019. Seed germination in *Abies nebrodensis* (Pinaceae). Flora Mediterranea

29:272–276.

Scialabba A, Schicchi R, Cordi R, others. 2005. Monitoraggio e analisi del polline. Pages 94–

98 Rendiconto sul progetto LIFE Natura" Conservazione in situ ed ex situ di Abies

nebrodensis (Lojac.) Mattei. ITA.

Scotti-Saintagne C, de Sousa Rodrigues A, Roig A, Fady B. 2023. A comprehensive strategy

for the conservation of forest tree genetic diversity: an example with the protected Pinus

nigra subsp. salzmannii (Dunal) Franco in France. Conservation Genetics.

https://doi.org/10.1007/s10592-023-01581-8

Stejskal J, Horák J, Typta J. 2016. Effect of hybridization in the firs: artificial hybridization

may lead to higher survival rate. European Journal of Forest Research **135**:1097–1105.

Su J, Yan Y, Song J, Li J, Mao J, Wang N, Wang W, Du FK. 2018. Recent fragmentation

may not alter genetic patterns in endangered long-lived species: Evidence from *Taxus*

cuspidata. Frontiers in Plant Science **9**:1571.

Thomas P. 2017. Abies nebrodensis. The IUCN Red List of Threatened Species:2012–2017.

Tinner W et al. 2016. Holocene vegetation and fire history of the mountains of Northern

Sicily (Italy). Vegetation History and Archaeobotany **25**:499–519.

Vanden Broeck A, Villar M, Van Bockstaele E, Van Slycken J. 2005. Natural hybridization

between cultivated poplars and their wild relatives: Evidence and consequences for

native poplar populations. Annals of Forest Science **62**:601-613.

Vendramin GG, Michelozzi M, Lelli L, Tognetti R. 1996. Genetic variation in *Abies*

nebrodensis: a case study for a highly endagered species. Forest Genetics **2**:171–175.

Venturella G, Mazzola P, Raimondo F. 1997. Strategies for the conservation and restoration

of the relict population of *Abies nebrodensis* (Lojac.) Mattei. Bocconea **7**:417–425.

Vicario F, Vendramin GG, Rossi P, Liò P, Giannini R. 1995. Allozyme, chloroplast DNA and

RAPD markers for determining genetic relationships between *Abies alba* and the relic

population of *Abies nebrodensis*. Theoretical and Applied Genetics **90**:1012–1018.

vonHoldt BM, Brzeski KE, Wilcove DS, Rutledge LY. 2018. Redefining the role of

admixture and genomics in species conservation. Conservation Letters **11**.

VonHoldt BM, Hinton JW, Shutt AC, Murphy SM, Karlin ML, Adams JR, Waits LP, Brzeski

KE. 2022. Reviving ghost alleles: Genetically admixed coyotes along the American Gulf

Coast are critical for saving the endangered red wolf. Science Advances **8**:eabn7731.

Yao X, Ye Q, Kang M, Huang H. 2007. Microsatellite analysis reveals interpopulation

differentiation and gene flow in the endangered tree *Changiostyrax dolichocarpa*

(Styracaceae) with fragmented distribution in central China. New Phytologist **176**:472–

480.

Zhang C, Dong SS, Xu JY, He WM, Yang TL. 2019. PopLDdecay: A fast and effective tool

for linkage disequilibrium decay analysis based on variant call format files.

Bioinformatics **35**:1786–1788.

Figure Legends

Figure 1. Violin plots display genetic diversity estimators based in 20,824 SNPs for silver

(orange), Greek (light yellow), and Sicilian (green) firs. Metrics include nucleotide diversity

(π), heterozygosity (Hs), private alleles, and inbreeding (F_{IS}).

Figure 2. Individual ancestry at $K=$ 2, the most supported numbers of clusters in Structure

analysis, for the 30 adult trees and the 118 seedlings in the Sicilian fir population. Adults are

sorted by proximity and arrows show seedlings origin from mother plants.

Figure 3. PCA plot displaying nine seedlings of uncertain origin from the natural Sicilian fir

population and other alien firs species. Colored dots represent Sicilian (green), silver (light

yellow), and Greek (orange) firs, and seedlings with ambiguous origins (red). Darker colors

highlight two adult trees and one seedling showing introgression.

Figure 4. Genetic population parameters based in 120 SNP-array for the adult Sicilian firs

(red) and seedlings (orange) from the natural population, the simulated AGF population

(green), and the AGF-next-generation population (blue). Violin plots display the distribution

of homozygosity (H_o), inbreeding (F_{IS}), nucleotide diversity (π), and effective size population

(N_e).

Table 1. Results of the genetic diversity estimations calculated for the 30 adult trees of the Sicilian fir.

HsExp (standardized heterozygosity based on the mean expected heterozygosity), *HL* (homozygosity

by loci), and *INBR* (inbreeding coefficient).

Adult tree	*HsExp*	*HL*	*INBR*
01M	0.60	0.74	0.25
02M	0.76	0.65	0.15
04M	0.72	0.67	0.25
06M	0.69	0.68	0.20
07M	0.69	0.69	0.20
08M	0.79	0.64	0.57
09M	0.76	0.66	0.12
10M	0.90	0.61	0.17
11M	0.83	0.61	0.11
12M	0.93	0.57	-0.01
13M	1.13	0.50	-0.20
14M	0.93	0.59	0.02
15M	0.74	0.67	0.08
16M	0.37	0.83	0.40
17M	0.83	0.63	0.10
18M	0.90	0.61	0.12
19M	0.42	0.83	0.79
20M	0.81	0.63	0.06
21M	0.58	0.75	0.55
22M	0.97	0.57	-0.02
23M	0.79	0.63	0.07
24M	0.74	0.66	0.13
25M	0.74	0.66	0.18
26M	0.93	0.58	0.04
27M	0.60	0.73	0.31
28M	0.65	0.71	0.17
29M	0.97	0.57	-0.01
30M	0.49	0.78	0.61
31M	0.32	0.86	0.65
32M	0.58	0.74	0.36

645 **Figure 1**
646
647

648　**Figure 2**

650　**Figure 3**

 Figure 4