Mating preferences act independently on individual elements of visual signals in *Heliconius* butterflies.

Smith, S. H.[1], Queste, L.[1], Wright, D.S. [1], Bacquet, C.N.[2] & Merrill, R.M.[1,3]

[1] Division of Evolutionary Biology, Ludwig-Maximilians-Universität München, Grosshaderner Str. 2, 82152 Planegg-Martinsried, Germany

[2] Universidad Regional Amazónica de Ikiam, Tena, Ecuador

[3] Smithsonian Tropical Research Institute, Gamboa, 0843-03092, Panama

Abstract

Mating cues are often comprised of several elements, which can act independently, or in concert to attract a suitable partner. Individual elements may also function in other contexts, such as predator defence or camouflage. In *Heliconius* butterflies, wing patterns comprise several individual colour pattern elements, which advertise the butterflies' toxicity to predators. These wing patterns are also mating cues, and males predominantly court females that possess the same wing pattern as their own. However, it is not known whether male preference is based on the full wing pattern or only individual pattern elements. We compared preferences of male *H. erato lativitta* between female models with the full wing pattern and those with some pattern elements removed. We found no differences in preference between the full wing pattern model and a model with pattern elements removed, indicating that the complete composition of all elements is not essential to the mating signal. Wing pattern preferences also contribute to pre-mating isolation between two other *Heliconius* taxa, *H. erato cyrbia* and *H. himera*, therefore, we next compared preferences for the same models in these species. *H. erato cyrbia* and *H. himera* strongly differed in preferences for the models, potentially providing a mechanism for how pre-mating isolation acts between these species. These findings suggest that contrasting levels of selective constraint act on elements across the wing pattern.

Introduction

Animal signals are highly diverse and often involve multiple elements, acting across various sensory modalities. Even within a single modality, different elements may target different receivers and/or communicate different information (Candolin 2003). For example, visual cues can include multiple discrete colour and pattern elements (Endler 1978), as well as ornamentation or movement (Fleishman 1988). These signals may be shaped by several sources of selection acting in conflict or co-operation, including predation and sexual selection. Determining how receiver behaviour responds to individual signal elements can therefore help us understand how the evolution of composite signals is constrained and/or contributes to diversification.

Incorporating multiple elements can enhance signalling function in several ways. Certain elements may be more effective when viewed in unison (Guilford & Dawkins 1991, Brooks & Caithness 1995). For example, in budgerigars (*Melopsittacus undulatus*) the high contrast generated by adjacent UV fluorescent and reflective plumage patches is more attractive to mates than either element in isolation (Arnold *et al.* 2002, Hausmann *et al.* 2003). Composite signals may also allow transmission over a broader range of sensory environments. Male *Schizocosa* wolf spiders use both vibratory and visual courtship signals, which they adjust to more effectively signal across different sensory backgrounds (Hebets & Papaj 2005, Gordon & Uetz 2011). Finally, individual elements can carry independent information. In American goldfinches (*Carduelis tristis*), carotenoid-based plumage and bill colouration transmit information about parasite load, whereas the melanin-based black cap signals social status (McGraw & Hill 2000).

Despite this, it is increasingly apparent that mate preferences are sometimes based on only a subset of the total signal elements. Male *Eleutherrodactylus coqui* tree frogs, for example, produce a distinctive two-note call to communicate with conspecifics, but females respond to only the second note (Narins & Capranica 1976). Similarly, female preferences in mallard ducks primarily depend on bill colour brightness, even though males display various other visual ornaments (Omland 1996). A study of male preference in Lycaenid butterflies showed that preference is based on an apparently very minor part of the overall signal. By subtly manipulating the wing patterns of female *Lycaenides melissa* to change the size of the hindwing spot and the associated 'aurorae' to mimic those of its sister species, *L. idas*, Fordyce *et al.* (2002) found that male *L. idas* preferred the manipulated wing patterns, even

though the rest of the wing matched the heterospecific pattern. Together these examples

highlight that different visual elements may serve different evolutionary functions, and

perhaps that selection may not be uniform across different elements.

Numerous studies have now shown that traits under divergent ecological selection are

additionally used as mating cues (Servedio *et al.* 2011). These types of traits may be

important during speciation, because they couple divergent selection acting on the ecological

trait to mating preferences, thereby promoting the evolution of premating barriers (Gavrilets,

2004; Servedio 2011). Warning patterns, advertising toxicity or unpalatability to potential

predators, may act in this way if they are additionally used as mating cues by conspecifics.

For example, the bright warning signals of the toxic strawberry poison dart frog, *Oophaga*

pumilio, are used also used by females during mate choice (Summers *et al.* 1999, Siddiqi *et*

al. 2004). However, while elements of mating cues can act independently from each other,

the function of aposematic cues relies on them being perceived in composite. This is because

a key feature of aposematic signals is visibility and recognisability, which is often achieved

by patterns of adjacent colour patches, contrasting strongly both with each other and the

background scene (Stevens & Ruxton 2012)**.** Given the multi-component nature of

aposematic signals, it is unclear how mate preference acts when the same elements that make

up aposematic signals are also used as mating signals.

The Neotropical *Heliconius* butterflies exhibit a huge diversity of bright warning patterns,

which are often associated with Müllerian mimicry (Merrill *et al.* 2015). These warning

patterns are both under strong positive frequency dependent selection (Mallet & Barton 1989;

Merrill et al 2012), and are also used as mating cues (Crane 1955, Jiggins *et al.* 2001).

Specifically, multiple studies have now shown that *Heliconius* males normally prefer to court

females that share their own colour pattern over that of closely related taxa, (e.g Jiggins *et al.*

2001, Jiggins *et al.* 2004, Estrada & Jiggins 2008; Merrill et al. 2014), and this contributes a

strong premating reproductive barrier between closely related species (Jiggins *et al.* 2001).

The extremely close mimicry exhibited between the distantly related species *H. erato* and *H.*

melpomene suggests that predators discriminate against deviations in pattern across the entire

wing. However, although both *H. erato* and *H. melpomene* males are known to discriminate

between females based on visual cues (Jiggins *et al.* 2004, Estrada & Jiggins 2008) relatively

little is known about which pattern elements trigger the behavioural response. Previously,

Estrada & Jiggins (2008) compared preferences between geographical populations of *H.*

erato with divergent warning patterns and found that males were more likely to be attracted

towards females with at least one of the wing pattern elements in common to their own, over females with more different warning patterns. Additionally, Jiggins and colleagues (2001) found little difference in the response of *H. melpomene* to visually near identical models made from females from populations with or without a yellow hindwing bar. Finkbeiner *et al.* (2014) also showed that colour is a more important determinant of male attraction towards *H. erato petiverana* females than the arrangement of these colours (i.e. pattern). However, no experiments have explicitly tested whether the whole composite wing pattern elicits male courtship behaviours, or just specific elements.

To determine how visual preferences are influenced by individual elements of composite warning patterns, we manipulated individual elements of the *H. erato lativitta* colour pattern and tested the corresponding male preferences. In order to understand how preference for pattern elements could impact divergence of populations, we also investigated how males from two other Erato clade populations, *H. himera* and *H. erato cyrbia*, responded to manipulated *H. erato lativitta* wing patterns. Overall, our results suggest that not all elements of the aposematic pattern are important for mate choice, and that differences in preference based on individual pattern elements could play a role in the divergence of closely related species.

Figure 1. (A) Topographical map of Ecuador, showing the Andes mountain range in the centre. Experiments took place at Universidad Regional Amazónica IKIAM (red diamond) and butterflies were collected at the sites surrounding the locations marked with images of the butterflies. (B) Layout of experimental cage, with circular frame suspended in the centre. Butterfly models were attached to the ends of the white sticks extending inwards from the frame.

Methods

Butterfly collection and rearing.

All experiments took place in insectaries at Universidad Regional Amazónica Ikiam, Tena, Ecuador (Figure 1, red diamond) between November 2021 and January 2022. *H. erato lativitta* were collected in the southeast of Napo province, Ecuador, whereas *H. erato cyrbia* and *H. himera* came from insectary populations established from individuals collected in southwest Ecuador (Figure 1). Butterfly stocks were maintained in 2 x 2 x 2m cages and provided with *Passiflora punctata* shoots for oviposition, nectar-bearing flowers, and ~20% sugar solution with supplementary pollen for additional nutrition (Gilbert 1972). Caterpillars from these stocks were reared indoors and fed *ad libitum* on *Passiflora punctata*.

Mounted female wing models.

To make the female wing models, wings from sacrificed *H. erato lativitta* females were removed and subsequently washed for 5 minutes in dichloromethane to remove pheromones (Darragh *et al.* 2017). Wings were then manipulated by colouring over a patch of colour on both sides of the wing with a black marker pen (COPIC Ciao Black 100 (Too Marker Products Inc.)) to generate one of 3 distinct wing patterns: 'intact', 'red only' or 'yellow only' as shown in Figure 2A-C. The remaining unmanipulated part of the wing was coloured with a colourless marker pen with the same solvent (COPIC Ciao Colourless Blender 0 (Too Marker Products Inc.). Wings were then glued to white card, using a thin paper strip to allow flapping motion, and mounted onto black foam strip that acted as the butterfly's body and allowed to dry for at least 2 hours before use.

Male preference experiments.

Trials took place between 08:00 and 17:00. Males were tested at least 5 days after eclosion. One male each of *H. erato lativitta*, *H. e. cyrbia* and *H. himera* were tested together in the same trial, and the combination of 3 males tested together was randomised between trials. Males were released into experimental cages at between 1 and 24 hours before the trial to acclimatise, during which they were provided with nectar-bearing flowers and sugar water solution, as above, but these were removed during the trial.

During 30-minute trials, males were presented with a set of three female models (intact, red only and yellow only) mounted on a circular frame, 90cm in diameter, suspended 1m from the ground (Figure 1B). The models were randomly placed at one of three positions on the

frame, equally spaced at 60cm away from each other. A string attached to the centre of the

frame was pulled continuously throughout the trials to simulate wing movement (Klein & de

Araújo 2010). Four different sets of models were used to account for individual variation

between models. At the beginning of each trial, we recorded the date, time, weather,

(recorded as 'clear', 'under 50% cloud cover', over 50% cloud cover' or 'overcast'), model

set and temperature.

Two types of behavioural interactions between the males and the models were recorded

during the trial, using the Field Notes app (Neukadye). These were 'approach' and

'courtship'. We defined approach as a male directing its flight towards and coming within

10cm of the model. Courtship was defined as 'approach' but additionally included hovering

above the model for at least 1 second. Accordingly, if a male courted the model, this was also

recorded as 'approach'. Both behaviours were scored independently once per minute, for

each model type and male, as having occurred (1) or not occurred (0).

Preference trials for each male were repeated 5 times for each model set to account for

individual variation. Each male was tested on multiple model sets when possible, to account

for variation in female model quality, though this was often not possible and most males

(60%) were tested with only one set. Each model set was used for testing at least 2 different

males each of species. Males were tested a maximum of 2 times per day and remained in

experimental cages between the 5 repeated trials per model set, in the presence of other

males.

Statistical analysis.

To determine whether males responded differently to the three model types, we fitted

generalised linear mixed models (GLMM) with binomial error distribution, implemented in

the *lme4* package (Bates *et al.* 2015) in R (R Core Team 2021). The dependent variable was

either 'approach' or 'courtship' encoded as a binary response variable for each minute of the

trial (*i.e.* 0 or 1, see above). Saturated models included male species, model type and their

interaction as dependent variables. Weather was also included as a fixed factor with four

levels: '100% overcast', '>50% overcast', '<50% overcast' and 'clear' which correlated

strongly with temperature (see supplementary materials). Male ID and trial nested within

model set were included as random variables in all models to account for repeated measures.

We implemented model optimiser BOBYQA (Powell 2009) to overcome convergence

warnings associated with the complex structure of the models. We subsequently checked for

deviation between log likelihoods of different optimisers using the command *allFit* from the

lme4 package (SM Table 1). Significance of fixed factors was determined using likelihood

ratio tests (LRTs) by comparing models with and without individual terms. Effect sizes were

calculated as the estimated marginal means (EMM) of the minimum adequate model using

the *emmeans* package (Lenth 2021). We generated ternary plots of the raw data to visualise

differences in approach and courtship by each male type, towards the three models. All

analysis scripts are included in the supplementary materials.

Figure 2. (A – C) Model types used to test mate preference based on wing pattern elements. (A) the
full *H. erato lativitta* wing pattern, used for the 'intact' model; (B) 'Red only' model, featuring only the
red dennis-ray elements of the *H. erato lativitta* wing pattern; (C) 'Yellow only' model, featuring only
the yellow forewing band of the *H. erato lativitta* wing pattern.. Figures are a graphical representation
of the model types. (D) *H. erato cyrbia* wing pattern. (E) *H. himera* wing pattern. Figure shows only
the left dorsal wing pattern, but patterns were identical on all sides of the wing.

Figure 3. Ternary plots showing the proportional preference responses of minutes in which individual males directed preference behaviours towards either of the three female wing pattern models. A-C shows approach behaviours in (A) *H. erato lativitta*, (B) *H. erato cyrbia*, and (C) *H. himera*. D-F shows hovering courtship behaviours in (D) *H. erato lativitta*, (E) *H. erato cyrbia*, and (F) *H. himera*. Left ternary axis shows the proportion of minutes in which males directed preference towards the intact model, bottom ternary axis, towards the red only model, and right axis towards the yellow only model. Each point represents the mean responses of males across all trials, and point size represents the cumulative activity level of that male across all trials. Males which did not show any courtship responses are not represented in the ternary plots. Butterfly images in the top left of each plot show the wing pattern of the male species of the plot.

Table 1. Contrast estimates showing effect size and direction of differences in estimated marginal means of preference behaviours**.** Values indicate difference in probability of approach (above diagonal, rows minus columns) or courtship (below diagonal, columns minus rows) with 95% confidence intervals**.** Values in bold indicate significant differences for each population in their approach or courtship towards the compared model types.

A) *H. erato lativitta*	Intact	Red only	Yellow only
Intact	–	-0.006 (-0.014 – 0.001)	**0.008 (0.0004 – 0.015)**
Red only	-0.002 (-0.0042 – 0.0008)	–	**0.014 (0.003 – 0.025)**
Yellow only	**0.002 (0.00004 – 0.005)**	**0.004 (0.0006 – 0.007,)**	–

B) *H. erato cyrbia*	Intact	Red only	Yellow only
Intact	–	-0.004 (-0.014 – 0.007)	**0.036 (0.010 – 0.061)**
Red only	0.0008 (-0.003 – 0.004)	–	**0.039 (0.012 – 0.067)**
Yellow only	**0.006 (0.001 – 0.012)**	**0.006 (0.001 – 0.01)**	–

C) *H. himera*	Intact	Red only	Yellow only
Intact	–	0.002 (-0.003 – 0.008)	-0.004 (-0.011 – 0.002)
Red only	0.0007 (-0.001 – 0.003)	–	-0.007 (-0.014 – 0.0003)
Yellow only	0.001 (-0.0006 – 0.003)	0.0007 (-0.0009 – 0.002)	–

Results

In total, we measured preferences of 21 *H. erato lativitta*, 20 *H. erato cyrbia* and 21 *H. himera* males over 122 behavioural trials. The three male types responded differently to the female models, both in terms of the number of 'approaches' and number of 'courtships' (Figure 3), as indicated by a significant reduction in deviance when the interaction between male species and model type was removed from our GLMMs for both 'approach' (LRT: $2\Delta\ln L = 147.9$, d.f. = 4, $p < 2.2e^{-16}$) and 'courtship' (LRT: $2\Delta\ln L = 24.4$, d.f. = 4, $p = 6.8e^{-5}$), indicating that the males respond differently to different pattern elements. Weather strongly predicted male approach behaviours (LRT: $2\Delta\ln L = 13.73$, d.f. = 3, $p < 0.01$) but not courtship behaviours (LRT: $2\Delta\ln L = 5.32$, d.f. = 3, $p = 0.15$); however, there was no significant difference between the EMM of male approach probability towards each model in each weather category (SM Figures 1-3; SM Table 2). Nevertheless, we retained weather in all models because it was shown to influence likelihood of a male approaching the model,

and all courtship behaviours were preceded by a male approaching a model. As such, for both approach and courtship the minimum adequate model included the male species*model type interaction term, as well as male species, model type and weather as fixed effects. To further investigate differences in preference for wing pattern across the three species, we calculated the EMM of proportion of minutes with 'approaches' and 'courtships' towards each model type separately for each species. *H. erato lativitta* did not distinguish between the intact and red only model but was more likely to 'approach' and 'court' the intact and red only model than the yellow only model (Table 1A). Looking across all three male types, we found similar preference (both in terms of 'approach' and 'courtship') between *H. erato lativitta* and *H. erato cyrbia*, which both preferred the intact and red only models most, and the yellow only model least, while *H. himera* showed no model preference in either 'approach' or 'courtship' behaviours (Table 1).

Discussion

We found that experimentally removing individual warning colour pattern elements significantly influenced male behaviours; however, this effect differed between males of closely related species. For *H. erato lativitta,* individual colour pattern elements influenced male courtship response, but the full pattern model was not preferred most of all. Male *H. erato lativitta* preferred to court the model with the red dennis-ray pattern only over the model that lacked any red elements, but there was no difference in preference between the intact wing pattern model, and the model without the yellow forewing element. It would appear that the red dennis-ray elements are more attractive to male *H. erato lativitta* than yellow elements, and that this part of the pattern may act independently in signalling to courting males, if at all. At first these results might seem surprising: the survival of these butterflies relies heavily on predators recognising their wing patterns as toxic, reinforced by mimicry between co-occurring species. *H. erato lativitta* is part of a mimicry ring with several *Heliconius* species found in the Ecuadorian Amazon, as well as some non-*Heliconius* and even some moths, all of which have both the red dennis-ray pattern and yellow patch in their warning pattern (Jiggins 2016). This would predict strong selection for males to prefer females with this shared warning pattern, to maintain the recognised signal in their offspring.

The lack of strong preference in *H. erato lativitta* for females with all the components of its

own colour pattern shows that some elements may be redundant to the *H. erato lativitta*

visual mating signal. This may indicate that there is variation in the exact information

communicated across the wing. These findings contribute to the collection of studies showing

that individual wing pattern elements serve a variety of functions, and not all are used in mate

preference (Candolin 2003). For example, a study by Omland (1996) found that in mallard

ducks (*Anas platyrhynchos*), females base preference predominantly on the brightness of the

males' bill, although other signals apparently act as sexual ornaments. Similarly, in a study of

plumage evolution in hummingbirds, Beltrán et al (2021) proposed that females receive

different information from different plumage patches; those which are similar across species

could be used for mate quality evaluation, while patches which differ were more likely to be

used in species discrimination (Beltrán *et al.* 2021). While is not possible to conclude the

exact functions of individual *H. erato lativitta* pattern elements from this study, these results

do indicate that individual wing pattern elements can act independently as signals and are

likely under different selective pressures.

When comparing between species of *Heliconius*, it appears that most of the differences were

associated with presence or absence of red pattern elements. Despite very different wing

patterns, both *H. erato* subspecies were similar in approaching the yellow only model

significantly less than the intact or red only models, while *H. himera* showed no model

preference. This difference is noteworthy considering evidence of premating isolation

between *H. himera* and *H. erato cyrbia* based on wing pattern (Merrill *et al.* 2014). *H. himera*

and *H. erato cyrbia* are sister species, in which assortative mating is maintained at a rate of

90% across their hybrid zone at least in part by visual premating isolation alone (Jiggins *et al.*

1996, Merrill *et al.* 2014). The *H. himera* wing pattern features a yellow forewing patch,

similar in appearance to the yellow patch on the *H. erato lativitta* pattern, while *H. erato*

cyrbia features a red forewing patch and no yellow forewing elements (Figure 2). It could be

that *H. erato cyrbia* males preferentially avoid the yellow only model because it gave the

same mating signal as the wing pattern of *H. himera*. However, *H. himera* males did not

show preference for the models with the yellow forewing patch, despite this similarity. In

fact, *H. himera* males exhibited no differences in preference between any of the models,

which could be driven by lower investment in visual sensitivity in *H. himera* compared to *H.*

erato (Montgomery & Merrill 2017). It is striking to find that male *H. erato cyrbia* showed

preference for red wing pattern elements even in a very different wing pattern, however this

phenomenon has also been observed in Lycaenid butterflies (Fordyce *et al.* 2002). Although

subtle, these differences may demonstrate that premating isolation between *H. erato cyrbia*

and *H. himera* is maintained by preference in *H. erato cyrbia* for red forewing elements, and

lack of visual sensitivity leading to weaker visual preference in *H. himera*.

We also found no overall significant differences in preference between weather conditions,

although weather did interact with model type in our statistical model of approach behaviour.

This is in contrast to Hausmann AE *et al.* (2021), who found that *Heliconius* butterflies are

more likely to court their own colour pattern with brighter light environment. Across our

study, over 80% of our trials occurred in overcast or highly cloudy conditions, therefore it is

likely that we did not experience enough days of bright light while conducting our trials to

detect strong differences in preference. This could also explain why we did not detect strong

preference differences within *H. himera*, as the low light conditions resulted in less active

males, fewer courtships, and less strong discrimination between wing patterns models by

males.

Visual mate choice signals play a role in the early stages of divergence in *Heliconius*

butterflies (Merrill *et al.* 2015). It would seem that *H. himera* has diverged in preference

based on wing pattern components from the two *H. erato* included in this study. This hints as

to the process of speciation with wing pattern divergence in these species; a complete

overhaul of the full wing pattern is not necessary, rather finer scale changes to individual

mating signal elements while maintaining key preference elements, alongside changes in

preferences, are enough to drive divergence. For species such as *Heliconius* butterflies,

whose morphological evolution is constrained by selection for aposematic warning signals,

this illuminates how wing pattern diversification can proceed within the constraints imposed

by predation. By enacting mating preferences on only particular elements of the mating

signal, selection for aposematic warning patterns and mating cues are balanced across the full

wing pattern. The differences in preference between *H. erato* and *H. himera* illustrate this

idea, showing that strong differences preference for colours of the forewing pattern may be

enough to drive assortative mating. Previous studies have found that mate preferences in

Heliconius are commonly restricted to discrimination between forewing patterns (Jiggins *et*

al. 2001, Kronforst *et al.* 2006). This releases the hind wing pattern from the constraining

selection of mate preference and allows the pattern to be more flexible to selection by

aposematism and mimicry. Whether predators impose differential selection pressures on

different wing elements, or different wings, is yet to be explored in *Heliconius*.

Acknowledgements. We thank José Borerro, Kimberly Gavilanes, Michelle Guachamin, Andrei Flores de Valgas for help in the insectaries and collecting butterflies. We are grateful to the Ministerio del Ambiente, Agua y Transición Ecológica for permission to collect and work with butterflies in Ecuador (MAATE-DBI-CM-2021-0176; MAATE-DNB-CM-2021-0176). This work was funded by an ERC Starting grant (Grant number: 851040) awarded to RMM.

Literature cited

Arnold KE, Owens IPF, Marshall NJ. 2002. Fluorescent Signaling in Parrots. Science 295: 92–92.

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using {lme4}. Journal of Statistical Software 67: 1–48.

Beltrán DF, Shultz AJ, Parra JL. 2021. Speciation rates are positively correlated with the rate of plumage color evolution in hummingbirds. Evolution 75: 1665–1680.

Brooks R, Caithness N. 1995. Manipulating a seemingly non-preferred male ornament reveals a role in female choice. Proceedings of the Royal Society of London Series B: Biological Sciences 261: 7–10.

Candolin U. 2003. The use of multiple cues in mate choice. Biological Reviews of the Cambridge Philosophical Society 78: 575–595.

Crane J. 1955. Imaginal behavior of a Trinidad butterfly, Heliconius erato hydara Hewitson, with special reference to the social use of color. Zoologica : scientific contributions of the New York Zoological Society 40: 167–196.

Darragh K, Vanjari S, Mann F, Gonzalez-Rojas MF, Morrison CR, Salazar C, Pardo-Diaz C, Merrill RM, McMillan WO, Schulz S, Jiggins CD. 2017. Male sex pheromone components in Heliconius butterflies released by the androconia affect female choice. PeerJ 5: e3953.

Endler JA. 1978. A Predator's View of Animal Color Patterns. In: Hecht MK, Steere WC, Wallace B (ed.). Evolutionary Biology, pp. 319–364. Springer US, Boston, MA.

Estrada C, Jiggins CD. 2008. Interspecific sexual attraction because of convergence in warning colouration: is there a conflict between natural and sexual selection in mimetic species? Journal of Evolutionary Biology 21: 749–760.

Finkbeiner SD, Briscoe AD, Reed RD. 2014. Warning signals are seductive: Relative contributions of color and pattern to predator avoidance and mate attraction in Heliconius butterflies. Evolution 68: 3410–3420.

Fleishman LJ. 1988. Sensory influences on physical design of a visual display. Animal Behaviour 36: 1420–1424.

Fordyce JA, Nice CC, Forister ML, Shapiro AM. 2002. The significance of wing pattern
diversity in the Lycaenidae: mate discrimination by two recently diverged species.
Journal of Evolutionary Biology 15: 871–879.

Gavrilets, S. 2004. Fitness landscapes and the Origin of Species. Princeton, NJ, USA:
Princeton University Press.

Gilbert LE. 1972. Pollen Feeding and Reproductive Biology of Heliconius Butterflies.
Proceedings of the National Academy of Sciences 69: 1403–1407.

Gordon SD, Uetz GW. 2011. Multimodal communication of wolf spiders on different
substrates: evidence for behavioural plasticity. Animal Behaviour 81: 367–375.

Guilford T, Dawkins MS. 1991. Receiver psychology and the evolution of animal signals.
Animal Behaviour 42: 1–14.

Hausmann AE, Kuo C-Y, Freire M, Rueda-M N, Linares M, Pardo-Diaz C, Salazar C,
Merrill RM. 2021. Light environment influences mating behaviours during the early
stages of divergence in tropical butterflies. Proceedings of the Royal Society B:
Biological Sciences 288: 20210157.

Hausmann F, Arnold KE, Marshall NJ, Owens IPF. 2003. Ultraviolet signals in birds are
special. Proceedings of the Royal Society B: Biological Sciences 270: 61–67.

Hebets E, Papaj D. 2005. Complex signal function: developing a framework of testable
hypotheses. Behavioral Ecology and Sociobiology 57: 197–214.

Jiggins CD. 2016. The Ecology and Evolution of Heliconius Butterflies. doi
10.1093/acprof:oso/9780199566570.001.0001.

Jiggins CD, Estrada C, Rodrigues A. 2004. Mimicry and the evolution of premating isolation
in Heliconius melpomene Linnaeus. Journal of Evolutionary Biology 17: 680–691.

Jiggins CD, McMillan WO, Neukirchen W, Mallet J. 1996. What can hybrid zones tell us
about speciation? The case of *Heliconius erato* and *H. himera* (Lepidoptera:
Nymphalidae). Biological Journal of the Linnean Society 59: 221–242.

Jiggins CD, Naisbit RE, Coe RL, Mallet J. 2001. Reproductive isolation caused by colour
pattern mimicry. Nature 411: 302–305.

Klein AL, de Araújo AM. 2010. Courtship behavior of Heliconius erato phyllis (Lepidoptera,
Nymphalidae) towards virgin and mated females: conflict between attraction and
repulsion signals? Journal of Ethology 28: 409–420.

Kronforst MR, Young LG, Kapan DD, McNeely C, O'Neill RJ, Gilbert LE. 2006. Linkage of
butterfly mate preference and wing color preference cue at the genomic location of
wingless. Proceedings of the National Academy of Sciences 103: 6575–6580.

Lenth RV. 2021. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Mallet J, Barton NH. 1989. Strong natural selection in a warning-color hybrid zone.
Evolution 43(2): 421-431.

McGraw KJ, Hill GE. 2000. Differential effects of endoparasitism on the expression of
carotenoid- and melanin-based ornamental coloration. Proceedings of the Royal
Society B: Biological Sciences 267: 1525–1531.

Merrill RM, Dasmahapatra KK, Davey JW, Dell'Aglio DD, Hanly JJ, Huber B, Jiggins CD,
Joron M, Kozak KM, Llaurens V, Martin SH, Montgomery SH, Morris J, Nadeau NJ,
Pinharanda AL, Rosser N, Thompson MJ, Vanjari S, Wallbank RWR, Yu Q. 2015.
The diversification of Heliconius butterflies: what have we learned in 150 years?
Journal of Evolutionary Biology 28: 1417–1438.

Merrill, R.M., Chia, A., Nadeau, N. 2014. Divergent warning patterns contribute to
assortative mating between incipient *Heliconius* species. Ecology and Evolution 4:
911-917

Merrill, R.M., Wallbank, R.W., Bull, V., Salazar, P., Mallet J., Stevens, M. & Jiggins C.D.
2012. Disruptive ecological selection on a mating cue. Proceedings of the Royal
Society of London Series B: Biological Sciences 279: 4907-4913

Montgomery SH, Merrill RM. 2017. Divergence in brain composition during the early stages
of ecological specialization in Heliconius butterflies. Journal of Evolutionary Biology
30: 571–582.

Narins PM, Capranica RR. 1976. Sexual Differences in the Auditory System of the Tree Frog
Eleutherodactylus coqui. Science 192: 378–380.

Neukadye. Field Notes Journal - Neukadye. WWW document: neukadye.com/#home.
Accessed 14 February 2022.

Omland KE. 1996. Female Mallard Mating Preferences for Multiple Male Ornaments: II.
Experimental Variation. Behavioral Ecology and Sociobiology 39: 361–366.

Powell MJD. 2009. The BOBYQA algorithm for bound constrained optimization without
derivatives. University of Cambridge

R Core Team RF for SC. 2021. R: A Language and Environment for Statistical Computing.

Servedio MR, Doorn GSV, Kopp M, Frame AM, Nosil P. 2011. Magic traits in speciation:
'magic' but not rare? Trends in Ecology & Evolution 26: 389–397.

Siddiqi A, Cronin TW, Loew ER, Vorobyev M, Summers K. 2004. Interspecific and
intraspecific views of color signals in the strawberry poison frog Dendrobates
pumilio. The Journal of Experimental Biology 207: 2471–2485.

Stevens M, Ruxton GD. 2012. Linking the evolution and form of warning coloration in
nature. Proceedings of the Royal Society B: Biological Sciences 279: 417–426.

Summers K, Symula R, Clough M, Cronin T. 1999. Visual mate choice in poison frogs.
Proceedings of the Royal Society of London Series B: Biological Sciences 266: 2141–
2145.

Too Marker Products Inc. COPIC. WWW document: https://copic.jp/en/. Accessed 14
February 2022.

Supplementary Materials

We tested for the relationship between our four weather variables ('100% overcast', '>50% overcast', '<50% overcast' and 'clear') and temperature, and the effect of these variables on both approach and courtship. First, to test how temperature and weather relate, we fitted a linear regression of temperature against the categorical variables of weather, and then tested the significance of this relationship in an ANOVA. Overall, 51% of trials occurred in overcast weather, 32% occurred in high cloud weather, 11% in low cloud weather and 6% in clear weather. The mean temperature of the trials was 27.5°C (SD = 4.1°C), and average temperature decreased with increasing cloud cover (SM Figure 1; one-way ANOVA: F = 15478, D.F. = 3, $p < 2.2e^{-16}$). This showed that temperature and weather are highly correlated, so then we tested how male activity level varied with weather category. For this we used the binomial counts of approach or courtship behaviour as in our analysis of preference differences, but fitted these separately in a GLMM with only weather as an independent variable. We then use LRT to test for a significant difference in deviance when weather was removed, and estimated the EMMs of the activity level across each weather category.

 SM Table 1 Results of model optimisation comparisons

Dependent variable	Independent variables	Random effects	Log likelihood of model optimisers				
			Nelder-Mead	bobyqa	nlminbwrap	nloptwrap. NLOPT_LN_NELDERMEAD	nloptwrap. NLOPT_LN_BOBYQA
Approach	weather + model_type * male_species	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-4417.102	-4417.101	-4417.101	-4417.105	-4417.102
Approach	weather + model_type + male_species	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-4491.041	-4491.035	-4491.035	-4491.035	-4491.035
Approach	weather + model_type	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-4492.21	-4492.21	-4492.21	-4492.21	-4492.21
Approach	weather + male_species	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-4533.723	-4533.723	-4533.726	-4533.723	-4533.723
Courtship	weather + model_type * male_species	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-1191.709	-1191.697	-1191.697	-1191.697	-1191.697
Courtship	weather + model_type + male_species	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-1203.878	-1203.876	-1203.876	-1203.876	-1203.876
Courtship	weather + model_type	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-1205.237	-1205.237	-1205.237	-1205.237	-1205.237
Courtship	weather + male_species	(1 | male_id) + (1 | model_set) + (1 | model_set:trial_id)	-1236.995	-1236.995	-1236.995	-1236.995	-1236.995

SM Figure 1 Number of approach behaviours recorded per male per trial, per weather category, plotted against the temperature recorded during the trial. Points are coloured by the weather category the trial occurred in, and a line of best fit for each weather category is fitted to the graph.

SM Figure 2 Number of courtship behaviours recorded per male per trial, per weather category, plotted against the temperature recorded during the trial. Points are coloured by the weather category the trial occurred in, and a line of best fit for each weather category is fitted to the graph.

480 **SM Figure 3** Approach behaviour of males during trials of each weather category, shown by the estimated
481 marginal means of proportion of minutes during the trial in which males of all species approached the
482 models. Error bars show the 95% confidence intervals, based on the analysis. Numbers above the error bars
483 show the number of trials conducted in each weather category.