rsofun v4.4.1: A model-data integration framework for simulating ecosystem processes

Josefa Arán Paredes[1,2], Koen Hufkens[1,2], Benjamin D. Stocker[1,2]

[1]Institute of Geography, University of Bern, Hallestrasse 12, 3012 Bern, Switzerland
[2] Oeschger Center for Climate Change Research (OCCR), University of Bern, Hochschulstrasse 6, 3012 Bern, Switzerland

Correspondence to: Benjamin D. Stocker (benjamin.stocker@unibe.ch)

Abstract. Mechanistic vegetation models serve to estimate terrestrial carbon fluxes and climate impacts on ecosystems across diverse biotic and abiotic conditions. Systematically informing them with data is key for enhancing their predictive accuracy and estimate uncertainty. Here we present the Simulating Optimal FUNctioning {rsofun} R package, providing a computationally efficient and parallelizable implementation of the P-model for site-scale simulations of ecosystem photosynthesis, complemented with functionalities for Bayesian model-data integration and estimation of parameters and uncertainty. We provide a use case to demonstrate the package functionalities for modelling ecosystem gross CO_2 uptake at one flux measurement site, including model sensitivity analysis, Bayesian parameter calibration, and prediction uncertainty estimation. {rsofun} lowers the bar of entry to ecosystem modelling and model-data integration and serves as an open access resource for model development and dissemination.

1 Introduction

The modelling of land ecosystem processes and structure, water, and carbon fluxes relies on both mechanistic and statistical approaches (Dietze et al., 2018; Hartig et al., 2012; Van Oijen et al., 2005). Mechanistic models are formulated as mathematical descriptions of functional relationships between the abiotic environment and ecosystem states and rates. These descriptions reflect available theory and general empirical patterns and provide a means for "translating" hypotheses about governing principles and causal relationships into testable predictions (Marquet et al., 2014), and for upscaling model-based estimates in geographical space and to novel environmental conditions. However, also mechanistic models rely on statistical (empirical) descriptions of processes at varying levels of abstraction.

Mechanistic models have model parameters that are either specified directly or fitted to data. A great advantage of mechanistic models is that they explicitly link known physical constants with process representations (e.g., molecular mass of CO_2 for diffusion and assimilation, or the gravitational constant and viscosity of water for its transport and transpiration). Other parameters may be specified based on independent measurements under controlled conditions (e.g., the activation energy of Arrhenius-type metabolic rates), or represent (more or less directly) measurable plant functional traits, taken as constant over

time and within plant functional types (PFTs - commonly the basic units in mechanistic vegetation models). Both types of parameters have traditionally been specified in models directly ('direct parameterization' (Hartig et al., 2012)). Yet other model parameters may not be directly observable and describe processes that are not explicitly resolved in models, but parameterized at a higher level of abstraction. Such parameters are often fitted to data such that the agreement between one (or several) related model predictions and observations is optimised. Parameter optimization/estimation for mechanistic vegetation models typically employs generic optimization algorithms or Bayesian statistical approaches and is often used for specifying different types of parameters (except for universal physical constants). Bayesian methods have the advantage that they enable a systematic assessment of the correlation structure among multiple fitted parameters, provide a means for considering uncertainty in observations and available prior information, and generate probabilistic parameter estimations and model predictions (Dietze et al., 2018; Hartig et al., 2012; Van Oijen et al., 2005).

State-of-the-art mechanistic vegetation models, used for global simulations and as components in Earth System Models, are complex. They describe a large number of interrelated state variables, treat multiple feedbacks between them, and rely on a large number of parameters for multiple PFTs and other ecosystem components and properties (e.g., soil, microbes, biophysical and hydrological properties). This makes them "data-hungry" (Hartig et al., 2012). However, their complexity and computational demand poses a limitation for systematic model-data integration and Bayesian parameter estimation.

Eco-evolutionary optimality (EEO) principles have been proposed for reducing model complexity and for a robust grounding of models in governing principles (Franklin et al., 2020; Harrison et al., 2021). They enable parameter-sparse representations, limit the distinction of separate PFTs, and may enable better model generalisations to novel environmental regimes. EEO principles make predictions of plant functional traits that would otherwise have to be prescribed as temporally fixed model parameters. However, although parameter-sparse, EEO-based vegetation models are not devoid of model parameters. Ideally, remaining parameters represent known physical constants or quantities that can be measured independently. But remaining parameters in optimality models typically also represent quantities that are not directly measurable - e.g., the marginal cost of water in (Cowan and Farquhar, 1977), or the unit cost ratio in (Prentice et al., 2014). These are considered to be more universally valid (e.g., without distinctions between PFTs), but still have to be fitted to data.

The P-model (Prentice et al., 2014; Stocker et al., 2020; Wang et al., 2017b) is an example of an EEO-guided model for terrestrial photosynthesis and its acclimation. It avoids the requirement for prescribing PFT-specific parameters of photosynthesis and stomatal regulation, but instead predicts them from universal EEO principles for the full range of environmental conditions across the Earth's (C_3 photosynthesis-dominated) biomes. However, not directly observable parameters of the underlying EEO theory and of additional empirical parameterizations employed in the P-model (Stocker et al., 2020) remain and have to be fitted to data.

Here, we provide a solution for this challenge, acknowledging that the data is an integral part of the modelling process (Dietze et al., 2013) - even in theory-based models of ecosystem processes. We show how the most important parameters contributing to uncertainty in the P-model can be estimated inversely, using observations of ecosystem-level photosynthetic CO_2 uptake (gross primary productivity, GPP). We provide a brief description of the theory embodied in the P-model and introduce the P-

model implementation in the Simulating Optimal FUNctioning {rsofun} version v4.4 modelling framework, made available as an R package. A more comprehensive P-model description can be found in (Stocker et al., 2020). We demonstrate the functionalities of {rsofun} through a case study for simulating GPP at a single flux measurement site. The case study includes sensitivity analysis, Bayesian model calibration, and inference - the prediction of GPP with an estimation of its uncertainty. This paper presents the calibration to GPP observations only, but the package allows calibration to multiple targets simultaneously, including fluxes and leaf traits.

2 P-model description

The P-model predicts the acclimation of leaf-level photosynthesis to a (slow varying) environment based on EEO principles and thereby yields a parameter-sparse representation of ecosystem-level quantities, generalising across (C_3 photosynthesis-dominated) vegetation types and biomes. It combines the established theory for C_3 photosynthesis following the Farquhar-von Caemmerer-Berry (FvCB, (Farquhar et al., 1980)) model with the Least-Cost hypothesis for the optimal balancing of water loss and carbon gain (Prentice et al., 2014), and the coordination hypothesis (Wang et al., 2017a), which states that the light and Rubisco-limited assimilation rates (as described by the FvCB model) are equal for representative daytime environmental conditions. Based on these theoretical foundations, gross primary productivity (GPP) can be modelled as the product of absorbed photosynthetically active radiation, specified by the locally measured photosynthetically active radiation (PAR) and remotely sensed fAPAR, and the theory-based prediction of the ecosystem-level light use efficiency (LUE, (Stocker et al., 2020; Wang et al., 2017b). LUE acclimates to preceding environmental conditions with a characteristic, empirically determined (calibrated), time scale τ. Two latent (not directly observable) parameters govern the optimality-guided water-carbon trade-off: the unit cost ratio β (governing the balancing of maintaining the carboxylation capacity versus the transpiration stream) and c^* (the marginal cost of maintaining the electron transport rate). These have previously been calibrated separately to data and specified as fixed model parameters in the P-model (Stocker et al., 2020; Wang et al., 2017b). The theory for predicting acclimated LUE then requires only the atmospheric environment to be specified (meteorological variables and CO_2). A set of corollary predictions, physically and physiologically consistent with the simulated LUE, follow. These include the acclimated base rates of photosynthetic capacities in the FvCB model (V_{cmax25} and J_{max25}), the acclimated average ratio of leaf internal-to-ambient CO_2 concentration (c_i:c_a), acclimated average daytime stomatal conductance (g_s), and the acclimated base rate of leaf dark respiration (R_{d25}). Physical constants and additional parameters that determine the instantaneous temperature dependence of V_{cmax}, J_{max}, R_d, and parameters in the FvCB model are prescribed and held fixed in the P-model (Tab. A2 in (Stocker et al., 2020)).

For simulating GPP, the P-model is conceived as a single-big-leaf model (Fig. 1). While describing leaf-level quantities at relatively high mechanistic detail, the link between the leaf and the canopy-scale is not explicitly resolved. Instead, an empirical approach for leaf-to-canopy scaling CO_2 uptake is employed by treating the quantum yield parameter φ_0 to be

representative for the canopy-scale and allowing it to be calibrated to ecosystem-level CO_2 flux data. The implementation of
the P-model in the {rsofun} package (version v4.4) further includes an empirical parameterization of the temperature
dependency of the quantum yield φ_0, generalising the approach taken in (Stocker et al., 2020)), and an empirical soil
moisture stress function, generalising the approach taken in (Stocker et al., 2020)) (Appendix A). Acclimating quantities are
derived by employing the P-model theory to gradually varying environmental conditions where variations are damped and
lagged by a characteristic (calibrated) time scale τ. The continuous treatment of the acclimation time scale is different from
(Stocker et al., 2020). The temperature dependency of φ_0 can be turned off by setting a (see Tab. 1 and Appendix A) to 0
('ORG' model setup in (Stocker et al., 2020)). The soil moisture stress function can be turned off by setting β_0 (see Tab. 1
and Appendix A) to 1 ('BRC' model setup in (Stocker et al., 2020)). Taken together, the P-model approach is guided by
EEO and thus yields predictions for quantities that otherwise have to be prescribed and fitted to data. Yet, a small set of
model parameters, related to empirical parametrizations and latent quantities remain and have to be calibrated to data. See
Tab. 1 for a list of all (calibratable) model parameters of the P-model implementation in the {rsofun} package.

Figure 1: P-model inputs, outputs, and target observations for the calibration. The model takes as inputs static site information
(longitude, latitude, elevation and soil water holding capacity), time series of meteorological forcing (daily values of temperature, VPD,
PPFD, net radiation, atmospheric pressure, precipitation, cloud coverage, CO_2, fAPAR), simulation parameters (spinup years, recycle
period length) and, common for all sites, model parameters (listed in Tab. 1). The simulation returns a time series of several ecosystem
fluxes, acclimating leaf traits and ecosystem water balance quantities. By comparing these outputs to field measurements and flux data in a
Bayesian calibration routine, model parameters can be estimated.

Table 1: Calibratable parameters in the {rsofun} P-model implementation. We list the fixed parameter values for non-calibrated
variables used in Sec. 4.2 and, marked with *, the MAP estimates for the calibrated parameters. In parentheses, we list the lower and upper
bounds used in the sensitivity analysis (Sec. 4.1) and for the uniform priors in the Bayesian calibration (Sec 4.2).

Symbol	Parameter name	Description	Units	Values
φ_0	kphio	Quantum yield φ_0 (at optimal temperature)	mol mol^{-1}	0.05797586* (0.03, 0.15)
a	kphio_par_a	Shape parameter for the temperature dependence of φ_0	°C^{-2}	-0.0025 (-0.004, 0.001)
b	kphio_par_b	Optimal temperature for the quantum yield	°C	20 (10, 30)
θ^*	soilm_thetastar	Threshold plant-available soil water content in the soil moisture stress function	mm	144 (0, 240)
β_0	soilm_betao	Stress factor at low soil moisture, intercept for the soil moisture stress function	unitless	0.45964652* (0, 1)
β	beta_unitcostratio	Unit cost ratio of carboxylation (maintenance of V_{cmax}) to transpiration	unitless	146 (50, 200)
b_0	rd_to_vcmax	Ratio of temperature-normalised dark respiration (R_{d25}) to the temperature-normalised maximum carboxylation rate V_{cmax25} (eq. C8 in Stocker et al. 2020)	unitless	0.014 (0.01, 0.1)
τ	tau_acclim	Acclimation time scale of photosynthesis	days	30 (7, 60)
c^*	kc_jmax	Unit cost of electron transport (maintenance of J_{max})	unitless	0.54131889* (0.2, 0.8)

3 The {rsofun} model framework

{rsofun} implements the P-model (Stocker et al., 2020) and provides off-the-shelf methods for Bayesian (probabilistic) parameter and prediction uncertainty estimation. {rsofun} is distributed as an R package and also implements the BiomeE vegetation demography model (Weng et al., 2015). The latter is not further described here and is implemented at an experimental stage in {rsofun} version v4.4. The P-model implementation in {rsofun} is designed for time series simulations by accounting for temporal dependencies in the acclimation to a continuously varying environment. Function wrappers in R make the simulation workflow user-friendly and all functions and input forcing data structures are comprehensively documented (https://geco-bern.github.io/rsofun/).

In {rsofun}, model parameters can be calibrated using a calibration function `calib_sofun()`, providing two modes of calibration, one based on generalised simulated annealing ({GenSA} R package) for global optimization (Xiang et al., 2013) and one based on the {BayesianTools} R package, giving access to a wide variety of Bayesian methods (Hartig et al., 2023). The former being fast, while the latter provides more informed parameter optimization statistics (Clark, 2004; Dietze et al.,

2013). This gives the option for both exploratory and more in-depth analysis of estimated parameters. A set of standard cost

functions are provided for the calibration, facilitating the exploration of various metrics or target variables, and the

specification of calibrated model parameters. Furthermore, the vignettes accompanying the package (https://geco-

bern.github.io/rsofun/articles/) explain how to customise the calibration cost functions and interpret the calibration results.

4 Case study

We use {rsofun} to model GPP as estimated from ecosystem flux measurements taken at one site using the eddy-covariance

technique. The site, selected here for demonstration purposes, is Puéchabon (Rambal et al., 2004) - an evergreen

Mediterranean forest, dominated by *Quercus ilex*, growing on relatively shallow soil on karstic bedrock, and located in

southern France. The climate is governed by a distinct seasonality in solar radiation and temperature, peaking in summer,

and a dry period during summer months with recurrent ecosystem water limitation in late summer (Rambal et al., 2004).

Data used here cover years 2007-2012. Both the model target data (GPP) and model forcing data (see Fig. 1) were measured

at the site and obtained from the FLUXNET2015 dataset (Pastorello et al., 2020). Data for years 2013 and 2014 were

available from FLUXNET2015 but removed here due to unexpectedly low GPP measurements (*pers. comm. Jean-Marc*

Limousin).

4.1 Sensitivity analysis

The P-model implementation in {rsofun} has a total of nine model parameters that are accessible for calibration (Tab. 1).

Due to the relatively high computational cost of simultaneously calibrating all nine model parameters, we start by

performing a sensitivity analysis to determine the most influential parameters on the model fit and exclude the least

influential parameters for the subsequent calibration step. Here, we apply the Morris method for global sensitivity analysis

(Morris, 1991) from the {sensitivity} R package (Looss et al., 2023) and compute the sensitivity metrics μ^*, indicating the

magnitude of the overall influence of a given parameter on the prediction, and σ, a measure of the heterogeneity of a

parameter's influence on the prediction across the parameter space. We assume an additive and normally distributed model

error term for the GPP prediction by the P-model (Trotsiuk et al., 2020) and express the fit to observed data via the Gaussian

log-likelihood. The sensitivity analysis result (Fig. 2) indicates that the model error term is the most influential parameter,

followed by the quantum yield intercept parameter φ_0, the unit cost of electron transport c^* and the soil moisture stress

intercept parameter β_0. The remaining parameters have relatively little influence on the evaluation of daily GPP predictions

from the single site considered here and b_0, the ratio of dark respiration to the temperature-normalised maximum

carboxylation rate, has no influence on GPP predictions, which follows from the model structure. These parameters will be

held constant and thus excluded from the Bayesian calibration below. Note that we chose the threshold of influential and less

influential (and thus fixed) parameters for demonstration purposes here and thus arbitrarily.

4.2 Bayesian calibration

We simultaneously calibrate a subset of the model parameters that have been identified as particularly influential (Sec. 4.1.).

These include the model error term, φ_0, c^*, and β_0. By letting φ_0 and β_0 to be calibrated and setting a to a fixed value of -

0.0025 °C $^{-2}$, the simulations consider a temperature-dependence of φ_0 and a soil moisture stress effect on GPP (see

Appendix A) - similar to the model setup 'FULL' in (Stocker et al., 2020). Note that the model error is simultaneously

calibrated following (Dietze, 2017, p.201). We use the DEzs MCMC sampler (Ter Braak and Vrugt, 2008) as implemented

in {BayesianTools} (Hartig et al., 2023) to estimate the posterior distribution of the calibrated parameters (Fig. 3). All

parameters are given a uniform prior with bounds informed by their physical interpretations (Tab. 1). GPP is assumed to be

normally distributed, as in the Morris analysis. On a 12th Gen Intel Core i7-1270P processor, it took 552.854 sec. to run 3

independent MCMC chains, each with 3 internal chains, and 2000 iterations of individual {rsofun} P-model runs per internal

chain. The algorithm converged with a scale reduction factor (Gelman and Rubin, 1992) of 1.05 (≤1.1). More detail,

examples, and explanations of calibration diagnostics are provided in the vignette (https://geco-

bern.github.io/rsofun/articles/sensitivity_analysis.html). The calculation of the log-likelihood is implemented in the function

`cost_likelihood_pmodel()`, enabling custom calibrations, also for multiple target variables that are considered

simultaneously during model calibration (examples provided in https://geco-
bern.github.io/rsofun/articles/new_cost_function.html).

Figure 3: Prior and posterior distributions of the calibrated model parameters and error term. The maximum a posteriori (MAP)
estimate for the gaussian error is 1.22210684 and the estimates for each calibrated parameter are given in Tab. 1.

4.3. Inference and prediction uncertainty estimation

The parameter sets generated by the MCMC chains provide the basis for inference (model prediction) and prediction
uncertainty estimation, allowing us to get insights into the sources of uncertainty and assess the potential risks associated
with the predicted outcomes in a changing environment. We consider a simple representation of the uncertainty, split
between the parameter uncertainty and the model error (Dietze, 2017).

Figure 4: Predicted and observed GPP time series. The comparison is provided for the first year of GPP observations (black) at the site
FR-Pue (Rambal et al., 2004) against GPP predictions (orange line), calculated as the median of the posterior distribution. A light green
band indicates the 90% credible interval for GPP predictions, which captures parameter uncertainty, while the 90% predictive interval for
GPP predictions (in orange) captures model uncertainty.

We take 600 MCMC samples from the multivariate parameter posterior distribution (estimated during the calibration) and
run the P-model for each set of parameters to predict GPP. The credible interval is computed from the posterior distribution
of predicted GPP (at each time step). The predictive credible interval for GPP is computed by adding generated Gaussian
error, with standard deviation err_gpp, to the predicted GPP. Fig. 4 shows that the uncertainty ascribable to the parameters
(in green) is much smaller than the uncertainty due to the model error (in orange). A point estimate of GPP at each time step
is calculated as the median of the posterior distribution of GPP (in orange).

5 Discussion

The {rsofun} R package provides a user-friendly and fast implementation of the P-model. It implements off-the-shelf model-data assimilation routines with simple function calls, while maintaining flexibility for future experiments and the further development of model uncertainty estimation. The {rsofun} vegetation modelling framework is designed to strike a balance between integration and flexibility, enabled by extensible and user-defined calibration specifications and the parallelizable and fast low-level code implementation in Fortran 90. Its high computational efficiency offers the potential for effective model parameter and uncertainty estimation using Bayesian statistical methods - as demonstrated here.

With the function `calib_sofun()`, {rsofun} provides a blueprint for model-data assimilation and an implementation of the likelihood function with flexibility in selecting among a predefined list of model parameters and target observations. We have demonstrated here how ecosystem flux measurements of GPP from a single site can be used to estimate model parameters of the P-model and generate estimations of prediction uncertainty. The approach taken in previous studies with the P-model was to specify latent parameters directly, based on independent observations. In (Wang et al., 2017b), β and c^* were determined from observations of V_{cmax}, J_{max}, and c_i:c_a. These were then used as constants for modelling GPP, while additional parameters related to empirical photosynthesis stress parameterizations were calibrated to GPP observations. Here, we simultaneously calibrate both types of parameters to GPP data. We have followed a simplified setup - used here for demonstration purposes. We note that future applications may use V_{cmax}, J_{max}, or c_i:c_a data directly as additional calibration targets to provide complementary constraints. Note also that the FvCB photosynthesis model contains additional parameters (see Tab. A2 in Stocker et al., 2020) that are treated as constants here - as in previous publications (Bloomfield et al., 2023; Mengoli et al., 2022; Stocker et al., 2020; Tan et al., 2021; Wang et al., 2017b).

Our model implementation as an R package takes inspiration from the {r3PG} forest model (Trotsiuk et al., 2020), and our implementation of model-data integration on the basis of ecosystem data serves similar, yet reduced, aims and functionalities compared to PEcAn (https://pecanproject.github.io/index.html) (LeBauer et al., 2013). {rsofun} is designed to be minimally reliant on package dependencies and connections to specific data, while limiting the implementation and usability of a predefined set of process models (currently P-model, BiomeE at an experimentation stage). Note that the {rpmodel} R package (available on CRAN) also provides an implementation of the P-model, but is written fully in R and in the form of a function of a given environment - without a treatment of temporal dependencies and without the functionalities for model-data integration.

6 Conclusions

We have demonstrated how important parameters contributing to uncertainty in GPP predictions by the P-model can be estimated inversely, using observations of ecosystem-level photosynthetic CO_2 uptake flux time series. The Bayesian

approach to model-data integration enables a probabilistic prediction of GPP and estimation of model parameters. The

{rsofun} model implementation as an R package makes it possible to leverage a set of methods and complementary libraries

for parameter estimation, sensitivity analysis, and calibration diagnostics - as demonstrated here. Its low-level code in

Fortran 90 is geared towards computational efficiency. Provision of {rsofun} as an open-access library aims at lowering the

bar of entry to modelling for both field ecologists and computational ecologists and serves as an Open Science resource for

future model development and experimentation and the further development of model uncertainty estimation.

Appendix A: Calibrated empirical functions

A new formulation of the temperature dependency of the quantum yield efficiency φ_0 is introduced in the {rsofun} package,

allowing more flexibility than Eq. 18 in (Stocker et al., 2020). It is expressed as follows:

$\varphi_0(T) = c(1 + a(T - b)^2)$ if $0 < c(1 + a(T - b)^2) < 1,$

$\varphi_0(T) = 0$ if $c(1 + a(T - b)^2) \leq 0,$ and

$\varphi_0(T) = 1$ if $c(1 + a(T - b)^2) \geq 1.$

Where T stands for temperature, c is the quantum yield efficiency at optimal temperature (in mol mol^{-1}), a is a unitless

shape parameter and b is the optimal temperature. Whenever $a = 0$, the quantum yield efficiency is kept constant at $\varphi_0 = c$.

A possible improvement for the model would be to use a peaked Arrhenius function instead of a parabola (Medlyn et al.,

2002). Furthermore, the soil moisture stress function follows (Stocker et al., 2020), but the parameters considered for

calibration there differ from the calibratable parameters in the package.

$\beta(\theta) = \frac{\beta_0 - 1}{\theta^{*2}}(\theta - \theta^*)^2 + 1$ if $0 \leq \theta \leq \theta^*,$

$\beta(\theta) = 1$ if $\theta > \theta^*.$

In the equation above, θ stands for the plant-available soil water (in mm) and θ^* for the threshold indicating when the plants

start being water stressed. The intercept β_0 is the β reduction at low soil moisture, that is $\beta_0 = \beta(0)$. This intercept is now

calibrated directly, rather than expressed as a function of mean aridity (see eq. 20 in (Stocker et al., 2020)).

Code availability

The {rsofun} R package can be installed from GitHub and its source code is publicly available on https://github.com/geco-

bern/rsofun under an AGPLv3 licence.

Data availability

Data used here for model calibration and evaluation is in the {rsofun} package and is based on publicly available FLUXNET2015 data for the site FR-Pue.

Author contribution

B.D.S. wrote the Fortran code and designed the study. K.H. initiated the implementation of the calibration procedure and wrote the initial draft of the manuscript. J.A.P. documented the package, implemented the Bayesian calibration, sensitivity analysis and uncertainty estimation, and contributed to manuscript writing.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We thank Florian Hartig for advice on uncertainty modelling and the maintenance of the BayesianTools package, Volodymyr Trotsiuk for an initial template of the R package, and Colin Prentice for comments on the manuscript. B.D.S. was funded by the Swiss National Science Foundation grant PCEFP2_181115. This work is a contribution to the LEMONTREE (Land Ecosystem Models based On New Theory, obseRvations and ExperimEnts) project, funded through the generosity of Eric and Wendy Schmidt by recommendation of the Schmidt Futures program (K.H., B.D.S.).

References

Bloomfield, K. J., van Hoolst, R., Balzarolo, M., Janssens, I. A., Vicca, S., Ghent, D., and Prentice, I. C.: Towards a General Monitoring System for Terrestrial Primary Production: A Test Spanning the European Drought of 2018, Remote Sensing, 15, 1693, https://doi.org/10.3390/rs15061693, 2023.

Clark, J. S.: Why environmental scientists are becoming Bayesians: Modelling with Bayes, Ecology Letters, 8, 2–14, https://doi.org/10.1111/j.1461-0248.2004.00702.x, 2004.

Cowan, I. R. and Farquhar, G. D.: Stomatal function in relation to leaf metabolism and environment. In: Integration of Activity in the Higher Plant, 471–505, 1977.

Dietze, M. C.: Ecological Forecasting, Princeton University Press, https://doi.org/10.2307/j.ctvc7796h, 2017.

Dietze, M. C., Lebauer, D. S., and Kooper, R.: On improving the communication between models and data, Plant, Cell & Environment, 36, 1575–1585, https://doi.org/10.1111/pce.12043, 2013.

Dietze, M. C., Fox, A., Beck-Johnson, L. M., Betancourt, J. L., Hooten, M. B., Jarnevich, C. S., Keitt, T. H., Kenney, M. A., Laney, C. M., Larsen, L. G., Loescher, H. W., Lunch, C. K., Pijanowski, B. C., Randerson, J. T., Read, E. K., Tredennick, A. T., Vargas, R., Weathers, K. C., and White, E. P.: Iterative near-term ecological forecasting: Needs, opportunities, and challenges, Proc. Natl. Acad. Sci. U.S.A., 115, 1424–1432, https://doi.org/10.1073/pnas.1710231115, 2018.

Farquhar, G. D., Caemmerer, S., and Berry, J.: A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species, Planta, 149, 78–90, 1980.

Franklin, O., Harrison, S. P., Dewar, R., Farrior, C. E., Brännström, Å., Dieckmann, U., Pietsch, S., Falster, D., Cramer, W., Loreau, M., Wang, H., Mäkelä, A., Rebel, K. T., Meron, E., Schymanski, S. J., Rovenskaya, E., Stocker, B. D., Zaehle, S., Manzoni, S., Van Oijen, M., Wright, I. J., Ciais, P., Van Bodegom, P. M., Peñuelas, J., Hofhansl, F., Terrer, C., Soudzilovskaia, N. A., Midgley, G., and Prentice, I. C.: Organizing principles for vegetation dynamics, Nat. Plants, 6, 444–453, https://doi.org/10.1038/s41477-020-0655-x, 2020.

Gelman, A. and Rubin, D. B.: Inference from Iterative Simulation Using Multiple Sequences, Statist. Sci., 7, https://doi.org/10.1214/ss/1177011136, 1992.

Harrison, S. P., Cramer, W., Franklin, O., Prentice, I. C., Wang, H., Brännström, Å., de Boer, H., Dieckmann, U., Joshi, J., Keenan, T. F., Lavergne, A., Manzoni, S., Mengoli, G., Morfopoulos, C., Peñuelas, J., Pietsch, S., Rebel, K. T., Ryu, Y., Smith, N. G., Stocker, B. D., and Wright, I. J.: Eco-evolutionary optimality as a means to improve vegetation and land-surface models, New Phytologist, 231, 2125–2141, https://doi.org/10.1111/nph.17558, 2021.

Hartig, F., Dyke, J., Hickler, T., Higgins, S. I., O'Hara, R. B., Scheiter, S., and Huth, A.: Connecting dynamic vegetation models to data - an inverse perspective: Dynamic vegetation models - an inverse perspective, J. Biogeogr., 39, 2240–2252, https://doi.org/10.1111/j.1365-2699.2012.02745.x, 2012.

Hartig, F., Minunno, F., Paul, S., Cameron, D., Ott, T., and Pichler, M.: BayesianTools: General-Purpose MCMC and SMC Samplers and Tools for Bayesian Statistics, 2023.

LeBauer, D. S., Wang, D., Richter, K. T., Davidson, C. C., and Dietze, M. C.: Facilitating feedbacks between field measurements and ecosystem models, Ecological Monographs, 83, 133–154, 2013.

Marquet, P. A., Allen, A. P., Brown, J. H., Dunne, J. A., Enquist, B. J., Gillooly, J. F., Gowaty, P. A., Green, J. L., Harte, J., Hubbell, S. P., O'Dwyer, J., Okie, J. G., Ostling, A., Ritchie, M., Storch, D., and West, G. B.: On Theory in Ecology, BioScience, 64, 701–710, https://doi.org/10.1093/biosci/biu098, 2014.

Medlyn, B. E., Dreyer, E., Ellsworth, D., Forstreuter, M., Harley, P. C., Kirschbaum, M. U. F., Le Roux, X., Montpied, P., Strassemeyer, J., Walcroft, A., Wang, K., and Loustau, D.: Temperature response of parameters of a biochemically based model of photosynthesis. II. A review of experimental data, Plant, Cell & Environment, 25, 1167–1179, https://doi.org/10.1046/j.1365-3040.2002.00891.x, 2002.

Mengoli, G., Agustí-Panareda, A., Boussetta, S., Harrison, S. P., Trotta, C., and Prentice, I. C.: Ecosystem Photosynthesis in Land-Surface Models: A First-Principles Approach Incorporating Acclimation, Journal of Advances in Modeling Earth Systems, 14, e2021MS002767, https://doi.org/10.1029/2021MS002767, 2022.

Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K. J., Cinti, B. D., Grandcourt, A. de, Ligne, A. D., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., Tommasi, P. di, Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., et al.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, Sci Data, 7, 225, https://doi.org/10.1038/s41597-020-0534-3, 2020.

Prentice, I. C., Dong, N., Gleason, S. M., Maire, V., and Wright, I. J.: Balancing the costs of carbon gain and water transport: testing a new theoretical framework for plant functional ecology, Ecol Lett, 17, 82–91, https://doi.org/10.1111/ele.12211, 2014.

Rambal, S., Joffre, R., Ourcival, J. M., Cavender-Bares, J., and Rocheteau, A.: The growth respiration component in eddy CO_2 flux from a Quercus ilex mediterranean forest, Global Change Biology, 10, 1460–1469, https://doi.org/10.1111/j.1365-2486.2004.00819.x, 2004.

Stocker, B. D., Wang, H., Smith, N. G., Harrison, S. P., Keenan, T. F., Sandoval, D., Davis, T., and Prentice, I. C.: P-model v1.0: an optimality-based light use efficiency model for simulating ecosystem gross primary production, Geoscientific Model Development, 13, 1545–1581, https://doi.org/10.5194/gmd-13-1545-2020, 2020.

Tan, S., Wang, H., Prentice, I. C., and Yang, K.: Land-surface evapotranspiration derived from a first-principles primary

production model, Environ. Res. Lett., 16, 104047, https://doi.org/10.1088/1748-9326/ac29eb, 2021.

Ter Braak, C. J. F. and Vrugt, J. A.: Differential Evolution Markov Chain with snooker updater and fewer chains, Stat Comput, 18, 435–446, https://doi.org/10.1007/s11222-008-9104-9, 2008.

Trotsiuk, V., Hartig, F., and Forrester, D. I.: r3PG – An R package for simulating forest growth using the 3-PG process-based model, Methods Ecol. Evol., 11, 1470–1475, https://doi.org/10.1111/2041-210X.13474, 2020.

Van Oijen, M., Rougier, J., and Smith, R.: Bayesian calibration of process-based forest models: bridging the gap between models and data, Tree Physiology, 25, 915–927, https://doi.org/10.1093/treephys/25.7.915, 2005.

Wang, H., Prentice, I. C., Davis, T. W., Keenan, T. F., Wright, I. J., and Peng, C.: Photosynthetic responses to altitude: an explanation based on optimality principles, New Phytol, 213, 976–982, https://doi.org/10.1111/nph.14332, 2017a.

Wang, H., Prentice, I. C., Keenan, T. F., Davis, T. W., Wright, I. J., Cornwell, W. K., Evans, B. J., and Peng, C.: Towards a universal model for carbon dioxide uptake by plants, Nature Plants, 3, 734–741, https://doi.org/10.1038/s41477-017-0006-8, 2017b.

Weng, E. S., Malyshev, S., Lichstein, J. W., Farrior, C. E., Dybzinski, R., Zhang, T., Shevliakova, E., and Pacala, S. W.: Scaling from individual trees to forests in an Earth system modeling framework using a mathematically tractable model of height-structured competition, Biogeosciences, 12, 2655–2694, https://doi.org/10.5194/bg-12-2655-2015, 2015.

Xiang, Y., Gubian, S., Suomela, B., and Hoeng, J.: Generalized Simulated Annealing for Global Optimization: The GenSA Package, The R Journal, 5, 13, https://doi.org/10.32614/RJ-2013-002, 2013.