Serotonergic mediation of orienting and defensive responses in zebrafish

Bianca Gomes do Nascimento[a], Jeane Rodrigues Rodrigues[b,c], Maria Clara Rosa Silva[c,d], Bruna Patrícia Dutra Costa[a,b], Marissol Leite[c,d], Maryana Pereira Pyterson[a], Hadda Tercya Lima Silva[d], Loanne Valeria Xavier Bruce de Souza[a,e], Diógenes Henrique de Siqueira Silva[b,c,d], Caio Maximino[a,b,e*]

[a] Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff", Faculdade de Psicologia, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil; [b] Rede de Biotecnologia e Biodiversidade da Amazônia Legal, Marabá/PA, Brazil; [c] Programa de Pós-Graduação em Reprodução Animal na Amazônia (ReproAmazon), Universidade Federal Rural da Amazônia / Universidade Federal do Pará, Belem/PA, Brazil; [d] Grupo de Estudos da Reprodução de Peixes Amazônicos, Faculdade de Ciências Biológicas, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil; [e] Programa de Pós-Graduação em Neurociências e Comportamento, Universidade Federal do Pará, Belem/PA, Brazil

*corresponding author:

Caio Maximino
Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff",
Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do
Pará - Unidade III (Cidade Universitária), Rodovia BR-230, Av. Paulo Fonteles Filho,
s/n.º, Cidade Jardim Marabá/PA, Brazil

cmaximino@unifesspa.edu.br

Serotonergic mediation of orienting and defensive responses in zebrafish

Serotonin (5-HT) is involved in arousal and defensive responses, both of which represent modulators of attention and orienting. Orienting responses (ORs), considered as a unit of attentional processing, are elicited by novel innocuous stimuli, differ from defense reflexes (DR)s, as the latter are elicited by painful and threatening stimuli. When zebrafish (Danio rerio) are exposed to a conspecific alarm substance (CAS), visual stimuli elicit a DR instead of ORs, suggesting a state of hypervigilance. CAS elicited an OR-to-DR shift, as shown both by the shift from an approach to an escape response when the stimulus was turned on and by the changes in directional focus. Both pCPA and 5-HTP attenuated the CAS-elicited OR-to-DR shift . While pCPA did not alter ORs in animals which were not exposed to CAS, 5-HTP abolished those responses, suggesting that the serotonergic tone is important in regulating the general arousal levels in zebrafish.

Keywords: serotonergic tone; visual attention; general arousal; emotion-cognition interface.

Introduction

Orienting responses (ORs), originally described by Ivan Pavlov in the beginning of the 20th century as the "what is that?" reflex, are elicited by novel innocuous stimuli and can be considered a the unit of attentional processing (Barry, 2009). The OR is a whole-body reflex, involving adjustments in a large range of physiological systems, including visuomotor reflexes, electrodermal activity, respiratory and cardiovascular changes, and changes in whole-brain activity (Sokolov, 1960, 1963a, 1963b). While innocuous stimuli elicit ORs, painful and threatening stimuli elicit a separate class of responses, the defense reflex (Sokolov, 1960, 1963a, 1963b). In spite of that, situations of potential or distal threat can lead to hypervigilance, in which innocuous stimuli can lead to

defense reflexes (DRs). For example, in zebrafish (*Danio rerio*), exposure to an alarm

substance – an olfactory stimulus that represents a distal threat – leads to a shift from

ORs to DRs elicited by an innocuous visual stimulus (do Nascimento et al., 2020).

Arousal plays an important role in this OR-to-DR shift. The arousal level of an

organism can serve as an "amplifying factor", decreasing habituation (Sokolov, 1963a),

or, at higher levels, lead to "false alarms" in a state of hypervigilance. The neural bases

of this OR-to-DR shift probably involve the overlap between circuits for visual attention

and defensive behavior. For example, in a preparation of eye-brain-spinal cord of

lampreys (both *Lampetra fluviatilis* and *Petromyzon marinus*), recordings of the ventral

roots were able to discriminate between a unilateral burst response corresponding to

orientation of the head toward slowly expanding or moving stimuli, particularly within

the anterior visual field, that corresponds to a OR, and a unilateral or bilateral burst

response triggering avoidance in response to rapidly expanding looming stimuli or

moving bars, corresponding to a DR (Suzuki et al., 2019). These fictive responses were

enhanced by pharmacological blockade of brainstem-projecting neurons in the deep

layers of the optic tectum (OT)(Suzuki et al., 2019). Similarly, in goldfish (*Carassius*

auratus), unilateral tectal ablation abolishes the OR towards the contralateral hemifield,

while electrical stimulation of the OT evokes contraversive eye movements (Torres et

al., 2005). Habituation of ORs appear to depend on telencephalic input in goldfish, as

electrical stimulation of the dorsomedial (Dm) and dorsocentral (Dc) regions of the

telencephalon – the first being homologous to the associative amygdaloid nuclei of

mammals (Maximino et al., 2013) – evoke cardiac and ventilatory responses, as well as

bodily movements consistent with ORs (Quick & Laming, 1988). Likewise,

telencephalic ablation decreased OR habituation in goldfish (Rooney & Laming, 1988),

and animals with lesions in supracommissural (Vs) and postcommissural (Vp) regions

of the telencephalon showed DRs to innocuous stimuli instead of ORs (Rooney &

Laming, 1986). Thus, in fish, regions of the telencephalon that are part of the aversive

brain system (do Carmo Silva, Lima-Maximino, et al., 2018) project to regions of the

OT that organize ORs and DRs, modulating whether the animal will emit orienting or

defense responses.

Both the deeper layers of the OT that project to the brainstem and the

telencephalic regions involved in ORs and DRs show extensive serotonergic projections

in fish (Lillesaar et al., 2009; Yokogawa et al., 2012) and mammals (Mooney et al.,

1996). In rats, electrical stimulation of the superior colliculus (mammalian homologue

of the OT) elicits head movements at lower current intensities and running and jumping

responses at higher intensities; concurrent administration of the 5-HT reuptake blocker

fluoxetine or the 5-HT1A receptor antagonist WAY100,635 decreases these responses,

and administration of 5-HT in the midbrain, but not in the frontal cortex, produced the

same effect (Dringenberg et al., 2003). In zebrafish larvae, a brief exposure to water

flow elevates activity and sensitivity to behaviorally relevant visual cues that resemble

an OR (Yokogawa et al., 2012). These effects are accompanied by increased activity in

superior raphe (SR) neurons; genetic ablation of these cells abolished the increased

sensitivity to visual cues (Yokogawa et al., 2012). In a similar paradigm, exposure to a

conspecific alarm substance (CAS) induces a persistence hypervigilance in adult

zebrafish, as indexed by lowered threshold to respond to stimuli of different sensory

modalities; this state appears to be mediated by a synchronized state of neuronal

dynamics in the dorsocentral and dorsodorsal telencephali (putative homologues of the

mammalian cortex; Ganz et al., 2015; T. Mueller et al., 2011; Yáñez et al., 2022) and

CAS leads to an intense activation of neurons in the SR (Zhao et al., 2024). Transgenic

zebrafish lacking *tph2*, the rate-limiting enzyme in serotonin synthesis, do not show

CAS-elicited neuronal synchrony nor hypervigilance after CAS (Zhao et al., 2024).

Thus, serotonergic neurons of the SR appear to be involved in arousal-elicited

amplifications in vigilance in zebrafish.

We have previously shown that an innocuous visual stimulus elicits an orienting

response zebrafish, characterized by approaching the stimulus and projecting the body

towards it (do Nascimento et al., 2020). We have also determined that, during the OR,

the stimulus is investigated primarily with the left eye (do Nascimento et al., 2020),

consistent with lateralization of novelty processing in this species (Miklósi et al., 1997).

However, when animals were exposed to CAS – a mixture released in the water when

club cells in the skin are damaged, signaling threat to conspecifics in a shoal (Maximino

et al., 2019) –, this OR changes to a defense reflex, characterized by increased erratic

swimming and escaping the stimulus. We have previously demonstrated a role for

serotonin in other components of the alarm response: after exposure to CAS,

extracellular 5-HT levels are elevated in zebrafish brains (Maximino et al., 2014) and

monoamine oxidase activity is decreased (Lima-Maximino et al., 2020). Zhao et al.

(2024) observed serotonin release in the dorsocentral telencephalon (Dc) after CAS

exposure. Moreover, some components of the alarm response could be blocked by

treatment with *para*-chlorophenylalanine (pCPA), a false substrate of tryptophan

hydroxylase (TPH) which blocks serotonin synthesis, or metergoline, a non-selective

antagonist at 5-HT receptors (Lima-Maximino et al., 2020). Similar effects were

observed in transgenic zebrafish which do not express *tph2* (Zhao et al., 2024). The

present work attempts to unravel the roles of 5-HT in this arousal-elicited OR-to-DR

shift in adult zebrafish. In order to do that, animals were treated with either *para*-

chlorophenylalanine (pCPA), a false substrate of tryptophan hydroxylase which blocks

serotonin synthesis, or 5-hydroxytryptophan (5-HTP), a 5-HT precursor. It was

hypothesized that pCPA-treated animals would display an attenuated effect of CAS,

while 5-HTP would potentiate this effect.

Methods

Animals and housing

48 adult wild-type zebrafish from the longfin phenotype were used in the pCPA

experiments, and 40 animals were used in the 5-HTP experiments. Sample sizes were

not calculated prior to the experiments; however, *a posteriori* power calculations were

made, and are reported with each statistical analysis. The populations used are expected

to better represent the natural populations in the wild, due to its heterogeneous genetic

background (Parra et al., 2009; Speedie & Gerlai, 2008). Animals were bought from a

commercial vendor (Belém/PA) and collectively maintained in 40 L tanks for at least

two weeks before the onset of experiments. The sex proportion was estimated to be of 1

male:1 female, as judged by body morphology; however, sex was not determined either

through body nor gonadal morphology for each individual in the experiments, and

therefore was not included as a variable. Animals were acquired from commercial

vendors (Fernando Peixes, Belém/PA, and PisciculturaPower Fish, Itaguaí/RJ), and

delivered to the laboratory with an approximate age of 3 months (standard length = 12.9

± 1.6 mm), being quarantined for at least two weeks before experiments begun.

Throughout experiments, animals were estimated to be 4 months old (standard length =

21.7 ± 3.1 mm) The animals were fed daily with fish flakes. The tanks were kept at

constant temperature (28 °C), oxygenation, light cycle (14:10 LD photoperiod) and a

pH of 7.0-8.0, according to standards of care for zebrafish (Lawrence, 2007). Animals

were used for only one experiment to reduce interference from apparatus exposure.

Potential suffering of animals was minimized by controlling for the aforementioned

environmental variables. Furthermore, in the all experiments the animals used were

handled, anesthetized and sacrificed according to the norms of the Brazilian Guideline

for the Care and Use of Animals for Scientific and Didactic Purposes (CONCEA,

2017). Since, in all cases, animals were individually stressed and/or individually treated,

experimental unit was a single animal, and therefore all sample sizes refer to number of

animals. The experimental protocols were approved by the Animal Use Ethics

Committee of UNIFESSPA (Universidade Federal do Sul e do Sudeste do Pará)

protocol number 23479.019576/2022-43.

Drug treatments

For drug treatment, animals were injected intraperitoneally with either vehicle (DMSO

10%), pCPA (150 mg/kg), or 5-HTP (5 mg/kg), in a repeated dosing schedule; animals

were injected once per day for two days, followed by 24 h without drug injection, after

which they were subjected to the behavioural tests. This treatment schedule is typically

used in studies using pCPA to induce serotonin depletion in the central nervous system,

and has been shown to reduce 5-HT levels in the zebrafish brain to ~10% of controls

(Maximino et al. 2013). After injection, animals were returned to collective tanks

(maximum of 12 animals/tank); as a result, at least 4 different tanks were used per

experiment, which helped to prevent pseudo-replication. Tank number was added in

statistical analysis as a dummy variable, but showed no effect at $p < 0.05$. Animals were

randomly allocated to groups using a random number generator

(http://www.jerrydallal.com/random/random_block_size_r.htm), with each subject

randomized to a single (vehicle/drug) treatment using random permuted blocks. For

each experiment, animals were treated and tested in the order of allocation (i.e.,

randomly). In all experiments, experimenters and data analysts were blinded to drugs by

using coded vials (with the same code used for randomization); blinding was removed

only after data analysis.

Exposure to alarm substance (CAS)

For the exposure stage, each animal was transferred individually to a container (2 L)

where, after 3 minutes of acclimatization, it was carefully exposed to 7 ml of alarm

substance (CAS), extracted using a standardized protocol (do Carmo Silva, Rocha, et

al., 2018). Briefly, donor animals (size-matched to animals which were exposed to

CAS) were cold-anesthetized, sacrificed by spinal section, and CAS was extracted by

making 15 shallow cuts on each size of the carcass, avoiding drawing blood, and

washing the cuts with 10 ml double distilled water. CAS from a single donor was used

per animal, in a nominal concentration of 7 ml CAS/5 L water; CAS was always fresh-

prepared, and delivered directly into the apparatus water. As a negative control, a group

with the same amount of animals was exposed to the same volume of distilled water,

according to the protocol of Lima-Maximino et al. (2020). The animals remained

exposed for the entire session. All stages of the experiment were performed under

constant white Gaussian noise, producing an average of 58 dB above the tank. Light

levels above the tanks were measured using a handheld light meter, and ranged from

251 to 280 lumens (coefficient of variation = 3.399% between subjects).

Animals were randomly allocated to groups using a random number generator

(http://www.jerrydallal.com/random/random_block_size_r.htm), with each subject

randomized to a single treatment using random permuted blocks. For each experiment,

animals were treated and tested in the order of allocation (i.e., randomly). In all

experiments, experimenters and data analysts were blinded to drugs and treatment by

using coded vials (with the same code used for randomization); thus, results were coded

for both drug treatment (using randomly assigned letters for each drug) and exposure

(using randomly assigned numbers for either water or CAS). Blinding was removed

only after data analysis. Experiments were always run between 08:00AM and 02:00

PM. After experiments, animals were sacrificed by prolonged bath in ice-cold water (<

12 ºC), followed by spinal transection (Matthews & Varga, 2012).

Stimulus presentation

A blue circle (R: 0; G: 0; B: 128; maximum 256) was used, based on the ability of

zebrafish to discriminate blue from green stimuli in a simple discrimination paradigm

(K. P. Mueller & Neuhauss, 2012). Stimulus size was 72º of horizontal visual angle,

calculated considering the diameter of the stimulus in the screen when it is at the front

of the animal, and the animal is in the centre of the tank. The size was based on previous

work which show that this size is insufficient to produce an escape response in adult

zebrafish that were not exposed to CAS (do Nascimento et al., 2020).

Animals were transferred to the tank and left to acclimate for 3 min before the

presentation of the stimuli. The computer screen was turned on for the entire

experiment, including acclimatization time. Using an animation based on LibreOffice

Impress (v. 6.0.7.3), the stimulus was presented in 1-min intervals ("trials") interspersed

with 1-min stimulus-free periods. In trials in which the stimulus was on, the stimulus

was presented for the entire duration of the trial. 10 trials were made; total session

length was 22 minutes per animal. The number of trials and intertrial durations were

chosen based on previous work on arousal in goldfish (Laming & Savage, 1980).

Quantification of orienting and defensive responses

The experimental setup consisted of two $13 \times 13 \times 17$ (depth × length × depth cm each)

glass tanks, positioned side by side, and filled with 2 L tank water. A mobile phone

(Redmi Note® 10S, camera resolution 1080p at 30 fps) was positioned above the tanks

to allow recording and offline behavioral scoring. All focal fish behaviors were

therefore tracked from a top-down perspective, using both TheRealFishTracker

(http://www.dgp.toronto.edu/~mccrae/projects/FishTracker/) and fishtracker

(https://github.com/joseaccruz/fishtracker). For each behavioral video, a 2D region

(arena) was defined for tracking, comprising the inner area of the tank. A region-of-

interest (ROI) of 13 x 8.5 cm was determined as "stimulus zone". The following

variables were recorded, using TheRealFishTracker:

• Total time in the stimulus zone (s);

• Absolute turn angle (º), a measure of erratic swimming (Lima-Maximino et al.,

2020);

• Speed (cm/s)

Following Abril-de-Abreu, Cruz, & Oliveira (2015), a region of interest (ROI)

of 13 x 3 cm (approximately 25% of the tank), corresponding to the width of the tank

and the mean body length of an adult zebrafish, was defined in the area of the arena

closer to the computer screen in fishtracker. Animal orientation is defined as a mean

projection vector R_{proj} (Abril-de-Abreu et al., 2015), determined by the transformation of

each orientation angle in a vector $r_i = (cos\alpha_i, sin\alpha_i)$, where αi represents the angle

formed by the direction of the centroid-head axis relative to the horizontal axis in each

frame. The resulting mean vector, with length $R = \|r\|$, is a measure of directional focus,

and varies between 0 and 1. The projection of this vector R in the stimulus direction

(arbitrarily defined as 180º) is defined as $R_{proj} = -Rcos\alpha$. Since R_{proj} is a mean vector –

that is, it takes projection angles in each frame and averages then through the whole 1-

min trial –, if the animal is consistently orienting throughout most of the trial, values

tend to be positive; if the animal is consistently facing away from the stimulus, values

tend to be negative; and if the animal is orienting towards many different directions

during the trial, values tend to be close to 0. Thus, positive values indicate direction

towards the stimulus, while negative values indicate direction away from the stimulus,

and values close to 0 indicate absence of orienting.

Eye use was estimated by analysing projection angles averaged across trials in

which the stimulus was on and trials in which the stimulus was off, with angles between

150º and 179º recorded as "left eye use" ("L"), angles between 181º and 210º recorded

as "right eye use" ("R"), and average angles above 210º and below 150º recorded as "no

preference". Thus, eye use was defined as orienting the head in an angle of up to 30º,

consistent with what is observed in other experiments on eye use in zebrafish (do

Nascimento et al., 2020; Miklósi et al., 1997; Miklósi & Andrew, 1999).

Statistical analysis

For all data except directional angle (R), three-way (treatment X drug X trial type)

analyses of variance were applied. When p-values were smaller than 0.05, Tukey's HSD

post-hoc tests were applied. For directional angle data, circular means were analyzed

using projected normal circular generalized linear models; data were compared based on

the posterior distribution of the circular mean for each condition, based on lower and

upper intervals of the highest posterior density interval (HPD)(Cremers & Klugkist,

2018). Data and scripts can be found at

Results

Experiment 1

Significant main effects of trial type, treatment, and drug were found for the time near

the stimulus (Table 1; Figure 1A). In general, animals spent almost half of the each 1-

min. trial near the stimulus when it was turned on (24.07 ± 15.78 s with stimulus on vs.

12.54 ± 8.65 s with stimulus off [mean $\pm$ S.D.]); thus, when the stimulus was turned on,

animals approached it. Significant interactions were also found between drug and

treatment, drug and trial type, treatment and trial type, and a three-way interaction

between all variables (Table 1). Post-hoc Tukey's HSD tests found that the approach

response to the stimulus was abolished in the presence of CAS ($d = 1.13$, $p < 0.001$); in

fact, it is apparent that the approach response is turned into an escape response in the

presence of CAS, as individuals in this condition spend less time near the stimulus when

it was turned on than when it was turned off. pCPA abolished this effect, as individuals

treated with the drug and exposed to CAS spent more time near the stimulus when it

was turned on than when it was turned off ($d = -4.81$, $p < 0.001$). In summary, when the

stimulus was turned on, control animals approached it, and CAS-exposed animals

escaped it, and this last effect was prevented by pCPA.

Significant main effects of treatment, and drug were found for the erratic

swimming (Table 1; Figure 1B). In general, animals exposed to CAS showed higher

erratic swimming than non-exposed individuals at both trial types ($20.87º \pm 6.66$ for

controls vs. $26.49º \pm 9.51$ for CAS-exposed animals), producing a medium-sized effect

($d = -0.75$, $p < 0.001$). A significant interaction between drug and treatment was also

found (Table 1). Vehicle-treated, CAS-exposed animals showed increased erratic

swimming in relation to controls (d = -1.56, p < 0.001), while pCPA-treated animals did

not show differences in relation to controls when not exposed to CAS (d = -0.12, p =

0.59), and showed lower erratic swimming in relation to vehicle-treated, CAS-exposed

animals (d = 1.44, p < 0.001). In summary, CAS increased erratic swimming, an effect

that was prevented by pCPA and independent of stimulus presentation.

Significant effects of treatment and drug were found for swimming speed, but

interaction effects were absent (Table 1, Figure 1C). Treatment with pCPA generally

decreased swimming speed (d = 0.17, p = 0.01), while exposure to CAS increased it (d

= -2.85, p = 0.004).

By examining the lower and upper bounds of the 95% highest posterior density

interval (HPD) in Table 2, it is apparent that the HPD intervals of the circular mean of

all conditions overlap, and thus we conclude that there is not enough evidence to reject

the null hypothesis that the circular means do not differ and that there is no effect of

treatment, drug, or trial type on the average directional angle (Figure 1D). In summary,

stimulus presentation (on trials) elicited an orientation towards the stimulus in control

animals and away from it in CAS-exposed animals, and this latter effect was prevented

by pCPA.

Significant main effects of trial type and treatment were found for directional

focus (Table 1, Figure 1D). Significant treatment X drug, trial type X drug, trial type X

treatment, and three-way interactions were also found (Table 1). In general, directional

focus increased when the stimulus was turned on (d = -0.5, p = 0.017). When vehicle-

injected animals were exposed to CAS, directional focus was negative when the

stimulus was turned on (d = 1.7, p = 0.002 vs. stimulus off; d = 3.09, p < 0.00 vs.

CTRL). Treatment with pCPA abolished this effect.

Finally, significant associations between eye use and trial type, treatment, and

drug were found (χ^2[df = 14] = 85.033, p < 0.0001). No association was found when the

stimulus was turned off, but significant associations were found when the stimulus was

turned on. In these latter cases, control-exposed individuals treated with vehicle used

more the left eye, and control-exposed individuals treated with pCPA used more the

right eye; CAS-exposed individuals treated with vehicle showed no eye preference, and

CAS-exposed individuals treated with pCPA used more the left eye. In summary,

control individuals preferred to inspect the stimulus with the left eye, while pCPA

switches this preference for the right eye. pCPA also prevented the tendency in CAS-

exposed individuals to not display eye preferences.

Experiment 2

Main effects of trial type, treatment, and drug were found for the time spent near the

stimulus zone in Experiment 2 (Table 3, Fig 1A). Interactions between stimulus type

and treatment, drug and treatment, and three-way interactions were found. In general,

CAS-exposed animals treated with vehicle showed decreased time in the stimulus area

when the stimulus was turned on (d = 0.6, p = 0.003), while control animals treated with

vehicle showed more time in the stimulus area when the stimulus was turned on (d = -

0.93, p < 0.001). Again, presentation of the stimulus elicited an approach response in

control animals, and an escape response in CAS-exposed animals. This effect was

attenuated by treatment with 5-HTP (d =-0.4, p = 0.356 vs. CAS-exposed animals with

stimulus on); however, 5-HTP also appeared to abolish the OR in animals which were

not exposed to CAS (d = 0.18, p = 0.93, CTRL animals treated with 5-HTP, stimulus off

vs. on). In summary, while the effects of trial type and CAS from Experiment 1 were

replicated, the effects of 5-HTP on both controls and CAS-exposed individuals is

consistent with an attenuation in responsiveness.

Main effects of treatment, but not drug nor trial type, were found for erratic

swimming (Table 3, Figure 2B). Drug X treatment interactions were also found. In

general, CAS-exposed animals had lower erratic swimming than non-exposed animals

(d = 0.34, p < 0.001), an effect that appears to be mainly driven by the higher levels of

erratic swimming in non-exposed, 5-HTP-treated animals regardless of stimulus type.

Significant main effects of drug and treatment were found for swimming speed

(Table 3, Figure 2C). Significant interactions were also found between drug and

treatment. 5-HTP-treated animals showed consistently lower swimming speeds than all

other groups, regardless of stimulus type.

Main effects of trial type and drug were found for directional focus (Table 4),

and interactions between trial type and drug and trial type and treatment were also

found. By examining the lower and upper bounds of the 95% highest posterior density

interval (HPD) in Table 4, no overlap in the HPD intervals of the circular mean of on

and off trials in CTRL animals (both treated with vehicle and 5-HTP), and thus we

conclude that, for animals not exposed to CAS, the average directional angles were

different in these conditions, with the body directed towards the stimulus in on trials

(Figure 2D).

Finally, significant associations between eye use and trial type, treatment, and

drug were found (χ^2[df = 14] = 23.87, p < 0.0001). No association was found when the

stimulus was turned off, but significant associations were found when the stimulus was

turned on. In these cases, control-exposed individuals treated with vehicle used more

the left eye, and control-exposed individuals treated with 5-HTP lost this preference;

CAS-exposed individuals treated with vehicle showed no eye preference, and CAS-

exposed individuals treated with 5-HTP used more the right eye.

Discussion

The present work tested the participation of serotonin on orienting responses of

zebrafish, as well as in the shift from orienting response to defense reaction (OR-to-DR

shift) elicited by a conspecific alarm substance (CAS). pCPA blocked CAS-elicited OR-

to-DR shift, as shown both by the shift from an approach to an escape response when

the stimulus was turned on and by the changes in directional focus. pCPA had no effects

on erratic swimming; importantly, erratic swimming was not significantly altered by

trial type, only by CAS, suggesting again that this stimulus induces an aversive state

that is unrelated to visual stimulation, while simultaneously leading to the OR-to-DR

shift. 5-HTP, on the other hand, attenuated the CAS-elicited OR-to-DR shift and

abolished ORs in non-exposed animals, suggesting that the serotonergic tone is more

important in regulating the general arousal levels. These results are consistent with the

prediction that pCPA would attenuate the OR-to-DR shift, but not the prediction that 5-

HTP would potentiate it.

The OR observed in the present experiments essentially replicates previous

findings from our group (do Nascimento et al., 2020). The OR was characterized by an

approach response (more time spent near the stimulus when it was turned on) and a

body oriented towards the stimulus (higher directional focus), with predominant use of

the left eye to inspect the stimulus. We also replicated our previous observation that

CAS elicits an OR-to-DR shift in zebrafish (do Nascimento et al., 2020). This shift was

characterized by avoiding the stimulus zone when the visual stimulus was turned off

and reduced directional focus. These effects are consistent with a defensive reaction,

with CAS leading to an attentional bias in which the animal "misinterprets" the

innocuous visual stimulus as a threatening stimulus. Evidence for an aversive state

elicited by CAS is also present at the increased, stimulus-independent elevation in

erratic swimming.

Importantly, we found that pCPA abolished the OR-to-DR shift in CAS-exposed

animals. This effect was also accompanied by the abolishing of the effect of CAS on

erratic swimming, suggesting that, while the serotonergic tone is permissive to ORs

elicited by visual non-threatening stimuli, it is also necessary to mount a defensive

response. This is consistent with previous findings with zebrafish, in which pCPA

blocked the effects of CAS on responses to distal and proximal threat (Lima-Maximino

et al., 2020). These findings are also consistent with previous work which shows that

transgenic zebrafish that do not express *tph2* are not sensitive to arousal-eliciting effects

of CAS (Zhao et al., 2024). The OR-attenuating effect of pCPA was also observed in

rats, in which the drug significantly reduced the responding to the first presentation of a

tone, but did not alter the rate of habituation (File, 1975). On the other hand, pCPA has

been shown to enhance startle responses in rats (Carlton & Advokat, 1973), suggesting

that the role of serotonin in orienting and defensive responses can be opposite.

We initially hypothesized that 5-HTP would produce an opposite effect of pCPA

in the OR-to-DR shift. However, we found that 5-HTP attenuated both the OR (in

control animals) and the OR-to-DR shift (in CAS-exposed animals). 5-HTP also

decreased swimming speed across treatments and exposure regimens, suggesting either

a psychomotor effect or decreased responsiveness. Indeed, serotonin has been shown to

suppress locomotion in zebrafish (Gabriel et al., 2009; Tran et al., 2016). An hypothesis

to explain these effects is that the vigilance-promoting effects of serotonin are

dependent on baseline levels of this transmitter, and increasing these levels above a

certain threshold do not promote more arousal. In rhesus macaques, 5-HTP increases

attention in animals that exhibit low baseline attention, while decreasing attention in

animals that exhibit high baseline attention; importantly, this effect is associated with

baseline concentrations of 5-HTP in the cerebrospinal fluid, with high baseline attention

being associated with lower 5-HTP levels (Weinberg-Wolf et al., 2018). While our

design does not allow to separate between baseline and post-drug effects, it is possible

that the serotonergic tone is permissive to ORs elicited by non-threatening visual

stimuli, while phasic stimulation of serotonin are less important in that regard. We have

previously suggested that, for threatening stimuli, phasic serotonergic signals inhibit

fear-like responses, while the serotonergic tone tracks aversive expectation values

(Lima-Maximino et al., 2020). A similar mechanism might be involved in tracking

novelty and expectation values in orienting responses.

Surprisingly, pCPA affected eye use during the presentation of the stimulus. Our

results replicate previous results which show preference for the left eye in control

animals during the presentation of the stimulus, while CAS-exposed animals showed no

preference (do Nascimento et al., 2020). Moreover, we found that non-exposed

individuals treated with pCPA showed a preference for the right eye when viewing the

stimulus, while non-exposed animals treated with 5-HTP lost preference. When exposed

to CAS, pCPA-treated animals shifted preference to the left eye, while 5-HTP-treated

animals used more the right eye. In zebrafish, novel objects were investigated mainly

with right frontal field, while in a second trial left frontal viewing tended to be used

instead (Miklósi et al., 1997; Miklósi & Andrew, 1999). This is unlikely to be due to

asymmetry in the habenular nuclei, which has been linked to preference in eye use

(Dadda et al., 2010), because habenular projections to zebrafish raphe are not

asymmetrical (Amo et al., 2010). Miklósi et al. (1997) suggested that the right eye is

used when it is necessary to inhibit premature response in conditions of decision-

making, and the left is used when it is necessary to track familiar objects. While we

were not able to ascertain whether eye use changed during trials, it is possible that

serotonin helps to track novelty, which could explain the effects of pCPA and 5-HTP on

this variable.

Despite our findings that pCPA and 5-HTP altered the CAS-elicited OR-to-DR

shift, a more thorough investigation of the specific mechanisms by which phasic and

tonic serotonin modulate arousal and defensive responses to non-threatening visual

stimuli is still needed. Given the high number of different serotonin receptors and the

wide distribution of serotonergic innervation across the brain, the specific participation

of receptors and brain regions is far from known. Nonetheless, these results point to an

important serotonergic regulation of an emotion-cognition interface in vertebrates,

which could represent a mechanism through which individuals respond to threats.

Acknowledgments

This work was supported by a Conselho Nacional de Desenvolvimento Científico e Tecnológico

(CNPq/Brazil) grant (Edital Universal 2016, #400726/2016-6).

References

Abril-de-Abreu, R., Cruz, J., & Oliveira, R. F. (2015). Social Eavesdropping in

Zebrafish: Tuning of Attention to Social Interactions. Scientific Reports, 5, Artigo 1.

https://doi.org/10.1038/srep12678

Amo, R., Aizawa, H., Takahoko, M., Kobayashi, M., Takahashi, R., Aoki, T., &

Okamoto, H. (2010). Identification of the zebrafish ventral habenula as a

homolog of the mammalian lateral habenula. Journal of Neuroscience, 30, 1566–

1574. https://doi.org/10.1523/JNEUROSCI.3690-09.2010

Barry, R. J. (2009). Habituation of the orienting reflex and the development of

Preliminary Process Theory. Neurobiology of Learning and Memory, 92(2),

235–242. https://doi.org/10.1016/j.nlm.2008.07.007

Carlton, P. L., & Advokat, C. (1973). Attenuated Habituation Due to

Parachlorophenylalanine. 1, 657–663.

Cremers, J., & Klugkist, I. (2018). One Direction? A Tutorial for Circular Data Analysis

Using R With Examples in Cognitive Psychology. Frontiers in Psychology, 9,

2040. https://doi.org/10.3389/fpsyg.2018.02040

Dadda, M., Domenichini, A., Piffer, L., Argenton, F., & Bisazza, A. (2010). Early

differences in epithalamic left–right asymmetry influence lateralization and

personality of adult zebrafish. Behavioural Brain Research, 206(2), 208–215.

https://doi.org/10.1016/j.bbr.2009.09.019

Diretriz brasileira para o cuidado e a utilização de animais para fins científicos e

didáticos - DBCA. Anexo I. Peixes mantidos em instalações de instituições de

ensino ou pesquisa científica, Resolução Normativa CONCEA no 34 218

(2017).

https://www.mctic.gov.br/mctic/opencms/legislacao/outros_atos/resolucoes/

Resolucao_Normativa_CONCEA_n_34_de_27072017.html

do Carmo Silva, R. X., Lima-Maximino, M. G., & Maximino, C. (2018). The aversive

brain system of teleosts: Implications for neuroscience and biological

psychiatry. Neuroscience & Biobehavioral Reviews, 95, 123–135.

https://doi.org/10.1016/j.neubiorev.2018.10.001

do Carmo Silva, R. X., Rocha, S. P., Lima-Maximino, M. G., & Maximino, C. (2018).

Extraction of alarm substance in zebrafish. Protocols.Io.

https://doi.org/10.17504/protocols.io.tr3em8n

do Nascimento, B. G., Oliveira, H. S. T. O. e., Silva, H. T. L., de Siqueira-Silva, D. H.,

Lima-Maximino, M., & Maximino, C. (2020). A model to study orienting

responses in zebrafish, and applications towards the emotion–cognition

interaction. Animal Cognition, 23(5), 965–972. https://doi.org/10.1007/s10071-

020-01403-1

Dringenberg, H. C., Dennis, K. E. B., Tomaszek, S., & Martin, J. (2003). Orienting and

defensive behaviors elicited by superior colliculus stimulation in rats: Effects of

5-HT depletion, uptake inhibition, and direct midbrain or frontal cortex

application. Behavioural Brain Research, 144(1), 95–103.

https://doi.org/10.1016/S0166-4328(03)00065-2

File, S. E. (1975). Effects of parachlorophenylalanine and amphetamine on habituation

of orienting. Pharmacology Biochemistry and Behavior, 3(6), 979–983.

https://doi.org/10.1016/0091-3057(75)90005-2

Gabriel, J. P., Mahmood, R., Kyriakatos, A., Söll, I., Hauptmann, G., Calabrese, R. L.,

& Manira, A. E. (2009). Serotonergic Modulation of Locomotion in Zebrafish—

Endogenous Release and Synaptic Mechanisms. Journal of Neuroscience, 29,

10387–10395. https://doi.org/10.1523/JNEUROSCI.1978-09.2009

Ganz, J., Kroehne, V., Freudenreich, D., Machate, A., Geffarth, M., Braasch, I., Kaslin,

J., & Brand, M. (2015). Subdivisions of the adult zebrafish pallium based on

molecular marker analysis. F1000Research, 3, 308.

https://doi.org/10.12688/f1000research.5595.2

Laming, P. R., & Savage, G. E. (1980). Physiological changes observed in the goldfish

(Carassius auratus) during behavioral arousal. Behavioral and Neural Biology,

29, 255–275.

Lawrence, C. (2007). The husbandry of zebrafish (Danio rerio): A review. Aquaculture,

269(1), 1–20. https://doi.org/10.1016/j.aquaculture.2007.04.077

Lillesaar, C., Stigloher, C., Tannhäuser, B., Wullimann, M. F., & Bally-Cuif, L. (2009).

Axonal projections originating from raphe serotonergic neurons in the

developing and adult zebrafish, Danio rerio, using transgenics to visualize

raphe-specific pet1 expression. Journal of Comparative Neurology, 512(2), 158–

182. https://doi.org/10.1002/cne.21887

Lima-Maximino, M., Pyterson, M. P., Silva, R. X. do C., Gomes, G. C. V., Rocha, S. P.,

Herculano, A. M., Rosemberg, D. B., & Maximino, C. (2020). Phasic and tonic

serotonin modulate alarm reactions and post-exposure behavior in zebrafish.

Journal of Neurochemistry, 153(4), 495–509. https://doi.org/10.1111/jnc.14978

Matthews, M., & Varga, Z. M. (2012). Anesthesia and Euthanasia in Zebrafish. ILAR

Journal, 53(2), 192–204. https://doi.org/10.1093/ilar.53.2.192

Maximino, C., do Carmo Silva, R. X., Campos, K. dos S., Oliveira, J. S. de, Rocha, S.

P., Pyterson, M. P., Souza, D. P. dos S., Feitosa, L. M., Ikeda, S. R., Pimentel,

A. F. N., Ramos, P. N. F., Costa, B. P. D., Herculano, A. M., Rosemberg, D. B.,

Siqueira-Silva, D. H., & Lima-Maximino, M. (2019). Sensory ecology of

ostariophysan alarm substances. Journal of Fish Biology, 95(1), 274–286.

https://doi.org/10.1111/jfb.13844

Maximino, C., Lima, M. G., Costa, C. C., Guedes, I. M. L., & Herculano, A. M. (2014).

Fluoxetine and WAY 100,635 dissociate increases in scototaxis and analgesia

induced by conspecific alarm substance in zebrafish (Danio rerio Hamilton

1822). Pharmacology Biochemistry and Behavior, 124, 425–433.

https://doi.org/10.1016/j.pbb.2014.07.003

Miklósi, A., & Andrew, R. J. (1999). Right eye use associated with decision to bite in

zebrafish. Behavioural Brain Research, 105, 199–205.

https://doi.org/10.1016/S0166-4328(99)00071-6

Miklósi, A., Andrew, R. J., & Savage, H. (1997). Behavioural Lateralisation of the

Tetrapod Type in the Zebrafish (Brachydanio Rerio). Physiology & Behavior,

63, 127–135. https://doi.org/10.1016/S0031-9384(97)00418-6

Mooney, R. D., Huang, X., Shi, M.-Y., Bennett-Clarke, C. A., & Rhoades, R. W.

(1996). Chapter 4 Serotonin modulates retinotectal and corticotectal

convergence in the superior colliculus. Em M. Norita, T. Bando, & B. E. Stein

(Orgs.), Progress in Brain Research (Vol. 112, p. 57–69). Elsevier.

https://doi.org/10.1016/S0079-6123(08)63320-8

Mueller, K. P., & Neuhauss, S. C. F. (2012). Automated visual choice discrimination

learning in zebrafish (Danio rerio). Journal of Integrative Neuroscience, 11, 73–

85. https://doi.org/10.1142/S0219635212500057

Mueller, T., Dong, Z., Berberoglu, M. A., & Guo, S. (2011). The dorsal pallium in

zebrafish, Danio rerio (Cyprinidae, Teleostei). Brain Research, 1381, 95–105.

https://doi.org/10.1016/j.brainres.2010.12.089

Parra, K. V., Adrian, J. C., & Gerlai, R. (2009). The synthetic substance hypoxanthine

3-N-oxide elicits alarm reactions in zebrafish (Danio rerio). Behavioural Brain

Research, 205(2), 336–341. https://doi.org/10.1016/j.bbr.2009.06.037

Quick, I. A., & Laming, P. R. (1988). Cardiac, ventilatory and behavioural arousal

responses evoked by electrical brain stimulation in the goldfish (Carassius

auratus). Physiology & Behavior, 43, 715–727.

Rooney, D. J., & Laming, P. R. (1986). Localization of telencephalic regions concerned
with habituation of cardiac and ventilatory responses associated with arousal in
the goldfish (Carassius auratus). Behavioral Neuroscience, 100, 45–50.

Rooney, D. J., & Laming, P. R. (1988). Effects of telencephalic ablation on habituation
of arousal responses, within and between daily training sessions in goldfish.
Behavioral and Neural Biology, 49, 83–96.

Sokolov, E. N. (1960). Neuronal models and the orienting reflex. Em M. A. B. Brazier,
The central nervous system and behaviour (p. 187–276). Macey.

Sokolov, E. N. (1963a). Higher Nervous Functions: The Orienting Reflex. Annual
Review of Physiology, 25(1), 545–580.
https://doi.org/10.1146/annurev.ph.25.030163.002553

Sokolov, E. N. (1963b). Perception and the conditioned reflex. Pergamon Press.

Speedie, N., & Gerlai, R. (2008). Alarm substance induced behavioral responses in
zebrafish (Danio rerio). Behavioural Brain Research, 188(1), 168–177.
https://doi.org/10.1016/j.bbr.2007.10.031

Suzuki, D. G., Pérez-Fernández, J., Wibble, T., Kardamakis, A. A., & Grillner, S.
(2019). The role of the optic tectum for visually evoked orienting and evasive
movements. Proceedings of the National Academy of Sciences, 116, 15272–
15281. https://doi.org/10.1073/pnas.1907962116

Torres, B., Luque, M. Á., Pérez-Pérez, M. P., & Herrero, L. (2005). Visual orienting
response in goldfish: A multidisciplinary study. Brain Research Bulletin, 66,
376–380. https://doi.org/10.1016/j.brainresbull.2005.02.002

Tran, S., Nowicki, M., Muraleetharan, A., Chatterjee, D., & Gerlai, R. (2016).
Neurochemical factors underlying individual differences in locomotor activity
and anxiety-like behavioral responses in zebrafish. Progress in Neuro-
Psychopharmacology and Biological Psychiatry, 65, 25–33.
https://doi.org/10.1016/j.pnpbp.2015.08.009

Weinberg-Wolf, H., Fagan, N. A., Anderson, G. M., Tringides, M., Dal Monte, O., &
Chang, S. W. C. (2018). The effects of 5-hydroxytryptophan on attention and
central serotonin neurochemistry in the rhesus macaque.
Neuropsychopharmacology, 43, 1589–1598. https://doi.org/10.1038/s41386-
017-0003-7

Yáñez, J., Folgueira, M., Lamas, I., & Anadón, R. (2022). The organization of the

zebrafish pallium from a hodological perspective. Journal of Comparative

Neurology, 530, 1164–1194. https://doi.org/10.1002/cne.25268

Yokogawa, T., Hannan, M. C., & Burgess, H. A. (2012). The dorsal raphe modulates

sensory responsiveness during arousal in zebrafish. Journal of Neuroscience, 32,

15205–15215. https://doi.org/10.1523/JNEUROSCI.1019-12.2012

Zhao, Y., Huang, C.-X., Gu, Y., Zhao, Y., Ren, W., Wang, Y., Chen, J., Guan, N. N., &

Song, J. (2024). Serotonergic modulation of vigilance states in zebrafish and

mice. Nature Communications, 15, 2596. https://doi.org/10.1038/s41467-024-

47021-0

Table 1. Summary of statistical analysis on the variables *time near stimulus, erratic*

swimming, and *directional focus* for Experiment 1.

Endpoint	Independent variable	df	F	p	ω²	Power
Time near stimulus	Trial type (on vs. off)	1, 949	514.88	< 0.001	0.17	0.98
	Treatment (CTRL vs. CAS)	1, 949	11.3	< 0.001	0.00	0.1
	Drug (vehicle vs. pCPA)	1, 949	148.57	< 0.001	0.05	0.78
	Treatment x Drug	1, 949	303.64	< 0.001	0.10	0.97
	Trial type x Drug	1, 949	369.29	< 0.001	0.13	0.99
	Trial type x Treatment	1, 949	32.11	< 0.001	0.01	0.22
	Trial type x Treatment x Drug	1, 949	664.12	< 0.001	0.22	0.99
Erratic swimming	Trial type	1, 949	0.81	0.369	-0.01	0.1
	Treatment	1, 949	133.83	< 0.001	0.10	0.53
	Drug	1, 949	95.9	< 0.001	0.07	0.11
	Treatment x Drug	1, 949	152.03	< 0.001	0.11	0.74
	Trial type x Drug	1, 949	1.94	0.164	0.0	0.05
	Trial type x Treatment	1, 949	0.07	0.785	-0.0	0.05
	Trial type x Treatment x Drug	1, 949	0.03	0.868	-0.0	0.05
Swimming speed	Trial type	1, 949	0.01	0.918	-0.0	0.05
	Treatment	1, 949	8.11	0.004	0.01	0.05
	Drug	1, 949	6.63	0.01	0.01	0.05
	Treatment x Drug	1, 949	2.39	0.123	0.0	0.05
	Trial type x Drug	1, 949	0.7	0.403	-0.0	0.05
	Trial type x Treatment	1, 949	0.99	0.321	-0.0	0.05
	Trial type x Treatment x Drug	4, 949	0.54	0.461	-0.0	0.05

Directional focus (R_{proj})	Trial type	1, 88	5.89	0.017	0.03	0.43
	Treatment	1, 88	6.76	0.011	0.03	0.26
	Drug	1, 88	3.42	0.068	0.01	0.15
	Treatment x Drug	1, 88	8.91	0.004	0.05	0.37
	Trial type x Drug	1, 88	22.54	< 0.001	0.11	0.53
	Trial type x Treatment	1, 88	24.87	< 0.001	0.14	0.84
	Trial type x Treatment x Drug	4, 88	20.81	< 0.001	0.11	0.74

Table 2. Posterior estimates of the circular means of the projection angle for Experiment
1.

Treatment	Drug	Trial type	Mode	Mean ± S.D.	LB HPD	UB HPD
CTRL	VEH	off	1.873	1.858 ± 0.849	-0.214	3.779
		on	0.321	0.492 ± 2.282	-2.278	3.574
CAS	VEH	off	2.156	1.47 ± 2.731	-2.322	3.652
		on	-0.157	1.203 ± 2.503	2.714	2.284
CTRL	pCPA	off	2.662	2.797 ± 2.66	-1.146	4.701
		on	1.774	1.256 ± 2.662	-1.521	4.343
CAS	pCPA	off	3.201	2.115 ± 2.547	0.137	6.035
		on	1.021	1.246 ± 2.513	-1.689	4.126

Table 3. Summary of statistical analysis on the variables *time near stimulus*, *erratic
swimming,* and *directional focus* for Experiment 2.

Endpoint	Independent variable	df	F	p	ω^2	Power
Time near stimulus	Trial type	1, 664	16.28	< 0.001	0.02	0.13

	Treatment	1, 664	48.71	< 0.001	0.07	0.36
	Drug	1, 664	23.81	< 0.001	0.03	0.22
	Treatment x Drug	1, 664	4.95	0.026	0.01	0.05
	Trial type x Drug	1, 664	3.11	0.078	0.0	0.05
	Trial type x Treatment	1, 664	181.65	< 0.001	0.21	0.85
	Trial type x Treatment x Drug	1, 664	8.76	0.003	0.01	0.05
Erratic swimming	Trial type	1, 606	0.19	0.662	-0.0	0.05
	Treatment	1, 606	17.2	< 0.001	0.03	0.18
	Drug	1, 606	2.93	0.087	0.0	0.05
	Treatment x Drug	1, 606	8.15	0.004	0.01	0.05
	Trial type x Drug	1, 606	0.09	0.767	-0.0	0.05
	Trial type x Treatment	1, 606	0.07	0.785	-0.0	0.05
	Trial type x Treatment x Drug	1, 606	0.01	0.934	-0.0	0.05
Swimming speed	Trial type	1, 685	2.21	0.138	0.0	0.05
	Treatment	1, 685	15.38	< 0.001	0.02	0.13
	Drug	1, 685	78.9	< 0.001	0.09	0.49
	Treatment x Drug	1, 685	70.55	< 0.001	0.08	0.45
	Trial type x Drug	1, 685	1.0	0.317	0.0	0.05
	Trial type x Treatment	1, 685	1.52	0.217	0.0	0.05
	Trial type x Treatment x Drug	1, 685	0.0	0.97	-0.0	0.05
Directional focus (R_{proj})	Trial type	1, 88	5.502	0.0212	0.06	0.58
	Treatment	1, 88	3.033	0.085	0.03	0.31
	Drug	1, 88	25.501	< 0.001	0.225	0.98
	Treatment x Drug	1, 88	1.049	0.309	0.012	0.15

Trial type x Drug	1, 88	14.964	< 0.001	0.145	0.96
Trial type x Treatment	1, 88	6.091	0.015	0.065	0.62
Trial type x Treatment x Drug	1, 88	0.031	0.861	0.0	0.05

Table 4. Posterior estimates of the circular means of the directional angle for
Experiment 2.

Treatment	Drug	Trial type	Mode	Mean ± S.D.	LB HPD	UB HPD
CTRL	VEH	off	-1.543	-1.989 ± 2.574	0.522	0.066
		on	-1.449	-1.305 ± 0.616	-2.403	0.004
CAS	VEH	off	-2.147	-2.126 ± 0.24	-2.6	-1.62
		on	-0.703	-0.782 ± 0.745	-2.321	0.698
CTRL	5-HTP	off	-2.414	-2.552 ± 1.34	0.257	5.70
		on	-0.638	-0.677 ± 0.653	-2.009	0.68
CAS	5-HTP	off	-1.61	-1.57 ± 1.49	2.11	1.35
		on	-2.349	-2.795 ± 1.265	0.891	5.987

Figure 1. pCPA blocked CAS-elicited OR-to-DR shift. (A) Presenting zebrafish with a visual stimulus elicits approach to the stimulus zone, as shown by peaks in the time near stimulus in trials in which the stimulus was on. CAS inverts this approach response, decreasing time near the stimulus when the stimulus is turned on. pCPA treatment prevents the effects of CAS. (B) The visual stimulus does not alter erratic swimming, but CAS increases it, an effect that is attenuated by pCPA. (C) No effects of the visual stimulus, CAS, or pCPA were observed on swimming speed. (D) Zebrafish project their bodies towards the stimulus (position indicated by the letter "S") when it is on, while CAS leads animals to project their body away from the stimulus; pCPA abolishes the effects of CAS on the OR. In (A-C), red lines represent group averages for vehicle-treated animals, with pink ribbons representing 95% confidence intervals; blue lines

represent group averages for pCPA-treated animals, with light blue ribbons representing

95% confidence intervals. In (D), each dot represents average values for a single

individual, with the distance from the center of the polar plot representing the

directional focus R_{proj}, and its position in the circular plot represents the angle (in º) of

orientation; both variables are average across trial types (on vs. off).

Figure 2. 5-HTP attenuates OR responses in control zebrafish. (A) Presenting zebrafish

with a visual stimulus elicits approach to the stimulus zone, as shown by peaks in the

time near stimulus intrials in which the stimulus was on. CAS inverts this approach

response, decreasing time near the stimulus when the stimulus is turned on. 5-HTP

treatment decreases the approach in both control and CAS-exposed zebrafish. (B) The

visual stimulus does not alter erratic swimming, but CAS increases it, an effect that is

not altered by 5-HTP. (C) No effects of the visual stimulus or CAS were observed on

swimming speed, but 5-HTP treatment decreased swimming speed. (D) Zebrafish

project their bodies towards the stimulus (position indicated by the letter "S") when it is

on, while CAS leads animals to project their body away from the stimulus; 5-HTP

attenuates or abolishes this OR. In (A-C), red lines represent group averages for vehicle-

treated animals, with pink ribbons representing 95% confidence intervals; blue lines

represent group averages for pCPA-treated animals, with light blue ribbons representing

95% confidence intervals. In (D), each dot represents average values for a single

individual, with the distance from the center of the polar plot representing the

directional focus R_{proj}, and its position in the circular plot represents the angle (in º) of

orientation; both variables are average across trial types (on vs. Off).