Developmental trajectories and differences in functional brain network properties of preterm and at-term neonates

López-Guerrero N[a], Alcauter S[a]*

[a] Instituto de Neurobiología, Universidad Nacional Autónoma de México, Querétaro, México.

Abstract

Premature infants, born before 37 weeks of gestation can have consequences on neurodevelopment and cognition, even when no anatomical lesions are evident. Resting-state functional neuroimaging of naturally sleeping babies has shown altered connectivity patterns, but there is limited evidence on the developmental trajectories of the functional organization in preterm neonates. By using a large dataset (n=597) from the developing Human Connectome Project, we explored the differences in graph theory properties between at-term and preterm neonates at term-equivalent age, considering the age subgroups proposed by the World Health Organization for premature birth. Leveraging the longitudinal follow-up for some preterm participants, we characterized the developmental trajectories for preterm and at-term neonates. We found significant differences between groups for connectivity strength, clustering coefficient, characteristic path length and global efficiency. Specifically, at term-equivalent ages, higher connectivity, clustering coefficient and efficiency are identified for neonates born at higher postmenstrual ages. Similarly, the characteristic path length showed the inverse pattern. These results were consistent for a variety of connectivity thresholds and both at the global (whole brain) and local level (brain regions). The brain regions with the greatest differences between groups include primary sensory and motor regions and the precuneus which may relate with the risk factors for sensorimotor and behavioral deficits associated with premature birth.Our results also show non-linear developmental trajectories for premature neonates, but decreased integration and segregation even at term-equivalent age. Overall, our results confirm altered functional connectivity, integration and segregation properties of the premature brain despite showing rapid maturation after birth.

Keywords

Brain network; Connectome; Development; Preterm; Graph theory.

Introduction

According to the World Health Organization (WHO), preterm birth is a live birth that occurs before 37 weeks of gestational age (GA). The WHO classifies preterm birth according to weeks of gestation (WG): extremely preterm, born < 28 WG; very preterm, between 28 to 32 WG; and moderate to late preterm: 32 to 37 WG (*Preterm Birth*, 2022). In 2020, an estimated 13.4 million preterm infants were born, a slightly lower rate compared to 2010, where it was 13.8 million. However, over the last decade, the global annual rate of reduction is estimated at -0.14% and very few countries have achieved this reduction (Geneva: World Health & Organization, 2023; Ohuma et al., 2023). Mortality and morbidity risks increase as a function of the degree of prematurity, including the risk for neurodevelopmental deficits (Perin et al., 2022; World Health Organization, 2012,2023).

The developmental consequences of premature birth can be neurological, cognitive, behavioral, and emotional, even when no anatomical brain lesions are evident; the severity increases with decreasing gestational age at birth, and the sequelae can last into adulthood (Larsen et al., 2022; Ream & Lehwald, 2018; Rogers et al., 2018). A premature infant is likely to survive from 25 weeks of GA at a critical time of development and vulnerability to the extrauterine environment, which could influence the child's later development. Among the most frequent alterations are attention deficit hyperactivity disorder (ADHD), depression, anxiety, autism spectrum disorder (ASD), cerebral palsy (CP), antisocial personality, auditory and visual deficits, and an increased risk for cognitive, sensory, motor and language deficits (Hee Chung et al., 2020)

In recent years, neuroimaging methods have allowed the exploration of the brain *in vivo*; in particular, resting state functional Magnetic Resonance Imaging (rsfMRI) has made it possible to study the functional organization of the brain in fetus, preterm and term neonates (Smyser et al., 2016; Smyser & Neil, 2015; Thomason et al., 2015). Most rsfMRI studies in preterm infants have focused on characterizing the main brain networks in the resting state, showing that the primary networks are similar to those of

adults. In contrast, there appears to be no complete precursor of the default mode network (DMN) in premature infants at term-equivalent age (Doria et al., 2010; Eyre et al., 2021). When characterizing the brain as a complex network, empirical evidence shows that from 20 WG, the fetal brain follows a modular organization with the centers being densely interconnected, forming a rich club organization; with modules representing the areas that will give rise to the motor, visual, auditory, and the DMN, among others (Thomason et al., 2014). By 31 WG, the network organization starts to become more efficient, specialized, and integrated with a stronger connection between modules than towards the interior of modules; there is a decrease in local efficiency as age increases the clustering coefficient and shortest path length (De Asis-Cruz et al., 2021; Thomason et al., 2014; Turk et al., 2019). Modular structure, clustering coefficient, and shortest path length continuously change during the neonatal period (Gao et al., 2011). These network properties are also observed in premature infants at term-equivalent age, showing however, reduced rich club organization, less segregated networks, less integration and reduced global efficiency (Bouyssi-Kobar et al., 2019; Scheinost et al., 2016). Nevertheless, there is limited documentation regarding the functional organization in preterm neonates, and the characterization of developmental trajectories for this population has been challenging mainly due to small sample sizes.

In this study, leveraging a large public dataset, we examined whether premature infants, at term-equivalent age, exhibit brain functional organization similar to that of at-term neonates, both at the global and local levels. By using data from the Developing Human Connectome Project (Hughes et al., 2017), including longitudinal follow ups for some premature, we delved into the developmental trajectories of these network properties in both premature and at-term neonates.

Methods

Participants

We analyzed 597 preprocessed structural-functional datasets from the developing Human Connectome Project (Hughes et al., 2017, third open access data),

corresponding to data acquired between 26-45 weeks postmenstrual age (PMA). We extracted two samples; the first cross-sectional sample consisted of 393 scans taken at term-equivalent age, 336 considered at-term with a mean GA of 40.07 weeks (IQR=1.86), scanned at an average PMA of 41.55 weeks (IQR=2.57) and 74 preterms with a mean GA of 32.04 weeks (IQR=5.43) scanned at an average PMA of 41.21 weeks (IQR=2.15). The second, consisted of a longitudinal sample, two scans per subject, the first at a few days after birth (average PMA = 34.10 weeks, IQR=2.43) and the second at the term-equivalent age (average PMA = 41.34 weeks (IQR=1.85)).

Image acquisition and preprocessing

The infants were recruited at St Thomas' Hospital, London, and the images were taken at the Evelina Newborn Imaging Centre, Centre for the Developing Brain, King's College London, United Kingdom. The study was approved by the UK Health Research Authority Ethics Committee (14/LO/1169) and written informed consent was obtained from the parents to collect and disclose the images, as described in Hughes et al. (2017). Images were obtained in natural sleep in the MRI room located within the neonatal intensive care unit using a Philips Achieva 3T system (using a modified R3.2.2 software), with a 32-channel receiver head coil, acoustic hood, optimized transport system, positioning devices and hearing protection specially designed for this population, allowing optimum head position control and increased comfort. Infants were fed, swaddled, and placed in a vacuum sheath; scans were supervised by a neonatal nurse or pediatrician who monitored oxygen saturation, heart rate, and body temperature throughout the study (Hughes et al., 2017).

The fMRIs were acquired using a 9x accelerated multiband (MB) echo-planar imaging sequence over 15 minutes and 3 seconds, with a repetition time (TR)= 392 ms, echo time (TE) = 38 ms, voxel size = 2.15 x 2.15 x 2.15 mm^3, flip angle = 34°, 45 slices and a total of 2300 volumes. T1-weighted images had a reconstructed voxel size = 0.8 x 0.8 x 0.8 mm^3, a field of view = 145 x 122 x 100 mm^3, TR = 4795 ms and T2-weighted images had reconstructed voxel size = 0.8 x 0.8 x 0.8 mm^3, a field of view = 145 x 145 x 108 mm, TR = 12 s, TE = 156 ms (Edwards et al., 2022). A radiologist evaluated all scans

with scores from 1 to 5, with 4 and 5 showing significant lesions in the white matter, cortex, cerebellum, or basal ganglia. The most common incidentals included punctate white matter lesions (PWML) (12%) and caudothalamic subependymal cysts (10%) (Carney et al., 2021). Only datasets with a radiologist score between 1 and 3, *i.e.* with no significant incidental findings, were included in this study.

Data processing

Individual fMRI were preprocessed with a pipeline made explicitly for the dHCP neonatal data to minimize artifacts in the BOLD signal (for details, refer to Fitzgibbon et al., 2020). Summarizing, preprocessing included field distortion, intra- and inter-volume motion corrections, removal of structured noise and cardiorespiratory artifacts, and regression of physiological noise and motion related estimates, including frame-wise displacement and the spatial standard deviation of successive difference images DVARS (Power et al., 2014). Finally, fmri datasets were registered to the Infant Brain Atlas from Neonates brain template using linear and nonlinear transformations. We further delimited the fmri datasets exclusively to gray matter voxels according to the dHCP atlas.

Network measures

A graph is a non-empty set of nodes or vertices and a set of edges or links between pairs of nodes. The brain network may be represented as a graph where the nodes are defined brain regions and the links, in this case, are the value of functional connectivity between them, defined as the correlation between their BOLD signals (Friston, 1994). For each subject, we estimated the correlation between the BOLD time-series of all possible pairs of the 90 regions within the neonate AAL atlas (Shi et al., 2011), the resulting matrix represents the brain functional network for each subject. To minimize the number of variables and keep the number of connections constant between subjects and groups, the correlation matrices were thresholded keeping a fixed proportion of the strongest connections. To avoid biases by selecting a single threshold, various proportion thresholds (also known as cost values) were tested from 0.05 to 0.50 in steps

of 0.05 (this is, keeping 5 to 50 % of the strongest connections in steps of 5 %). From these thresholded matrices, we estimated network properties as described below.

Network properties can be studied at the local (nodes) or global (the whole network) level. To compare the properties of the brain networks of preterm and at-term infants, we used four properties: 1) node strength (S), 2) clustering coefficient (CC), 3) length of the shortest path (d) and 4) global efficiency (E_{global}) using the Brain Connectivity Toolbox (Rubinov & Sporns, 2010). Detailed definitions are provided below, according to Rubinov & Sporns (2010).

Through these parameters, we can study two fundamental properties to understand the brain functional organization: segregation, which refers to specialized densely interconnected groups of brain regions, and integration, which estimates the potential to share information globally. Higher CC indicates a segregated network and shorter l and higher GE indicates high integration.

The node strength (S), also known as weighted degree or connectivity strength, is defined as the sum of the weights w of all edges connected to a node i:

$$S_i = \sum_j w_{ij} \; ,$$

(1)

The clustering coefficient (CC) measures the average probability that two neighbors of node i are also connected, i.e. the proportion of triangles around a single node at the network level, is defined as the average value of C_i over all nodes (n) in the network where N is the total number of nodes (Watts y Strogatz, 1998):

$$CC = \frac{1}{n}\sum_{i \in N} C_i = \frac{1}{n}\sum_{i \in N} \frac{2t_i}{D_i(D_i - 1)},$$

(2)

where the degree of connectivity of a node i is denoted as Di and t_i is the number of triangles in the graph that includes all the nodes that node i forms with all the other nodes in N.

The average length of the shortest path (d) is also called the characteristic path length, d_{ij} is defined as the length of the shortest path going from nodes i to j (Watts & Strogatz, 1998):

$$d_{ij}^{w} = \sum_{auv \in g_{i \leftrightarrow j}^{w}} f(w_{uv}),$$

(3)

where f is a mapping from weight to length and $g_{i \leftrightarrow j}^{w}$ is the shortest weighted path between nodes i and j.

The global efficiency (E_{global}) is the average of the inverse of the shortest path length from a node to all other nodes i other than j, measures the efficiency of parallel communication when all nodes exchange information with each other at the same time (Liu):

$$E_{global} = \frac{1}{n} \sum_{i \in N} \frac{\sum_{j \in N, j \neq i} (d_{ij}^{w})^{-1}}{n-1},$$

(4)

where d_{ij}^{w} is the shortest path length between node i and node j.

Division of groups

Figure 1 shows the total sample of neonates that we grouped based on the classification proposed by the WHO. Specifically, preterm infants are classified according to PMA as follows:

- Extremely preterm: < 28 WG
- Very preterm: 28 to 32 WG
- Moderate to late preterm: 32 to 37 WG

In at-term infants, variability has been demonstrated in a variety of domains depending on the gestational week of delivery within a 5-week range - e.g.respiratory morbidity - and therefore, the American College of Obstetricians and Gynecologists and the Society for Maternal-Fetal Medicine (2012) proposed a more detailed classification that we also adopted for grouping the at-term neonates. Specifically, the following grouping was implemented:

- Early term: 37 to 38 WG
- Full term: 39 to 40 WG
- Late term: 41 WG
- Postterm: > 42 WG

Figure 1. PMA of the sample. PMA at birth and at scan for the sample here explored (lines represent longitudinal data for the same participant).

To characterize the developmental trajectories of the graph theory properties here explored, linear, quadratic, and log-linear mixed models were constructed with gestational age at scan as an independent fixed-effect variable. Random effects were added for the intercept and subject ID. Significance was defined as p < 0.05, and the model with the lowest Akaike Information Criterion (AIC) was selected as the best model to describe the data.

Results

First, we observed the connectivity values for each group as histograms. Figure 2 shows the distribution of the correlation values by group, with the average values between 0 and 0.5 for all groups. However, the average values are lower for the premature groups, increasing with PMA at birth.

Figure 2. Frequency of the correlation values per group. The average correlation values increase with the PMA at birth (vertical lines indicate the average for each group).

When contrasting the network properties between groups, we observed that all cost values show significant differences between groups for all graph theory measures (ANCOVA tests; P<0.05, FDR corrected, Figure 3).

Figure 3. Graph theory properties as a function of the connection density. A) Clustering coefficient. B) Node strength. C) Characteristic path length. D) Global efficiency. Significant differences are shown for all costs (percent of total connections) between all groups (P<0.05, FDR corrected). On the x-axis, the cost values are shown from 0.05 to 0.50 in steps of 0.05. On the y-axis, the values of the graph theory measures are shown.

For simplicity, we further detail differences between groups considering only the cost value of 0.10 (strongest 10% of all the connections).

1. Cross-sectional dataset, differences at term-equivalent age

To investigate whether preterm infants scanned at term-equivalent age exhibited the same global network properties as infants born at-term, a sample of 393 infants acquired between 37-44 weeks of postmenstrual age (PMA) and without radiological signs of white matter lesions was explored (Figure 4).

Figure 4. PMA at birth and at scan in cross-sectional data.

Using a one-way ANCOVA, controlling for PMA at the time of scan and sex, we compared graph theory measures among seven groups of infants as a function of their PMA at birth. A significant effect between groups was identified (Table 1, Figure 5), with post hoc tests (Tukey's HSD) showing significant differences between preterm and term infants.

As expected, the values for the clustering coefficient CC, connectivity strength S and global efficiency E_{global} increase as PMA at birth increases, controlling for age at scanning and sex. Additionally, the characteristic path length d decreases as PMA at birth increases. Figures 5A and 5B for CC and S show significant differences between preterm and term infants, while in Figures 5C and 5D d and E_{global} show significant differences between preterm infants and at least one group of at-term infants.

Table 1.

One-way ANCOVA test for graph theory properties

	Df	F value	Pr(>F)
Clustering coefficient	6	16.14	$<2\times10^{-16}$ ***
Node strength	6	14.029	1.89×10^{-14}***
Shortest path length	6	6. 530	1.41×10^{-06}***
Global efficiency	6	7.751	6.96×10^{-08}***

The table shows the results of the 1-way ANCOVA test that compares graph theory measures between seven groups of infants based on their PMA at birth, controlling for PMA at MRI session and sex. Significant between group effect values for the four graph theory measures are shown.(*** $p < 0.001$).

Figure 5. Graph theory measures. Significant differences between preterm and term infants are shown. (Lines at the top represent comparisons with p < 0.05, Tukey HSD post hoc tests). A) Mean clustering coefficient. B) Mean node strength. C) Shortest path length. D) Global efficiency. Clustering coefficient, node strength and global efficiency increase as PMA at birth increases, controlling for age at scanning and sex, while shortest path length decreases as PMA at birth increases (* p < 0.05, ** p < 0.01, *** p < 0.001).

2. Local effects (brain regions)

In general, the results show that even at term-equivalent age, the higher the PMA at birth is associated with better global properties of the network. Then, the question if these findings are true for all the brain regions arises. Figures 6 to 8 show the graph

theory parameters estimated for each brain region for the seven groups and the corresponding F value obtained by performing a one-way ANCOVA between groups, controlling for PMA at the time of scan and sex.

Figure 6 shows that the largest effects CC are located in the precuneus, precentral, postcentral and paracentral gyri. But a large portion of the brain show significant differences between groups, including primary sensory regions.

Figure 6. Average clustering coefficient per group. The first six rows show the average CC value per group; the higher the PMA, the higher the CC value in the central, parietal and sensorimotor areas. The seventh row shows the *F values* of the ANCOVA test

between groups by brain regions; the higher the value, the more significant the difference between the groups. The red line indicates the correction with FDR at q < 0.05 (p < 0.026, f value = 2.477).

For the characteristic path length the most significant effects are also observed in the precuneus, the precentral and middle frontal gyri, with other regions of primary sensory regions also showing significant differences between groups (Figure 7).

Figure 7. Average node strength per group. The first six rows show the mean S value per group; the higher the PMA, the higher the S value in the central areas. The seventh row shows the *F values* of the ANCOVA test between groups; the higher the value, the

more significant the difference between the regions. The red line indicates the correction with FDR at q < 0.05 (p < 0.021, f value = 2.568).

The efficiency of a node was estimated on the subgraph created by its neighbors, this is defined as the local efficiency (E_{local}). Again the brain regions with the largest differences in the E_{local} are the precuneus, the precentral, postcentral and paracentral gyri. But significant differences are identified in several brain regions, particularly including primary sensory regions (Figure 8).

Average local efficiency per group

Figure 8. Average local efficiency per group. The first six rows show the average local efficiency E_{local}, per group, the higher the PMA, the higher the E_{local} value in the central and parietal areas. The seventh row shows the *F values* of the ANCOVA test between groups; the higher the value, the more significant the difference between groups. The red line indicates the correction with FDR at q < 0.05 (p < 0.03, f value = 2.534).

3. Developmental trajectories

When exploring the developmental trajectories of such functional organization properties. The best fitting models showed non-linear trajectories for all the properties in preterm neonates and two of them in at-term neonates. Specifically, in at-term neonates, clustering coefficient showed quadratic fit and node strength showed a logarithmic trajectory, while the shortest path length and global efficiency showed linear trajectories (Figure 9). However, it is worth noting that all models for at-term neonates showed very similar fitting, with just marginal differences in their AIC (Table 2).

Figure 9. Best fitting models in at-term infants. A) Clustering coefficient. B) Node strength. C) Shortest path length. D) Global efficiency. Clustering coefficient, node strength and global efficiency increase as gestational age scan increases.

Table 2.

Model fitting for graph theory measures in at-term neonates.

	Clustering coefficient			Node strength			Shortest path length			Global efficiency		
	F value	Pr(>F)	AIC	F value	Pr(>F)	AIC	F value	Pr(>F)	AIC	F value	Pr(>F)	AIC
Lineal model	7.94	5.14×10^{-03}	-1117.0615	0.0002	9.99×10^{-01}	887.5415	0.76	3.84×10^{-01}	**902.8121**	7.02	8.42×10^{-03}	**-825.5566**
Quadratic model	8.01	4.92×10^{-03}	**-1119.3961**	0.0002	9.99×10^{-01}	887.8418	0.76	3.84×10^{-01}	904.0926	7.00	8.52×10^{-03}	-823.5578
Logarithmic model	8.20	4.45×10^{-03}	-1117.3221	0.002	9.62×10^{-01}	**887.5338**	0.72	3.97×10^{-01}	902.8408	6.99	8.58×10^{-03}	-825.5487

The table compares the linear, quadratic, and logarithmic models for each graph theory measure in at-term neonates. The lower the AIC value indicates the best fitting model, highlighted in bold for each case.

For the premature neonates, the best fitting models showed non-linear trajectories for all the explored properties. Specifically, the clustering coefficient, node strength and global efficiency showed logarithmic trajectories, while the characteristic path length

showed the quadratic as the best fitting model (Figure 9; Table 3).

Figure 10. Best fitting models for the network properties in preterm infants. A) Clustering coefficient. B) Node strength. C) Shortest path length. D) Global efficiency. Clustering coefficient, node strength and global efficiency increase as gestational age scan increases.

Table 3.

Model fitting for graph theory measures in premature neonates.

	Clustering coefficient			Node strength			Shortest path length			Global efficiency		
	F value	Pr(>F)	AIC	F value	Pr(>F)	AIC	F value	Pr(>F)	AIC	F value	Pr(>F)	AIC
Lineal model	57.99	6.53×10^{-11}	-287.4677	47.01	1.93×10^{-09}	433.3717	36.30	8.99×10^{-08}	752.5673	26.59	2.09×10^{-06}	-143.1496
Quadratic model	11.66	9.05×10^{-04}	-279.2732	14.09	2.87×10^{-04}	433.2890	24.00	3.84×10^{-06}	**740.8478**	15.79	1.31×10^{-04}	-140.2883
Logarithmic model	61.09	2.08×10^{-11}	**-296.8879**	49.99	6.43×10^{-10}	**423.9279**	40.15	2.27×10^{-08}	742.4612	28.92	8.18×10^{-07}	**-152.2713**

The table compares the linear, quadratic, and logarithmic fit models for each graph theory measure in preterm. The lower the AIC value indicates the best fitting model; it is highlighted in bold.

4. Longitudinal dataset

Subsequently, we wanted to investigate how global network characteristics change with age in the subjects with more than one acquisition. For this, we used the longitudinal subsample of 37 infants with two MRI scans each and without radiological signs of white matter lesions (Figure 11).

Figure 11. PMA at birth and at scan for the longitudinal data. Lines represent longitudinal data for the same participant.

The following results are also shown with a connection density of 10%. Using a Wilcoxon test for repeated measures, we compared the values of the graph theory measures between the first and second scan in the extrem preterm, very preterm and late preterm groups.

There is a tendency to improve values in all measures of graph theory in the second scan (Figure 12); however, there is noticeable variability among preterm infants, with some infants showing no changes with time or even the opposite change to that of the group. The number of extremely preterm infants is very small and this may limit the finding of significant differences due to small statistical power, however it shows a tendency to have improved values in the second imaging session (at term-equivalent age) for all the measures. On the other hand, the sample of late preterm infants only shows significant differences for clustering coefficient, potentially due to the few days between scanning sessions. Figure 12 shows the results for the four measurements, the figure 12A is for CC, no significant differences are observed in extremely preterms V = 1, p-value = 0.25, while in very preterm V = 19, p-value = 0.01807 and late preterm infants V = 28, p-value = 0.01041 there are significant differences. As expected, CC increases with increasing PMA, the network seems to be more segregated in the second resonance at term-equivalent age.

Figure 12. Graph theory measures longitudinal data. A) Longitudinal changes clustering coefficient. B) Longitudinal changes in node strength. C) Longitudinal changes shortest path length. D) Longitudinal changes in global efficiency. Lines and the star at the top represent comparisons with p < 0.05.

Similarly, the connectivity strength also shows significant differences only for very preterm infants (extremely preterm V = 0, p-value = 0.125; very preterm V = 19, p-value = 0.01807; late preterm V = 58, p-value = 0.2462; Figure 12B). The same is true for the characteristic path length d (extremely preterm V = 10, p-value = 0.125; very preterm V = 97, p-value = 0.03534; late preterm V = 70, p-value = 0.5226; Figure 12C).

Finally, for the global efficiency E_{global}, no significant differences were observed for any group (extremely preterm V = 0, p-value = 0.125; very preterm V = 32, p-value = 0.1205; late preterm V = 91, p-value = 0.8317; Figure 12D).

Discussion

In this study, we performed a detailed analysis of the functional organization of the brain and its developmental trajectory in preterm and at-term neonates.

We performed a detailed exploration on the effects of PMA at birth, first dividing the sample in a total of seven groups: three preterm groups categorized by the WHO criteria and four at-term groups categorized according to the guidelines of the American College of Obstetricians and Gynecologists and the Society for Maternal-Fetal Medicine. The results show significant and robust differences for the premature neonates even at term-equivalent age, with an evident gradient of effects depending on the PMA at birth. The results are reproduced for a large variety of connectivity thresholds, here density thresholds ranging from the top 5% higher connectivity values to the 50% with 5% intervals, evidencing that these results are very robust independently of the connectivity threshold.

Based on the classification of the American College of Obstetricians and Gynecologists and the Society for Maternal-Fetal Medicine, there has been a debate whether the early term group presents sufficient fetal maturity to avoid the neurological comorbidities of prematurity. Recent research on this specific group of early term subjects has revealed risk factors for physiological comorbidities, such as respiratory deficiency and diabetes that seem to persist into adulthood (Chen et al., 2022; Odibo et al., 2016). However, our results on the brain functional organization suggest that this group is more similar to the other at-term groups rather than to the premature groups. Specifically, the early terms showed significant differences with at least one of the preterm groups in all the properties here explored, but showed no significant difference with any of the other three at-term groups.

When exploring such properties at the local level (brain regions), the precuneus, mid cingulate, primary motor and somatosensory cortices, parietal gyrus, orbitofrontal cortex, superior and inferior frontal gyrus, temporal gyrus, occipital gyrus, supramarginal gyrus, Heschl gyrus left, Rolandic operculum, lingual gyrus, cingulate gyrus and insula showed the greatest differences between groups. It is not surprising that these regions

include the primary somatosensory and primary motor regions, as premature birth is usually associated with sensorimotor, auditory, visual, cognitive and language deficits, antisocial personality, ADHD, ASD and CP (Do et al., 2019; Hee Chung et al., 2020). As such, most of the prospective interventions in preterm infants include sensorimotor enrichment (Harmony et al., 2016), with overall benefits in behavioral and cognitive domains. In general, the results show that the regions that will be part of the primary systems are the ones that show the most significant differences between the preterm and term groups. This evidence is consistent with the higher prevalence rates of visual and auditory disorders/deficits in premature. In particular, the auditory cortex showed the greatest effect in the left hemisphere, which is largely recognized as the commonly dominant cortex for language acquisition (Olulade et al., 2020). This result provides the potential neurofunctional substrate for the language acquisition deficits identified in preterm infants (Gozzo et al., 2009; Myers et al., 2010). In addition, the largest differences between groups were identified for the precuneus (Figures 6-8), which is one of the main hubs of the brain network and part of the DMN. Alterations in the connectivity of this network have been associated with ADHD (Castellanos & Aoki, 2016; Elton et al., 2014; Gracia-Tabuenca et al., 2020), visuospatial abilities (Fernandez-Baizan et al., 2021) episodic memory (Stedall et al., 2023), affective disorders (Guilherme Monte Cassiano et al., 2016) and social cognition (Pereira et al., 2017). Our results suggest a link between these brain functional alterations and the vulnerability to develop such behavioral disorders.

To characterize the developmental trajectories of the graph theory properties here explored, linear, quadratic, and logarithmic models were fitted using gestational age at the scanning session as a fixed effect and sex as confounding variable. The best fitting models for preterms were logarithmic, except for the characteristic path length that followed a quadratic trajectory (Table 3). In at-term neonates, clustering coefficient showed quadratic fit and node strength showed a logarithmic trajectory, while the characteristic path length and global efficiency showed linear trajectories. However, we must note that for the at-term neonates the AIC was very similar for all the tested models for each property (Table 2). These results are relevant as we identified no previous description of such trajectories for these properties, with only one recent study

suggesting non-linear trajectories for the development of resting state networks in the same sample (Kim et al., 2023).

Finally, using the longitudinal subsample of preterm infants, we identified a significant increase with age from ages lower than 37 WG to at term-equivalent ages, mainly for the very preterm group in all properties, except for the global efficiency (Figure 12). Global efficiency is considered a measure of network integration, which in the brain is understood to be supported by the establishment of long-range connections (REFs). The emergence of these connections has been associated with the development of the white matter microstructural properties, including myelination (REFs). However, premature birth is known to alter the production of premyelinating oligodendrocytes during gestation, which consequently alters myelin production in later stages of development (Volpe, 2009). This interruption in myelin maturation may be associated with the lack of significant changes in the global efficiency of the network (Figure 12). In contrast, extremely preterm infants showed no significant changes potentially due to the small sample size for this group. Meanwhile, the late preterm infants showed significant changes only for the clustering coefficient, probably due to the shortest period between both scans, preventing the observation of more remarkable changes. In general, it can be observed a high degree of variability in the developmental changes in this longitudinal subsample, with some of the subjects showing no change or even opposing effects to the group average change. This variability may be associated with the varying prevalence rates for a diversity of developmental outcomes. However, results from cognitive-behavioral assessments and demographic data were not available to explore these associations.

Overall, our results confirm that the functional connectivity, the integration and segregation properties of the premature brain are significantly decreased at-term equivalent age, despite showing rapid maturation after birth.

Conclusions

Preterm birth is associated with significant alterations of the functional organization of the brain network, even in the absence of evident brain injuries. Dividing the sample into

finer groups allowed us to observe with greater detail how the smaller the PMA, the greater the alterations in the functional network even at term-equivalent ages. However, the premature infants do show dramatic improvement in the brain network from birth to at term-equivalent age, but with great variability among subjects.

Acknowledgments

Nelsiyamid López-Guerrero is a doctoral student from the Programa de Doctorado en Ciencias Biomédicas, at the Universidad Nacional Autónoma de México (UNAM) and received the fellowship 823584 from CONAHCYT, Mexico. This work was supported by the grant UNAM-PAPIIT IN208622 to Sarael Alcauter. We thank Leopoldo González Santos and the Unidad de Cómputo del Instituto de Neurobiología, UNAM, for their technical assistance. Data was provided by the developing Human Connectome Project, KCL-Imperial-Oxford Consortium funded by the European Research Council under the European Union Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement no. [319456]. We are grateful to the families who generously supported this trial and the dHCP for sharing the datasets.

References

Blencowe, H., Cousens, S., Oestergaard, M. Z., Chou, D., Moller, A.-B., Narwal, R., Adler, A., Vera Garcia, C., Rohde, S., Say, L., & Lawn, J. E. (2012). National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: A systematic analysis and implications. *Lancet (London, England)*, *379*(9832), 2162–2172. https://doi.org/10.1016/S0140-6736(12)60820-4

Börner, K., Sanyal, S., & Vespignani, A. (2007). Network Science. *Annual Review of Information Science & Technology*, *41*, 537–607.

Bouyssi-Kobar, M., De Asis-Cruz, J., Murnick, J., Chang, T., & Limperopoulos, C. (2019). Altered Functional Brain Network Integration, Segregation, and Modularity in Infants Born Very Preterm at Term-Equivalent Age. *The Journal of Pediatrics*, *213*, 13-21.e1.

https://doi.org/10.1016/j.jpeds.2019.06.030

Carney, O., Hughes, E., Tusor, N., Dimitrova, R., Arulkumaran, S., Baruteau, K. P., Collado, A. E., Cordero-Grande, L., Chew, A., Falconer, S., Allsop, J. M., Rueckert, D., Hajnal, J., Edwards, A. D., & Rutherford, M. (2021). Incidental findings on brain MR imaging of asymptomatic term neonates in the Developing Human Connectome Project. *eClinicalMedicine*, *38*, 100984. https://doi.org/10.1016/j.eclinm.2021.100984

Castellanos, F. X., & Aoki, Y. (2016). Intrinsic Functional Connectivity in Attention-Deficit/Hyperactivity Disorder: A Science in Development. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *1*(3), 253–261. https://doi.org/10.1016/j.bpsc.2016.03.004

Chen, Z., Xiong, C., Liu, H., Duan, J., Kang, C., Yao, C., Chen, K., Chen, Y., Liu, Y., Liu, M., & Zhou, A. (2022). Impact of early term and late preterm birth on infants' neurodevelopment: Evidence from a cohort study in Wuhan, China. *BMC Pediatrics*, *22*(1), 251. https://doi.org/10.1186/s12887-022-03312-3

Do, C. H. T., Kruse, A. Y., Wills, B., Sabanathan, S., Clapham, H., Pedersen, F. K., Pham, T. N., Vu, P. M., & Børresen, M. L. (2019). Neurodevelopment at 2 years corrected age among Vietnamese preterm infants. *Archives of Disease in Childhood*, archdischild-2019-316967. https://doi.org/10.1136/archdischild-2019-316967

Doria, V., Beckmann, C. F., Arichi, T., Merchant, N., Groppo, M., Turkheimer, F. E., Counsell, S. J., Murgasova, M., Aljabar, P., Nunes, R. G., Larkman, D. J., Rees, G., & Edwards, A. D. (2010). Emergence of resting state networks in the preterm human brain. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(46), 20015–20020. https://doi.org/10.1073/pnas.1007921107

Edwards, A. D., Rueckert, D., Smith, S. M., Abo Seada, S., Alansary, A., Almalbis, J., Allsop, J., Andersson, J., Arichi, T., Arulkumaran, S., Bastiani, M., Batalle, D., Baxter, L., Bozek, J., Braithwaite, E., Brandon, J., Carney, O., Chew, A., Christiaens, D., … Hajnal, J. V.

(2022). The Developing Human Connectome Project Neonatal Data Release. *Frontiers in Neuroscience*, *16*. https://www.frontiersin.org/articles/10.3389/fnins.2022.886772

Elton, A., Alcauter, S., & Gao, W. (2014). Network connectivity abnormality profile supports a categorical‑dimensional hybrid model of ADHD. *Human Brain Mapping*, *35*(9), 4531–4543. https://doi.org/10.1002/hbm.22492

Eyre, M., Fitzgibbon, S. P., Ciarrusta, J., Cordero-Grande, L., Price, A. N., Poppe, T., Schuh, A., Hughes, E., O'Keeffe, C., Brandon, J., Cromb, D., Vecchiato, K., Andersson, J., Duff, E. P., Counsell, S. J., Smith, S. M., Rueckert, D., Hajnal, J. V., Arichi, T., … Edwards, A. D. (2021). The Developing Human Connectome Project: Typical and disrupted perinatal functional connectivity. *Brain*, *144*(7), 2199–2213. https://doi.org/10.1093/brain/awab118

Fernandez-Baizan, C., Caunedo-Jimenez, M., Martinez, J. A., Arias, J. L., Mendez, M., & Solis, G. (2021). Development of visuospatial memory in preterm infants: A new paradigm to assess short-term and working memory. *Child Neuropsychology*, *27*(3), 296–316. https://doi.org/10.1080/09297049.2020.1847264

Fitzgibbon, S. P., Harrison, S. J., Jenkinson, M., Baxter, L., Robinson, E. C., Bastiani, M., Bozek, J., Karolis, V., Cordero Grande, L., Price, A. N., Hughes, E., Makropoulos, A., Passerat-Palmbach, J., Schuh, A., Gao, J., Farahibozorg, S.-R., O'Muircheartaigh, J., Ciarrusta, J., O'Keeffe, C., … Andersson, J. (2020). The developing Human Connectome Project (dHCP) automated resting-state functional processing framework for newborn infants. *NeuroImage*, *223*, 117303. https://doi.org/10.1016/j.neuroimage.2020.117303

Gao, W., Gilmore, J. H., Giovanello, K. S., Smith, J. K., Shen, D., Zhu, H., & Lin, W. (2011). Temporal and Spatial Evolution of Brain Network Topology during the First Two Years of Life. *PLoS ONE*, *6*(9), e25278. https://doi.org/10.1371/journal.pone.0025278

Geneva: World Health, & Organization. (2023). *Born too soon: Decade of action on preterm birth.* https://www.who.int/publications/i/item/9789240073890

Gozzo, Y., Vohr, B., Lacadie, C., Hampson, M., Katz, K. H., Maller-Kesselman, J., Schneider, K. C., Peterson, B. S., Rajeevan, N., Makuch, R. W., Constable, R. T., & Ment, L. R. (2009). Alterations in neural connectivity in preterm children at school age. *NeuroImage*, *48*(2), 458–463. https://doi.org/10.1016/j.neuroimage.2009.06.046

Gracia-Tabuenca, Z., Díaz-Patiño, J. C., Arelio, I., & Alcauter, S. (2020). Topological Data Analysis Reveals Robust Alterations in the Whole-Brain and Frontal Lobe Functional Connectomes in Attention-Deficit/Hyperactivity Disorder. *Eneuro*, *7*(3), ENEURO.0543-19.2020. https://doi.org/10.1523/ENEURO.0543-19.2020

Guilherme Monte Cassiano, R., Gaspardo, C. M., Cordaro Bucker Furini, G., Martinez, F. E., & Martins Linhares, M. B. (2016). Impact of neonatal risk and temperament on behavioral problems in toddlers born preterm. *Early Human Development*, *103*, 175–181. https://doi.org/10.1016/j.earlhumdev.2016.09.015

Harmony, T., Barrera-Reséndiz, J., Juárez-Colín, M. E., Carrillo-Prado, C., Del Consuelo Pedraza-Aguilar, M., Asprón Ramírez, A., Hinojosa-Rodríguez, M., Fernández, T., & Ricardo-Garcell, J. (2016). Longitudinal study of children with perinatal brain damage in whom early neurohabilitation was applied: Preliminary report. *Neuroscience Letters*, *611*, 59–67. https://doi.org/10.1016/j.neulet.2015.11.013

Hee Chung, E., Chou, J., & Brown, K. A. (2020). Neurodevelopmental outcomes of preterm infants: A recent literature review. *Translational Pediatrics*, *9*(Suppl 1), S3–S8. https://doi.org/10.21037/tp.2019.09.10

Hughes, E. J., Winchman, T., Padormo, F., Teixeira, R., Wurie, J., Sharma, M., Fox, M., Hutter, J., Cordero-Grande, L., Price, A. N., Allsop, J., Bueno-Conde, J., Tusor, N., Arichi, T., Edwards, A. D., Rutherford, M. A., Counsell, S. J., & Hajnal, J. V. (2017). A dedicated neonatal brain imaging system. *Magnetic Resonance in Medicine*, *78*(2), 794–804. https://doi.org/10.1002/mrm.26462

Kim, J.-H., De Asis-Cruz, J., Krishnamurthy, D., & Limperopoulos, C. (2023). Toward a more

informative representation of the fetal–neonatal brain connectome using variational autoencoder. *eLife*, *12*, e80878. https://doi.org/10.7554/eLife.80878

Larsen, M. L., Wiingreen, R., Jensen, A., Rackauskaite, G., Laursen, B., Hansen, B. M., Hoei-Hansen, C. E., & Greisen, G. (2022). The effect of gestational age on major neurodevelopmental disorders in preterm infants. *Pediatric Research*, *91*(7), 1906–1912. https://doi.org/10.1038/s41390-021-01710-4

Myers, E. H., Hampson, M., Vohr, B., Lacadie, C., Frost, S. J., Pugh, K. R., Katz, K. H., Schneider, K. C., Makuch, R. W., Constable, R. T., & Ment, L. R. (2010). Functional connectivity to a right hemisphere language center in prematurely born adolescents. *NeuroImage*, *51*(4), 1445–1452. https://doi.org/10.1016/j.neuroimage.2010.03.049

Odibo, I. N., Bird, T. M., McKelvey, S. S., Sandlin, A., Lowery, C., & Magann, E. F. (2016). Childhood Respiratory Morbidity after Late Preterm and Early Term Delivery: A Study of Medicaid Patients in S outh C arolina. *Paediatric and Perinatal Epidemiology*, *30*(1), 67–75. https://doi.org/10.1111/ppe.12250

Ohuma, E. O., Moller, A.-B., Bradley, E., Chakwera, S., Hussain-Alkhateeb, L., Lewin, A., Okwaraji, Y. B., Mahanani, W. R., Johansson, E. W., Lavin, T., Fernandez, D. E., Domínguez, G. G., De Costa, A., Cresswell, J. A., Krasevec, J., Lawn, J. E., Blencowe, H., Requejo, J., & Moran, A. C. (2023). National, regional, and global estimates of preterm birth in 2020, with trends from 2010: A systematic analysis. *The Lancet*, *402*(10409), 1261–1271. https://doi.org/10.1016/S0140-6736(23)00878-4

Olulade, O. A., Seydell-Greenwald, A., Chambers, C. E., Turkeltaub, P. E., Dromerick, A. W., Berl, M. M., Gaillard, W. D., & Newport, E. L. (2020). The neural basis of language development: Changes in lateralization over age. *Proceedings of the National Academy of Sciences*, *117*(38), 23477–23483. https://doi.org/10.1073/pnas.1905590117

Pereira, S. A., Pereira Junior, A., Costa, M. F. D., Monteiro, M. D. V., Almeida, V. A. D., Fonseca Filho, G. G. D., Arrais, N., & Simion, F. (2017). A comparison between preterm and

full‑term infants' preference for faces. *Jornal de Pediatria (Versão em Português)*, *93*(1), 35–39. https://doi.org/10.1016/j.jpedp.2016.07.011

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*, *84*, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Preterm birth. (2022). https://www.who.int/en/news-room/fact-sheets/detail/preterm-birth

Ream, M. A., & Lehwald, L. (2018). Neurologic Consequences of Preterm Birth. *Current Neurology and Neuroscience Reports*, *18*(8), 48. https://doi.org/10.1007/s11910-018-0862-2

Rogers, C. E., Lean, R. E., Wheelock, M. D., & Smyser, C. D. (2018). Aberrant structural and functional connectivity and neurodevelopmental impairment in preterm children. *Journal of Neurodevelopmental Disorders*, *10*(1), 38. https://doi.org/10.1186/s11689-018-9253-x

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: Uses and interpretations. *NeuroImage*, *52*(3), 1059–1069. https://doi.org/10.1016/j.neuroimage.2009.10.003

Scheinost, D., Kwon, S. H., Shen, X., Lacadie, C., Schneider, K. C., Dai, F., Ment, L. R., & Constable, R. T. (2016). Preterm birth alters neonatal, functional rich club organization. *Brain Structure and Function*, *221*(6), 3211–3222. https://doi.org/10.1007/s00429-015-1096-6

Shi, F., Yap, P.-T., Wu, G., Jia, H., Gilmore, J. H., Lin, W., & Shen, D. (2011). Infant Brain Atlases from Neonates to 1- and 2-Year-Olds. *PLoS ONE*, *6*(4), e18746. https://doi.org/10.1371/journal.pone.0018746

Smyser, C. D., & Neil, J. J. (2015). Use of resting state functional MRI to study brain development and injury in neonates. *Seminars in Perinatology*, *39*(2), 130–140. https://doi.org/10.1053/j.semperi.2015.01.006

Smyser, C. D., Snyder, A. Z., Shimony, J. S., Mitra, A., Inder, T. E., & Neil, J. J. (2016).

Resting-State Network Complexity and Magnitude Are Reduced in Prematurely Born Infants. *Cerebral Cortex (New York, N.Y.: 1991)*, *26*(1), 322–333. https://doi.org/10.1093/cercor/bhu251

Stedall, P. M., Spencer-Smith, M. M., Lah, S., Doyle, L. W., Spittle, A. J., Burnett, A. C., & Anderson, P. J. (2023). Episodic and prospective memory difficulties in 13-year-old children born very preterm. *Journal of the International Neuropsychological Society*, *29*(3), 257–265. https://doi.org/10.1017/S1355617722000170

Thomason, M. E., Brown, J. A., Dassanayake, M. T., Shastri, R., Marusak, H. A., Hernandez-Andrade, E., Yeo, L., Mody, S., Berman, S., Hassan, S. S., & Romero, R. (2014). Intrinsic Functional Brain Architecture Derived from Graph Theoretical Analysis in the Human Fetus. *PLoS ONE*, *9*(5), e94423. https://doi.org/10.1371/journal.pone.0094423

Thomason, M. E., Grove, L. E., Lozon, T. A., Vila, A. M., Ye, Y., Nye, M. J., Manning, J. H., Pappas, A., Hernandez-Andrade, E., Yeo, L., Mody, S., Berman, S., Hassan, S. S., & Romero, R. (2015). Age-related increases in long-range connectivity in fetal functional neural connectivity networks in utero. *Developmental Cognitive Neuroscience*, *11*, 96–104. https://doi.org/10.1016/j.dcn.2014.09.001

Volpe, J. J. (2009). Brain injury in premature infants: A complex amalgam of destructive and developmental disturbances. *The Lancet Neurology*, *8*(1), 110–124. https://doi.org/10.1016/S1474-4422(08)70294-1

Watts, D. J., & Strogatz, S. H. (1998). *Collective dynamics of 'small-world' networks. 393*.

World Health Organization. (2012). *Born too soon: The global action report on preterm birth*. 112.