1 **Energy Landscape near Glioma Decipher the Role of Glioma Stem Cell within Glioma**
2 **Hierarchy**
3

4 Mengchao Yao[1], Yang Su[1], Ruiqi Xiong[1], Xile Zhang[1], Xiao-Mei Zhu[2], Yong-Cong Chen[1], Ping
5 Ao[3*]
6

[1] Shanghai Center for Quantitative Life Sciences and Physics Department, Shanghai University,
Shanghai, China;
[2] Shanghai Key Laboratory of Modern Optical System, School of Optical-Electrical and
Computer Engineering, University of Shanghai for Science and Technology, Shanghai, China;
[3] College of Biomedical Engineering, Sichuan University, Chengdu, Sichuan Province, China.
[*] Correspondence: aoping@sjtu.edu.cn

Abstract

Glioma stem cells (GSCs) have been recognized as key players in glioma recurrence and
therapeutic resistance, presenting a promising target for novel treatments. However, limited
understanding of the role GSCs play in the hierarchy of glioma has drawn controversy and
hindered research translation into therapies. Despite significant advances in our
understanding of gene regulatory networks, the dynamics of these networks and their
implications for glioma remain elusive. This study employs a systemic theoretical perspective
to integrate empirical knowledge into a core endogenous network model for glioma, thereby
elucidating its energy landscape through network dynamics computation. The model
identifies two stable states corresponding to astrocytoma and oligodendroglioma, connected
by a transitional state associated with GSCs, indicating the instability of GSCs *in vivo* and
providing negative proof on their role as the cellular origin of glioma. We also obtained
various stable states further indicative of glioma's multicellular origins and uncovered a group
of transition states that could potentially induce tumor heterogeneity and therapeutic
resistance. This research proposes that the transitional states linking both glioma stable states
are central to glioma heterogeneity and resistance, suggesting a combination of apoptosis-
inducing and differentiation-promoting therapies as a potential strategic. Our approach holds
promise for advancing glioma therapy by providing a new lens through which to view the
complex landscape of glioma biology.

Keywords: glioma, glioma stem cell, gene regulatory network, network dynamics, endogenous
network theory, energy landscape

Introduction

Years ago, glioma cells with the characteristics of neural/glia stem cells were identified as cancer stem cells in glioma or glioma stem cells (GSCs) (Hemmati et al. 2003; Ignatova et al. 2002; Galli et al. 2004). GSCs play crucial roles in the recurrence of glioma and the resistance to radiation and chemotherapy (Biserova et al. 2021; Lathia et al. 2015) and thus are considered promising therapeutic targets for glioma.

However, limited understanding of GSCs has drawn controversy (Rahman et al. 2011; Capp 2019) and hindered research translation into therapies (Lathia et al. 2015). Whether GSCs are the cells of glioma origin remains unclear (Biserova et al. 2021; Lathia et al. 2015), the intermediate states between somatic glia cells and glioma cells are still hiding behind haze, even the roles of GSCs and non-stem tumor cells (NSTCs) are not unified in the hierarchy of glioma (Lathia et al. 2015).

Gliomas and the normal glial system may follow the same differentiation rules (Laug, Glasgow, and Deneen 2018), understanding the principle behind the carcinogenesis of glioma and the development process of the glial system, that is, the gene regulatory network and the landscape determined by it (Ao 2008; Huang, Ernberg, and Kauffman 2009; Raju and Siggia 2023; Feinberg and Levchenko 2023), may lead to the enlightenments of the role of GSCs in glioma hierarchy, thereby helping to find insights for new therapies (Lathia et al. 2015; Uthamacumaran 2022; Prokop 2022).

Currently, data-driven studies are attempting to obtain the gene regulatory network and landscape of glioma and the glial system based on high-throughput data (Csermely et al. 2015; Niu et al. 2019; Qiu et al. 2022; Feinberg and Levchenko 2023). Notably, Hoang *et al.* (Hoang et al. 2020) have driven gene regulatory networks of retinal regeneration from data, Suvà and Tirosh (Suvà and Tirosh 2020) have obtained single cell scale distribution of different glioma samples in a time section way. Even so, these data-based studies currently still face the difficulty of insufficient quality and quantity of data (Kirk et al. 2015; Uthamacumaran 2022).

These data-driven studies are a continuation of the data explosion brought by the technological revolution witnessed in the past few decades. From the late 20th century to now, a vast amount of cancer-related knowledge was discovered by pioneers in cancer research, which was then summarized as a set of hierarchical hallmarks (Hanahan and Weinberg 2011) as a milestone. However, one of these pioneers asserted that cancer research is currently in false prosperity and urgently requires new theoretical foundations and mathematical tools (Weinberg 2014). The establishment of better theories and theory-experiment coupling, rather than solely relying on data that cannot generate true knowledge, was also called loudly in various fields of life science (MacArthur 2023; Smaldino 2019; Nurse 2021).

Such theories have long been budding. Wright (Wright 1932) and Waddington (Waddington

1957) introduced the metaphoric concept of adaptive/epigenetic landscape in evolution and development, respectively. Kauffman (Kauffman 1969) introduced Boolean dynamics to simulate gene regulatory processes, demonstrating the feasibility of using dynamics processes to model gene regulation and initiating the quantification of landscapes. Decades later, with the accumulation of necessary knowledge and data, the first real-data-based quantified landscape was obtained (X.-M. Zhu et al. 2004). After that, the Endogenous Network Theory (ENT) (Ao et al. 2008) was proposed, which argued that cancer and other complex diseases are endogenous states in the landscape emerging from the dynamics of cellular-molecular network shaped by evolution, serving as a constitutive structure at the cellular-molecular level of the evolution dynamics (Ao 2005; Buchanan 2013), which then addressed several questions in cancer and development (Yuan et al. 2017). Lord *et al.* (Lord et al. 2019) provided experimental evidence connecting the theory of gene network dynamics and experimentally observed phenotypes. Studies in early brain development (Giacomantonio and Goodhill 2010; Goodhill 2018) as well as in hematopoietic system development (Olariu and Peterson 2019; Rockne et al. 2020) have also been implemented based on a similar systemic perspective.

Herein, we adapt the systemic perspective aforementioned to construct a core endogenous network for glioma based on experimental evidence. The energy landscape near glioma depicted by network dynamics enables us to elucidate the functional role of GSCs within the glioma hierarchy, in order to provide new perspectives for developing novel glioma therapies.

Results

Construction of the endogenous network for glioma

The existence and importance of gene regulatory networks no longer needs further debate,
but how to obtain a dynamically workable and biologically rational network remains in
controversy (Kim et al. 2023; Badia-i-Mompel et al. 2023; Prokop 2022; Uthamacumaran
2022). As the first step, we chose a bottom-up way based on causality knowledge rather than
a top-down data driven way to construct the network.

The core function modules and signaling pathways of glioma have been already well
summarized (J. Chen, McKay, and Parada 2012; Sedo and Mentlein 2014). We selected the
following functional modules and signaling pathways to construct the core endogenous
network for glioma (Fig. 1A): RTK pathway (Berezowska and Schlegel 2011; Annenkov 2014;
Lemmon and Schlessinger 2010; Zeng, Cui, and Gao 2015; Newbern and Birchmeier 2010),
NF-κB pathway (Perkins 2012; Taniguchi and Karin 2018; Sun 2017; Zhang, Lenardo, and
Baltimore 2017), Ras pathway (Perkins 2012; Taniguchi and Karin 2018; Sun 2017; Zhang,
Lenardo, and Baltimore 2017), Akt pathway (Manning and Toker 2017; Vanhaesebroeck,
Stephens, and Hawkins 2012; Janku, Yap, and Meric-Bernstam 2018; Fruman and Rommel
2014), HIF pathway (Choudhry and Harris 2018; Gonzalez, Xie, and Jiang 2018; Nordgren and
Tavassoli 2011; Rankin and Giaccia 2016; Eltzschig, Bratton, and Colgan 2014), p53 pathway
(Agostini, Melino, and Bernassola 2018; Wade, Li, and Wahl 2013; Joerger and Fersht 2016;
Kastenhuber and Lowe 2017; Hafner et al. 2019), cell cycle (Banfalvi 2011; Gérard and
Goldbeter 2016; Otto and Sicinski 2017; Harashima, Dissmeyer, and Schnittger 2013; Bertoli,
Skotheim, and De Bruin 2013), cellular senescence (Wiley and Campisi 2016; Kritsilis et al.
2018; Calcinotto et al. 2019; Gorgoulis et al. 2019; Muñoz-Espín and Serrano 2014), apoptosis
(Singh, Letai, and Sarosiek 2019; Man and Kanneganti 2016; Spencer and Sorger 2011; Lavrik
2010; Hata, Engelman, and Faber 2015; Czabotar et al. 2014; Fricker et al. 2018; Zmasek and
Godzik 2013) and glia differentiation (Kyrousi, Lygerou, and Taraviras 2017; Hidalgo and
Logan 2017; Tchieu et al. 2019; K. S. Chen et al. 2017; Kamakura et al. 2004; Kang et al. 2012;
Klum et al. 2018; Kuhlbrodt et al. 1998).

The network consists of 10 function modules and pathways mentioned above, comprising 25
nodes and 75 edges, including 44 activatory edges and 31 inhibitory edges (Fig. 1A,
Supplementary Table 1). For each node, we employed an equation incorporating both
activatory and inhibitory effects to construct a 25-dimension equation set (Fig. 1B,
Supplementary Table 1).

Network dynamics results are robust under different computation frameworks

The Boolean dynamics was introduced into gene network dynamics by Kauffman in the 1960s
(Kauffman 1969). As a discrete method, Boolean dynamics can quickly capture the overall
structure. In this work, we used threshold functions equivalent to Boolean algebra to perform
calculations (Fig. 1B, Supplementary Table 1). A Boolean equation set consisting of 25 Boolean

equations was obtained for the network constructed above (Supplementary Table 1). With
10^7 random initial vectors, we obtained 62 attractors, including 16 point attractors and 46
linear attractors, resulting in a total of 116 states after expanding the linear attractors (Fig. 2A,
Supplementary Table 1). By increasing the number of random initial vectors to 10^8, we
obtained the same number and types of attractors (Supplementary Table 1). Additionally, by
exhaustively considering all initial values ($2^{25}=33554432$), we obtained the same number
and types of attractors (Supplementary Table 1). These results indicate that we have obtained
all attractors under the Boolean dynamics.

While Boolean dynamics, as a discrete method, can quickly produce preliminary results, it fails
to capture transitional states and simulate the consequence of small perturbations (Bornholdt
2008), therefore unable to determine the specific paths of transition between attractors.
Experiments in biology often involve time-slicing, capturing features at the moment of death
or detection, thus cannot guarantee that the system has reached equilibrium. Even if the
system has reached a macroscopic equilibrium, noise-driven state transitions may occur at
the microscopic level. Without transition states, it is impossible to determine how cell fate
transitions occur. Stem cells cultured *in vitro* also exhibit characteristics of unstable
equilibrium states that are prone to differentiation under small perturbations. Therefore,
calculations that do not include transition states are incomplete. To address this issue, we
need to use continuous computation frameworks.

The Langevin equation or stochastic differential equation (SDE) naturally corresponds to
biological processes (Ao 2005). However, solving a n-dimensional nonlinear SDE, in this work
n=25, would require excessive computation power, let alone there is currently no relevant
knowledge for the random terms in this work. Moreover, the two commonly used stochastic
integrals in applied mathematics and physics, Ito and Stratonovich, yield inconsistent results
on the extreme points between SDE and ordinary differential equation (ODE) (Yuan, Tang,
and Ao 2017). However, based on the mathematical tools developed previously, the extreme
points of SDEs and ODEs are equal (Yuan, Tang, and Ao 2017). Therefore, solving a high-
dimensional nonlinear SDE can be reduced to finding special solutions of high-dimensional
nonlinear ODE, which are sufficient to depict the main features of the energy landscape of
the system.

We chose the Hill equation for model construction in this work (Fig. 1B; Supplementary Table
1). Here, x represents the concentration/activity of core nodes, which is normalized to [0, 1].
The degradation rate is normalized to 1 in this model, so the degradation amount per
iteration is -x. h and k represent the Hill coefficient and the inverse of the apparent
dissociation constant, respectively. Since the equation has been already normalized, to ensure
that the S part of the sigmoid curve is as close to the center as possible (i.e., $f(x) \approx 0.5$ when
x=0.5), h and k need to satisfy the relationship $k \approx 2^h$. Based on previous experience, we
chose h=3 and k=10 initially (Fig. 1B).

We obtained a set of 25 ODEs (Supplementary Table 1). We performed numerical calculations
using uniformly distributed random initial vectors: generating a set of 25-dimensional

random vectors corresponding to the concentration/activity of each node in the network as
the starting point for each iteration. Both Newton's and Euler's method was parallelly applied
to avoid bias.
195
We performed preliminary calculations with 10^4 random initial vectors. With Newton's
method, we obtained 10 stable states, and 14 with Euler's method. To avert information loss
due to insufficient sampling, we sequentially increased the number of random initial vectors
tenfold until reached 10^8. The results converged rapidly on 10^5 random initial vectors to
20 stable states with Euler's method compared with Newton's method which the same 20
stable states were obtained on 10^7 random initial vectors. With 10^8 random initial vectors,
Newton's method and Euler's method yield the same 20 stable states under the 10^7
condition (Fig. 2B, D; Supplementary Table 1). We also identified transition states similarly, the
results converged to 97 transition states on 10^7 initial random vectors with Newton's
method and remained the same under 10^8 initial random vectors (Fig. 2C; Supplementary
Table 1). Based on these results, it can be concluded that we have identified the vast majority,
if not all, of the stable and transition states under ODE.
208
Furthermore, we explored the parameter space to ensure the parameter robustness. The
results indicated the presence of bifurcations between h=3, k=8, and h=3, k=10, but 85% of
stable states and 82% of transition states were already obtained under the h=3, k=8 parameter
set. The quantity and types of fixed points obtained were consistent among h=4, 5, $k=2^h$,
and h=3, k=10 (Fig. 2E; Supplementary Table 1). Considering that core nodes in gene
regulatory networks of vertebrates often exhibit high interconnection (Gil-Gálvez et al. 2022),
it can be speculated that significant cooperative effects exist in the core endogenous network
of vertebrates. These results indicate that the network we constructed is robust against
parameter alteration.
218
219 We obtained 116 solutions with spread linear attractors under Boolean dynamics; and 117
220 fixed points under ODE, yielding almost identical quantities. Hierarchical clustering and the
221 Euclidean distances show that every attractor under Boolean dynamics corresponds to at least
222 one fixed point in ODE (Fig. 2F; Supplementary Table 1). These results indicate that the
223 network we constructed is robust against dynamics methods.
224
225 In conclusion, the dynamics results we obtained are acceptable enough for further analysis.
226 Therefore, we will continue our work based on results obtained under the condition of h=3,
227 k=10, 10^7 random initial vectors.
228
229
230 **Potential energy landscape of glia**
231
In previous work, we discovered a universal construction method for energy functions
(Kwon, Ao, and Thouless 2005). Subsequently, energy functions in various nonlinear systems
were constructively obtained (Ma et al. 2014; Yuan et al. 2013; Y. C. Chen et al. 2020) and
the existence of energy functions in nonlinear systems was mathematically proved in two

dimension condition (Gan, Wang, and Ao 2021). Therefore, we believe that for the system
studied in this work, an energy function and related energy landscape exists. For a
hypothetical energy landscape demonstrated in Fig. 3 A, the fixed points in the equation of
motion correspond to the extremal points (red dots "peak" as unstable fixed
points/transition states and blue dots "bottom" as stable fixed points/stable states) in its
energy landscape (Fig. 3 B). The topological connections between these stable and unstable
fixed points are effectively enough to describe the primary features of its energy landscape
(Fig. 3 C) which bypass the task of obtaining a global energy function of such a high-
dimensional nonlinear system (Yuan, Tang, and Ao 2017; Yuan et al. 2017).
245
Specifically, we applied 10,000 small random perturbations to each transition state obtained
earlier to simulate the system's evolution from transition states to stable states, thereby
obtaining connections between transition states and stable states which describe the main
features of the energy landscape. We transformed the results into a topological connection
graph of fixed points, where larger nodes represent stable states, smaller nodes represent
transition states, the shape of the nodes represents the level of the apoptosis module, the
color of the nodes represents the differentiation type, and the color of the node labels
represents the activation status of the cell cycle (Fig. 4). It should be emphasized that the
position of each node does not represent the energy.
255
Arranging the nodes according to their differentiation type, we obtained seven clusters of
fixed points. One cluster of fixed points possesses high levels of stemness signatures, which
consist of only transition states that will evolve into stable states with small perturbations,
compliant with the characteristics of stem cells cultured in vitro, and was categorized as glia
stem cells. Directly connected to this cluster of transition states are two clusters of fixed
points, one possessed high level of both oligodendrocyte and stemness signatures and was
categorized as oligodendrocyte precursor cells (OPCs), another possessed high level of both
astrocyte and stemness signatures was categorized as of astrocyte precursor cells (APCs). A
cluster of fixed points exhibiting low levels of all genes in the differentiation module was
identified as non-(macro)glia. Two clusters of fixed points with the lowest levels of stemness
and high levels of either astrocyte or oligodendrocyte differentiation signatures were
identified as astrocytes or oligodendrocytes. Notably, there is a cluster of transition states
that exhibits slightly high level of astrocytes-related signatures and slightly low level of
stemness and oligodendrocytes-related signatures, which is capable of transitioning into
multiple different differentiation types with small perturbations. We refer to this cluster as
astrocyte-like unstable progenitor cells, which might correspond to NG2+ cells (Fig. 4).
272
273 After the arrangement, we found that the potential energy landscape of glia obtained
274 encompasses the known glial differentiation hierarchy (Kriegstein and Alvarez-Buylla 2009;
275 Laug, Glasgow, and Deneen 2018; Allen and Lyons 2018), indicating that the overall results
276 up to this step under all mentioned modeling and computation are biologically acceptable.
277 We then further explored the glioma-related implications based on these results.
278
279

Locating the glioma states

To locate the glioma states, we validated the results at the module and pathway level at first. For a single-factor module or pathway, in this work, HIF, Ras, cellular senescence, and RTK, the value of the sole factor itself represents the level of the module or pathway. For a module or pathway that has both inhibitory and activatory nodes, with the hypothesis that each node in the core network as well as both the inhibitory and activatory part is equally important, we averaged the activation and inhibition levels and subtracted the inhibition level from the activation level, then normalizing the results to [0, 1] to obtain the module level results for multi-node modules. 0 represents complete inhibition, 1 represents complete activation, and values around 0.5 represent activation/inhibition balancing.

Stable states sta_2, sta_5, and transition state tra_6 exhibit high activation of the RTK, AKT, RAS, cell cycle, and HIF, as well as moderate activation of the NF-κB in module and pathway level. The p53, cellular senescence, and apoptosis modules are inhibited. This is fully consistent with the existing knowledge of glioma (J. Chen, McKay, and Parada 2012) (Fig. 5A, B). sta_2 exhibits a high level of Sox9, Notch, Pros, and NFIA, indicating a feature of low differentiated astrocytes; sta_5 exhibits a high level of Olig, Sox10, Notch, and Pros, indicating a feature of low differentiated oligodendrocytes; tra_6 exhibits a high level of Notch and Pros, as well as an intermediate level of Olig, Sox10, Sox9, and NFIA, indicating a feature of neuro/glia stem cells that have undergone gliogenic switch but have not yet differentiated into specific glial cell types (Elbaz and Popko 2019; Reiprich and Wegner 2015; Laug, Glasgow, and Deneen 2018; Wahlbuhl et al. 2012; Kato et al. 2011; Wang et al. 2020; Weider and Wegner 2017; Klum et al. 2018; J. S. Lee, Hoxha, and Song 2017; Glasgow et al. 2014; Samanta and Kessler 2004; Tchieu et al. 2019). Based on these results, we consider sta_2, sta_5, and tra_6 as glioma candidates, with sta_2 corresponding to astrocytoma cells, sta_5 corresponding to oligodendroglioma cells, and tra_6 corresponding to a type of glioma cell that embodied the feature of high stemness (Fig. 5A, B).

To make further confirmation of the correspondence of these three candidate states with glioma, considering that low-throughput methods, PCR, western blot, IHC, e.g., are still the gold standards for result verification in molecular biology, we then compared low-throughput experimental data with these three candidates at the molecular level. We obtained transcriptional and protein functional levels data on genes related to our model by reviewing literature from patient samples or cell lines(Chow et al. 2011; L. et al. 2012; Marumoto et al. 2009; Jin et al. 2018; Tanaka et al. 2011; J. S. Lee et al. 2014; G. Lee et al. 2016; Thomas et al. 2001; Y. Zhu et al. 2005; Rollbrocker et al. 1996; Sato et al. 2011; Kfoury et al. 2018; Reilly et al. 2000; Guo et al. 2011; Zheng et al. 2008; Rajasekhar et al. 2003; Simon et al. 1999) . After discretizing both our computational results and low-throughput data, we found that the consistency between the glioma states and the low-throughput data exceeded 95%, further confirming these candidates: sta_2, sta_5, and tra_6 as glioma states (Fig. 5C, D).

A single low-throughput experiment rarely covers all relevant genes in this model. Therefore, we further used high-throughput datasets to verify the computational results. We aligned

RNA-seq datasets of LGG and GBM from TCGA with our candidate glioma states. Our computational results normalized the final activity of the factors, with 0 representing the lowest possible level of coarse-grained gene function in the system, and 1 representing the highest possible level. Since there are no normal brain tissues in this dataset, genes that may be highly expressed in LGG and GBM may not reflect their actual levels after normalization. Therefore, we jointly analyzed the TCGA data with the cortex RNA-seq dataset from GTEx. We selected genes related to the network in this work and performed double-normalization. After normalization, the glioma candidate states are organized into the same clustering branch with TCGA data rather than GTEx data after hierarchical clustering. We then made a more detailed comparison by using the distribution of every selected gene from the above datasets (Fig. 5 F). The comparison results for each module are as follows: In the RTK module, EGFR, FGFR1, and PDGFRA are consistent with the calculation results; In the RAS pathway, KRAS and NRAS are consistent with the calculation results; In the AKT pathway, AKT1 is consistent with the calculation results, while AKT2 and PTEN have multiple peaks in the experimental data, which are partially consistent with the calculation results; In the NF-κB pathway, NFKBIA, NFKBIE, NFKB1, and NFKB2 are consistent with the calculation results; In the cell cycle module, CDK4 and CDK6 are consistent with the calculation results, while the proliferation-suppressing genes CDKN1 (p21) and RB-related genes are completely opposite to our calculation results; In the P53 pathway, the experimental data of the classic "tumor suppressor gene" TP53 is completely opposite to our calculation results, while MDM2 and MDM4 are consistent with our calculation results; In the characterization of cell senescence, CDKN2 family genes, CDKN2A-C data all showed a double peak, the calculation results are only consistent with some sample data, only the expression data of CDKN2D is consistent with our calculation results; In the HIF pathway, HIF1A is completely consistent with the calculation results; In the apoptosis module, the expression of BAD and XIAP genes is consistent with our calculation results, BCL2 and CASP9 are partially consistent with our calculation results, and the expression of BAX and CASP3 is completely opposite to our calculation results; In the differentiation module, the results of NOTCH1-3, PROS1, PROX1 are consistent with the calculation results, SOX9 and NFIA data both have multiple peaks, each peak value can correspond to the calculation results, OLIG2 is extremely high in TCGA-LGG, and in TCGA-GBM is distributed uniformly, judged as partially consistent, SOX10 is not fully present among all datasets thus cannot be normalized nor been compared. We then integrated the comparison results to the core node level, for genes with consistent subtypes with core nodes, we set the comparison result to 1; for genes with opposite results, we set them to 0; for partially consistent results, we set them to 0.5; if the comparison cannot be made, they are not included in the statistics, and the average value is calculated for each module. The final average value of each module is used as the overall consistency rate, with a result of 76.5% (Fig. 5 F).

Since the above datasets came from different databases, to avoid bias caused by batch effects, we further used the dataset GSE151352 which consists of paired para-tumor and tumor tissue from GBM patients for comparison. We selected the core node-related genes and obtained the fold change of the relevant genes in each pair of para-tumor and tumor tissues, normalizing them to [0, 1]. Results from this comparison are not entirely consistent with the

results obtained from TCGA-GTEx datasets. Among these: in the RTK module, EGFR and
FGFR1 are consistent with the computational results; in the Akt pathway, AKT1, AKT2, and
PTEN are consistent with the computational results; in the Ras pathway, HRAS and NRAS are
consistent with the computational results; in the HIF pathway, HIF1A is consistent with the
computational results; the marker of cell senescence, CDKN2A and CDKN2D in the CDKN2
family are consistent with the computational results; all subtypes in the NF-kB pathway are
consistent with the computational results; the p53 pathway is opposite to the computational
results; in the cell cycle module, all are consistent with the computational results except for
CDKN1A; in the apoptosis module, all are inconsistent with the computational results except
for BCL2 and XIAP; in the differentiation module, all NOTCH subtypes and PROS1 are
consistent with the computational results, while NFIA, SOX9, SOX10, and Olig1-3 are partially
consistent with the computational results. The overall consistency rate is 77.2% (Fig. 5 E).
380
Since the computational results in this work represent the functional level of the core nodes
related gene, multi-omics analysis has already revealed that the correlation between the
transcriptome and proteome is weak (Williams et al. 2016), further considering the inevitable
cell type heterogeneity in tissue samples (Brat and Meir 2004; Suvà and Tirosh 2020), thus a
perfect consistency between our results and the transcriptome cannot be expected. The
consistency rate (76.5%, 77.2%) of high-throughput data validation is acceptable.
387
388   In summary, we can confidently assert that the glioma candidate states: sta_2, sta_5, and tra_6
389   correspond to glioma.
390
391
392   **Landscape near glioma**
393
In the results aforementioned, we have portraited the energy landscape of the glial system
and located the glioma states. To address the questions mentioned at the beginning, we
need to take a close look at the landscape near glioma. By selecting the 1st neighboring
stable states which connect to the glioma states through only one transition state as well as
the transition states connecting these stable states (Fig. 6 A, B), we obtained the energy
landscape near glioma, consisting of a total of 12 stable states and 22 transition states (Fig.
6 C, D).
401
402
403   **GSC state is the energy barrier separating astrocytoma and oligodendroglioma**
404
405   All glioma states we acquired possess the feature of high stemness (Fig. 4; Fig. 5 C, D), which
406   has been already observed by the analysis of single-cell sequencing data that most highly
407   proliferative cells in samples of low grade glioma as well as glioblastoma exhibit high stemness
408   (Suvà and Tirosh 2020).
409
Among these glioma states, a transition state, tra_6, is present. It possesses a signature of
activated cell cycle, and high levels of glia stemness-related genes, Notch and Pros, as the

mediated level of astrocyte differentiation-related genes, NFIA and Sox9, as well as
oligodendrocyte differentiation-related genes, Olig and Sox10, representing low-
differentiated glial cells that have undergone a gliogenic switch but remain high stemness
(Laug, Glasgow, and Deneen 2018; Klum et al. 2018; Reiprich and Wegner 2015). tra_6 is an
unstable fixed point, a transition state. With a small perturbation, it will and can only evolve
towards the two glioma stable states, sta_2 and sta_5, consists with the knowledge of the
functional features of GSCs (Lathia et al. 2015) (Fig. 7 A).
419
Further, the landscape indicates that a stable glioma state can evolve into another via tra_6
following significant targeted intervention or accumulated minor perturbations. When such
transitions occur in large numbers, the transition state tra_6 can be clearly observed. This
has also been observed experimentally, where a portion of rat GBM cell line C6
spontaneously exhibited stem-cell-like phenotype (De et al. 2020), as well as a
developmental cell type "outer radial glia" being re-activated in GBM (Bhaduri et al. 2020).
426
A stable state needs sufficient time to evolve into another under the accumulation of minor
perturbations, and the states that have crossed the energy barrier are exponentially
distributed compared to non-transitioned states, meaning there are far fewer crossed cells
than non-crossed ones. Suvà and Tirosh observed that the number of differentiated cells
consistent with histological grading far exceeded those of another differentiated cell type
(Suvà and Tirosh 2020), which again aligns with our theoretical results.
433
434   Based on the comparison between our theoretical results and the experimental observation
435   mentioned above, only tra_6 in the entire glia landscape meets the features of glioma stem
436   cells, thus we considered tra_6 to be the GSCs state (Fig. 7 B, C).
437
438
439   **Stable states near glioma are cells of glioma origin**
440
In the energy landscape near glioma, all non-glioma stable states only need to traverse one
transition state to transform into a glioma state (Fig. 6 A, B), meaning there is only one energy
barrier to prevent these non-glioma stable states correlated cells from the fate of
carcinogenesis, thus these states are all considered to be the most likely to transform into
glioma states among the global energy landscape, in other words, cells correspond with these
stable states are the cells of glioma origin, except the all-zero death state sta_17 (Fig. 6 C).
447
448   Sorted by differentiation module, they can be divided into groups of high stemness astrocytes
449   or APCs (sta_1), high stemness oligodendrocytes OPCs (sta_6), differentiated
450   oligodendrocytes (sta_11, sta_13, sta_18, and sta_19), and non-(macro) glia states that cannot
451   be identified based on this study, which may be ependymal cells, neurons or mesothelial cells
452   (sta_14, sta_16, sta_20) (Fig. 6 A, C, D).
453

454 60% of these stable states exhibit the feature of activated apoptosis module (sta_1, sta_6,
455 sta_11, sta_13, sta_16, and sta_20), indicate the possibility that cells entered apoptosis
456 program may also have the potential to enter the carcinogenesis progress (Fig. 6 C, D).
457
458 Previous researches reported that cellular senescence promotes glioma progression (Salam
459 et al. 2023; Chojak et al. 2023), which also observed in our results. 60% of the glioma origin
460 states (sta_1, sta_6, sta_11, sta_14, sta_19, and sta_20) exhibit activated cellular senescence
461 module represented by the gene of CDKN2A which translates into p16^{INK4} (Fig. 6 C, D).
462
Experiments suggest that NSCs, mature astrocytes and OPCs could all potentially be cells of
glioma origin (Zong, Verhaak, and Canolk 2012). Similarly, our results showed that astrocytes
with high stemness and undergoing apoptosis are one type of cells of glioma origin, while
differentiated astrocytes are not the cells of glioma origin. Oligodendrocytes with high
stemness and undergoing apoptosis are also cells of glioma origin. Differentiated
oligodendrocytes may also be cells of glioma origin, as well as non-(macro) glia cells which
differentiation type are undistinguished based on this study solely.
470
Our results show that only tra_6 is the GSCs state in the landscape of glia, and tra_6, as a
transition state, is unstable. That is, any small noise will cause it to evolve towards its
neighboring stable states, in which only two glioma states exist (Fig. 7 B). Due to its unstable
nature, the only condition of its long-term existence is always maintaining a subtle balance
to stay in static equilibrium in the *in vivo* environment where noises constantly exist (Chang
et al. 2008), otherwise it will soon transform into a stable state with less stemness. After that,
it would only flash up during the path of the transformation from one stable glioma state to
another. Thus, GSCs are unlikely to be the cells of glioma origin.
479
480
481 **Heterogeneity Inducing Transition States are Potential Mechanism Behind Therapy**
482 **Resistance and Glioma Recurrence**
483
In the energy landscape near glioma, all transition states (except tra_6) connect glioma origin
states and glioma states, serving as potential routes for tumorigenesis and also as
intermediate states between somatic glia and glioma (Fig. 6 C, D). These transition states can
be divided into two categories based on their connections to glioma states:
The first type of transition states connects only one glioma state, such as those connecting
only sta_2 and somatic states (tra_4, tra_13, tra_14, tra_21, tra_23, tra_38, tra_48) and those
connecting only sta_5 and somatic states (tra_7, tra_34, tra_37, tra_39, tra_45, tra_52, tra_54)
(Fig. 7 D, E, F);
The second type of transition states simultaneously connects both types of glioma states,
including tra_6, tra8, tra9, tra15, tra18, tra31, tra33 and tra_51 (Fig. 7 G, H, I). Interestingly, all
heterogeneity inducing transition states exhibit high stemness molecularly.
495
When driven from one glioma state through the first type of transition states, the system will
only transfer into somatic stable states, which can be considered as potential cures (Fig. 6 C,

LEFT & RIGHT; Fig. 7 D, E, F). However, if driven towards the second type of transition states,
there is the possibility that the system transfers from one type of glioma stable state to
another as well as multiple somatic states, resulting in tumor heterogeneity (Fig. 6 C, MIDDLE,
Fig. 7 G, H, I). As long as the paths toward these heterogeneity inducing transition states exist,
it would be difficult to reach a true cure, these treatment behaviors may further lead to an
increase in the clinical grade of glioma.

504
505
506 **Conclusion**

507
508 In this work, by taking a systemic theoretical perspective and integrating experiment-based
509 knowledge, we constructed an endogenous network for glioma and depicted the energy
510 landscape of the glioma system through network dynamics computation. By comparing
511 theoretical computations with multi-level knowledge and data, we were able to locate the
512 glioma states.

513
514 Our results revealed two stable states corresponding to astrocytoma and
515 oligodendroglioma, interconnected by a transition state corresponding to GSC. This
516 suggests that GSCs are unstable in vivo and are unlikely to be the cellular origin of glioma.
517 The landscape near glioma presented several stable states of diverse differentiation types,
518 further confirming the multicellular origins of glioma theoretically.

519
520 In the landscape near glioma, a group of transition states that simultaneously connected
521 two types of glioma stable states displayed characteristics of non-proliferation and high
522 stemness. If the system is driven through these transition states, this would lead to increased
523 tumor heterogeneity. If these heterogeneity inducing transition states are not blocked
524 during treatment, the true cure of glioma will be difficult to achieve.

525
526
527 **Discussion**

528
Differentiation therapy for glioma has already been called for (Laug, Glasgow, and Deneen,
2018; Suvà and Tirosh, 2020). Here, our theoretical results preliminarily suggest that the
transitional states connecting two glioma stable states are responsible for glioma
heterogeneity and resistance to radio-chemotherapy. These heterogeneity inducing
transition states all demonstrate features of high stemness and low proliferation. Therefore,
apoptosis-promoting therapies and differentiation-promoting therapies combined could be
promising strategy in the treatment of glioma.

536
The model we used in this study is an "average field" model for the "average person," and
does not consider the genetic polymorphisms existing within populations. The network
structure and parameters can be adjusted to further accommodate such genetic
polymorphisms. Strategies for molecular therapies already target the core genes in our model
(Rajesh et al., 2017). Our method can serve as an *in silico* navigation tool for developing

personal cocktail therapies. However, we must concede that our 25-node model is overly
coarse-grained, and future work should refine and enlarge the model.
544
Experimental validation of the theoretical results in this study is all retrospective. The following
prospective experiments, such as investigating the presence of GSCs in the purified glioma
cell line or implant tumors originated from differentially-purified glioma cells, can be used to
verify our prediction that glioma stem cells serve as transition states connecting glioma stable
states as well as that the heterogeneity and treatment resistance are mediated by
heterogeneity inducing transition states:
For glioma cell lines or primary glioma cells, knock-in fluorescent protein markers following
differentiation-related transcription factors such as the Sox family, NFIA, Olig, Pros, and Notch
family mentioned in this work. Using flow cytometry to purify astrocytoma-like tumor cells as
well as oligodendroglioma-like tumor cells. Observing the fluorescence signal during *in vitro*
culturing and passaging. We expect to observe homogeneous fluorescence signals becoming
heterogeneous during culture and passage with ground noise. Using microfluidic chips for
long-term *in vitro* observation. We expect to observe the cell fate switching as observed in
the system described by Lord et al. (Lord et al. 2019). Performing tumor implant experiments
with purified cells. We expect to observe more LGG phenotypes than GBM phenotypes
(implant tumor group); if low-dose apoptosis-inducing therapy is applied, the incidence of
GBM phenotypes will increase (implant tumor + low dose treatment group); if glioma
differentiation is promoted during low-dose apoptosis-inducing therapy, the incidence of
GBM phenotypes will be lower than in the implant tumor + low-dose treatment group.
564
Research on gene regulatory networks has significantly advanced our understanding in recent
decades. However, the integration of this vast array of static knowledge into a systemic
framework that can yield dynamic insights has remained a challenge. In this study, we have
attempted to bridge this gap by integrating experimental knowledge to construct a core
endogenous network for glioma. By incorporating the concept of the landscape, rooted in
biological principles and extensively applied in statistical physics, we have sought to generate
novel insights. Our theoretical findings, while promising, represent a preliminary step that calls
for further experimental validation. They offer a glimpse into the complex interplay between
theoretical models and experimental research, a relationship characterized by mutual
reinforcement and continuous dialogue. As we move forward, it is with cautious optimism
that we envision our approach contributing to a deeper understanding of glioma biology and,
ultimately, to the development of more effective therapeutic interventions.
577
578
Methods
Network dynamics
Boolean Dynamics
Boolean dynamics is a useful and relatively swift tool to obtain architectural information of
the network dynamics results (Bornholdt 2008). The typical Boolean function is shown in figure
1 using Akt as an example: each variable has a discrete value 0 or 1 representing low/high
level separately; t denotes the current state of the variables, t + 1 for the next state; NOT,

AND, and OR are the usual logic operators. An equivalent formulation with threshold function
is used in Matlab for our computation, details of the algorithms have been described in Yuan
et al. 2017, the functions used in this work can be found in Supplementary Table 1.
ODE
We use two different algorithms to calculate stable states, details of the algorithms have been
described in Yuan et al. 2017, the functions used in this work can be found in Supplementary
Table 1.
Algorithms 1 (Euler's method): We initiate the procedure by generating a random starting
point, denoted as the initial vector. Utilizing the dynamic system's rules, we enter a loop of
iterative calculations to progressively update the vector using a time step multiplication factor.
The iterative process is designed to continue until we reach a predefined number of steps,
ensuring the vector's convergence to stability, which is ascertained through specific
convergence criteria involving preset thresholds. Once a candidate for stability meets the
convergence criteria, we classify and record it as a stable state.
Algorithms 2 (Newton's method):
we employed a numerical method to solve the nonlinear equation representing the rate of
change of the system variable over time. Similarly, our process began by initializing a random
vector as a starting point for the iterative procedure. Using the fsolve function in MATLAB, we
applied Newton's method to locate the solution vector that zeroes the function, which
signifies a potential stable state. To ascertain the stability of this solution, we evaluated the
eigenvalues of the Jacobian matrix at the solution point, considering only those solutions as
stable states where all eigenvalues have negative real parts, and those solutions where
positive eigenvalues exist are considered as transition states. This procedure was
systematically repeated to compile a comprehensive list of stable states.
610
State interconnection analysis
Our algorithm is intended to generate trajectories leading from each unstable state to its
corresponding stable states. To facilitate this, we introduced minor perturbations to the
system via small, random vectors while it was positioned at an identified unstable state. This
enabled us to trace the pathways from the unstable state to each linked stable state. The
details can be seen in Supplementary table 1.
617
Modelized the ODE results
In our study, the variable 'x' in the ordinary differential equations (ODE) represents the
concentration or biological activity of nodes within the network. We have translated
computational results to the module level to align with established biological knowledge.
Following equation was used to determine the module status:

$$Module\ Status = 0.5 \times \left(\frac{\sum_n x_{act}}{n} - \frac{\sum_n x_{inh}}{n} + 1 \right)$$

Nodes within functional modules and signaling pathways were categorized as either
activatory (x_{act}) or inhibitory (x_{inh}), with the exception of the differentiation module. The final
module status is determined by threshold values: an 'OFF' state is indicated when the Module
Status is below 0.33, signifying inhibition; an 'ON' state is indicated when the Module Status
exceeds 0.66, denoting activation; and a 'Stable' (STA) state is assigned when the Module

Status falls between 0.33 and 0.66, reflecting a balance between activatory and inhibitory influences within the module. In a module with internal negative feedback, taking NF-κB pathway as an example: a STA state might suggest a balance similar to a 2:1 ratio of NF-κB subunits to their inhibitor pIkB forming a complex; an 'ON' state would imply a preponderance of free NF-κB subunits not complexed with pIkB and thus free to enter the nucleus and exert their function; an 'OFF' state would indicate a dominance of inhibitory genes, such as a high level of IkB relative to NF-κB subunits.

High-Throughput Data Preprocessing

TCGA-GTEx
We acquired RNAseq data (TPM) from the GTEx cortex samples and the TCGA GBMLGG dataset. Randomly selecting 150 samples each from the cortex, GBM, and LGG collections, we performed a two-step scaling on the datasets. Initially, we computed the z-score for each sample to mitigate batch effects, and subsequently, we calculated the z-score for each gene. Values exceeding one standard deviation were capped to this threshold, followed by normalization to the [0, 1] range. The continuous distribution of gene expression was plotted by ggplot2 package in R.

GSE151352
Data from GSE151352 was retrieved from GEO database, converted into an exponential format (e), and its standard error was computed. We determined an effective information threshold for this dataset by averaging the standard errors of the genes with the top 200 smallest standard errors ($\approx 30\%$). The original dataset was manipulated by subtracting the para-cancerous sample values from the tumor sample values to derive the fold change. A high expression threshold was set at $\ln(1.3)$, capping fold changes above this threshold to $\ln(1.3)$; a low expression threshold was set at $\ln(1/1.3)$, with values below this threshold adjusted to $\ln(1/1.3)$. Subsequently, the data was normalized to the range [0, 1].

Reference
Agostini, Massimiliano, Gerry Melino, and Francesca Bernassola. 2018. "The P53 Family in Brain Disease." *Antioxidants and Redox Signaling* 29 (1): 1–14. https://doi.org/10.1089/ars.2017.7302.
Allen, Nicola J, and David A Lyons. 2018. "Glia as Architects of Central Nervous System Formation and Function." *Science* 362 (6411): 181–85. https://doi.org/10.1126/science.aat0473.
Annenkov, Alexander. 2014. "Receptor Tyrosine Kinase (RTK) Signalling in the Control of Neural Stem and Progenitor Cell (NSPC) Development." *Molecular Neurobiology* 49 (1): 440–71. https://doi.org/10.1007/s12035-013-8532-5.
Ao, Ping. 2005. "Laws in Darwinian Evolutionary Theory." *Physics of Life Reviews* 2 (2): 117–56. https://doi.org/10.1016/j.plrev.2005.03.002.
———. 2008. "Article Commentary: Borges Dilemma, Fundamental Laws, and Systems Biology." *Bioinformatics and Biology Insights* 2 (January): 117793220800200. https://doi.org/10.1177/117793220800200002.

Ao, Ping, David Galas, Leroy Hood, and Xiaomei Zhu. 2008. "Cancer as Robust Intrinsic State of
Endogenous Molecular-Cellular Network Shaped by Evolution." *Medical Hypotheses* 70 (3):
678–84. https://doi.org/10.1016/j.mehy.2007.03.043.

Badia-i-Mompel, Pau, Lorna Wessels, Sophia Müller-Dott, Rémi Trimbour, Ricardo O. Ramirez
Flores, Ricard Argelaguet, and Julio Saez-Rodriguez. 2023. "Gene Regulatory Network
Inference in the Era of Single-Cell Multi-Omics." *Nature Reviews Genetics* 24 (November):
739–54. https://doi.org/10.1038/s41576-023-00618-5.

Banfalvi, Gaspar. 2011. "Chapter 2 Synchronization of Mammalian Cells and Nuclei by Centrifugal
Elutriation," 25–45. https://doi.org/10.1007/978-1-61779-182-6.

Berezowska, Sabina, and Jurgen Schlegel. 2011. "Targeting ErbB Receptors in High-Grade
Glioma." *Current Pharmaceutical Design* 17 (23): 2468–87.
https://doi.org/10.2174/138161211797249233.

Bertoli, Cosetta, Jan M. Skotheim, and Robertus A.M. De Bruin. 2013. "Control of Cell Cycle
Transcription during G1 and S Phases." *Nature Reviews Molecular Cell Biology* 14 (8): 518–
28. https://doi.org/10.1038/nrm3629.

Bhaduri, Aparna, Elizabeth Di Lullo, Diane Jung, Sören Müller, Elizabeth Erin Crouch, Carmen
Sandoval Espinosa, Tomoko Ozawa, et al. 2020. "Outer Radial Glia-like Cancer Stem Cells
Contribute to Heterogeneity of Glioblastoma." *Cell Stem Cell* 26 (1): 48-63.e6.
https://doi.org/10.1016/j.stem.2019.11.015.

Biserova, Karina, Arvids Jakovlevs, Romans Uljanovs, and Ilze Strumfa. 2021. "Cancer Stem Cells:
Significance in Origin, Pathogenesis and Treatment of Glioblastoma." *Cells* 10 (3): 1–20.
https://doi.org/10.3390/cells10030621.

Bornholdt, Stefan. 2008. "Boolean Network Models of Cellular Regulation: Prospects and
Limitations." *Journal of the Royal Society Interface* 5 (SUPPL. 1).
https://doi.org/10.1098/rsif.2008.0132.focus.

Brat, Daniel J, and Erwin G Van Meir. 2004. "Vaso-Occlusive and Prothrombotic Mechanisms
Associated with Tumor Hypoxia , Necrosis , and Accelerated Growth in Glioblastoma."
Laboratory Investigation, no. 84: 397–405. https://doi.org/10.1038/labinvest.3700070.

Buchanan, Mark. 2013. "Change Is Good." *Nature Physics* 9 (11): 690.
https://doi.org/10.1038/nphys2803.

Calcinotto, Arianna, Jaskaren Kohli, Elena Zagato, Laura Pellegrini, Marco Demaria, and Andrea
Alimonti. 2019. "Cellular Senescence: Aging, Cancer, and Injury." *Physiological Reviews* 99
(2): 1047–78. https://doi.org/10.1152/physrev.00020.2018.

Capp, Jean Pascal. 2019. "Cancer Stem Cells: From Historical Roots to a New Perspective."
Journal of Oncology 2019. https://doi.org/10.1155/2019/5189232.

Chang, Hannah H., Martin Hemberg, Mauricio Barahona, Donald E. Ingber, and Sui Huang. 2008.
"Transcriptome-Wide Noise Controls Lineage Choice in Mammalian Progenitor Cells."
Nature 453 (7194): 544–47. https://doi.org/10.1038/nature06965.

Chen, Jian, Renée M. McKay, and Luis F. Parada. 2012. "Malignant Glioma: Lessons from
Genomics, Mouse Models, and Stem Cells." *Cell* 149 (1): 36–47.
https://doi.org/10.1016/j.cell.2012.03.009.

Chen, Kok Siong, Jonathan W.C. Lim, Linda J. Richards, and Jens Bunt. 2017. "The Convergent
Roles of the Nuclear Factor I Transcription Factors in Development and Cancer." *Cancer
Letters* 410: 124–38. https://doi.org/10.1016/j.canlet.2017.09.015.

Chen, Yong Cong, Chunxiao Shi, J. M. Kosterlitz, Xiaomei Zhu, and Ping Ao. 2020. "Global Potential, Topology, and Pattern Selection in a Noisy Stabilized Kuramoto-Sivashinsky Equation." *Proceedings of the National Academy of Sciences of the United States of America* 117 (38): 23227–34. https://doi.org/10.1073/pnas.2012364117.

Chojak, Rafał, Jawad Fares, Edgar Petrosyan, and Maciej S. Lesniak. 2023. "Cellular Senescence in Glioma." *Journal of Neuro-Oncology*. https://doi.org/10.1007/s11060-023-04387-3.

Choudhry, Hani, and Adrian L. Harris. 2018. "Advances in Hypoxia-Inducible Factor Biology." *Cell Metabolism* 27 (2): 281–98. https://doi.org/10.1016/j.cmet.2017.10.005.

Chow, Lionel M.L., Raelene Endersby, Xiaoyan Zhu, Sherri Rankin, Chunxu Qu, Junyuan Zhang, Alberto Broniscer, David W. Ellison, and Suzanne J. Baker. 2011. "Cooperativity within and among Pten, P53, and Rb Pathways Induces High-Grade Astrocytoma in Adult Brain." *Cancer Cell* 19 (3): 305–16. https://doi.org/10.1016/j.ccr.2011.01.039.

Csermely, Peter, János Hódsági, Tamás Korcsmáros, Dezso Módos, Áron R. Perez-Lopez, Kristóf Szalay, Dániel V. Veres, Katalin Lenti, Ling Yun Wu, and Xiang Sun Zhang. 2015. "Cancer Stem Cells Display Extremely Large Evolvability: Alternating Plastic and Rigid Networks as a Potential Mechanism. Network Models, Novel Therapeutic Target Strategies, and the Contributions of Hypoxia, Inflammation and Cellular Senescence." *Seminars in Cancer Biology* 30: 42–51. https://doi.org/10.1016/j.semcancer.2013.12.004.

Czabotar, Peter E., Guillaume Lessene, Andreas Strasser, and Jerry M. Adams. 2014. "Control of Apoptosis by the BCL-2 Protein Family: Implications for Physiology and Therapy." *Nature Reviews Molecular Cell Biology* 15 (1): 49–63. https://doi.org/10.1038/nrm3722.

De, Arpan, Dilshan H. Beligala, Vishal P. Sharma, Christian A. Burgos, Angelia M. Lee, and Michael E. Geusz. 2020. "Cancer Stem Cell Generation during Epithelial-Mesenchymal Transition Is Temporally Gated by Intrinsic Circadian Clocks." *Clinical and Experimental Metastasis*, no. 0123456789. https://doi.org/10.1007/s10585-020-10051-1.

Elbaz, Benayahu, and Brian Popko. 2019. "Molecular Control of Oligodendrocyte Development." *Trends in Neurosciences*. https://doi.org/10.1016/j.tins.2019.01.002.

Eltzschig, Holger K., Donna L. Bratton, and Sean P. Colgan. 2014. "Targeting Hypoxia Signalling for the Treatment of Ischaemic and Inflammatory Diseases." *Nature Reviews Drug Discovery* 13 (11): 852–69. https://doi.org/10.1038/nrd4422.

Feinberg, Andrew P., and Andre Levchenko. 2023. "Epigenetics as a Mediator of Plasticity in Cancer." *Science* 379 (6632). https://doi.org/10.1126/science.aaw3835.

Fricker, Michael, Aviva M. Tolkovsky, Vilmante Borutaite, Michael Coleman, and Guy C. Brown. 2018. "Neuronal Cell Death." *Physiological Reviews* 98 (2): 813–80. https://doi.org/10.1152/physrev.00011.2017.

Fruman, David A., and Christian Rommel. 2014. "PI3K and Cancer: Lessons, Challenges and Opportunities." *Nature Reviews Drug Discovery* 13 (2): 140–56. https://doi.org/10.1038/nrd4204.

Galli, R., E. Binda, U. Orfanelli, B. Cipelletti, A. Gritti, S. De Vitis, R. Fiocco, C. Foroni, F. Dimeco, and A. Vescovi. 2004. "Erratum: Isolation and Characterization of Tumorigenic, Stem-like Neural Precursors from Human Glioblastoma (Cancer Research (October 2004) 64 (7011-7021)." *Cancer Research* 64 (21): 8130.

Gan, Xiaoliang, Haoyu Wang, and Ping Ao. 2021. "Existence of a Smooth Lyapunov Function for Any Smooth Planar Dynamical System with One Limit Cycle." *Nonlinear Dynamics* 105 (4):

3117–30. https://doi.org/10.1007/s11071-021-06775-8.

Gérard, Claude, and Albert Goldbeter. 2016. "Dynamics of the Mammalian Cell Cycle in

Physiological and Pathological Conditions." *Wiley Interdisciplinary Reviews: Systems*

Biology and Medicine 8 (2): 140–56. https://doi.org/10.1002/wsbm.1325.

Giacomantonio, Clare E., and Geoffrey J. Goodhill. 2010. "A Boolean Model of the Gene

Regulatory Network Underlying Mammalian Cortical Area Development." *PLoS*

Computational Biology 6 (9). https://doi.org/10.1371/journal.pcbi.1000936.

Glasgow, Stacey M., Wenyi Zhu, C. Claus Stolt, Teng Wei Huang, Fuyi Chen, Joseph J. LoTurco,

Jeffrey L. Neul, Michael Wegner, Carrie Mohila, and Benjamin Deneen. 2014. "Mutual

Antagonism between Sox10 and NFIA Regulates Diversification of Glial Lineages and

Glioma Subtypes." *Nature Neuroscience* 17 (10): 1322–29. https://doi.org/10.1038/nn.3790.

Gonzalez, Frank J., Cen Xie, and Changtao Jiang. 2018. "The Role of Hypoxia-Inducible Factors in

Metabolic Diseases." *Nature Reviews Endocrinology* 15 (1): 21–32.

https://doi.org/10.1038/s41574-018-0096-z.

Goodhill, Geoffrey J. 2018. "Theoretical Models of Neural Development." *IScience* 8: 183–99.

https://doi.org/10.1016/j.isci.2018.09.017.

Gorgoulis, Vassilis, Peter D. Adams, Andrea Alimonti, Dorothy C. Bennett, Oliver Bischof, Cleo

Bishop, Judith Campisi, et al. 2019. "Cellular Senescence: Defining a Path Forward." *Cell* 179

(4): 813–27. https://doi.org/10.1016/j.cell.2019.10.005.

Guo, Deliang, Felicia Reinitz, Mary Youssef, Cynthia Hong, David Nathanson, David Akhavan,

Daisuke Kuga, et al. 2011. "An LXR Agonist Promotes Glioblastoma Cell Death through

Inhibition of an EGFR/AKT/ SREBP-1/LDLR-Dependent Pathway." *Cancer Discovery* 1 (5):

442–56. https://doi.org/10.1158/2159-8290.CD-11-0102.

Hafner, Antonina, Martha L. Bulyk, Ashwini Jambhekar, and Galit Lahav. 2019. "The Multiple

Mechanisms That Regulate P53 Activity and Cell Fate." *Nature Reviews Molecular Cell*

Biology 20 (4): 199–210. https://doi.org/10.1038/s41580-019-0110-x.

Hanahan, Douglas, and Robert A. Weinberg. 2011. "Hallmarks of Cancer: The next Generation."

Cell. Elsevier Inc. https://doi.org/10.1016/j.cell.2011.02.013.

Harashima, Hirofumi, Nico Dissmeyer, and Arp Schnittger. 2013. "Cell Cycle Control across the

Eukaryotic Kingdom." *Trends in Cell Biology* 23 (7): 345–56.

https://doi.org/10.1016/j.tcb.2013.03.002.

Hata, Aaron N., Jeffrey A. Engelman, and Anthony C. Faber. 2015. "The BCL2 Family: Key

Mediators of the Apoptotic Response to Targeted Anticancer Therapeutics." *Cancer*

Discovery 5 (5): 475–87. https://doi.org/10.1158/2159-8290.CD-15-0011.

Hemmati, Houman D., Ichiro Nakano, Jorge A. Lazareff, Michael Masterman-Smith, Daniel H.

Geschwind, Marianne Bronner-Fraser, and Harley I. Kornblum. 2003. "Cancerous Stem Cells

Can Arise from Pediatric Brain Tumors." *Proceedings of the National Academy of Sciences*

of the United States of America 100 (25): 15178–83.

https://doi.org/10.1073/pnas.2036535100.

Hidalgo, Alicia, and Ann Logan. 2017. "Go and Stop Signals for Glial Regeneration." *Current*

Opinion in Neurobiology 47: 182–87. https://doi.org/10.1016/j.conb.2017.10.011.

Hoang, Thanh, Jie Wang, Patrick Boyd, Fang Wang, Clayton Santiago, Lizhi Jiang, Sooyeon Yoo,

et al. 2020. "Gene Regulatory Networks Controlling Vertebrate Retinal Regeneration."

Science 370 (6519). https://doi.org/10.1126/science.abb8598.

Huang, Sui, Ingemar Ernberg, and Stuart Kauffman. 2009. "Cancer Attractors: A Systems View of Tumors from a Gene Network Dynamics and Developmental Perspective." *Seminars in Cell and Developmental Biology* 20 (7): 869–76. https://doi.org/10.1016/j.semcdb.2009.07.003.

Ignatova, Tatyana N., Valery G. Kukekov, Eric D. Laywell, Oleg N. Suslov, Frank D. Vrionis, and Dennis A. Steindler. 2002. "Human Cortical Glial Tumors Contain Neural Stem-like Cells Expressing Astroglial and Neuronal Markers in Vitro." *Glia* 39 (3): 193–206. https://doi.org/10.1002/glia.10094.

Janku, Filip, Timothy A. Yap, and Funda Meric-Bernstam. 2018. "Targeting the PI3K Pathway in Cancer: Are We Making Headway?" *Nature Reviews Clinical Oncology* 15 (5): 273–91. https://doi.org/10.1038/nrclinonc.2018.28.

Jin, Peng, Seung Hyun Shin, Yang Sook Chun, Hyun Woo Shin, Yong Jae Shin, Yeri Lee, Donggeon Kim, Do Hyun Nam, and Jong Wan Park. 2018. "Astrocyte-Derived CCL20 Reinforces HIF-1-Mediated Hypoxic Responses in Glioblastoma by Stimulating the CCR6-NF-KB Signaling Pathway." *Oncogene* 37 (23): 3070–87. https://doi.org/10.1038/s41388-018-0182-7.

Joerger, Andreas C., and Alan R. Fersht. 2016. "The P53 Pathway: Origins, Inactivation in Cancer, and Emerging Therapeutic Approaches." *Annual Review of Biochemistry* 85 (1): 375–404. https://doi.org/10.1146/annurev-biochem-060815-014710.

Kamakura, Sachiko, Koji Oishi, Takeshi Yoshimatsu, Masato Nakafuku, Norihisa Masuyama, and Yukiko Gotoh. 2004. "Hes Binding to STAT3 Mediates Crosstalk between Notch and JAK-STAT Signalling." *Nature Cell Biology* 6 (6): 547–54. https://doi.org/10.1038/ncb1138.

Kang, Peng, Hyun Kyoung Lee, Stacey M. Glasgow, Meggie Finley, Tataka Donti, Zachary B. Gaber, Brett H. Graham, et al. 2012. "Sox9 and NFIA Coordinate a Transcriptional Regulatory Cascade during the Initiation of Gliogenesis." *Neuron* 74 (1): 79–94. https://doi.org/10.1016/j.neuron.2012.01.024.

Kastenhuber, Edward R., and Scott W. Lowe. 2017. "Putting P53 in Context." *Cell* 170 (6): 1062–78. https://doi.org/10.1016/j.cell.2017.08.028.

Kato, Kentaro, Manuel G. Forero, Janine C. Fenton, and Alicia Hidalgo. 2011. "The Glial Regenerative Response to Central Nervous System Injury Is Enabled by Pros-Notch and Pros-NFκB Feedback." *PLoS Biology* 9 (8): 6–9. https://doi.org/10.1371/journal.pbio.1001133.

Kauffman, S. A. 1969. "Metabolic Stability and Epigenesis in Randomly Constructed Genetic Nets." *Journal of Theoretical Biology* 22 (3): 437–67. https://doi.org/10.1016/0022-5193(69)90015-0.

Kfoury, Najla, Tao Sun, Kwanha Yu, Nathan Rockwell, Kelsey L. Tinkum, Zongtai Qi, Nicole M. Warrington, et al. 2018. "Cooperative P16 and P21 Action Protects Female Astrocytes from Transformation." *Acta Neuropathologica Communications* 6 (1): 12. https://doi.org/10.1186/s40478-018-0513-5.

Kim, Daniel, Andy Tran, Hani Jieun Kim, Yingxin Lin, Jean Yee Hwa Yang, and Pengyi Yang. 2023. "Gene Regulatory Network Reconstruction: Harnessing the Power of Single-Cell Multi-Omic Data." *Npj Systems Biology and Applications* 9 (1): 51. https://doi.org/10.1038/s41540-023-00312-6.

Kirk, P D W, A C Babtie, M P H Stumpf, S. Brenner, E. U. Azeloglu, R. Iyengar, I. G. Johnston, et al. 2015. "SYSTEMS BIOLOGY. Systems Biology (Un)Certainties." *Science (New York, N.Y.)* 350

(6259): 386–88. http://www.ncbi.nlm.nih.gov/pubmed/26494748.

Klum, Susanne, Cécile Zaouter, Zhanna Alekseenko, Åsa K Björklund, Daniel W Hagey, Johan
Ericson, Jonas Muhr, and Maria Bergsland. 2018. "Sequentially Acting SOX Proteins
Orchestrate Astrocyte- and Oligodendrocyte-specific Gene Expression." *EMBO Reports* 19
(11): e46635. https://doi.org/10.15252/embr.201846635.

Kriegstein, Arnold, and Arturo Alvarez-Buylla. 2009. "The Glial Nature of Embryonic and Adult
Neural Stem Cells." *Annual Review of Neuroscience* 32 (1): 149–84.
https://doi.org/10.1146/annurev.neuro.051508.135600.

Kritsilis, Marios, Sophia V. Rizou, Paraskevi N. Koutsoudaki, Konstantinos Evangelou, Vassilis G.
Gorgoulis, and Dimitrios Papadopoulos. 2018. "Ageing, Cellular Senescence and
Neurodegenerative Disease." *International Journal of Molecular Sciences* 19 (10).
https://doi.org/10.3390/ijms19102937.

Kuhlbrodt, Kirsten, Beate Herbarth, Elisabeth Sock, Irm Hermans-Borgmeyer, and Michael
Wegner. 1998. "Sox10, a Novel Transcriptional Modulator in Glial Cells." *The Journal of*
Neuroscience 18 (1): 237–50. https://doi.org/10.1523/JNEUROSCI.18-01-00237.1998.

Kwon, Chulan, Ping Ao, and David J Thouless. 2005. "Structure of Stochastic Dynamics near Fixed
Points" 102 (37): 13029–33. https://doi.org/10.1073/pnas.0506347102.

Kyrousi, Christina, Zoi Lygerou, and Stavros Taraviras. 2017. "How a Radial Glial Cell Decides to
Become a Multiciliated Ependymal Cell." *Glia* 65 (7): 1032–42.
https://doi.org/10.1002/glia.23118.

L., Cen, Carlson B.L., Schroeder M.A., Ostrem J.L., Kitange G.J., Mladek A.C., Fink S.R., et al. 2012.
"P16-Cdk4-Rb Axis Controls Sensitivity to a Cyclin-Dependent Kinase Inhibitor PD0332991
in Glioblastoma Xenograft Cells." *Neuro-Oncology* 14 (7): 870–81.
https://doi.org/10.1093/neuonc/nos114 LK   -
http://vu.on.worldcat.org/atoztitles/link?sid=EMBASE&issn=15228517&id=doi:10.1093%2Fn
euonc%2Fnos114&atitle=P16-Cdk4-Rb+axis+controls+sensitivity+to+a+cyclin-
dependent+kinase+inhibitor+PD0332991+in+glioblastoma+xenograft+cells&stitle=Neuro
-Oncology&title=Neuro-
Oncology&volume=14&issue=7&spage=870&epage=881&aulast=Cen&aufirst=Ling&auin
it=L.&aufull=Cen+L.&coden=NEURJ&isbn=&pages=870-
881&date=2012&auinit1=L&auinitm=.

Lathia, Justin D., Stephen C. Mack, Erin E. Mulkearns-Hubert, Claudia L.L. Valentim, and Jeremy
N. Rich. 2015. "Cancer Stem Cells in Glioblastoma." *Genes & Development* 29 (12): 1203–
17. https://doi.org/10.1101/gad.261982.115.

Laug, Dylan, Stacey M. Glasgow, and Benjamin Deneen. 2018. "A Glial Blueprint for
Gliomagenesis." *Nature Reviews Neuroscience* 19 (7): 393–403.
https://doi.org/10.1038/s41583-018-0014-3.

Lavrik, Inna N. 2010. "Systems Biology of Apoptosis Signaling Networks." *Current Opinion in*
Biotechnology 21 (4): 551–55. https://doi.org/10.1016/j.copbio.2010.07.001.

Lee, Gina, Brenda Auffinger, Donna Guo, Tanwir Hasan, Marc Deheeger, Alex L. Tobias, Jeong
Yeon Kim, et al. 2016. "Dedifferentiation of Glioma Cells to Glioma Stem-like Cells By
Therapeutic Stress-Induced HIF Signaling in the Recurrent GBM Model." *Molecular Cancer*
Therapeutics 15 (12): 3064–76. https://doi.org/10.1158/1535-7163.mct-15-0675.

Lee, Jun Sung, Edlira Hoxha, and Hae Ri Song. 2017. "A Novel NFIA-NFκB Feed-Forward Loop

Contributes to Glioblastoma Cell Survival." *Neuro-Oncology* 19 (4): 424–34.
https://doi.org/10.1093/neuonc/now233.

Lee, Jun Sung, Jiping Xiao, Parita Patel, Jake Schade, Jinhua Wang, Benjamin Deneen, Anat
Erdreich-Epstein, and Hae Ri Song. 2014. "A Novel Tumor-Promoting Role for Nuclear
Factor IA in Glioblastomas Is Mediated through Negative Regulation of P53, P21, and
PAI1." *Neuro-Oncology* 16 (2): 191–203. https://doi.org/10.1093/neuonc/not167.

Lemmon, Mark A., and Joseph Schlessinger. 2010. "Cell Signaling by Receptor Tyrosine Kinases."
Cell 141 (7): 1117–34. https://doi.org/10.1016/j.cell.2010.06.011.

Lord, Nathan D, Thomas M Norman, Ruoshi Yuan, Somenath Bakshi, Richard Losick, and Johan
Paulsson. 2019. "Stochastic Antagonism between Two Proteins Governs a Bacterial Cell
Fate Switch." *Science* 366 (6461): 116–20. https://doi.org/10.1126/science.aaw4506.

Ma, Yian, Qijun Tan, Ruoshi Yuan, Bo Yuan, and Ping Ao. 2014. "Potential Function in a
Continuous Dissipative Chaotic System: Decomposition Scheme and Role of Strange
Attractor." *International Journal of Bifurcation and Chaos* 24 (2).
https://doi.org/10.1142/S0218127414500151.

MacArthur, Ben D. 2023. "Stem Cell Biology Needs a Theory." *Stem Cell Reports* 18 (1): 3–5.
https://doi.org/10.1016/j.stemcr.2022.11.005.

Man, Si Ming, and Thirumala Devi Kanneganti. 2016. "Converging Roles of Caspases in
Inflammasome Activation, Cell Death and Innate Immunity." *Nature Reviews Immunology*
16 (1): 7–21. https://doi.org/10.1038/nri.2015.7.

Manning, Brendan D., and Alex Toker. 2017. "AKT/PKB Signaling: Navigating the Network." *Cell*
169 (3): 381–405. https://doi.org/10.1016/j.cell.2017.04.001.

Marumoto, Tomotoshi, Ayumu Tashiro, Dinorah Friedmann-Morvinski, Miriam Scadeng, Yasushi
Soda, Fred H. Gage, and Inder M. Verma. 2009. "Development of a Novel Mouse Glioma
Model Using Lentiviral Vectors." *Nature Medicine* 15 (1): 110–16.
https://doi.org/10.1038/nm.1863.

Muñoz-Espín, Daniel, and Manuel Serrano. 2014. "Cellular Senescence: From Physiology to
Pathology." *Nature Reviews Molecular Cell Biology* 15 (7): 482–96.
https://doi.org/10.1038/nrm3823.

Newbern, Jason, and Carmen Birchmeier. 2010. "Nrg1/ErbB Signaling Networks in Schwann Cell
Development and Myelination." *Seminars in Cell and Developmental Biology* 21 (9): 922–
28. https://doi.org/10.1016/j.semcdb.2010.08.008.

Niu, Bing, Chaofeng Liang, Yi Lu, Manman Zhao, Qin Chen, Yuhui Zhang, Linfeng Zheng, and
Kuo Chen Chou. 2019. "Glioma Stages Prediction Based on Machine Learning Algorithm
Combined with Protein-Protein Interaction Networks." *Genomics*, no. April.
https://doi.org/10.1016/j.ygeno.2019.05.024.

Nordgren, Ida Karin, and Ali Tavassoli. 2011. "Targeting Tumour Angiogenesis with Small
Molecule Inhibitors of Hypoxia Inducible Factor." *Chemical Society Reviews* 40 (8): 4307–17.
https://doi.org/10.1039/c1cs15032d.

Nurse, Paul. 2021. "Biology Must Generate Ideas as Well as Data." *Nature* 597 (7876): 305–305.
https://doi.org/10.1038/d41586-021-02480-z.

Olariu, Victor, and Carsten Peterson. 2019. "Kinetic Models of Hematopoietic Differentiation."
Wiley Interdisciplinary Reviews: Systems Biology and Medicine 11 (1): e1424.
https://doi.org/10.1002/wsbm.1424.

Otto, Tobias, and Piotr Sicinski. 2017. "Cell Cycle Proteins as Promising Targets in Cancer
Therapy." *Nature Reviews Cancer* 17 (2): 93–115. https://doi.org/10.1038/nrc.2016.138.
Perkins, Neil D. 2012. "The Diverse and Complex Roles of NF-KB Subunits in Cancer." *Nature*
Reviews Cancer 12 (2): 121–32. https://doi.org/10.1038/nrc3204.
Prokop, Aleš. 2022. "Towards the First Principles in Biology and Cancer: New Vistas in
Computational Systems Biology of Cancer." *Life* 12 (1).
https://doi.org/10.3390/life12010021.
Qiu, Xiaojie, Yan Zhang, Jorge D. Martin-Rufino, Chen Weng, Shayan Hosseinzadeh, Dian Yang,
Angela N. Pogson, et al. 2022. "Mapping Transcriptomic Vector Fields of Single Cells." *Cell*
185 (4): 690-711.e45. https://doi.org/10.1016/j.cell.2021.12.045.
Rahman, Maryam, Loic Deleyrolle, Vinata Vedam-Mai, Hassan Azari, Muhammad Abd-El-Barr,
and Brent A. Reynolds. 2011. "The Cancer Stem Cell Hypothesis: Failures and Pitfalls."
Neurosurgery 68 (2): 531–45. https://doi.org/10.1227/NEU.0b013e3181ff9eb5.
Rajasekhar, Vinagolu K, Agnes Viale, Nicholas D Socci, Martin Wiedmann, Xiaoyi Hu, and Eric C
Holland. 2003. "Oncogenic Ras and Akt Signaling Contribute to Glioblastoma Formation by
Differential Recruitment of Existing MRNAs to Polysomes." *Molecular Cell* 12 (4): 889–901.
http://www.ncbi.nlm.nih.gov/pubmed/14580340.
Raju, Archishman, and Eric D. Siggia. 2023. "A Geometrical Perspective on Development."
Development Growth and Differentiation 65 (5): 245–54.
https://doi.org/10.1111/dgd.12855.
Rankin, Erinn B., and Amato J. Giaccia. 2016. "Hypoxic Control of Metastasis." *Science* 352 (6282):
175–80. https://doi.org/10.1126/science.aaf4405.
Reilly, Karlyne M., Dagan A. Loisel, Roderick T. Bronson, Margaret E. McLaughlin, and Tyler Jacks.
2000. "Nf1;Trp53 Mutant Mice Develop Glioblastoma with Evidence of Strain-Specific
Effects." *Nature Genetics* 26 (1): 109–13. https://doi.org/10.1038/79075.
Reiprich, Simone, and Michael Wegner. 2015. "From CNS Stem Cells to Neurons and Glia: Sox for
Everyone." *Cell and Tissue Research* 359 (1): 111–24. https://doi.org/10.1007/s00441-014-
1909-6.
Rockne, Russell C., Sergio Branciamore, Jing Qi, David E. Frankhouser, Denis O'Meally, Wei-Kai
Hua, Guerry Cook, et al. 2020. "State-Transition Analysis of Time-Sequential Gene
Expression Identifies Critical Points That Predict Development of Acute Myeloid Leukemia."
Cancer Research 80 (15): 3157–69. https://doi.org/10.1158/0008-5472.CAN-20-0354.
Rollbrocker, Britta, Andreas Waha, David N. Louis, Otmar D. Wiestler, and Andreas Von Deimling.
1996. "Amplification of the Cyclin-Dependent Kinase 4 (CDK4) Gene Is Associated with
High Cdk4 Protein Levels in Glioblastoma Multiforme." *Acta Neuropathologica* 92 (1): 70–
74. https://doi.org/10.1007/s004010050491.
Salam, Rana, Alexa Saliou, Franck Bielle, Mathilde Bertrand, Christophe Antoniewski, Catherine
Carpentier, Agusti Alentorn, et al. 2023. "Cellular Senescence in Malignant Cells Promotes
Tumor Progression in Mouse and Patient Glioblastoma." *Nature Communications* 14 (1): 1–
21. https://doi.org/10.1038/s41467-023-36124-9.
Samanta, Jayshree, and John A. Kessler. 2004. "Interactions between ID and OLIG Proteins
Mediate the Inhibitory Effects of BMP4 on Oligodendroglial Differentiation." *Development*
131 (17): 4131–42. https://doi.org/10.1242/dev.01273.
Sato, Atsushi, Jun Sunayama, Ken Ichiro Matsuda, Shizuka Seino, Kaori Suzuki, Eriko Watanabe,

Ken Tachibana, Arata Tomiyama, Takamasa Kayama, and Chifumi Kitanaka. 2011. "MEK-ERK Signaling Dictates DNA-Repair Gene MGMT Expression and Temozolomide Resistance of Stem-like Glioblastoma Cells via the MDM2-P53 Axis." *Stem Cells* 29 (12): 1942–51. https://doi.org/10.1002/stem.753.

Sedo, Aleksi, and Rolf Mentlein, eds. 2014. *Glioma Cell Biology*. Vienna: Springer Vienna. https://doi.org/10.1007/978-3-7091-1431-5.

Simon, Matthias, Gertraud Köster, Anil G Menon, and Johannes Schramm. 1999. "Functional Evidence for a Role of Combined CDKN2A (P16-P14 ARF)/ CDKN2B (P15) Gene Inactivation in Malignant Gliomas." *Acta Neuropathologica* 98 (5): 444–52. https://doi.org/10.1007/s004010051107.

Singh, Rumani, Anthony Letai, and Kristopher Sarosiek. 2019. "Regulation of Apoptosis in Health and Disease: The Balancing Act of BCL-2 Family Proteins." *Nature Reviews Molecular Cell Biology* 20 (3): 175–93. https://doi.org/10.1038/s41580-018-0089-8.

Smaldino, Paul. 2019. "Better Methods Can't Make up for Mediocre Theory." *Nature* 575 (7781): 9–9. https://doi.org/10.1038/d41586-019-03350-5.

Spencer, Sabrina L., and Peter K. Sorger. 2011. "Measuring and Modeling Apoptosis in Single Cells." *Cell* 144 (6): 926–39. https://doi.org/10.1016/j.cell.2011.03.002.

Sun, Shao Cong. 2017. "The Non-Canonical NF-KB Pathway in Immunity and Inflammation." *Nature Reviews Immunology* 17 (9): 545–58. https://doi.org/10.1038/nri.2017.52.

Suvà, Mario L., and Itay Tirosh. 2020. "The Glioma Stem Cell Model in the Era of Single-Cell Genomics." *Cancer Cell* 37 (5): 630–36. https://doi.org/10.1016/j.ccell.2020.04.001.

Tanaka, Kazuhiro, Ivan Babic, David Nathanson, David Akhavan, Deliang Guo, Beatrice Gini, Julie Dang, et al. 2011. "Oncogenic EGFR Signaling Activates an MTORC2-NF-KB Pathway That Promotes Chemotherapy Resistance." *Cancer Discovery* 1 (6): 524–38. https://doi.org/10.1158/2159-8290.CD-11-0124.

Taniguchi, Koji, and Michael Karin. 2018. "NF-B, Inflammation, Immunity and Cancer: Coming of Age." *Nature Reviews Immunology* 18 (5): 309–24. https://doi.org/10.1038/nri.2017.142.

Tchieu, Jason, Elizabeth L. Calder, Sudha R. Guttikonda, Eveline M. Gutzwiller, Kelly A. Aromolaran, Julius A. Steinbeck, Peter A. Goldstein, and Lorenz Studer. 2019. "NFIA Is a Gliogenic Switch Enabling Rapid Derivation of Functional Human Astrocytes from Pluripotent Stem Cells." *Nature Biotechnology* 37 (3): 267–75. https://doi.org/10.1038/s41587-019-0035-0.

Thomas, C, G Ely, C D James, R Jenkins, M Kastan, A Jedlicka, P Burger, and R Wharen. 2001. "Glioblastoma-Related Gene Mutations and over-Expression of Functional Epidermal Growth Factor Receptors in SKMG-3 Glioma Cells." *Acta Neuropathol* 101 (6): 605–15. internal-pdf://142.210.225.158/Thomas-2001-Acta Neuropathol.pdf%5Cnhttp://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11515790 http://link.springer.com/content/pdf/10.1007%2Fs004010000332.

Uthamacumaran, Abicumaran. 2022. "Dissecting Cell Fate Dynamics in Pediatric Glioblastoma through the Lens of Complex Systems and Cellular Cybernetics." *Biological Cybernetics* 116 (4): 407–45. https://doi.org/10.1007/s00422-022-00935-8.

Vanhaesebroeck, Bart, Len Stephens, and Phillip Hawkins. 2012. "PI3K Signalling: The Path to Discovery and Understanding." *Nature Reviews Molecular Cell Biology* 13 (3): 195–203.

1025   https://doi.org/10.1038/nrm3290.

1026 Waddington, Conrad Hal. 1957. *The Strategy of the Genes: A Discussion of Some Aspects of*
1027   *Theoretical Biology*. New York: MacMillan Company.

1028 Wade, Mark, Yao Cheng Li, and Geoffrey M. Wahl. 2013. "MDM2, MDMX and P53 in
1029   Oncogenesis and Cancer Therapy." *Nature Reviews Cancer* 13 (2): 83–96.
1030   https://doi.org/10.1038/nrc3430.

1031 Wahlbuhl, Mandy, Simone Reiprich, Michael R Vogl, Michael R. Bösl, and Michael Wegner. 2012.
1032   "Transcription Factor Sox10 Orchestrates Activity of a Neural Crest-Specific Enhancer in the
1033   Vicinity of Its Gene." *Nucleic Acids Research* 40 (1): 88–101.
1034   https://doi.org/10.1093/nar/gkr734.

1035 Wang, Zilai, Daochun Sun, Yu Jung Chen, Xuanhua Xie, Yufeng Shi, Viviane Tabar, Cameron W.
1036   Brennan, et al. 2020. "Cell Lineage-Based Stratification for Glioblastoma." *Cancer Cell* 38
1037   (3): 366-379.e8. https://doi.org/10.1016/j.ccell.2020.06.003.

1038 Weider, Matthias, and Michael Wegner. 2017. "SoxE Factors: Transcriptional Regulators of Neural
1039   Differentiation and Nervous System Development." *Seminars in Cell and Developmental*
1040   *Biology* 63: 35–42. https://doi.org/10.1016/j.semcdb.2016.08.013.

1041 Weinberg, Robert A. 2014. "Coming Full Circle - From Endless Complexity to Simplicity and Back
1042   Again." *Cell* 157 (1): 267–71. https://doi.org/10.1016/j.cell.2014.03.004.

1043 Wiley, Christopher D., and Judith Campisi. 2016. "From Ancient Pathways to Aging Cells -
1044   Connecting Metabolism and Cellular Senescence." *Cell Metabolism* 23 (6): 1013–21.
1045   https://doi.org/10.1016/j.cmet.2016.05.010.

1046 Williams, Evan G, Yibo Wu, Pooja Jha, Sébastien Dubuis, Peter Blattmann, Carmen A. Argmann,
1047   Sander M. Houten, et al. 2016. "Systems Proteomics of Liver Mitochondria Function."
1048   *Science* 352 (6291). https://doi.org/10.1126/science.aad0189.

1049 Wright, Sewall. 1932. "The Roles of Mutation,Inbreeding,Crossbreeding, and Selection in
1050   Evolution." In *Proceedings of the Sixth International Congress of Genetics 1*, edited by D. F.
1051   Jones, 356–366.

1052 Yuan, Ruoshi, Ying Tang, and Ping Ao. 2017. "SDE Decomposition and A-Type Stochastic
1053   Interpretation in Nonequilibrium Processes." *Frontiers of Physics* 12 (6): 120201.
1054   https://doi.org/10.1007/s11467-017-0718-2.

1055 Yuan, Ruoshi, Xinan Wang, Yian Ma, Bo Yuan, and Ping Ao. 2013. "Exploring a Noisy van Der Pol
1056   Type Oscillator with a Stochastic Approach." *Physical Review E - Statistical, Nonlinear, and*
1057   *Soft Matter Physics* 87 (6): 1–7. https://doi.org/10.1103/PhysRevE.87.062109.

1058 Yuan, Ruoshi, Xiaomei Zhu, Gaowei Wang, Site Li, and Ping Ao. 2017. "Cancer as Robust Intrinsic
1059   State Shaped by Evolution: A Key Issues Review." *Reports on Progress in Physics* 80 (4):
1060   042701. https://doi.org/10.1088/1361-6633/aa538e.

1061 Zeng, Tao, Daming Cui, and Liang Gao. 2015. "Glioma: An Overview of Current Classifications,
1062   Characteristics, Molecular Biology and Target Therapies." *Frontiers in Bioscience -*
1063   *Landmark* 20 (7): 1104–15. https://doi.org/10.2741/4362.

1064 Zhang, Qian, Michael J. Lenardo, and David Baltimore. 2017. "30 Years of NF-ĸB: A Blossoming
1065   of Relevance to Human Pathobiology." *Cell* 168 (1–2): 37–57.
1066   https://doi.org/10.1016/j.cell.2016.12.012.

1067 Zheng, Hongwu, Haoqiang Ying, Haiyan Yan, Alec C. Kimmelman, David J. Hiller, An Jou Chen,
1068   Samuel R. Perry, et al. 2008. "P53 and Pten Control Neural and Glioma Stem/Progenitor

Cell Renewal and Differentiation." *Nature* 455 (7216): 1129–33.

https://doi.org/10.1038/nature07443.

Zhu, X.-M., L. Yin, L. Hood, and P. Ao. 2004. "Robustness, Stability and Efficiency of Phage

Lambda Genetic Switch: Dynamical Structure Analysis." *Journal of Bioinformatics and*

Computational Biology 02 (04): 785–817. https://doi.org/10.1142/S0219720004000946.

Zhu, Yuan, Frantz Guignard, Dawen Zhao, Li Liu, Dennis K. Burns, Ralph P. Mason, Albee

Messing, and Luis F. Parada. 2005. "Early Inactivation of P53 Tumor Suppressor Gene

Cooperating with NF1 Loss Induces Malignant Astrocytoma." *Cancer Cell* 8 (2): 119–30.

https://doi.org/10.1016/j.ccr.2005.07.004.

Zmasek, Christian M., and Adam Godzik. 2013. "Evolution of the Animal Apoptosis Network."

Cold Spring Harbor Perspectives in Biology 5 (3).

https://doi.org/10.1101/cshperspect.a008649.

Zong, Hui, Roel G.W. Verhaak, and Peter Canolk. 2012. "The Cellular Origin for Malignant Glioma

and Prospects for Clinical Advancements." *Expert Review of Molecular Diagnostics* 12 (4):

383–94. https://doi.org/10.1586/erm.12.30.

1084

1085

Fig. 1. Endogenous network and network dynamics frameworks for glioma. (A) The core endogenous network for glioma. This network, constructed from experimentally driven causality knowledge, comprises 10 core functional modules or molecular pathways, including the RTK pathway, NF-κB pathway, Ras pathway, Akt pathway, HIF pathway, p53 pathway, cell cycle, cellular senescence, apoptosis, and glia differentiation. A closed a 25-node network with 75 edges formed by selecting key agents and interactions from modules and pathways, which includes 44 activatory interactions (indicated with red arrows) and 31 inhibitory interactions (indicated with blue T-shaped lines). **(B)** From the network to the dynamics. Two dynamic frameworks, Boolean dynamics and ordinary differential equations (ODE), were applied to this network. The dynamics equations for Akt are presented as an example. Here, x denotes the concentration/activity of core nodes, h represents the Hill coefficient, and k is the inverse of the apparent dissociation constant. A normalized degradation rate of 1 is used in this model, implying a degradation amount of -x per iteration. The full sets of equations are available in Supplementary Table 1.

Fig. 2. Dynamics results. (A) Attractor domains obtained using Boolean dynamics, with a random initial vector number of 1e8. States are represented with red for the active (1) and cyan for the inhibition (0). The x-axis corresponds to the identifier of each attractor domain, and the y-axis lists the core nodes. **(B) (C)** Fixed points obtained under the ODE framework. Fixed points, derived using ODE with a random initial vector number of 1e7, are hierarchically clustered by columns. Colors indicate the state, with red for active (1) and cyan for inhibition (0). The x-axis lists the identifier of each attractor domain, and the y-axis displays the core nodes. (B) Depicts stable states (stable fixed points). (C) Depicts transition states (unstable fixed points). **(D)** A comparison of the number of stable states obtained using different ODE algorithms. The count of stable states reached using Newton's method (fsolve in MATLAB) and Euler's method across various random initial vectors is plotted. The x-axis represents the count of random initial vectors, and the y-axis shows the number of stable states, with blue indicating results from Newton's method and red from Euler's method. Parameters used were h=3, k=10. **(E)** Comparison of the count of stable and transition states obtained under different parameters. The x-axis represents the parameters used, and the y-axis the counted number of fixed points. These results obtained under 1e7 random initial vectors and the Newton's method. **(F)** Comparison between Boolean dynamics and ODE. After averaging the linear attractors from Boolean dynamics, they are hierarchically clustered by column alongside fixed points obtained from ODE. The x-axis denotes the identifier of the dynamic results, and the y-axis the core nodes. Grid colors indicate computation results, with red for active (1) and cyan for inhibition (0). Annotations above columns signify the computational framework, with red for ODE and cyan for Boolean dynamics.

Fig. 3. Schematic illustration: Using the topology structure of fixed points to depict the potential energy landscape. (A) A three-dimensional potential energy landscape schematic. The x-y plane represents a two-dimensional phase space, and the z-axis is potential energy. The "mountainous" surface depicts the potential energy surface, akin to Waddington's epigenetic landscape used to describe development. White curves mark intersections between two orthogonal planes and the potential energy surface. Small dots on the curves

represent fixed points, which will be further exemplified. **(B)** A two-dimensional projection of the potential energy landscape. White curves from (A) are mapped onto (B). (UP) indicates mapping of the y-z plane curve, while (DOWN) corresponds to the x-z plane curve. Small dots represent fixed points, corresponding one-to-one with those in (A). **(C)** A topological network mapping of the potential energy landscape. Small dots denote fixed points, correlating one-to-one with those in (A and B). Arrows point from higher energy fixed points towards lower energy ones. Red dots and cyan dots both signify fixed points, with red for unstable fixed points—those having at least one positive real part in the eigenvalues of the Jacobian matrix—and cyan for stable fixed points—those with all negative eigenvalues.

Fig. 4. The global potential energy landscape of glia. The network topology of fixed points illustrates the global potential energy landscape for the glial system. Each node signifies a fixed point; large nodes indicate stable fixed points (stable states), and small nodes represent unstable fixed points (transition states). Node colors code for differentiation lineage, with yellow for astrocytes, blue for oligodendrocytes, and gray for glia stem cells or non-glia. Node shapes signal the apoptotic module state, with circles for apoptosis inhibition, octagons for balanced activation and inhibition of apoptosis, and triangles for apoptosis activation. Node labels in font color indicate the cell cycle module, with red for proliferation activation and black denoting proliferation inhibition. Arrows point in the direction of lower energy. Spatial relationships between nodes do not reflect relative energy levels.

Fig. 5. Comparisons between theory and experiment. (A) (B) Theoretical results compared with experimental data at the functional modules and signaling pathways level. (A) Knowledge about glioma summarized from the literature at the level of functional modules and pathways. (B) Theoretical calculations at the module level (ODE, Newton's method, h=3, k=10, 1e7, same below). The far-left of the clustering tree displays glioma candidate states. **(C) (D)** Theoretical results compared with low-throughput data. (C) (LEFT) Core node-associated gene expression/activity data compiled from the literature, with columns representing individual reports. 1 denotes activation, 0 denotes inhibition, and na indicates the gene was not covered in that report. (BOTTOM-RIGHT) Typical states of transcription factors regulating glial cell differentiation summarized from the literature. (D) 1st Neighbouring States of glioma candidate states as determined by theoretical calculations. **(E) (F)** Theoretical results compared with high-throughput experimental data. (E) (LEFT) Expression profiles of core node gene families from GSE151352, presented as fold changes between tumor and adjacent non-tumorous samples, normalized to the [0, 1]. (RIGHT) Theoretical calculation results for glioma candidate states. (F) Comparison of glioma candidate states with GTEx-TCGA expression profiles. TPM data from the cortex in GTEx and GBM and LGG samples in TCGA, with 150 randomly selected from each, normalized to the [0, 1] and plotted for the distribution of core node-associated gene expression. The x-axis represents normalized expression levels, and the y-axis shows distribution density; vertical lines indicate the level of that core node in glioma candidate states.

Fig. 6. Landscape near glioma. (A) Locating the landscape near glioma within the global landscape of the glial system. The transparent overlay represents the global landscape of the

glial system, with the highlighted red network indicating the landscape near glioma. **(B)** A two-dimensional representation of the 1st Neighbouring States. Blue curves sketch the potential energy surface, with red and blue nodes corresponding to unstable or stable fixed points, respectively. Transition states connecting glioma stable states and those transition states connected to stable states constitute the landscape near glioma. **(C)** The energy landscape near glioma. The 1st Neighbouring States of glioma, rearranged according to differentiation type and connections to glioma states, yield the landscape near glioma. **(D)** Profile of the energy landscape near glioma. Activities of core genes in all fixed points within the landscape near glioma are hierarchically clustered by column. The x-axis lists fixed point identifiers, and the y-axis core genes. Red signifies active (1), and cyan inhibition (0).

Fig. 7. Transition states within the landscape near glioma. (A) Theoretical versus experimental knowledge comparison for GSC states. **(B)** The GSC state is a transition state connecting exclusively to glioma stable states. (UP) Blue curves sketch the potential energy surface, with red and blue nodes corresponding to unstable or stable fixed points, respectively, same below. (DOWN) The connection of GCS state and glioma stable states. large nodes indicate stable fixed points (stable states), and small nodes represent unstable fixed points (transition states). Node colors code for differentiation lineage, with yellow for astrocytes, blue for oligodendrocytes, and gray for glia stem cells or non-glia. Node shapes signal the apoptotic module state, with circles for apoptosis inhibition, octagons for balanced activation and inhibition of apoptosis, and triangles for apoptosis activation. Node labels in font color indicate the cell cycle module, with red for proliferation activation and black denoting proliferation inhibition. Arrows point in the direction of lower energy. Spatial relationships between nodes do not reflect relative energy levels. Same below. **(C)** Theoretical calculation results of glioma states. The x-axis lists the core nodes, and the y-axis corresponds to the identifier of each attractor domain. Same below. **(D)** Schematic of potential cure transition states. **(E)** Profile of potential cure transition states. **(F)** Potential cure transition states in the landscape near glioma. **(G)** Schematic of heterogeneity-inducing transition states. **(H)** Profile of heterogeneity-inducing transition states. **(I)** Heterogeneity-inducing transition states in the landscape near glioma.

A

NF-kB
Pathway

HIF
Pathway

RTK
Pathway

Ras
Pathway

Cell Cycle

EGFR

Ras

CDK4

Rb

P21

NF-kB

IkB

HIF

Akt Pathway

PTEN

Akt

Cellular
Senescence

CDKN2A

Sox9

Pros

MDM2

P53

Olig

Sox10

Notch

NFIA

BAD

CASP9

CASP3

XIAP

Bcl2

BAX

Neuroglia Differentiation

P53 Pathway

Apoptosis

MODULES &
PATHWAYS:

•RTK Pathway

•NF-kB Pathway

•Ras Pathway

•Akt Pathway

•HIF Pathway

•Cell Cycle

•P53 Pathway

•Cellular
  Senescence

•Apoptosis

•Differentiation

B

Akt

Ras

CDKN2A

PTEN

EGFR

Boolean dynamics:

$$x(Akt)_{t+1} = (x(EGFR)_t \ OR \ x(Ras)_t) \ NOT(x(PTEN)_t \ OR \ x(CDKN2A)_t)$$

Differential equation:

$$\frac{dx(Akt)}{dt} = -x(Akt) + \frac{k*(x(EGFR)^h + x(Ras)^h)}{1+k*(x(EGFR)^h + x(Ras)^h)}$$

$$*\frac{1}{1+k*(x(PTEN)^h + x(CDKN2A)^h)}$$

A Boolean Dynamics Results
(1e8, Spread Linear Attractors)

B ODE Results (1e7, Stable States)

C ODE Results (1e7, Transition States)

D

E

F Boolean Dynamics & ODE, Comparison

A — Potential energy landscape (showed in 3-dimension)

B — Potential energy landscape (sections showed in 2-dimension)

C — Topology structure of fixed points depicts main feature of potential energy landscape

Fixed Points
$$\frac{\mathrm{d}\boldsymbol{x}}{\mathrm{d}t} = 0$$

Transition State
$$\exists \mathrm{Re}\{\lambda_i[\boldsymbol{J}(\boldsymbol{x})]\} > 0$$
$$i = 1, 2, 3, \ldots, n$$

Stable State
$$\forall \mathrm{Re}\{\lambda_i[\boldsymbol{J}(\boldsymbol{x})]\} < 0$$
$$i = 1, 2, 3, \ldots, n$$

A

B

The Potential Energy
Landscape near Glioma

C

Landscape near Glioma

D

Glioma 1st
Neighbouring
States

A

Functional
characteristics of GSCs:

· Being glioma

· Self-renewable

· Giving rise to
differentiated progeny

Features of **tra_6**:

· Being glioma

· High stemness
molecularly

· Evolving to two
differentiated states
after small
perturbation

D

Potential Cure
Transition State

Treatment

Glioma   Apoptosis

Apoptosis   Senescence

G

Heterogeneity
Inducing
Transition State

Treatment

Glioma   Glioma

Apoptosis