Enhanced reproductive outcomes under worst-case climate change scenarios in Red-sided

Garter snakes (*Thamnophis sirtalis parietalis*)

David L Hubert[1,*], Ehren J Bentz[2], and Robert T Mason[1]

[1] Department of Integrative Biology, Oregon State University, Corvallis, Oregon. USA, 97333

[2] Department of Psychology, Cornell University, Ithaca, New York, USA

* Corresponding author: E-mail: hubertd@oregonstate.edu

ABSTRACT

As global temperatures continue to increase, understanding the impacts of warming

environments is increasingly relevant. Temperature is especially relevant for ectothermic

organisms which depend upon consistent and predictable annual temperature cycles for

reproduction and development. However, additional research is required in this area to elucidate

the potential long-term impacts of climate change. To understand how projected increases in

environmental temperatures may impact reproductive outcomes within natural populations of

ectothermic vertebrates, we manipulated ambient temperatures during gestation in Red-sided

garter snakes (*Thamnophis sirtalis parietalis*). Wild snakes were collected in the Interlake region

of Manitoba, Canada during their spring mating season and allowed to mate in controlled

conditions. For the duration of gestation, mated females were placed into one of two ambient

thermal conditions: temperatures emulating those found in the species' natural habitat or

temperatures with a consistent 5 °C increase to match end-of-century climate change projections.

We recorded observations for each litter and all neonates resulting from controlled mating trials.

We observed no difference in litter sizes or birth rates between thermal conditions. Interestingly,

we observed both a significant reduction in gestation length and a significant increase in neonate

body size associated with increased ambient temperatures. These results suggest that increased

temperatures during gestation may confer reproductive benefits for the northern populations of

this species even under the most extreme current modeled warming predictions. We discuss the

broader implications of this effect, including possible negative ecological outcomes.

1. Introduction

In a time of rapidly changing environmental temperatures, it is increasingly important to

elucidate how altered thermal conditions impact the coordination of important life history stages

and essential events such as mating, gestation, and foraging. Because reproductive success is a

defining feature of individual fitness, factors that greatly impact reproduction among individuals

within a population have the potential to dramatically affect the long-term success and survival

of that population (Brommer et al, 2002; Sherman and Runge, 2002; Rabaiotti et al., 2023).

Ectotherms are influenced particularly strongly by environmental conditions, especially

temperature, which is frequently an important cue used to coordinate behavior and physiological

processes essential to reproductive success including courtship behavior and successful

incubation or gestation (Huey and Stevenson, 1979; Huey, 1982; Miles et al., 2000; Lourdias et

al., 2013). Ectotherms from terrestrial ecosystems demonstrate existing plastic physiological and

behavioral mechanisms to maintain homeostasis during large seasonal and acute temperature

fluctuations (Huey et al., 2001; Gunderson et al., 2017; Noer et al., 2022). These extreme

temperature fluctuations have been increasingly exacerbated by recent climate change (Masson-

Delmotte et al., 2021). Therefore, investigating how terrestrial ectotherms respond to these new

conditions provides an excellent opportunity to explore the broader impacts of a warming

climate.

The challenge of maintaining ideal thermal conditions during embryonic development is

central to reproductive success among many ectotherms, though overcoming this challenge is

accomplished using different strategies. Oviparous (egg laying) ectotherms must ensure

consistent thermal conditions for their eggs throughout the entire period of embryonic

development by utilizing complex behaviors such as careful nest building, precise nest site

selection, or by maintaining thermal conditions through monitored incubation of clutches (Du

and Shine, 2014). Typically, these nest sites are abandoned by the parent shortly after egg

deposition, allowing the nest itself to maintain environmental conditions conducive to proper

embryonic development while the mother is free to forage and rebuild depleted energy reserves

(Shine, 1988; Gans et al., 1996; Huang and Pike, 2013; Doody and Refsnider, 2022). In contrast,

viviparous (live-bearing) ectotherms must alter their behavior while gravid to maintain optimal

internal thermal conditions throughout gestation, even to the extent of risking their own survival

to prioritize ideal thermal conditions for developing embryos (Seigel et al., 1987; Charland and

Gregory, 1995; Gregory et al., 1999; Lorioux et al., 2013). This behavioral tradeoff serves to

illustrate the importance of an organism's thermal environment, and that maintaining proper

temperature is vital for the survival and optimal development of offspring and therefore directly

impacts the individual fitness of reproducing adults. Temperatures experienced during gestation

are often observed to impact neonate size in viviparous squamates (Mathies and Andrews, 1997;

Lourdias et al., 2004; Gao et al., 2010; King, 2022). Both greater neonate body length (Brown

and Shine, 2004; Kissner and Weatherhead, 2005; Addis et al., 2017) and greater body mass

(Jayne and Bennett, 1990; Bronikowski, 2000; Kissner and Weatherhead, 2005) are associated

with significantly increased rates of survival to adulthood in viviparous snakes. Here, we use the

term "body condition" to refer specifically to the ratio of body mass to snout-vent length (SVL),

which is frequently used as a proximate measure for the nutritional state and overall health of an

individual.

Changes to environmental thermal conditions during gestation have been observed to

impact both physiology and behavior of viviparous ectotherms to alter characteristics such as

maternal energy use and basking and foraging behavior (Lourdias et al., 2002; Ladyman et al.,

2003; Wang et al., 2017), gestation length (Blanchard and Blanchard, 1940; Ji et al., 2006; Wang

et al., 2017; King, 2022), litter size (Lourdias et al., 2004; Zhang et al., 2010; Tang et al., 2012),

and body condition and viability of neonates (Lourdias et al., 2004; Wang et al., 2017; King,

2022). Although there is often agreement regarding the influence of warmer temperatures on

gestation, neonate survival, and body condition, the specific effects observed differ between

studies and depend on the biological system examined. Wang et al. (2017) reported that

increased ambient temperature was associated with significantly shorter gestation length,

increased neonate mass, and a decrease in general activity during gestation for Toad-headed

lizards (*Phrynocephalus putajatia*). Beuchat (1988), found that increased environmental

temperatures during gestation were associated with a decrease in gestation length in Yarrow's

spiny lizards (*Sceloporus jarrovi*), but this study also reported that sustained increased

temperatures resulted in a detrimental effect on the body conditions of both mother and neonate.

This latter general trend was further supported, as Tang et al. (2012) reported an increase in litter

size and a decrease in gestation length associated with higher temperatures for Multi-ocellated

racerunner lizards (*Eremias multiocellata*), but that these potentially beneficial effects were lost

as temperatures continued to increase. Lourdias et al. (2004) demonstrated that increased

ambient temperature resulted in shorter gestation length, increased embryo viability, and caused

phenotypic changes, including changes to scale patterns, in neonate European asps (*Vipera aspis*)

within a population living at the northern boundary of its range. Each of these studies

demonstrate that, even among animals that are able to effectively behaviorally thermoregulate,

increased ambient temperatures significantly impact both the behavior of gravid females and the

resulting reproductive outcomes in viviparous species.

Red-sided garter snakes (*Thamnophis sirtalis parietalis)* are viviparous ectotherms and

one of the most northerly living squamate species in North America (Aleksuik and Stewart,

1971). The focal population of this study is found in the Interlake region of Manitoba, Canada

near the northern extreme of the species' range. The Interlake region is particularly well suited

for studies examining the effects of temperature on ectothermic vertebrates, as it is characterized

by major seasonal changes to the thermal landscape. Unsurprisingly, ectotherms living within

this region are well adapted to this extreme environment and demonstrate highly coordinated

annual behavioral cycles primarily driven by seasonal temperature changes (Aleksiuk and

Stewart, 1971; Gregory, 1974; Lutterschmidt et al., 2006). Indeed, Red-sided garter snakes spend

the majority of their life in a hibernation-like state of winter dormancy known as "brumation,"

during which the entire population retreats into subterranean hibernacula for up to 8 months each

year (Aleksiuk and Stewart, 1971; Costanzo, 1985). Brumation is a behavioral and physiological

state characterized by extremely low levels of activity and greatly reduced metabolic rates in

ectotherms. This state is driven by low temperatures and serves as an essential strategy for

surviving extremely cold and resource-limited winter conditions (Gregory, 1982; Geiser, 2013).

Red-sided garter snakes remain relatively inactive throughout brumation and do not feed for

approximately 8-9 months each year (O'Donnell, 2004). Thus, they must rely solely upon stored

energy to survive winter dormancy as well as the intense and energetically costly activities of

mating and migration immediately following emergence from the hibernacula in early spring

(Aleksiuk and Gregory, 1974; Aleksiuk, 1976). These energy stores must be obtained during an

extremely brief summer feeding season lasting as little as 10 weeks, during which an individual

must acquire all of the energy necessary for an entire year of growth, development, reproduction,

and survival (Crews et al., 1987; O'Donnell et al., 2004). Moreover, the challenge of building

energy reserves becomes even more difficult for reproducing females as this brief feeding period

coincides with gestation – a period that is both behaviorally and energetically demanding. Thus,

this limited window of time is critically important and has the potential to dramatically impact

individual fitness, as a reproducing female must carefully balance activities and energy demands

to simultaneously ensure survival while maximizing reproductive success.

Although climate change-induced warming has resulted in a global average temperature

increase of approximately 0.6 ºC over the last 60 years, the Interlake region has been

disproportionately impacted, experiencing nearly triple the global rate with recorded

temperatures showing an increase of 1.7 ºC over the same period (Zhang et al., 2000; Houghton

et al., 2001; Environment Canada, 2014; Bush and Lemmen, 2019). Global temperatures are

predicted to continue to increase by as much as 5.3 ºC by the end of the century, which is well

beyond the findings of previous work where increases of as little as 2 ºC significantly impacted

gestation in viviparous ectotherms (Houghton et al., 2001; Kingsolver et al., 2013; Wang et al.,

2017; Masson-Delmotte et al., 2021). With current regional warming levels so much higher than

the global average, it is reasonably likely that future regional warming will be at the higher end

of global climate change model projections.

Currently, the daily mean environmental temperatures experienced by Red-sided garter

snakes during the summer feeding and gestation period range from 13.8 ⁰C in late September to

20.1 ⁰C in July, and mean maximum daily temperatures reach 25 ⁰C (https://en.climate-

data.org/north-america/canada/manitoba/inwood-108217/). Previous research focused on the

Red-sided garter snakes and other members of the genus *Thamnophis* have shown that both free-

ranging and captive gravid females behaviorally maintain daytime body temperatures between

26 ⁰C and 31 ⁰C – a range that typically exceeds ambient temperatures (Gibson and Falls, 1979;

Arnold and Peterson, 2002; Hughes et al., 2019).

In the present study, we manipulated the thermal environment of gestating female Red-

sided garter snakes by increasing ambient temperatures by an ecologically relevant 5 ⁰C and

observed the impact of this increase on gestation length and neonate size and body condition. We

predicted an increased ambient temperature of 5 ⁰C would significantly reduce gestation length.

However, we also predicted that this thermal increase would exceed beneficial thresholds and

individual gains would be lost resulting in a reduction in neonate body condition or a reduction

of the number of offspring per litter (Beuchat, 1988; Tang et al., 2012).

2. Materials and methods

All experimental protocols presented were approved by the Oregon State University

Institutional Animal Care and Use Committee (Animal Care and Use Protocol 5058), in

accordance with the Association for the Assessment and Accreditation of Laboratory Animal

Care. This research was conducted under the authority of Manitoba Wildlife Scientific Permit

No. WB20333.

3. *Animal Collection*

We collected free-ranging male and female Red-sided garter snakes (male *n* = 80, female

n = 80) from the natural population at over-wintering hibernacula near the town of Inwood,

Manitoba, Canada during the spring mating season between 1 May and 4 May 2018. This site

contains several communal hibernacula that collectively shelter a population estimated at

approximately 35,000 snakes (Shine et al., 2006). Male snakes were selected and captured after

being observed actively exhibiting courtship behavior. Non-mated females were selected and

captured only if their initial emergence from the hibernaculum was observed, and they were

being actively courted by males. To minimize the effects of body size variation on litter size, all

females selected for this study were approximately the same size, with SVLs and body masses of

approx. 60 cm and 100 g, respectively. To confirm that the females had not yet mated that

season, we visually inspected their cloaca for copulatory plugs upon collection. Copulatory plugs

are large, gelatinous spermatophores deposited by a male inside of a female's cloaca during

copulation. Copulatory plugs are substantial in size, persist for several days, and represent an

easily identifiable and reliable indication that a female has recently mated (Whittier et al., 1985;

Friesen et al., 2013). Any female presenting a copulatory plug, or any indication of potentially

having mated, was immediately released. All animals were collected by hand and placed in cloth

bags for transportation to the Chatfield Research Station, Chatfield, Manitoba. Snakes were

separated by sex and housed outdoors in specially designed 1 m^3 nylon enclosures containing

natural stones, hide boxes, and water dishes. Water was provided *ad libitum* for the duration of

captivity, but food was not provided in the field or at the research station, as this species is

aphageous during the spring mating season (O'Donnell et al., 2004, Crews et al., 1987). Capture,

handling, and transportation are not likely to have impacted results, as previous studies have

consistently shown that these activities do not adversely affect survival, reproduction, or

courtship behavior (Friesen et al., 2013; Friesen et al., 2014).

4. Mating Trials

To control mating conditions and timing, we used courtship arenas (opaque, cylindrical,

nylon enclosures 0.5 m in diameter and 1 m tall) with heat lamps (above) and electric heating

pads (below) to maintain homogenous thermal environments at ≥ 13.0 ºC – optimal temperatures

to stimulate vigorous courtship behavior. Female snakes were placed individually into arenas and

paired with 5 randomly selected males. All snakes were allowed to freely behave under constant

observation until successful copulation occurred. Successful copulation was confirmed by visual

inspection and identification of a copulatory plug. Females with copulatory plugs were separated

from the males and transferred to a 1 m^3 outdoor nylon enclosure containing only mated females.

Each male was allowed to mate only once. Males that successfully mated were removed from the

remainder of the study to minimize any potential bias that may arise from allowing a male to sire

more than a single litter of offspring. All males were marked dorsally with indelible ink to

prevent recapture and released at the site of capture within 5 days of collection. All controlled

matings took place within the same 48-hour period.

4.1 Experimental Conditions and Data Collection

We transported the mated females ($n = 50$) by car to the laboratory facilities at Oregon

State University. Animals were transported in cloth bags, housed in a thermally controlled

environment (air temperatures between 20 ºC and 22 ºC), and given water via a damp sponge to

prevent dehydration for the duration of the trip (< 30 h). Females were sorted by size and

systematically assigned to one of two groups in order to ensure that each group contained snakes

with an equivalent distribution of body sizes. Each group was then randomly assigned to

different thermal regimes representing treatment and control conditions. Once assigned to a

group, and for the remainder of the study, snakes were housed within microprocessor-controlled

environmental chambers (Environmental Growth Chambers, Chagrin Falls, OH). The control

group was housed within a chamber emulating the thermal environment found in this species'

natural range during the summer feeding period, which consists of a daytime temperature of 22

⁰C and a nighttime temperature of 15 ⁰C. The treatment group was housed in an identical

chamber set to maintain ambient temperatures increased by 5 ⁰C, which consists of a daytime

temperature of 27 ⁰C and a nighttime temperature of 20 ⁰C. Females were housed individually in

45 liter glass aquaria containing approximately 2.5cm of CareFresh™ paper fiber substrate, a

water bowl, a hide box, a rock to facilitate shedding of skin, and a heat lamp situated at one end

to allow behavioral thermoregulation within the constraints of the experiment. Photoperiods were

identical for both treatments and emulated that of the summer feeding period in Manitoba (16 h:8

h light:dark cycle). Basking lamps provided additional heat for 4 hours per day in the center of

the daylight photoperiod. Humidity was maintained at 65 % for both treatments. Maximum

temperatures under the basking lamps were approximately 31 ⁰C for control group and 40 ⁰C for

the treatment group. Water was provided *ad libitum*, and food (a mixture of earthworms and

salmon fry supplemented with calcium) was provided once per week, during which time all

snakes were allowed to eat to satiation. Snakes were maintained in these conditions for the

duration of the experiment.

Each female was inspected daily to identify births, and offspring were collected within no

more than 24 hours of birth. Dates of birth and number of offspring were recorded for each litter.

Neonates were weighed and measured immediately after collection. Masses were recorded to the

nearest 0.01 g and SVLs were recorded to the nearest 1 mm. Body mass was divided by SVL to

calculate a body condition index (BCI; g/cm) for each neonate. BCI is frequently used as a

measure of approximate overall health and nutritional state as it accounts for both body mass and

length for the same individual, allowing for a better description of the individual than comparing

those values separately (Madsen and Shine, 2002). Measurements were collected for every litter

and for every neonate until the last female gave birth in late September. All adult snakes were

maintained in captivity until the following spring then returned to the location of capture.

Offspring were maintained in the lab for further data collection facilitating additional studies

aiming to characterize growth and development.

5. Results

Gestation duration, litter size, neonate SVL, neonate mass, and neonate body condition

index (BCI) were all checked for normal distributions using the Shapiro-Wilk Normality Test,

and all were found to be significantly different from a normal distribution ($P < 0.005$ for all

measures). Because the data were not normally distributed, we used a non-parametric two-

sample Wilcoxon rank sum test for each measure to test for significant differences between the

treatment and control conditions. The median values for gestation length and neonate mass, SVL,

BCI, were all found to be significantly different between groups (Figure 1, Supplementary Table

1). Median mass, SVL and BCI for neonates in the 27 ºC treatment group were all significantly

247 greater than those in the 22 ℃ control group ($P < 0.001$ for all measures) (Figure 1: D-F). The

248 median gestation length was significantly shorter for the 27 ℃ treatment group compared to the

249 median for the 22 ℃ control group ($P < 0.001$) (Figure 1: A). Litter size was not significantly

250 different between thermal conditions ($P > 0.74$) (Figure 1: E). Additionally, there was no

251 significant difference in birth rates measured as the proportion of females that successfully gave

252 birth divided by the total number of females within each group (two proportions z-test $P > 0.77$)

253 (Figure 1: F).

254 **6. Discussion**

255   Our observations of gestation times accurately agreed with our initial prediction that a 5

256 ℃ increase in temperature during gestation would result in a reduction in gestation time. Indeed,

257 we found that gestation lengths were significantly reduced in the 27 ℃ treatment conditions,

258 even to the extent that the resulting ranges of birth dates did not overlap between the treatment

259 and control groups. The median gestation length for the 27 ℃ treatment was 21 days shorter than

260 that of the 22 ℃ control group representing a 17.8 % decrease in median gestation length.

261 Shorter gestation lengths may increase the female's survival as gravidity imposes energetic costs

262 as well as reduced locomotory and foraging capabilities and requires increased amounts of time

263 engaged in active thermoregulation – all of which may ultimately result in an increased exposure

264 and risk of predation for the gravid female (Seigel et al., 1987; Charland and Gregory, 1995;

265 Gregory et al., 1999). As these costs in Red-sided garter snakes are imposed simultaneously with

266 the already extremely limited opportunities for summer foraging, a reduction in gestation length

267 would allow for both increased amount of time to forage each year and more frequent prey

268 capture during this period ultimately resulting in increased opportunity to replenish depleted

269 energy stores after giving birth. Because the summer feeding period is so limited, even small

increases in annual energy intake may have a large impact facilitating improved winter survival

and potentially a shorter interlude between reproductive bouts, thus increasing overall fecundity

(Aleksiuk and Stewart, 1971; Costanzo, 1985).

Although we observed no differences in either birth rates or average litter sizes between

treatments ($P > 0.77$ and $P > 0.74$ respectively), we did observe a significant difference in the

overall size and body condition of neonates. SVL and mass were both significantly greater for

the 27 ⁰C treatment group compared to the 22 ⁰C control group ($P < 0.001$ for both), and this

difference remained significant when combined to calculate individual BCI values ($P < 0.001$).

Surprisingly, a 5 ⁰C increase ambient temperature both significantly decreased gestation length

and, contrary to our initial predictions, significantly increased neonate size and body condition.

Because larger size at birth is associated with both improved short term and lifetime survival in

snakes, offspring gestated in warmer conditions will likely have increased overall fitness

resulting in better parental reproductive outcomes (Bronikowski, 2000; Brown and Shine, 2004;

Kissner and Weatherhead, 2005; Addis et al., 2017). However, this increase in ambient

temperature had no impact on successful birth rate or the number of offspring in each litter. The

observed decrease in gestation length is consistent with those previously observed among a

number of other viviparous ectotherms (Blanchard and Blanchard, 1940; Ji et al., 2006; Wang et

al., 2017; King, 2022).

Though not observed during this study, a common observation resulting from a moderate

increase in ambient temperature is an increase in average litter size (Lourdias et al., 2004; Zhang

et al., 2010; Tang et al., 2012). However, larger increases in ambient temperature have been

shown to have a detrimental impact on the body conditions of the neonates, as well as potentially

reversing any potentially positive impact caused by a moderate temperature increase (Beuchat,

1988; Yan et al., 2011; Tang et al., 2012). The 5 ⁰C increase in ambient temperature treatment

used in the present study did not result in any observed detrimental impact. This suggests that a 5

⁰C increase in ambient temperature is below the threshold at which any detrimental effects would

begin to arise within this population. This further suggests that there may be additional potential

for increased beneficial effects as temperatures increase in excess of 5 ⁰C, but before

temperatures reach this critical, but unknown, threshold. Further research exploring a wider

range of thermal conditions during gestation would be interesting, as this would expand our

perspective allowing for increased ability to predict the effects of future increases in ambient

temperatures and would potentially pinpoint the maximum ambient temperature threshold at

which beneficial results began to reverse and detrimental impacts begin to arise.

This study did not identify that critical threshold, but these results suggest interesting

implications for future warming associated with continued climate change. The response

variables examined here were limited and did not directly address or account for other

phenotypic affects that may have arisen from increased temperatures during gestation. While it

has been well documented in oviparous ectotherms that increased thermal conditions can result

in significant changes to sex ratios among offspring (Hawkes et al., 2007; Holleley et al., 2015;

Noble et al., 2018), temperature likely plays little to no role in sex determination in viviparous

snakes, such as Red-sided garter snakes, where sex is determined chromosomally rather than

being driven by environmental temperatures (Dunlap and Lang, 1990; Matsubara et al., 2006;

Viana et al., 2020; reviewed in Singchat et al., 2020). However, evidence suggests that many

other developmental processes may be influenced by altered thermal conditions, and increased

temperatures may result in changes to offspring phenotype such as differences in scale and bone

development (Arnold and Peterson, 2002; Lourdias et al., 2004). Additionally, the impacts of

temperature during gestation on offspring traits such as the aforementioned changes to

morphology, body size, and body condition are frequently associated with long-lasting impacts

on survival after birth (Bronikowski, 2000; Brown and Shine, 2004; Kissner and Weatherhead,

2005; Noble et al., 2018).

Current climate change models predict future global average temperature increases of up

to 5.3 ºC by the end of the century (Houghton et al., 2001; Masson-Delmotte et al., 2021). Thus,

the results of this experiment show that an increase in average temperature of this degree,

specifically during the brief yet crucial opportunity for feeding and gestation, is not likely to

cause direct detrimental effects on the reproductive success of northern populations of Red-sided

garter snakes but will potentially result in improved neonate fitness. However, the extended

impacts of increasing temperature are likely to result in other adverse effects on this population.

Thermal stress caused by exposure to increased temperature extremes (acute high or low

temperatures) is known to result in a number of detrimental effects including misaligned

behavioral timing, altered development, reduced or altered activity, and higher mortality rates

(Houghton et al., 2001; Jentsch et al., 2009; Paaijmans et al., 2013; Stott, 2016). The greater

thermal instability associated with climate change conditions may also lead to thermally induced

alterations to energy use and metabolic rate (Friesen et al., 2015; Wilson, 2020; Holden et al.,

2021). The combination of physiological and environmental changes due to increased

environmental temperatures are also known to increase pathogen exposure and increase

individual susceptibility to infection (Walther et al., 2001; Harvell et al., 2009; Gallana et al.,

2013; Nnadi and Carter, 2021). Additionally, many species have been observed to alter seasonal

migration patterns and shift territorial ranges to mitigate changes to their traditional ecosystems,

ultimately resulting in disruption of ecosystems arising from novel interspecific competition and

alterations to the availability of prey species or trophic effects (Parmesan, 2006; Gilman et al.,

2010; Urban et al., 2012; Salinas-Ramos et al., 2020). Although we observed effects of increased

temperatures that appeared likely beneficial to offspring, it is possible that that the lifetime

reproductive potential of mothers would in fact be reduced as a result of long-term exposure to

increased temperatures or repeated bouts of rapid gestation. Thus, future research regarding

changes to ambient temperatures during gestation may benefit from examining both the

immediate and long-term health impacts of mothers, such as impacts on body condition, future

fecundity, parasite load, and subsequent winter survival, in addition to the long-term survival and

impacts on the physiology and long-term reproductive potential of offspring as well.

Higher temperatures caused by future warming have the potential to change the shape of

this population, resulting in offspring that not only develop faster, but are born larger. In the

present study, we observed no apparent tradeoff or cost to the overall number of offspring in

each litter. This has the potential to dramatically increase the survival rates of future generations,

while increasing the time each year spent foraging for both the mothers and the resulting

offspring (Seigel et al., 1987; Charland and Gregory, 1995; Brown and Shine, 2004; Kissner and

Weatherhead, 2005). This species already has extremely large population sizes, with this den site

alone numbering in the tens of thousands, and the potential for rapid increases in population

could result in large impacts on their ecosystem spanning trophic levels (Shine et al., 2006).

The myriad impacts of climate change on an ecosystem are overwhelmingly complicated

and interconnected. Thus, it is exceedingly difficult to develop models that accurately predict

comprehensive, species-specific outcomes in response to altered thermal landscapes. The present

study examined the effect of altered temperatures during gestation within a single large

population and thus represents only a single piece of the complex overall puzzle. As global

temperatures continue to rise, additional research examining the effects of increasing temperature

on the behavior and physiology of individual species, as well as studies encompassing entire

ecosystems, will provide the invaluable insight necessary to further our understanding of how

temperature affects ecosystems and, more importantly, will enhance our ability to predict the

impacts of future climate change on the global ecosystem.

7. Conclusions

Our research demonstrated that, even under the worst-case scenario climate change

model predictions, Red-sided garter snakes may see improved reproductive outcomes at the

northern edge of their range. Reduced gestation length has the potential to facilitate increased

fitness for both mother and offspring with no observed cost to reproductive outcomes.

References

ADDIS, E. A., E. J. GANGLOFF, M. G. PALACIOS, K. E. CARR, AND A. M. BRONIKOWSKI. 2017.

Merging the "morphology–performance–fitness" paradigm and life-history theory in the

Eagle Lake garter snake research project. Integrative and Comparative Biology 57:423-

435.

ALEKSIUK, M. 1976. Reptilian hibernation: evidence of adaptive strategies in *Thamnophis sirtalis*

parietalis. Copeia 12:170-178.

ALEKSIUK, M. AND P. T. GREGORY. 1974. Regulation of seasonal mating behavior in *Thamnophis*

sirtalis parietalis. Copeia 18:681-689.

ALEKSIUK, M. AND K. W. STEWART. 1971. Seasonal changes in the body composition of the

garter snake (*Thamnophis sirtalis parietalis*) at northern latitudes. Ecology 52:485-490.

ARNOLD, S. J., C. R. PETERSON. 2002. A model for optimal norms: the case of the pregnant garter

snake and her temperature sensitive embryos. American Naturalist 160:306-616.

BEUCHAT, C. 1988. Temperature effects during gestation in a viviparous lizard. Journal of

Thermal Biology 13:135-142.

BLANCHARD, F. N. AND F. C. BLANCHARD. 1940. Factors determining time of birth in the garter

snake *Thamnophis sirtalis sirtalis* (Linnaeus). Papers from the Michigan Academy of

Science, Arts and Letters 26:161–176.

BROMMER, J. E., J. MERILÄ, AND H. KOKKO. 2002. Reproductive timing and individual fitness.

Ecology Letters 5:802-810.

BRONIKOWSKI, A. M. 2000. Experimental evidence for the adaptive evolution of growth rate in

the garter snake *Thamnophis elegans*. Evolution 54:1760-1767.

BROWN, G. P. AND R. SHINE. 2004. Maternal nest-site choice and offspring fitness in a tropical

snake (*Tropidonophis mairii, Colubridae*). Ecology 85:1627-1634.

BUSH, E AND D. S. LEMMEN. 2019. Canada's Changing Climate Report. Government of Canada

Press, Canada.

CHARLAND, M. B. AND P. T. GREGORY. 1995. Movements and habitat use in gravid and

nongravid female garter snakes (Colubridae: *Thamnophis*). Journal of Zoology 236:543-

561.

Climate-data.org. Inwood climate: Weather Inwood & temperature by month. Available at

https://en.climate-data.org/north-america/canada/manitoba/inwood-108217/

COSTANZO, J. P. 1985. The bioenergetics of hibernation in the Eastern garter snake *Thamnophis*

sirtalis sirtalis. Physiological Zoology 58:682-692.

CREWS, D., M. GRASSMAN, W. GARSTKA, A. HALPERT, AND B. CAMAZINE. 1987. Sex and

seasonal differences in metabolism in the Red-sided garter snake, *Thamnophis sirtalis*

parietalis. Canadian Journal of Zoology 65:2362-2368.

DOODY, J.S. AND J.M. REFSNIDER. 2022. Nesting in reptiles: Natural and anthropogenic threats

and evolutionary responses. Frontiers in Ecology and Evolution 10:1103193.

DU, W. G. AND R. SHINE. 2015. The behavioural and physiological strategies of bird and reptile

embryos in response to unpredictable variation in nest temperature. Biological Reviews

90:19-30.

DUNLAP, K. D. AND J. W. LANG. 1990. Offspring sex ratio varies with maternal size in the

common garter snake, *Thamnophis sirtalis*. Copeia 1990:568-570.

ENVIRONMENT CANADA. 2014. Canada's sixth national report on climate change 2014 : actions to

meet commitments under the United Nations Framework Convention on Climate Change.

Pp. 120-121. Environment Canada, Canada.

FRIESEN, C. R., R. SHINE, R. W. KROHMER, AND R. T. MASON. 2013. Not just a chastity belt: the

functional significance of mating plugs in garter snakes, revisited. Biological Journal of

the Linnean Society 109:893-907.

FRIESEN, C. R., D. R. POWERS, P. E. COPENHAVER, AND R. T. MASON. 2015. Size dependence in

non-sperm ejaculate production is reflected in daily energy expenditure and resting

metabolic rate. Journal of Experimental Biology 218:1410-1418.

FRIESEN, C. R., R. SHINE, R. W. KROHMER, AND R. T. MASON. 2013. Not just a chastity belt: the

functional significance of mating plugs in garter snakes, revisited. Biological Journal of

the Linnean Society 109:893-907.

FRIESEN, C. R., M. K. SQUIRE, AND R. T. MASON. 2014. Intrapopulational variation of ejaculate

traits and sperm depletion in Red-sided garter snakes. Journal of Zoology 292:192-201.

GALLANA, M., M. P. RYSER-DEGIORGIS, T. WAHLI, AND H. SEGNER. 2013. Climate change and

infectious diseases of wildlife: altered interactions between pathogens, vectors and hosts.

Current Zoology 59:427-437.

GANS, C. 1996. An overview of parental care among the Reptilia. Advances in the Study of

Behavior. 25:145-157.

GAO, J. F., Y. F. QU, L. G. LUO, AND X. JI. 2010. Evolution of reptilian viviparity: a test of the

maternal manipulation hypothesis in a temperate snake, *Gloydius brevicaudus*

(*Viperidae*). Zoological Science 27:248-255.

GEISER, F. 2013. Hibernation. Current Biology 23:188-193.

GIBSON, A. R. AND J. B. FALLS. 1979. Thermal biology of the common garter snake *Thamnophis*

sirtalis (L.). I. Temporal variation, environmental effects and sex differences. Oecologia

1:79-97

GILMAN, S. E., M. C. URBAN, J. TEWKSBURY, G. W. GILCHRIST, AND R. D. HOLT. 2010. A

framework for community interactions under climate change. Trends in Ecology &

Evolution 25:325-331.

GREGORY, P. T. 1974. Patterns of spring emergence of the Red-sided garter snake (*Thamnophis*

sirtalis parietalis) in the Interlake region of Manitoba. Canadian Journal of Zoology

52:1063-1069.

GREGORY, P. T. 1982. Reptilian hibernation. 1982. Biology of the Reptilia. Pp. 53-154.

Academic Press, USA.

GREGORY, P. T., L. H. CRAMPTON, AND K. M. SKEBO. 1999. Conflicts and interactions among

reproduction, thermoregulation and feeding in viviparous reptiles: are gravid snakes

anorexic? Journal of Zoology 248:231-241.

GUNDERSON, A.R., M.E. DILLON, AND J.H. STILLMAN. 2017. Estimating the benefits of plasticity

in ectotherm heat tolerance under natural thermal variability. Functional Ecology

31:1529-1539.

HARVELL, D., S. ALTIZER, I. M. CATTADORI, L. HARRINGTON, AND E. WEIL. 2009. Climate

change and wildlife diseases: when does the host matter the most? Ecology 90:912-920.

HAWKES, L. A., A. C. BRODERICK, M. H. GODFREY, AND B. J. GODLEY. 2007. Investigating the

potential impacts of climate change on a marine turtle population. Global Change

Biology 13:923-932.

HOLDEN, K. G., E. J. GANGLOFF, E. GOMEZ-MANCILLAS, K. HAGERTY, AND A. M. BRONIKOWSKI.

2021. Surviving winter: Physiological regulation of energy balance in a temperate

ectotherm entering and exiting brumation. General and Comparative Endocrinology

307:113758.

HOLLELEY, C. E., D. O'MEALLY, S. D. SARRE, J. A. MARSHALL GRAVES, T. EZAZ, AND K.

MATSUBARA. 2015. Sex reversal triggers the rapid transitions from genetic to

temperature-dependent sex. Nature 523:79-82.

HOUGHTON, T., Y. DING, J. GRIGGS, N. NOGUER, P. VAN DER LINDEN, D. XIAOSU, K. MASKELL

AND A. JOHNSON. 2001. Climate change 2001: The scientific basis. Contribution of

Working Group I to the Third Assessment Report of the Intergovernmental Panel on

Climate Change. Quarterly Journal of the Royal Meteorological Society 128:1038-1039.

HUANG, W.S. AND D.A. PIKE. 2013. Testing cost-benefit models of parental care evolution using

lizard populations differing in the expression of maternal care. PloS one 8:e54065.

HUEY, R. B. AND R. D. STEVENSON. 1979. Integrating thermal physiology and ecology of

ectotherms: a discussion of approaches. American Zoologist 19:357-366.

HUEY, R. B. 1982. Temperature, physiology, and the ecology of reptiles. *Biology of the Reptilia,*

Physiology. Gans C and Pough FH (*Eds*.). London, Academic Press 12:25-91.

HUEY, R.B. AND D. BERRIGAN. 2001. Temperature, demography, and ectotherm fitness. The

American Naturalist 158:204-210.

HUGHES, D. F., P. R. DELIS, AND W. E. MESHAKA. 2019. Comparison of body temperatures

across physiological states in syntopic snake species (*Thamnophis sirtalis* and *Nerodia*

sipedon) from Pennsylvania. Northeastern Naturalist 26:749-760.

JAYNE, B. C. AND A. F. BENNETT. 1990. Selection on locomotor performance capacity in a

natural population of garter snakes. Evolution 44:1204-1229.

JENTSCH, A., J. KREYLING, J. E. BOETTCHER-TRESCHKOW, AND C. BEIERKUHNLEIN. 2009.

Beyond gradual warming: extreme weather events alter flower phenology of European

grassland and heath species. Global Change Biology 15:837-849.

JI, X., L. H. LIN, L. G. LUO, H. L. LU, J. F. GAO, AND J. HAN. 2006. Gestation temperature affects

sexual phenotype, morphology, locomotor performance, and growth of neonatal Brown

forest skinks, *Sphenomorphus indicus*. Biological Journal of the Linnean Society 88:453-

463.

KING, R. B. 2022. Temperature-induced multi-species cohort effects in sympatric snakes.

Ecology and Evolution 12:e8601.

KINGSOLVER, J. G., S. E. DIAMOND, AND L. B. BUCKLEY. 2013. Heat stress and the fitness

consequences of climate change for terrestrial ectotherms. Functional Ecology 27: 1415-

1423.

KISSNER, K. J. AND P. J. WEATHERHEAD. 2005. Phenotypic effects on survival of neonatal

Northern watersnakes *Nerodia sipedon*. Journal of Animal Ecology 74:259-265.

LADYMAN, M., X. BONNET, O. LOURDAIS, D. BRADSHAW, AND G. NAULLEAU. 2003. Gestation,

thermoregulation, and metabolism in a viviparous snake, *Vipera aspis*: evidence for

fecundity-independent costs. Physiological and Biochemical Zoology 76:497-510.

LORIOUX, S., H. LISSE, AND O. LOURDAIS. 2013. Dedicated mothers: predation risk and physical

burden do not alter thermoregulatory behaviour of pregnant vipers. Animal Behaviour

86:401-408.

LOURDAIS, O., X. BONNET, R. SHINE, D. DENARDO, G. NAULLEAU, AND M. GUILLON. 2002.

Capital-breeding and reproductive effort in a variable environment: a longitudinal study

of a viviparous snake. Journal of Animal Ecology 71:470-479.

LOURDAIS, O., R. SHINE, X. BONNET, M. GUILLON AND G. NAULLEAU. 2004. Climate affects

embryonic development in a viviparous snake *Vipera aspis*. Oikos 104:551-560.

LOURDAIS, O., M. GUILLON, D. DENARDO, AND G. BLOUIN-DEMERS. 2013. Cold climate

specialization: adaptive covariation between metabolic rate and thermoregulation in

pregnant vipers. Physiology & Behavior 119:149-155.

LUTTERSCHMIDT, D. I., M. P. LEMASTER, AND R. T. MASON. 2006. Minimal overwintering

temperatures of Red-sided garter snakes (*Thamnophis sirtalis parietalis*): a possible cue

for emergence? Canadian Journal of Zoology 84:771-777.

MADSEN, T. AND R. SHINE. 2002. Short and chubby or long and slim? food intake, growth and

body condition in free-ranging pythons. Austral Ecology 27:672-680.

MASSON-DELMOTTE, V., P. ZHAI, A. PIRANI, S.L. CONNORS, C. PÉAN, S. BERGER, N. CAUD, Y.

CHEN, L. GOLDFARB, M. I. GOMIS, AND M. HUANG. 2021 Climate change 2021: the

physical science basis. Contribution of working group I to the sixth assessment report of

the intergovernmental panel on climate change. Cambridge University Press, UK.

MATHIES, T. AND R. M. ANDREWS. 1997. Influence of pregnancy on the thermal biology of the

lizard, *Sceloporus jarrovi*: why do pregnant females exhibit low body temperatures?

Functional Ecology 11:498-507.

MATSUBARA, K., H. TARUI, M. TORIBA, K. YAMADA, C. NISHIDA-UMEHARA, K. AGATA, AND Y.

MATSUDA. 2006. Evidence for different origin of sex chromosomes in snakes, birds, and

mammals and step-wise differentiation of snake sex chromosomes. Proceedings of the

National Academy of Sciences, USA 103:18190-18195.

MICHEL, C. L., J. H. PASTORE, AND X. BONNET. 2013. Impact of cool versus warm temperatures

on gestation in the Aspic viper (*Vipera aspis*). Comparative Biochemistry and Physiology

Part A: Molecular & Integrative Physiology 165:338-342.

MILES, D. B., B. SINERVO, AND W. A. FRANKINO. 2000. Reproductive burden, locomotor

performance, and the cost of reproduction in free-ranging lizards. Evolution 54:1386-

1395.

NNADI, N. E. AND D. A. CARTER. 2021. Climate change and the emergence of fungal pathogens.

PLoS Pathogens 17:e1009503.

NOER, N. K., M. ØRSTED, M. SCHIFFER, A. A. HOFFMANN, S. BAHRNDORFF, AND T. N.

KRISTENSEN. 2022. Into the wild—a field study on the evolutionary and ecological

importance of thermal plasticity in ectotherms across temperate and tropical regions.

Philosophical Transactions of the Royal Society B 377:20210004.

O'DONNELL, R., R. SHINE, AND R. T. MASON. 2004. Seasonal anorexia in the male Red-sided

garter snake, *Thamnophis sirtalis parietalis*. Behavioral Ecology and Sociobiology

56:413-419.

PAAIJMANS, K. P., R. L. HEINIG, R. A. SELIGA, J. I. BLANFORD, S. BLANFORD, C. C. MURDOCK,

AND M. B. THOMAS. 2013. Temperature variation makes ectotherms more sensitive to

climate change. Global Change Biology 19:2373-2380.

PARMESAN, C. 2006. Ecological and evolutionary response to recent climate change. Annual

Review of Ecology, Evolution, and Systematics 37:637-669.

RABAIOTTI, D., T. COULSON, AND R. WOODROFFE. 2023. Climate change is predicted to cause

population collapse in a cooperative breeder. Global Change Biology 29:6002-6017.

SALINAS-RAMOS, V. B., L. ANCILLOTTO, L. BOSSO, V. SÁNCHEZ-CORDERO, AND D. RUSSO. 2020.

Interspecific competition in bats: state of knowledge and research challenges. Mammal

Review 50:68-81.

SEIGEL, R. A., M. M. HUGGINS, AND N. B. FORD. 1987. Reduction in locomotor ability as a cost

of reproduction in gravid snakes. Oecologia 73:481-485.

SHERMAN, P. W. AND M. C. RUNGE. 2002. Demography of a population collapse: the northern

Idaho ground squirrel (*Spermophilus brunneus brunneus*). Ecology 83:2816–2831.

SHINE, R. 1988. Parental care in reptiles. Biology of the Reptilia, Ecology B: defense and life

history 16:275-329.

SHINE, R., T. LANGKILDE, M. WALL, AND R. T. MASON. 2006. Temporal dynamics of emergence

and dispersal of garter snakes from a communal den in Manitoba. Wildlife Research

33:103–111.

SINGCHAT, W., S. F. AHMAD, N. LAOPICHIENPONG, A. SUNTRONPONG, T. PANTHUM, D. K.

GRIFFIN, AND K. SRIKULNATH. 2020. Snake W sex chromosome: The shadow of ancestral

amniote super-sex chromosome. Cells 9:2386.

STOTT, P. 2016. How climate change affects extreme weather events. Science 352:1517-1518.

TANG, X., F. YUE, X. YAN, D. ZHANG, Y. XIN, C. WANG, AND Q. CHEN. 2012. Effects of gestation

temperature on offspring sex and maternal reproduction in a viviparous lizard (*Eremias*

multiocellata) living at high altitude. Journal of Thermal Biology 37:438-444.

URBAN, M. C., J. J. TEWKSBURY, AND K. S. SHELDON. 2012. On a collision course: competition

and dispersal differences create no-analogue communities and cause extinctions during

climate change. Proceedings of the Royal Society B: Biological Sciences 279:2072-2080.

VIANA, P. F., T. EZAZ, M. DE BELLO CIOFFI, T. LIEHR, A. AL-RIKABI, L. G. GOLL, A. M. ROCHA,

AND E. FELDBERG. 2020. Landscape of snake'sex chromosomes evolution spanning 85

MYR reveals ancestry of sequences despite distinct evolutionary trajectories. Scientific

Reports 10:1-4.

WANG, Z., L. MA, M. SHAO, AND X. JI. 2017. Are viviparous lizards more vulnerable to climate

warming because they have evolved reduced body temperature and heat tolerance?

Oecologia 185:573-582.

WALTHER, G. R., C. A. BURGA, P. J. EDWARDS, AND P. J. EDWARDS (Eds.) 2001. "Fingerprints"

of climate change: adapted behaviour and shifting species. Springer Science & Business

Media

WHITTIER, J.M., R. T. MASON, AND D. CREWS. 1985. Mating in the red-sided garter snake,

Thamnophis sirtalis parietalis: differential effects on male and female sexual behavior.

Behavioral Ecology and Sociobiology 16:257-261.

WILSON, R. C. 2020. Energy Stores and Life-history Transitions in Red-Sided Garter Snakes

(*Thamnophis sirtalis parietalis*). Doctoral dissertation, Portland State University, USA.

YAN, X. F., X. L. TANG, F. YUE, D. J. ZHANG, Y. XIN, C. WANG, AND Q. CHEN. 2011. Influence of

ambient temperature on maternal thermoregulation and neonate phenotypes in a

viviparous lizard, *Eremias multiocellata*, during the gestation period. Journal of Thermal

Biology 36:187-192.

ZHANG, D. J., X. L. TANG, F. YUE, Z. CHEN, AND Q. CHEN. 2010. Effect of gestation temperature

on sexual and morphological phenotypes of offspring in a viviparous lizard, *Eremias*

multiocellata. Journal of Thermal Biology 35:129-133.

ZHANG, X., L. A. VINCENT, W. D. HOGG AND A. NIITSOO. 2000. Temperature and precipitation

trends in Canada during the 20th century. Atmosphere-ocean 38:395-429.

Supplementary Table 1: Median effect of a 5 ˚C increase in ambient temperature on gestation

length and neonate body condition in Red-sided garter snakes.

	Mass (g)	SVL (cm)	BCI (g/cm)	Gestation (days)	Litter size
Treatment (27 ⁰C)	1.47 (SD 0.23)	15.78 (SD 0.76)	0.095 (SD 0.01)	97.0 (SD 3.2)	21.0 (SD 6.5)
Control (22 ⁰C)	1.33 (SD 0.21)	15.25 (SD 1.10)	0.085 (SD 0.01)	118.0 (SD 7.1)	18.0 (SD 8.7)
Wilcoxon (*P*-value)	<0.001	<0.001	<0.001	<0.001	>0.74 (ns)

Fig. 1. Reproductive outcomes under a 5 ⁰C increase in ambient temperature during gestation.

Plots A-E show comparison between control (ambient daytime temperature 22 ⁰C) and treatment

(ambient daytime temperature 27 ⁰C) conditions. All comparisons made using a nonparametric

two-sample Wilcoxon rank test except for plot C. A: Gestation length in days as measured from

the day of confirmed copulation to the day of birth ($P < 0.001$). B: Litter size for each female

that successfully gave birth ($P > 0.74$). C: The proportion of mated females that gave birth were

compared using a two proportions z-test ($P > 0.77$). D: Neonate snout-vent lengths in cm ($P <$

0.001). E: Neonate mass in grams ($P < 0.001$). F: Neonate body condition index of mass to SVL

ratio ($P < 0.001$).