Increased offspring size and accelerated gestation in a temperate ectothermic vertebrate

under temperatures predicted by a worst-case climate change model.

David L Hubert[1,*], Ehren J Bentz[2], and Robert T Mason[1]

[1] *Department of Integrative Biology, Oregon State University, Corvallis, Oregon. USA, 97333*

[2] *Department of Psychology, Cornell University, Ithaca, New York, USA*

[*] *Corresponding author. E-mail address: hubertd@oregonstate.edu*

ABSTRACT

As global temperatures continue to increase, understanding the impacts of warming

environments is increasingly relevant. Temperature is especially relevant for ectothermic

organisms which depend upon consistent and predictable annual temperature cycles for

reproduction and development. However, additional research is required in this area to elucidate

the potential impacts of climate change on future generations. To understand how projected

increases in environmental temperatures may impact reproductive outcomes within natural

populations of ectothermic vertebrates, we manipulated minimum ambient temperatures during

gestation in Red-sided garter snakes (*Thamnophis sirtalis parietalis*). Wild snakes were collected

in the Interlake region of Manitoba, Canada during their spring mating season and allowed to

mate in controlled conditions. For the duration of gestation, mated females were placed into one

of two ambient thermal conditions: temperatures emulating those found in the species' natural

habitat or temperatures with a consistent 5 ⁰C increase to match end-of-century climate change

projections. We recorded observations for each litter and all neonates resulting from controlled

mating trials. We observed no difference in litter sizes or birth rates between thermal conditions.

However, we observed both a significant reduction in gestation length and a significant increase

in neonate body size associated with increased ambient temperatures. These results suggest that

increased minimum temperatures during gestation may confer reproductive benefits for the

northern populations of this species even under the most extreme current modeled warming

predictions. We discuss the broader implications of this effect, including possible negative

ecological outcomes.

Key words: Climate change, Reproduction, Gestation, Ectotherm, Viviparous, Thamnophis

sirtalis parietalis

1. Introduction

In a time of rapidly changing environmental temperatures, it is increasingly important to

elucidate how altered thermal conditions impact the coordination of important life history stages

and essential events such as mating, gestation, and foraging. Because reproductive success is a

defining feature of individual fitness, factors that greatly impact reproduction among individuals

within a population have the potential to dramatically affect the long-term success and survival

of that population (Brommer et al, 2002; Sherman and Runge, 2002; Rabaiotti et al., 2023).

Ectotherms are influenced particularly strongly by environmental conditions, especially

temperature, which is frequently an important cue used to coordinate behavior and physiological

processes essential to reproductive success including courtship behavior and successful

incubation or gestation (Huey and Stevenson, 1979; Huey, 1982; Miles et al., 2000; Lourdais et

al., 2013). Ectotherms from terrestrial ecosystems demonstrate existing plastic physiological and

behavioral mechanisms to maintain homeostasis during large seasonal and acute temperature

fluctuations (Huey et al., 2001; Gunderson et al., 2017; Noer et al., 2022). These extreme

temperature fluctuations have been increasingly exacerbated by recent climate change (Masson-

Delmotte et al., 2021). Therefore, investigating how terrestrial ectotherms respond to these new

conditions provides an excellent opportunity to explore the broader impacts of a warming

climate.

The challenge of maintaining optimal thermal conditions during embryonic development

is central to reproductive success among many ectotherms, though overcoming this challenge is

accomplished using different strategies. Oviparous (egg laying) ectotherms must ensure

consistent thermal conditions for their eggs throughout the entire period of embryonic

development by utilizing complex behaviors such as careful nest building, precise nest site

selection, or by maintaining thermal conditions through monitored incubation of clutches (Du

and Shine, 2014). Typically, these nest sites are abandoned by the parent shortly after egg

deposition, allowing the nest itself to maintain environmental conditions conducive to proper

embryonic development while the mother is free to forage and rebuild depleted energy reserves

(Shine, 1988; Gans et al., 1996; Huang and Pike, 2013; Doody and Refsnider, 2022). In contrast,

viviparous (live-bearing) ectotherms must alter their behavior while gravid to maintain optimal

internal thermal conditions throughout gestation, even to the extent of risking their own survival

to prioritize ideal thermal conditions for developing embryos (Seigel et al., 1987; Charland and

Gregory, 1995; Gregory et al., 1999; Lorioux et al., 2013). This behavioral tradeoff serves to

illustrate the importance of an organism's thermal environment, and that maintaining proper

temperature is vital for the survival and optimal development of offspring and therefore directly

impacts the individual fitness of reproducing adults. Temperatures experienced during gestation

are often observed to impact neonate size in viviparous squamates (Mathies and Andrews, 1997;

Lourdais et al., 2004; Gao et al., 2010; King, 2022). Both greater neonate body length (Brown

and Shine, 2004; Kissner and Weatherhead, 2005; Addis et al., 2017) and greater body mass

(Jayne and Bennett, 1990; Bronikowski, 2000; Kissner and Weatherhead, 2005) are associated

with significantly increased rates of survival to adulthood in viviparous snakes.

Changes to environmental thermal conditions during gestation have been observed to

impact both physiology and behavior of viviparous ectotherms to alter characteristics such as

maternal energy use and basking and foraging behavior (Lourdais et al., 2002; Ladyman et al.,

2003; Wang et al., 2017), gestation length (Blanchard and Blanchard, 1940; Ji et al., 2006; Wang

et al., 2017; King, 2022), litter size (Lourdais et al., 2004; Zhang et al., 2010; Tang et al., 2012),

and body condition and viability of neonates (Lourdais et al., 2004; Wang et al., 2017; King,

2022). Although there is often agreement regarding the influence of warmer temperatures on

gestation, neonate survival, and body condition, the specific effects observed differ between

studies and depend on the biological system examined. Wang et al. (2017) reported that

increased ambient temperature was associated with significantly shorter gestation length,

increased neonate mass, and a decrease in general activity during gestation for Toad-headed

lizards (*Phrynocephalus putajatia*). Beuchat (1988), found that increased environmental

temperatures during gestation were associated with a decrease in gestation length in Yarrow's

spiny lizards (*Sceloporus jarrovi*), but this study also reported that sustained increased

temperatures resulted in a detrimental effect on the body conditions of both mother and neonate.

This latter general trend was further supported, as Tang et al. (2012) reported an increase in litter

size and a decrease in gestation length associated with higher temperatures for Multi-ocellated

racerunner lizards (*Eremias multiocellata*), but that these potentially beneficial effects were lost

as temperatures continued to increase. Lourdais et al. (2004) demonstrated that increased

ambient temperature resulted in shorter gestation length, increased embryo viability, and caused

phenotypic changes, including changes to scale patterns, in neonate European asps (*Vipera aspis*)

within a population living at the northern boundary of its range. Each of these studies

demonstrate that, even among animals that are able to effectively behaviorally thermoregulate,

increased ambient temperatures significantly impact both the behavior of gravid females and the

resulting reproductive outcomes in viviparous species.

Red-sided garter snakes (*Thamnophis sirtalis parietalis)* are viviparous ectotherms and

one of the most northerly living squamate species in North America (Aleksuik and Stewart,

1971). The focal population of this study is found in the Interlake region of Manitoba, Canada

near the northern extreme of the species' range which spans as far South as Northern Texas. The

Interlake region is particularly well suited for studies examining the effects of temperature on

ectothermic vertebrates, as it is characterized by major seasonal changes to the thermal

landscape. Unsurprisingly, ectotherms living within this region are well adapted to this extreme

environment and demonstrate highly coordinated annual behavioral cycles primarily driven by

seasonal temperature changes (Aleksiuk and Stewart, 1971; Gregory, 1974; Lutterschmidt et al.,

2006). Indeed, Red-sided garter snakes spend the majority of their life in a hibernation-like state

of winter dormancy known as "brumation," during which the entire population retreats into

subterranean hibernacula for up to 8 months each year (Aleksiuk and Stewart, 1971; Costanzo,

1985). Brumation is a behavioral and physiological state characterized by extremely low levels

of activity and greatly reduced metabolic rates in ectotherms. This state is driven by low

temperatures and serves as an essential strategy for surviving extremely cold and resource-

limited winter conditions (Gregory, 1982; Geiser, 2013). Red-sided garter snakes remain

relatively inactive throughout brumation and do not feed for approximately 8-9 months each year

(O'Donnell, 2004). Thus, they must rely solely upon stored energy to survive winter dormancy

as well as the intense and energetically costly activities of mating and migration immediately

following emergence from the hibernacula in early spring (Aleksiuk and Gregory, 1974;

Aleksiuk, 1976). These energy stores must be obtained during an extremely brief summer

feeding season lasting as little as 10 weeks, during which an individual must acquire all of the

energy necessary for an entire year of growth, development, reproduction, and survival (Crews et

al., 1987; O'Donnell et al., 2004). Moreover, the challenge of building energy reserves becomes

even more difficult for reproducing females as this brief feeding period coincides with gestation

– a period that is both behaviorally and energetically demanding. Unlike the southern

populations, the northern populations optimal body temperatures are much higher relative to

environmental conditions, suggesting minimum ambient temperatures may play a large role in

the behavior and reproductive outcomes for snakes undergoing gestation in this region. Thus,

this limited window of time is critically important and has the potential to dramatically impact

individual fitness, as a reproducing female must carefully balance activities and energy demands

to simultaneously ensure survival while maximizing reproductive success.

Although climate change-induced warming has resulted in a global average temperature

increase of approximately 0.6 ºC over the last 60 years, the Interlake region has been

disproportionately impacted, experiencing nearly triple the global rate with recorded

temperatures showing an increase of 1.7 ⁰C over the same period (Zhang et al., 2000; Houghton

et al., 2001; Environment Canada, 2014; Bush and Lemmen, 2019). Global temperatures are

predicted to continue to increase by as much as 5.3 ⁰C by the end of the century, which is well

beyond the findings of previous work where increases of as little as 2 ⁰C significantly impacted

gestation in viviparous ectotherms (Houghton et al., 2001; Kingsolver et al., 2013; Wang et al.,

2017; Masson-Delmotte et al., 2021). With current regional warming levels so much higher than

the global average, it is reasonably likely that future regional warming will be at the higher end

of global climate change model projections. While increased temperature in this region will

likely not result in thermal conditions that are significantly warmer than those currently

experienced by this species at the southern end of their range, increases to minimum

temperatures of this degree have a large potential to alter behavioral thermal regulation and

reproductive outcomes for snakes at the colder northern end of their range. This is especially true

for increases in nighttime temperatures, where even during summer they currently remain well

below optimal gestation temperatures.

Currently, the daily mean environmental temperatures experienced by Red-sided garter

snakes during the summer feeding and gestation period range from 13.8 ⁰C in late September to

20.1 ⁰C in July, with mean maximum daily temperatures 25 ⁰C, and mean minimum temperatures

15.9 ⁰C (https://en.climate-data.org/north-america/canada/manitoba/inwood-108217/). Previous

research focused on the Red-sided garter snakes and other members of the genus *Thamnophis*

have shown that both free-ranging and captive gravid females behaviorally maintain daytime

body temperatures between 26 ⁰C and 31 ⁰C – a range that typically exceeds ambient

temperatures (Gibson and Falls, 1979; Arnold and Peterson, 2002; Hughes et al., 2019). Though

often overlooked, nighttime temperatures are rising faster than daytime temperatures, resulting in

a potentially larger impact to minimum temperatures than maximum temperatures, especially for

organisms living at the coldest edges of their range (Alexander et al., 2006; Rutschmann et al.,

2023)

In the present study, we manipulated the thermal environment of gestating female Red-

sided garter snakes by increasing minimum diurnal and nocturnal ambient temperatures by an

ecologically relevant 5 ⁰C and observed the impact of this increase on gestation length and the

neonate and litter-wide traits at birth. We predicted an increased minimum ambient temperature

of 5 ⁰C would significantly reduce gestation length, with potential increases to neonate mass,

body condition, and parturition success. However, we also predicted that if this thermal increase

exceeds the beneficial thresholds of this population, potential individual gains would be lost

resulting in a reduction in neonate body condition, reduced offspring per litter, increases to

stillbirths and severe deformities (Beuchat, 1988; Tang et al., 2012).

169 2. Materials and methods

170   All experimental protocols presented were approved by the Oregon State University

171 Institutional Animal Care and Use Committee (Animal Care and Use Protocol 5058), in

172 accordance with the Association for the Assessment and Accreditation of Laboratory Animal

173 Care. This research was conducted under the authority of Manitoba Wildlife Scientific Permit

174 No. WB20333.

 Animal Collection

We collected free-ranging male and female Red-sided garter snakes (approximately 100

of each) from the natural population at over-wintering hibernacula near the town of Inwood,

Manitoba, Canada during the spring mating season between 1 May and 4 May 2018. This site

contains several communal hibernacula that collectively shelter a population estimated at

approximately 35,000 snakes (Shine et al., 2006). Male snakes were selected and captured after

being observed actively exhibiting courtship behavior. Non-mated females were selected and

captured only if their initial emergence from the hibernaculum was observed, and they were

being actively courted by males. To minimize the effects of body size variation on litter size,

only females of the same approximate size were selected, with snout-vent length (SVL) and body

masses of approx. 60 cm and 100 g, respectively. Due to the extremely large population and ease

of collection, we were able to limit collection to this target size while easily maintaining a

sufficient sample size.   To confirm that the females had not yet mated that season, we visually

inspected their cloaca for copulatory plugs upon collection. Copulatory plugs are large,

gelatinous spermatophores deposited by a male inside of a female's cloaca during copulation.

Copulatory plugs are substantial in size, persist for several days, and represent an easily

identifiable and reliable indication that a female has recently mated (Whittier et al., 1985; Friesen

et al., 2013). Any female presenting a copulatory plug, or any indication of potentially having

mated, was immediately released. All animals were collected by hand and placed in cloth bags

for transportation to the Chatfield Research Station, Chatfield, Manitoba. Snakes were separated

by sex and housed outdoors in specially designed 1 m^3 nylon enclosures containing natural

stones, hide boxes, and water dishes. Water was provided *ad libitum* for the duration of captivity,

but food was not provided in the field or at the research station, as this species is aphageous

during the spring mating season (O'Donnell et al., 2004, Crews et al., 1987). Capture, handling,

and transportation are not likely to have impacted results, as previous studies have consistently

shown that these activities do not adversely affect survival, reproduction, or courtship behavior

(Friesen et al., 2013; Friesen et al., 2014).

Mating Trials

To control mating conditions and timing, we used courtship arenas (opaque, cylindrical,

nylon enclosures 0.5 m in diameter and 1 m tall) with heat lamps (above) and electric heating

pads (below) to maintain homogenous thermal environments at $\geq 13.0\ ^\circ$C – optimal temperatures

to stimulate vigorous courtship behavior. Female snakes were placed individually into arenas and

paired with 5 randomly selected males. All snakes were allowed to freely behave under constant

observation until successful copulation occurred. Successful copulation was confirmed by visual

inspection and identification of a copulatory plug. Females with copulatory plugs were separated

from the males and transferred to a 1 m^3 outdoor nylon enclosure containing only mated females.

Each male was allowed to mate only once. Males that successfully mated were removed from the

remainder of the study to minimize any potential bias that may arise from allowing a male to sire

more than a single litter of offspring. All males were marked dorsally with indelible ink to

prevent recapture and released at the site of capture within 5 days of collection. All controlled

matings took place within the same 48-hour period, concluding when 80 successful matings

occurred. Similar experiments have reported that 23-64% of mated female Red-sided garter

snakes give birth in a laboratory setting (Friesen et al., 2014; Blakemore 2017; Dayger et al.,

2018), as such our sample size should result in 10-20 successful litters per treatment.

Experimental Conditions and Data Collection

We transported the mated females ($n = 80$) by car to the laboratory facilities at Oregon

State University. Animals were transported in cloth bags, housed in a thermally controlled

environment (air temperatures between 20 ⁰C and 22 ⁰C), and given water via a damp sponge to

prevent dehydration for the duration of the trip (< 30 h). Though females were all approximately

the same size as a result of carefully targeted collection, further efforts were made to ensure any

remaining size variation was accounted for between treatment and control groups. Female snakes

were visually sorted by SVL, then size matched into pairs based on BCI estimate. Size matched

pairs were then randomly split into either the treatment or control condition, from largest to

smallest to ensure an even distribution of size ranges in each group of 40 snakes. Snakes were

marked ventrally via scale clipping to indicate both the experimental condition and the relative

size ranking within each treatment. Once assigned to a group, and for the remainder of the study,

snakes were housed within microprocessor-controlled environmental chambers (Environmental

Growth Chambers, Chagrin Falls, OH). The control group was housed within a chamber

emulating the thermal environment found in this species' natural range during the summer

feeding period, which consists of a daytime temperature of 22 ⁰C and a nighttime temperature of

15 ⁰C. The treatment group was housed in an identical chamber set to maintain ambient

temperatures increased by 5 ⁰C, which consists of a daytime temperature of 27 ⁰C and a

nighttime temperature of 20 ⁰C. Females were housed individually in 45 liter glass aquaria

containing approximately 2.5cm of CareFresh™ paper fiber substrate, a water bowl, a hide box,

a rock to facilitate shedding of skin, and a heat lamp situated at one end to produce a thermal

gradient allowing behavioral thermoregulation within the constraints of the experiment.

Maximum temperatures under the basking lamps were approximately 31 ⁰C for control group

and 40 ⁰C for the treatment group. Minimum temperatures never dropped below the ambient

temperature for each treatment (15-22 ⁰C or 20-27 ⁰C). Thus, this study tightly controlled the

minimum ambient temperatures snakes were able to access during gestation while allowing

snakes to freely thermoregulate and access higher temperatures if preferred, mimicking thermal

opportunities experienced in the wild where minimum temperatures are reached at night but

behavioral thermal regulation during the day allows for higher diurnal body temperatures.

Photoperiods were identical for both treatments and emulated that of the summer feeding period

in Manitoba (16 h:8 h light:dark cycle). Humidity was maintained at 65 % for both treatments.

Water was provided *ad libitum*, and food (a mixture of earthworms and salmon fry supplemented

with calcium) was provided once per week, during which time all snakes were allowed to eat to

satiation. Snakes were maintained in these conditions for the duration of the experiment.

Each female was inspected daily to identify births, and offspring were collected within no

more than 24 hours of birth. Dates of birth and number of offspring were recorded for each litter.

Neonates were weighed and measured immediately after collection. Masses were recorded to the

nearest 0.01 g and SVLs were recorded to the nearest 1 mm. To assess the overall health of

neonates, a Body Condition Index (BCI) was calculated using the residuals from a linear

regression model fit to the relationship between neonate SVL and neonate mass, allowing for a

standardized measure of body condition. BCI is frequently used as a measure of approximate

overall health and nutritional state as it accounts for both body mass and SVL for the same

individual, allowing for a better description of the individual than comparing those values

separately (Madsen and Shine, 2002). Measurements were collected for every litter and for every

neonate until the last female gave birth in late September (see Supplemental Table 1 for full

data). All adult snakes were maintained in captivity until the following spring then returned to

the location of capture. Offspring were maintained in the lab for further data collection

facilitating additional studies aiming to characterize growth and development.

Statistical methods

To evaluate the effect of treatment on reproductive outcomes, a linear mixed effects

model (LMM) was used to evaluate the effect of treatment (treatment or control) on traits

including neonate mass (g), SVL (cm), BCI, as well as gestation length (days) using the

following:

272
$$Y_i = \beta_0 + \beta_1 X + bZ + \varepsilon$$

Where Y_i is trait of interest, β_0 is the intercept, β_1 is the coefficient for the fixed effect of

X (treatment), b is the coefficient for the random effect of Z (mother), and ε is the error term.

Using an LMM allows us to account for random minor differences in physiology, behavior, and

environmental conditions by treating the mother as a random effect. Additionally, LMMs are

robust to violations of normality and independence, allowing us to make full use of the dataset

without limiting analyses to mean litter values for each trait, as well as enabling the use of non-

transformed data to generate results that can be directly interpreted (Bates et al., 2015; Schielzeth

et al., 2020). A model that also included relative maternal size as fixed effect, as well as a further

reduced model that only accounted for the random effects of mother were compared using the

Akaike information criterion (AIC), however the model above resulted in the best fit

(Supplemental Table 2). All LMM analyses were conducted using the lme4 package (Bates et al.,

2015) in R (R Core Team, 2023). All figures created using the ggplot2 package in R (Wickham,

2016).

To further evaluate the effect of treatment on reproductive outcomes we used a linear

model (LM) to compare litter outcomes between treatments (See Supplemental Table 3 for litter

wide traits). Treatment (treatment or control) was treated as a fixed effect on traits including total

litter mass (g), average neonate mass (g), average neonate SVL (cm), average neonate BCI, litter

size, number of stillborn, and number of deformed neonates per litter, using the following

equation:

292
$$Y_i = \beta_0 + \beta_1 X + \varepsilon$$

Where Y_i is the trait of interest for each litter, β_0 is the intercept, β_1 is the coefficient for

the predictor X (treatment), and ε is the error term. All LM analyses were conducted using the

stats package (version 4.3.1) in R. All figures created using the ggplot2 package in R.

To explore the potential relationships between traits measured, an LM that is similar to

the one above, however the X variable now represented one of the traits, was used as described

above. Traits compared this way include neonate mass, SVL, BCI, gestation length, litter size.

Additional analyses were conducted to explore the relationship between relative maternal size

and those same traits, however, due to the ranked nature of the relative maternal size, Kendall's

Tau correlation was calculated using the ggpubr (version 0.6.0) package in R. All correlation

figures were visualized using ggplot2.

A two proportions z test with continuity correction was used to evaluate the proportion of

females that gave birth out of the total number of mated females for each treatment using the

prop.test function of the stats package in R.

3. Results

A total of 26 mated females reached parturition, with 12 females from the treatment and

14 from the control group successfully giving birth. The proportion of mated females that

reached parturition was well within the reported range for a laboratory setting, with 35% of

control and 30% of treatment snakes successfully giving birth. When compared, there was no

significant difference in birthrate between treatments ($z^2 = 0.06$, $p > 0.8$, $n_{treatment} = 12$, $n_{control} =$

14) (Figure 1L).

To evaluate the effect of treatment on the reproductive outcomes we employed LMMs to

estimate the effect of treatment on traits including neonate mass, SVL, BCI, and gestation length.

Neonate mass, BCI, and gestation length were all found to be significantly affected by the

treatment condition ($p < 0.05$), with no difference present in SVL between treatments ($p > 0.5$)

(Figure 1A-D and Table 1). The effect of treatment increased both neonate mass and BCI, with

an estimated increase of 0.19 g and 0.16 respectively, and decreased gestation length by an

estimated 24 days. This shows that not only did the treatment condition reduce gestation length

by approximately 20%, but also increased neonate mass by approximately 14% while

maintaining neonate SVL.

To further evaluate the effects of treatment on reproductive outcomes, we used LMs to

estimate the effect of treatment on litter-wide outcomes including total litter mass (g), mean

neonate mass per litter (g), mean neonate SVL per litter (cm), mean neonate BCI per litter, litter

size, number of stillbirths per litter, and number of significant developmental deformities per

litter. Only mean neonate mass per litter and mean neonate BCI per litter were significantly

different between treatments ($p < 0.05$), with the estimated effect of treatment consistent with

those found in the LMM results (0.19 g and 0.16 respectively) (Figure 1E-K and Table 2).

A strong positive relationship was found between neonate mass and neonate SVL for

both treatment ($F_{[1, 211]} = 290$, $p < 0.001$, $R^2 = 0.58$, $n = 213$) and control ($F_{[1, 220]} = 294$, $p <$

0.001, $R^2 = 0.57$, $n = 222$) (Figure 2B), however the contribution of neonate SVL and neonate

mass to neonate BCI was less clear. Neonate mass had a strong positive relationship with

neonate BCI for treatment ($F_{[1, 211]} = 348$, $p < 0.001$, $R^2 = 0.62$, $n = 213$), but this positive

relationship was much weaker in control ($F_{[1, 220]} = 33.4$, $p < 0.001$, $R^2 = 0.13$, $n = 222$) (Figure

2C). Neonate SVL had a weak negative relationship with neonate BCI in control ($F_{[1, 220]} = 27.8$,

$p < 0.001$, $R^2 = 0.11$, $n = 222$), and a very weak positive relationship in treatment ($F_{[1, 211]} = 9.0$,

$p < 0.005$, $R^2 = 0.04$, $n = 213$) (Figure 2A). Taken together, this suggests that while there was no

difference in SVL between treatments (Figure 1C), neonates with a higher SVL were able to

build more mass in the treatment group compared to the control group, where neonates had

difficulty maintaining BCI for longer individuals.

Litter size had little to no relationship with gestation length, neonate SVL, neonate mass

or neonate BCI for the treatment group, but a moderately negative relationship was found

between litter size and neonate mass ($F_{[1, 220]} = 101$, $p < 0.001$, $R^2 = 0.31$, $n = 222$), neonate SVL

($F_{[1, 220]} = 182$, p <0.001, $R^2 = 0.45$, n = 222), and gestation length ($F_{[1, 220]} = 164$, p <0.001, $R^2 =$

0.43, n = 222) for the control group (Supplemental Figure 1 and Supplemental Table 4). This

suggests that cost associated with larger litters for the treatment group was generally very small,

but controls with larger litters had moderate reductions to neonate mass and SVL as well as

shorter gestation length.

A similar pattern emerged for the relationship between gestation length and neonate

traits, with weak relationships between gestation length and neonate traits for the treatment

group, but stronger relationships for the control group (Supplemental Figure 2 and Supplemental

Table 4). Interestingly, we see further evidence for the control groups having difficulty putting

on weight for longer neonates, as we see a positive relationship between gestation length and

neonate SVL ($F_{[1, 220]} = 65.1$, p < 0.001, $R^2 = 0.23$, n = 222), no real relationship between

gestation length and neonate mass ($F_{[1, 220]} = 5.4$, p < 0.05, $R^2 = 0.02$, n = 222), and a negative

relationship between gestation length and neonate BCI ($F_{[1, 220]} = 58.4$, p < 0.001, $R^2 = 0.21$, n =

222). This shows that neonate mass was maintained regardless of gestation length, but longer

gestion length resulted on longer, but skinnier neonates in the control condition.

When comparing relative maternal size to neonate traits, we see several relationships that

are significant, however all of these relationships show a very weak effect (Supplemental figure

3 and Supplemental Table 5). We believe that this shows that while maternal physiology plays a

large role in neonate physiology, the weakness of the relationships in our data reflects our efforts

to control for differences in maternal size during our initial sample selection.

All code used for the statistical analysis above and for all figures presented can be found

at https://github.com/hubertdl/Tsp_Gestation_and_ClimateChange.

4. Discussion

Our observations of gestation times accurately agreed with our initial prediction that a 5 °C increase in minimum temperature during gestation would result in a reduction in gestation length. Indeed, we found that gestation lengths were significantly reduced in the treatment condition, such that the extent of the resulting ranges of birth dates did not overlap between the treatment and control groups. The average gestation length for the treatment was 24 days shorter than that of the control group representing a 20 % decrease in average gestation length. Shorter gestation lengths may increase the female's survival as gravidity imposes energetic costs as well as reduced locomotory and foraging capabilities and requires increased amounts of time engaged in active thermoregulation – all of which may ultimately result in an increased exposure and risk of predation for the gravid female (Seigel et al., 1987; Charland and Gregory, 1995; Gregory et al., 1999). As these costs in Red-sided garter snakes are imposed simultaneously with the already extremely limited opportunities for summer foraging, a reduction in gestation length would allow for both increased amount of time to forage each year and more frequent prey capture during this period ultimately resulting in increased opportunity to replenish depleted energy stores after giving birth. Additionally, it has been shown that higher temperatures increase the ability to build mass during the summer season in adult females, allowing them to further make use of potentially expanded foraging in these hotter conditions to accelerate postpartum recovery (Halliday and Blouin-Demers, 2018).

However, it must be considered that any increase in foraging opportunity may be counterbalanced by potential decreases in prey availability, as increased temperatures will undoubtedly affect the whole ecosystem in ways that are difficult to predict. Additionally,

climate change results in more frequent and extreme temperature swings, which will likely result

in an increased occurrence of conditions that are outside the thermal range needed for normal

activity, potentially reducing the time available for foraging per day (Masson-Delmotte et al.,

2021). Despite this uncertainty, any increase in the time available for foraging will potentially

increase the ability to overcome the challenges that result from reductions to suitable foraging

conditions and prey availability. While warming conditions will likely alter seasonal timing,

resulting in some extension to the spring and fall that are available for activity, winter brumation

conditions are unlikely to change in a meaningful way under current climate change predictions,

as current mean temperatures are well below freezing with sustained snow coverage from late

October to mid-April each year (https://en.climate-data.org/north-

america/canada/manitoba/inwood-108217/). While the longest period of brumation is unlikely to

change, even a marginal shift that reduces brumation duration at the margins will add an

additional variable that can alter the impact of a reduced gestation length.

We found that, along with the reduced gestation length, the treatment condition also

resulted in both increased neonate mass and BCI at birth but did not affect SVL. This was most

pronounced when comparing neonates with larger SVL measurements, suggesting that control

neonates had a harder time building mass for longer neonates. Generally, larger neonate size at

birth is associated with both improved short term and lifetime survival in snakes (Bronikowski,

2000; Brown and Shine, 2004; Kissner and Weatherhead, 2005; Addis et al., 2017). Based on

this metric alone, the offspring gestated in warmer conditions may have increased overall fitness

resulting in better parental reproductive outcomes. However, there are potential trade-offs

associated with accelerated development that can have long-term consequences, such as

reductions in longevity, stress resistance, and future fecundity, any of which might overshadow

the benefits gained by their larger size at birth (Chippindale et al., 2004; Shalev and Belsky,

2016). Long term studies tracking lifetime outcomes would be necessary to fully determine if the

potential short-term survival benefits are outweighed by long-term costs of accelerated

development, however it takes up to 3 years to reach sexual maturity, and a lifespan in the wild

that can exceed 10 years, making this extremely logistically difficult to conduct, and far outside

the scope of this study (Gregory, 1977; Bronikowski, 2008; Rollings et al., 2017).

418   Interestingly we observed no differences in birth rates, average litter sizes, total litter

mass, the number of stillborn or severely deformed neonates between treatments. It has been

shown that increased temperatures during gestation can reach a threshold, after which improved

outcomes observed under moderate temperature increases begin to be overtaken by detrimental

outcomes for developing offspring or gestating mothers, such as reductions to body conditions of

the neonates, reduced litter sizes, and increased stillbirths (Beuchat, 1988; Yan et al., 2011; Tang

et al., 2012). During this study, by artificially increasing the minimum ambient temperatures

available to mothers with developing offspring, we prevented them from thermoregulating at a

temperature below such a threshold. This serves to simulate and evaluate the effect of general

warming in an environment with thermal conditions that are lower than preferred body

temperatures. Under these conditions, our observations suggest improved reproductive outcomes

at the time of birth, with no directly observed negative results associated with the treatment

condition. Thus, we conclude that an increase of 5 ºC to the minimum ambient temperatures

during gestation likely did not exceed the threshold at which detrimental outcomes overcome the

benefits gained. Generally, this suggests that there is a potential for additional beneficial effects

from an increased minimum ambient temperature beyond 5 ⁰C during gestation, as the benefits

presented here suggest that they have not yet reached their detrimental thermal threshold.

Though it is possible that some benefits may be present with a more moderate increase in

temperature, such as increased litter size or birthrates, and may have already been lost to some

degree at this higher temperature. Further research exploring a wider range of thermal conditions

during gestation would be interesting, as this would expand our perspective allowing for

increased ability to predict the effects of future increases in ambient temperatures and would

potentially pinpoint the maximum ambient temperature threshold at which beneficial results

began to reverse and detrimental impacts begin to arise.

While our experimental design increased both minimum diurnal and minimum nocturnal

temperatures by 5 ⁰C, we believe that the treatment effect was driven primarily by the increased

nocturnal minimum temperature. In fact, the large thermal differences experienced by

populations at different geographic regions are likely overcome using behavioral plasticity, as

they exhibit the same diurnal thermal preference regardless of large differences in latitude and

climate (Rosen, 1991). During the day, gravid females were allowed to use basking to

behaviorally thermoregulate, mimicking behavior exhibited in natural settings, however the

minimum temperature experienced during the night cannot be overcome using this strategy. This

should result in relatively little difference in body temperature during the day but would allow

for body temperatures to remain closer to optimal gestation temperatures for the treatment group

at night. Because higher maternal body temperatures accelerate embryonic development, a

higher minimum temperature experienced during the night has the potential to increase the rate

of development at the time it is the slowest (Beuchat, 1988; Lorioux et al., 2013). Further studies

looking specifically at increased nocturnal temperatures in this system would be valuable in

deciphering the impact of this increasingly relevant change in thermal conditions (Rutschmann et

al., 2023)

This study did not identify that critical threshold, but these results suggest interesting

implications for future warming associated with continued climate change. The response

variables examined here were limited and did not directly address or account for other

phenotypic affects that may have arisen from increased temperatures during gestation. However,

evidence suggests that many other developmental processes may be influenced by altered

thermal conditions, and increased temperatures may result in changes to offspring phenotype

such as differences in scale and bone development (Arnold and Peterson, 2002; Lourdais et al.,

2004). Additionally, the impacts of temperature during gestation on offspring traits such as the

aforementioned changes to morphology, body size, and body condition are frequently associated

with both beneficial and detrimental long-lasting impacts on survival after birth (Bronikowski,

2000; Brown and Shine, 2004; Kissner and Weatherhead, 2005; Noble et al., 2018).

Current climate change models predict future global average temperature increases of up

to 5.3 ºC by the end of the century (Houghton et al., 2001; Masson-Delmotte et al., 2021). Thus,

the results of this experiment show that an increase in average temperature of this degree,

specifically during the brief yet crucial opportunity for feeding and gestation, is not likely to

cause direct detrimental effects on the reproductive outcomes measured in this study for northern

populations of Red-sided garter snakes and have the potential to improve neonate fitness. In fact,

because this species spans much of North America and different populations experience a wide

range of thermal conditions, the populations at the colder high latitudes are likely able to survive

much warmer conditions similar to those at the southern edge of their range (Fitch, 1965;

Aleksiuk and Stewart 1971).

However, the extended impacts of increasing temperature are likely to result in other

adverse effects on this population. Thermal stress caused by exposure to increased temperature

extremes (acute high or low temperatures) is known to result in a number of detrimental effects

including misaligned behavioral timing, altered development, reduced or altered activity, and

higher mortality rates (Houghton et al., 2001; Jentsch et al., 2009; Paaijmans et al., 2013; Stott,

2016). The greater thermal instability associated with climate change conditions may also lead to

thermally induced alterations to energy use and metabolic rate (Friesen et al., 2015; Wilson,

2020; Holden et al., 2021). The combination of physiological and environmental changes due to

increased environmental temperatures are also known to increase pathogen exposure and

increase individual susceptibility to infection (Walther et al., 2001; Harvell et al., 2009; Gallana

et al., 2013; Nnadi and Carter, 2021). Additionally, many species have been observed to alter

seasonal migration patterns and shift territorial ranges to mitigate changes to their traditional

ecosystems, ultimately resulting in disruption of ecosystems arising from novel interspecific

competition and alterations to the availability of prey species or trophic effects (Parmesan, 2006;

Gilman et al., 2010; Urban et al., 2012; Salinas-Ramos et al., 2020). Although we observed

effects of increased temperatures that appeared likely beneficial to offspring, it is possible that

that the lifetime reproductive potential of mothers would in fact be reduced as a result of long-

term exposure to increased temperatures or repeated bouts of rapid gestation. Thus, future

research regarding changes to ambient temperatures during gestation may benefit from

examining both the immediate and long-term health impacts of mothers, such as impacts on body

condition, future fecundity, parasite load, and subsequent winter survival, in addition to the long-

term survival and impacts on the physiology and long-term reproductive potential of offspring.

Higher temperatures caused by future warming have the potential to change the shape of

this population, resulting in offspring that not only develop faster, but are born larger. In the

present study, we observed no apparent tradeoff or cost to the overall number of offspring in

each litter. This has the potential to dramatically increase the survival rates of future generations,

while increasing the time each year spent foraging for both the mothers and the resulting

offspring, which may serve to ameliorate potential decreased foraging success caused by reduced

prey availability or suitable foraging temperatures (Seigel et al., 1987; Charland and Gregory,

1995; Brown and Shine, 2004; Kissner and Weatherhead, 2005).

The myriad impacts of climate change on an ecosystem are overwhelmingly complicated

and interconnected. Thus, it is exceedingly difficult to develop models that accurately predict

comprehensive, species-specific outcomes in response to altered thermal landscapes. The present

study examined the effect of altered minimum temperatures during gestation within a single

large population and thus represents only a single piece of the complex overall puzzle. As global

temperatures continue to rise, additional research examining the effects of increasing temperature

on the behavior and physiology of individual species, as well as studies encompassing entire

ecosystems, will provide the invaluable insight necessary to further our understanding of how

temperature affects ecosystems and, more importantly, will enhance our ability to predict the

impacts of future climate change on the global ecosystem.

5. Conclusions

Our research demonstrated that, even under a worst-case scenario climate change model prediction, increased minimum temperatures may lead to improved reproductive outcomes for terrestrial ectotherms living at the northern edge of their range. Reduced gestation length has the potential to facilitate increased fitness for both mother and offspring, and while further work is needed to determine the long-term consequences of this altered gestation, this study found no observed cost to reproductive outcomes.

References

ADDIS, E. A., E. J. GANGLOFF, M. G. PALACIOS, K. E. CARR, AND A. M. BRONIKOWSKI. 2017. Merging the "morphology–performance–fitness" paradigm and life-history theory in the Eagle Lake garter snake research project. Integrative and Comparative Biology 57:423-435.

ALEKSIUK, M. 1976. Reptilian hibernation: evidence of adaptive strategies in *Thamnophis sirtalis parietalis*. Copeia 12:170-178.

ALEKSIUK, M. AND P. T. GREGORY. 1974. Regulation of seasonal mating behavior in *Thamnophis sirtalis parietalis*. Copeia 18:681-689.

ALEKSIUK, M. AND K. W. STEWART. 1971. Seasonal changes in the body composition of the garter snake (*Thamnophis sirtalis parietalis*) at northern latitudes. Ecology 52:485-490.

ALEXANDER, L.V., ZHANG, X., PETERSON, T.C., CAESAR, J., GLEASON, B., KLEIN TANK, A.M.G., HAYLOCK, M., COLLINS, D., TREWIN, B., RAHIMZADEH, F. AND A. TAGIPOUR. 2006.

Global observed changes in daily climate extremes of temperature and precipitation.

Journal of Geophysical Research: Atmospheres, *111*(D5).

ARNOLD, S. J., C. R. PETERSON. 2002. A model for optimal norms: the case of the pregnant garter

snake and her temperature sensitive embryos. American Naturalist 160:306-616.

BATES, D., MÄCHLER, M., BOLKER, B. M., AND S. C. WALKER. 2015. Fitting linear mixed-effects

models using lme4. *Journal of Statistical Software*, *67*(1).

BEUCHAT, C. 1988. Temperature effects during gestation in a viviparous lizard. Journal of

Thermal Biology 13:135-142.

BLANCHARD, F. N. AND F. C. BLANCHARD. 1940. Factors determining time of birth in the garter

snake *Thamnophis sirtalis sirtalis* (Linnaeus). Papers from the Michigan Academy of

Science, Arts and Letters 26:161–176.

BLAKEMORE, L.A., 2017. Sex and Survival: Reproduction and anti-microbial defense in the Red-

sided garter snake (*Thamnophis sirtalis parietalis*). Oregon State University.

BROMMER, J. E., J. MERILÄ, AND H. KOKKO. 2002. Reproductive timing and individual fitness.

Ecology Letters 5:802-810.

BRONIKOWSKI, A. M. 2000. Experimental evidence for the adaptive evolution of growth rate in

the garter snake *Thamnophis elegans*. Evolution 54:1760-1767.

BRONIKOWSKI, A.M. 2008. The evolution of aging phenotypes in snakes: a review and synthesis

with new data. *Age*, *30*(2), pp.169-176.

BROWN, G. P. AND R. SHINE. 2004. Maternal nest-site choice and offspring fitness in a tropical

snake (*Tropidonophis mairii, Colubridae*). Ecology 85:1627-1634.

BUSH, E AND D. S. LEMMEN. 2019. Canada's Changing Climate Report. Government of Canada

Press, Canada.

CHARLAND, M. B. AND P. T. GREGORY. 1995. Movements and habitat use in gravid and

nongravid female garter snakes (Colubridae: *Thamnophis*). Journal of Zoology 236:543-

561.

Climate-data.org. Inwood climate: Weather Inwood & temperature by month. Available at

https://en.climate-data.org/north-america/canada/manitoba/inwood-108217/

COSTANZO, J. P. 1985. The bioenergetics of hibernation in the Eastern garter snake *Thamnophis*

sirtalis sirtalis. Physiological Zoology 58:682-692.

CREWS, D., M. GRASSMAN, W. GARSTKA, A. HALPERT, AND B. CAMAZINE. 1987. Sex and

seasonal differences in metabolism in the Red-sided garter snake, *Thamnophis sirtalis*

parietalis. Canadian Journal of Zoology 65:2362-2368.

DAYGER, C.A., LEMASTER, M.P. AND D.I. LUTTERSCHMIDT. 2018. Physiological correlates of

reproductive decisions: relationships among body condition, reproductive status, and the

hypothalamus-pituitary-adrenal axis in a reptile. *Hormones and behavior*, *100*, pp.1-11.

DOODY, J.S. AND J.M. REFSNIDER. 2022. Nesting in reptiles: Natural and anthropogenic threats

and evolutionary responses. Frontiers in Ecology and Evolution 10:1103193.

DU, W. G. AND R. SHINE. 2015. The behavioural and physiological strategies of bird and reptile

embryos in response to unpredictable variation in nest temperature. Biological Reviews

90:19-30.

ENVIRONMENT CANADA. 2014. Canada's sixth national report on climate change 2014 : actions to

meet commitments under the United Nations Framework Convention on Climate Change.

Pp. 120-121. Environment Canada, Canada.

FITCH, H.S. 1965. An ecological study of the garter snake, *Thamnophis sirtalis*. *Univ Kansas*

Publ Mus Nat Hist, *15*, pp.493-564.

FRIESEN, C. R., R. SHINE, R. W. KROHMER, AND R. T. MASON. 2013. Not just a chastity belt: the

functional significance of mating plugs in garter snakes, revisited. Biological Journal of

the Linnean Society 109:893-907.

FRIESEN, C. R., R. SHINE, R. W. KROHMER, AND R. T. MASON. 2013. Not just a chastity belt: the

functional significance of mating plugs in garter snakes, revisited. Biological Journal of

the Linnean Society 109:893-907.

FRIESEN, C. R., M. K. SQUIRE, AND R. T. MASON. 2014. Intrapopulational variation of ejaculate

traits and sperm depletion in Red-sided garter snakes. Journal of Zoology 292:192-201.

FRIESEN, C.R., MASON, R.T., ARNOLD, S.J. AND S. ESTES. 2014. Patterns of sperm use in two

populations of Red-sided Garter Snake (*Thamnophis sirtalis parietalis*) with long-term

female sperm storage. *Canadian Journal of Zoology*, *92*(1), pp.33-40.

FRIESEN, C. R., D. R. POWERS, P. E. COPENHAVER, AND R. T. MASON. 2015. Size dependence in

non-sperm ejaculate production is reflected in daily energy expenditure and resting

metabolic rate. Journal of Experimental Biology 218:1410-1418.

GALLANA, M., M. P. RYSER-DEGIORGIS, T. WAHLI, AND H. SEGNER. 2013. Climate change and

infectious diseases of wildlife: altered interactions between pathogens, vectors and hosts.

Current Zoology 59:427-437.

GANS, C. 1996. An overview of parental care among the Reptilia. Advances in the Study of

Behavior. 25:145-157.

GAO, J. F., Y. F. QU, L. G. LUO, AND X. JI. 2010. Evolution of reptilian viviparity: a test of the

maternal manipulation hypothesis in a temperate snake, *Gloydius brevicaudus*

(*Viperidae*). Zoological Science 27:248-255.

GEISER, F. 2013. Hibernation. Current Biology 23:188-193.

GIBSON, A. R. AND J. B. FALLS. 1979. Thermal biology of the common garter snake *Thamnophis*

sirtalis (L.). I. Temporal variation, environmental effects and sex differences. Oecologia

1:79-97

GILMAN, S. E., M. C. URBAN, J. TEWKSBURY, G. W. GILCHRIST, AND R. D. HOLT. 2010. A

framework for community interactions under climate change. Trends in Ecology &

Evolution 25:325-331.

GREGORY, P. T. 1974. Patterns of spring emergence of the Red-sided garter snake (*Thamnophis*

sirtalis parietalis) in the Interlake region of Manitoba. Canadian Journal of Zoology

52:1063-1069.

GREGORY PT. 1977. Life history parameters of the red-sided garter snake *(Thamnophis sirtalis*

pairetalis) in an extreme environment, the Interlake region of Manitoba Canada, pp. 1–

44. Ottawa, Canada: National Museum of Canada.

GREGORY, P. T. 1982. Reptilian hibernation. 1982. Biology of the Reptilia. Pp. 53-154.

Academic Press, USA.

GREGORY, P. T., L. H. CRAMPTON, AND K. M. SKEBO. 1999. Conflicts and interactions among

reproduction, thermoregulation and feeding in viviparous reptiles: are gravid snakes

anorexic? Journal of Zoology 248:231-241.

GUNDERSON, A.R., M.E. DILLON, AND J.H. STILLMAN. 2017. Estimating the benefits of plasticity

in ectotherm heat tolerance under natural thermal variability. Functional Ecology

31:1529-1539.

HALLIDAY, W.D. AND G. BLOUIN-DEMERS. 2018. Body temperature influences growth rates of

Common Gartersnakes (Thamnophis sirtalis). *The Canadian Field-Naturalist*, *132*(1),

pp.25-29.

HARVELL, D., S. ALTIZER, I. M. CATTADORI, L. HARRINGTON, AND E. WEIL. 2009. Climate

change and wildlife diseases: when does the host matter the most? Ecology 90:912-920.

HOLDEN, K. G., E. J. GANGLOFF, E. GOMEZ-MANCILLAS, K. HAGERTY, AND A. M. BRONIKOWSKI.

2021. Surviving winter: Physiological regulation of energy balance in a temperate

ectotherm entering and exiting brumation. General and Comparative Endocrinology

307:113758.

HOUGHTON, T., Y. DING, J. GRIGGS, N. NOGUER, P. VAN DER LINDEN, D. XIAOSU, K. MASKELL

AND A. JOHNSON. 2001. Climate change 2001: The scientific basis. Contribution of

Working Group I to the Third Assessment Report of the Intergovernmental Panel on

Climate Change. Quarterly Journal of the Royal Meteorological Society 128:1038-1039.

HUANG, W.S. AND D.A. PIKE. 2013. Testing cost-benefit models of parental care evolution using

lizard populations differing in the expression of maternal care. PloS one 8:e54065.

HUEY, R. B. AND R. D. STEVENSON. 1979. Integrating thermal physiology and ecology of

ectotherms: a discussion of approaches. American Zoologist 19:357-366.

HUEY, R. B. 1982. Temperature, physiology, and the ecology of reptiles. *Biology of the Reptilia,*

Physiology. Gans C and Pough FH (*Eds*.). London, Academic Press 12:25-91.

HUEY, R.B. AND D. BERRIGAN. 2001. Temperature, demography, and ectotherm fitness. The

American Naturalist 158:204-210.

HUGHES, D. F., P. R. DELIS, AND W. E. MESHAKA. 2019. Comparison of body temperatures

across physiological states in syntopic snake species (*Thamnophis sirtalis* and *Nerodia*

sipedon) from Pennsylvania. Northeastern Naturalist 26:749-760.

JAYNE, B. C. AND A. F. BENNETT. 1990. Selection on locomotor performance capacity in a

natural population of garter snakes. Evolution 44:1204-1229.

JENTSCH, A., J. KREYLING, J. E. BOETTCHER-TRESCHKOW, AND C. BEIERKUHNLEIN. 2009.

Beyond gradual warming: extreme weather events alter flower phenology of European

grassland and heath species. Global Change Biology 15:837-849.

JI, X., L. H. LIN, L. G. LUO, H. L. LU, J. F. GAO, AND J. HAN. 2006. Gestation temperature affects

658   sexual phenotype, morphology, locomotor performance, and growth of neonatal Brown

659   forest skinks, *Sphenomorphus indicus*. Biological Journal of the Linnean Society 88:453-

660   463.

661   KING, R. B. 2022. Temperature-induced multi-species cohort effects in sympatric snakes.

662   Ecology and Evolution 12:e8601.

663   KINGSOLVER, J. G., S. E. DIAMOND, AND L. B. BUCKLEY. 2013. Heat stress and the fitness

664   consequences of climate change for terrestrial ectotherms. Functional Ecology 27: 1415-

665   1423.

666   KISSNER, K. J. AND P. J. WEATHERHEAD. 2005. Phenotypic effects on survival of neonatal

667   Northern watersnakes *Nerodia sipedon*. Journal of Animal Ecology 74:259-265.

668   LADYMAN, M., X. BONNET, O. LOURDAIS, D. BRADSHAW, AND G. NAULLEAU. 2003. Gestation,

669   thermoregulation, and metabolism in a viviparous snake, *Vipera aspis*: evidence for

670   fecundity-independent costs. Physiological and Biochemical Zoology 76:497-510.

671   LORIOUX, S., H. LISSE, AND O. LOURDAIS. 2013. Dedicated mothers: predation risk and physical

672   burden do not alter thermoregulatory behaviour of pregnant vipers. Animal Behaviour

673   86:401-408.

674   LORIOUX, S., VAUGOYEAU, M., DENARDO, D.F., CLOBERT, J., GUILLON, M. AND O. LOURDAIS.

675   2013. Stage dependence of phenotypical and phenological maternal effects: insight into

676   squamate reptile reproductive strategies. *The American Naturalist*, *182*(2), pp.223-233.

LOURDAIS, O., X. BONNET, R. SHINE, D. DeNARDO, G. NAULLEAU, AND M. GUILLON. 2002.

Capital-breeding and reproductive effort in a variable environment: a longitudinal study

of a viviparous snake. Journal of Animal Ecology 71:470-479.

LOURDAIS, O., R. SHINE, X. BONNET, M. GUILLON AND G. NAULLEAU. 2004. Climate affects

embryonic development in a viviparous snake *Vipera aspis*. Oikos 104:551-560.

LOURDAIS, O., M. GUILLON, D. DeNARDO, AND G. BLOUIN-DEMERS. 2013. Cold climate

specialization: adaptive covariation between metabolic rate and thermoregulation in

pregnant vipers. Physiology & Behavior 119:149-155.

LUTTERSCHMIDT, D. I., M. P. LEMASTER, AND R. T. MASON. 2006. Minimal overwintering

temperatures of Red-sided garter snakes (*Thamnophis sirtalis parietalis*): a possible cue

for emergence? Canadian Journal of Zoology 84:771-777.

MADSEN, T. AND R. SHINE. 2002. Short and chubby or long and slim? food intake, growth and

body condition in free-ranging pythons. Austral Ecology 27:672-680.

MASSON-DELMOTTE, V., P. ZHAI, A. PIRANI, S.L. CONNORS, C. PÉAN, S. BERGER, N. CAUD, Y.

CHEN, L. GOLDFARB, M. I. GOMIS, AND M. HUANG. 2021 Climate change 2021: the

physical science basis. Contribution of working group I to the sixth assessment report of

the intergovernmental panel on climate change. Cambridge University Press, UK.

MATHIES, T. AND R. M. ANDREWS. 1997. Influence of pregnancy on the thermal biology of the

lizard, *Sceloporus jarrovi*: why do pregnant females exhibit low body temperatures?

Functional Ecology 11:498-507.

MICHEL, C. L., J. H. PASTORE, AND X. BONNET. 2013. Impact of cool versus warm temperatures

on gestation in the Aspic viper (*Vipera aspis*). Comparative Biochemistry and Physiology

Part A: Molecular & Integrative Physiology 165:338-342.

MILES, D. B., B. SINERVO, AND W. A. FRANKINO. 2000. Reproductive burden, locomotor

performance, and the cost of reproduction in free-ranging lizards. Evolution 54:1386-

1395.

NNADI, N. E. AND D. A. CARTER. 2021. Climate change and the emergence of fungal pathogens.

PLoS Pathogens 17:e1009503.

NOBLE, D.W., STENHOUSE, V., AND L.E. SCHWANZ. 2018. Developmental temperatures and

phenotypic plasticity in reptiles: A systematic review and meta-analysis. *Biological*

Reviews, *93*(1), pp.72-97.

NOER, N. K., M. ØRSTED, M. SCHIFFER, A. A. HOFFMANN, S. BAHRNDORFF, AND T. N.

KRISTENSEN. 2022. Into the wild—a field study on the evolutionary and ecological

importance of thermal plasticity in ectotherms across temperate and tropical regions.

Philosophical Transactions of the Royal Society B 377:20210004.

O'DONNELL, R., R. SHINE, AND R. T. MASON. 2004. Seasonal anorexia in the male Red-sided

garter snake, *Thamnophis sirtalis parietalis*. Behavioral Ecology and Sociobiology

56:413-419.

PAAIJMANS, K. P., R. L. HEINIG, R. A. SELIGA, J. I. BLANFORD, S. BLANFORD, C. C. MURDOCK,

AND M. B. THOMAS. 2013. Temperature variation makes ectotherms more sensitive to

climate change. Global Change Biology 19:2373-2380.

PARMESAN, C. 2006. Ecological and evolutionary response to recent climate change. Annual

Review of Ecology, Evolution, and Systematics 37:637-669.

RABAIOTTI, D., T. COULSON, AND R. WOODROFFE. 2023. Climate change is predicted to cause

population collapse in a cooperative breeder. Global Change Biology 29:6002-6017.

ROLLINGS, N., UHRIG, E.J., KROHMER, R.W., WAYE, H.L., MASON, R.T., OLSSON, M.,

WHITTINGTON, C.M. AND C.R. FRIESEN. 2017. Age-related sex differences in body

condition and telomere dynamics of red-sided garter snakes. *Proceedings of the Royal*

Society B: Biological Sciences, *284*(1852), p.20162146.

ROSEN, P.C. 1991. Comparative field study of thermal preferenda in garter snakes (*Thamnophis*).

Journal of Herpetology, pp.301-312.

RUTSCHMANN, A., PERRY, C., LE GALLIARD, J.F., DUPOUÉ, A., LOURDAIS, O., GUILLON, M.,

BRUSCH IV, G., COTE, J., RICHARD, M., CLOBERT, J., AND D.B. MILES. 2024. Ecological

responses of squamate reptiles to nocturnal warming. *Biological Reviews*, *99*(2), pp.598-

621.

SALINAS-RAMOS, V. B., L. ANCILLOTTO, L. BOSSO, V. SÁNCHEZ-CORDERO, AND D. RUSSO. 2020.

Interspecific competition in bats: state of knowledge and research challenges. Mammal

Review 50:68-81.

SCHIELZETH, H., DINGEMANSE, N.J., NAKAGAWA, S., WESTNEAT, D.F., ALLEGUE, H., TEPLITSKY,

C., RÉALE, D., DOCHTERMANN, N.A., GARAMSZEGI, L.Z. AND Y.G. ARAYA-AJOY. 2020.

Robustness of linear mixed-effects models to violations of distributional assumptions.

Methods in ecology and evolution, *11*(9), pp.1141-1152.

SEIGEL, R. A., M. M. HUGGINS, AND N. B. FORD. 1987. Reduction in locomotor ability as a cost

of reproduction in gravid snakes. Oecologia 73:481-485.

SHALEV, I. AND BELSKY, J., 2016. Early-life stress and reproductive cost: A two-hit

developmental model of accelerated aging? *Medical Hypotheses*, *90*, pp.41-47.

SHERMAN, P. W. AND M. C. RUNGE. 2002. Demography of a population collapse: the northern

Idaho ground squirrel (*Spermophilus brunneus brunneus*). Ecology 83:2816–2831.

SHINE, R. 1988. Parental care in reptiles. Biology of the Reptilia, Ecology B: defense and life

history 16:275-329.

SHINE, R., T. LANGKILDE, M. WALL, AND R. T. MASON. 2006. Temporal dynamics of emergence

and dispersal of garter snakes from a communal den in Manitoba. Wildlife Research

33:103–111.

STOTT, P. 2016. How climate change affects extreme weather events. Science 352:1517-1518.

TANG, X., F. YUE, X. YAN, D. ZHANG, Y. XIN, C. WANG, AND Q. CHEN. 2012. Effects of gestation

temperature on offspring sex and maternal reproduction in a viviparous lizard (*Eremias*

multiocellata) living at high altitude. Journal of Thermal Biology 37:438-444.

URBAN, M. C., J. J. TEWKSBURY, AND K. S. SHELDON. 2012. On a collision course: competition

and dispersal differences create no-analogue communities and cause extinctions during

climate change. Proceedings of the Royal Society B: Biological Sciences 279:2072-2080.

WANG, Z., L. MA, M. SHAO, AND X. JI. 2017. Are viviparous lizards more vulnerable to climate

warming because they have evolved reduced body temperature and heat tolerance?

Oecologia 185:573-582.

WALTHER, G. R., C. A. BURGA, P. J. EDWARDS, AND P. J. EDWARDS (Eds.) 2001. "Fingerprints"

of climate change: adapted behaviour and shifting species. Springer Science & Business

Media

WHITTIER, J.M., R. T. MASON, AND D. CREWS. 1985. Mating in the red-sided garter snake,

Thamnophis sirtalis parietalis: differential effects on male and female sexual behavior.

Behavioral Ecology and Sociobiology 16:257-261.

WICKHAM, H. 2016. ggplot2 Elegant Graphics for Data Analysis (Use R!). *Springer*.

WILSON, R. C. 2020. Energy Stores and Life-history Transitions in Red-Sided Garter Snakes

(*Thamnophis sirtalis parietalis*). Doctoral dissertation, Portland State University, USA.

YAN, X. F., X. L. TANG, F. YUE, D. J. ZHANG, Y. XIN, C. WANG, AND Q. CHEN. 2011. Influence of

ambient temperature on maternal thermoregulation and neonate phenotypes in a

viviparous lizard, *Eremias multiocellata*, during the gestation period. Journal of Thermal

Biology 36:187-192.

ZHANG, D. J., X. L. TANG, F. YUE, Z. CHEN, AND Q. CHEN. 2010. Effect of gestation temperature

on sexual and morphological phenotypes of offspring in a viviparous lizard, *Eremias*

multiocellata. Journal of Thermal Biology 35:129-133.

ZHANG, X., L. A. VINCENT, W. D. HOGG AND A. NIITSOO. 2000. Temperature and precipitation

trends in Canada during the 20th century. Atmosphere-ocean 38:395-429.

Figure 1. Differences in reproductive outcomes for snakes held at different minimum
temperatures during gestation. Treatment condition represents snakes housed a constant 5 ˚C
higher minimum temperature relative to control for the duration of gestation. Boxes indicate the
median (horizontal line in box), the first (Q1) and third (Q3) quartiles (bottom and top of box)
and mean (blue diamond), whiskers represent the largest or smallest value within 1.5 x
interquartile range (IQR). Barplot indicates the proportion of mated females that reached
parturition for each condition. Statistical results from linear mixed-effects models (A-D), linear
models (E-K), and a two proportions z-test (L). Gestation length was significantly reduced (A)
and neonate mass and body condition (BCI) were significantly higher in the treatment condition
(B and D respectively). There was no difference in neonate snout-vent length (SVL) between
conditions (C). There was no significant effect caused by treatment at the litter level for total
litter mass (E), number of stillbirths (F), number of developmental deformities (G), litter size
(H), or average (mean) neonate SVL per litter (I). Both average (mean) neonate mass per litter
(J) and average (mean) neonate BCI per litter (K) were significantly higher in the treatment
condition. There was no significant difference in the birthrates (L) between treatments.

Statistical significance is indicated as: ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, and ns ($p > 0.05$).

Figure 2. Relationship between neonate traits. Lines indicate liner regression model fits, color

coded by treatment, with the shaded areas representing the standard error. R^2 is displayed for

each model, by treatment. Neonate snout-vent length (SVL) was weakly related to body

condition index (BCI) in both treatment conditions, with a positive relationship in the treatment

conditions and a negative relationship n the control (A). A strong positive relationship between

neonate SVL and neonate mass for both treatment conditions (B). A positive relationship was

found between neonate mass and BCI for both treatment conditions, however this was weakly

related for the control conditon and strongly related in the treatment (C). See Supplemental Table

4 for full statistical results.

808 Tables

809 **Table 1.** The effect of treatment on neonate traits at birth

Trait (Y_i)	Intercept (β_0)	Estimate (β_1)	Std. Error	df	t-value	p-value	Var. (b)	Std. Dev. (b)	Var. (ε)	Std. Dev. (ε)
Mass (g)	1.36	0.19	0.08	24.08	2.30	**0.0302**	0.044	0.210	0.009	0.095
SVL (cm)	15.61	0.19	0.34	23.79	0.57	0.574	0.721	0.850	0.203	0.450
BCI	-0.08	0.16	0.05	24.02	2.93	**0.0073**	0.017	0.132	0.008	0.088
Gestation (days)	42.16	-24.02	2.55	24.00	-9.40	**1.61 x 10^{-9}**	42.2	6.50	0.036	0.189

811 Results from linear mixed-effects models estimating the effect of the treatment condition on

812 neonate traits. Bold values indicate a significant impact on the trait caused by the treatment

813 condition.

815 **Table 2.** The effect of treatment on traits of neonates at birth by litter

Trait (Y_i)	Intercept (β_0)	Estimate (β_1)	Std. Error	df	t-value	p-value
Total litter mass (g)	20.70	6.42	4.07	24	1.58	0.128
Average neonate mass (g)	1.36	0.19	0.08	24	2.29	**0.0314**
Number of stillborn	1.86	-1.61	1.13	24	-1.42	0.1678
Number of deformed	0.07	0.01	0.11	24	0.11	0.914
Litter size	15.86	1.89	3.25	24	0.58	0.566
Average neonate SVL (cm)	15.64	0.17	0.34	24	0.50	0.621
Average neonate BCI	-0.08	0.16	0.05	24	2.99	**0.00635**

817 Results from linear models estimating the effect of the treatment condition on litter wide neonate

818 traits. Bold values indicate a significant impact on the trait caused by the treatment condition.