Floral scent emission of *Epiphyllum oxypetalum*: identification of a novel cytosol-localized geraniol biosynthesis pathway

Yiyang Zhang [1, *], Yuhan Zhang [1, *], Qiurui Tian [1], Likun Wei [1], Ting Zhu [1], Zhiwei Zhou [1], Jiaqi Wang [1], Zhibin Liu [1], Wei Tang [2], Haijun Xiao [1], Mingchun Liu [1], Tao Li [1], Qun Sun [1]

[1] Key Laboratory of Bio-resources and Eco-environment Ministry of the Education, College of Life Sciences, Sichuan University, Chengdu, Sichuan 610064, PR China

[2] Sichuan Academy of Botanical Engineering, Sichuan Academy of Agricultural Sciences, Zizhong, Sichuan 641200, PR China

[*] These authors contributed equally to this work

Authors for correspondence:

Qun Sun: qunsun@scu.edu.cn

Tao Li: tao.li@scu.edu.cn

Total word count (excluding summary, References and legends):	5085	No. of Figures:	7 (Fig. 1-7 in color)
Summary	188	No. of Tables:	0
Introduction	740	No. of Supporting Information files:	11 (Fig. S1; Table S1-S9, Video S1)
Materials and methods	1508		
Results	1633		
Discussion	1204		
Acknowledgements:	89		

ORCID：

Qun Sun, https://orcid.org/0000-0002-4372-8865

Tao Li, https:/orcid.org/0000-0003-1462-5135

Yiyang Zhang, https://orcid.org/0000-0002-2159-6451

Yuhan Zhang, https://orcid.org/0009-0009-3562-2452

Qiurui Tian, https://orcid.org/0009-0002-3782-5594

Haijun Xiao, https://orcid.org/0000-0002-9587-8101

Zhibin Liu, https://orcid.org/0000-0003-2053-5136

Mingchun Liu, https://orcid.org/0000-0001-8004-1758

Summary

•   *Epiphyllum oxypetalum*, a renowned ornamental species in Cactaceae, releases attractive

fragrance during its infrequent, transient and nocturnal flowering, the nature and biosynthesis

of the volatiles for this floral scent, however, remained unexplored.

•   Employing volatilomic, transcriptomic and biochemical approaches, we systematically

characterized the composition, emission dynamics and biosynthesis of *E. oxypetalum* floral

scent.

•   Floral scent composition was highly dynamic, with trans-geraniol comprising 72.54% of the

total emission at full bloom, followed by benzyl alcohol (12.96%) and methyl salicylate

(3.75%), which predominantly emanated from petals and sepals. Transcriptomic analysis and

inhibition assay using pathway-specific inhibitors revealed the mevalonate (MVA) pathway

as the precursor source for trans-geraniol biosynthesis. Together with elevated expression of

both cytosol-localized geraniol pyrophosphate synthase EoGDPS and geraniol synthase

EoTPSa1, we elucidated an unusual cytosolic biosynthesis route for geraniol in *E.*

oxypetalum petals.

•   Our study on *E. oxypetalum* scent emission and its biosynthesis offers a comprehensive

profile of cactus floral scent profiles and presents a rare case of cytosolic geraniol

biosynthesis using MVA pathway-derived substrates. These findings underline the unique

metabolism of cactus flower volatiles, which hold a promise to be applied in the

development of novel perfumes.

Key words: floral scent, VOCs, emission dynamics, transcriptome, terpene biosynthesis, MVA

pathway.

Introduction

Angiosperm flowers emit diverse and complex fragrance of volatile organic compounds

(VOCs) to attract pollinators and repel florivores (Dudareva *et al.*, 2013; Dötterl & Gershenzon,

2023). The composition and emission patterns of floral VOCs exhibit considerable spatial and

temporal variability (Zhang *et al.*, 2022), creating distinct and characteristic fragrance. Species of

Cactaceae, often cultivated as ornamental plants, emit diverse floral scents with pollinator-related

chemical composition and temporal patterns. Usually, bee- or bird-pollinated cacti have scentless

diurnal flowers, while nocturnal-blooming cacti emit a contrasting spectrum of volatiles. The

bat-pollinated flowers usually produce sulfur-containing VOCs, causing unpleasant cabbage-like

odor. Whereas moth-pollinated ornamental species usually emit complex blends of terpenoids and

benzenoids volatiles during their transient nocturnal flowering, creating attractive cactus scent

(Kaiser & Tollsten, 1995). *E. oxypetalum*, a popular nocturnal flowering ornamental species from

Cactaceae, emits a strong and pleasant scent that last only for a few hours during nocturnal

blooming. During its explosive floral VOC emission, *E. oxypetalum* starts with a slightly phenolic

odor followed by a strong fruity scent. Previous studies have identified several monoterpene

volatiles from moth-pollinating cacti, some of which have been widely used in fragrance

industries for their fruity scent (Chen & Viljoen, 2010). However, these sporadic reports usually

neglected the highly dynamic composition and biosynthetic origin of cactus scent, limiting our

understanding of this highly specialized taxa.

Terpenoid volatiles contribute greatly to floral scent diversity (Abbas *et al.*, 2017). Despite

their great structural possibility, all terpenoid volatiles are directly derived from common

iso-pentyl diphosphate precursors, which are condensed from the common C_5 building blocks

isopentyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) (Muhlemann *et al.*, 2014).

Two alternative and compartmentally separated pathways synthesize IPP and DMAPP in plants.

The mevalonate (MVA) pathway, located in the cytosol and peroxisome, synthesizes these C_5

building blocks from acetyl-CoA, while the plastidic methylerythritol phosphate (MEP) pathway

synthesizes IPP and DMAPP from pyruvic acid and glyceraldehyde triphosphate. It has been long

accepted that these two pathways preferentially provide substrate for different classes of

terpenoids (Dudareva *et al.*, 2005). In general, the MVA pathway provides substrate for

sesquiterpene synthesis, while monoterpene synthesis utilizes MEP-derived precursors in plastids.

Recent work suggests that these two pathways are connected by species- and tissue-specific

metabolic cross-talk (Vranová *et al.*, 2012), providing additional possibilities for terpenoid

metabolism.

The monoterpene geraniol is widely distributed in floral scent, and it has been intensively

applied in cosmetic and fragrance industries for its sweet rouse-like scent. Specifically, geraniol is

synthesized from geranyl pyrophosphate (GPP), the direct product of IPP and DMAPP

condensation. Two different routes have been proposed for geraniol formation in previous studies.

Usually, a plastid-localized geraniol synthase (GES) from the canonical terpene synthase (TPS)

family (Chen *et al.*, 2011) catalyzes the formation of geraniol utilizing the plastid MEP derived

GPP (Dudareva *et al.*, 2005) via carbocation mechanism (Christianson, 2017) (MEP-TPS case).

This case was first established in snapdragon flowers (Dudareva *et al.*, 2005) and confirmed later

in numerous other plant species. Interestingly, a recent study in several Rosaceae species

demonstrated that the cytosolic MVA derived GPP (Conart *et al.*, 2023) was directly hydrolyzed

by nudix hydrolase (NUDX) (Magnard *et al.*, 2015) and phosphatase to form geraniol

(MVA-NUDX case). Moreover, a transgenic study in *Nicotiana benthamiana* suggests that the

geraniol biosynthesis potential of cytosol mainly rely on the local GPP pool and geraniol synthase

abundance (Dong *et al.*, 2016). These cases raise the possibility of cytosolic geraniol biosynthesis

utilizing MVA-derived GPP and TPS geraniol synthase (MVA-TPS case), instead of a special

nudix hydrolase. However, almost all GES characterized so far are plastid-localized (Bartram *et*

al., 2006; Gutensohn *et al.*, 2013; Opitz *et al.*, 2014; Mendoza-Poudereux *et al.*, 2015; Abbas *et*

al., 2019), preventing this MVA-TPS case from validation.

In this study, we used the widely-loved ornamental cactus *E. oxypetalum* as a model system

to investigate the emission dynamics and biosynthetic route of cactus floral scent. We

demonstrated the temporal composition change during its transient nocturnal flowering and

proved that both petal and sepal were the major source of floral VOCs emission. By combining

transcriptomic and biochemical evidences, we functionally characterized a cytosol localized

geraniol synthase *in vitro*. Pathway-specific inhibitor assay showed that MVA pathway mainly

provided the substrate for geraniol biosynthesis, while biochemical quantification of metabolites

indicated that starch was used as carbon source for the synthesis of terpenoids volatiles. Together,

our results proved a rare MVA-TPS case of geraniol biosynthesis, and broadened our

understanding of volatile synthesis in cactus family.

Materials and methods

Plant materials and chemicals

Plant materials of *E. oxypetalum* were collected from 5-years old cottage seedlings at the *E.*

oxypetalum planting demonstration base of Sichuan Yuanlan Agricultural Development Co., Ltd.

in Zizhong, Sichuan, China (29°40′N, 104°45′E). The seedlings were grown under semi-natural

conditions with pine needle mulch soil. To aid in the development of healthy flowers, a specialized

phosphate-potash-amino acid mixture was applied as a foliar fertilizer. Shade roofs were used

during summer to protect plants from excessive sunlight. For RNA-seq, qRT-PCR, molecular

cloning and biochemical experiments, samples from various parts of the flowers were collected in

September, immediately frozen in liquid nitrogen and stored at -80□. The materials used for

SPME-GC/MS were collected into glass headspace bottles, immediately frozen and stored in dry

ice. Floral scent volatile samples for DHC-GC/MS were collected in October with headspace

adsorption. All chemical reagents used were purchased from Shanghai Aladdin Biochemical

Technology Co. Ltd. (www.aladdin-e.com).

Floral VOCs collection and GC-MS analysis

For *in situ* VOCs collection, floral scent was sampled hourly from 19:00 p.m. to 7:00 a.m. the

following day using a push-pull headspace sampling system. Briefly, precleaned (120□ for 1h)

polyethylene terephthalate (PET) bags (35 × 43 cm) were securely attached to the pedicel to

enclose the flower. Filtered air was circulated through the bag at a flow rate of 1,000 ml min^{-1}

using a battery-operated pump connected to Teflon tubing. The incoming air was purified by an

activated charcoal filter to remove particles and VOCs, and a copper tube coated with potassium

iodide to scrub for ozone. Air was sucked out of the bag at a flow rate of 200 ml min^{-1} through a

stainless-steel adsorbent cartridge packed with 155 mg of Tenax TA, 66 mg of Carbopack B and

75 mg of Carbopack X. VOCs were collected from three flowers simultaneously, each from a

different plant. After sampling, cartridges were sealed with Teflon-coated brass caps and stored at

4□ until analysis. Blank measurements from empty PET bags were done to characterize impurities

originating from sampling or analysis system.

VOCs trapped in adsorbent cartridges were thermally desorbed (TD100-xr, Markes

International Ltd, Llantrisant, UK) at 250□ for 10 min and analyzed on a gas

chromatography-mass spectrometry (GC-MS) (7890A GC, 5975C VL MSD, Agilent Technologies,

Santa Clara, California, USA). The VOCs were first cryofocused at -10 □ and then injected into

HP-5MS capillary column (30 m × 0.25 mm, film thickness 0.25 μm), using helium as carrier gas

at a flow rate of 1.0 ml min^{-1} with temperature program as follows: 40□, held 1 min; raised at 5□

min^{-1} to 210°C, raised at 20°C min^{-1} to 250°C, held 8 min.

For flower organ VOCs extraction, VOCs were sampled using SPME fiber (SUPELCO

57328-U, PDMS) for 40 min at 50 °C. VOCs adsorbed by SPME fibers were thermally desorbed at

230 °C before analyzed on GC-MS (SHIMADZU GCMS-QP2010) with column Rtx-5. Helium

was used as carrier gas at a flow rate of 1.0 ml min^{-1} using temperature program as follows: 50 °C,

held 3 min; raised at 6 °C min^{-1} to 200 °C, raised at 10 °C min^{-1} to 240 °C, held 6 min.

Chromatograms were analyzed using PARADISe v.6.0.1 (Quintanilla-Casas *et al.*, 2023).

Compounds were identified using pure standards when available, or tentatively identified using

NIST20 library. The VOC concentrations in blanks were subtracted from those in the samples.

Inhibition assay for MVA and MEP pathways

In the inhibition experiment, we administered 100µM solution of the MEP-specific inhibitor

fosmidomycin or the MVA-specific inhibitor mevinolin (Dudareva *et al.*, 2005) by uniformly

injecting 1ml of the solution into the junction of petals and perianth tubes at three specific time

points: 9:00 p.m. the day before bloom, 9:00 a.m. and 7:00 p.m. The control group was injected

with ddH$_2$O. Floral scent samples were collected at 0:00 a.m. the next day. Throughout the

experiment, the chosen inhibitors did not cause any noticeable effects on the appearance of the

flowers. For MVA inhibitor assay, the lactone hydrolysis of mevinolin were performed following

previous report (Kita *et al.*, 1980).

RNA-seq and analysis

Total RNAs were extracted from *E. oxypetalum* petal using an improved hexadecyl trimethyl

ammonium bromide (CTAB) method. At each sampling stages: pre-bloom (9:00 a.m.), blooming

(S06) and post-bloom (S13), three replicates of DNBSEQ RNA-seq libraries were prepared and

quality-checked. The libraries were sequenced using the MISEQ-2000 platform by BGI, resulting

in 150 bp paired-end reads. Subsequently, the raw sequencing reads underwent a quality control

and cleaning process (Bolger *et al.*, 2014), involving the removal of adaptors, ambiguous

N-containing reads, and low-quality reads. The resulting clean data were then *de novo* assembled

into transcripts using Trinity (Grabherr *et al.*, 2011; Haas *et al.*, 2013).

For sequence annotation, all assembled transcripts were compared to NCBI Nr, swiss-prot,

GO (The Gene Ontology Consortium *et al.*, 2023) and KEGG databases (Kanehisa & Goto, 2000)

with an E-value threshold at 1e^{-5} for BLAST operations (Altschul *et al.*, 1990). The Pfam protein

database (Mistry *et al.*, 2021) was searched using HMMER program (Eddy, 2011) with an E-value

threshold set at $1e^{-5}$. Annotation integration and functional categorization by GO terms were

performed by Trinotate (Duarte *et al.*, 2021). Later, abundance estimation and TMM (Trimmed

mean of M-values) was calculated using RSEM (Li & Dewey, 2011). Differential expression

analysis was conducted using DESeq2 (Love *et al.*, 2014), with a false discovery rate (FDR)

threshold of < 0.01 and an absolute value of $\log_2$ ratio ≥2 to identify differentially expressed genes.

GO term enrichment analysis was performed using GOseq R package (Young *et al.*, 2010) based

on the Wallenius non-central hyper-geometric distribution. Sequence clustering was carried out

using MUSCLE (Edgar, 2004), ModelTest-NG (Darriba *et al.*, 2020), RAxML-NG (Kozlov *et al.*,

2019) and iTOL (Letunic & Bork, 2021).

189　qRT-PCR for genes responsible for trans-geraniol synthesis

Specific primers for qRT-PCR were designed with NCBI. Total RNAs were extracted from *E.*

oxypetalum petals with Sangon Biotech Plant Total RNA Isolation Kit (Sangon Company,

https://www.sangon.com). Full-length cDNA was then reverse transcribed using Takara Prime

Script™ RT reagent Kit (Perfect Real Time) (https://www.Takarabiomed.com.cn). qRT-PCR was

performed with Takara RR430B TB Green Fast qPCR system, using independent RNA

preparations as biological replicates. *Ubqln* gene transcripts were used as control. The relative

expression levels of target genes were calculated by $2^{-\Delta\Delta Ct}$. The specificity of each primer pair was

verified by agarose gel electrophoresis, melting curve analysis and sequence homology with local

script. All relevant primer sequences were given in supplementary materials (Table S8).

199　Subcellular localization of EoTPSa1 and EoGDPS

Putative sequences of *EoTPSa1* and *EoGDPS* were selected by hmmscan with Pfam models

(PF01397.2, PF03936.19, PF19083.3) against the transcriptome assembly. Alignments were

performed using MUSCLE (Edgar, 2004), and sequence clustering was performed with

RAxML-ng. Primers were designed to amplify specific cDNA fragment with Takara

PrimeSTAR® Max PCR system. We utilized SignalP 6.0 for subcellular localization predicting.

Later, the transient gene expression in *N. benthamiana* leaves following established procedures

with pSuper1300 and *Agrobacterium tumefaciens* strain GV3101. After infiltration, the plants

were kept in the dark for 24 hours. 2.5 days after infiltration, confocal images *N. benthamiana*

leaves was taken using Zeiss CELL Observer SD.

The enzymatic activity validation of EoTPSa1

Structure prediction and molecular docking of EoTPSa1 were established with Alphafold2 (Jumper *et al.*, 2021) and AutoDock (Morris *et al.*, 2009) respectively. For *in vitro* enzymatic analysis, the cDNA of *EoTPSa1* was amplified and ligated into the multiple cloning sites of pET-28a expression vector utilizing Takara. In-Fusion Snap Assembly Master Mix. The constructed vector was then transformed into *Escherichia coli Rosseta-gami 2 (DE3)* cells. Positive clone incubation, protein induction, extraction and purification were all carried out under the guidance of Bacterial Protein Extraction Kit (Sangon) and Ni-NTA spin columns (Sangon) according to manufacturer's instructions. Purified EoTPSa1 protein was examined by 12.5% (w/v) SDS-PAGE.

The enzymatic assay was conducted in 0.5ml reaction buffer (50 mM Tris–HCl, 1 mM $MgCl_2$, and 0.1 mM $MnCl_2$) supplemented with 100 μM of geranyl diphosphate. To the reaction buffer, 0.5μg of purified EoTPSa1 protein was added. After overlaid with 100μl hexane, the reaction mixtures were incubated at 21 ☐ for 1 min. Reaction products were extracted by vigorously vertexing for 10s, keeping on ice for 30s, and centrifugation at 12000×g for 30s at 4 ☐ to separate the supernatant hexane. 10μl of the dehydrated hexane supernatant was collected for GC-MS analysis. Verification and quantification of the reaction products were achieved by comparing them with the NIST17 library and authentic standards.

Biochemical characterization of the acetyl-CoA generating metabolic pathway

The samples were collected from petals at different time points on T01-T08 (Fig. S1). The biochemical experiments were conducted using reagent kits purchased from Sangon Biotech company: reducing sugar (D799393), triglyceride (D799795), free fatty acids (D799793), and acetyl-CoA (D751001). Amylum and soluble polysaccharides were separated via centrifugation at 10000 rpm for 10 minutes after homogenization. The amylum fraction was washed with ddH_2O, while soluble polysaccharides were precipitated using 80% ethanol. Both components were then tested with the Amylum Content Assay Kit (D799325). Absorbance in each well was measured at the required wavelength using an Agilent BioTek Synergy H1 Multimode Reader.

Results

Trans-geraniol dominates *E. oxyptalum* floral scent and follows the unimodal emission

pattern.

Unlike many other plant species that bloom gradually over an extended period, *E.*

oxypetalum flowers rapidly blooms during few hours. In our experiments, the flowers typically

began opening after 20:00 local time, and reached full bloom around 23:00 (Fig. 1a, 1b, Video S1).

To investigate the temporal patterns of floral scent emission from *E. oxypetalum*, we collected

floral VOCs at 13 different stages using dynamic headspace collection, covering the entire

flowering period from 19:00 to 07:00 the next morning (Fig. 1b, 1e). In total, we identified 49

VOCs, including 26 terpenoids, 5 fatty acid derivatives and 18 benzenoids (Table S1). Among

them, trans-geraniol, benzyl alcohol and methyl salicylate were the dominant VOCs, collectively

constituting 90.25% of the total emission (165.45 μg/flower/h) at the peak emission stage (02:00

local time, Stage 08) (trans-geraniol, 72.54%; benzyl alcohol, 13.72%; methyl salicylate, 4.55%)

(Fig. 1b, Table S2). These three VOCs were also key variables in distinguishing between

blooming and non-blooming stages, as revealed by the OPLS-DA analysis (Fig. 1c).

Total floral VOCs emission followed the unimodal pattern, in synchrony with flower

openness (Fig. 1b, 1d). However, different compounds or compound classes exhibited varying

temporal patterns. Initially, benzenoid esters dominated, accounting for 92.46% of the total

emission shortly after flower opening at 21:00 (Stage 03). Terpenoid emission began around 20:00.

(Stage 04) and became the dominant VOCs class at 01:00 (Stage 07), constituting 81.33% of the

total emission. As terpenoid emission increased, benzenoid alcohols gradually surpassed

benzenoid esters as the primary benzenoid VOCs, reaching their peak emission at 03:00 (Stage

09), slightly after the terpenoid peak. Subsequently, as the flowers started to wither, the emission

of terpenoids and benzenoid alcohols decreased. However, from 05:00 (Stage 11), esters again

constituted the majority of VOCs till the flower fully withered (Fig. 1e). Notably, trans-geraniol

and benzyl alcohol displayed the unimodal emission pattern, while methyl salicylate exhibited a

bimodal pattern (Fig. 1b).

The floral scent is mainly emitted from petal and sepal of *E. oxypetalum*.

To determine the primary source for floral VOCs biosynthesis and emission, we analyzed

VOCs profiles of five organs, namely petal, sepal, floral tube, stamen and pistil (Fig. 2a) at three

flower developmental stages: pre-bloom (S1), blooming (S8) and post-bloom (S13) using solid

phase micro-extraction. The VOCs blends varied substantially depending on the flower

developmental stages and the specific flower organs. The most pronounced differences among

flower organs were observed at blooming, where the VOCs blends of petals and sepals were

similar to each other but clearly distinguishable from those of pistils, stamens and floral tubes (Fig.

2b, Table S3). Samples other than petals and sepals from the blooming stage, as well as pistils

from all stages, were clustered together. Trans-geraniol and benzyl alcohol were primarily emitted

from petals and sepals, followed by pistils, floral tubes and stamens, whereas methyl salicylate

was mainly emitted from pistils, followed by sepals and petals (Fig. 2c-e). Considering that petals

and sepals accounted for the majority of the whole flower biomass, these organs, particularly

petals, served as the primary source for trans-geraniol emission (Fig. 2f, 2g). Therefore, our

subsequent analyses focused on trans-geraniol biosynthesis and subcellular compartmentalization

in petal.

A cytosol-localized subfamily-a terpene synthase is responsible for trans-geraniol

biosynthesis

A total of 242.13 million reads were generated from nine libraries, resulting in the assembly

of 164,996 unigenes and 299,340 transcripts. The average contig N50 was 2,339 bp. 64.20% of

the transcripts were annotated with uniport and nr database. 45.52% of the transcripts were

functionally tagged by GO database (Table S5, S6). TMM values were utilized for gene

expression quantification. To identify the terpene synthases (TPS) responsible for trans-geraniol

biosynthesis, we focused on transcripts annotated with Pfam TPS models (PF01397.2,

PF03936.19, PF19083.3). A total of 84 unique peptides from 23 unigenes were identified as

putative TPSs, with eight of them showing significant high expression levels (Fig. 3a). Sequence

clustering analysis on these 23 putative TPSs from *E. oxypetalum* and 342 TPS sequences (Table

S7) from 9 other species yielded 7 subfamilies, with 17 *E. oxypetalum* TPSs belonging to

subfamily-a, 1 TPS to subfamily-b, 1 to subfamily-c, 1 to subfamily-e and 3 to subfamily-g (Fig.

3c). Among all the putative TPSs identified, a transcript *DN1377.2.1.2* from the subfamily-a

exhibited significantly higher expression levels. This particular transcript, designated as *EoTPSa1*,

exhibited approximately a 63-fold up-regulation during the blooming stage, which coincided with

the peak emission of trans-geraniol.

The predicted sequence of *EoTPSa1* had an Open Reading Frame (ORF) of 1638 bp,

resulting in a protein with 545 amino acids and a molecular weight of 63.29 kDa. Alignment of

this protein with other functionally validated geraniol synthase (GES) revealed the presence of

conserved Aspartate-rich DDxxD and NSE/DTE motifs, which were important for the

magnesium-dependent prenyl diphosphate ionization in class-I mechanisms for terpene synthases.

Additionally, a DxDD-like motif, commonly observed in class-II terpene synthases, was also

identified (Fig. 3d). Protein structure prediction indicated the characteristic α-β domain structure

of monoterpene synthases. Key amino acid residues, such as Asp^{294}, Asp^{298}, Asn^{440} and Glu^{448}

were found to be appropriately positioned for the binding of magnesium ions (Fig. 3b). The alpha

helixes in the α domain form a porous pocket that facilitates substrate binding and enables water

quenching of carbon cation intermediates for trans-geraniol formation. Molecular docking

simulations positioned magnesium ions and geraniol pyrophosphate accurately in the active center

of the enzyme. The catalytic ability of EoTPSa1 was further verified through an *in vitro* enzymatic

assay using geraniol pyrophosphate as the substrate. Notably, trans-geraniol was detected as the

sole reaction product above detection limit, suggesting a high specificity of EoTPSa1. (Fig. 4).

Additionally, EoTPSa1 was predicted to be localized in the cytosol, alongside the highly

expressed geranyl pyrophosphate synthase in *E. oxypetalum*, namely EoGDPS. We further

validated the cytosolic localization of EoTPSa1 and EoGDPS through *Nicotiana* transient

expression system. We detected strong GFP signals in the cytosols of transformed leaf epidermis

for all EoTPSa1-GFP, EoGDPS-GFP and pure GFP constructs. The GFP signal and the red

chloroplast fluorescence distribution did not exhibit overlapping patterns, suggesting the absence

of protein sorting activities (Fig. 5). These findings distinguish EoTPSa1 from other characterized

geraniol synthases, which are typically localized in plastids.

MVA pathway mainly provides precursors for trans-geraniol biosynthesis

To validate the cytosol-localized monoterpene biosynthesis and investigate the biosynthetic

origin of trans-geraniol precursors IPP and DMAPP, we examined the expression of genes

involved in the MVA and MEP pathways. Interestingly, homologous genes from the MVA pathway

experienced upregulation at the blooming stage, contrary to the uniformly low expression of MEP

pathway genes across all three time points (Fig. 6a, Table S9). Specifically, the *HMG* gene, which

encodes a putative hydroxymethylglutaryl-CoA reductase and acts as the rate-limiting enzyme of

the MVA pathway, showed a 69-fold upregulation at the blooming stage compared to the

pre-bloom stage. Another gene, *MVAD*, which encodes a putative diphosphate-mevalonate

decarboxylase, also exhibited a two-fold upregulation. On average, during the blooming stage, the

expression of the MVA pathway genes were found to be 33 times higher than that of the MEP

pathway. To further investigate the predominant role of the MVA pathway in the biosynthesis of

trans-geraniol, we employed pathway specific inhibitors, mevinolin for *HMG* and fosmidomycin

for *DXPS*, to selectively block the MVA and MEP pathways, respectively. We then collected

samples from the petal and sepal for VOCs characterization. While the VOCs of the

fosmidomycin-treated flowers and untreated flowers showed similarities, there was a noticeable

difference in the volatile components of the mevinolin-treated flowers (Fig. 6b, c, Table S4). This

distinction was mainly driven by the emission of monoterpenes, particularly trans-geraniol, which

showed 83.4% decrease in mevinolin-treated flowers but a mild increase in fosmidomycin-treated

groups (Fig. 6d). In addition, mevinolin-treated flowers emitted higher amounts of ethyl salicylate

but lower amounts of benzyl alcohols and methyl salicylate, whereas fosmidomycin-treated

flowers emitted higher amounts of benzyl alcohol and methyl salicylate (Fig. 6e-g). This suggests

that blocking either the MVA or MEP pathways may divert the carbon flow to other biosynthetic

pathways, such as the benzenoid biosynthetic pathways.

Amyloplasts provide carbon source for floral scent biosynthesis

To explore the origin of the carbon source for floral VOCs biosynthesis in *E. oxypetalum*, we

investigated the expression patterns of homologous genes involved in starch degradation,

glycolysis, triglyceride degradation and fatty acid β-oxidation. These genes showed inconsistent

expression patterns during the pre-blooming stage compared to the blooming and post-blooming

stages. Notably, the beta amylase *BAM*, which catalyzes the initial step of starch degradation,

showed a 6.8-fold upregulation at the full-blooming stage, while the genes involved in triglyceride

degradation and fatty acid β-oxidation did not exhibit a similar trend (Fig. 7a).

Later, our biochemical quantification showed that the degradation of starch began in the

T02-T03 stage, leading to the accumulation of reducing sugars and a sharp increase in the content

of acetyl CoA, which serves as a direct substrate for the MVA pathway. Over the next half day

from T03 to T05, as metabolic activity intensified, starch degradation further increased; the

concentration of reducing sugars remained consistently high, while acetyl-CoA continuously

consumed. Although the total amount of pyruvate, an intermediate metabolite, was relatively low,

it exhibited a significant upward trend, indicating increased physiological activity in *E.*

oxypetalum. By the T05-T06 stage, due to active metabolism, starch concentration reached its

lowest level, the content of pyruvate reached its peak, while acetyl-CoA was consumed to its

minimum due to intense physiological processes. After the blooming stage, the acetyl-CoA content

returned to its initial lowest level (Fig. 7b-e). However, not only triglycerides and free fatty acids

were detected at negligible levels from T01-T07, but the trend of their changes was also

independent of acetyl-CoA (Fig. 7f, 7g). Together, these findings suggest that the carbon source

for trans-geraniol biosynthesis in *E. oxypetalum* originates from starch granules within the

amyloplasts.

Discussion

The biosynthesis and emission pattern of floral volatiles is highly diverse among different

taxa. The Cactaceae family contains several well-known ornamental species for their unique

morphological characteristics and attractive flowers (Mauseth, 2006). Extensive work has been

focused on their pollination mode and ecological relevance, but the biosynthesis of their floral

scent remains mostly unexplored. Here, using the famous *E. oxypetalum* as a model system, we

investigated the composition and biosynthesis of its floral scent in the great detail. We first

characterized its highly dynamic scent composition and then identified the petal and sepal to be

the major source of VOC emission. By exploring the biosynthesis of its major component

trans-geraniol, we finally showed a novel MVA-TPS case for geraniol biosynthesis in *E.*

oxypetalum.

Highly dynamic composition of *E. oxypetalum* floral scent volatiles during transient
blooming

E. oxypetalum has impressive huge flowers and strong scent, both of which are widely

distributed traits across Cactaceae. Previous investigation categorized cactus flowers into different

classes according to their pollination mode, each with distinct floral VOCs composition and

emission dynamics (Kaiser & Tollsten, 1995). Similar to those nocturnal blooming cacti, *E.*

oxypetalum emits excessively high content of terpenoid and benzenoid volatiles during its

transient blooming, which lasts for one night only. The nocturnal flowering and strong scent

emission may be both due to its coevolution with insects (Guerrero *et al.*, 2019) and a legacy of

drought adaptation when ancient cactus first thrived in central Andeans (Hernández-Hernández *et*

al., 2014). Interestingly, in our study, the composition of *E. oxypetalum* scent did not keep the

same even between the short 1-hour sampling interval. Trans-geraniol was the primary compound

responsible for its floral scent, and exhibited a unimodal emission pattern peaking at full blooming.

The two top benzenoids in *E. oxypetalum*, benzyl alcohol and methyl salicylate, exhibited their

possible substrate competition (Schuurink *et al.*, 2006), as benzyl alcohol peaked around

full-bloom while methyl salicylate peaked before and after full-bloom. The biosynthesis of these

VOCs primarily occurred in the petal and sepal tissue, both with high surface-to-volume ratio to

promote the overall scent emission. Also, trans-geraniol, benzyl alcohol and methyl salicylate are

all known to convey chemical signals between plants and insects, especially trans-geraniol for

hawk-moth (Foster And & Harris, 1997) , and thus they may have played crucial roles in the

interaction between *E. oxypetalum* and pollinators. Whether these compounds, either individually

or in specific ratios, during flowering convey specific chemical messages to pollinators remains to

be investigated (Raguso, 2008).

Geraniol biosynthesis in MVA-TPS pathway

Generally, monoterpenes volatiles are synthesized with plastid-localized monoterpene

synthase utilizing MEP-derived GPP (MEP-TPS case) (Dudareva *et al.*, 2005). Interestingly,

recent work in rose proved a special nudix hydrolase that catalyzed the hydrolysis of GPP into

geraniol (MVA-NUDX case) (Magnard *et al.*, 2015; Conart *et al.*, 2023); However, apart from

these two canonical cases, some recent reports provided several exceptions. Several

cytosolic-localized geraniol synthases were characterized in *Fragaria ananassa* (Aharoni *et al.*,

2004), *Lippia dulcis* (Dong *et al.*, 2013), *N. benthamiana* (Wu *et al.*, 2006; Dong *et al.*, 2016) and

Solanum lycopersicum (Davidovich-Rikanati *et al.*, 2008; Zhou & Pichersky, 2020). Moreover,

isotope labeling experiments proved MEP-dependence of sesquiterpene formation in snapdragon,

along with MVA-dependent linalool formation in raspberry (Hampel *et al.*, 2007). These results

suggest a possible new case for monoterpene biosynthesis, in which a cytosolic TPS might utilize

MVA-derived GPP for monoterpene production (MVA-TPS case). This MVA-TPS case might be

more ubiquitous than the special MVA-NUDX case for geraniol biosynthesis, as a nudix hydrolase

capable of hydrolyzing GPP into geraniol was only characterized for rose hybrids and

Pelargonium graveolens (Bergman *et al.*, 2021) so far, while both TPS and MVA pathway are

widely distributed in higher plants.

In the current study, we have provided both biochemical and transcriptomic evidence that a

cytosolic-localized TPS utilizes MVA-derived GPP to synthesize trans-geraniol in *E. oxypetalum*

flowers. The cytosol localization of geraniol synthase EoTPSa1 and significantly reduced

terpenoid content during mevinolin treatment provided the solid evidence for the MVA-TPS case.

The special geraniol synthase EoTPSa1, which contributed most of the total scent emission at full

bloom, was classified into subfamily-a during sequence clustering. Previous study has suggested

sesquiterpene preference for subfamily-a TPSs (Chen *et al.*, 2011), and the characterized geraniol

synthases so far has diverse phylogenetic preference. This observation concurs with previous

knowledge about active gene neo-functionalization during terpenoid evolution, and preference for

utilizing existing enzymes for specialized metabolism with catalytic promiscuity (Weng, 2014).

Incorporation of cytosolic MVA pathways rather than plastid originated substrate coupled

downstream transportation can improve catalysis efficiency significantly (Dong *et al.*, 2016). This

special MVA pathway incorporation might be due to the elevated HMG expression at full bloom,

which might share the common regulation scheme with eoTPSa1.

In our transcriptome assembly, *EoTPSa1* was the only terpene synthase encoding gene that

exhibited significant expression. *In vitro* enzymatic assay identified EoTPSa1 with significant

trans-geraniol biosynthesis activity. Interestingly, both isomers trans-geraniol and cis-geraniol,

along with their respective oxidized products trans- and cis-citral were detected in floral scent.

While the existence of citral suggests the possible dehydrogenase activity (Zhao *et al.*, 2023),

cis-geraniol formation indicates an uncommon lack of stereoselectivity during GPP biosynthesis.

Usually, during the nucleophilic attack of IPP on the isoprene carbocation generated by the

dissociation of pyrophosphate in DMAPP, the proton abstracted by the newly formed tertiary

carbocation often generates an trans-structured double bond (Walsh & Tang, 2017). The formation

of less common cis-structured double bonds in isopentenyl chain extension is only sometimes

observed in bactoprenols formation (Walsh & Wencewicz, 2016). In order to form a cis-structured

double bound, the basic group that catalyzes the departure of hydrogen ion must be positioned on

the opposite side of the reaction site, which is evident by the difference between trans- and

cis-stereoselective isopentyl diphosphate synthase (Guo *et al.*, 2004). However, our results showed

that only one GPP synthase might be active during *E. oxypetalum* trans- and cis-GPP synthesis, suggesting that the GPPS identified might be capable to form both trans- and cis-structured double bound simultaneously, and the geraniol synthase to hydrolyze these two isomers at the same time.

Starch as primary carbon source for the synthesis of *E. oxypetalum* floral VOCs

MVA pathway utilizes acetyl-CoA as substrate, which can derive from both starch and lipid degradation. Transcriptome assembly and quantifications identified the active expression at blooming stage for both β-amylase (BAM) and TAGL, the initial steps for these two degradation pathways. Biochemical quantification revealed the existence of excessively high content of starch (up to 30% dry weight) and reducing sugars in petal, but almost complete lack of triglyceride and free fatty acids, indicating that starch may act as the primary carbon source. The drop in starch content suggests that starch degradation initiated in the morning on the day of blooming, creating hoarded acetyl-CoA for active metabolism during flower opening and volatile biosynthesis. Thus, our findings suggest that an adequate supply of substrate, the strict regulation of enzymes, and the selection of the cytoplasm as the site for geraniol synthesis collectively constitute the mechanism that allows *E. oxypetalum* to release a substantial number of volatile compounds in such a short flowering period, and distinguishing it from other plants in the cactus family with usually longer flowering length and less fragrance emission.

Acknowledgements

This work was supported by the National Nature Science Foundation of China (32271692, 32100326), and the Department of Science and Technology of Sichuan Province of China (2022ZHXC0009, 2023YFSY0054). We further wish to thank Yudong Ma, Wei Wu and Jin Dai for their assistance in sample collection, and Bolei Jiao, Xinrui Liu and Chuanfang Wu for sharing the *N. benthamiana* plants and pSuper1300, pET-28a plastids. We also would like to thank Qijun Deng and Wei Ni from Sichuan Bowu Cultural Development Co. Ltd. for the drawing of *E. oxypetalum* flowers.

Competing interests

None declared

Author contributions

Yiyang Zhang, Yuhan Zhang, LT and QS designed the experiments. The top seven authors

and WT jointly completed the collection and preparation of samples. Yiyang Zhang and Yuhan

Zhang quantified VOCs content with GCMS. YiYang Zhang analyzed GCMS and transcriptome

data. Yuhan Zhang and QT performed molecular cloning, enzymatic assay and localization studies.

LW performed biochemical quantifications. Yiyang Zhang plotted the figures and wrote the first

draft. Yuhan Zhang, QS, TL, HX, QT, ML and ZL revised the manuscript. Yiyang Zhang and

Yuhan Zhang contributed equally to this work.

Data Availability

All data needed to evaluate the conclusions in the paper are present inthe paper and/or the

Supplementary Materials. The transcriptome raw reads for 9 individuals in this study have been

deposited in the National Genomics Data Center (https://ngdc.cncb.ac.cn) under accession number

PRJCA024038.

References

Abbas F, Ke Y, Yu R, Fan Y. 2019. Functional characterization and expression analysis of two

terpene synthases involved in floral scent formation in Lilium 'Siberia'. *Planta* **249**: 71–93.

Abbas F, Ke Y, Yu R, Yue Y, Amanullah S, Jahangir MM, Fan Y. 2017. Volatile terpenoids: multiple

functions, biosynthesis, modulation and manipulation by genetic engineering. *Planta* **246**:

803–816.

Aharoni A, Giri AP, Verstappen FWA, Bertea CM, Sevenier R, Sun Z, Jongsma MA, Schwab W,

Bouwmeester HJ. 2004. Gain and loss of fruit flavor compounds produced by wild and cultivated

strawberry species. *The Plant Cell* **16**: 3110–3131.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool.

Journal of Molecular Biology **215**: 403–410.

Bartram S, Jux A, Gleixner G, Boland W. 2006. Dynamic pathway allocation in early terpenoid

biosynthesis of stress-induced lima bean leaves. *Phytochemistry* **67**: 1661–1672.

Bergman ME, Bhardwaj M, Phillips MA. 2021. Cytosolic geraniol and citronellol biosynthesis

require a Nudix hydrolase in rose-scented geranium (Pelargonium graveolens). *The Plant Journal:*

For Cell and Molecular Biology **107**: 493–510.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**: 2114–2120.

Chen F, Tholl D, Bohlmann J, Pichersky E. 2011. The family of terpene synthases in plants: a
mid-size family of genes for specialized metabolism that is highly diversified throughout the
kingdom: Terpene synthase family. *The Plant Journal* **66**: 212–229.

Chen W, Viljoen AM. 2010. Geraniol — A review of a commercially important fragrance material.
South African Journal of Botany **76**: 643–651.

Christianson DW. 2017. Structural and Chemical Biology of Terpenoid Cyclases. *Chemical Reviews*
117: 11570–11648.

Conart C, Bomzan DP, Huang X-Q, Bassard J-E, Paramita SN, Saint-Marcoux D, Rius-Bony A,
Hivert G, Anchisi A, Schaller H, *et al*. 2023. A cytosolic bifunctional geranyl/farnesyl diphosphate
synthase provides MVA-derived GPP for geraniol biosynthesis in rose flowers. *Proceedings of the*
National Academy of Sciences **120**: e2221440120.

Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. 2020. ModelTest-NG: A New
and Scalable Tool for the Selection of DNA and Protein Evolutionary Models. *Molecular Biology*
and Evolution **37**: 291–294.

Davidovich-Rikanati R, Lewinsohn E, Bar E, Iijima Y, Pichersky E, Sitrit Y. 2008. Overexpression of
the lemon basil alpha-zingiberene synthase gene increases both mono- and sesquiterpene
contents in tomato fruit. *The Plant Journal: For Cell and Molecular Biology* **56**: 228–238.

Dong L, Jongedijk E, Bouwmeester H, Van Der Krol A. 2016. Monoterpene biosynthesis potential
of plant subcellular compartments. *New Phytologist* **209**: 679–690.

Dong L, Miettinen K, Goedbloed M, Verstappen FWA, Voster A, Jongsma MA, Memelink J, van
der Krol S, Bouwmeester HJ. 2013. Characterization of two geraniol synthases from Valeriana
officinalis and Lippia dulcis: similar activity but difference in subcellular localization. *Metabolic*
Engineering **20**: 198–211.

Dötterl S, Gershenzon J. 2023. Chemistry, biosynthesis and biology of floral volatiles: roles in
pollination and other functions. *Natural Product Reports*.

Duarte GT, Volkova PYu, Geras'kin SA. 2021. A Pipeline for Non-model Organisms for de novo
Transcriptome Assembly, Annotation, and Gene Ontology Analysis Using Open Tools: Case Study

with Scots Pine. *Bio-protocol* **11**: e3912.

Dudareva N, Andersson S, Orlova I, Gatto N, Reichelt M, Rhodes D, Boland W, Gershenzon J.

2005. The nonmevalonate pathway supports both monoterpene and sesquiterpene formation in

snapdragon flowers. *Proceedings of the National Academy of Sciences* **102**: 933–938.

Dudareva N, Klempien A, Muhlemann JK, Kaplan I. 2013. Biosynthesis, function and metabolic

engineering of plant volatile organic compounds. *New Phytologist* **198**: 16–32.

Eddy SR. 2011. Accelerated Profile HMM Searches (WR Pearson, Ed.). *PLoS Computational*

Biology **7**: e1002195.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Research **32**: 1792–1797.

Foster And SP, Harris MO. 1997. BEHAVIORAL MANIPULATION METHODS FOR INSECT

PEST-MANAGEMENT. *Annual Review of Entomology* **42**: 123–146.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,

Raychowdhury R, Zeng Q, *et al.* 2011. Trinity: reconstructing a full-length transcriptome without

a genome from RNA-Seq data. *Nature biotechnology* **29**: 644–652.

Guerrero PC, Majure LC, Cornejo-Romero A, Hernández-Hernández T. 2019. Phylogenetic

Relationships and Evolutionary Trends in the Cactus Family. *Journal of Heredity* **110**: 4–21.

Guo R-T, Kuo C-J, Chou C-C, Ko T-P, Shr H-L, Liang P-H, Wang AH-J. 2004. Crystal structure of

octaprenyl pyrophosphate synthase from hyperthermophilic Thermotoga maritima and

mechanism of product chain length determination. *The Journal of Biological Chemistry* **279**:

4903–4912.

Gutensohn M, Orlova I, Nguyen TTH, Davidovich-Rikanati R, Ferruzzi MG, Sitrit Y, Lewinsohn E,

Pichersky E, Dudareva N. 2013. Cytosolic monoterpene biosynthesis is supported by

plastid-generated geranyl diphosphate substrate in transgenic tomato fruits. *The Plant Journal* **75**:

351–363.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li

B, Lieber M, *et al.* 2013. De novo transcript sequence reconstruction from RNA-Seq: reference

generation and analysis with Trinity. *Nature protocols* **8**: 10.1038/nprot.2013.084.

Hampel D, Swatski A, Mosandl A, Wüst M. 2007. Biosynthesis of monoterpenes and

norisoprenoids in raspberry fruits (Rubus idaeus L.): the role of cytosolic mevalonate and

plastidial methylerythritol phosphate pathway. *Journal of Agricultural and Food Chemistry* **55**:

9296–9304.

Hernández-Hernández T, Brown JW, Schlumpberger BO, Eguiarte LE, Magallón S. 2014. Beyond

aridification: multiple explanations for the elevated diversification of cacti in the New World

Succulent Biome. *New Phytologist* **202**: 1382–1397.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,

Žídek A, Potapenko A, *et al.* 2021. Highly accurate protein structure prediction with AlphaFold.

Nature **596**: 583–589.

Kaiser R, Tollsten L. 1995. An introduction to the scent of cacti. *Flavour and Fragrance Journal* **10**:

153–164.

Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids*

Research **28**: 27–30.

Kita T, Brown MS, Goldstein JL. 1980. Feedback regulation of 3-hydroxy-3-methylglutaryl

coenzyme A reductase in livers of mice treated with mevinolin, a competitive inhibitor of the

reductase. *The Journal of Clinical Investigation* **66**: 1094–1100.

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAxML-NG: a fast, scalable and

user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**: 4453–4455.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. *Nucleic Acids Research* **49**: W293–W296.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC bioinformatics* **12**: 323.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* **15**: 550.

Magnard J-L, Roccia A, Caissard J-C, Vergne P, Sun P, Hecquet R, Dubois A, Hibrand-Saint Oyant

L, Jullien F, Nicolè F, *et al.* 2015. Biosynthesis of monoterpene scent compounds in roses. *Science*

349: 81–83.

Mauseth JD. 2006. Structure-Function Relationships in Highly Modified Shoots of Cactaceae.

Annals of Botany **98**: 901–926.

Mendoza-Poudereux I, Kutzner E, Huber C, Segura J, Eisenreich W, Arrillaga I. 2015. Metabolic

cross-talk between pathways of terpenoid backbone biosynthesis in spike lavender. *Plant*

Physiology and Biochemistry **95**: 113–120.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE,

Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families database in 2021. *Nucleic*

Acids Research **49**: D412–D419.

Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, Olson AJ. 2009.

AutoDock4 and AutoDockTools4: Automated Docking with Selective Receptor Flexibility. *Journal*

of computational chemistry **30**: 2785–2791.

Muhlemann JK, Klempien A, Dudareva N. 2014. Floral volatiles: from biosynthesis to function.

Plant Cell and Environment **37**: 1936–1949.

Opitz S, Nes WD, Gershenzon J. 2014. Both methylerythritol phosphate and mevalonate

pathways contribute to biosynthesis of each of the major isoprenoid classes in young cotton

seedlings. *Phytochemistry* **98**: 110–119.

Quintanilla-Casas B, Bro R, Hinrich JL, Davie-Martin CL. 2023. *Tutorial on PARADISe:*

PARAFAC2-based Deconvolution and Identification System for processing GC–MS data. Protocol

Exchange.

Raguso RA. 2008. Wake Up and Smell the Roses: The Ecology and Evolution of Floral Scent.

Annual Review of Ecology, Evolution, and Systematics **39**: 549–569.

Schuurink RC, Haring MA, Clark DG. 2006. Regulation of volatile benzenoid biosynthesis in

petunia flowers. *Trends in Plant Science* **11**: 20–25.

The Gene Ontology Consortium, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ,

Ebert D, Feuermann M, Gaudet P, Harris NL, et al. 2023. The Gene Ontology knowledgebase in

2023. *Genetics* **224**: iyad031.

Vranová E, Coman D, Gruissem W. 2012. Structure and dynamics of the isoprenoid pathway

network. *Molecular Plant* **5**: 318–333.

Walsh CT, Tang Y. 2017. *Natural Product Biosynthesis: Chemical Logic and Enzymatic Machinery*.

The Royal Society of Chemistry.

Walsh C, Wencewicz T. 2016. *Antibiotics: Challenges, Mechanisms, Opportunities, 2nd Edition*.

Washington DC: ASM Press.

Weng J-K. 2014. The evolutionary paths towards complexity: a metabolic perspective. *New*

Phytologist **201**: 1141–1149.

Wu S, Schalk M, Clark A, Miles RB, Coates R, Chappell J. 2006. Redirection of cytosolic or

plastidic isoprenoid precursors elevates terpene production in plants. *Nature Biotechnology* **24**:

1441–1447.

Young MD, Wakefield MJ, Smyth GK, Oshlack A. 2010. Gene ontology analysis for RNA-seq:

accounting for selection bias. *Genome Biology* **11**: R14.

Zhang W, Jiang Y, Chen F, Guan Z, Wei G, Chen X, Zhang C, Köllner TG, Chen S, Chen F, *et al*. 2022.

Dynamic regulation of volatile terpenoid production and emission from Chrysanthemum

morifolium capitula. *Plant Physiology and Biochemistry* **182**: 11–21.

Zhao Y, Chen Y, Gao M, Wang Y. 2023. Alcohol dehydrogenases regulated by a MYB44

transcription factor underlie Lauraceae citral biosynthesis. *Plant Physiology*: kiad553.

Zhou F, Pichersky E. 2020. The complete functional characterisation of the terpene synthase

family in tomato. *New Phytologist* **226**: 1341–1360.

Figure legend

Fig. 1 Temporal patterns of flower opening and volatile organic compounds (VOCs) emission

dynamics in *Epiphyllum oxypetalum.*

Fig. 2 Emission patterns of five floral tissues during pre-bloom, blooming, and post-bloom stages.

Fig. 3 Identification of the terpene synthase for trans-geraniol biosynthesis.

Fig. 4 Biochemical characterization of EoTPSa1 as trans-geraniol synthase.

Fig. 5 Subcellular localization of EoTPSa1 and EoGDPS.

Fig. 6 Preference of the trans-geraniol biosynthesis pathway in *Epiphyllum oxypetalum* petal.

Fig. 7 Carbon source for trans-geraniol biosynthesis in Epiphyllum oxypetalum petal.

Supporting information

Table S1. Content matrix of compounds detected in *Epiphyllum oxypetalum* floral scent.

Table S2. Absolute content of top eight volatiles detected in floral scent.

Table S3. Content matrix of compounds detected in samples of 5 floral tissues at pre-bloom,

655   blooming and post-bloom.

Table S4. Content matrix of compounds detected in pathway inhibitor assay.

Table S5. Transcriptome expression matrix of samples at pre-bloom, blooming and post-bloom.

Table S6. Transcriptome annotation matrix of samples at pre-bloom, blooming and post-bloom.

Table S7. Protein sequences used for TPS family sequence clustering.

Table S8. Primers used in this study.

Table S9. Results matrix for qRT-PCR measurement of genes involved in MVA, MEP and geraniol

662   synthesis.

663   Figure S1. Sampling time for photography, transcriptome, headspace and biochemical

664   experiments.

665   Video S1. Time-lapse video for the nocturnal blooming of *Epiphyllum oxypetalum*

Fig. 1 Temporal patterns of flower opening and volatile organic compounds (VOCs) emission

dynamics in *Epiphyllum oxypetalum*.

a: Time-lapse photos showing the rapid blooming of *E. oxypetalum* flowers.

b: VOC emission rate of the categories of trans-geraniol, benzyl alcohol, methyl salicylate and

total emission across all sampling stages.

c: VIP value of top 20 VOCs from OPLS-DA analysis. The three major VOCs with VIP values

higher than threshold were colored in brighter red.

d: PCA clustering of floral VOCs content. Dots were shaped and colored according to sampling

stages, time tags inside arrow rings corresponds to sampling time of nearby dots.

e: Relative concentration of terpenoids, fatty-acid derivatives, benzenoid alcohols, esters and other

678   VOCs across all sampled stages. Note that the legend is presented in c.

Fig. 2 Emission patterns of five floral tissues during pre-bloom, blooming, and post-bloom stages.

a: Schematic diagram of the longitudinal section of *Epiphyllum oxypetalum* floral tissues, with the five types of tissues colored differently.

b: PCA clustering of volatile organic compound (VOC) content of peal, sepal, floral tube, stamen and pistil at pre-bloom, blooming and post-bloom stage.

687   c-e: Dry weight-based content of trans-geraniol, benzyl alcohol and methyl salicylate of five floral

688   tissues at blooming stage.

689   f: Dry weight of five tissues sampled at blooming stage.

690   g: Relative content of five major classes of VOCs at blooming stage, not adjusted by dry mass.

Fig. 3 Identification of the terpene synthase for trans-geraniol biosynthesis.

a: Expression patterns of eight terpene synthases identified with significant expression during

pre-bloom, blooming and post-bloom stages, with each block colored according to TMM value.

b: Structure prediction and molecular docking of EoTPSa1. Protein structure colored according to

domain architecture and catalyzing motifs with the residues of Asp^{294}, Asp^{298}, Asn^{440} and Glu^{448}

shown.

699   c: Clustering of identified *Epiphyllum oxypetalum* TPSs with TPSs from other plant species. The

700   major TPSs in *E. oxypetalum* petal marked with dashed line.

701   d: Protein alignment of EoTPSa1 and GESs from other six species, with protein sequence around

702   DxDD, DDxxD and NSE/DTE motifs colored according to amino acid properties. Subcellular

703   localizations predicted with signal P 6.0 were also given.

Fig. 4 Biochemical characterization of EoTPSa1 as trans-geraniol synthase.

a: TIC comparison between active enzyme and inactive control from 14.5 min to 20 min RT, showing huge trans-geraniol peak at 19.045 RT.

b-c: Mass spectral fragmentation pattern of the trans-geraniol peak (RT 19.045) catalyzed with EoTPSa1 (b) and the trans-geraniol standard (c).

713 **Fig. 5 Subcellular localization of EoTPSa1 and EoGDPS.**

714 a-c: Subcellular localization of EoTPSa1-GFP, EoGDPS-GFP, and the control GFP, respectively.

715 Confocal image was taken at AF488 green band for GFP fluorescence, and AF568 red band for

716 chloroplast autofluorescence.

Fig. 6 Preference of the trans-geraniol biosynthesis pathway in *Epiphyllum oxypetalum* petal.

a: Expression pattern of MVA and MEP pathway genes during pre-bloom, blooming and post-bloom stages, with each block colored according to TMM value.

b: PCA clustering of volatile organic compound (VOC) content of petals with mevinolin, fosmidomycin treatment group and control group at blooming. Dots were shaped and colored according to different treatments.

c: VIP value of top 20 VOCs from OPLS-DA analysis. VOCs with VIP values higher than threshold were colored in orange.

d-g: VOC content of trans-geraniol, benzyl alcohol, methyl salicylate and ethyl salicylate in petal

of mevinolin, fosmidomycin and control group.

Fig. 7 Carbon source for trans-geraniol biosynthesis in *Epiphyllum oxypetalum* petal.

a: Expression pattern of starch degradation, glycolysis and β-oxidation pathway genes at

pre-bloom (T03), blooming (T06) and post-bloom (T07), with each block colored according to

TMM value.

b-g: Concentration of starch, reducing sugar, pyruvate, acetyl-CoA, free fatty acid and triglyceride

in petal during T01-T08.