Mycobacterium smegmatis NucS-promoted DNA Mismatch Repair involves limited resection by a 5'-3' exonuclease and is independent of homologous recombination and NHEJ

Iris V. Rivera-Flores , Emily X. Wang, and Kenan C. Murphy

Department of Microbiology and Physiological Systems

University of Massachusetts Chan Medical School

Worcester, MA 01605

ABSTRACT

The MutSL mismatch repair system in bacteria and eukaryotes correct mismatches made at the replication fork to maintain genome stability. Despite lacking the mutSL genes, most actinobacterial species, including *Mycobacterium tuberculosis,* exhibit spontaneous mutation rates similar to MutSL-bearing bacteria, suggesting the existence of an alternative mismatch repair (MMR) system. It has been shown previously that the EndoMS/NucS protein from Actinobacterium *Corynebacterium glutamicum* is a mismatch-specific endonuclease that recognizes and creates dsDNA breaks in vitro. Additionally, the structure of the NucS protein from the archaeal species P. *abyssi* reveals high similarity to the structures of restriction endonucleases, leading to a model that NucS mismatch repair involves a dsDNA break in the repair process.   In this report, a genetic analysis shows that an M. *smegmatis* Δ*nucS* strain could be complemented by expression of wild type NucS protein, but not by one deleted of its last five amino acids, a region predicted to be critical for binding to the β-clamp at the replication fork. Oligo-recombineering promoted by Che9 phage RecT annealase was then leveraged to directly deliver defined mismatches to a defective hygromycin resistance gene on the *M. smegmatis* chromosome. We find that NucS recognizes and repairs G-G, G-T, and T-T mismatches in vivo, consistent with the recognition of these same mismatches in C. *glutamicum* in vitro, as well as mutation accumulation studies done in M. smegmatis. Three residues in *M. smegmatis* NucS, predicted to be catalytic residues based on the restriction enzyme-like structure of *P. abyssi* NucS, were mutated to alanine; all three mutants were defective for mismatch repair. Finally, an assay the employs one oligo that promotes the generation of two mismatches in close proximity on the chromosome shows that a NucS-promoted cut in dsDNA is processed by a 5' – 3' exonuclease, but is

independent of both RecA, RadA, and NHEJ functions Ku and LigD, suggesting that the mechanism of DNA mismatch repair does not involve a homologous recombination event or non-homologous end-joining.

INTRODUCTION

High fidelity of DNA replication in all organisms is of paramount importance to minimize mutation accumulation and maintain genome stability. Most cells correct the misincorporation of the wrong nucleotide during DNA replication by two distinct mechanisms. The first line of defense is provided by the proofreading functions of DNA polymerases that immediately detect and exonucleolytically remove an incorrect nucleotide in the nascent DNA strand (1-3). Should this system fail, however, DNA mismatch repair (MMR) systems recognize mismatched base pairs in the wake of the replication fork, identify which base of the mismatch is the incorrect one, and processes its removal (4,5). Resynthesis of the exposed template and religation to the undamaged preexisting DNA strand completes the repair process. The canonical MMR pathway is represented by the MutSL system, which was first identified in *E. coli* (6,7). The E. coli MutS protein recognizes the mismatched bases, and together with MutL promotes a platform for nicking of the newly synthesized strand by MutH endonuclease, governed by the hemi-methylation status of the newly synthesized strand. In most bacteria, however, *mutH* is not present and a latent MutL endonuclease is responsible for cutting the replicative strand (for reviews, see (8-11)). Homologous MutSL systems are found ubiquitously in both prokaryotic and eukaryotic organisms, including humans (5).

Mutations of these MMR genes lead to hyper-mutagenic genetic backgrounds, increased rates of homeologous recombination, and cancer.

Most archaeal and many Actinobacterial genomes (including mycobacteria) are devoid of the canonical MutSL MMR system. It was believed for many years that perhaps these organisms did not possess an MMR system. However, these species do not show higher rates of mutagenesis relative to bacteria that contain MutSL systems, suggesting that DNA replication fidelity in these bacteria rely solely on an efficient polymerase proofreading mechanism, or that an alternative MMR system exist in these species. The first hint of such a repair system came from the studies of Ishino et al (12) that described an endonuclease from the hyperthermophilic archaeon *Pyrococcus furiosus* that specifically recognizes and cuts mismatched dsDNA, which they called EndoMS (endonuclease specific for mismatch-specific DNA). EndoMS is homologous to a protein previously isolated from *Pyrococcus abyssi*, that was identified as a nuclease specific for ssDNA, called NucS (13), which is the name used for the protein hereafter.

Ichino *et al* (12) went on to study the NucS protein from *Thermococcus kodakarensis* (TKO), a more genetically tractable thermophilic archaea. They found that the TKO NucS specifically binds to and cuts both strands of dsDNA species that contained G/T, G/G and T/T mismatches. Higher concentrations of the enzyme were required to cut dsDNA containing T/C and A/C mismatches, though such substrates failed to bind TKO NucS in an electrophoretic mobility shift assay. DNA substrates containing C/C, A/C and A/A showed no binding or cutting of mismatched substrates.

The NucS proteins have no homology to any of the components of the mutSL systems, and thus was the first protein from an archaeon that suggested the existence of an alternative MMR system. Enzymatically, TKO NucS cuts on both sides of the mismatch leaving 5'-P and 3-OH ends with 5' protruding overhangs, reminiscent of the action of restriction enzymes. Indeed, the structure of the TKO NucS bound to 15-mer mismatched-containing dsDNA was solved by Nakae *et al* (14) showing that NucS protein to be remarkably similar to the structure of type II restriction enzymes, with the N-terminal domain promoting dimerization of NucS and the C-terminal domain binding to regions flanking the mismatched bases. The two bases of the mismatch were flipped-out of the helix into binding sites within the protein, not unlike the recognition sequence of the Ecl18kl restriction enzyme (CCNGG), where the central base pair is flipped out in a manner like NucS acting on a mismatch. Unlike restriction enzymes, however, NucS does not have a recognition sequence, in that it does not contact any bases at the cut site except for those contained within the mismatch. The recognition and cutting of mismatches by TKO NucS highlight the difference between this novel MMR system and that of MutSL, where MutS recognizes the mismatch, but a single-stranded DNA nick is made either by MutH directed to GATC sites on the unmethylated strand of the replication fork (*e.g.,* in *E. coli*), or in other bacteria and eukaryotes, by a β-clamp activation of a latent MutL endonuclease that nicks the newly synthesized strand. For recent reviews on NucS endonuclease function and strand discrimination in MMR, see (15,16).

Of particular interest is whether the novel NucS MMR system exhibited in the archaea described above is active as an anti-mutator in Actinobacteria where the

gene is present, including the clinically relevant bacterium *Mycobacterium*

tuberculosis. The question was addressed by two labs (17,18) that studied

actinobacterium *Corynebacterium glutamicum* and found that deletion of the *nucS*

homolog (NCgl-1168, hereafter called $NucS_{Cg}$) resulted in a strain with a mutator

phenotype. Purification of the $NucS_{Cg}$ protein showed that it bound to DNA

containing base pair mismatches, and that cutting of these substrates was activated

by an interaction with the β-clamp from *C. glutamicum*, $DnaN_{Cg}$. The mismatch

recognition and cutting specificities of $NucS_{Cg}$ were the same as those observed for

TKO NucS and was consistent with the mutation spectrum of *C. glutamicum* in

mutational accumulation (MA) studies done in the *nucS*-deficient background, *i.e.,* a

high levels of transitions observed relative to wild type (18).

This question of whether NucS from a mycobacteria species could bind

mismatched DNA was addressed by Castañeda-Garcia et al (19) who examined

thousands of transposon mutants in *Mycobacterium smegmatis* for high rates of

spontaneous resistance to rifampicin. They identified MSMEG_4923 as a gene

responsible for generating a mutator phenotype in *M. smegmatis*; this gene product

has 27% sequence identity with NucS from *P. abyssi* and was named NucS.

However, isolation and characterization of the *M. smegmatis* NucS protein showed

that while it bound to ssDNA and not dsDNA (like its *P. abyssi* homolog), it did not

demonstrate binding activity to mismatch containing dsDNA substrates in vitro.

This result contrasts with those from the NucS proteins from *P. furiosus* and *T.*

kodakarensis in vitro*,* raising the question of whether the mycobacterial NucS acts in

vivo in a similar way to how these proteins function in vivo.

We have addressed this question here genetically by examining if *M. smegmatis* NucS requires an interaction with the β-clamp in vivo, as was seen previously with the NucS protein from *T. kodakarensis* in vitro.   Our results of complementation studies with plasmid-encoded wild type NucS and a derivative containing a C-terminal 5 amino acid truncation suggest an interaction with the *M. smegmatis* β-clamp is required for MMR activity in vivo. Secondly, we examined the mismatch specificity of *M. smegmatis* NucS in vivo and compared it to the mismatch specificity of *T. kodakarensis* reported in vitro. We used Che9 RecT-promoted oligo recombineering in *M. smegmatis* to directly deliver all possible types of base mismatches (except C/C) to a hygromycin-resistant (Hyg^R) gene reporter construct integrated into the chromosome. We find that the specificity of *M. smegmatis* mismatch repair in vivo matches that found previously for *T. kodakarensis* in vitro, suggesting a commonality in the NucS-promoted MMR systems in these bacteria. Finally, we used oligo-mediated recombineering to delivery defined mismatches to the defective hygromycin-resistant target in the chromosome and find that a cut induced by NucS at a G-G mismatch is processed by a 5'-3'exonuclease in vivo, which is limited in extent to 8-10 base pairs, and that repair is independent of RecA, RadA, and non-homologous end joining functions Ku and LigD.

RESULTS

Mutator phenotype of *M. smegmatis* nucS strain and testing of the PIP domain

Using the recently describe ORBIT method for the generation of chromosomal modifications in mycobacteria (see Materials and Methods), a deletion mutant of *M. smegmatis nucS* gene (MSMEG_4923) was constructed. The frequency of spontaneous

mutation to rifampicin resistance was measured in both wild type *M. smegmatis* and its

ΔnucS derivative (see Fig. 1). As reported previously (19), an *M. smegmatis* strain

deleted of *nucS* exhibited a mutator phenotype, with a greater than 100-fold increase in

the frequency of spontaneous mutations when compared to wild type.

In complementation experiments, the expression of NucS from pIR540

reduced the mutation frequency considerably (about ~12-fold), close to that observed

in WT cells. However, a similar construct expressing a NucS protein deleted of the last

five amino acids from the C-terminus (NucS-ΔC5) failed to complement the *nucS* strain,

resulting in high levels of mutagenesis (see Fig. 1B).   This region of *M. smegmatis* NucS

is homologous to the C-terminal region in *P. abyssi* NucS that has been shown to

interact with the PCNA sliding clamp (thus called the PIP domain, for PCNA interacting

protein) (12). This region is also identical to the C-terminal 5 amino acids of NucS

from *C. glutamicans*, which required an interaction with β-sliding clamp to promote

dsDNA cutting of mismatched-containing substrates in vitro (17,18). The failure of the

M. smegmatis NucSΔC5 to suppress the mutagenic phenotype of the ΔnucS mutant

suggests that an interaction of the C-terminal PIP domain in *M. smegmatis* NucS with

the β-sliding clamp at the replication fork is also required for efficient NucS-promoted

cutting in vivo, and thus subsequent MMR.

It is possible that the loss of the C-terminal 5 amino acids causes *M. smegmatis*

NucS protein to become unstable resulting in a mutagenic phenotype. To address this

issue, we performed ORBIT to tag *nucS* to be able to follow the protein and its

truncated version by Western analysis. However, C-terminal chromosomal fusions of

wild type *nucS* with either Flag-DAS or GFP tags resulted in strains with mutagenic

phenotypes, presumably by interfering with the interaction of the NucS PIP domain

and the β-clamp. Tagging of chromosomal *nucS* at the N-terminus was then performed using a 6x-His tag, but this modification also resulted in a strain that exhibited a mutagenic phenotype, suggestion the His tag may have interfered with dimerization of NucS or the binding of a protein partner in vivo. These results indicate that an interaction between a free-ended C-terminus PIP domain of *M. smegmatis* NucS and the β-clamp is required for efficient mismatch repair in vivo, and likely explains the inability of purified *M. smegmatis* NucS alone to recognize and cut mismatched dsDNA substrates in vitro, as was observed in a previous study (19).

Mismatch DNA repair specificity in vivo

Previous mutational accumulation studies in both *C. glutamicans* and *M. smegmatis* have shown that the absence of the NucS MMR system results in an a greatly elevated rate of transitions, both G:C > A:T and A:T > G:C (the latter one being the higher of the two, opposite to what is observed in WT cells). In particular, Castañeda-García *et al* (20) used an integrated plasmid reporter system to measure the mutational rates (and specificities) of both transitions and transversions using mutant versions of *aph* (kan[R]) alleles. With their reporter strains, the fold increase in mutation rates (ΔnucS/WT) was highest (65-fold) for transitions that required repair of a G-T mismatch to generate a Kan[R] cell. In contrast, the fold increase in transversion mutations on their reported plasmids was low, including ones that are predicted to generate G-G and T-T mismatches (2.4 and 3.6 -fold respectively), mismatches that are recognized and cut efficiently by *C. glutamicans* NucS in vitro. The fold rates, while low, are higher than those seen with plasmids that generate G-A and T-C mismatches (1.0 and 1.5 respectively), which are not recognized by NucS

in vitro. The low rate of NucS-dependent activity on transversion repair, despite efficient cutting of G-G and T-T mismatches in vitro, may be simply due to the low levels of G-G and T-T mismatches that arise during DNA replication in mycobacteria, in a supposed background containing high levels of G-T mismatches. Or it may mean that NucS does not act on these mismatches at a high efficiency in vivo, despite highly efficient cutting of them in vitro.

We were thus encouraged to set up a similar assay to examine the mycobacterial mismatch repair system in vivo, to compare the specificity of mismatch recognition to what has been reported in vitro. We thus tested the ability of Che9 RecT-mediated oligonucleotide recombineering to generate defined mismatches in a way that would allow us to follow the fate of mismatches by simple plating assays. A first test of this system was to deliver mismatches to the replication fork within the *rpsL* gene (21,22). Oligos were designed to generate either K43R or K43N changes to the *rpsL* ribosomal protein, both of which are known to confer resistance to streptomycin. The K43R oligo creates a C/A mismatch at the replication fork, while the K43N oligo generate at G/G mismatch (see Fig. 2). According to mismatch specificity reported for *T. kodakarensis* NucS in vitro, the K43R oligo would create a mismatch in *M. smegmatis* that would not be subject to NucS-mediated repair (C/A) and thus generate high levels of streptomycin-resistant colonies. Along the same lines, the K43N oligo would generate a mismatch that is subject to repair (G/G) and thus fail to generate streptomycin-resistant colonies. These predictions were fulfilled by the data shown in Table 1, showing that recombineering can be used to deliver mismatches to test

in vivo specificities, and that *M. smegmatis* shows a similar mismatch repair specificity as exhibited in vitro by *Actinobacteria*, but do so in vivo. The same experiment was performed in *M. tuberculosis* with similar results (Fig. S2).

To further test the in vivo specificities of mismatch repair, a similar assay was designed to study all the other types of mismatches. For this purpose, we inserted a stop codon (TAG) in place of the glycine codon (GGA) at position 110 in the hygromycinB phosphotransferase gene (*hyg*) in plasmid pIR542, a Giles integrating plasmid. The G110 residue was selected because of its presence in an extended loop on the surface of the protein (see Fig. S1A). In addition, the G-110 residue is not conserved among *hyg* gene from other bacterial species (Fig. S1B), highlighting this position as one that is likely tolerant to multiple types of amino acid substitutions.

Oligos were designed to convert the stop codon in integrated plasmid pIR542 to various sense codons, allowing for full translation of the *hyg* gene. Each oligo generates a different type of mismatch (see Table S1). If a mismatch escapes repair, it leads to a recombinant that expresses a functional phosphotransferase and generates Hyg[R] colonies; on the contrary, if the mismatch is repaired, no or few Hyg[R] transformants would be observed.

M. smegmatis containing pKM402 (RecT producer) and pIR542 were electroporated with the oligos listed in Table S1 and the outgrowths were plated on LB plates with and without hygromycin, as described in Materials and Methods; the results are shown in Fig 3. In otherwise wild type *M. smegmatis,* oligos that generated G-G, G-T and T-T mismatches failed to generate Hyg[R] colonies beyond the

frequency shown by a control transformation with no oligo, indicative of repair of these mismatches in vivo. All the other mismatches tested escaped MMR, generating Hyg[R] colonies at a frequency 100-fold greater than the control transformation (Fig. 3A). When the same experimental protocol was performed in the *M. smegmatis* Δ*nucS* strain, transformation with all the oligos generated high frequencies of Hyg[R] colonies, revealing that in the absence of MMR, all mismatches generated were incorporated into the chromosome (Fig. 3B).   These results reveal that *M. smegmatis* shows the same mismatch repair specificity in vivo as the archaeal TKO strain exhibited in vitro, indicating that the two systems recognize identical types of mismatches and likely behave by similar mechanisms. Additionally, the G-T mismatch recognition specificity revealed here is consistent with the mutational specificity of *M. smegmatis* reported by Castaneda-Garcia et al. (20), who showed a high rate of transitions when the reporter plasmids generated G-T mismatches in order to restore kanamycin resistance. On the other hand, G-G and T-T (which if unrepaired lead to transversions), do not show high levels of NucS-dependent repair in mutational accumulation studies, even though NucS recognized and repaired G-G and T-T mismatches in vivo at the same rate as G-T mismatches (Fig. 3A), This result is further evaluated in the Discussion section

Inability of *M. smegmatis* to NucS to repair small indels.

A test to examine the in vivo repair capacity of *M. smegmatis* NucS on small indels was done with our recombineering assay, examining the ability to remove 1-2 bp insertions and deletions in the *leuB* gene. Such indels inactivate *leuB* activity and recombineering with oligos that removed these indels allow this strain to grow on

7H10 plates without leucine supplementation. If NucS is inactive on DNA containing 1-2 bp indels, such oligo-mediated leuB⁺ strains will be easily observed.   This was, in fact, the case, as insertions and deletions between 1-2 bp were easily removed from these leuB frameshift mutants using oligos the contained the wild type *leuB* sequence, restoring the defective *leuB* allele back to wild type at high frequencies (see Fig. 4). The lack of activity of mycobacterial NucS on indels in this recombineering assay is in agreement with the MA experiments and reporter plasmids previously described (20). This result, and those in Fig 3, reveal that delivery of mismatch by oligo-mediated recombineering faithfully mimics bona fide DMA mismatches in vivo.

Identification of likely NucS nucleolytic active site residues

Nakae *et al* (14) solved the structure of the NucS protein from *P. abyssi* and identified three amino acid residues that may be involved in nucleolytic activity (Fig. 5A). In comparison with the *P. abyssi* NucS protein sequence, the corresponding amino acids in *M smegmatis* NucS are D138, E152 and K154. Using Che9 RecT-mediated oligo recombineering, each of these residues were separately changed to alanine and the subsequent strains were tested for mutagenic activity by plating on 7H10 plates containing rifampicin. The D138A, E152A, and K154A mutations each conferred a mutagenic phenotype to *M. smegmatis*, implicating all these residues in NucS endonucleolytic activity (Fig. 5B). Confirmation of these residues as nucleolytic active site residues will require biochemical analysis of the mutant proteins.

Resection of NucS-promoted cuts in vivo involves limited processing by a 5'-3' exonuclease

The identification of nucleolytic active site residues in *M. smegmatis* NucS that lead to high mutability suggests that cutting in vivo likely occurs in the processing of mismatches in mycobacteria. Whether this involves a ssDNA nick (as seen with the MutSLH system) or a dsDNA break (as suggested by in vitro results from *C. glutamicum* NucS$_{Cg}$) has not currently been established in vivo. In either case, we sought to examine the processing of such cuts by endogenous exonucleases in vivo by delivering two mismatches (with one oligo) to the chromosome via recombineering, with one mismatch (C-T) generating Hyg^R mutants (as described above) and the other mismatch (G-T) being recognized and cut by NucS. The oligo sequence is shown in blue in Fig. 6B and generates mismatches that are 32 bp apart on the lagging strand. It is assumed that exonucleolytic processing would occur from the NucS-induced cut at codon 99 to codon 110 of the Hyg^R gene. If processing by a 5'-3'exonuclease proceeds past codon 110 of the Hyg^R gene (Fig. 6C), subsequent repair by resynthesis of the gap (or recombination repair) would lead to loss of the C-T mismatch and restoration of the stop codon. This, in turn, would lead to loss of Hyg^R transformants relative to the use of an oligo that did not generate a repairable mismatch (control oligo), but still generates the C-T mismatch. On the contrary, limited exonucleolytic processing from the G-T mismatch at codon 99 that did not proceed to codon 110 would not affect the level of Hyg^R transformants compared to the control oligo (Fig. 6D).

A first test of this two-mismatch-promoting oligo design showed that it could generate Hyg^R transformants at the same frequency as a control oligo that generated the

C-T mismatch but not the G-G mismatch ($\sim 5 \times 10^{-4}$ Hyg^R transformants/total cell titer).

Sequencing of PCR fragments containing this region from 8 Hyg^R colonies found that all

the G-T mismatches were repaired (resulting in a "C" in the replicating strand). This

result shows that exonucleolytic processing from the NucS-promoted cut at the G-T

mismatch did not extend 32 bases to the C-T mismatch, leaving the frequency of Hyg^R

colony formation intact relative to the control oligo (no G-G mismatch).

Given the result above, we performed the same test with a series of oligos the

varied the distance between a repairable mismatch (this time G-G) and the C-T

mismatch that confers Hyg^R. Oligos containing both a G-G and C-T mismatch were

electroporated into *M. smegmatis* containing the defective *hyg* gene; see Fig. 7A for

positions of the G-G mismatches relative to the C-T mismatch in each of the oligos.

We followed the frequency of Hyg^R colonies from the outgrowth when the two

mismatches were relatively close (4-5 bp) or further apart (16 or 20 bp), as well as

oligos where the repairable mismatch was downstream of the C-T mismatch at

variable distances (see Fig.7A). At a distance 20 bp upstream of codon 110, a G-G

mismatch-guided NucS-promoted cut did not lead to loss of Hyg^R colonies relative to

the oligo with no G-G mismatch. This result suggests that 5'-3' processing of the

predicted dsDNA cut in the top strand did not extend 20 bp to the C-T mismatch,

which would have resulted in loss of Hyg^R colonies relative to the oligo with no G-G

mismatch (similar to the result described above). On the other hand, when the G-G

mismatch was only 5 bp away from the C-T mismatch, there was a 100-fold drop the

frequency of Hyg^R colonies. This result suggests that a dsDNA break at the G-G site

close to the C-T mismatch resulted in 5'-3' exonucleolytic processing of the top

strand that extended to the C-T mismatch, removing the "C" base, with eventual

restoration of an "A" at this position to restore the stop codon (leading to loss of Hyg^R colonies) (see Fig 7B).

When the position of the G-G mismatch was designed to be downstream of the C-T mismatches (three positions shown on Fig. 7A), the assay can follow the exonucleolytic processing of the 5' overhang in the bottom strand (Fig 7B and 7D). In this case, exonucleolytic processing of the bottom strand (toward the C-T mismatch) would remove the "T" base, leaving the "C" in the top strand, securing the replacement of the stop codon with a TCG codon (Arg), leading to high levels of Hyg^R. Thus, in this configuration, one would expect no (or less) loss of Hyg^R colonies with any oligo containing a G-G mismatch downstream of the C-T mismatch. This outcome is what was observed for oligos 8 and 15 bp downstream of the C-T mismatch, and partially true when the G-G mismatch was positioned only 4 bp downstream from the C-T mismatch (see Fig. 7B and 7D). In this latter case, there was a ~10-fold drop in Hyg^R colonies, instead of the 100-fold drop seen when the G-G mismatch was located on the other side of the C-T mismatch)

The key observation from the above assay is that processing of the break induced by NucS is principally performed by a 5'-3' exonuclease. This can be seen with the 8-fold higher loss of Hyg^R colonies with G-G at the -5 position (relative to C-T) compared to the +4 position. If processing were performed by 3'-5' exonucleases, the opposite would have been expected to be observed. Furthermore, given the 5' overhangs predicted from a NucS-promoted dsDNA cut made at the G-G mismatch at the +4 position, nibbling of only 2 bp of the 3' end would result in loss of Hyg^R transformants. However, processing of the G-G mismatch at the -5 position requires at least 8 base pairs to be excised before loss of any Hyg^R colonies is observed.

Nonetheless, Hyg^R colony loss occurs at an 8-fold higher frequency at the -5 position relative to the +4 position.

These results support a model of 5'-3' exonucleolytic processing of the dsDNA break at the site of a NucS-induced cut at a mismatch in vivo. What is also revealing is that while there is a gradual increase in Hyg^R colonies as the G-G mismatch is moved from the -5 to the -13 position, at -20 bases away the frequency of Hyg^R transformants is equal to the frequency seen with the control oligo (*i.e.*, no G-G mismatch). This result suggests that while there is 5'-3' exonucleolytic processing from the NucS-induced cut, it is very limited to a small patch of DNA and inconsistent with the at least ~200 bp to 1 kb of processing of ssDNA that usually accompanies RecA-promoted dsDNA break repair events. Such short patch repair is reminiscent of a similar pathway of mismatch repair events promoted by MutS in *E. coli*. While these results have been presented in Figure 7 in the context of NucS making a dsDNA cut at the G-G mismatch (consistent with in vitro data of actinobacterial NucS$_{Cg}$ protein), the same conclusions can be drawn if NucS promoted a nick in the replicative strand (blue strand in Fig. 6B).

Repair of NucS-promoted ds-DNA breaks is not dependent on RecA- or RadA-promoted homologous recombination, or NHEJ

A second test of the involvement of homologous recombination in NucS-promoted MMR is to examine if the double-mismatch-promoting oligo described above leads to Hyg^R transformants in a *recA* mutant host. If a dsDNA cut is indeed generated in vivo at the G-G mismatch, and recombination repair acts on that break, the absence of RecA protein would be expected to lead to an unrepaired dsDNA break and death of that

cell, resulting in loss of Hyg^R transformants relative to a WT host. Along the same line of reasoning, if the processing of the dsDNA cut (or nick) at the G-G mismatch was impaired by a mutation in a resident 5'-3' dsDNA or flap endonuclease, it likely would lead to some level of interruption of the *hyg* reading frame by loss of base pairs other than 3 (or multiples of 3), again leading to loss of Hyg^R transformants relative to WT. Thus, mutant strains of *M. smegmatis* in genes suspected of being involved in MMR were transformed with the defective *hyg* gene and electroporated with a double mismatch-generating oligo where the C-T and G-G mismatches were separated by 32 base pairs (oligonucleotide *Hyg-repair-32* in Table S2). Loss of Hyg^R transformants in a mutant strain, compared to a control oligo not generating the G-G mismatch, would indicate involvement of that gene in mycobacterial MMR.

Besides RecA, the *M. smegmatis* RadA was also tested in this assay. RadA is an archaeal analog of the RecA recombinase and can form nucleoprotein filaments on DNA and catalyze DNA pairing and strand exchange (23,24). RadA has a role in stimulating branch migration that extends heteroduplex formations in RecA-mediated strand transfer reactions (25,26). Mutants of RadA in *E. coli* have strong synergistic phenotypes with *recG* mutants, a branch migration enzyme, and show synthetic genetic interactions for survival to AZT when combined with several known helicases including PriA, RuvAB, UvrD, and others (27). B. subtilis RadA has also been shown to possess a 5'-3' helicase activity capable of unwinding recombinational intermediates (28). The *M. smegmatis radA* gene seemed especially relevant to a study of *nucS* function since both these genes are of archaeal origin and in one genus of archaea (*Thermococcus*), *nucS* is co-transcribed with *radA* (12,13). Other functions in *M. smegmatis* were also tested in this assay and included a strain

mutant in both Ku and LigD, to see if NHEJ had a role in mycobacterial MMR, and the

MSMEG_3883 gene which encodes *fenA*, a flap endonuclease involved in processing

of 5' overhangs. Results of this experiment are shown in Fig. 8.

In support of the data of limited resection shown in Fig. 7, neither *recA*,

radA, or the double mutant had any effect on the level of Hyg^R transformants

following transfer of the oligo generating the G-G mismatch relative to the control

oligo (where a G-G mismatch is not generated). Thus, recombinational repair is not

involved NucS-promoted MMR. The results shown in Fig 8 also suggests that

neither NHEJ functions LigD and Ku (MSMEG_5570 and MSMEG-5580) or the FenA

flap endonuclease nuclease (MSMEG_3883) are involved in mycobacterial mismatch

repair, as none of these functions lead to loss of Hyg^R transformants with the oligo

that generates the G-G mismatch relative to the control oligo. Other functions

thought to be important players in NucS-promoted MMR in mycobacteria, such as

other dsDNA exonucleases and helicases, are currently being examined using this

double-mismatch oligo recombineering assay.

DISCUSSION

The discovery of the EndoMS/NucS functions, principally found in archaea

and Actinobacter species, revealed that a new pathway for DNA mismatch repair

had been identified that did not follow the nearly ubiquitous paradigm of MutS DNA

mismatch recognition, followed by MutL (or MutH) nicking of the duplex strand

containing the "wrong" base. Instead, in vitro characterizations of the NucS proteins

from the archaeal species *T. kodakarensis* and the Actinobacter bacterium *C.*

glutamicans revealed that while they recognized substrates containing mismatched substrates, these proteins proceed to make cuts in both strands in DNA mismatched-containing substrates. The cutting of dsDNA containing mismatches with *T. kodakarensis* NucS showed that the cut left five nucleotide 5'-overhangs and recessed 3'-OH ends leaving ligateable sticky ends characteristic of restriction enzymes. Further support of the dsDNA cutting capability of NucS comes from the Xray structural analysis of the NucS protein from the *T. kodakarensis* which clearly has structural characteristics of a restriction enzyme: a homodimeric structure consisting a two domains, an N-terminal domain acting as a dimerization function that recognizes the mismatched bases that are flipped out (like restriction enzyme Ecl18kl), and a C-terminal domain that contains the active site residues for cutting dsDNA in a manner reminiscent of type II restriction enzymes (14). These studies strongly suggest that NucS acting in archaea and actinobacterial species act on mismatched bases by creating dsDNA break at the site of the mismatch, though this cutting has not been demonstrated in vivo.

A first step to examine the capability of NucS in mycobacteria to promote such a dsDNA break was investigated by Castenda-Garcia et al (19), who showed that an *M. smegmatis* ΔnucS strain showed a mutagenic phenotype when plated on rifampicin plates, which we confirmed here. However, their biochemical analysis revealed that while NucS bound to ssDNA, it did not bind to or cut substrates containing DNA mismatches, leaving open the possibility that NucS from mycobacteria might not behave in a similar manner to NucS from *C. glutamicans.* The most likely scenario for their failure to see cutting by *M. smegmatis* NucS was

that, like NucS from *C. glutamicans,* it requires an interaction with the β-clamp. To this end, our ability to complement the mutagenic phenotype of *M. smegmatis* ΔnucS with full length NucS, but not one missing the last 5 amino acids containing the PIP domain, is consistent with this supposition. Verification of this interaction between *M. smegmatis* NucS and its β-clamp will require biochemical analysis of these functions.

We then leveraged the oligo-mediated recombineering capabilities of *M. smegmatis* to deliver defined mismatches to directly to the mycobacterial chromosome in order to examine the mismatch specificity of NucS MMR. We found that mismatches containing G-T, G-G, and T-T were specifically repaired using the NucS MMR pathway, while none of the other mismatches were corrected. The mismatch recognition specificities revealed here *in vivo* are consistent with previous specificities shown in *C. glutamicans* NucS$_{Cg}$ in vitro. (17,18). They are also consistent with the *mutational* specificity of *M. smegmatis* reported by Castaneda-Garcia et al (20), who showed a high rate of both G:C > A:T and A:T > G:C transitions in *M. smegmatis nucS* strains (relative to wild type strains) using mutational accumulation experiments. This is consistent with the recognition of G-T mismatches by NucS, which would prevent the accumulation of transitions mutation in wild type cells.

What remains somewhat of a mystery is that while large increases in transversions are not observed in nucS mutant MA studies (3-fold increase at best relative to wild type), mismatches of G-G and T-T that would lead to such transversion mutants are recognized and cut by NucS both in vitro and in vivo at

essentially the same efficiency at G-T mismatches. This discrepancy may be resolved by the supposition that G-T base pairs are by far the most highly generated mismatch, the repair of which has evolved to be the most efficient, thus leading to NucS targeting G-T as the primary mismatch to deal with. It is only when G-G and T-T mismatches are generated in the absence of multitudes of G-T mismatches (such as in vitro or here using a recombineering methodology) that NucS has the opportunity to efficient recognize and cut them.

To start to unravel mechanistic details of NucS MMR, we used oligo-mediated recombineering to examine exonucleolytic processing that takes place in the chromosome following the cutting of the chromosome at a repairable mismatch. This was done by delivering an oligo containing at G-G mismatch by recombineering and measuring either loss or retainment of a nearby mismatch (C-T) as determined by the frequency of Hyg^R colonies following plating of the outgrowth. The assay, however, cannot distinguish between whether a nick or a dsDNA cut is made at the site of the mismatch, as this recombineering approach use can only measure the activity of endogenous exonucleases on the strand containing the mispaired bases supplied by the oligo (the top strand of the oligo highlighted in blue in Fig. 6B).

However, by comparing the number of bases needed to process that top strand by either a 5' end or a 3' end, it is clear that the 5' end is processed more efficiently than the 3' end. This can be seen by recognition that the 5' end in Fig. 7C needs 8 bases removed from the top strand before it reaches the T-C mismatch (and thus capable of lowering the frequency of Hyg^R colonies; see the -5 oligo in Fig 7B). Significantly, it is down more than 100-fold in Hyg^R frequency relative to a 3' end digestion of the top strand with mismatches generated by oligo +8 in Fig. 7D, which

requires only 6 base pairs to reach the C-T mismatch. Thus, 5'-3' exonucleolytic

processing is heavily favored at the site of a NucS-promoted nick or dsDNA break.

However, what is more telling in this experiment is the limited nature of the

processing of the 5'-3' exonuclease following the cut made by NucS at a mismatch.

By examining the extent of loss of Hyg[R] transformants among the oligos that

generate mismatches that confer Hyg[R] (C-T) and a substrate for NucS cutting (G-G),

oligos "-5" to "-20"in Fig.7B, one can observe that processing of the NucS-induced

cut by a 5'-3' exonuclease in most cases extends to 8 bases from the cut site

(including the five base 5' overhang). Digestion of an additional 3 bases (11 total) is

diminished 75-fold, as evidenced by a 75-fold increase in the number of Hyg[R]

transformants seen with oligo "-8" in Fig. 7B.   Ultimately, the use of an oligo where

the distance between the C-T and G-G mismatches generated at the replication fork

is 20 bases (oligo "-20"), no loss of Hyg[R] transformants is observed relative to the

control oligo. These results suggest that digestion of the 5'-3' strand from the site of

a NucS cut (or nick) involves at least 8 base pairs, but not more than 20 base pairs.

This result means that if a dsDNA cut is made by NucS at a mismatch in vivo, it is not

further processed by RecA-promoted homologous recombination events, as such

recombination requires much larger sequences of homology (~200 bases) relative

to what is observed in here.

The use of a double-mismatch generating oligo was also used to test the

genetic dependency of NucS MMR. It is reasoned that if a NucS-promoted dsDNA cut

is made in vivo at a mismatch, then inactivating mutations of any downstream

functions would lead to non-repair of the dsDNA cut, and subsequent death of the

cell. In a similar vein, if only a nick is made at the mismatch by NucS, any functions

required for exonucleolytic processing of the nick would likely lead to some

frequency of indel formation and thus to loss of Hyg^R transformants. Our results

showed that neither RecA nor RadA is involved in downstream processing of the cut

made by NucS, consistent with the limited processing of cuts made in vivo shown in

Fig. 7. We next examined if the non-homologous end-joining functions Ku and LigD

might somehow be involved MMR in M. smegmatis. Since mycobacteria exhibit the

uniqueness of possessing both the novel NucS MMR system and NHEJ functions

(neither present in most bacteria), it was hypothesized that NHEJ, together with

restriction enzyme-like NucS, might work together to repair mismatches following a

NucS promoted dsDNA cut, in a way the exhibits higher fidelity of dsDNA breaks

independent of NucS. However, we found no effect on the frequency of Hyg^R

transformants in the double-mismatch generating oligo in a strain deleted of both

Ku and LigD, showing that NHEJ functions are not involved in MMR in mycobacteria.

Finally, the limited processing of the cut made in vivo was highly suggestive of a flap

endonuclease, where a cut is made at the ssDNA-dsDNA junction to remove a 5'

single-stranded DNA flap. In this scenario, one imagines that NucS cutting in vivo is

limited to only the replicative strand (by some unknown function) and that

unwinding from the nick is limited to the generation of a 5' flap containing the

mispaired base. Subsequent cutting of the flap by a resident endonuclease was

suspected. However, the deletion of the gene that encodes the FenA 5- flap

endonuclease (MSMEG_3883) had no effect on the frequency of Hyg^R transformants

in our double-mismatch generating oligo assay, ruling out a role of *fenA* in NucS-

promoted MMR, or that some other exonuclease in vivo can substitute for FenA in

its absence. In these types of cases, it may require double (or multiple) mutants to

be generated to see if processing of the NucS-induced break can be carried out by overlapping functions.

Thus, downstream components of the NucS MMR pathway remain elusive, as we continue to use this assay to examine the role of other nucleases and helicases known to exist in the mycobacterial genome that may have a role in high fidelity repair of mismatched bases in the wake of the replication fork.

MATERIAL AND METHODS

Bacterial strains and media

The *M. smegmatis* strains used in this study were derived from mc^{2}155; the *M. tuberculosis* strains was derived from H37Rv. *M. smegmatis* and *M. tuberculosis* were grown in 7H9 broth with 0.05% Tween 80, 0.2% glycerol, and OADC (oleic acid-albumin-dextrose-catalase; Becton, Dickinson); transformants were selected on 7H10 plates with 0.5% glycerol and OADC. Deletion mutants of *M. smegmatis* were constructed using ORBIT (29); oligonucleotides used in the ORBIT procedure are listed in Table S2. Verification of each ORBIT-promoted deletion were done by colony PCR by amplification of both junctions between the inserted plasmid and the chromosomal targets site, as well as the absence of an amplicon using PCR primers internal to the target gene (compared to WT cells). *M. smegmatis* strains MGM199 (Δ*recA*) and MGM156 (ΔKu and Δ*ligD*) were gifts kindly received from Michael Glickman (30). A list of mutants constructed and used in this study is shown in Table 2. Construction of *leuB* mutants containing single or two base pair indels and active site mutants of *nucS* were done using oligo-mediated recombineering as described previously (21). *M. smegmatis* strains that contained a defective

chromosomal hygromycin-resistant gene were constructed by either transformation

with a zeocin-resistant Giles integrating vector pIR542 or by transformation with

the zeocin-resistant ORBIT integration vectors pKM611, pKM614, or pKM585.

When needed, the following antibiotics concentrations were added to 7H9 media or

7H10 plates: kanamycin (20 μg/ml), streptomycin (20 μg/ml), hygromycin (50

μg/ml), and zeocin (25 μg/ml).

Plasmids

Oligo-recombineering plasmid pKM402, Che9 RecT and Bxb1 Integrase plasmid

pKM461, and Orbit integration plasmids have been described previously (29,31).

Plasmid pIR540 is a Giles integration vector that expresses full length *M. smegmatis*

nucS gene (MSMEG_4923) from the PgroEL promoter. Plasmid pIR541 is similar to

pIR540 except that is missing the last five codons of the *nucS* gene. Plasmid pIR542

is a Giles integration vector (Zeo^R) that contains a defective hygromycin-resistant

gene where the codon of glycine 110 (GGA) has been altered to stop codon (TGA).

Plasmid pKM611 is an Bxb1 *attB*-containing ORBIT-integration vector that contains

the same defective hygromycin-resistant gene from pIR542; pKM614 is the same as

pKM611, but with the defective-hygromycin-resistant gene in the opposite direction

relative to the Bxb1 *attB* site; pKM585 is a streptomycin version of pKM611.

Details of plasmid constructions are available upon request.

Oligonucleotides

Oligonucleotides used for the determination of mismatch specificity ate listed in

Table S1. Oligonucleotides used for mutant construction by ORBIT are listed in

Table S2 and were obtained from IDT as Ultramers at a concentration of 100 μM

(delivered in 96-well plates). They were supplied desalted with no further

purification and diluted 10-fold in 10 mM Tris-HCl (pH 8.0).   Final concentrations

(250 to 350ng/ml) were determined by absorbance maxima at 260 nm (Abs_{260}).

Oligos for plasmid constructions and recombineering experiments were obtained

from Life Science Technologies and were used at concentrations as described in the

figure legends; they are listed in Table S2.

Transformations

Details of the transformation procedure using the ORBIT technology for mutant

strain construction has been described previously (29). Electroporations for

oligonucleotide-promoted recombineering experiments have been described

previously (21). Briefly, 150 ul of fresh overnight culture of *M. smegmatis* was

added to 20 ml of 7H9-OADC-tween containing 20 μg/mL of kanamycin and swirled

at 37^{o}C overnight. The next morning, when the culture reached an absorbance of

0.5, anhydrotetracycline (Atc) was added to a final concentration of 500 ng/ml. The

culture was allowed to grow for an additional three hours or until the O.D. reached

1.0. Cells were then collected by centrifugation, placed on ice, and washed twice

with 20 mL of cold 10% glycerol. After the final centrifugation, the cells were

resuspended in 2 ml 10% glycerol. Oligonucleotides (2 μg) were placed in

Eppendorf tubes and 380 μL of electrocompetent cells were mixed with oligos then

transferred to 0.2 cm electroporation cuvettes. Cells were shocked and transferred

to 2 ml 7H9-OADC-tween and grown overnight at 37^{o}C. Appropriate dilutions of

overnight cultures were plated on 7H10-OADC plates (+/- 50 ug/ml hygromycin) to determine the fraction of Hyg^R transformants.

Mutagenicity Assay

The lack of NucS repair activity in M. *smegmatis* was determined by plating 200 μL of a saturated culture (O.D. of 1.0-1.5) on 7H10 plates containing 150 ug/ml of rifampicin; total cell numbers were determined by plating on 7H10 plates. Plates were incubated at 4-5 days at 37°C. The frequency of spontaneous mutation was determined by the titer of rifampicin-resistant colonies divided by the total cell titer.

Recombineering with oligonucleotides generating two mismatches

To examine the extent of resection after NucS cutting, we designed oligos to create G-G and C-T mismatches separated by 4 to 32 base pairs along the chromosome, allowing us to measure the extent of processing of the C-T mismatch as a function of the distance (and position) from the NucS-proposed dsDNA cut (or nick) at the G-G mismatch. Oligos containing these two base pair changes were electroporated into *M. smegmatis* containing pKM461 following induction with anyhydrotetracycline, as described above. After overnight growth, portions of the culture were plated on 7H10 and 7H10 plates with 50 μg/ml hygromycin to determine the frequency of Hyg^R transformants.

Mutant studies

Strains containing deletions of *radA* and *fenA* were constructed by ORBIT using attB-containing integrating plasmids that also contained a defective *hyg* gene (either pKM611 or pKM614). In each case, the *hyg* genes are positioned in the chromosome

so that the oligo shown in Fig. 7A would always target the lagging strand template of the replication fork. These strains retained the pKM461 plasmid used in the ORBIT procedure, which is proficient for the subsequent oligo-mediated recombineering experiment.

ACKNOWLEDGEMENTS

We wish to acknowledge Michael Glickman for the *recA* knockout strain and the *ku-ligD* double mutant. We also like to thank Christopher Sassetti and Kadamba Papavinasasundaram for material and managerial support.

FUNDING

This work was supported in part by the Bill and Melinda Gates Foundation grant # INV-040487.

CONFLICT OF INTEREST

Authors declare no conflicts of interest.

REFERENCES

1.   Bębenek, A. and Ziuzia-Graczyk, I. (2018) Fidelity of DNA replication-a matter of proofreading. *Curr Genet*, **64**, 985-996.
2.   Echols, H. and Goodman, M.F. (1991) Fidelity mechanisms in DNA replication. *Annu Rev Biochem*, **60**, 477-511.
3.   Ganai, R.A. and Johansson, E. (2016) DNA Replication-A Matter of Fidelity. *Mol Cell*, **62**, 745-755.
4.   Modrich, P. and Lahue, R. (1996) Mismatch repair in replication fidelity, genetic recombination, and cancer biology. *Annu Rev Biochem*, **65**, 101-133.
5.   Kunkel, T.A. and Erie, D.A. (2015) Eukaryotic Mismatch Repair in Relation to DNA Replication. *Annu Rev Genet*, **49**, 291-313.
6.   Lu, A.L., Clark, S. and Modrich, P. (1983) Methyl-directed repair of DNA base-pair mismatches in vitro. *Proc Natl Acad Sci U S A*, **80**, 4639-4643.
7.   Fishel, R.A. and Kolodner, R. (1984) An Escherichia coli cell-free system that catalyzes the repair of symmetrically methylated heteroduplex DNA. *Cold Spring Harb Symp Quant Biol*, **49**, 603-609.
8.   Jiricny, J. (2013) Postreplicative mismatch repair. *Cold Spring Harb Perspect Biol*, **5**, a012633.
9.   Kunkel, T.A. and Erie, D.A. (2005) DNA mismatch repair. *Annu Rev Biochem*, **74**, 681-710.
10.   Lenhart, J.S., Pillon, M.C., Guarné, A., Biteen, J.S. and Simmons, L.A. (2016) Mismatch repair in Gram-positive bacteria. *Res Microbiol*, **167**, 4-12.
11.   Marinus, M.G. (2012) DNA Mismatch Repair. *EcoSal Plus*, **5**.
12.   Ishino, S., Nishi, Y., Oda, S., Uemori, T., Sagara, T., Takatsu, N., Yamagami, T., Shirai, T. and Ishino, Y. (2016) Identification of a mismatch-specific endonuclease in hyperthermophilic Archaea. *Nucleic Acids Res*, **44**, 2977-2986.
13.   Ren, B., Kuhn, J., Meslet-Cladiere, L., Briffotaux, J., Norais, C., Lavigne, R., Flament, D., Ladenstein, R. and Myllykallio, H. (2009) Structure and function of a novel endonuclease acting on branched DNA substrates. *EMBO J*, **28**, 2479-2489.
14.   Nakae, S., Hijikata, A., Tsuji, T., Yonezawa, K., Kouyama, K.I., Mayanagi, K., Ishino, S., Ishino, Y. and Shirai, T. (2016) Structure of the EndoMS-DNA Complex as Mismatch Restriction Endonuclease. *Structure*, **24**, 1960-1971.
15.   Putnam, C.D. (2021) Strand discrimination in DNA mismatch repair. *DNA Repair (Amst)*, **105**, 103161.
16.   Zhang, L., Jiang, D., Wu, M., Yang, Z. and Oger, P.M. (2020) New Insights Into DNA Repair Revealed by NucS Endonucleases From Hyperthermophilic Archaea. *Front Microbiol*, **11**, 1263.
17.   Ishino, S., Skouloubris, S., Kudo, H., l'Hermitte-Stead, C., Es-Sadik, A., Lambry, J.C., Ishino, Y. and Myllykallio, H. (2018) Activation of the mismatch-specific endonuclease EndoMS/NucS by the replication clamp is required for high fidelity DNA replication. *Nucleic Acids Res*, **46**, 6206-6217.
18.   Takemoto, N., Numata, I., Su'etsugu, M. and Miyoshi-Akiyama, T. (2018) Bacterial EndoMS/NucS acts as a clamp-mediated mismatch endonuclease to

prevent asymmetric accumulation of replication errors. *Nucleic Acids Res*, **46**, 6152-6165.

19. Castaneda-Garcia, A., Prieto, A.I., Rodriguez-Beltran, J., Alonso, N., Cantillon, D., Costas, C., Perez-Lago, L., Zegeye, E.D., Herranz, M., Plocinski, P. *et al.* (2017) A non-canonical mismatch repair pathway in prokaryotes. *Nat Commun*, **8**, 14246.

20. Castaneda-Garcia, A., Martin-Blecua, I., Cebrian-Sastre, E., Chiner-Oms, A., Torres-Puente, M., Comas, I. and Blazquez, J. (2020) Specificity and mutagenesis bias of the mycobacterial alternative mismatch repair analyzed by mutation accumulation studies. *Sci Adv*, **6**, eaay4453.

21. Murphy, K.C. (2021) Oligo-Mediated Recombineering and its Use for Making SNPs, Knockouts, Insertions, and Fusions in Mycobacterium tuberculosis. *Methods Mol Biol*, **2314**, 301-321.

22. van Kessel, J.C. and Hatfull, G.F. (2008) Efficient point mutagenesis in mycobacteria using single-stranded DNA recombineering: characterization of antimycobacterial drug targets. *Mol Microbiol*, **67**, 1094-1107.

23. Sandler, S.J., Satin, L.H., Samra, H.S. and Clark, A.J. (1996) recA-like genes from three archaean species with putative protein products similar to Rad51 and Dmc1 proteins of the yeast Saccharomyces cerevisiae. *Nucleic Acids Res*, **24**, 2125-2132.

24. Seitz, E.M., Brockman, J.P., Sandler, S.J., Clark, A.J. and Kowalczykowski, S.C. (1998) RadA protein is an archaeal RecA protein homolog that catalyzes DNA strand exchange. *Genes Dev*, **12**, 1248-1253.

25. Cooper, D.L. and Lovett, S.T. (2016) Recombinational branch migration by the RadA/Sms paralog of RecA in Escherichia coli. *Elife*, **5**.

26. Marie, L., Rapisarda, C., Morales, V., Bergé, M., Perry, T., Soulet, A.L., Gruget, C., Remaut, H., Fronzes, R. and Polard, P. (2017) Bacterial RadA is a DnaB-type helicase interacting with RecA to promote bidirectional D-loop extension. *Nat Commun*, **8**, 15638.

27. Cooper, D.L., Boyle, D.C. and Lovett, S.T. (2015) Genetic analysis of Escherichia coli RadA: functional motifs and genetic interactions. *Mol Microbiol*, **95**, 769-779.

28. Torres, R., Serrano, E. and Alonso, J.C. (2019) Bacillus subtilis RecA interacts with and loads RadA/Sms to unwind recombination intermediates during natural chromosomal transformation. *Nucleic Acids Res*, **47**, 9198-9215.

29. Murphy, K.C., Nelson, S.J., Nambi, S., Papavinasasundaram, K., Baer, C.E. and Sassetti, C.M. (2018) ORBIT: a New Paradigm for Genetic Engineering of Mycobacterial Chromosomes. *mBio*, **9**.

30. Stephanou, N.C., Gao, F., Bongiorno, P., Ehrt, S., Schnappinger, D., Shuman, S. and Glickman, M.S. (2007) Mycobacterial nonhomologous end joining mediates mutagenic repair of chromosomal double-strand DNA breaks. *J Bacteriol*, **189**, 5237-5246.

31. Murphy, K.C., Papavinasasundaram, K. and Sassetti, C.M. (2015) Mycobacterial recombineering. *Methods Mol Biol*, **1285**, 177-199.

32. Snapper, S.B., Melton, R.E., Mustafa, S., Kieser, T. and Jr, W.R.J. (1990) Isolation and characterization of efficient plasmid transformation mutants of Mycobacterium smegmatis. *Molecular Microbiology*, **4**, 1911-1919.

Figure Legends

Figure 1

Mutagenic phenotype of *M. smegmatis* Δ*nucS* strain. **(A)** Wild and a Δ*nucS* derivative of *M. smegmatis* were grown in 7H9-OADC-tween to saturation (O.D. of 1.5) and 200 ul of cultures were spread on 7H10-OADC-tween plates containing 150 ug/ml rifampicin.

(B) *M. smegmatis* wild type and ΔnucS strains containing no plasmids, pIR540 (expressing full length NucS), or pIR541 (expressing NucS containing a 5 amino acid truncation from the C-terminus – NucSDC5) were grown to saturation and plated on 7H10-OADC-tween rifampicin plates as described in above. The mutation frequency is the titer of rifampicin resistant mutants divided by the total number of cells plated. Unpaired t-test statistical analysis was performed,

Figure 2

Oligo-mediated recombineering of the *rpsL* locus of *M. smegmatis*. (Left) Sequences of the oligo-generated mismatch of rpsL region resulting in a mismatch (A-C) predicted not to be recognized by mycobacterial NucS, and a mismatch (G-G) expected to be acted upon by NucS, as predicted from in vitro results of *C.. glutamificans*. (Right) Following replication of mismatched-containing DNA, both wild type and mutant versions of *rpsL* are generated. Both mutants are known to cause streptomycin resistance in mycobacteria.

Figure 3

Mismatch recognition specificity of *M. smegmatis* NucS in vivo. Oligo-mediated

recombineering experiments were carried out in WT **(A)** or ∆nucS **(B)** cells

containing a defective hygromycin gene, where codon-110 for glycine (GGA) had

been altered to a stop codon (TGA). The oligos tested generated 7 different

mismatches for restoration of the *hyg* gene. Repair and non-repair of the

mismatches is evident by the appearance of low and high frequencies of

hygromycin-resistant colonies, respectively. Electroporations were carried out in

triplicate (standard deviations shown)

Figure 4

NucS does not repair indels. Oligo-mediated recombineering was performed in

strains where the *leuB* gene had been deactivated by the deletion of 1 or 2 base

pairs, and by the insertion of 1 base pair in the codon that encodes for the active site

residue Arg-216. An oligo containing the wild type *leuB* sequence was

electroporated into these cells which expresses the RecT annealase. LeuB+

recombinants were measured by plating the cells on 7H10-OADC-tween plates

without leucine supplementation; total cells were tittered on 7H10-OADC-tween

plates with 40 ug/ml leucine. For comparison, a *leuB* strain where the active site

codon was disable could be efficiently restored with an oligo that creates a

repairable G-G mismatch in ∆nucS host, but not wild type cells.

Figure 5

Mutations suspected of being critical for exonucleolytic activity in NucS are defective of MMR. (A) The structure of the NucS protein from *Thermococcus kodakarensis* highlighting residues thought to be critical for exonucleolytic activity. (B) The corresponding residues in *M. smegmatis* NucS were changed to alanine by recombineering and tested for levels of spontaneous mutation to rifampicin. The *T. kodakarensis* NucS structure reprinted from Nakae et. al. (14) with permission.

Figure 6

Description of the double mismatch-generating oligos. (A) The dsDNA sequence of the defective hygromycin-resistant gene (*hyg*). The GGA-110 codon has been altered to stop codon TAG (highlighted in green). (B). After annealing of the oligo (in blue type) to the target site in the *M. smegmatis* chromosome, the oligo creates two mismatches: a C-T mismatch that (left unrepaired) leads to high levels of Hyg^R colonies and a T-G mismatch that is acted upon by NucS in vivo. (C) NucS is predicted to make a dsDNA cut at the mismatch. If the ends of the breaks are acted upon by a 5'-3' exonuclease, Hyg^R recombinants will be lost if the processing extends to the C-T mismatch or beyond and removes the "C". (D) If the processing by a 5'-3' exonuclease is limited and does not extend to the C-T mismatch, HygR transformants will be generated.

Figure 7

5'-3' exonuclease processing of NucS-induced breaks. **(A)** The target region of the defective hyg gene is shown; the C-T mismatch leading to Hyg^R is shown in green. The

top strand is the sequence of the recombineering oligo (replicating lagging strand) and positions where nine different oligos produce a repairable G-G mismatch are shown in red. **(B).** Oligo recombineering was caried out in WT *M. smegmatis* carrying the defective *hyg* gene. Each of nine oligos used in this assay generates a C-T mismatch at a fixed position (*i.e.,* the stop codon) and a G-G mismatch at variable positions as described in panel A. The names of the oligos on the X- axis defines the number of base pairs between the two mismatches, and if the G-G mismatch is upstream (-) or downstream (+) of the C-T mismatch. The frequency of Hyg^R colonies and total cell number was determined by plating on 7H10-OADC-tween plate with and without hygromycin, respectively. **(C)** Diagram of a dsDNA cut made by oligo (-5) *i.e.,* 5 base pairs upstream of the C-T mismatch that conveys Hyg^R. **(D)** Diagram of a dsDNA cut made by oligo (+4) *i.e.*, 4 base pairs downstream of the C-T mismatch that conveys Hyg^R.

Figure 8

Testing of functions suspected in *M. smegmatis* NucS-promoted MMR repair

Oligos generating C-T mismatches that promote Hyg^R and generated no other mismatch (Hyg-repair-0), or a G-G mismatch 32 bp away from the C-T mismatch (Hyg repair-32), were electroporated into *M. smegmatis* strains containing deletion mutants for a variety of genes suspected to be involved with nucS-promoted MMR (see legend). All strains contained the RecT-producing plasmid pKM461. Disruption of genes involved in NucS-promoted MMR are expected to generate low frequencies of HygR with oligos containing the G-G mismatch (G-G), compared to high frequencies of HygR expected with oligos not containing the G-G mismatch (no mismatch).

Table 1

Frequencies* of oligonucleotide-mediated

Streptomycin resistance in *M. smegmatis*

Strain	K43R oligo (A/C mismatch)	K43Q oligo (G/G mismatch)
Wild type	1.59	0.0097
ΔnucS	1.05	1.4

*   Titer of streptomycin resistant colonies/total colonies

Table 2
Strains used and constructed for this study

Strain Name	Genotype	Description	Markers	Source
MC^2-155	wild type	Efficient plasmid transformation (ept)	none	(32)
MGM199	ΔrecA	Deletion of *recA*		(30)
MGM153	ΔKu, ΔligD	Deletion of *Ku* and *ligD*	none	(30)
KM272	Wild type Defective Hyg^R gene	pKM461 (RecT); pKM614 (defective hyg gene) inserted into the MSMEG_1397-*rpsL* intergenic region by ORBIT	Kan^R Zeo^R	This study
KM274	ΔrecA Defective Hyg^R gene	MGM199/KM461 (RecT); pKM614 (defective hyg gene) inserted into the MSMEG-rpsL intergenic region by ORBIT	Kan^R Zeo^R	This study
KM295	ΔKu, ΔligD Defective Hyg^R gene	MGM153/KM461 (RecT); pKM614 (defective hyg gene) inserted into the MSMEG-rpsL intergenic region by ORBIT	Kan^R Zeo^R	This study
KM165	ΔnucS::pKM464	MC^2-155 deleted of MSMEG_4923 by Orbit	Hyg^R	This study
KM183	ΔnucS::attP	KM165 - cured of pKM464.	none	This study
VF20	Wild type Defective Hyg^R gene	MC^2-155/pKM461 (RecT); pIR542 (defective *hyg* gene) inserted into phage Giles site.	Kan^R Zeo^R	This study
VF22	ΔnucS:attP Defective Hyg^R gene	KM196/pKM461 (RecT); pIR542 (defective *hyg* gene) inserted into phage Giles site.	Kan^R Zeo^R	This study
KM209	ΔfenA::pKM464	MC^2-155 deleted of MSMEG_3883 by ORBIT	Hyg^R	This study
KM210	ΔfenA::attP	KM209 - cured of pKM464.	none	This study
KM211	ΔfenA:pKM611 Defective Hyg^R gene	KM210/pKM461 (RecT); pKM611 (defective *hyg* gene) inserted Δ*fenA::attP* site	Kan^R Zeo^R	This study

KM215	ΔradA::pKM464	MC²-155 deleted of MSMEG_6079 by ORBIT	Kan^R Zeo^R	This study
KM284	ΔradA::attP	KM215 - cured of pKM464.	none	
KM287	ΔradA:pKM611 Defective Hyg^R gene	KM284/pKM461 (RecT); pKM611 (defective *hyg* gene) inserted Δ*radA::attP* site	Hyg^R	This study
KM287	ΔrecA, ΔradA:pKM611 Defective Hyg^R gene	MGM199/pKM461 (RecT); pKM611 (defective *hyg* gene) inserted Δ*radA* gene by ORBIT	none	This study
KM229	MSMEG_2379 (leuB frameshift)	Two bp deletion in the arginine-101 codon; pKM461; pKM585	Kan^R Strep^R	This study
KM230	MSMEG_2379 (leuB frameshift)	One bp deletion in the arginine-101 codon; pKM461: pKM585	Kan^R Strep^R	This study
KM231	MSMEG_2379 (leuB frameshift)	One bp insertion in the arginine-1; pKM46101 codon; pKM585	Kan^R Strep^R	This study
KM234	MSMEG_2379 (leuB frameshift)	One bp insertion in the arginine-1; cured of ORBIT plasmid pKM461 and pKM585.	none	This study
KM262	NucS-D138A	Active site mutant of *M. smegmatis* NucS	none	This study
KM263	NucS-E152A	Active site mutant of *M. smegmatis* NucS	none	This study
KM264	NucS-K154A	Active site mutant of *M. smegmatis* NucS	none	This study
Mtb-25	MtbΔnucS::pKM488	pKM461/H37Rv deleted of Rv1321 by ORBIT; cured of pKM461	Hyg^R	This study
Mtb-26	Mtb ΔnucS::attP	Mtb-25 - cured of pKM488.	none	This study

A.

Wild type *M. smegmatis*   *M. smegmatis (ΔnucS)*

B.

Figure 1

Figure 2

A.

B.

Figure 3

Figure 4

A

E179
D165A
K181 C

Y41
N76
W77

Thermococcus kodakarensis NucS

B.

Figure 5

A.   Target dsDNA (defective *hyg* gene):

GACGCCCCGGTGCCGGTGCCCCGCCTCCTCGGCCGCGGCGAGCTGCGGCCCGGCACC<mark>TAG</mark>GCCTGGCCGTGGCCCTACCTGGTG
CTGCGGGGCCACGGCCACGGGGCGGAGGAGCCGGCGCCGCTCGACGCCGGGCCGTGGATCCGGACCGGCACCGGGATGGACCAC

B.   Oligo (top strand) annealed to lagging strand template:

Mismatch at codon 99   Mismatch at codon 110
Repairable Mismatch   Not recognized by NucS
Cut by NucS   Restores Hyg[R]

C.   NucS-induced dsDNA cut followed by **extensive** 5'-3' exonucleolytic processing leads to **loss** Hyg[R] transformants.

GACGCCCCGGTGCCGGTGCCCCGC–3'   5'–TGGCCCTACCTGGTG
CTG–5'   3'–GCCGGCGCCGCTCGACGCCGGGCCGTGGA<mark>T</mark>CCGGACCGGCACCGGGATGGACCAC

D.   NucS-induced dsDNA cut followed by **limited** 5'-3' exonucleolytic processing leads to **preservation** of Hyg[R] transformants.

GACGCCCCGGTGCCGGTGCCCCGC–3'   5'CTGCGGCCCGGCACCT<mark>C</mark>GGCCTGGCCGTGGCCCTACCTGGTG
CTGCGGGGC–5'   3'–GCCGGCGCCGCTCGACGCCGGGCCGTGGA<mark>T</mark>CCGGACCGGCACCGGGATGGACCAC

Figure 6

A.
ACGCCCCGGTGCCGGTGCCCCGCCTTCTCGGCCGCGG**G**GAG**G**TG**G**GG**G**C**G**GG**G**ACCT**C**GGC**G**TGG**G**CGTGGCC**G**TACCTGGTG
TGCGGGGCCACGGCCACGGGGCGGAGGAGCCGGCGCC**G**CTC**G**AC**G**CC**G**GG**G**CC**G**TGGA**T**CCG**G**ACC**G**GCACCGG**G**ATGGACCAC

B. Exo processing of NucS-induced breaks

C.

-5
(G-G mismatch 5 bp
upstream of C-T mismatch)

….GGGCGGG**G**ACCT**C**GGCG…
……CCCGCCC**G**TGGA**T**CCGC…

….GGGCG-3'   5'-GG**G**ACCT**C**GGCG…
CCCGCCC**G**TG-5'   3'-GA**T**CCGC…

D.

+4
(G-G mismatch 4 bp
downstream of C-T mismatch)

…. GACCT**C**GGC**G**TGGGGGTG…
…… CTGGA**T**CCG**G**ACCCCCAC

… GACCT**C**G-3'   5'-GC**G**TGGGGGTG…
…CTGGA**T**CCG**G**AC-5'   3'- CCCCCAC

Figure 7

Figure 8