Aging impairs skeletal muscle regeneration by promoting fibro/fatty degeneration and inhibiting inflammation resolution via fibro-adipogenic progenitors

Francisco Garcia-Carrizo[1,2], Sabrina Gohlke[1], Georgia Lenihan-Geels[1], Anne-Marie Jank[1], Marina Leer[1,2], George A. Soultoukis[1], Masoome Oveisi[1,2], Catrin Herpich[3,4,5], Claudia A. Garrido[6,7], Georgios Kotsaris[8], Sophie Pöhle-Kronawitter[1,8], Arnold Tsamo-Tetou[1], Antonia Graja[1], Mario Ost[9], Laura Villacorta[10], Raphael S. Knecht[6], Susanne Klaus[9], Annette Schürmann[2,12], Sigmar Stricker[8], Katharina Schmidt-Bleek[6,11], Amaia Cipitria[6,7,13,14], Georg N. Duda[6,11], Vladimir Benes[10], Ursula Müller-Werdan[4,15], Kristina Norman[3,4,5], Tim J. Schulz[1,2,5,*]

1. Department of Adipocyte Development and Nutrition, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

2. German Center for Diabetes Research (DZD), München-Neuherberg, Germany

3. Department of Nutrition and Gerontology, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

4. Department of Geriatrics and Medical Gerontology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität Zu Berlin, Berlin, Germany

5. Institute of Nutritional Science, University of Potsdam, Potsdam-Rehbrücke, 14558 Nuthetal, Germany

6. Julius Wolff Institute, Berlin Institute of Health at Charité - Universitätsmedizin Berlin, Berlin, Germany

7. Department of Biomaterials, Max Planck Institute for Colloids and Interfaces, Potsdam, Germany

8. Institute of Chemistry and Biochemistry, Musculoskeletal Development and Regeneration Group, Freie Universität Berlin, Thielallee 63, 14195 Berlin, Germany

9. Department of Physiology of Energy Metabolism, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

10. Genomics Core Facility, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany

11. Berlin Institute of Health Centre for Regenerative Therapies (BCRT), Berlin Institute of Health at Charité - Universitätsmedizin Berlin, Berlin, Germany

12. Department of Experimental Diabetology, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

13. Current address: Group of Bioengineering in Regeneration and Cancer, Biogipuzkoa Health Research Institute,

San Sebastián, Spain

14. Current address: IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

15. Protestant Geriatric Center Berlin (EGZB), Berlin, Germany

*Corresponding author and Lead Contact

Tim J. Schulz, PhD

German Institute of Human Nutrition

114-116, Arthur-Scheunert Allee

Potsdam-Nuthetal, D-14558, Germany

Phone: +49-33200-88-2110;

Email: tim.schulz@dife.de

Summary

Skeletal muscle regeneration depends on the function of fibro/adipogenic progenitors (FAPs). Here we show that aging

impairs myogenic stem cells by disrupting the integration of extracellular matrix and immunomodulatory functions

within the stem cell niche, thereby promoting fibro/fatty degeneration. We identify the FAP-secreted protein Periostin

as a niche factor that is decreased in aged muscle and in circulation of aged humans with low-exercise lifestyle.

Periostin controls FAP-expansion after injury and its depletion fate-regulates FAPs towards adipogenesis. This leads

to delayed pro- to anti-inflammatory macrophage transition during regeneration. Transplantation of young FAPs with

high Periostin secretion, but not Periostin-deficient FAPs, into aged muscle restores inflammation resolution and

successful regeneration. Mechanistically, Periostin activates Focal adhesion kinase- and AKT-signaling in

macrophages via integrins to promote an anti-inflammatory profile, which synchronizes matrix-derived

mechanosensory signaling and immunomodulation. These results uncover a novel role of FAP-based regulation that

orchestrates successful muscle regeneration and prevents fibro/fatty degeneration.

Keywords: Aging, muscle stem cell niche, ECM-remodeling, FAPs-Periostin, satellite cells, macrophages

Introduction

Aging is characterized by a gradual loss of muscle mass and strength that leads to sarcopenia as part of the frailty syndrome, a pathology that constitutes a serious health problem associated with adverse outcomes in elderly individuals[1,2]. This age-related decline in muscle function is not only associated with a reduction in muscle stem cell (MuSC) numbers and a reduced regenerative potential but also the accumulation of fibrotic tissue and ectopic adipocytes within the skeletal muscle, which is referred to as fibro/fatty infiltration or fatty degeneration[3-7]. MuSCs also interact with their local microenvironment and a key component of the muscle stem cell niche is the extracellular matrix (ECM) which experiences extensive remodeling during regeneration and throughout life[8-12]. Fibro/adipogenic progenitor cells (FAPs) are considered the main ECM-source, they regulate its remodeling and are indispensable for long-term muscle maintenance and MuSCs' functionality during regeneration[13-15]. However, under pathological conditions, such as muscular dystrophies, obesity, and type 2 diabetes mellitus (T2DM), subsets of FAPs are thought to generate fibrotic scar tissue and differentiate into adipocytes that progressively replace functional muscle tissue, resulting in fibro/fatty degeneration and the age-related metabolic dysfunction of muscle tissue[6,16-20].

Besides their role in ECM remodeling, FAPs secrete trophic factors that control MuSC activation through different mechanisms[14,21-23] and aged FAPs display an aberrant secretory profile of WISP1, BMP3b or SMOC2, among others, that reduce MuSCs' regeneration capacity[10,11,24]. The contribution of aging to FAPs' dysregulation and fibro/fatty infiltration in skeletal muscle remains less well understood. In addition to the FAP-MuSCs niche crosstalk, FAPs interact with different immune cells[25-28]. For instance, pro-inflammatory macrophages regulate the number of FAPs via TNF-mediated apoptosis[28]. A failure to resolve the initial pro-inflammatory macrophage response, however, reduces myogenic regeneration and promotes chronic inflammation in aged muscles[29-32].

Periostin (*Postn*) is a matricellular protein that interacts with ECM components to drive fibrillogenesis and matrix scaffold formation. It also displays signaling activity by binding integrin receptors, mainly αvβ3 and αvβ5 heterodimers, to promote cell adhesion, migration and proliferation in different tissues and in the context of regeneration[33-35]. Expression of *Postn* is transiently induced in FAPs during muscle regeneration and initial data suggest a role in regeneration by coupling myogenesis and angiogenesis[36-38]. Omics-based analyses also list *Postn* among the potentially dysregulated genes in aged muscles[11,22]. In addition to its tissue/stem cell-specific function, Periostin is discussed as a circulating biomarker of type-2 inflammation in asthma, breast cancer, cardiovascular disease and liver steatosis[39-42], as well as a potential biomarker of aging in serum and across several murine tissues[22,43].

We here investigate the role of age-related changes of the muscle stem cell nice, which result in increased fibro/fatty

degeneration of skeletal muscle at the expense of healthy tissue regeneration. We identify a mechanism that

synchronizes several key ECM-functions, including mechano-signaling and immunomodulation, between the different

cell populations of the muscle microenvironment to orchestrate successful muscle regeneration. This niche factor-

mediated integration is impaired during aging and we identify the matricellular protein Periostin as an important

contributor to these processes. Loss of Periostin phenocopies muscle aging, resulting in delayed regeneration, ectopic

adipocyte accumulation and fibrosis, which contribute to delayed inflammation resolution during injury repair.

Reconstitution of Periostin through FAP-transplantation recovers young-like regeneration in aged muscle.

Mechanistically, Periostin targets mechanosensory pathways in niche-resident macrophages, thereby promoting

resolution of the inflammatory microenvironment during regeneration. Periostin may serve as sarcopenia biomarker

as circulating levels positively correlate with muscle mass and function at different ages, but are reduced with age in

humans.

Results

Loss of Periostin in aged muscle disrupts extracellular matrix function and the stem cell niche

To examine the contribution of aging to myogenic niche disruption, we compared young, 2-months-old male and

female mice to sex-matched animals aged 15-18 months. This age was chosen to reduce the likelihood of other,

potentially confounding, pathologies that may occur in even older mice. At 15 months, muscles display a gene

expression profile and morphology consistent with fibro/fatty degeneration, suggesting such changes are relatively

early hallmarks of muscle aging (Figures 1A, S1A-S1C). Fluorescence-activated cell sorting (FACS) showed increased

numbers of muscle-resident FAPs with age while MuSCs decreased in male and female mice (Figures 1B, S1D-S1H).

Primary FAPs isolated from muscle of 15-months-old mice displayed spontaneous lipid droplet accumulation, while

aged MuSCs showed reduced myogenic potential compared to respective young-muscle cell cultures (Figures 1C and

1D). Given the marked increase in adipogenic potential in aged FAPs, we decided to further examine this cell

population. Microarray analysis of freshly sorted FAPs showed a marked reduction in many genes and pathways linked

to the ECM (Table S1, Figure S1I). The top down-regulated gene, *Postn*, which encodes for Periostin, is a secreted

matricellular protein[35] and could thus act as paracrine niche factor (Figure 1E). Aging muscles displayed reduced

Periostin expression on protein and mRNA levels in male and female mice, and cultured young FAPs readily released

Periostin into the supernatant, while aged FAPs only secreted minor amounts (Figures 1F-1H, S1J and S1K).

We next tested whether muscle injury would elicit a similar phenotype and applied an established model of tibialis anterior (TA) muscle injury by intramuscular injection of glycerol, a protocol that induces ectopic adipogenesis and fibrosis at the injury site[44]. Injury-induced expansion of FAPs was more pronounced in aged TA compared to young post-injury muscle in male and female mice. Unexpectedly, MuSC expansion was also induced in aged muscle of female mice and to some extent in male mice Figures 1I, 1J and S1L-S1N). Clearance of FAPs is critical to prevent fibro/fatty infiltration[28], and accordingly we detected increased adipogenesis and fibrosis areas in injured TA of aged compared to young male mice which was accompanied by an age-related reduction in myofiber cross-sectional areas (CSA) at the injury site and reduced expression of myogenesis markers (Figures 1K-1M and S1O-S1Q). To further explore the involvement of Periostin in muscle healing, we evaluated its expression profile on multiple days post-injury (dpi). In line with other publications[36-38,45], we observed an increase of Periostin in the interstitial space between regenerating fibers and corresponding dynamic mRNA and protein expression patterns that peaked between 5- and 8-dpi in muscle of young mice, all of which was markedly attenuated in aged muscle (Figures 1N-1P). We verified disrupted FAP-functions in regenerating aged muscles by applying single cell-RNA sequencing (scRNA-seq) of FAPs isolated from 4-dpi muscle. Uniform manifold approximation and projection (UMAP) visualization and differentially expressed gene (DEG) analysis revealed partial transcriptional separation of FAPs of the two age groups (Figure 1Q; Table S2). This analysis confirmed *Postn* as one of the most significantly down-regulated genes and the enrichment of ECM-related pathways when assessing all significantly downregulated genes (Figures 1R-1T). Up-regulated pathways and a subsequent analysis of predicted secretome genes showed a potential impact of FAP-aging on the modulation of inflammatory responses (Figures S1R and S1S). These findings taken together indicate that aging negatively influences niche functions of FAPs which contribute to fibro/fatty degeneration of aged muscles.

Genetic inactivation of *Postn* recapitulates an aging-like phenotype and promotes fibro/fatty degeneration

To determine whether loss of Periostin could indeed contribute to a decline of muscle regeneration as observed in aged mice, we obtained a mouse model with genetic inactivation of the *Postn* gene by insertion of a LacZ cassette, leading to loss of the *Postn* gene product and expression of a β-galactosidase reporter under control of the native *Postn* promoter (Figures S2A-S2D). Periostin protein was completely absent in muscles of *Postn*[-/-] mice whereas the reporter was expressed in the interstitial space between myofibers, consistent with the localization of FAPs (Figure S2B). No differences in myofiber CSAs or frequencies of FAPs and MuSCs were observed in 2-3 months-old muscles of male or female *Postn*[-/-] mice without injury. However, we noticed that dissected muscles displayed a significant reduction

in mechanical stiffness (Figures S2E to S2G). Similar to aged muscle, isolated *Postn*$^{-/-}$-FAPs displayed significantly

enhanced spontaneous adipogenesis in culture whereas cultured MuSCs had significantly lower myogenic potential

(Figures 2A and 2B). Applying the glycerol-injection regeneration model to *Postn*$^{-/-}$ mice and littermate wildtype

controls using both male and female offspring showed a phenotype closely resembling the phenotype seen in aged

muscle: We consistently observed increased expansion of FAPs and MuSCs in *Postn*$^{-/-}$ mice (Figure 2C, 2D, S2H and

S2I). Importantly, both sexes showed impaired regeneration in *Postn*$^{-/-}$ mice with significant accumulation of

adipocytes and fibrotic structures and reduced CSA of newly regenerated myofibers at 14-dpi, which was also reflected

by increased expression of adipocyte and fibrosis genes and reduced expression of myogenesis markers (Figures 2E-

2G and S2J-S2Q). In summary, genetic ablation of *Postn* recapitulates an aging-like disruption of regeneration in male

and female mice at the level of FAPs and MuSCs, resulting in fibro/fatty degeneration.

Periostin rescues stem cell function and muscle regeneration in aged mice

To better understand the function of Periostin as ECM-related niche factor, we tested the impact of a mouse

recombinant Periostin (mrPOSTN) on FACS-purified FAPs and MuSCs of aged and *Postn*-deficient mice. Both cell

types, aged and *Postn*$^{-/-}$-FAPs, displayed significantly increased proliferation when compared to the respective control

FAPs, whereas MuSCs displayed reduced proliferation rates in comparison to respective controls (Figures 3A-3D,

S3A and S3B). After 48h of culture in the presence of mrPOSTN the respective FAP- and MuSC-proliferation

phenotypes were fully reversed in *Postn*$^{-/-}$ and aged cells. Considering that FAPs are likely a main source of muscle

Periostin, these data suggest that FAPs auto-inhibit their own expansion while stimulating cell cycle in MuSCs through

paracrine Periostin (Figures 1H, 3A-3D). To determine whether FAP-derived Periostin influences the function of

MuSCs, we used an in vitro co-culture system. Consistent with our initial observation, lack of Periostin in MuSCs

significantly reduced fusion capacity and resulted in shorter myofibers during differentiation (Figures 3E-3G). Co-

cultures of wildtype, Periostin-secreting FAPs, but not *Postn*$^{-/-}$-FAPs, with *Postn*-deficient MuSCs resulted in a full

recovery of the impaired myogenesis phenotype (Figures 3E-3G). As cultured MuSCs themselves secrete very little

Periostin in comparison to FAPs, this interaction appears to be a key element in maintaining the in vitro myogenic

potential of the myogenic stem cell pool (Figure S3C).

To determine whether Periostin reconstitution has the capacity to enhance the regeneration process in aged skeletal

muscle, we sorted FAPs from a mouse model carrying a red fluorescent protein (RFP) expression allele in addition to

the *Postn*-knockout allele (Figures S3D and S3E) [46]. RFP$^+$ FAPs either with (FAP^{RFP}-WT) or without (FAP^{RFP}-KO)

Postn expression were transplanted into TA muscle of 15-months-old recipient mice one day after glycerol-injury.

Transplanted FAPs remained detectable based on RFP expression until at least 8-dpi (Figures 3H, S3D-S3F).

Immunofluorescence analysis at 8-dpi showed that FAP^RFP-WT transplants significantly reduced the number of

embryonic myosin heavy chain (eMYHC) positive cells and increased CSAs in regenerating fibers compared to aged

recipients transplanted with FAP^RFP-KO (Figures 3I-3K). Consistently, recipients transplanted with FAP^RFP-WT

maintained significantly increased CSAs at 14-dpi when compared to FAP^REF-KO transplants (Figures 3L and 3M).

Finally, we compared the CSAs average and fiber size range results at 14-dpi to the original experiments without FAP-

transplantation (i.e. CSA-means and fiber size range of data reported in Figure 1M). Aged mice receiving FAP^RFP-KO

transplants displayed significantly lower average CSA compared to young control mice and were at a similar level

compared to aged TA without FAP transplant, whereas Periostin-producing FAP^RFP-WT transplants significantly

increased regenerating myofibers' CSA compared to aged mice without transplanted FAPs or with FAP^RFP-KO

transplants (Figure 3N and S3G). Thus, reinstating Periostin as a niche factor ameliorates the age-related defect of

myogenic regeneration, indicating that Periostin is a critical ECM-protein to orchestrate muscle regeneration.

187 Periostin mediates immunomodulatory functions of the myogenic stem cell niche

Since regeneration is orchestrated by the interplay of multiple non-myofiber cell populations[47], we performed droplet-

based scRNA-seq of FACS-purified non-myofiber cells from 2.5- and 15-month-old female mice at 4-dpi, around the

time when FAPs frequency peaks during muscle injury repair (Figures S4A and S4B). UMAP visualization,

unsupervised Louvain-clustering and subsequent cell type annotation based on gene scoring of single cell

transcriptomes resulted in the identification of distinct clusters representing the main nine, non-myofiber cell

populations involved in muscle regeneration (Figures 4A, 4B, S4C and S4D; Tables S3 and S4) [48]. Analysis of the cell-

type proportions between age groups showed increased relative abundances of myeloid subsets, i.e. neutrophil (NP)

and macrophage (MC) subsets in aged muscle, while lymphoid subpopulations, i.e. T-lymphocytes (TC) and natural

killer cells (NK), showed lower relative frequencies (Figure 4C; Table S5). Considerable differences in gene expression

patterns were observed among multiple cell types and when comparing the 2.5- and 15-months-old samples. For

instance, analysis of enriched term annotations of downregulated DEGs from the FAP subset confirmed that aging

negatively influences ECM regulation (Figure S4E, Table S6). As before, *Postn* was mostly expressed in FAPs and its

expression was significantly reduced in aged FAPs (Figures 4D, 4E, and S4D; Table S4). To examine cell-cell

interactions (CCI) between Periostin-secreting FAPs and other niche cells, we took advantage of a previously published

murine atlas of age-related changes in intercellular communications, which uses an algorithm to infer CCI from

published gene expression databases[49]. When selecting Periostin as a FAP-emitted ligand, high CCI-scores were

identified for a potential autocrine FAP-FAP interaction and for paracrine FAP-endothelial cell (EC), FAP-MuSC and

FAP-MC interactions mediated through different Integrin receptors (*Itga5*, *Itgβ1*, and heterodimers *Itgαv/β5*, *Itgαv/β3*)

and the Epidermal growth factor receptor (*Egfr*; Figure 4F). Using aging as a regulatory factor of these interactions,

statistical significance levels for each interaction showed a significant down-regulation of Periostin-based FAP-FAP,

FAP-EC, FAP-MuSC and FAP-MC interactions, thus verifying our own experimental data on the auto- and paracrine

roles of Periostin (Figure 4G). The age-dependent down-regulation of the interaction between FAPs and MCs through

the predicted CCI between Periostin and the Itgαv/β5 heterodimer raised our interest as this was consistent with the

alterations in immune subsets in our scRNA-seq data and offered the possibility of an immunomodulatory niche effect

through the FAP-based release of Periostin (Figure 4G). We validated subset-specific expression patterns of all

potential Periostin receptors in our scRNA-seq data and found that the proposed Itgαv/β5 heterodimer showed an

expression pattern that would most closely reflect the effects of Periostin on the cell types that we also detected in our

experimental analyses, i.e. FAPs, MuSCs, and MCs (Figure 4H).

Loss of Periostin impairs FAP-based immunomodulation by attenuating anti-inflammatory macrophages

Corroborating the FAP-MC interaction through Periostin, FACS-analyses of MCs populations confirmed the increase

of MCs upon muscle injury in aged and also female *Postn*[-/-] mice compared to the respective control animals, while

only trends or no differences were observed in male *Postn*[-/-] mice (Figures S5A-S5E). Next, we examined MC subsets

representing functional changes from a pro-inflammatory (M1-like) profile to an anti-inflammatory (M2-like)

phenotype during the course of muscle healing[25,50]. As expected, injury led to a rapid increase of $CD45^+CD11b^+F4/80^+$

MCs, which over time shifted from a predominantly pro-inflammatory $Ly6C^+$ population, which peaked at 1- and 3-

dpi, to an anti-inflammatory $Ly6C^-$ population which peaked at 4- and 8-dpi (Figure 5A). Importantly, aging as well

as genetic ablation of *Postn* led to an attenuation of this shift in male and female mice, both of which resulted in

significantly elevated levels of pro-inflammatory MCs at earlier time points and significantly lower levels of anti-

inflammatory MCs at the later time points (Figures 5A-5C, S5F and S5G). We confirmed this delay in the formation

of an anti-inflammatory, pro-regenerative MC profile by assessing CD206, a surface marker commonly found on anti-

inflammatory MCs[51-53]. Aging and the deletion of *Postn* significantly reduced the accumulation of

$CD45^+CD11b^+F4/80^+CD206^+$ MCs in the late, regenerative phase at 3-, 4- and 8-dpi, regardless of gender (Figures

5D-5F, S5H and S5I). Lastly, we examined MCs in our FAP transplantation model (Figure 3H) and detected no

differences in overall MCs but a similar phenotype of significantly elevated CD206+ MCs in 8-dpi-injury sites of aged

mice transplanted with *Postn*-expressing FAPs compared to *Postn*-deficient transplants (Figures 5G and 5H). These

data reiterate the notion that loss of Periostin, similar to aging, delays the formation of a regeneration-supporting anti-

inflammatory niche.

Periostin promotes anti-inflammatory macrophages through integrin-FAK-AKT mechano-signal transduction

To investigate the mechanism of Periostin-dependent control of macrophage polarization, we co-cultured wildtype and

Postn$^{-/-}$-FAPs with bone marrow-derived macrophages (BMDM) in a trans-well system and compared them to known

pro- and anti-inflammatory stimuli. After 48h, FAPS-WT (*Postn$^{+/+}$*), but not FAPs-KO (*Postn$^{-/-}$*), significantly induced

expression of anti-inflammatory MC markers *Mrc1* and *Cd163* while other anti-inflammatory genes, *Il10*, and several

pro-inflammatory markers were not affected by the presence of FAPs (Figures 6A and S6A). Addition of species-

specific recombinant Periostin proteins (mrPOSTN, hrPOSTN) to murine BMDM- and human, THP-1 cell line-

derived macrophages resulted in a significant, dose-dependent increase in the expression of anti-inflammatory markers

Mrc1, *Cd163* and *Il10*, but did not affect pro-inflammatory gene expression (Figures 6B, S6B and S6D). Integrins, and

specifically Itgαv/β3, Itgαv/β5 heterodimers, have been linked to anti-inflammatory macrophage function[54,55] and our

own findings and published data indicate that Periostin as able to bind both integrin heterodimers[33,34]. Treatment of

BMDM and THP-1 macrophages with Cilengitide, an inhibitor of ITGαv/β3 and ITGαv/β5 heterodimers[56,57], resulted

in a significant reduction of rPOSTN-induced expression of anti-inflammatory surface protein CD206 and similar

effects on *Mrc1*, which encodes for CD206, and *Cd163* mRNA expression, whereas recombinant IL4-induced

expression of these genes was not affected by Cilengitide and little or no effect on other pro-inflammatory genes was

evident (Figures 6C-6E, S6C, S6E and S6F). Protein kinase B (AKT) and focal adhesion kinase (FAK) phosphorylation

have been linked to anti-inflammatory macrophage polarization[58,59] and both kinases are potential downstream targets

of the Periostin-ITGαvβ3 and -αvβ5 axis[33,34,60]. Accordingly, rPOSTN acutely induced the phosphorylation of FAK at

Tyr397 and of AKT at Ser437, which was reduced significantly after Cilengitide addition in BMDM and THP-1

macrophages (Figures 6F, 6G, S6G and S6H). Collectively, these results indicate that rPOSTN induces M2-like

polarization through integrin-dependent FAK-AKT signaling in macrophages.

Aging and fibrotic scar tissue formation during muscle regeneration have been associated with altered muscle stiffness,

somewhat resembling the reduced muscle stiffness we observed in our *Postn$^{-/-}$* mice (Figure S2F) [12,23]. During muscle

regeneration, a transient decrease in tissue stiffness at the injury site around 5-dpi, which subsequently rebounds to

normal stiffness levels in fully regenerated muscle, has been reported recently[61]. To investigate the relationship of

Periostin and the mechanical properties of the local niche, and their impact on macrophage polarization, BMDM were

encapsulated in a three-dimensional ECM microenvironment using different alginate-based hydrogels with either low

or high mechanical stiffness. Consistent with previous reports[62], low stiffness hydrogels contained significantly fewer

F4/80$^+$/CD206$^+$ anti-inflammatory MCs compared to high stiffness hydrogels, while the opposite was true for

F4/80$^+$/CD86$^+$ pro-inflammatory MCs, which were more abundant under low stiffness and practically absent under

high stiffness conditions. These data suggest that mechanical input from the microenvironment directly affects

macrophage polarization and imply that loss of *Postn* may exert a similar effect in the muscle niche. Importantly, the

addition of murine rPOSTN to the hydrogels resulted in a Cilengitide-sensitive, modulatory effect where pro-

inflammatory polarization induced by low stiffness was significantly reduced. Instead, an anti-inflammatory MC

profile was promoted under conditions of low stiffness, suggesting that Periostin acts as a niche factor that synchronizes

biomechanical and immunomodulatory processes within the muscle niche (Figures 6H, 6I and S7A-S7C).

274 Circulating Periostin may predict muscle dysfunction in geriatric patients

Circulating Periostin has been proposed as biomarker of several inflammatory, chronic and metabolic diseases[39-42], but

its role in relation to aging in mice and humans is only now emerging[22,43]. To explore the link between aging and

muscle dysfunction, we measured circulating human Periostin (hPOSTN) in serum samples from two separate human

study cohorts[63,64]. When comparing serum Periostin from young adults to healthy older, community-dwelling

individuals, significantly lower levels of circulating hPOSTN were observed in aged individuals. Additionally, we

found that increased physical activity in elderly individuals restored hPOSTN-concentrations to levels found in young

adults (Figure 7A). As previous studies had reported elevated levels of circulating Periostin due to certain pathologies[65],

we also assessed serum hPOSTN in a group of healthy older individuals compared to age-matched individuals living

in an acute geriatric hospital setting. No differences were found, indicating that circulating hPOSTN is not applicable

as a generalized marker of impaired health status (Figure 7B). Next, we performed correlations including young adults

to healthy older community-dwelling individuals only (from panel 7A), and we found an inverse correlation between

age and circulating hPOSTN serum levels in women, but not in men, which could be due to lower inclusion numbers

for men (Figure 7C). Using all individuals of this study cohort, we subsequently examined whether other metabolic

and functional muscle parameters would also correlate with circulating hPOSTN. Indeed, hPOSTN correlated

inversely with body mass index (BMI) and its serum concentration was positively corelated with skeletal muscle mass

and grip strength in women, but not men (Figures 7D-7F). These data suggest that circulating Periostin levels could be

used as a potential biomarker of skeletal muscle aging and the age-related sarcopenia syndrome.

Discussion

Muscle injury repair requires complex interactions between different cell types residing within the stem cell niche[47].

For instance, extensive evidence exists for FAP-MuSC interactions to ensure successful myogenic regeneration[10,13-

15,23,24,66]. Aging and other pathologies result in defective myogenesis due to stem cell niche dysfunction by inhibiting

different aspects of myogenic stem cell maintenance, activation and differentiation[4]. We here report on a niche-

dependent mechanism that integrates paracrine signals and mechano-signal transduction to impact on the levels of

MuSCs and the myogenesis-regulatory FAPs, and by exerting immunomodulatory effects regulating the inflammatory

profile of injury site-associated macrophages. De-synchronization of these processes creates a defective stem cell niche

that phenocopies the detrimental effects of aging on myogenic regeneration, which result in enhanced fatty

degeneration of muscle tissue instead.

Firstly, we show that loss of the matricellular protein, Periostin, which is secreted by muscle-resident FAPs, directly

impinges on MuSCs, attenuating their ability to expand in response to injury and to differentiate into fully functional

myofibers. These findings are consistent with recent studies showing that aging alters secretion of signaling molecules

like the WNT1-inducible signaling pathway protein 1 (WISP1), bone morphogenetic protein 3 (BMP3b) and SPARC-

related modular calcium-binding protein 2 (SMOC2) by FAPs in the stem cell niche, thereby reducing MuSCs'

regeneration capacity[10,11,67].

Previous reports have established that aging and chronic disease prime FAPs to form adipocytes that accumulate

between muscle fibers, leading to an impairment of muscle function[2,6,16]. In line with these observations, our animal

models with loss of Periostin due to age or genetic targeting display defective autoregulation of FAPs, resulting in

excessive FAP-expansion upon injury which feature an elevated commitment towards the adipogenic lineage at the

expense of their ability to maintain appropriate myogenic regeneration capacity. This process results in the formation

of a pro-fibro/adipogenic microenvironment with enhanced fibro/fatty infiltration instead of successful muscle repair.

A tight regulation of FAP expansion and clearance is essential to maintain tissue integrity and prevent fibro/fatty

degeneration. However, the mechanisms through which aging may affect FAPs' expansive capacities in homeostatic

and regenerating muscle tissues remain unclear. Previous reports have described a reduction of FAPs in uninjured

muscle during aging[10,24]. Upon injury the situation could depend on the specifics of the experimental layout,

physiological status and age, as age-related failure of FAPs to expand has been reported[10], while our own observations

suggest enhanced FAP expansion, which align with scRNA-seq and tissue-array data at steady-state in old muscle[8,68].

Intramuscular FAP transplants into mice with impaired myogenesis have been shown to rescue the normal

characteristics of healthy muscle under homeostatic conditions and improved regeneration capacity[10,14,21]. Our

FAP/MuSC co-cultures and FAPs-transplants into muscle injuries of aged mice help restore secretion of Periostin and

are sufficient to reconstitute myogenesis and muscle regeneration. Our findings therefore support the concept that

FAP-secreted ECM factors are necessary and sufficient to sustain myogenesis via the FAP-MuSC crosstalk.

Our work sheds additional light on the concept of ECM-linked factors that exert immunomodulatory effects by

targeting the pro- to anti-inflammatory transition of macrophages within the muscle injury site. This process integrates

mechanical properties of the ECM and its changes during muscle repair with mechanosensory signaling through AKT

and FAK pathways. When disrupted, for instance through loss of Periostin, this results in a delay of the pro- to anti-

inflammatory switch within the post-injury myogenic niche and is also observed in aging muscle. The crosstalk

between FAPs and muscle-resident immune cell subsets to sustain adequate muscle regeneration has been reported on

before[23,26-28]. Nevertheless, the exact mechanisms of interaction between FAPs and macrophage polarization associated

with muscle regeneration capacity, and their alterations during the aging process, remain largely unknown. We here

introduce the concept that niche factors like Periostin may help to synchronize the biological signaling processes of

immunomodulation and biomechanics during muscle regeneration. Previous reports have shown that FAPs coordinate

myeloid and lymphoid cell contributions to inflammatory processes during the regeneration process[23,26-28]. Our scRNA-

seq analysis of regeneration in young and aged muscle shows that myeloid and lymphoid subsets are the main clusters

affected by aging. In line with these observations, experimental reconstitution of macrophage-derived signaling

molecules growth differentiation factor 3 (GDF3) and mesencephalic astrocyte-derived neurotrophic factor (MANF)

improved regeneration[31,69]. These results also support the concept of reinstating a normal macrophage function to

ensure successful regeneration in the context of aging. This could be achieved by targeting regulatory FAPs to provide

appropriate immunomodulatory signals, such as Periostin, to orchestrate the pro- to anti-inflammatory macrophage

transition. In our hands, FAPs transplantation with functional Periostin expression into aged, post-injury muscle

restored a timely onset of anti-inflammatory niche formation and successful regeneration. These findings are in line

with previous results showing that Periostin could support a shift from a pro- towards an anti-inflammatory phenotype

to sustain heart, kidney and bladder tissue regeneration, although a molecular mechanism has so far not been reported[51-]

[53]. Thus, our data link FAPs-derived Periostin secretion to the muscle stem cell niche with synchronizing effects on

FAPs, MuSCs and macrophages during muscle healing. Periostin exerts its function through integrin receptors and

mainly through $\alpha v \beta 3$ and $\alpha v \beta 5$ heterodimers in different biological contexts[33-35]. Analyses of CCIs in the muscle stem

cell niche corroborate that Periostin is a FAPs-secreted ligand that mediates intercellular communication between

FAPs, MuSCs and macrophages. Interestingly, the Periostin:Itgαv/β5 interaction, which mediates the crosstalk

between FAPs and macrophages, declines with aging in the model, thus mirroring our own experimental evidence and

presenting a new mechanism of age-related macrophage dysfunction and niche deterioration where a relative lack of

Periostin fails to activate FAK-AKT signaling. This in turn attenuates the transition from a pro- towards an anti-

inflammatory milieu in the regenerating muscle niche and is in line with other studies that implicate Periostin in

regulation of macrophage function in other organs and physiological contexts[34,51,59]. Corresponding

immunomodulatory mechanisms have been described where the integrin-αvβ5 heterodimer regulates anti-

inflammatory macrophages via a PPAR-γ dependent mechanism[55], suggesting that this signaling pathway may also

integrate different immunomodulatory signals during myogenic regeneration.

In addition to its role as a signaling molecule, our data also suggest a link between Periostin and mechanosensory

signal transduction. The ability of matrices with different mechanical properties to regulate macrophage polarization

is well-documented in the literature[62]. Molecules like Periostin could offer an additional layer of regulatory input that

offsets or enhances the role of the local ECMs' mechanical profile on inflammatory processes, including macrophage

functions, thereby also impacting on myogenic stem cells and fatty degeneration of muscle. Aging increases muscle

stiffness[12], a mechanical property which would normally promote anti-inflammatory macrophages. It should be noted

that such a situation would only occur in aged muscle at steady-state, i.e. non-injury conditions, and has indeed been

reported[70]. In intact muscle, however, overall numbers of tissue macrophages are comparably low. Conversely, post-

injury muscles experience dynamic ECM-remodeling, marked by an initial, but transient decline of stiffness at 5-dpi[61],

which in fact coincides with the pro- to anti-inflammatory niche/ macrophage transition. As low stiffness would

preferentially stimulate pro-inflammatory macrophages, the presence of Periostin could mediate or at last accelerate

the transition to an anti-inflammatory microenvironment.

Lastly, circulating Periostin has been proposed as a potential biomarker in some instances such as for cardiovascular

risk, early cancer detection, or different type 2 inflammatory diseases[39-42]. Our own and other previous work suggests

that reduced gene expression levels of *Postn* are reliable predictors of aging and associated metabolic disorders[22,43].

Thus, it stands to reason that loss of circulating Periostin may also inform on the progression of muscle degeneration

as a result of age-related sarcopenia. It should be noted that increased levels of Periostin have also been proposed as a

biomarker of disease progression in older adults, including cognitive decline[65], suggesting that associations such as

those found in our cohorts need to be evaluated in more detail. In conclusion, our study presents a novel role of FAP-

based synchronization of the muscle stem cell niche, which links impaired regeneration to age-related niche

deterioration and fatty degeneration of skeletal muscle. Our results provide mechanistic insights into a Periostin-

dependent crosstalk between FAPs and macrophages that is impaired during aging, which integrates paracrine niche

effects and mechano-sensing to orchestrate the precisely timed steps of successful myogenic regeneration instead of

fibro-fatty infiltration.

Acknowledgments

The authors thank A. Kretschmer, S. Marg, S. Richter, A. Weber and N. Dittberner for expert technical assistance.

This work was supported by grants from the German Research Foundation (DFG) as individual grants and within the

Collaborative Research Centre CRC1444 (project IDs 323196138, 249509554, and 427826188), and by a grant from

the Leibniz Association (ID K398/2021, to TJS). Grants within the German Center for Diabetes Research (DZD)

funded by the German Ministry of Education and Research (BMBF) and the State of Brandenburg (DZD grant IDs

82DZD00302, 82DZD03D03 and 82DZD03C3G) are acknowledged.

Author Contributions

F.G.-C. and T.J.S. conceived the study and wrote the manuscript. F.G.-C. conducted the majority of experiments. S.G.,

G.L.-G., A.-M.J., C.G., S.P.-K., A.T., A.G., G.K. and M.Os., contributed research data to this study. G.S., L.V. und

V.B. performed scRNA-seq experiments. M.L. and M.Ov. performed computational analyses of scRNA-seq data.

C.H., U.M.-W., K.N., provided human serum samples and helped with data analysis. S.K., A.S., S.S., K.S.-B., A.C.,

R.S.K., and G.N.D. contributed valuable materials and expertise to the article.

Declaration of interests

The authors declare no conflict of interest.

References

1. Goodpaster, B.H.*, et al.* The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging

and body composition study. *J. Gerontol. A Biol. Sci. Med. Sci.* **61**, 1059-1064 (2006).

2. Marcus, R.L., Addison, O., Kidde, J.P., Dibble, L.E. & Lastayo, P.C. Skeletal muscle fat infiltration: impact of age,

inactivity, and exercise. *J. Nutr. Health Aging* **14**, 362-366 (2010).

3. Brack, A.S.*, et al.* Increased Wnt signaling during aging alters muscle stem cell fate and increases fibrosis. *Science*

317, 807-810 (2007).

4. Chakkalakal, J.V., Jones, K.M., Basson, M.A. & Brack, A.S. The aged niche disrupts muscle stem cell quiescence.

Nature **490**, 355-360 (2012).

5. Cosgrove, B.D.*, et al.* Rejuvenation of the muscle stem cell population restores strength to injured aged muscles.

Nat. Med. **20**, 255-264 (2014).

6. Farup, J.*, et al.* Human skeletal muscle CD90(+) fibro-adipogenic progenitors are associated with muscle

degeneration in type 2 diabetic patients. *Cell Metab.* **33**, 2201-2214 e2211 (2021).

7. Garcia-Prat, L.*, et al.* Autophagy maintains stemness by preventing senescence. *Nature* **529**, 37-42 (2016).

8. Lazure, F.*, et al.* Transcriptional reprogramming of skeletal muscle stem cells by the niche environment. *Nat.*

Commun. **14**, 535 (2023).

9. Lukjanenko, L.*, et al.* Loss of fibronectin from the aged stem cell niche affects the regenerative capacity of skeletal

muscle in mice. *Nat. Med.* **22**, 897-905 (2016).

10. Lukjanenko, L.*, et al.* Aging Disrupts Muscle Stem Cell Function by Impairing Matricellular WISP1 Secretion

from Fibro-Adipogenic Progenitors. *Cell Stem Cell* **24**, 433-446 e437 (2019).

11. Schuler, S.C.*, et al.* Extensive remodeling of the extracellular matrix during aging contributes to age-dependent

impairments of muscle stem cell functionality. *Cell Rep* **35**, 109223 (2021).

12. Stearns-Reider, K.M.*, et al.* Aging of the skeletal muscle extracellular matrix drives a stem cell fibrogenic

conversion. *Aging Cell* **16**, 518-528 (2017).

13. Joe, A.W.*, et al.* Muscle injury activates resident fibro/adipogenic progenitors that facilitate myogenesis. *Nat. Cell.*

Biol. **12**, 153-163 (2010).

14. Uezumi, A., Fukada, S., Yamamoto, N., Takeda, S. & Tsuchida, K. Mesenchymal progenitors distinct from satellite

cells contribute to ectopic fat cell formation in skeletal muscle. *Nat. Cell. Biol.* **12**, 143-152 (2010).

15. Wosczyna, M.N.*, et al.* Mesenchymal Stromal Cells Are Required for Regeneration and Homeostatic Maintenance

of Skeletal Muscle. *Cell Rep.* **27**, 2029-2035 e2025 (2019).

16. Hogarth, M.W., *et al.* Fibroadipogenic progenitors are responsible for muscle loss in limb girdle muscular

dystrophy 2B. *Nat. Commun.* **10**, 2430 (2019).

17. Kopinke, D., Roberson, E.C. & Reiter, J.F. Ciliary Hedgehog Signaling Restricts Injury-Induced Adipogenesis.

Cell **170**, 340-351 e312 (2017).

18. Madaro, L., *et al.* Denervation-activated STAT3-IL-6 signalling in fibro-adipogenic progenitors promotes

myofibres atrophy and fibrosis. *Nat. Cell Biol.* **20**, 917-927 (2018).

19. Perez-Diaz, S., *et al.* Obesity impairs skeletal muscle repair through NID-1 mediated extracellular matrix

remodeling by mesenchymal progenitors. *Matrix Biol.* **112**, 90-115 (2022).

20. Uezumi, A., *et al.* Fibrosis and adipogenesis originate from a common mesenchymal progenitor in skeletal muscle.

J. Cell Sci. **124**, 3654-3664 (2011).

21. Mozzetta, C., *et al.* Fibroadipogenic progenitors mediate the ability of HDAC inhibitors to promote regeneration

in dystrophic muscles of young, but not old Mdx mice. *EMBO Mol. Med.* **5**, 626-639 (2013).

22. Schaum, N., *et al.* Ageing hallmarks exhibit organ-specific temporal signatures. *Nature* **583**, 596-602 (2020).

23. Kotsaris, G., *et al.* Odd skipped-related 1 controls the pro-regenerative response of fibro-adipogenic progenitors.

NPJ Regen. Med. **8**, 19 (2023).

24. Uezumi, A., *et al.* Mesenchymal Bmp3b expression maintains skeletal muscle integrity and decreases in age-related

sarcopenia. *J. Clin. Invest.* **131**(2021).

25. Arnold, L., *et al.* Inflammatory monocytes recruited after skeletal muscle injury switch into antiinflammatory

macrophages to support myogenesis. *J. Exp. Med.* **204**, 1057-1069 (2007).

26. Heredia, J.E., *et al.* Type 2 innate signals stimulate fibro/adipogenic progenitors to facilitate muscle regeneration.

Cell **153**, 376-388 (2013).

27. Kuswanto, W., *et al.* Poor Repair of Skeletal Muscle in Aging Mice Reflects a Defect in Local, Interleukin-33-

Dependent Accumulation of Regulatory T Cells. *Immunity* **44**, 355-367 (2016).

28. Lemos, D.R., *et al.* Nilotinib reduces muscle fibrosis in chronic muscle injury by promoting TNF-mediated

apoptosis of fibro/adipogenic progenitors. *Nat. Med.* **21**, 786-794 (2015).

29. Cui, C.Y., *et al.* Skewed macrophage polarization in aging skeletal muscle. *Aging Cell* **18**, e13032 (2019).

30. Moiseeva, V., *et al.* Senescence atlas reveals an aged-like inflamed niche that blunts muscle regeneration. *Nature*

613, 169-178 (2023).

461 31. Patsalos, A., *et al.* In vivo GDF3 administration abrogates aging related muscle regeneration delay following acute

462   sterile injury. *Aging Cell* **17**, e12815 (2018).

463 32. Reidy, P.T., *et al.* Aging impairs mouse skeletal muscle macrophage polarization and muscle-specific abundance

464   during recovery from disuse. *Am. J. Physiol. Endocrinol Metab* **317**, E85-E98 (2019).

465 33. Kuhn, B., *et al.* Periostin induces proliferation of differentiated cardiomyocytes and promotes cardiac repair. *Nat.*

466   *Med.* **13**, 962-969 (2007).

467 34. Li, G., *et al.* Periostin mediates vascular smooth muscle cell migration through the integrins alphavbeta3 and

468   alphavbeta5 and focal adhesion kinase (FAK) pathway. *Atherosclerosis* **208**, 358-365 (2010).

469 35. Norris, R.A., *et al.* Periostin regulates collagen fibrillogenesis and the biomechanical properties of connective

470   tissues. *J. Cell Biochem.* **101**, 695-711 (2007).

471 36. Latroche, C., *et al.* Coupling between Myogenesis and Angiogenesis during Skeletal Muscle Regeneration Is

472   Stimulated by Restorative Macrophages. *Stem Cell Reports* **9**, 2018-2033 (2017).

473 37. Ozdemir, C., *et al.* Periostin is temporally expressed as an extracellular matrix component in skeletal muscle

474   regeneration and differentiation. *Gene* **553**, 130-139 (2014).

475 38. Scott, R.W., Arostegui, M., Schweitzer, R., Rossi, F.M.V. & Underhill, T.M. Hic1 Defines Quiescent

476   Mesenchymal Progenitor Subpopulations with Distinct Functions and Fates in Skeletal Muscle Regeneration. *Cell*

477   *Stem Cell* **25**, 797-813 e799 (2019).

478 39. Azharuddin, M., *et al.* Predictor for cardiovascular risk in patients with type-2 diabetes mellitus. *Clin. Epidemiol.*

479   *Glob. Health* **12** (2021).

480 40. Rachner, T.D., *et al.* High serum levels of periostin are associated with a poor survival in breast cancer. *Breast*

481   *Cancer Res. Treat.* **180**, 515-524 (2020).

482 41. Yang, Z., *et al.* Circulating periostin in relation to insulin resistance and nonalcoholic fatty liver disease among

483   overweight and obese subjects. *Sci. Rep.* **6**, 37886 (2016).

484 42. Zhao, S., *et al.* Periostin expression is upregulated and associated with myocardial fibrosis in human failing hearts.

485   *J. Cardiol.* **63**, 373-378 (2014).

486 43. Graja, A., *et al.* Loss of periostin occurs in aging adipose tissue of mice and its genetic ablation impairs adipose

487   tissue lipid metabolism. *Aging Cell* **17**, e12810 (2018).

488 44. Mahdy, M.A.A. Glycerol-induced injury as a new model of muscle regeneration. *Cell Tissue Res.* **374**, 233-241

489   (2018).

45. Ito, N., Miyagoe-Suzuki, Y., Takeda, S. & Kudo, A. Periostin Is Required for the Maintenance of Muscle Fibers

during Muscle Regeneration. *Int. J. Mol. Sci.* **22**(2021).

46. Muzumdar, M.D., Tasic, B., Miyamichi, K., Li, L. & Luo, L. A global double-fluorescent Cre reporter mouse.

Genesis **45**, 593-605 (2007).

47. Mashinchian, O., Pisconti, A., Le Moal, E. & Bentzinger, C.F. The Muscle Stem Cell Niche in Health and Disease.

Curr. Top. Dev. Biol. **126**, 23-65 (2018).

48. De Micheli, A.J.*, et al.* Single-Cell Analysis of the Muscle Stem Cell Hierarchy Identifies Heterotypic

Communication Signals Involved in Skeletal Muscle Regeneration. *Cell Rep.* **30**, 3583-3595 e3585 (2020).

49. Lagger, C.*, et al.* scAgeCom: a murine atlas of age-related changes in intercellular communication inferred with

the package scDiffCom. *BioRxiv* https://doi.org/10.1101/2021.08.13.456238 (2021).

50. Varga, T.*, et al.* Highly Dynamic Transcriptional Signature of Distinct Macrophage Subsets during Sterile

Inflammation, Resolution, and Tissue Repair. *J. Immunol.* **196**, 4771-4782 (2016).

51. Chen, Z.*, et al.* Periostin Attenuates Cyclophosphamide-induced Bladder Injury by Promoting Urothelial Stem Cell

Proliferation and Macrophage Polarization. *Stem Cells Transl. Med.* **11**, 659-673 (2022).

52. Kormann, R.*, et al.* Periostin Promotes Cell Proliferation and Macrophage Polarization to Drive Repair after AKI.

J. Am. Soc. Nephrol. **31**, 85-100 (2020).

53. Liao, Y.*, et al.* Cardiac Nestin(+) Mesenchymal Stromal Cells Enhance Healing of Ischemic Heart through

Periostin-Mediated M2 Macrophage Polarization. *Mol. Ther.* **28**, 855-873 (2020).

54. Antonov, A.S., Kolodgie, F.D., Munn, D.H. & Gerrity, R.G. Regulation of macrophage foam cell formation by

alphaVbeta3 integrin: potential role in human atherosclerosis. *Am. J. Pathol.* **165**, 247-258 (2004).

55. Yao, Q.*, et al.* Peroxisome proliferator-activated receptor gamma (PPARgamma) induces the gene expression of

integrin alpha(V)beta(5) to promote macrophage M2 polarization. *J. Biol. Chem.* **293**, 16572-16582 (2018).

56. Lin, X.*, et al.* Omentin-1 Modulates Macrophage Function via Integrin Receptors alphavbeta3 and alphavbeta5 and

Reverses Plaque Vulnerability in Animal Models of Atherosclerosis. *Front. Cardiovasc. Med.* **8**, 757926 (2021).

57. Zhou, W.*, et al.* Periostin secreted by glioblastoma stem cells recruits M2 tumour-associated macrophages and

promotes malignant growth. *Nat. Cell. Biol.* **17**, 170-182 (2015).

58. Po-Chun Chen, H.-C.C., John Wang, Shin-Wei Wang, Huai-Ching Tai, Chiao-Wen Lin, and Chih-Hsin Tang.

Prostate cancer-derived CCN3 induces M2 macrophage infiltration and contributes to angiogenesis in prostate

cancer microenvironment. *Oncotarget* (2014).

59. Vergadi, E., Ieronymaki, E., Lyroni, K., Vaporidi, K. & Tsatsanis, C. Akt Signaling Pathway in Macrophage

Activation and M1/M2 Polarization. *J. Immunol.* **198**, 1006-1014 (2017).

60. Ghatak, S*., et al.* Periostin induces intracellular cross-talk between kinases and hyaluronan in atrioventricular

valvulogenesis. *J. Biol. Chem.* **289**, 8545-8561 (2014).

61. Silver, J.S*., et al.* Injury-mediated stiffening persistently activates muscle stem cells through YAP and TAZ

mechanotransduction. *Sci. Adv.* **7**(2021).

62. Chen, M*., et al.* Substrate stiffness modulates bone marrow-derived macrophage polarization through NF-kappaB

signaling pathway. *Bioact. Mater.* **5**, 880-890 (2020).

63. Herpich, C*., et al.* Associations Between Serum GDF15 Concentrations, Muscle Mass, and Strength Show Sex-

Specific Differences in Older Hospital Patients. *Rejuvenation Res.* **24**, 14-19 (2021).

64. Herpich, C*., et al.* Postprandial dynamics and response of fibroblast growth factor 21 in older adults. *Clin. Nutr.*

40, 3765-3771 (2021).

65. Sanchez-Sanchez, J.L*., et al.* Periostin Plasma Levels and Changes on Physical and Cognitive Capacities in

Community-Dwelling Older Adults. *J. Gerontol. A. Biol. Sci. Med. Sci.* **78**, 424-432 (2023).

66. Mathew, S.J*., et al.* Connective tissue fibroblasts and Tcf4 regulate myogenesis. *Development* **138**, 371-384 (2011).

67. Wang, Y*., et al.* Functional regeneration and repair of tendons using biomimetic scaffolds loaded with recombinant

periostin. *Nat. Commun* **12**, 1293 (2021).

68. Wang, Y.X*., et al.* A single cell spatial temporal atlas of skeletal muscle reveals cellular neighborhoods that

orchestrate regeneration and become disrupted in aging. *bioRxiv* https://doi.org/10.1101/2022.06.10.494732

(2022).

69. Sousa, N.S*., et al.* Aging disrupts MANF-mediated immune modulation during skeletal muscle regeneration. *Nat*

Aging **3**, 585-599 (2023).

70. Wang, Y., Wehling-Henricks, M., Samengo, G. & Tidball, J.G. Increases of M2a macrophages and fibrosis in aging

muscle are influenced by bone marrow aging and negatively regulated by muscle-derived nitric oxide. *Aging Cell*

14, 678-688 (2015).

Figure legends

Figure 1: Aging impairs myogenic regeneration by disrupting FAPs-derived niche signals. (**A**) Representative

hematoxylin/eosin staining images of gastrocnemius muscle cross-sections showing inter-myocellular adipocyte

infiltration between myofibers from young (2 months) and aged (age: 15-18 months; pertains to all subsequent panels)

male mice (red arrows indicate adipocytes; scale bar: 100 µm). (**B**) Quantifications of flow cytometric analysis of

CD45$^-$CD31$^-$SCA1$^+$ FAPs and CD45$^-$CD31$^-$SCA1$^-$ITGA7$^+$ MuSCs per grams of murine muscle tissue (n=18/group).

White bars indicate young mice/cells (age: 2-2.5 months), blue bars indicate aged mice/cells (age: 15-18 months;

pertains to all subsequent panels). (**C**) Representative images and quantification of spontaneous adipogenic

differentiation of FAPs (green: BODIPY, lipid droplet; blue: DAPI, nuclei; n=6; 2 independent experiments; scale bar:

50 µm). (**D**) Representative images and fusion index quantifications in differentiated MuSC (green: Desmin,

myofibers; blue: DAPI-nuclei; n=6; 2 independent experiments; scale bar, 100 µm). (**E**) Heat map of top-20 down-

regulated ECM genes in microarray analysis comparing freshly sorted FAPs from uninjured skeletal muscle. (**F**)

Periostin protein detection and quantification by western blot (Actin: loading control; n=5/group). (**G**) *Postn* mRNA

expression in uninjured gastrocnemius (GS), tibialis anterior (TA) and soleus (SL) muscles (n=5/group). (**H**) Periostin

protein detection by western blot of cell culture supernatants of FAPs collected during adipogenic differentiation 48h

after last medium change (C+: mouse rPOSTN; TP: total protein loading control). (**I, J**) FACS-analysis of CD45$^-$

CD31$^-$SCA1$^+$ FAPs per grams of TA muscle comparing uninjured to injured TA at 4- and 8-dpi (n=3/group for

uninjured and 8-dpi; n=6/group for 4-dpi). (**K**) Representative hematoxylin/eosin staining and quantification of

adipocyte infiltration in injury sites of TA at 14-dpi (yellow arrows indicate adipocyte areas; n=3/group; scale bar: 50

µm). (**L**) Representative Sirius red staining and quantification of fibrotic tissue deposition in TA injury sites at 14-dpi

(blue arrows indicate fibrosis areas; n=3/group; scale bar: 50 µm). (**M**) Representative images and CSA quantification

of fiber size ranges in injury sites at 14-dpi (n=4/group; scale bar: 100 µm). (**N**) Representative images of Periostin

protein immunofluorescence in injury sites in TA at 8-dpi (red: Periostin; blue: DAPI, nuclei; scale bar: 50 µm). (**O**)

Postn mRNA levels in uninjured TA-muscles compared to TA at 3-, 5-, 8- and 14-dpi (n=4/group). (**P**) Periostin protein

western blots and quantification (Actin: loading control) in TA at 3-, 5-, and 8-dpi (n=4/group). (**Q**) UMAP

dimensionality reduction of scRNA-seq analysis of 287 CD45$^-$CD31$^-$SCA1$^+$ FAPs FACS-isolated at 4-dpi from young

(FAPS-2mo, 128 cells; blue) and aged (FAPs-15mo, 159 cells; red) mice generated using microfluidics-based

technology, and (**R**) volcano plot of DEGs (p<0.05 and |Log2|fold change ≥0.5 as cutoffs; black arrow indicates *Postn*).

(**S**) Violin plot depiction of *Postn* gene expression per cell in scRNA-seq dataset. (**T**) Enriched Gene Ontology terms

of biological process (BP) and cellular component (CC) of downregulated DEGs from scRNA-seq dataset. Data were

collected from male mice except in (J, female). All data are presented as mean ± standard error of the mean (SEM),

statistical significances are *p<0.05, **p<0.01, ***p<0.001.

Figure 2: Ablation of *Postn* gene recapitulates an aging-like phenotype in muscle. (**A**) Representative images and quantification of spontaneous adipogenic differentiation of FAPs from *Postn*-KO (black bar) compared to wildtype (WT) littermates (gray bar; green: BODIPY, lipid droplet; blue: DAPI, nuclei; n=8/group; 2 independent experiments; scale bar: 50 µm). (**B**) Representative images and fusion index quantification in differentiated MuSC from *Postn*[-/-] (KO; black bar) compared to WT littermates (gray bar; green: Desmin, myofibers; blue: DAPI, nuclei; n=3/group; scale bar: 100 µm). (**C, D**) Flow cytometric analysis of CD45[-]CD31[-]SCA1[+] FAPs (panel C) and CD45[-]CD31[-]SCA1[-] ITGA7[+] MuSCs (panel D) per grams of TA-muscle comparing uninjured to injured TA at 4- and 8-dpi (males: uninj n=4/3, 4-dpi n=4/3, 8-dpi n=4/4; females: uninj n=10/10, 4-dpi n=7/7, 8-dpi n=4/4). Data show comparisons of male WT (gray bars, black line) to KO (orange bars and line) and female WT (gray bars, black line) to KO (green bars and line) mice (also applies to subsequent panels). (**E**) Representative hematoxylin/eosin staining and quantification of adipocyte infiltration in injury sites of TA at 14-dpi (yellow arrows indicate adipocyte areas; males: n=3/group, females: n=6/7/group; scale bar: 50 µm). (**F**) Representative Sirius red staining and quantification of fibrotic tissue deposition in TA injury sites at 14-dpi (black arrows indicate fibrosis areas; males: n=3/group, females: n=3/4/group; scale bar: 50 µm). (**G**) Representative images, CSA quantification of fiber size ranges and average in injury site at 14-dpi (males: n=3/group; females: n=4/group; scale bar, 100 µm). All data are presented as mean ± standard error of the mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001.

Figure 3: FAPs-secreted Periostin rescues age-related defect of myogenic regeneration. (**A**-**D**) Proliferation analysis by KI67 immunostaining of FAPs and MuSCs FACS-isolated from young (Y, 2 months) and aged (O, 15 months) mice and young *Postn*[-/-] (KO) and littermate wildtype (WT) control mice and treated with mrPOSTN (0.5 µg/ml) for 48h (n=6-12; two independent experiments). (**E**) Representative images of co-cultures from FAPs-WT and FAPs-KO with MuSCs-KO co-cultured for 5 days in myogenic differentiation medium (green: PDGFRα, FAPs; red: Desmin, myofibers; blue: DAPI, nuclei; scale bar: 50µm). (**F, G**) Quantifications of fusion index (% of myofibers with >3 DAPI[+] nuclei) and myofiber length in co-cultures in panel E, compared to pure MuSC cultures from wildtype (MuSCs-WT) or *Postn*[-/-] mice (MuSCs-KO) (n=6 wells; two-independent experiments). (**H**) Experimental layout and timeline of FAPs transplants of young FAP[RFP]-WT and FAP[RFP]-KO into old recipient mice (TA-muscle, 15 months). (**I**) Representative immunofluorescence images of transplanted TA-muscle sections at 8-dpi (green: eMHC; red: Laminin; blue: DAPI, nuclei; scale bar: 100 µm). (**J, K**) Quantifications of eMHC (top) and CSA average (bottom) of

FAP[RFP]-WT and FAP[RFP]-KO transplants at 8-dpi (n=5/group). (**L, M**) Representative images and CSA quantification

of fiber size ranges in injury site at 14-dpi comparing 15-months TA-muscle transplanted with FAP[RFP]-WT or FAP[RFP]-

KO transplants (red: Laminin; blue: DAPI, nuclei; n=5-6/group; scale bar: 100 µm). (**N**) Quantification of CSA

averages in regenerating fibers comparing non-transplanted (as shown in Figure 1M) young (2 months; n=4) and old

(15 months; n=4) to transplanted 15-months old TA-muscles with either FAP[RFP]-WT (n=5) or FAP[RFP]-KO (n=6). All

data are presented as mean ± standard error of the mean (SEM), statistical significances are *p<0.05, **p<0.01,

***p<0.001.

Figure 4: Computational analysis uncovers a Periostin-dependent FAPs-macrophage interaction. (**A**) UMAP of

scRNA-seq data generated by droplet-based sequencing technology of all cells isolated from 4-dpi TA-muscle,

integrating single cell transcriptomic data of young (2 months; blue-green) and aged (15 month; reds) female C57BL/6J

mice. (**B, C**) Unsupervised Louvain clustering analysis and bar plot of population comparison in percentages of nine

distinct cell populations identified in regenerating muscle. (**D**) Violin plot showing *Postn* expression levels across the

nine cell populations. (**E**) Mean expression and percentage of cells expressing *Postn* in FAPs between 2- and 15-

months groups (p-value: Wilcoxon rank sum test). (**F, G**) Age-dependent cell-cell interaction (CCI) scoring of ligand

receptor interactions of Periostin with known receptors as a function of age (left panel, F) and interaction depiction of

age-related changes in ligand-receptor-driven CCIs linking log fold changes and adjusted p-values (right panel, G)

using a published input scRNA-seq dataset comparing young and aged female murine limb-muscle as described in[49].

Cell types within each CCI are color-indicated; red circles indicate statistically significant age-dependent changes in

CCI scores. (**H**) Violin plot showing expression for *Itgαv*, *Itga5*, I*tgβ5*, *Itgβ1*, *Itgβ3*, and *Egfr* receptors across

individual cells in the nine cell populations detected in our scRNA-seq dataset.

Figure 5: Aging and lack of Periostin impair macrophage polarization in regenerating muscle. (**A**)

Representative density plots from FACs analyses of macrophage populations comparing either young (MCs-2mo) to

aged (MCs-15mo), or young wildtype (MCs-WT) to littermate *Postn*[-/-] (MCs-KO) mice. Total MCs were initially gated

as CD45[+]CD11b[+]F4/80[+] and subsequently for Ly6C-expression to indicate Ly6C-high, pro-inflammatory (M1) vs.

Lyc6-low, anti-inflammatory (M2) MCs at 1-, 3-, 4-, and 8-dpi. (**B, C**) Quantitative analysis of MC frequencies as

described in panel A (n=3-4/group). Comparisons are young (MCs-2mo; white bars) to aged (MCs-15mo; blue bars)

and wildtype (MCs-WT; gray bars) to Postn[-/-] (MCs-KO; orange bars; also applies to panels E, F) mice. (**D**)

Representative density plots of anti-inflammatory F4/80$^+$CD206$^+$ MCs comparing groups as described in panels A-C.

MCs were initially gated as CD45$^+$CD11b$^+$. (**E**, **F**) Quantitative analysis of MCs frequencies as described in panel D

(n=3-4/group). (**G**, **H**) Representative immunofluorescence images and quantifications of F4/80$^+$ and F4/80$^+$CD206$^+$

MCs in injured site of 15-months old TA-muscle sections at 8-dpi transplanted with FAP^{RFP}-WT (blue bar) or FAP^{RFP}-

KO (brown bar) at 1-dpi (red: F4/80, macrophage-marker; green: CD206, M2-marker; blue: DAPI, nuclei; n=4/group;

yellow arrows indicate co-localization of F4/80$^+$ and CD206$^+$ MCs; scale bar: 20 µm). All animals were male, all data

are presented as mean ± standard error of the mean (SEM), statistical significances are *p<0.05, **p<0.01,

***p<0.001.

Figure 6: Periostin promotes anti-inflammatory niche transition through mechano-sensitive integrin-FAK-

AKT signaling. (**A**) *Mrc1* (Mannose receptor c-type 1) mRNA expression in differentiated BMDM exposed to pro-

and anti-inflammatory cytokines (mrIFNγ: 50 ng/mL; mrIL4: 10 ng/mL; n=8/treatment) or co-cultured with FAPs-WT

or FAPs-KO (n=11, co-culture group) for 48h compared to untreated BMDM control (n=6) in 3 independent

experiments. (**B**) *Mrc1* mRNA expression in differentiated BMDM treated with mrPOSTN at indicated concentrations

for 48h (n=3/group). (**C**, **D**) Representative immunofluorescence images and quantification in untreated, differentiated

BMDM control compared to a 48h treatment with mrPOSTN (1 µg/mL) alone or in combination with integrin receptor-

inhibitor Cilengitide (10 nM). mrIL4 (10 ng/mL) was used as positive control of M2-polarization (green: CD206, M2-

marker; red: F4/80, macrophage-marker; blue: DAPI, nuclei; n=3/treatment; scale bar: 50 µm). (**E**) *Mrc1* mRNA

expression in untreated BMDM control compared to BMDM 48h-exposed to Cilengitide (10nM), mrPOSTN (1

µg/mL), mrPOSTN + Cilengitide (1 µg/mL + 10nM), mrIL4 (10 ng/ml) or mrIL4 + Cilengitide (10 ng/mL + 10 nM)

(n=4/treatment). (**F, G**) Protein detection and quantification of phosphorylated Ser473-AKT (p-AKT) and total AKT

(panel F) and phosphorylated Tyr397-FAK (p-FAK) and total FAK (panel G) in untreated compared to treated BMDM

with mrPOSTN (1 µg/mL) alone or combined with Cilengitide (10 nM) for indicated times (Actin loading control;

n=4; 2 independent experiments in panels F, G). (**H, I**) Immunofluorescence quantifications of pro- F4/80$^+$CD86$^+$ and

anti-inflammatory F4/80$^+$CD206$^+$ MC polarization of differentiated BMDM encapsulated in soft or high stiffness

hydrogels untreated (control) or treated with mrPOSTN (1 µg/mL) and Cilengitide (10 nM) alone or combined

(mrPOSTN + Cilengitide) for 24h (n=12/group; 2 independent experiments). All data are presented as mean ± standard

error of the mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 7: Circulating Periostin functions as biomarker of muscle dysfunction in elderly patients. (**A**) Serum levels of human Periostin (hPOSTN) in young adults (white; n=59; n=16 men; n=43 women), healthy aged, community-dwelling adults (red; n=59; n=14 men; n=45 women), and aged community-dwelling adults with high activity levels (Aged/active; green; n=20; n=5 men; n=15 women). (**B**) Serum hPOSTN levels in community-dwelling aged adults (light blue; n=44), and geriatric hospital patients (Aged/patient; dark blue; n=34). (**C-F**) Pearson correlations of serum hPOSTN levels and individuals' age (panel C), body mass index (BMI, panel D), total skeletal muscle mass (panel E) and grip strength (panel F) separately in men (orange, left) and women (green, right), including young adults and aged community-dwelling individuals only, as described in panel A. All data are presented as mean ± standard error of the mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.