Aging impairs muscle regeneration by desynchronizing matrix mechano-signaling and macrophage immunomodulation via fibro-adipogenic progenitors

Francisco Garcia-Carrizo[1,2], Sabrina Gohlke[1], Georgia Lenihan-Geels[1], Anne-Marie Jank[1], Marina Leer[1,2], George A. Soultoukis[1], Masoome Oveisi[1,2], Catrin Herpich[3,4,5], Claudia A. Garrido[6,7], Georgios Kotsaris[8], Sophie Pöhle-Kronawitter[1,8], Arnold Tsamo-Tetou[1], Antonia Graja[1], Mario Ost[9], Laura Villacorta[10], Raphael S. Knecht[6], Susanne Klaus[9], Annette Schürmann[2,12], Sigmar Stricker[8], Katharina Schimdt-Bleek[6,11], Amaia Cipitria[6,7,13,14], Georg N. Duda[6,11], Vladimir Benes[10], Ursula Müller-Werdan[4,15], Kristina Norman[3,4,5], Tim J. Schulz[1,2,5,*]

1. Department of Adipocyte Development and Nutrition, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

2. German Center for Diabetes Research (DZD), München-Neuherberg, Germany

3. Department of Nutrition and Gerontology, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

4. Department of Geriatrics and Medical Gerontology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität Zu Berlin, Berlin, Germany

5. Institute of Nutritional Science, University of Potsdam, Potsdam-Rehbrücke, 14558 Nuthetal, Germany

6. Julius Wolff Institute, Berlin Institute of Health at Charité - Universitätsmedizin Berlin, Berlin, Germany

7. Department of Biomaterials, Max Planck Institute for Colloids and Interfaces, Potsdam, Germany

8. Institute of Chemistry and Biochemistry, Musculoskeletal Development and Regeneration Group, Freie Universität Berlin, Thielallee 63, 14195 Berlin, Germany

9. Department of Physiology of Energy Metabolism, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

10. Genomics Core Facility, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany

11. Berlin Institute of Health Centre for Regenerative Therapies (BCRT), Berlin Institute of Health at Charité - Universitätsmedizin Berlin, Berlin, Germany

12. Department of Experimental Diabetology, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany

13. Current address: Group of Bioengineering in Regeneration and Cancer, Biogipuzkoa Health Research Institute, San Sebastián, Spain

14. Current address: IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

15. Protestant Geriatric Center Berlin (EGZB), Berlin, Germany

*Corresponding author and Lead Contact

Tim J. Schulz, PhD

German Institute of Human Nutrition

114-116, Arthur-Scheunert Allee

Potsdam-Nuthetal, D-14558, Germany

Phone: +49-33200-88-2110;

Email: tim.schulz@dife.de

Summary

Skeletal muscle regeneration depends on the function of fibro/adipogenic progenitors (FAPs). Here we show that

aging impairs myogenic stem cells and immunomodulatory processes by disrupting extracellular matrix function,

thereby inhibiting regeneration in aged muscle. We identify the FAP-secreted matricellular protein Periostin as a

niche factor that is decreased in aged murine muscle and in circulation of aged humans with low-exercise lifestyle.

Periostin controls FAP-expansion after injury and its loss fate-regulates FAPs to contribute to fibro/fatty

infiltration. This causes stem cell defects and attenuates pro- to anti-inflammatory macrophage transition during

regeneration. Transplantation of young FAPs with high Periostin secretion, but not Periostin-deficient FAPs, into

aged muscle restores anti-inflammatory niche formation and regeneration. Mechanistically, Periostin activates

Focal adhesion kinase- and AKT-signaling in macrophages via integrin binding to promote an anti-inflammatory

profile, which synchronizes matrix-derived mechanosensory signaling and immunomodulation. These results

uncover a key role of FAP-based matrix regulation via niche factors that orchestrate successful muscle

regeneration.

Keywords: Aging, muscle stem cell niche, ECM-remodeling, FAPs-Periostin, satellite cells, macrophages

Introduction

Aging is characterized by a gradual loss of muscle mass and strength that leads to sarcopenia as part of the frailty syndrome, a pathology that constitutes a serious health problem associated with adverse outcomes in elderly individuals [1,2]. This age-related decline in muscle function is associated with a reduction in numbers and regenerative potential of muscle stem cells (MuSCs) by impacting on several cell-intrinsic signaling pathways [3-8]. MuSCs also interact with their local microenvironment, the muscle stem cell niche. These interactions play a critical role in regulating MuSCs behavior and affect the outcome of regeneration, which in the context of aging has been linked to fibro/fatty degeneration of muscle tissue [4,9-11].

A key component of the muscle stem cell niche is the extracellular matrix (ECM) which experiences extensive remodeling during regeneration and throughout life [9,10,12,13]. Fibro/adipogenic progenitor cells (FAPs) are considered the main ECM-source, they regulate its remodeling and are indispensable for long-term muscle maintenance and MuSCs' functionality during regeneration [14,15]. However, due to surgical interventions or in general under pathological conditions, such as muscular dystrophies, obesity, and type 2 diabetes mellitus (T2DM), subsets of FAPs are thought to generate fibrotic scar tissue and/or differentiate into adipocytes that progressively replace functional muscle tissue, resulting in fibro/fatty degeneration [16-21]. Besides their role in ECM remodeling, FAPs secrete trophic factors that control MuSC activation through different mechanisms [22-26] and aged FAPs display an aberrant secretory profile of WISP1, BMP3b or SMOC2, among others, that reduce MuSCs' regeneration capacity [11,12,14]. Conversely, the contribution of aging to FAPs' dysregulation and muscle fibro/fatty infiltration remains less well understood. In addition to the FAP-MuSCs niche crosstalk, FAPs interact with different immune cells [27-30]. For instance, pro-inflammatory macrophages regulate the number of FAPs via TNF-mediated apoptosis [30]. A failure to resolve the initial pro-inflammatory macrophage response, however, reduces myogenic regeneration and promotes chronic inflammation in aged muscles [31-34].

Periostin (*Postn*) is a matricellular protein that interacts with ECM components to drive fibrillogenesis and matrix scaffold formation. It also displays signaling ability by binding integrin receptors, mainly αvβ3 and αvβ5 heterodimers, to promote cell adhesion, migration and proliferation in different tissues and in the context of regeneration [35-37]. Expression of *Postn* is transiently induced in FAPs during muscle regeneration and initial data suggest a role in regeneration by coupling myogenesis and angiogenesis [38-40]. Omics-based analyses also show *Postn* among the potentially dysregulated genes in aged muscles [12,25]. In addition to its tissue/stem cell-specific function, Periostin is discussed as a circulating biomarker of type-2 inflammation in asthma, breast cancer, cardiovascular disease and liver steatosis [41-44], as well as a potential biomarker of aging in serum and across several murine tissues [25,45].

We here investigate the role of age-related changes of the ECM as a niche component in orchestrating stem cell-based muscle regeneration. Loss of Periostin phenocopies several aspects of muscle aging, including MuSC dysfunction, delayed inflammation resolution, and fibro/fatty degeneration after injury. Reconstitution of Periostin through FAP-transplantation recovers young-like regeneration in aged muscle. Mechanistically, Periostin targets mechanosensory pathways in myogenic niche-resident macrophages, thereby promoting resolution of the inflammatory microenvironment during regeneration. Periostin may serve as sarcopenia biomarker as circulating levels positively correlate with muscle mass and function at different ages, but are reduced with age in humans.

Results

Loss of Periostin in aged muscle disrupts extracellular matrix function and the stem cell niche

To examine the contribution of aging to myogenic niche disruption, we compared young, 2-months-old male and female mice to sex-matched animals aged 15-18 months. This age was chosen to reduce the likelihood of other, potentially confounding, pathologies that may occur in even older mice. At 15 months, muscles display a gene expression profile and histology consistent with fibro/fatty degeneration, suggesting such changes are relatively early hallmarks of muscle aging (Suppl. Figure S1A-S1C). Fluorescence-activated cell sorting (FACS) showed increased numbers of muscle-resident FAPs with age while MuSCs decreased in male and female mice (Figures 1A, 1B and S1D-S1H). Primary FAPs isolated from muscle of 15-month-old mice displayed spontaneous lipid droplet accumulation, while aged MuSCs showed reduced myogenic potential compared to respective young-muscle cell cultures (Figures 1C and 1D). Given the marked increase in adipogenic potential in aged FAPs, we decided to further examine this cell population. Microarray analysis of freshly sorted FAPs showed a marked reduction in many genes and pathways linked to the ECM (Table S1, Figure S1I). The top down-regulated gene, *Postn*, which encodes for Periostin, is a secreted matricellular protein [37] and could thus act as paracrine niche factor (Figure 1E). Aging muscles displayed reduced Periostin expression on protein and mRNA levels in male and female mice, and cultured young FAPs readily released Periostin into the supernatant, while aged FAPs only secreted minor amounts (Figures 1F-1H, S1J and S1K).

We next tested whether muscle injury would elicit a similar phenotype and applied an established model of tibialis anterior (TA) muscle injury by intramuscular injection of glycerol, a protocol known to induce ectopic adipogenesis and fibrosis at the injury site [46]. The expected injury-induced expansion of FAPs was more pronounced in aged TA compared to young post-injury muscle in male and female mice. Unexpectedly, MuSC expansion was also induced in aged muscle of female mice and to some extent in male mice Figures 1I, 1J and S1L-S1N). Clearance of FAPs is critical to prevent fibro/fatty infiltration [30], and accordingly we detected increased

adipogenesis and fibrosis areas in injured TA of aged compared to young male mice which was accompanied by

an age-related reduction in myofiber cross-sectional areas (CSA) at the injury site and reduced expression of

myogenesis markers (Figures 1K-1M and S1O-S1Q). To further explore the involvement of Periostin in muscle

healing, we evaluated its expression profile on multiple days post-injury (dpi). In line with other publications [38-

40,47], we observed an increase of Periostin in the interstitial space between regenerating fibers and corresponding

dynamic mRNA and protein expression patterns that peaked between 5- and 8-dpi in muscle of young mice, all of

which was markedly attenuated in aged muscle (Figures 1N-1P). We verified disrupted FAP-functions in

regenerating aged muscles by applying single cell-RNA sequencing (scRNA-seq) of FAPs isolated from 4-dpi

muscle. Uniform manifold approximation and projection (UMAP) visualization and differentially expressed gene

(DEG) analysis revealed partial transcriptional separation of FAPs of the two age groups (Figure 1Q; Table S2).

This analysis confirmed *Postn* as one of the most significantly down-regulated genes and the enrichment of ECM-

related pathways when assessing all significantly downregulated genes (Figures 1R-1T). Up-regulated pathways

and a subsequent analysis of predicted secretome genes showed a potential impact of FAP-aging on the modulation

of inflammatory responses (Figures S1R and S1S). These findings taken together indicate that aging negatively

influences niche functions of FAPs which contribute to fibro/fatty degeneration of aged muscles.

Genetic inactivation of *Postn* recapitulates an aging-like phenotype in muscle regeneration

To determine whether loss of Periostin could indeed contribute to a decline of muscle regeneration as observed in

aged mice, we obtained a mouse model with genetic inactivation of the *Postn* gene by insertion of a LacZ cassette,

leading to loss of the *Postn* gene product and expression of a β-galactosidase reporter under control of the native

Postn promoter (Figures S2A-S2D). Periostin protein was completely absent in muscles of *Postn$^{-/-}$* mice whereas

the reporter was expressed in the interstitial space between myofibers, consistent with the localization of FAPs

(Figure S2B). No differences in myofiber CSAs or frequencies of FAPs and MuSCs were observed in 2-3 months-

old muscles of male or female *Postn$^{-/-}$* mice without injury. However, we noticed that dissected muscles displayed

a significant reduction in mechanical stiffness (Figures S2E to S2G). Similar to aged muscle, isolated *Postn$^{-/-}$*-

FAPs displayed significantly enhanced spontaneous adipogenesis in culture whereas cultured MuSCs had

significantly lower myogenic potential (Figures 2A and 2B). Applying the glycerol-injection regeneration model

to *Postn$^{-/-}$* mice and littermate wildtype controls using both male and female offspring showed a phenotype closely

resembling the phenotype seen in aged muscle: We consistently observed increased expansion of FAPs and MuSCs

in *Postn$^{-/-}$* mice (Figure 2C, 2D, S2H and S2I). Importantly, both sexes showed impaired regeneration in *Postn$^{-/-}$*

mice with significant accumulation of adipocytes and fibrotic structures and reduced CSA of newly regenerated

myofibers at 14-dpi, which was also reflected by increased expression of adipocyte and fibrosis genes and reduced

expression of myogenesis markers (Figures 2E-2G and S2J-S2Q). In summary, genetic ablation of *Postn*

recapitulates an aging-like disruption of regeneration in male and female mice at the level of FAPs and MuSCs,

resulting in fibro/fatty degeneration.

Periostin rescues stem cell function and muscle regeneration in aged mice

To better understand the function of Periostin as ECM-related niche factor, we tested the impact of a mouse

recombinant Periostin (mrPOSTN) on FACS-purified FAPs and MuSCs of aged and *Postn*-deficient mice. Both

cell types, aged and $Postn^{-/-}$-FAPs, displayed significantly increased proliferation when compared to the respective

control FAPs, whereas MuSCs displayed reduced proliferation rates in comparison to respective controls (Figures

3A-3D, S3A and S3B). After 48h of culture in the presence of mrPOSTN the respective FAP- and MuSC-

proliferation phenotypes were fully reversed in $Postn^{-/-}$ and aged cells. Considering that FAPs are likely a main

source of muscle Periostin, these data suggest that FAPs auto-inhibit their own expansion while stimulating cell

cycle in MuSCs through paracrine Periostin (Figures 1H, 3A-3D). To determine whether FAP-derived Periostin

influences the function of MuSCs, we used an in vitro co-culture system. Consistent with our initial observation,

lack of Periostin in MuSCs significantly reduced fusion capacity and resulted in shorter myofibers during

differentiation (Figures 3E-3G). Co-cultures of wildtype, Periostin-secreting FAPs, but not $Postn^{-/-}$-FAPs, with

Postn-deficient MuSCs resulted in a full recovery of the impaired myogenesis phenotype (Figures 3E-3G). As

cultured MuSCs themselves secrete very little Periostin in comparison to FAPs, this interaction appears to be a

key element in maintaining the in vitro myogenic potential of the myogenic stem cell pool (Figure S3C).

To determine whether Periostin reconstitution has the capacity to enhance the regeneration process in aged skeletal

muscle, we sorted FAPs from a mouse model carrying a red fluorescent protein (RFP) expression allele in addition

to the *Postn*-knockout allele (Figures S3D and S3E) [48]. RFP[+] FAPs either with (FAP[RFP]-WT) or without (FAP[RFP]-

KO) *Postn* expression were transplanted into TA muscle of 15-months-old recipient mice one day after glycerol-

injury. Transplanted FAPs remained detectable based on RFP expression until at least 8-dpi (Figures 3H, S3D-

S3F). Immunofluorescence analysis at 8-dpi showed that FAP[RFP]-WT transplants significantly reduced the number

of embryonic myosin heavy chain (eMYHC) positive cells and increased CSAs in regenerating fibers compared

to aged recipients transplanted with FAP[RFP]-KO (Figures 3I-3K). Consistently, recipients transplanted with

FAP[RFP]-WT maintained significantly increased CSAs at 14-dpi when compared to FAP[REF]-KO transplants

(Figures 3L and 3M). Finally, we compared the CSAs average and fiber size range results at 14-dpi to the original

experiments without FAP-transplantation (i.e. CSA-means and fiber size range of data reported in Figure 1M).

Aged mice receiving FAP^{RFP}-KO transplants displayed significantly lower average CSA compared to young

control mice and were at a similar level compared to aged TA without FAP transplant, whereas Periostin-producing

FAP^{RFP}-WT transplants significantly increased regenerating myofibers' CSA compared to aged mice without

transplanted FAPs or with FAP^{RFP}-KO transplants (Figure 3N and S3G). Thus, reinstating Periostin as a niche

factor ameliorates the age-related defect of myogenic regeneration, indicating that Periostin is a critical ECM-

protein to orchestrate muscle regeneration.

Periostin mediates immunomodulatory functions of the myogenic stem cell niche

Since regeneration is orchestrated by the interplay of multiple non-myofiber cell populations [49], we performed

droplet-based scRNA-seq of FACS-purified non-myofiber cells from 2.5- and 15-month-old female mice at 4-dpi,

around the time when FAPs frequency peaks during muscle injury repair (Figures S4A and S4B). UMAP

visualization, unsupervised Louvain-clustering and subsequent cell type annotation based on gene scoring of single

cell transcriptomes resulted in the identification of distinct clusters representing the main nine, non-myofiber cell

populations involved in muscle regeneration (Figures 4A, 4B, S4C and S4D; Tables S3 and S4) [50]. Analysis of the

cell-type proportions between age groups showed increased relative abundances of myeloid subsets, i.e. neutrophil

(NP) and macrophage (MC) subsets in aged muscle, while lymphoid subpopulations, i.e. T-lymphocytes (TC) and

natural killer cells (NK), showed lower relative frequencies (Figure 4C; Table S5). Considerable differences in

gene expression patterns were observed among multiple cell types and when comparing the 2.5- and 15-months-

old samples. For instance, analysis of enriched term annotations of downregulated DEGs from the FAP subset

confirmed that aging negatively influences ECM regulation (Figure S4E, Table S6). As before, *Postn* was mostly

expressed in FAPs and its expression was significantly reduced in aged FAPs (Figures 4D, 4E, and S4D; Table

S4). To examine cell-cell interactions (CCI) between Periostin-secreting FAPs and other niche cells, we took

advantage of a previously published murine atlas of age-related changes in intercellular communications, which

uses an algorithm to infer CCI from published gene expression databases [51]. When selecting Periostin as a FAP-

emitted ligand, high CCI-scores were identified for a potential autocrine FAP-FAP interaction and for paracrine

FAP-endothelial cell (EC), FAP-MuSC and FAP-MC interactions mediated through different Integrin receptors

(*Itga5*, *Itgβ1*, and heterodimers *Itgαv/β5*, *Itgαv/β3*) and the Epidermal growth factor receptor (*Egfr*; Figure 4F).

Using aging as a regulatory factor of these interactions, statistical significance levels for each interaction showed

a significant down-regulation of Periostin-based FAP-FAP, FAP-EC, FAP-MuSC and FAP-MC interactions, thus

verifying our own experimental data on the auto- and paracrine roles of Periostin (Figure 4G). The age-dependent

down-regulation of the interaction between FAPs and MCs through the predicted CCI between Periostin and the

Itgαv/β5 heterodimer raised our interest as this was consistent with the alterations in immune subsets in our

scRNA-seq data and offered the possibility of an immunomodulatory niche effect through the FAP-based release

of Periostin (Figure 4G). We validated subset-specific expression patterns of all potential Periostin receptors in

our scRNA-seq data and found that the proposed Itgαv/β5 heterodimer showed an expression pattern that would

most closely reflect the effects of Periostin on the cell types that we also detected in our experimental analyses,

i.e. FAPs, MuSCs, and MCs (Figure 4H).

Loss of Periostin impairs FAP-based immunomodulation by attenuating anti-inflammatory macrophages

Corroborating the FAP-MC interaction through Periostin, FACS-analyses of MCs populations confirmed the

increase of MCs upon muscle injury in aged and also female *Postn*$^{-/-}$ mice compared to the respective control

animals, while only trends or no differences were observed in male *Postn*$^{-/-}$ mice (Figures S5A-S5E). Next, we

examined MC subsets representing functional changes from a pro-inflammatory (M1-like) profile to an anti-

inflammatory (M2-like) phenotype during the course of muscle healing [27,52]. As expected, injury led to a rapid

increase of CD45$^+$CD11b$^+$F4/80$^+$ MCs, which over time shifted from a predominantly pro-inflammatory Ly6C$^+$

population, which peaked at 1- and 3-dpi, to an anti-inflammatory Ly6C$^-$ population which peaked at 4- and 8-dpi

(Figure 5A). Importantly, aging as well as genetic ablation of *Postn* led to an attenuation of this shift in male and

female mice, both of which resulted in significantly elevated levels of pro-inflammatory MCs at earlier time points

and significantly lower levels of anti-inflammatory MCs at the later time points (Figures 5A-5C, S5F and S5G).

We confirmed this delay in the formation of an anti-inflammatory, pro-regenerative MC profile by assessing

CD206, a surface marker commonly found on anti-inflammatory MCs [53-55]. Aging and the deletion of *Postn*

significantly reduced the accumulation of CD45$^+$CD11b$^+$F4/80$^+$CD206$^+$ MCs in the late, regenerative phase at 3-,

4- and 8-dpi, regardless of gender (Figures 5D-5F, S5H and S5I). Lastly, we examined MCs in our FAP

transplantation model (Figure 3H) and detected no differences in overall MCs but a similar phenotype of

significantly elevated CD206$^+$ MCs in 8-dpi-injury sites of aged mice transplanted with *Postn*-expressing FAPs

compared to *Postn*-deficient transplants (Figures 5G and 5H). These data reiterate the notion that loss of Periostin,

similar to aging, delays the formation of a regeneration-supporting anti-inflammatory niche.

Periostin promotes anti-inflammatory macrophages through integrin-FAK-AKT mechano-signal

transduction

To investigate the mechanism of Periostin-dependent control of macrophage polarization, we co-cultured wildtype

and *Postn*$^{-/-}$-FAPs with bone marrow-derived macrophages (BMDM) in a trans-well system and compared them

to known pro- and anti-inflammatory stimuli. After 48h, FAPS-WT (*Postn^+/+^*), but not FAPs-KO (*Postn^-/-^*),

significantly induced expression of anti-inflammatory MC markers *Mrc1* and *Cd163* while other anti-

inflammatory genes, *Il10*, and several pro-inflammatory markers were not affected by the presence of FAPs

(Figures 6A and S6A). Addition of species-specific recombinant Periostin proteins (mrPOSTN, hrPOSTN) to

murine BMDM- and human, THP-1 cell line-derived macrophages resulted in a significant, dose-dependent

increase in the expression of anti-inflammatory markers *Mrc1*, *Cd163* and *Il10*, but did not affect pro-inflammatory

gene expression (Figures 6B, S6B and S6D). Integrins, and specifically Itgαv/β3, Itgαv/β5 heterodimers, have

been linked to anti-inflammatory macrophage function [56,57] and our own findings and published data indicate that

Periostin as able to bind both integrin heterodimers [35,36]. Treatment of BMDM and THP-1 macrophages with

Cilengitide, an inhibitor of ITGαv/β3 and ITGαv/β5 heterodimers [58,59], resulted in a significant reduction of

rPOSTN-induced expression of anti-inflammatory surface protein CD206 and similar effects on *Mrc1*, which

encodes for CD206, and *Cd163* mRNA expression, whereas recombinant IL4-induced expression of these genes

was not affected by Cilengitide and little or no effect on other pro-inflammatory genes was evident (Figures 6C-

6E, S6C, S6E and S6F). Protein kinase B (AKT) and focal adhesion kinase (FAK) phosphorylation have been

linked to anti-inflammatory macrophage polarization [60,61] and both kinases are potential downstream targets of the

Periostin-ITGαvβ3 and -αvβ5 axis [35,36,62]. Accordingly, rPOSTN acutely induced the phosphorylation of FAK at

Tyr397 and of AKT at Ser437, which was reduced significantly after Cilengitide addition in BMDM and THP-1

macrophages (Figures 6F, 6G, S6G and S6H). Collectively, these results indicate that rPOSTN induces M2-like

polarization through integrin-dependent FAK-AKT signaling in macrophages.

Aging and fibrotic scar tissue formation during muscle regeneration have been associated with altered muscle

stiffness, somewhat resembling the reduced muscle stiffness we observed in our *Postn^-/-^* mice (Figure S2F) [13,26].

During muscle regeneration, a transient decrease in tissue stiffness at the injury site around 5-dpi, which

subsequently rebounds to normal stiffness levels in fully regenerated muscle, has been reported recently [63]. To

investigate the relationship of Periostin and the mechanical properties of the local niche, and their impact on

macrophage polarization, BMDM were encapsulated in a three-dimensional ECM microenvironment using

different alginate-based hydrogels with either low or high mechanical stiffness. Consistent with previous reports

[64], low stiffness hydrogels contained significantly fewer F4/80^+^/CD206^+^ anti-inflammatory MCs compared to high

stiffness hydrogels, while the opposite was true for F4/80^+^/CD86^+^ pro-inflammatory MCs, which were more

abundant under low stiffness and practically absent under high stiffness conditions. These data suggest that

mechanical input from the microenvironment directly affects macrophage polarization and imply that loss of *Postn*

may exert a similar effect in the muscle niche. Importantly, the addition of murine rPOSTN to the hydrogels

resulted in a Cilengitide-sensitive, modulatory effect where pro-inflammatory polarization induced by low

stiffness was significantly reduced. Instead, an anti-inflammatory MC profile was promoted under conditions of

low stiffness, suggesting that Periostin acts as a niche factor that synchronizes biomechanical and

immunomodulatory processes within the muscle niche (Figures 6H, 6I and S7A-S7C).

Circulating Periostin may predict muscle dysfunction in geriatric patients

Circulating Periostin has been proposed as biomarker of several inflammatory, chronic and metabolic diseases [41-

44], but its role in relation to aging in mice and humans is only now emerging [25,45]. To explore the link between

aging and muscle dysfunction, we measured circulating human Periostin (hPOSTN) in serum samples from two

separate human study cohorts [65,66]. When comparing serum Periostin from young adults to healthy older,

community-dwelling individuals, significantly lower levels of circulating hPOSTN were observed in aged

individuals. Additionally, we found that increased physical activity in elderly individuals restored hPOSTN-

concentrations to levels found in young adults (Figure 7A). As previous studies had reported elevated levels of

circulating Periostin due to certain pathologies [67], we also assessed serum hPOSTN in a group of healthy older

individuals compared to age-matched individuals living in an acute geriatric hospital setting. No differences were

found, indicating that circulating hPOSTN is not applicable as a generalized marker of impaired health status

(Figure 7B). Next, we performed correlations including young adults to healthy older community-dwelling

individuals only (from panel 7A), and we found an inverse correlation between age and circulating hPOSTN serum

levels in women, but not in men, which could be due to lower inclusion numbers for men (Figure 7C). Using all

individuals of this study cohort, we subsequently examined whether other metabolic and functional muscle

parameters would also correlate with circulating hPOSTN. Indeed, hPOSTN correlated inversely with body mass

index (BMI) and its serum concentration was positively corelated with skeletal muscle mass and grip strength in

women, but not men (Figures 7D-7F). These data suggest that circulating Periostin levels could be used as a

potential biomarker of skeletal muscle aging and the age-related sarcopenia syndrome.

Discussion

Muscle injury repair requires complex interactions between different cell types residing within the stem cell niche

[49]. For instance, extensive evidence exists for FAP-MuSC interactions to ensure successful myogenic regeneration

[11,14,15,22,24,26,68]. Aging and other pathologies result in defective myogenesis due to stem cell niche dysfunction by

inhibiting different aspects of myogenic stem cell maintenance, activation and differentiation [4]. We here report on

a niche-dependent mechanism that integrates paracrine signals and mechano-signal transduction to impact on the

levels of MuSCs and the myogenesis-regulatory FAPs, and by exerting immunomodulatory effects regulating the

inflammatory profile of injury site-associated macrophages. De-synchronization of these processes creates a

defective stem cell niche that phenocopies the detrimental effects of aging on myogenic regeneration.

Firstly, we show that loss of the matricellular protein, Periostin, which is secreted by muscle-resident FAPs,

directly impinges on MuSCs, attenuating their ability to expand in response to injury and to differentiate into fully

functional myofibers. These findings are consistent with recent studies showing that aging alters secretion of

signaling molecules like the WNT1-inducible signaling pathway protein 1 (WISP1), bone morphogenetic protein

3 (BMP3b) and SPARC-related modular calcium-binding protein 2 (SMOC2) by FAPs in the stem cell niche,

thereby reducing MuSCs' regeneration capacity [11,12,69].

Previous reports have established that aging and chronic disease prime FAPs to form adipocytes that accumulate

between muscle fibers, leading to an impairment of muscle function [2,16,17]. In line with these observations, our

animal models with loss of Periostin due to age or genetic targeting display defective autoregulation of FAPs,

resulting in excessive FAP-expansion upon injury which feature an elevated commitment towards the adipogenic

lineage at the expense of their ability to maintain appropriate myogenic regeneration capacity. This process results

in the formation of a pro-fibro/adipogenic microenvironment with enhanced fibro/fatty infiltration instead of

successful muscle repair. A tight regulation of FAP expansion and clearance is essential to maintain tissue integrity

and prevent fibro/fatty degeneration. However, the mechanisms through which aging may affect FAPs' expansive

capacities in homeostatic and regenerating muscle tissues remain unclear. Previous reports have described a

reduction of FAPs in uninjured muscle during aging [11,14]. Upon injury the situation could depend on the specifics

of the experimental layout, physiological status and age, as age-related failure of FAPs to expand has been reported

[11], while our own observations suggest enhanced FAP expansion, which align with scRNA-seq and tissue-array

data at steady-state in old muscle [9,70]. Intramuscular FAP transplants into mice with impaired myogenesis have

been shown to rescue the normal characteristics of healthy muscle under homeostatic conditions and improved

regeneration capacity [11,23,24]. Our FAP/MuSC co-cultures and FAPs-transplants into muscle injuries of aged mice

help restore secretion of Periostin and are sufficient to reconstitute myogenesis and muscle regeneration. Our

findings therefore support the existing concept that FAP-secreted ECM factors are necessary and sufficient to

sustain myogenesis via the FAP-MuSC crosstalk. A less well-studied aspect of our work sheds additional light is

the concept of ECM-linked factors that exert immunomodulatory effects by targeting the pro- to anti-inflammatory

transition of macrophages within the muscle injury site. This process integrates mechanical properties of the ECM

and its changes during muscle repair with mechanosensory signaling through AKT and FAK pathways. When

disrupted, for instance through los of Periostin, this results in a delay of the pro- to anti-inflammatory switch within

the post-injury myogenic niche and is also observed in aging muscle. The crosstalk between FAPs and muscle-resident immune cell subsets to sustain adequate muscle regeneration has been reported on before [26,28-30]. Nevertheless, the exact mechanisms of interaction between FAPs and macrophage polarization associated with muscle regeneration capacity, and their alterations during the aging process, remain largely unknown. We here introduce the concept that niche factors like Periostin may help to synchronize the biological signaling processes of immunomodulation and biomechanics during muscle regeneration. Previous reports have shown that FAPs coordinate myeloid and lymphoid cell contributions to inflammatory processes during the regeneration process [26,28-30]. Our scRNA-seq analysis of regeneration in young and aged muscle shows that myeloid and lymphoid subsets are the main clusters affected by aging. The reduction of T cells in aged muscles has been linked to reduced secretion of IL33 by FAPs during the regeneration process, and a limited influence of NK cells on myogenic regeneration through IFNγ has been reported [29,71]. In line with our own observations, experimental reconstitution of macrophage-derived signaling molecules growth differentiation factor 3 (GDF3) and mesencephalic astrocyte-derived neurotrophic factor (MANF) improved regeneration [33,72]. These results also support the concept of reinstating a normal macrophage function to ensure successful regeneration in the context of aging. This could be achieved by targeting regulatory FAPs to provide appropriate immunomodulatory signals, such as Periostin, to orchestrate the pro- to anti-inflammatory macrophage transition. In our hands, FAPs transplantation with functional Periostin expression into aged, post-injury muscle restored a timely onset of anti-inflammatory niche formation and successful regeneration. These findings are in line with previous results showing that Periostin could support a shift from a pro- towards an anti-inflammatory phenotype to sustain heart, kidney and bladder tissue regeneration, although a molecular mechanism has so far not been reported [53-55]. Thus, our data link FAPs-derived Periostin secretion to the muscle stem cell niche with synchronizing effects on FAPs, MuSCs and macrophages during muscle healing. Periostin exerts its function through integrin receptors and mainly through αvβ3 and αvβ5 heterodimers in different biological contexts [35-37]. Analyses of CCIs in the muscle stem cell niche corroborate that Periostin is a FAPs-secreted ligand that mediates intercellular communication between FAPs, MuSCs and macrophages. Interestingly, the Periostin:Itgαv/β5 interaction, which mediates the crosstalk between FAPs and macrophages, declines with aging in the model, thus mirroring our own experimental evidence and presenting a new mechanism of age-related macrophage dysfunction and niche deterioration where a relative lack of Periostin fails to activate FAK-AKT signaling. This in turn attenuates the transition from a pro- towards an anti-inflammatory milieu in the regenerating muscle niche and is in line with other studies that implicate Periostin in regulation of macrophage function in other organs and physiological contexts [36,53,61]. Corresponding immunomodulatory mechanisms have been described where the integrin-αvβ5 heterodimer regulates anti-

inflammatory macrophages via a PPAR-γ dependent mechanism [57], suggesting that this signaling pathway may

also integrate different immunomodulatory signals during myogenic regeneration.

In addition to its role as a signaling molecule, our data also suggest a link between Periostin and mechanosensory

signal transduction. The ability of matrices with different mechanical properties to regulate macrophage

polarization is well-documented in the literature [64]. Molecules like Periostin could offer an additional layer of

regulatory input that offsets or enhances the role of the local ECMs' mechanical profile on inflammatory processes,

including macrophage functions, thereby also impacting on myogenic stem cells. Aging increases muscle stiffness

[13], a mechanical property which would normally promote anti-inflammatory macrophages. It should be noted that

such a situation would only occur in aged muscle at steady-state, i.e. non-injury conditions, and has indeed been

reported [73]. In intact muscle, however, overall numbers of tissue macrophages are comparably low. Conversely,

post-injury muscles experience dynamic ECM-remodeling, marked by an initial, but transient decline of stiffness

at 5-dpi [63], which in fact coincides with the pro- to anti-inflammatory niche/ macrophage transition. As low

stiffness would preferentially stimulate pro-inflammatory macrophages, the presence of Periostin could mediate

or at last accelerate the transition to an anti-inflammatory microenvironment.

Lastly, circulating Periostin has been proposed as a potential biomarker in some instances such as for

cardiovascular risk, early cancer detection, or different type 2 inflammatory diseases [41-44]. Our own and other

previous work suggests that reduced gene expression levels of *Postn* are reliable predictors of aging and associated

metabolic disorders [25,45]. Thus, it stands to reason that loss of circulating Periostin may also inform on the

progression of muscle degeneration as a result of age-related sarcopenia. However, it should be noted that

increased levels of Periostin have also been proposed as a biomarker of disease progression in older adults,

including cognitive decline [67], suggesting that associations such as those found in our cohorts need to be evaluated

in more detail. In conclusion, our study presents a key role of FAP-based Periostin secretion in the muscle stem

cell niche, which links age-related niche deterioration to the modulation of macrophage function and the

regeneration capacity of skeletal muscle. Our results provide mechanistic insights into a crosstalk between FAPs

and macrophages that is impaired during aging, which integrate paracrine niche effects and mechano-sensing.

Acknowledgments

The authors thank A. Kretschmer, S. Marg, S. Richter, A. Weber and N. Dittberner for expert technical assistance.

This work was supported by grants from the German Research Foundation (DFG) as individual grants and within

the Collaborative Research Centre CRC1444 (project IDs 323196138, 249509554, and 427826188), and by a grant

from the Leibniz Association (ID K398/2021, to TJS). Grants within the German Center for Diabetes Research

(DZD) funded by the German Ministry of Education and Research (BMBF) and the State of Brandenburg (DZD

grant IDs 82DZD00302, 82DZD03D03 and 82DZD03C3G) are acknowledged.

Author Contributions

F.G.-C. and T.J.S. conceived the study and wrote the manuscript. F.G.-C. conducted the majority of experiments.

S.G., G.L.-G., A.-M.J., C.G., S.P.-K., A.T., A.G., G.K. and M.Os., contributed research data to this study. G.S.,

L.V. und V.B. performed scRNA-seq experiments. M.L. and M.Ov. performed computational analyses of scRNA-

seq data. C.H., U.M.-W., K.N., provided human serum samples and helped with data analysis. S.K., A.S., S.S.,

K.S.-B., A.C., R.S.K., and G.N.D. contributed valuable materials and expertise to the article.

Declaration of interests

The authors declare no conflict of interest.

References

1. Goodpaster, B.H., Park, S.W., Harris, T.B., Kritchevsky, S.B., Nevitt, M., Schwartz, A.V., Simonsick, E.M.,

Tylavsky, F.A., Visser, M., and Newman, A.B. (2006). The loss of skeletal muscle strength, mass, and quality

in older adults: the health, aging and body composition study. J Gerontol A Biol Sci Med Sci *61*, 1059-1064.

10.1093/gerona/61.10.1059.

2. Marcus, R.L., Addison, O., Kidde, J.P., Dibble, L.E., and Lastayo, P.C. (2010). Skeletal muscle fat infiltration:

impact of age, inactivity, and exercise. J Nutr Health Aging *14*, 362-366. 10.1007/s12603-010-0081-2.

3. Brack, A.S., Bildsoe, H., and Hughes, S.M. (2005). Evidence that satellite cell decrement contributes to

preferential decline in nuclear number from large fibres during murine age-related muscle atrophy. J Cell Sci

118, 4813-4821. 10.1242/jcs.02602.

4. Chakkalakal, J.V., Jones, K.M., Basson, M.A., and Brack, A.S. (2012). The aged niche disrupts muscle stem

cell quiescence. Nature *490*, 355-360. 10.1038/nature11438.

5. Conboy, I.M., Conboy, M.J., Wagers, A.J., Girma, E.R., Weissman, I.L., and Rando, T.A. (2005).

Rejuvenation of aged progenitor cells by exposure to a young systemic environment. Nature *433*, 760-764.

10.1038/nature03260.

6. Cosgrove, B.D., Gilbert, P.M., Porpiglia, E., Mourkioti, F., Lee, S.P., Corbel, S.Y., Llewellyn, M.E., Delp,

S.L., and Blau, H.M. (2014). Rejuvenation of the muscle stem cell population restores strength to injured aged

muscles. Nat Med *20*, 255-264. 10.1038/nm.3464.

7. Garcia-Prat, L., Martinez-Vicente, M., Perdiguero, E., Ortet, L., Rodriguez-Ubreva, J., Rebollo, E., Ruiz-

Bonilla, V., Gutarra, S., Ballestar, E., Serrano, A.L., et al. (2016). Autophagy maintains stemness by preventing

senescence. Nature *529*, 37-42. 10.1038/nature16187.

8. Sousa-Victor, P., Gutarra, S., Garcia-Prat, L., Rodriguez-Ubreva, J., Ortet, L., Ruiz-Bonilla, V., Jardi, M.,

Ballestar, E., Gonzalez, S., Serrano, A.L., et al. (2014). Geriatric muscle stem cells switch reversible quiescence

into senescence. Nature *506*, 316-321. 10.1038/nature13013.

9. Lazure, F., Farouni, R., Sahinyan, K., Blackburn, D.M., Hernandez-Corchado, A., Perron, G., Lu, T., Osakwe,

A., Ragoussis, J., Crist, C., et al. (2023). Transcriptional reprogramming of skeletal muscle stem cells by the

niche environment. Nat Commun *14*, 535. 10.1038/s41467-023-36265-x.

10. Lukjanenko, L., Jung, M.J., Hegde, N., Perruisseau-Carrier, C., Migliavacca, E., Rozo, M., Karaz, S., Jacot,

G., Schmidt, M., Li, L., et al. (2016). Loss of fibronectin from the aged stem cell niche affects the regenerative

capacity of skeletal muscle in mice. Nat Med *22*, 897-905. 10.1038/nm.4126.

11. Lukjanenko, L., Karaz, S., Stuelsatz, P., Gurriaran-Rodriguez, U., Michaud, J., Dammone, G., Sizzano, F.,

Mashinchian, O., Ancel, S., Migliavacca, E., et al. (2019). Aging Disrupts Muscle Stem Cell Function by

Impairing Matricellular WISP1 Secretion from Fibro-Adipogenic Progenitors. Cell Stem Cell *24*, 433-446

e437. 10.1016/j.stem.2018.12.014.

12. Schuler, S.C., Kirkpatrick, J.M., Schmidt, M., Santinha, D., Koch, P., Di Sanzo, S., Cirri, E., Hemberg, M.,

Ori, A., and von Maltzahn, J. (2021). Extensive remodeling of the extracellular matrix during aging contributes

to age-dependent impairments of muscle stem cell functionality. Cell Rep *35*, 109223.

10.1016/j.celrep.2021.109223.

13. Stearns-Reider, K.M., D'Amore, A., Beezhold, K., Rothrauff, B., Cavalli, L., Wagner, W.R., Vorp, D.A.,

Tsamis, A., Shinde, S., Zhang, C., et al. (2017). Aging of the skeletal muscle extracellular matrix drives a stem

cell fibrogenic conversion. Aging Cell *16*, 518-528. 10.1111/acel.12578.

14. Uezumi, A., Ikemoto-Uezumi, M., Zhou, H., Kurosawa, T., Yoshimoto, Y., Nakatani, M., Hitachi, K.,

Yamaguchi, H., Wakatsuki, S., Araki, T., et al. (2021). Mesenchymal Bmp3b expression maintains skeletal

muscle integrity and decreases in age-related sarcopenia. J Clin Invest *131*. 10.1172/JCI139617.

15. Woszczyna, M.N., Konishi, C.T., Perez Carbajal, E.E., Wang, T.T., Walsh, R.A., Gan, Q., Wagner, M.W., and

Rando, T.A. (2019). Mesenchymal Stromal Cells Are Required for Regeneration and Homeostatic

Maintenance of Skeletal Muscle. Cell Rep *27*, 2029-2035 e2025. 10.1016/j.celrep.2019.04.074.

16. Farup, J., Just, J., de Paoli, F., Lin, L., Jensen, J.B., Billeskov, T., Roman, I.S., Comert, C., Moller, A.B.,

Madaro, L., et al. (2021). Human skeletal muscle CD90(+) fibro-adipogenic progenitors are associated with

muscle degeneration in type 2 diabetic patients. Cell Metab *33*, 2201-2214 e2211. 10.1016/j.cmet.2021.10.001.

17. Hogarth, M.W., Defour, A., Lazarski, C., Gallardo, E., Diaz Manera, J., Partridge, T.A., Nagaraju, K., and

Jaiswal, J.K. (2019). Fibroadipogenic progenitors are responsible for muscle loss in limb girdle muscular

dystrophy 2B. Nat Commun *10*, 2430. 10.1038/s41467-019-10438-z.

18. Kopinke, D., Roberson, E.C., and Reiter, J.F. (2017). Ciliary Hedgehog Signaling Restricts Injury-Induced

Adipogenesis. Cell *170*, 340-351 e312. 10.1016/j.cell.2017.06.035.

19. Madaro, L., Passafaro, M., Sala, D., Etxaniz, U., Lugarini, F., Proietti, D., Alfonsi, M.V., Nicoletti, C., Gatto,

S., De Bardi, M., et al. (2018). Denervation-activated STAT3-IL-6 signalling in fibro-adipogenic progenitors

promotes myofibres atrophy and fibrosis. Nat Cell Biol *20*, 917-927. 10.1038/s41556-018-0151-y.

20. Perez-Diaz, S., Koumaiha, Z., Borok, M.J., Aurade, F., Pini, M., Periou, B., Rouault, C., Baba-Amer, Y.,

Clement, K., Derumeaux, G., et al. (2022). Obesity impairs skeletal muscle repair through NID-1 mediated

extracellular   matrix   remodeling   by   mesenchymal   progenitors.   Matrix   Biol   *112*,   90-115.

10.1016/j.matbio.2022.08.006.

21. Uezumi, A., Ito, T., Morikawa, D., Shimizu, N., Yoneda, T., Segawa, M., Yamaguchi, M., Ogawa, R., Matev,

M.M., Miyagoe-Suzuki, Y., et al. (2011). Fibrosis and adipogenesis originate from a common mesenchymal

progenitor in skeletal muscle. J Cell Sci *124*, 3654-3664. 10.1242/jcs.086629.

22. Joe, A.W., Yi, L., Natarajan, A., Le Grand, F., So, L., Wang, J., Rudnicki, M.A., and Rossi, F.M. (2010).

Muscle injury activates resident fibro/adipogenic progenitors that facilitate myogenesis. Nat Cell Biol *12*, 153-

163. 10.1038/ncb2015.

23. Mozzetta, C., Consalvi, S., Saccone, V., Tierney, M., Diamantini, A., Mitchell, K.J., Marazzi, G., Borsellino,

G., Battistini, L., Sassoon, D., et al. (2013). Fibroadipogenic progenitors mediate the ability of HDAC

inhibitors to promote regeneration in dystrophic muscles of young, but not old Mdx mice. EMBO Mol Med *5*,

626-639. 10.1002/emmm.201202096.

24. Uezumi, A., Fukada, S., Yamamoto, N., Takeda, S., and Tsuchida, K. (2010). Mesenchymal progenitors

distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle. Nat Cell Biol *12*, 143-

152. 10.1038/ncb2014.

25. Schaum, N., Lehallier, B., Hahn, O., Palovics, R., Hosseinzadeh, S., Lee, S.E., Sit, R., Lee, D.P., Losada, P.M.,

Zardeneta, M.E., et al. (2020). Ageing hallmarks exhibit organ-specific temporal signatures. Nature *583*, 596-

602. 10.1038/s41586-020-2499-y.

26. Kotsaris, G., Qazi, T.H., Bucher, C.H., Zahid, H., Pohle-Kronawitter, S., Ugorets, V., Jarassier, W., Borno, S.,

Timmermann, B., Giesecke-Thiel, C., et al. (2023). Odd skipped-related 1 controls the pro-regenerative

response of fibro-adipogenic progenitors. NPJ Regen Med *8*, 19. 10.1038/s41536-023-00291-6.

27. Arnold, L., Henry, A., Poron, F., Baba-Amer, Y., van Rooijen, N., Plonquet, A., Gherardi, R.K., and Chazaud,

B. (2007). Inflammatory monocytes recruited after skeletal muscle injury switch into antiinflammatory

macrophages to support myogenesis. J Exp Med *204*, 1057-1069. 10.1084/jem.20070075.

28. Heredia, J.E., Mukundan, L., Chen, F.M., Mueller, A.A., Deo, R.C., Locksley, R.M., Rando, T.A., and Chawla,

A. (2013). Type 2 innate signals stimulate fibro/adipogenic progenitors to facilitate muscle regeneration. Cell

153, 376-388. 10.1016/j.cell.2013.02.053.

29. Kuswanto, W., Burzyn, D., Panduro, M., Wang, K.K., Jang, Y.C., Wagers, A.J., Benoist, C., and Mathis, D.

(2016). Poor Repair of Skeletal Muscle in Aging Mice Reflects a Defect in Local, Interleukin-33-Dependent

Accumulation of Regulatory T Cells. Immunity *44*, 355-367. 10.1016/j.immuni.2016.01.009.

30. Lemos, D.R., Babaeijandaghi, F., Low, M., Chang, C.K., Lee, S.T., Fiore, D., Zhang, R.H., Natarajan, A.,

Nedospasov, S.A., and Rossi, F.M. (2015). Nilotinib reduces muscle fibrosis in chronic muscle injury by

promoting TNF-mediated apoptosis of fibro/adipogenic progenitors. Nat Med *21*, 786-794. 10.1038/nm.3869.

31. Cui, C.Y., Driscoll, R.K., Piao, Y., Chia, C.W., Gorospe, M., and Ferrucci, L. (2019). Skewed macrophage

polarization in aging skeletal muscle. Aging Cell *18*, e13032. 10.1111/acel.13032.

32. Moiseeva, V., Cisneros, A., Sica, V., Deryagin, O., Lai, Y., Jung, S., Andres, E., An, J., Segales, J., Ortet, L.,

et al. (2023). Senescence atlas reveals an aged-like inflamed niche that blunts muscle regeneration. Nature *613*,

169-178. 10.1038/s41586-022-05535-x.

33. Patsalos, A., Simandi, Z., Hays, T.T., Peloquin, M., Hajian, M., Restrepo, I., Coen, P.M., Russell, A.J., and

Nagy, L. (2018). In vivo GDF3 administration abrogates aging related muscle regeneration delay following

acute sterile injury. Aging Cell *17*, e12815. 10.1111/acel.12815.

34. Reidy, P.T., McKenzie, A.I., Mahmassani, Z.S., Petrocelli, J.J., Nelson, D.B., Lindsay, C.C., Gardner, J.E.,

Morrow, V.R., Keefe, A.C., Huffaker, T.B., et al. (2019). Aging impairs mouse skeletal muscle macrophage

polarization and muscle-specific abundance during recovery from disuse. Am J Physiol Endocrinol Metab *317*,

E85-E98. 10.1152/ajpendo.00422.2018.

35. Kuhn, B., del Monte, F., Hajjar, R.J., Chang, Y.S., Lebeche, D., Arab, S., and Keating, M.T. (2007). Periostin

induces proliferation of differentiated cardiomyocytes and promotes cardiac repair. Nat Med *13*, 962-969.

10.1038/nm1619.

36. Li, G., Jin, R., Norris, R.A., Zhang, L., Yu, S., Wu, F., Markwald, R.R., Nanda, A., Conway, S.J., Smyth, S.S.,

and Granger, D.N. (2010). Periostin mediates vascular smooth muscle cell migration through the integrins

alphavbeta3 and alphavbeta5 and focal adhesion kinase (FAK) pathway. Atherosclerosis *208*, 358-365.

10.1016/j.atherosclerosis.2009.07.046.

37. Norris, R.A., Damon, B., Mironov, V., Kasyanov, V., Ramamurthi, A., Moreno-Rodriguez, R., Trusk, T., Potts,

J.D., Goodwin, R.L., Davis, J., et al. (2007). Periostin regulates collagen fibrillogenesis and the biomechanical

properties of connective tissues. J Cell Biochem *101*, 695-711. 10.1002/jcb.21224.

38. Latroche, C., Weiss-Gayet, M., Muller, L., Gitiaux, C., Leblanc, P., Liot, S., Ben-Larbi, S., Abou-Khalil, R.,

Verger, N., Bardot, P., et al. (2017). Coupling between Myogenesis and Angiogenesis during Skeletal Muscle

Regeneration Is Stimulated by Restorative Macrophages. Stem Cell Rep *9*, 2018-2033.

10.1016/j.stemcr.2017.10.027.

39. Ozdemir, C., Akpulat, U., Sharafi, P., Yildiz, Y., Onbasilar, I., and Kocaefe, C. (2014). Periostin is temporally

expressed as an extracellular matrix component in skeletal muscle regeneration and differentiation. Gene *553*,

130-139. 10.1016/j.gene.2014.10.014.

40. Scott, R.W., Arostegui, M., Schweitzer, R., Rossi, F.M.V., and Underhill, T.M. (2019). Hic1 Defines Quiescent

Mesenchymal Progenitor Subpopulations with Distinct Functions and Fates in Skeletal Muscle Regeneration.

Cell Stem Cell *25*, 797-813 e799. 10.1016/j.stem.2019.11.004.

41. Azharuddin, M., Kapur, P., Mishra, R., Saleem, S., Gupta, A.K., Adil, M., and Sharma, M. (2021). Predictor

for cardiovascular risk in patients with type-2 diabetes mellitus. Clin Epidemiol Glob Health *12*.

10.1016/j.cegh.2021.100870.

42. Rachner, T.D., Gobel, A., Hoffmann, O., Erdmann, K., Kasimir-Bauer, S., Breining, D., Kimmig, R., Hofbauer,

L.C., and Bittner, A.K. (2020). High serum levels of periostin are associated with a poor survival in breast

cancer. Breast Cancer Res Treat *180*, 515-524. 10.1007/s10549-020-05570-0.

43. Yang, Z., Zhang, H., Niu, Y., Zhang, W., Zhu, L., Li, X., Lu, S., Fan, J., Li, X., Ning, G., et al. (2016).

Circulating periostin in relation to insulin resistance and nonalcoholic fatty liver disease among overweight

and obese subjects. Sci Rep *6*, 37886. 10.1038/srep37886.

44. Zhao, S., Wu, H., Xia, W., Chen, X., Zhu, S., Zhang, S., Shao, Y., Ma, W., Yang, D., and Zhang, J. (2014).

Periostin expression is upregulated and associated with myocardial fibrosis in human failing hearts. J Cardiol

63, 373-378. 10.1016/j.jjcc.2013.09.013.

45. Graja, A., Garcia-Carrizo, F., Jank, A.M., Gohlke, S., Ambrosi, T.H., Jonas, W., Ussar, S., Kern, M.,

Schurmann, A., Aleksandrova, K., et al. (2018). Loss of periostin occurs in aging adipose tissue of mice and

its genetic ablation impairs adipose tissue lipid metabolism. Aging Cell *17*, e12810. 10.1111/acel.12810.

46. Mahdy, M.A.A. (2018). Glycerol-induced injury as a new model of muscle regeneration. Cell Tissue Res *374*,

233-241. 10.1007/s00441-018-2846-6.

47. Ito, N., Miyagoe-Suzuki, Y., Takeda, S., and Kudo, A. (2021). Periostin Is Required for the Maintenance of

Muscle Fibers during Muscle Regeneration. Int J Mol Sci *22*. 10.3390/ijms22073627.

48. Muzumdar, M.D., Tasic, B., Miyamichi, K., Li, L., and Luo, L. (2007). A global double-fluorescent Cre

reporter mouse. Genesis *45*, 593-605. 10.1002/dvg.20335.

49. Mashinchian, O., Pisconti, A., Le Moal, E., and Bentzinger, C.F. (2018). The Muscle Stem Cell Niche in Health

and Disease. Curr Top Dev Biol *126*, 23-65. 10.1016/bs.ctdb.2017.08.003.

50. De Micheli, A.J., Laurilliard, E.J., Heinke, C.L., Ravichandran, H., Fraczek, P., Soueid-Baumgarten, S., De

Vlaminck, I., Elemento, O., and Cosgrove, B.D. (2020). Single-Cell Analysis of the Muscle Stem Cell

Hierarchy Identifies Heterotypic Communication Signals Involved in Skeletal Muscle Regeneration. Cell Rep

30, 3583-3595 e3585. 10.1016/j.celrep.2020.02.067.

51. Lagger, C., Ursu, E., Equey, A., Avelar, R.A., Pisco, A.O., Tacutu, R., and de Magalhães, J.P. (2021).

scAgeCom: a murine atlas of age-related changes in intercellular communication inferred with the package

scDiffCom. BioRxiv. 10.1101/2021.08.13.456238.

52. Varga, T., Mounier, R., Horvath, A., Cuvellier, S., Dumont, F., Poliska, S., Ardjoune, H., Juban, G., Nagy, L.,

and Chazaud, B. (2016). Highly Dynamic Transcriptional Signature of Distinct Macrophage Subsets during

Sterile Inflammation, Resolution, and Tissue Repair. J Immunol *196*, 4771-4782. 10.4049/jimmunol.1502490.

53. Chen, Z., Liu, L., Chen, Y., Liu, M., Peng Xiang, A., Deng, C., and Jiang, M.H. (2022). Periostin Attenuates

Cyclophosphamide-induced Bladder Injury by Promoting Urothelial Stem Cell Proliferation and Macrophage

Polarization. Stem Cells Transl Med *11*, 659-673. 10.1093/stcltm/szac025.

54. Kormann, R., Kavvadas, P., Placier, S., Vandermeersch, S., Dorison, A., Dussaule, J.C., Chadjichristos, C.E.,

Prakoura, N., and Chatziantoniou, C. (2020). Periostin Promotes Cell Proliferation and Macrophage

Polarization to Drive Repair after AKI. J Am Soc Nephrol *31*, 85-100. 10.1681/ASN.2019020113.

55. Liao, Y., Li, G., Zhang, X., Huang, W., Xie, D., Dai, G., Zhu, S., Lu, D., Zhang, Z., Lin, J., et al. (2020).

Cardiac Nestin(+) Mesenchymal Stromal Cells Enhance Healing of Ischemic Heart through Periostin-Mediated

M2 Macrophage Polarization. Mol Ther *28*, 855-873. 10.1016/j.ymthe.2020.01.011.

56. Antonov, A.S., Kolodgie, F.D., Munn, D.H., and Gerrity, R.G. (2004). Regulation of macrophage foam cell
formation by alphaVbeta3 integrin: potential role in human atherosclerosis. Am J Pathol *165*, 247-258.
10.1016/s0002-9440(10)63293-2.

57. Yao, Q., Liu, J., Zhang, Z., Li, F., Zhang, C., Lai, B., Xiao, L., and Wang, N. (2018). Peroxisome proliferator-
activated receptor gamma (PPARgamma) induces the gene expression of integrin alpha(V)beta(5) to promote
macrophage M2 polarization. J Biol Chem *293*, 16572-16582. 10.1074/jbc.RA118.003161.

58. Lin, X., Sun, Y., Yang, S., Yu, M., Pan, L., Yang, J., Yang, J., Shao, Q., Liu, J., Liu, Y., et al. (2021). Omentin-
1 Modulates Macrophage Function via Integrin Receptors alphavbeta3 and alphavbeta5 and Reverses Plaque
Vulnerability in Animal Models of Atherosclerosis. Front Cardiovasc Med *8*, 757926.
10.3389/fcvm.2021.757926.

59. Zhou, W., Ke, S.Q., Huang, Z., Flavahan, W., Fang, X., Paul, J., Wu, L., Sloan, A.E., McLendon, R.E., Li, X.,
et al. (2015). Periostin secreted by glioblastoma stem cells recruits M2 tumour-associated macrophages and
promotes malignant growth. Nat Cell Biol *17*, 170-182. 10.1038/ncb3090.

60. Po-Chun Chen, H.-C.C., John Wang, Shin-Wei Wang, Huai-Ching Tai, Chiao-Wen Lin, and Chih-Hsin Tang
(2014). Prostate cancer-derived CCN3 induces M2 macrophage infiltration and contributes to angiogenesis in
prostate cancer microenvironment. Oncotarget *5*,1595-1608. 10.18632/oncotarget.1570.

61. Vergadi, E., Ieronymaki, E., Lyroni, K., Vaporidi, K., and Tsatsanis, C. (2017). Akt Signaling Pathway in
Macrophage Activation and M1/M2 Polarization. J Immunol *198*, 1006-1014. 10.4049/jimmunol.1601515.

62. Ghatak, S., Misra, S., Norris, R.A., Moreno-Rodriguez, R.A., Hoffman, S., Levine, R.A., Hascall, V.C., and
Markwald, R.R. (2014). Periostin induces intracellular cross-talk between kinases and hyaluronan in
atrioventricular valvulogenesis. J Biol Chem *289*, 8545-8561. 10.1074/jbc.M113.539882.

63. Silver, J.S., Gunay, K.A., Cutler, A.A., Vogler, T.O., Brown, T.E., Pawlikowski, B.T., Bednarski, O.J.,
Bannister, K.L., Rogowski, C.J., McKay, A.G., et al. (2021). Injury-mediated stiffening persistently activates
muscle stem cells through YAP and TAZ mechanotransduction. Sci Adv *7*. 10.1126/sciadv.abe4501.

64. Chen, M., Zhang, Y., Zhou, P., Liu, X., Zhao, H., Zhou, X., Gu, Q., Li, B., Zhu, X., and Shi, Q. (2020).
Substrate stiffness modulates bone marrow-derived macrophage polarization through NF-kappaB signaling
pathway. Bioact Mater *5*, 880-890. 10.1016/j.bioactmat.2020.05.004.

65. Herpich, C., Franz, K., Ost, M., Otten, L., Coleman, V., Klaus, S., Muller-Werdan, U., and Norman, K. (2021).
Associations Between Serum GDF15 Concentrations, Muscle Mass, and Strength Show Sex-Specific
Differences in Older Hospital Patients. Rejuvenation Res *24*, 14-19. 10.1089/rej.2020.2308.

66. Herpich, C., Hass, U., Kochlik, B., Franz, K., Laeger, T., Klaus, S., Bosy-Westphal, A., and Norman, K. (2021).

Postprandial dynamics and response of fibroblast growth factor 21 in older adults. Clin Nutr *40*, 3765-3771.

10.1016/j.clnu.2021.04.037.

67. Sanchez-Sanchez, J.L., Ader, I., Jeanson, Y., Planat-Benard, V., Vellas, B., Casteilla, L., and de Souto-Barreto,

P. (2023). Periostin Plasma Levels and Changes on Physical and Cognitive Capacities in Community-Dwelling

Older Adults. J Gerontol A Biol Sci Med Sci *78*, 424-432. 10.1093/gerona/glac226.

68. Mathew, S.J., Hansen, J.M., Merrell, A.J., Murphy, M.M., Lawson, J.A., Hutcheson, D.A., Hansen, M.S.,

Angus-Hill, M., and Kardon, G. (2011). Connective tissue fibroblasts and Tcf4 regulate myogenesis.

Development *138*, 371-384. 10.1242/dev.057463.

69. Wang, Y., Jin, S., Luo, D., He, D., Shi, C., Zhu, L., Guan, B., Li, Z., Zhang, T., Zhou, Y., et al. (2021).

Functional regeneration and repair of tendons using biomimetic scaffolds loaded with recombinant periostin.

Nat Commun *12*, 1293. 10.1038/s41467-021-21545-1.

70. Wang, Y.X., Holbrook, C.A., Hamilton, J.N., Garoussian, J., Afshar, M., Su, S., Schürch, C.M., Lee, M.Y.,

Goltsev, Y., Kundaje, A., et al. (2022). A single cell spatial temporal atlas of skeletal muscle reveals cellular

neighborhoods   that   orchestrate   regeneration   and   become   disrupted   in   aging.   BioRxiv.

10.1101/2022.06.10.494732.

71. Babaeijandaghi, F., Paiero, A., Long, R., Tung, L.W., Smith, S.P., Cheng, R., Smandych, J., Kajabadi, N.,

Chang, C.K., Ghassemi, A., et al. (2022). TNFalpha and IFNgamma cooperate for efficient pro- to anti-

inflammatory transition of macrophages during muscle regeneration. Proc Natl Acad Sci U S A *119*,

e2209976119. 10.1073/pnas.2209976119.

72. Sousa, N.S., Bras, M.F., Antunes, I.B., Lindholm, P., Neves, J., and Sousa-Victor, P. (2023). Aging disrupts

MANF-mediated immune modulation during skeletal muscle regeneration. Nat Aging *3*, 585-599.

10.1038/s43587-023-00382-5.

73. Wang, Y., Wehling-Henricks, M., Samengo, G., and Tidball, J.G. (2015). Increases of M2a macrophages and

fibrosis in aging muscle are influenced by bone marrow aging and negatively regulated by muscle-derived

nitric oxide. Aging Cell *14*, 678-688. 10.1111/acel.12350.

Figure legends

Figure 1: Aging impairs myogenic regeneration by disrupting FAPs-derived niche signals. (**A**) Flow cytometric separation and (**B**) quantification of CD45$^-$CD31$^-$SCA1$^+$ FAPs and CD45$^-$CD31$^-$SCA1$^-$ITGA7$^+$ MuSCs per grams of murine muscle tissue (n=18/group). White bars indicate young mice/cells (age: 2-2.5 months), blue bars indicate aged mice/cells (age: 15-18 months; pertains to all subsequent panels). (**C**) Representative images and quantification of spontaneous adipogenic differentiation of FAPs (green: BODIPY, lipid droplet; blue: DAPI, nuclei; n=6; 2 independent experiments; scale bar: 50 µm). (**D**) Representative images and fusion index quantifications in differentiated MuSC (green: Desmin, myofibers; blue: DAPI-nuclei; n=6; 2 independent experiments; scale bar, 100 µm). (**E**) Heat map of top-20 down-regulated ECM genes in microarray analysis comparing freshly sorted FAPs from uninjured skeletal muscle. (**F**) Periostin protein detection and quantification by western blot (Actin: loading control; n=5/group). (**G**) *Postn* mRNA expression in uninjured gastrocnemius (GS), tibialis anterior (TA) and soleus (SL) muscles (n=5/group). (**H**) Periostin protein detection by western blot of cell culture supernatants of FAPs collected during adipogenic differentiation 48h after last medium change (C+: mouse rPOSTN; TP: total protein loading control). (**I, J**) FACS-analysis of CD45$^-$CD31$^-$SCA1$^+$ FAPs per grams of TA muscle comparing uninjured to injured TA at 4- and 8-dpi (n=3/group for uninjured and 8-dpi; n=6/group for 4-dpi). (**K**) Representative hematoxylin/eosin staining and quantification of adipocyte infiltration in injury sites of TA at 14-dpi (yellow arrows indicate adipocyte areas; n=3/group; scale bar: 50 µm). (**L**) Representative Sirius red staining and quantification of fibrotic tissue deposition in TA injury sites at 14-dpi (blue arrows indicate fibrosis areas; n=3/group; scale bar: 50 µm). (**M**) Representative images and CSA quantification of fiber size ranges in injury sites at 14-dpi (n=4/group; scale bar: 100 µm). (**N**) Representative images of Periostin protein immunofluorescence in injury sites in TA at 8-dpi (red: Periostin; blue: DAPI, nuclei; scale bar: 50 µm). (**O**) *Postn* mRNA levels in uninjured TA-muscles compared to TA at 3-, 5-, 8- and 14-dpi (n=4/group). (**P**) Periostin protein western blots and quantification (Actin: loading control) in TA at 3-, 5-, and 8-dpi (n=4/group). (**Q**) UMAP dimensionality reduction of scRNA-seq analysis of 287 CD45$^-$CD31$^-$SCA1$^+$ FAPs FACS-isolated at 4-dpi from young (FAPS-2mo, 128 cells; blue) and aged (FAPs-15mo, 159 cells; red) mice generated using microfluidics-based technology, and (**R**) volcano plot of DEGs (p<0.05 and |Log2|fold change ≥0.5 as cutoffs; black arrow indicates *Postn*). (**S**) Violin plot depiction of *Postn* gene expression per cell in scRNA-seq dataset. (**T**) Enriched Gene Ontology terms of biological process (BP) and cellular component (CC) of downregulated DEGs from scRNA-seq dataset. Data were collected from male mice except in (J, female). All data are presented as mean ± standard error of the mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001.

Figure 2: Ablation of *Postn* gene recapitulates an aging-like phenotype in muscle. (**A**) Representative images and quantification of spontaneous adipogenic differentiation of FAPs from *Postn*-KO (black bar) compared to wildtype (WT) littermates (gray bar; green: BODIPY, lipid droplet; blue: DAPI, nuclei; n=8/group; 2 independent experiments; scale bar: 50 µm). (**B**) Representative images and fusion index quantification in differentiated MuSC from *Postn*$^{-/-}$ (KO; black bar) compared to WT littermates (gray bar; green: Desmin, myofibers; blue: DAPI, nuclei; n=3/group; scale bar: 100 µm). (**C**, **D**) Flow cytometric analysis of CD45$^-$CD31$^-$SCA1$^+$ FAPs (panel C) and CD45$^-$CD31$^-$SCA1$^-$ITGA7$^+$ MuSCs (panel D) per grams of TA-muscle comparing uninjured to injured TA at 4- and 8-dpi (males: uninj n=4/3, 4-dpi n=4/3, 8-dpi n=4/4; females: uninj n=10/10, 4-dpi n=7/7, 8-dpi n=4/4). Data show comparisons of male WT (gray bars, black line) to KO (orange bars and line) and female WT (gray bars, black line) to KO (green bars and line) mice (also applies to subsequent panels). (**E**) Representative hematoxylin/eosin staining and quantification of adipocyte infiltration in injury sites of TA at 14-dpi (yellow arrows indicate adipocyte areas; males: n=3/group, females: n=6/7/group; scale bar: 50 µm). (**F**) Representative Sirius red staining and quantification of fibrotic tissue deposition in TA injury sites at 14-dpi (black arrows indicate fibrosis areas; males: n=3/group, females: n=3/4/group; scale bar: 50 µm). (**G**) Representative images, CSA quantification of fiber size ranges and average in injury site at 14-dpi (males: n=3/group; females: n=4/group; scale bar, 100 µm). All data are presented as mean ± standard error of the mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001.

Figure 3: FAPs-secreted Periostin rescues age-related defect of myogenic regeneration. (**A-D**) Proliferation analysis by KI67 immunostaining of FAPs and MuSCs FACS-isolated from young (Y, 2 months) and aged (O, 15 months) mice and young *Postn*$^{-/-}$ (KO) and littermate wildtype (WT) control mice and treated with mrPOSTN (0.5 µg/ml) for 48h (n=6-12; two independent experiments). (**E**) Representative images of co-cultures from FAPs-WT and FAPs-KO with MuSCs-KO co-cultured for 5 days in myogenic differentiation medium (green: PDGFRα, FAPs; red: Desmin, myofibers; blue: DAPI, nuclei; scale bar: 50µm). (**F**, **G**) Quantifications of fusion index (% of myofibers with >3 DAPI$^+$ nuclei) and myofiber length in co-cultures in panel E, compared to pure MuSC cultures from wildtype (MuSCs-WT) or *Postn*$^{-/-}$ mice (MuSCs-KO) (n=6 wells; two-independent experiments). (**H**) Experimental layout and timeline of FAPs transplants of young FAP^{RFP}-WT and FAP^{RFP}-KO into old recipient mice (TA-muscle, 15 months). (**I**) Representative immunofluorescence images of transplanted TA-muscle sections at 8-dpi (green: eMHC; red: Laminin; blue: DAPI, nuclei; scale bar: 100 µm). (**J**, **K**) Quantifications of eMHC (top) and CSA average (bottom) of FAP^{RFP}-WT and FAP^{RFP}-KO transplants at 8-dpi (n=5/group). (**L**, **M**) Representative images and CSA quantification of fiber size ranges in injury site at 14-dpi comparing 15-months

TA-muscle transplanted with FAP^RFP-WT or FAP^RFP-KO transplants (red: Laminin; blue: DAPI, nuclei; n=5-6/group; scale bar: 100 μm). (**N**) Quantification of CSA averages in regenerating fibers comparing non-transplanted (as shown in Figure 1M) young (2 months; n=4) and old (15 months; n=4) to transplanted 15-months old TA-muscles with either FAP^RFP-WT (n=5) or FAP^RFP-KO (n=6). All data are presented as mean ± standard error of the mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001.

Figure 4: Computational analysis uncovers a Periostin-dependent FAPs-macrophage interaction. (**A**) UMAP of scRNA-seq data generated by droplet-based sequencing technology of all cells isolated from 4-dpi TA-muscle, integrating single cell transcriptomic data of young (2 months; blue-green) and aged (15 month; reds) female C57BL/6J mice. (**B, C**) Unsupervised Louvain clustering analysis and bar plot of population comparison in percentages of nine distinct cell populations identified in regenerating muscle. (**D**) Violin plot showing *Postn* expression levels across the nine cell populations. (**E**) Mean expression and percentage of cells expressing *Postn* in FAPs between 2- and 15-months groups (p-value: Wilcoxon rank sum test). (**F, G**) Age-dependent cell-cell interaction (CCI) scoring of ligand receptor interactions of Periostin with known receptors as a function of age (left panel, F) and interaction depiction of age-related changes in ligand-receptor-driven CCIs linking log fold changes and adjusted p-values (right panel, G) using a published input scRNA-seq dataset comparing young and aged female murine limb-muscle as described in [51]. Cell types within each CCI are color-indicated; red circles indicate statistically significant age-dependent changes in CCI scores. (**H**) Violin plot showing expression for *Itgαv*, *Itga5*, I*tgβ5*, *Itgβ1*, *Itgβ3*, and *Egfr* receptors across individual cells in the nine cell populations detected in our scRNA-seq dataset.

Figure 5: Aging and lack of Periostin impair macrophage polarization in regenerating muscle. (**A**) Representative density plots from FACs analyses of macrophage populations comparing either young (MCs-2mo) to aged (MCs-15mo), or young wildtype (MCs-WT) to littermate *Postn*^-/- (MCs-KO) mice. Total MCs were initially gated as CD45^+CD11b^+F4/80^+ and subsequently for Ly6C-expression to indicate Ly6C-high, pro-inflammatory (M1) vs. Lyc6-low, anti-inflammatory (M2) MCs at 1-, 3-, 4-, and 8-dpi. (**B, C**) Quantitative analysis of MC frequencies as described in panel A (n=3-4/group). Comparisons are young (MCs-2mo; white bars) to aged (MCs-15mo; blue bars) and wildtype (MCs-WT; gray bars) to Postn^-/- (MCs-KO; orange bars; also applies to panels E, F) mice. (**D**) Representative density plots of anti-inflammatory F4/80^+CD206^+ MCs comparing groups as described in panels A-C. MCs were initially gated as CD45^+CD11b^+. (**E, F**) Quantitative analysis of MCs frequencies as described in panel D (n=3-4/group). (**G, H**) Representative immunofluorescence images and

quantifications of F4/80$^+$ and F4/80$^+$CD206$^+$ MCs in injured site of 15-months old TA-muscle sections at 8-dpi

transplanted with FAP^{RFP}-WT (blue bar) or FAP^{RFP}-KO (brown bar) at 1-dpi (red: F4/80, macrophage-marker;

green: CD206, M2-marker; blue: DAPI, nuclei; n=4/group; yellow arrows indicate co-localization of F4/80$^+$ and

CD206$^+$ MCs; scale bar: 20 µm). All animals were male, all data are presented as mean ± standard error of the

mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001.

Figure 6: Periostin promotes anti-inflammatory niche transition through mechano-sensitive integrin-FAK-

AKT signaling. (**A**) *Mrc1* (Mannose receptor c-type 1) mRNA expression in differentiated BMDM exposed to

pro- and anti-inflammatory cytokines (mrIFNγ: 50 ng/mL; mrIL4: 10 ng/mL; n=8/treatment) or co-cultured with

FAPs-WT or FAPs-KO (n=11, co-culture group) for 48h compared to untreated BMDM control (n=6) in 3

independent experiments. (**B**) *Mrc1* mRNA expression in differentiated BMDM treated with mrPOSTN at

indicated concentrations for 48h (n=3/group). (**C, D**) Representative immunofluorescence images and

quantification in untreated, differentiated BMDM control compared to a 48h treatment with mrPOSTN (1 µg/mL)

alone or in combination with integrin receptor-inhibitor Cilengitide (10 nM). mrIL4 (10 ng/mL) was used as

positive control of M2-polarization (green: CD206, M2-marker; red: F4/80, macrophage-marker; blue: DAPI,

nuclei; n=3/treatment; scale bar: 50 µm). (**E**) *Mrc1* mRNA expression in untreated BMDM control compared to

BMDM 48h-exposed to Cilengitide (10nM), mrPOSTN (1 µg/mL), mrPOSTN + Cilengitide (1 µg/mL + 10nM),

mrIL4 (10 ng/ml) or mrIL4 + Cilengitide (10 ng/mL + 10 nM) (n=4/treatment). (**F, G**) Protein detection and

quantification of phosphorylated Ser473-AKT (p-AKT) and total AKT (panel F) and phosphorylated Tyr397-FAK

(p-FAK) and total FAK (panel G) in untreated compared to treated BMDM with mrPOSTN (1 µg/mL) alone or

combined with Cilengitide (10 nM) for indicated times (Actin loading control; n=4; 2 independent experiments in

panels F, G). (**H, I**) Immunofluorescence quantifications of pro- F4/80$^+$CD86$^+$ and anti-inflammatory

F4/80$^+$CD206$^+$ MC polarization of differentiated BMDM encapsulated in soft or high stiffness hydrogels untreated

(control) or treated with mrPOSTN (1 µg/mL) and Cilengitide (10 nM) alone or combined (mrPOSTN +

Cilengitide) for 24h (n=12/group; 2 independent experiments). All data are presented as mean ± standard error of

the mean (SEM), statistical significances are *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 7: Circulating Periostin functions as biomarker of muscle dysfunction in elderly patients. (**A**) Serum

levels of human Periostin (hPOSTN) in young adults (white; n=59; n=16 men; n=43 women), healthy aged,

community-dwelling adults (red; n=59; n=14 men; n=45 women), and aged community-dwelling adults with high

activity levels (Aged/active; green; n=20; n=5 men; n=15 women). (**B**) Serum hPOSTN levels in community-

dwelling aged adults (light blue; n=44), and geriatric hospital patients (Aged/patient; dark blue; n=34). (**C-F**)

Pearson correlations of serum hPOSTN levels and individuals' age (panel C), body mass index (BMI, panel D),

total skeletal muscle mass (panel E) and grip strength (panel F) separately in men (orange, left) and women (green,

right), including young adults and aged community-dwelling individuals only, as described in panel A. All data

are presented as mean $\pm$ standard error of the mean (SEM), statistical significances are *$p<0.05$, **$p<0.01$,

***$p<0.001$, ****$p<0.0001$.