Temporal stability of sex ratio distorter prevalence in natural populations of the isopod *Armadillidium vulgare*

Sylvine Durand[1], Isabelle Giraud[1], Anaïs Loisier[1], Nicolas Bech[1], Frédéric Grandjean[1], Thierry Rigaud[2], Jean Peccoud[1] and Richard Cordaux[1,3,*]

[1] Laboratoire Ecologie et Biologie des Interactions, Equipe Ecologie Evolution Symbiose, Université de Poitiers, UMR CNRS 7267, Bât. B31, 3 rue Jacques Fort, TSA 51106, 86073 Poitiers Cedex 9, France

[2] Laboratoire Biogéosciences, Université Bourgogne Franche-Comté, UMR CNRS 6282, 6 boulevard Gabriel, 21000 Dijon, France

[3] Université Paris-Saclay, CNRS, IRD, UMR Évolution Génomes Comportement Écologie, 91190 Gif-sur-Yvette, France

[*] Corresponding author: richard.cordaux@universite-paris-saclay.fr

Running title: Temporal stability of sex ratio distorter prevalence in *A. vulgare*

Word count: 3386

Abstract

In the terrestrial isopod *Armadillidium vulgare*, many females produce progenies with female-biased

sex ratios due to two feminizing sex ratio distorters (SRD): *Wolbachia* endosymbionts and a nuclear

non-mendelian locus called the *f* element. To investigate the potential impact of these SRD on the

evolution of sex determination, we analyzed their temporal distribution in 6 *A. vulgare* populations

sampled up to 6 times over 12 years, for a total of 29 time points. SRD distribution was

heterogeneous among populations despite their close geographic locations, so that when one SRD

was frequent in a population, the other SRD was rare. In contrast with spatial heterogeneity, our

results overall did not reveal substantial temporal variability in SRD prevalence within populations,

suggesting equilibria in SRD evolutionary dynamics may have been reached or nearly so. SRD

temporal stability was also reflected in mitochondrial and nuclear variation, with two exceptions.

One population was affected by anthropogenic alteration at the sampling spot between the earliest

and second time points, likely explaining a rapid shift in genetic composition. However, subsequent

global stability suggested that population dynamics equilibrium may be reached in a few generations.

In another population, a *Wolbachia* strain replacement coincided with changes in mitochondrial

composition but no change in nuclear composition, thus constituting a typical example of

mitochondrial sweep caused by endosymbiont rise in frequency. Rare incongruence between

Wolbachia strains and mitochondrial haplotypes further highlighted the occurrence of intraspecific

horizontal transmission, making it a biologically relevant parameter for *Wolbachia* evolutionary

dynamics in *A. vulgare*.

Keywords

Sex ratio distorter, endosymbiont, *Wolbachia*, *f* element, sex determination, temporal dynamics

Introduction

Sex ratio distorters (SRD) are selfish genetic elements located on sex chromosomes or transmitted by a single sex, which skew the proportion of males and females in progenies towards the sex that enhances their own vertical transmission (Beukeboom and Perrin, 2014). They are found in a wide range of animal and plant species, of which they tremendously impact the ecology and evolution (Burt and Trivers, 2006; Werren, 2011). SRD are particularly well documented in arthropods, among which is the emblematic bacterial endosymbiont *Wolbachia* (Werren *et al.*, 2008; Kaur *et al.*, 2021). *Wolbachia* is a cytoplasmic, maternally inherited alpha-proteobacterium that often acts as a reproductive parasite by manipulating host reproduction in favor of infected females, thereby conferring itself a transmission advantage. In particular, *Wolbachia* has evolved the ability to induce female-biased sex ratios in host progenies through male killing, thelytokous parthenogenesis and feminization (Werren *et al.*, 2008; Cordaux *et al.*, 2011; Hurst and Frost, 2015; Kaur *et al.*, 2021).

Feminization, causing infected (and non-transmitting) genetic males to develop into (transmitting) phenotypic females, is mostly documented in terrestrial isopods (Martin *et al.*, 1973). In the well-studied *Armadillidium vulgare*, chromosomal sex determination follows female heterogamety (ZZ males and ZW females) (Juchault and Legrand, 1972; Chebbi *et al.*, 2019; Cordaux *et al.*, 2021). In some *A. vulgare* populations, sex ratio is biased by *Wolbachia* bacteria or by a nuclear locus called the *f* element (Rigaud *et al.*, 1997; Cordaux *et al.*, 2011; Cordaux and Gilbert, 2017). Three *Wolbachia* strains are known to naturally occur in *A. Vulgare*: *w*VulC, *w*VulM and *w*VulP (Rigaud *et al.*, 1991; Cordaux *et al.*, 2004; Verne *et al.*, 2007). The *f* element results from the horizontal transfer of a large portion of a feminizing *Wolbachia* genome in the *A. vulgare* genome (Leclercq *et al.*, 2016). The *f* element induces female development, as a W chromosome does, and it shows non-Mendelian inheritance, making it an SRD (Legrand and Juchault, 1984; Leclercq *et al.*, 2016). It has been suggested that SRD may have contributed to the frequent turnovers of sex chromosomes recorded in terrestrial isopods (Juchault and Mocquard, 1993; Juchault and Rigaud, 1995; Becking *et al.*, 2017, 2019; Russell *et al.*, 2021).

To elucidate the potential impact of SRD on the evolution of sex determination in terrestrial isopods, it is essential to clarify the intraspecific evolutionary dynamics of SRD such as *Wolbachia* and the *f* element. Previous studies have shown that: (i) *Wolbachia* and the *f* element are present at variable frequencies in *A. vulgare* populations, (ii) the *f* element is overall more frequent than *Wolbachia*, (iii) the two SRD usually do not co-occur at high frequency in populations, and (iv) mitochondrial haplotypes are tightly linked to *Wolbachia* strains (suggesting stable maternal transmission), but not

to the *f* element (Juchault *et al.*, 1993; Durand *et al.*, 2023). While these results have provided

insights into the spatial distribution of *Wolbachia* and the *f* element in *A. vulgare*, in the present

study we investigate their temporal dynamics. This issue has previously been tackled by a single

study from 1992 (Juchault *et al.*, 1992), in which *Wolbachia* prevalence was found to decrease

concomitantly to an increase of *f* element prevalence in an *A. vulgare* population from Niort (western

France) sampled at three time points over a period of 23 years (1963, 1973 and 1986). However, a

single population was included in the study, thus limiting the breadth of its conclusions. Here we

report an analysis of *Wolbachia* and *f* element distribution in 6 *A. vulgare* populations sampled up to

6 times over a period of 12 years, representing a total of 889 individuals from 29 time points. The

studied populations were sampled in a narrow geographic area in western France, within 70 km of

the Niort population (Juchault *et al.*, 1992), to control for spatial dynamics. Analyzed in the context of

mitochondrial and nuclear variation, our results highlight an overall temporal stability of SRD

distribution in *A. vulgare*, with few exceptions.

Materials and Methods

We analyzed 889 *A. vulgare* individuals from 6 natural populations sampled in western France

between 2003 and 2017. DNA samples from Beauvoir-2017, Chizé-2017, Coulombiers-2017, Gript-

2017, La Crèche-2017 and Poitiers-2015 were available from (Durand *et al.*, 2023). All other

individuals were collected by hand. Individuals were sexed and stored in alcohol or at -20°C prior to

DNA extraction. Total genomic DNA of samples collected between 2003 and 2013 was extracted from

gonads using phenol and chloroform (Kocher *et al.*, 1989) and DNA of samples collected between

2014 and 2016 was extracted from the head and legs, as described previously (Leclercq *et al.*, 2016).

We used four molecular markers to assess the presence of *Wolbachia* and the *f* element in DNA

extracts: *Jtel* (Leclercq *et al.*, 2016), *wsp* (Braig *et al.*, 1998), *recR* (Badawi *et al.*, 2014) and *ftsZ*

(Werren *et al.*, 1995) (Table S1). While *Jtel* is specific to the *f* element, *wsp* and *recR* are specific to

Wolbachia, and *ftsZ* is present in both the *f* element and *Wolbachia* (Leclercq *et al.*, 2016). We

assessed the presence or absence of these markers by PCR, as described previously (Leclercq *et al.*,

2016). Different amplification patterns were expected for individuals with *Wolbachia* only (*Jtel-,*

wsp+, recR+, ftsZ+), the *f* element only (*Jtel+, wsp-, recR-, ftsZ+*), both *Wolbachia* and the *f* element

present (*Jtel+, wsp+, recR+, ftsZ+*) or both *Wolbachia* and the *f* element lacking (*Jtel-, wsp-, recR-,*

ftsZ-). The few individuals exhibiting other amplification patterns were classified as "undetermined

status".

To characterize *Wolbachia* strain diversity, *wsp* PCR products were purified and Sanger sequenced using both forward and reverse primers by GenoScreen (Lille, France). Sequences from forward and reverse reads were assembled using Geneious v.7.1.9 to obtain one consensus sequence per individual.

To evaluate mitochondrial diversity, we amplified by PCR a ~700 bp-long portion of the Cytochrome Oxidase I (*COI*) gene in all individuals (Folmer *et al.*, 1994) (Table S1). PCR products were purified and Sanger sequenced as described above. Haplotype networks were built using the *pegas* package for R (Paradis, 2010).

To evaluate nuclear diversity, all individuals were genotyped at 22 microsatellite markers (Verne *et al.*, 2006; Giraud *et al.*, 2013) distributed in five multiplexes (Multiplex 1: Av1, Av2, Av4, Av5, Av9; Multiplex 2: Av6, Av3, Av8; Multiplex 3: AV0023, AV0056, AV0085, AV0086, AV0096; Multiplex 4: AV0002, AV0016, AV0018, AV0032, AV0099; Multiplex 5: AV0061, AV0063, AV0089, AV0128) (Table S2). PCR was performed using fluorescence-marked forward primers, as described previously (Durand *et al.*, 2017). PCR fragments were separated by electrophoresis on an ABI 3730XL automated sequencer by Genoscreen (Lille, France). Alleles were scored using the software GeneMapper 3.7 (Applied Biosystems), each genotype being independently read by two people.

Of the 22 amplified microsatellite markers, Av4 and Av5 could not be scored because of multiple peaks, AV0096 and AV0128 did not amplify consistently, and AV0023 and AV0061 were monomorphic across the dataset. The *Genepop* package for R (Rousset, 2008) detected no linkage disequilibrium between the 16 remaining loci. We tested for the presence of null alleles using a combination of software, as recommended previously (Dąbrowski *et al.*, 2014). We used Micro-Checker (Van Oosterhout *et al.*, 2004), Cervus (Kalinowski *et al.*, 2007) and ML-NullFreq (Kalinowski and Taper, 2006). As a result, AV0099 was discarded because it consistently presented hints of null alleles in many populations and sampling years. For the following analyses, we removed individuals whose genotypes were available for fewer than 13 out of the 15 remaining markers.

Hardy-Weinberg equilibrium was tested for each locus, locality and sampling year with an exact test using Markov chain with the *Genepop* package for R. We used the Fstat software to calculate allelic richness (based on a minimum of 3 individuals). To test whether the same genetic populations have been sampled each year, we searched for genetic clusters without a priori with a Bayesian method using the software Structure (Pritchard *et al.*, 2000). We selected the admixture model as well as the option of correlated allele frequencies. We varied the number of clusters (K) from 2 to 9. For each value of K, we carried out 20 independent runs, as recommended in (Evanno *et al.*, 2005), with a total number of 100,000 iterations and a burn-in of 100,000. To determine the most likely value of K,

we applied the method described in (Evanno *et al.*, 2005) and implemented in Structure Harvester version 0.6.9 (Earl and vonHoldt, 2012). In addition, we performed a Discriminant Analysis of Principal Components (DAPC) (Jombart *et al.*, 2010) on populations according to their sampling locality and year with the *adegenet* package (Jombart, 2008) to search for potential discrepancies across time points within populations.

All analyses were performed with R v.3.6.0 (R Development Core Team, 2013). When multiple statistical tests of the same hypothesis were performed, we corrected p-values with the Benjamini-Yekutieli procedure (Benjamini and Yekutieli, 2001). Figures were realized with *ggplot2* (Wickham *et al.*, 2020). Results from Structure were processed with the program Distruct (Rosenberg, 2003) for graphical representation.

Results

We tested the presence of *Wolbachia* and the *f* element in 889 individuals (627 females and 262 males) from 6 populations sampled at various time points between 2003 and 2017, representing a total of 29 sampling points (Tables 1 and S3). We found that 29.9% of individuals carried only the *f* element, 15.2% carried only *Wolbachia*, 0.6% carried both SRD and 54.3% carried none. While both SRD were mostly found in females, they were also sometimes present in males, as previously reported (Durand *et al.*, 2023). *Wolbachia*-infected individuals carried one of the three previously known *Wolbachia* strains of *A. vulgare*: *w*VulC (n=22), *w*VulM (n=25) or *w*VulP (n=83).

Overall, both SRD were found in at least one time point in all 6 populations (Figure 1). However, the populations displayed substantial variation in the distribution of the two SRD. In Beauvoir, Chizé, Coulombiers and La Crèche populations, the *f* element was predominant (29-86% frequency) and *Wolbachia* rare (2-10%). Conversely, *Wolbachia* was frequent (63%) and the *f* element very rare (2%) in Poitiers. Finally, both SRD were very rare in Gript (2-3%). Remarkably, SRD prevalence was qualitatively stable across time points (spanning up to 12 years) within all populations (Figure 1). Yet, in Poitiers where *Wolbachia* remained globally frequent overtime, the *w*VulC strain progressively decreased in frequency while the *w*VulP strain exhibited the opposite trend.

Sequencing of the *COI* mitochondrial gene of 884 individuals identified 12 haplotypes, 9 of which have previously been detected in *A. vulgare* populations (named I to VIII, and XII) (Durand *et al.*, 2023) and three are newly described haplotypes (XXIV to XXVI; GenBank accession numbers OR074129 to OR074131, respectively) (Figure 2 and Table S3). There was an excellent congruence

between *Wolbachia* strains and mitochondrial haplotypes, as previously reported (Verne *et al.*, 2012; Durand *et al.*, 2023). Indeed, individuals carrying *w*VulC were associated with either haplotype V or its close relatives (XII and XXVI), those carrying *w*VulP were associated with haplotype VII and those carrying *w*VulM were associated with haplotype II. Exceptions included two individuals infected by *w*VulM, which were associated with the distantly related haplotypes I and III. By contrast, the *f* element was found in 8 different mitochondrial backgrounds (I to VI, VIII and XXV) distributed across the mitochondrial network.

All time points considered, there was between 3 (in Chizé) and 6 (in Poitiers) haplotypes per population (Figures 3 and S1). Haplotype frequencies were globally stable across time points within populations (e.g. Coulombiers, Figure 3A), with the notable exceptions of La Crèche and Poitiers populations. In La Crèche, a shift in major haplotypes occurred between 2005 and 2012, with the rarefaction of haplotype III and V being concomitant with the rise in frequency of haplotypes I and VIII, followed by stability since 2012 (Figure 3B). In Poitiers, the rise in frequency of haplotype VII (associated with *Wolbachia* strain *w*VulP) coincided with a relative decrease of *w*VulC-associated haplotypes V, XII and XXVI (Figure 3C).

To test if the patterns of SRD and mitochondrial variation were also reflected in the nuclear genome, we examined variation in 667 individuals with genotype information available for at least 13 out of the 15 retained microsatellite markers (see Materials and Methods), representing a total of 28 sampling points (Poitiers-2015 was discarded due to low genotyping success) (Table S3). Allelic richness ranged from 1 to 4.78 (Table S4) and all loci were at Hardy-Weinberg equilibrium for all sampling points.

To evaluate whether the same genetic populations had been sampled at each time point, we delineated genetic clusters. A Delta-K analysis (Evanno *et al.*, 2005) inferred that the best fit to the data was obtained for K=2 genetic clusters (Figure S2). Indeed, the major structuration of the data grouped Poitiers and Coulombiers on the one hand, and Beauvoir, Chizé, La Crèche and Gript on the other hand in a Structure analysis (Figure 4). No obvious change in genetic structure was apparent between time points within populations in the Structure analysis. The DAPC confirmed the major population structuration, as the first component separated Poitiers and Coulombiers from the other populations (Figure 5). It also supported overall homogeneity of populations across time points, apart from La Crèche-2005, which was separated from the other La Crèche time points in the second component of the DAPC.

Discussion

Wolbachia and *f* element distributions are highly heterogeneous among populations, despite their modest geographic distance of at most 80 km. The emerging trend is that when one SRD is frequent in a population, the other SRD is rare, as noted previously (Durand *et al.*, 2023). As it causes a stronger bias toward females, *Wolbachia* is expected to prevail over the *f* element in *A. vulgare* populations (Rigaud, 1997; Cordaux and Gilbert, 2017). Yet, the *f* element was the dominant SRD in 4 out of 6 populations we studied and was rising in frequency in Niort (Juchault *et al.*, 1992). Overall, the *f* element is more widespread than *Wolbachia* in *A. vulgare* populations (Juchault *et al.*, 1993; Durand *et al.*, 2023). Previously proposed explanations include a higher fitness cost entailed by *Wolbachia* relative to the *f* element and occasional paternal transmission of the *f* element (which some males carry, Figure 1) enabled by masculinizing epistatic alleles (Juchault *et al.*, 1992; Rigaud and Juchault, 1993; Rigaud, 1997; Rigaud and Moreau, 2004; Cordaux and Gilbert, 2017).

In contrast with spatial heterogeneity, our sampling scheme with up to 6 sampling time points spanning 12 years per population highlights a global temporal stability in SRD prevalence within populations. At first glance, this qualitative pattern differs from that previously reported for the Niort population, in which *Wolbachia* prevalence was found to decrease concomitantly to an increase of *f* element prevalence over a period of 23 years (Juchault *et al.*, 1992). Given *A. vulgare*'s generation time of one year, our study might have spanned too few generations (12) to capture variation in SRD prevalence, which the Niort study spanning 23 generations did. Yet, the time scale of our study enabled us to detect variation in *Wolbachia* strain prevalence, as well as mitochondrial and nuclear variation within and between populations, suggesting that lack of resolution is not an issue. Alternatively, most of the populations we studied may reflect some relatively stable equilibrium with respect to SRD evolutionary dynamics, an equilibrium that the Niort population might not have reached. Indeed, theoretical models have indicated that when feminizing factors are in competition, the one that induces the strongest bias toward females is expected to spread in the population (Taylor, 1990). Thus, a single SRD should remain in the population at equilibrium. Consistently, in most of the populations we analyzed, a single SRD occurs at high frequency, suggesting that these populations may be at or near equilibrium for an SRD. It has also been suggested that an apparent stability could be due to hidden processes such as population structure (including extinction-recolonization processes), intragenomic conflicts and coevolutionary processes (Hatcher, 2000).

The temporal stability of SRD in most of *A. vulgare* populations is also reflected in host mitochondrial and nuclear variation, with two notable singularities. The first is La Crèche population in 2005, which differs from the other time points (2012 to 2017) on both mitochondrial and nuclear grounds.

Interestingly, the sampling spot in La Crèche has been altered by land remodeling between 2005 and

2012. This anthropogenic activity may have caused the reduction or collapse of the historic *A.*

vulgare population and the introduction of new individuals as part of the addition of materials during

the remodeling (e.g., soil from another location). Such an extinction-recolonization scenario may

explain the loss of *Wolbachia* (present at low frequency in 2005) and the increase in *f* element

frequency in 2012. It is noteworthy that from 2012 on, SRD, mitochondrial and nuclear variation have

been stable, suggesting that stabilization of the population dynamics may be reached in a few years,

i.e., in a few generations.

The second case of instability is the Poitiers population, which is stable with respect to nuclear

variation but not to both mitochondrial and SRD variation. Poitiers is the only population in our

dataset in which *Wolbachia* is the dominant SRD across time points, thus highlighting a qualitative

pattern of temporal stability. However, our results indicate that the rise in frequency of the *w*VulP

strain correlates with a decrease of the *w*VulC strain, suggesting a *Wolbachia* strain replacement in

this population. The *w*VulP strain is characterized by a recombination event involving *w*VulC (Verne

et al., 2007), indicating that *w*VulC is older than *w*VulP, which is consistent with the situation

recorded in Poitiers. Assuming the driver of this replacement is *Wolbachia* and not another

cytoplasmic element (like the mitochondrion), replacement of *w*VulC by *w*VulP could be due to the

latter strain having a transmission advantage over the former strain. Unfortunately, the *w*VulP strain

is not very well characterized, and while feminization induction is likely (Verne *et al.*, 2007), it has not

been formally demonstrated and compared to feminization induced by wVulC (Rigaud *et al.*, 1991;

Cordaux *et al.*, 2004). The respective costs of these two *Wolbachia* strains has not been investigated

neither. In any event, because *Wolbachia* and mitochondria are co-inherited cytoplasmic entities,

changes in *Wolbachia* strains associated with different mitochondrial haplotypes are expected to

lead to concomitant changes in mitochondrial variation, but no change in nuclear variation.

Therefore, our observations in Poitiers may constitute a typical example of mitochondrial sweep

caused by endosymbiont rise in frequency (Galtier *et al.*, 2009).

Wolbachia dynamics in Poitiers also illustrates that transovarial, maternal transmission is the main

transmission mode of *Wolbachia* in *A. vulgare*. However, non-maternal transmission also appears to

occur, as testified by two individuals with *w*VulM from Beauvoir and La Crèche. These individuals

carry mitochondrial haplotypes I and III, respectively, unlike all other *w*VulM-infected individuals

which carry haplotype II. As haplotypes I, II and III are distantly related, the most likely explanation is

that the two unusual individuals have acquired *Wolbachia* by horizontal transfer. Horizontal transfer

of *Wolbachia* is largely documented in arthropods (O'Neill *et al.*, 1992; Werren *et al.*, 1995; Heath *et*

al., 1999; Vavre *et al.*, 1999), including terrestrial isopods (Bouchon *et al.*, 1998; Cordaux *et al.*, 2001,

2012). Potential mechanisms in isopods include contact between wounded individuals (Rigaud and

Juchault, 1995) and cannibalism/predation (Le Clec'h *et al.*, 2013). In total, we infer that 2 out of 136

Wolbachia-infected individuals may have acquired their symbionts by horizontal transmission. This

may be an underestimate, as horizontal transfers between individuals carrying the same

mitochondrial haplotype cannot be detected with our approach. Although it is apparently rare,

horizontal transmission occurs at a measurable rate in *A. vulgare*, suggesting that it is a parameter of

importance in *Wolbachia* evolutionary dynamics in this species.

Acknowledgments

We thank Sébastien Verne and Victorien Valette for assistance with sample collection. This work was

funded by Agence Nationale de la Recherche Grants ANR-15-CE32-0006 (CytoSexDet) to RC and TR,

and ANR-20-CE02-0004 (SymChroSex) to JP, and intramural funds from the CNRS and the University

of Poitiers.

Author contribution statement

RC, JP and TR designed the experiments. FG, NB and JP sampled populations. SD, IG and AL

performed laboratory work. SD performed data analyses. RC and SD wrote the first draft of the

manuscript. TR, JP, NB and FG amended the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Data archiving

Mitochondrial haplotypes are available in GenBank under accession numbers OR074129 to

OR074131. All other data are provided in the supplementary information.

References

Badawi M, Giraud I, Vavre F, Grève P, Cordaux R (2014). Signs of Neutralization in a Redundant Gene
Involved in Homologous Recombination in *Wolbachia* Endosymbionts. *Genome Biol Evol* **6**:
2654–2664.

Becking T, Chebbi MA, Giraud I, Moumen B, Laverré T, Caubet Y, *et al.* (2019). Sex chromosomes
control vertical transmission of feminizing Wolbachia symbionts in an isopod. *PLOS Biol* **17**:
e3000438.

Becking T, Giraud I, Raimond M, Moumen B, Chandler C, Cordaux R, *et al.* (2017). Diversity and
evolution of sex determination systems in terrestrial isopods. *Sci Rep* **7**: 1–14.

Benjamini Y, Yekutieli D (2001). The control of the false discovery rate in multiple testing under
dependency. *Ann Stat* **29**: 1165–1188.

Beukeboom LW, Perrin N (2014). *The evolution of sex determination*. Oxford University Press: Oxford.

Bouchon D, Rigaud T, Juchault P (1998). Evidence for widespread Wolbachia infection in isopod
crustaceans: molecular identification and host feminization. *Proc Biol Sci* **265**: 1081–90.

Braig HR, Zhou WG, Dobson SL, O'Neill SL (1998). Cloning and characterization of a gene encoding the
major surface protein of the bacterial endosymbiont Wolbachia pipientis. *J Bacteriol* **180**:
2373–2378.

Burt A, Trivers R (2006). *Genes in conflict*. The Belknap Press of Harvard University Press: Cambridge,
Massachusetts.

Chebbi MA, Becking T, Moumen B, Giraud I, Gilbert C, Peccoud J, *et al.* (2019). The Genome of
Armadillidium vulgare (Crustacea, Isopoda) Provides Insights into Sex Chromosome Evolution
in the Context of Cytoplasmic Sex Determination. *Mol Biol Evol* **36**: 727–741.

Cordaux R, Bouchon D, Greve P (2011). The impact of endosymbionts on the evolution of host sex-
determination mechanisms. *Trends Genet* **27**: 332–41.

Cordaux R, Chebbi MA, Giraud I, Pleydell DRJ, Peccoud J (2021). Characterization of a Sex-
Determining Region and Its Genomic Context via Statistical Estimates of Haplotype
Frequencies in Daughters and Sons Sequenced in Pools. *Genome Biol Evol* **13**: evab121.

Cordaux R, Gilbert C (2017). Evolutionary Significance of Wolbachia-to-Animal Horizontal Gene
Transfer: Female Sex Determination and the f Element in the Isopod Armadillidium vulgare.
Genes **8**: 186.

Cordaux R, Michel Salzat A, Bouchon D (2001). *Wolbachia* infection in crustaceans: novel hosts and
potential routes for horizontal transmission. *J Evol Biol* **14**: 237–243.

Cordaux R, Michel-Salzat A, Frelon-Raimond M, Rigaud T, Bouchon D (2004). Evidence for a new
feminizing Wolbachia strain in the isopod Armadillidium vulgare: evolutionary implications.
Heredity **93**: 78–84.

Cordaux R, Pichon S, Hatira HB, Doublet V, Greve P, Marcade I, *et al.* (2012). Widespread Wolbachia
infection in terrestrial isopods and other crustaceans. *Zookeys* **176**: 123–31.

Dąbrowski MJ, Pilot M, Kruczyk M, Żmihorski M, Umer HM, Gliwicz J (2014). Reliability assessment of
null allele detection: inconsistencies between and within different methods. *Mol Ecol Resour*
14: 361–373.

Durand S, Cohas A, Braquart-Varnier C, Beltran-Bech S (2017). Paternity success depends on male
genetic characteristics in the terrestrial isopod *Armadillidium vulgare*. *Behav Ecol Sociobiol*
71: 90.

Durand S, Lheraud B, Giraud I, Bech N, Grandjean F, Rigaud T, *et al.* (2023). Heterogeneous
distribution of sex ratio distorters in natural populations of the isopod Armadillidium vulgare.
Biol Lett **19**: 20220457.

Earl DA, vonHoldt BM (2012). STRUCTURE HARVESTER: a website and program for visualizing
STRUCTURE output and implementing the Evanno method. *Conserv Genet Resour* **4**: 359–
361.

Evanno G, Regnaut S, Goudet J (2005). Detecting the number of clusters of individuals using the
software structure: a simulation study. *Mol Ecol* **14**: 2611–2620.

Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994). DNA primers for amplification of
mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar*
Biol Biotechnol **3**: 294–299.

Galtier N, Nabholz B, Glémin S, Hurst GDD (2009). Mitochondrial DNA as a marker of molecular
diversity: a reappraisal. *Mol Ecol* **18**: 4541–4550.

Giraud I, Valette V, Bech N, Grandjean F, Cordaux R (2013). Isolation and characterization of
microsatellite loci for the isopod crustacean *Armadillidium vulgare* and transferability in
terrestrial isopods. *Plos One* **8**: e76639.

Hatcher MJ (2000). Persistence of selfish genetic elements: population structure and conflict. *Trends*
Ecol Evol **15**: 271–277.

Heath BD, Butcher RDJ, Whitfield WGF, Hubbard SF (1999). Horizontal transfer of Wolbachia
between phylogenetically distant insect species by a naturally occurring mechanism. *Curr Biol*
9: 313–316.

Hurst GDD, Frost CL (2015). Reproductive parasitism: maternally inherited symbionts in a biparental
world. *Cold Spring Harb Perspect Biol* **7**: a017699.

Jombart T (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics **24**: 1403–1405.

Jombart T, Devillard S, Balloux F (2010). Discriminant analysis of principal components: a new
method for the analysis of genetically structured populations. *BMC Genet* **11**: 94.

Juchault P, Legrand JJ (1972). Croisement de néo-mâles experimentaux chez Armadillidium vulgare
Latr. (Crustace, Isopode, Oniscoide). Mise en evidence d'une hétérogamétie femelle. *C R*
Acad Sci Paris **274**: 1387–1389.

Juchault P, Mocquard JP (1993). Transfer of a parasitic sex factor to the nuclear genome of the host:
A hypothesis on the evolution of sex-determining mechanisms in the terrestrial isopod
Armadillidium vulgare Latr. *J Evol Biol* **6**: 511–528.

Juchault P, Rigaud T (1995). Evidence for female heterogamety in two terrestrial crustaceans and the
problem of sex chromosome evolution in isopods. *Heredity* **75**: 466–471.

Juchault P, Rigaud T, Mocquard J-P (1992). Evolution of sex-determining mechanisms in a wild
population of *Armadillidium vulgare* Latr. (Crustacea, Isopod) : competition between two
feminizing parasitic sex factors. *Heredity* **69**: 382–390.

Juchault P, Rigaud T, Mocquard J-P (1993). Evolution of sex determination and sex ratio variability in
wild populations of *Armadillidium vulgare* (Latr.) (Crustacea, Isopoda): a case study in conflict
resolution. *Acta Oecologica* **14**: 547–562.

Kalinowski ST, Taper ML (2006). Maximum likelihood estimation of the frequency of null alleles at
microsatellite loci. *Conserv Genet* **7**: 991–995.

Kalinowski ST, Taper ML, Marshall TC (2007). Revising how the computer program cervus
accommodates genotyping error increases success in paternity assignment. *Mol Ecol* **16**:
1099–1106.

Kaur R, Shropshire JD, Cross KL, Leigh B, Mansueto AJ, Stewart V, *et al.* (2021). Living in the
endosymbiotic world of Wolbachia: A centennial review. *Cell Host Microbe* **29**: 879–893.

Kocher TD, Thomas WK, Meyer A, Edwards SV, Paabo S, Villablanca FX, *et al.* (1989). Dynamics of
mitochondrial-DNA evolution in animals: amplification and sequencing with conserved
primers. *Proc Natl Acad Sci U S A* **86**: 6196–6200.

Le Clec'h W, Chevalier FD, Genty L, Bertaux J, Bouchon D, Sicard M (2013). Cannibalism and Predation
as Paths for Horizontal Passage of Wolbachia between Terrestrial Isopods. *PLOS ONE* **8**:
e60232.

Leclercq Sb, Thézé J, Chebbi MA, Giraud I, Moumen B, Ernenwein L, *et al.* (2016). Birth of a W sex
chromosome by horizontal transfer of *Wolbachia* bacterial symbiont genome. *Proc Natl Acad*
Sci U S A **113**: 15036–15041.

Legrand JJ, Juchault P (1984). Nouvelles données sur le déterminisme génétique et épigénétique de
la monogénie chez le crustacés isopodes terrestres *Armadillidium vulgare* Latr. *Génét Sél Evol*
16: 57–84.

Martin G, Juchault P, Legrand JJ (1973). Mise en évidence d'un micro-organisme intracytoplasmique
symbiote de l'Oniscoide *Armadillidium vulgare* L. dont la présence accompagne
l'intersexualité ou la féminisation totale des mâles génétiques de la lignée thélygène.
Comptes Rendus Académie Sci Paris **276**: 2313–2316.

O'Neill SL, Giordano R, Colbert AM, Karr TL, Robertson HM (1992). 16S rRNA phylogenetic analysis of
the bacterial endosymbionts associated with cytoplasmic incompatibility in insects. *Proc Natl*
Acad Sci **89**: 2699–2702.

Paradis E (2010). pegas: an R package for population genetics with an integrated–modular approach.
Bioinformatics **26**: 419–420.

Pritchard JK, Stephens M, Donnelly P (2000). Inference of Population Structure Using Multilocus
420   Genotype Data. *Genetics* **155**: 945.

421   R Development Core Team (2013). *R: A language and environment for statistical computing.*

422   Rigaud T (1997). Inherited microorganisms and sex determination of arthropod hosts. In: O'Neill SL,
423   Hoffmann AA, Werren JH (eds) *Influential passengers: inherited microorganisms and*
424   *arthropod reproduction*, Oxford Univ. Press: Oxford, pp 81–101.

425   Rigaud T, Juchault P (1993). Conflict between feminizing sex ratio distorters and an autosomal
426   masculinizing gene in the terrestrial isopod Armadillidium vulgare Latr. *Genetics* **133**: 247–
427   252.

428   Rigaud T, Juchault P (1995). Success and failure of horizontal transfers of feminizing Wolbachia
429   endosymbionts in woodlice. *J Evol Biol* **8**: 249–255.

430   Rigaud T, Juchault P, Mocquard JP (1997). The evolution of sex determination in isopods crustaceans.
431   *Bioessays* **19**: 409–416.

432   Rigaud T, Moreau M (2004). A cost of Wolbachia-induced sex reversal and female-biased sex ratios:
433   decrease in female fertility after sperm depletion in a terrestrial isopod. *Proc R Soc B-Biol Sci*
434   **271**: 1941–1946.

435   Rigaud T, Souty Grosset C, Raimond R, Mocquard JP, Juchault P (1991). Feminizing endocytobiosis in
436   the terrestrial crustacean *Armadillidium vulgare* Latr. (Isopoda): Recent acquisitions.
437   *Endocytobiosis Cell Res* **7**: 259–273.

438   Rosenberg NA (2003). DISTRUCT: a program for the graphical display of population structure. *Mol*
439   *Ecol Notes* **4**: 137–138.

440   Rousset F (2008). GENEPOP'007: a complete re-implementation of the GENEPOP software for
441   Windows and Linux. *Mol Ecol Resour* **8**: 103–106.

442   Russell A, Borrelli S, Fontana R, Laricchiuta J, Pascar J, Becking T, *et al.* (2021). Evolutionary transition
443   to XY sex chromosomes associated with Y-linked duplication of a male hormone gene in a
444   terrestrial isopod. *Heredity* **127**: 266–277.

445   Taylor DR (1990). Evolutionary consequences of cytoplasmic sex ratio distorters. *Evol Ecol* **4**: 235–
446   248.

447   Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004). Micro-checker: software for
448   identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Notes* **4**: 535–
449   538.

450   Vavre F, Fleury F, Lepetit D, Fouillet P, Boulétreau M (1999). Phylogenetic evidence for horizontal
451   transmission of Wolbachia in host-parasitoid associations. *Mol Biol Evol* **16**: 1711–1723.

452   Verne S, Johnson M, Bouchon D, Grandjean F (2007). Evidence for recombination between feminizing
453   Wolbachia in the isopod genus Armadillidium. *Gene* **397**: 58–66.

454   Verne S, Johnson M, Bouchon D, Grandjean F (2012). Effects of parasitic sex-ratio distorters on host
455   genetic structure in the Armadillidium vulgare-Wolbachia association. *J Evol Biol* **25**: 264–76.

Verne S, Puillandre N, Brunet G, Gouin N, Samollow PB, Anderson JD, *et al.* (2006). Characterization
of polymorphic microsatellite loci in the terrestrial isopod *Armadillidium vulgare*. *Mol Ecol*
Notes **6**: 328–330.

Werren JH (2011). Selfish genetic elements, genetic conflict, and evolutionary innovation. *Proc Natl*
Acad Sci U S A **108 Suppl 2**: 10863–10870.

Werren JH, Baldo L, Clark ME (2008). Wolbachia: master manipulators of invertebrate biology. *Nat*
Rev Microbiol **6**: 741–51.

Werren JH, Zhang W, Guo LR (1995). Evolution and phylogeny of Wolbachia: reproductive parasites
of arthropods. *Proc Biol Sci* **261**: 55–63.

Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, *et al.* (2020). ggplot2: Create
Elegant Data Visualisations Using the Grammar of Graphics.

Figure legends

Figure 1. Prevalence of sex ratio distorters in *Armadillidium vulgare* males (m) and females (f) from

six populations sampled at different time points.

Figure 2. Haplotype network of 12 mitochondrial variants from six *Armadillidium vulgare* populations

sampled at different time points. Each circle represents one haplotype and circle diameter is

proportional to the number of individuals carrying the haplotype. Branch lengths connecting circles

are proportional to divergence between haplotypes.

Figure 3. Variation in haplotype counts across years for three *Armadillidium vulgare* populations

from (A) Coulombiers, (B) La Crèche and (C) Poitiers. Prevalence of sex ratio distorters is color-coded.

Figure 4. Assignment of individuals from six *Armadillidium vulgare* populations sampled at different

time points to one of two genetic clusters (blue and pink colors) following Bayesian analysis. Each bar

represents an individual, and the proportion of each color represents the probability of assignment

to the corresponding cluster.

Figure 5. Discriminant Analysis of Principal Components scatterplot. Dots represent individuals. Each

of the 29 sampling points presents a 95% inertia ellipse and is labeled with two letters indicating the

population and the two last digits of the sampling year.

Table 1. Prevalence of *Wolbachia* and *f* element sex ratio distorters in 6 populations of *Armadillidium vulgare*.

Population	Sampling year	Sample size	Sex	n	No *f* element, no *Wolbachia*	Only *f* element	Only *Wolbachia* wVulC	wVulM	wVulP	Undetermined	Both wVulM and *f* element	Both wVulC and *f* element	Undetermined status
Beauvoir	2017	31	Males	6	5	1							
			Females	25	9	14		1					1
	2016	25	Males	4	3	1							
			Females	21	8	11		1					1
	2015	21	Males	6	6								
			Females	15	6	7		1					1
	2013	37	Males	11	10			1					
			Females	26	7	14	1	4					
	2012	20	Males	10	7	1		2					
			Females	10	3	5		2					
	2005	34	Males	16	16								
			Females	18	3	11	1	2					1
Chizé	2017	52	Males	8	2	6							
			Females	44	3	36		2			3		
	2016	26	Males	6	3	3							
			Females	20	1	19							
Coulombiers	2017	24	Males	4	2	2							
			Females	20	6	13	1						
	2016	48	Males	23	13	9							1
			Females	25	10	13							2
	2015	27	Males	7	5	2							
			Females	20	4	14	2						
	2014	14	Males	6	5	1							
			Females	8	3	5							
	2005	29	Males	7	5	2							
			Females	22	2	17				1		2	
Gript	2017	45	Males	15	15								
			Females	30	26	2	2						
	2016	16	Males	4	4								
			Females	12	12								

	2015	30	Males	5	5								
			Females	25	24								1
	2014	35	Males	4	4								
			Females	31	30	1							
	2013	22	Males	3	3								
			Females	19	17	2							
	2005	34	Males	10	9								1
			Females	24	18	1	2						3
La Crèche	2017	58	Males	21	19	2							
			Females	37	23	13	1						
	2015	16	Males	4	3								1
			Females	12	7	5							
	2013	17	Males	13	11	2							
			Females	4	3	1							
	2012	20	Males	10	3	7							
			Females	10	2	8							
	2005	39	Males	17	14								3
			Females	22	11	3	1	1					6
Poitiers	2015	23	Males	4	4								
			Females	19	10	1			4	4			
	2014	41	Males	8	8								
			Females	33	5	1	1	3	23				
	2013	68	Males	19	17								2
			Females	49	4	1		2	41				1
	2010	21	Males	0									
			Females	21	1			5	13				2
	2003	16	Males	11	8			1					2
			Females	5				2	2	1			
Total males				262	209	39	1	3	0	0	0	0	10
Total females				627	258	218	19	19	83	6	3	2	19
Total				889	467	257	20	22	83	6	3	2	29

Figure 1

Figure 2

SRD
- no *f*, no *Wolbachia*
- *f*
- *w*VulC strain
- *w*VulM strain
- *w*VulP strain
- unknown strain
- both *f* and *w*VulC strain
- both *f* and *w*VulM strain

Figure 3

Figure 4

Figure 5