Anti- and pro-inflammatory milieu differentially regulate differentiation and immune functions of oligodendrocyte progenitor cells

Omri Zveik[1,2], Ariel Rechtman[1,2], Livnat Brill[1,2], and Adi Vaknin-Dembinsky[1,2]

1. Faculty of Medicine, Hebrew University of Jerusalem, 91120, Israel;
2. The Department of Neurology and Laboratory of Neuroimmunology, The Agnes-Ginges Center for Human Neurogenetics, Hadassah-Hebrew University Medical Center, Jerusalem 91120, Israel

Corresponding author:

Adi Vaknin-Dembinsky MD, PhD

Adembinsky@gmail.com

Hadassah Medical Center, Ein–Kerem

P.O.B. 12000, Jerusalem 91120, Israel

Abstract

Oligodendrocyte progenitor cells (OPC) were regarded for years solely for their regenerative role; however, their immune-modulatory roles have gained much attention recently. Despite extensive studies on OPCs, there are limited data elucidating the interactions between their intrinsic regenerative and immune functions, as well as their relationship with the inflamed central nervous system (CNS) environment. We examined the effects of pro-inflammatory cytokines, represented by interferon (IFN)-γ and tumor necrosis factor (TNF)-α, as well as anti-inflammatory cytokines, represented by interleukin(IL)-4 and IL-10, on OPC differentiation and immune characteristics. Using primary cultures, ELISA, and immunofluorescence stainings, we assesed differentiation capacity, phagocytic activity, major histocompatibility complex (MHC)-II expression, and cytokine secretion. We observed that the anti-inflammatory milieu (IL4 and IL10) reduced both OPC differentiation and immune functions. Conversely, exposure to TNF-α led to intact differentiation, increased phagocytic activity, high levels of MHC-II expression, and cytokines secretion. Those effects were attributed to signaling via TNF-receptor-2 (TNFR2) and counteracted the detrimental effects of IFNγ on OPC differentiation. Our findings suggest that a pro-regenerative, permissive inflammatory environment is needed for OPCs to execute both regenerative and immune-modulatory functions.

Keywords: multiple sclerosis, oligodendrocyte, OPC, remyelination, TNF, anti-inflammatory cytokines, pro-inflammatory cytokines

Introduction

Multiple sclerosis (MS) is a chronic autoimmune disorder with a heterogeneous range of clinical manifestations. These manifestations span from a relapsing course (relapsing-MS; rMS) to a progressively deteriorating clinical course (progressive-MS; pMS). The rMS form is characterized by a systemic adaptive autoimmune reaction against the myelin of the central nervous system (CNS), and peripheral immune cell infiltration, resulting in demyelination[1–3]. Conversely, in pMS the inflammation is mainly compartmentalized behind a relatively closed blood-brain barrier (BBB)[2–4].

Remyelination failure is one of the main obstacles in MS and is considered a major therapeutic challenge[5,6]. For remyelination to be effective, oligodendrocyte progenitor cells (OPC) need to be activated, migrate into the demyelinated lesions, and differentiate into mature myelinating oligodendrocytes[7]. For many years, OPCs were exclusively regarded for their pro-myelinating potential[8,9]. However, the recognition of the immune-modulatory properties of OPCs nowadays is well-established[10–12]. OPC can secrete cytokines and chemokines such as C-C Motif Chemokine Ligand 2 (CCL2) and Tumor necrosis factor α (TNFα) in response to injury[13,14], they can present antigen via major histocompatibility complex(MHC)-I and -II [12,14,15], affect T-cell proliferation[12,14,16], and phagocytose myelin debris[12].

While OPCs' immune functions are now acknowledged, few studies have explored the impact of an inflammatory environment on these functions, with the majority primarily concentrating on their pro-regenerative roles. Evidence indicates OPCs' pro-regenerative effects in anti-inflammatory settings: interleukin(IL)4 delivery enhanced white matter integrity post-stroke[17], and OPCs' treatment with conditioned media from microglia exposed to IL13 or IL10, but not interferon-γ (IFNγ) and Lipopolysaccharide (LPS), significantly enhanced oligodendrocyte differentiation[18]. In contrast, pro-inflammatory conditions, such as IFNγ are known to impede OPC differentiation[19]. TNFα has been linked to OPCs' differentiation via TNF receptor-2 (TNFR2) signaling[20–22]. Ablation of oligodendroglial TNFR2 exacerbated experimental autoimmune encephalomyelitis (EAE), increased axonal and myelin damage, and reduced remyelination[22]. This also influenced OPC immune responses, as seen in gene and cytokine expression changes[21].

Our previous work indicated that cerebrospinal fluid (CSF) of pMS patients reduced the differentiation and immune functions of OPCs compared to CSF of rMS patients where these functions remained intact[14].

Here, we aimed to examine how pro- and anti-inflammatory environments affect OPCs' immune and differentiation capacities. Based on our prior findings, we hypothesized that a pro-inflammatory milieu would enhance differentiation and immune functions compared to an anti-inflammatory one. Characterizing the distinct effects of these environments *in vitro* could deepen our understanding of MS pathology and facilitate the development of novel therapeutic strategies for MS patients.

Materials and methods

Approvals

The research reported in this study complied with all relevant ethical regulations for animal testing and research and was approved by the Hebrew University Institutional Animal Care and Use Committee(MD-20-16227-1;MD-22-17079-1).

Primary OPC cultures

Primary OPC cultures were isolated from naïve P0-1 neonatal C57/BL6 mice cortices as previously described[23]. After shaking, fibroblast growth factor-2(FGF-2;20ng/mL;R&D systems) and Platelet-derived growth factor(PDGF-AA;20ng/mL;R&D systems) were added to cultures daily unless otherwise specified. Assessments of OPC marker NG2 on day two revealed 91.7±1.6% NG2+ cells.

Cytokines assays

Mouse primary OPCs were supplemented with PDFG-AA and FGF-2 for the first 48 hours after shaking. Then, media were replaced, and recombinant mouse pro- or anti- inflammatory cytokines were added into the OPC media as specified in each assay. Recombinant mouse pro-inflammatory cytokines included IFNγ(Cat#315-05,PeproTech,10ng/ml) and TNFα(Cat#315-01A,PeproTech,200ng/ml). Recombinant mouse anti-inflammatory cytokines included IL4(Cat#214-14,PeproTech,10ng/ml) and IL10(Cat#210-10,PeproTech,100ng/ml).

Differentiation assays

48 hours after shaking, media were replaced, and combinations of Triiodothyronine(T3;60ng/mL;Sigma-Aldrich) and pro- or anti- inflammatory cytokines were added into the OPC media for seven days.

MHC assays

Myelin oligodendrocyte glycoprotein(MOG)-35-55 peptide(Sigma-Aldrich,100µg/ml), IFNγ(10ng/ml;Cat#315-05,PeproTech) and pro- or anti- inflammatory cytokines were supplemented into mouse primary OPC media for 72 hours to allow transcription and translation of MHC-II pathway mediators and permit for expression time, as described previously[14].

Latex beads phagocytosis

OPCs were stimulated using IFNγ(10ng/ml) in a combination of pro- and anti- inflammatory cytokines for 24 hours. Then, latex beads were added for two hours(Cat#L3030-1ML,3µl/1mL,Sigma-Aldrich). Following two hours, cells were immune stained for NG2 marker. Uptake analysis was quantified manually in a blinded manner.

TNFR2 neutralization assays

Mouse primary OPCs were treated with neutralizing antibody against TNFR2(anti-TNFR2; Cat#50128-RN204,RRID:AB_2860178,Sino Biological,6ug/mL) for 1 hour prior to exposure to pro-inflammatory cytokines, as previously described[24]. PBS served as control.

Immunostaining

For cell surface markers, staining was performed on living cells followed by fixation[25]. Anti-NG2(Cat#AB5320, RRID:AB_91789, Millipore, 1:100) was used to identify OPCs, anti-O1(Cat#MAB1327, RRID:AB_357618, R&D System, 1:100) for mature oligodendrocytes, and anti-IBL- 5/22(Cat#sc-59,322, RRID:AB_831551, Santa Cruz Biotechnology, 1:100) for evaluation of MHC-II expression. Goat anti-rabbit Alexa Fluor-488(Cat#A-11034, RRID:AB_2576217, Invitrogen, 1:200), goat anti-mouse Alexa Fluor-488(Cat#A-11001, RRID:AB_2534069, Invitrogen, 1:200), and goat anti-rat Alexa Fluor-555(Cat#A-21434, RRID:AB_141733, Invitrogen, 1:200) were used as secondary antibodies appropriately. Nuclei were counterstained with 4′,6-diamidino-2-phenylindole(DAPI; Cat#H-1200, RRID:AB_2336790, Vector Laboratories). Quantification was performed using ImageJ software(NIH,https://imagej.nih.gov/ij,RRID:SCR_003070) by measuring positively stained cells relative to total DAPI. Quantifications are represented as mean percentages from total DAPI+ cells ±SD and are from at least 15 random fields captured in 3 or more independent experiments.

Assessment of oligodendrocyte morphological complexity by fractal dimension and skeleton analyses

Fractal dimension(FD) analysis was performed to evaluate morphological complexity of oligodendrocytes following differentiation essays, as described previously[14]. A numerical value close to 1 represents cells with low morphological complexity. This procedure was performed for >25 cells per condition for each independent experiment.

Protein concentration measurements

Enzyme-linked immunosorbent-assay(ELISA) was used to measure proteins secretion in culture superannuants following the manufacturer's protocol. The following proteins were measured: CCL2(Cat#DY479-05, R&D System,), IL1b(Cat#DY401-05,R&D System), IL6(Cat#431304, BioLegend; a generous gift of Prof. Talma Brenner), and IL10(Cat#431414, BioLegend; a generous gift of Prof. Talma Brenner). IL1b levels were undetected among all groups.

Statistical analyses

Unpaired two-tailed Student's t-test, one-way ANOVA with Tukey's multiple comparisons post hoc test, and two-way ANOVA with Tukey's multiple comparisons post hoc test were performed. Specific tests are noted in figure legends with significance level annotations. Values are provided as mean ± SD, or as described for each figure. All error bars represent standard deviation. Each assay was repeated independently at least three times.

Results

1. OPC differentiation is impeded by anti-inflammatory cytokines and IFNγ, while preserved by TNFα

Initially, we studied the effect of pro- and anti-inflammatory mediators on OPCs' ability to differentiate. Mouse primary OPCs were cultured with T3(60ng/ml)[26] and pro- or anti-inflammatory cytokines for seven days. Differentiation was assayed using the mature oligodendrocyte marker O1[27](Figure1a-b). IL4 and IL10, representing anti-inflammatory mediators, significantly decreased OPC differentiation into mature oligodendrocytes(26.39±3.32% and 24.15±3.48%, respectively, Figure1b-c), compared to T3 alone(63.1±4.64%, p<0.0001, Figure1b-c). A similar inhibitory effect was observed upon exposure to IFNγ, a pro-inflammatory mediator(25.74±3.37%, Figure1b-c). The exposure of OPCs to TNFα led to intact differentiation levels(60.9±4.63%) compared to T3 alone(p=0.0634, Figure1b-c). Moreover, exposure of OPC to both TNFα and IFNγ together led to higher levels of differentiation(55.32±4.19%) compared to exposure to IFNγ alone(p<0.0001).

Analysis for NG2 expression, an OPC marker, revealed significantly higher expression upon exposure to IFNγ(65.63±3.96%), IL4(64.24±4.63%), or IL10(65.07±5.38%), compared with OPCs exposed to T3 alone, TNFα, or TNFα and IFNγ together(28.55±6.09%, 30.13±3.1%, and 35.43±6.18, respectively, p<0.0001, Figure1d).

Using fractal dimension and skeleton analyses, we performed combined quantification of the morphological pattern of O1+ oligodendrocytes upon culture with pro- and anti-inflammatory mediators for seven days. Oligodendrocytes cultured with pro-inflammatory mediators(TNFα and IFNγ) had lower morphological complexity, represented by a significantly lower FD index, than OPCs exposed to anti-inflammatory mediators(IL4 and IL10, Figure1e-k). Oligodendrocytes exposed to pro-inflammatory cytokines also had lower end points/cell index and branch length/cell compared to those exposed to anti-inflammatory cytokines(Figure1l-m). Remarkably, simultaneous exposure of OPCs to TNFα and IFNγ resulted in higher levels of end points/cell index and branch length/cell compared to their separate applications(Figure1f,g,j,l,m).

Our findings imply that an anti-inflammatory milieu might hinder OPC differentiation capabilities and impair the remyelination process in the inflamed CNS. Additionally, TNFα

seems to preserve OPC differentiation and may play an essential role in the delicate interplay between immunological function and regenerative abilities of OPCs in MS.

2. Anti-inflammatory cytokines impede OPC phagocytic capability, MHC-II expression, and cytokines secretion

2.1. Anti-inflammatory cytokines impede OPC phagocytic capability

The CNS environment of MS patients contains inflammatory mediators and myelin degradation products, which are known to influence MS inflammatory cortical activity, activation of CNS innate immune cells, and OPC remyelination ability. Immune cells, primarily microglia, are responsible for clearing myelin debris, which contains proteins that inhibit OPC differentiation. Our prior study demonstrated that the main differences in OPC exposed to CSF from rMS or pMS patients are related to pathways involved in immune response, specifically in phagocytosis[14]. This is further substantiated by a study demonstrating the phagocytic abilities of OPCs[12]. In this regard, we decided to investigate whether and which anti- or pro-inflammatory cytokines could affect this ability.

OPCs were initially stimulated with IFNγ(10ng/ml), then pro-inflammatory(TNFα) or anti-inflammatory cytokines(IL4 and IL10) were added for 24 hours. Afterward, latex beads were added to the cultures for 2 hours, followed by immunofluorescence staining(Figure2a). Simultaneous exposure of OPCs to TNFα and IFNγ significantly enhanced their phagocytic ability(36.29±2.81%) compared to IFNγ alone(28.88±2.86%), IL4 and IFNγ(23.61±2.51%), IL10 and IFNγ(17.37±2.13%), or both IFNγ, IL4, and IL10(19.12±3.78%, p<0.0001, Figure2b-c).

These results indicate that TNFα amplifies OPC-mediated clearance of myelin debris more than anti-inflammatory settings.

2.2. Reduced MHC-II expression in OPC exposed to anti-inflammatory cytokines

Antigen presentation is crucial for T-cell response. As we previously demonstrated that MHC-II expression in OPC is reduced upon exposure to CSF from pMS patients compared to rMS patients[14], we aimed to examine the influence of pro- and anti-inflammatory cytokines on this OPCs' ability to express MHC-II.

OPCs were cultured with MOG peptide(100ug/mL) and stimulated with IFNγ(10ng/ml). In addition, cells were simultaneously exposed to pro-inflammatory(TNFα) or

anti-inflammatory(IL4 and IL10) cytokines for 72 hours. We then assessed MHC-II expression(I-A/E regions in C57/BL6 mice) using immunofluorescence staining(Figure3a).

Exposure of OPCs to anti-inflammatory cytokines(IL4 and IL10) significantly reduced MHC-II expression(25.8±4.66% and 18.47±2.65%, respectively) compared to the exposure to IFNγ alone(33.63±3.6%, p<0.0001, Figure3b-c). Simultaneous exposure to IL4 and IL10 showed no synergistic effect(Figure3b). Notably, the simultaneous exposure of OPCs to two major pro-inflammatory mediators, TNFα and IFNγ, significantly enhanced MHC-II expression(55.23±3.22%) compared to IFNγ alone(33.63±3.6%), IL4(25.8±4.66%), or IL10(18.47±2.65%, p<0.0001, Figure3b-c).

2.3. Increased secretion of CCL2, IL6, and IL10 upon exposure of OPCs to TNFα

To better understand the effect of the environment on OPCs' immune phenotype, we studied the secretion of CCL2, IL6, and IL10 in OPCs' differentiation cultures upon exposure to pro- or anti-inflammatory cytokines for seven days(timeline is depicted in Figure1a).

First, we measured CCL2 levels in OPC cultures. Exposure of OPCs to TNFα led to higher levels of CCL2(953.75±87.46pg/mL) compared to T3 alone(302.84±86.32pg/mL, p<0.0001), IFNγ(799.18±46.99pg/mL, p=0.0281), IL10(610.37±154.64pg/mL, p<0.0001) or IL4(477.93±115.14pg/mL, p<0.0001, Figure4a). The simultaneous exposure of OPCs to TNFα and IFNγ led to higher levels of CCL2 secretion(1164.05±137.99pg/mL) compared to TNFα or IFNγ alone(p=0.0012 and p<0.0001, respectively, Figure4a).

The pro-inflammatory cytokine IL6 is known for its harmful effect on the inflamed CNS. The highest levels of IL6 were measured in cultures exposed simultaneously to TNFα and IFNγ(25.19±1.32pg/mL, Figure4b). The exposure to TNFα alone led to lower levels of IL6 secretion(13.63±1.76pg/mL) compared to IFNγ(20.86±2.13pg/mL, p<0.0001) but higher compared to the exposure to IL4, IL10, or T3 alone(5.02±1.13pg/mL, 3.74±2.07pg/mL, and 10.08±1.88pg/mL, respectively, p<0.0001, p<0.0001, and p<0.001, respectively, Figure4b).

IL10 was also measured in supernatants of OPC cultures upon exposure to a pro- or anti-inflammatory environment. The highest levels of IL10 were measured in supernatants of T3 alone cultures(54.5±6.14pg/mL, Figure4c). The exposure to TNFα led to higher levels of IL10 secretion(24.69±2.55pg/mL) compared to IFNγ(14.81±2.07pg/mL, p=0.0015). The simultaneous exposure of OPCs to TNFα and IFNγ led to higher levels of IL10 secretion(39.85±4.15pg/mL) compared to the exposure to TNFα or IFNγ alone(24.69±2.55pg/mL and 14.81±2.07pg/mL, respectively, p<0.0001, Figure4c).

These results suggest that TNFα potentially activates OPCs, promoting neuroprotection by enhancing the regenerative capabilities of OPCs and attracting immune cells to the lesion site.

3. TNFα-mediated differentiation, but not MHC-II expression, is dependent on TNFR2

TNFα, a multifaceted cytokine, is abundantly present in the serum, CSF, and active lesions of MS patients[28,29]. Signaling can occur through two receptors: TNFR1, which primarily promotes neurotoxicity, and TNFR2, which promotes neuroprotection and reparative effects[30]. TNFR2 plays a role in OPC differentiation and has been the focus of research in recent years[20–22]. To study the effects of TNFR2 on OPCs' immune and differentiation capabilities *in vitro*, we used a neutralizing antibody against TNFR2.

3.1. TNFR2 neutralization impairs TNFα-mediated OPC differentiation

We aimed to examine whether TNFR2 plays a role in the differentiation observed following TNFα exposure. Mouse primary OPCs were pre-treated with a neutralizing antibody directed to TNFR2 or PBS for one hour. Then, cultures were treated with T3[26] and pro-inflammatory cytokines for seven days.

In the anti-TNFR2 group, both exposure to TNFα alone and simultaneous exposure to TNFα and IFNγ led to significantly lower levels of differentiation compared to the control group(TNFα alone: 63.02±1.92% vs. 31.39±0.37%, p=0.0006; TNFα and IFNγ: 57.16±5.64% vs. 29.79±2.64%, p=0.0006, Figure5a,b). In OPCs exposed to IFNγ or T3 alone, there was no significant difference in O1+ cell percentages between the anti-TNFR2 group and the control group(Figure5a,b). Analysis for NG2 expression, an OPC marker, revealed significantly higher NG2 levels among OPCs cultured with TNFα alone or TNFα together with IFNγ, which were pre-treated with anti-TNFR2(Figure5c).

Upon stimulation with either TNFα or both TNFα and IFNγ simultaneously, anti-TNFR2 pre-treatment led to notably reduced morphological complexity. However, no such effect was observed in the T3 alone or IFNγ groups(Figure5d-j).

These observations support that TNFR2 signaling in OPC is essential for OPC differentiation.

3.2. TNFR2 neutralization does not alter MHC-II expression in OPC

To elucidate TNFR2's role in OPC immunomodulation, we assessed its effect on MHC-II expression in OPCs following simultaneous stimulation with TNFα and IFNγ. Mouse

primary OPCs were pre-treated with an anti-TNFR2 neutralizing antibody or PBS for one hour, followed by culture with MOG peptide, IFNγ, and TNFα.

Pre-treating OPCs with anti-TNFR2 resulted in a significant difference between exposure to IFNγ alone and combined exposure to TNFα and IFNγ($33.71\pm1.12\%$ vs. $55.42\pm0.46\%$, $p<0.0001$, Figure5k-l), mirroring the results in non-pre-treated OPCs(Figure3b-c, Figure 7k-l). MHC-II expression was consistent between control and pre-treated OPCs across all conditions, and no significant difference was observed(Figure3b-c, Figure5k-l).

3.3. TNFR2 neutralization increases the secretion of pro-inflammatory cytokines

Last, we aimed to investigate the role of TNFR2 in the reciprocal balance of immune function and the regenerative role of OPCs in the inflamed CNS. Thus, we measured CCL2, IL6, and IL10 levels in the superannuants of OPCs' differentiation cultures. Mouse primary OPCs were pre-treated with a neutralizing antibody directed to TNFR2 or PBS for one hour. Then, cultures were treated with T3[26] and pro-inflammatory cytokines for seven days. Then, superannuants were collected and evaluated using ELISA.

In the TNFR2-neutralized group, OPCs simultaneously exposed to TNFα and IFNγ secreted higher levels of CCL2 compared to the control group(anti-TNFR2: 1223.11 ± 43.07pg/mL; control: 1088.49 ± 55.63pg/mL; $p=0.0398$). Elevated secretion levels were similarly observed for IL6 and IL10 in the anti-TNFR2 group(Figure5m-o).

Together, these findings imply that TNFR2 plays a pivotal role in both the regenerative and immunomodulatory activities of OPCs within the inflamed CNS.

Discussion

In the current study, our objective was to delineate the influence of both pro- and anti-inflammatory environments on the immunological and differentiation functions of OPCs. We observed that exposure to an anti-inflammatory environment, represented by IL4 and IL10, resulted in decreased OPC differentiation. Interestingly, this effect was comparable to that induced by IFNγ, a predominant pro-inflammatory cytokine associated with MS. Moreover, exposure to IL4 and IL10 corresponded with reduced phagocytic activity, MHC-II expression, and cytokine secretion. Conversely, OPCs exposed to TNFα exhibited enhanced differentiation and robust immune functional capabilities. Those effects were attributed to signaling via TNFR2 and counteracted the detrimental effects of IFNγ on OPC differentiation. Our results support the hypothesis that the optimal functionality of OPCs requires a pro-regenerative inflammatory environment. Additionally, our results suggest the potential central role of TNFα and TNFR2 in orchestrating the equilibrium between the regenerative and immunological roles of OPCs within the inflamed CNS. These insights could lead to the development of novel targeted therapeutic approaches for MS.

For decades, the prevailing belief was that effective remyelination necessitates an anti-inflammatory environment. This belief has roots in some experimental findings. In an animal model examining stroke, the administration of IL4 was observed to enhance white matter integrity[17]. Additionally, oligodendrocyte differentiation was enhanced when exposed to conditioned media from microglia treated with anti-inflammatory cytokines(IL10 and IL13)[18]. Nonetheless, it is essential to consider that this effect was compared with the effects of IFNγ and LPS, both recognized as detrimental regulators of OPC differentiation. A critical observation here is that these studies offered indirect insights and were conducted under varying conditions, potentially obscuring a direct understanding of the effects on OPCs.

Our data suggest that a conducive permissive environment is essential for the effective functionality of the remyelinating cell. This may encompass pro-regenerative inflammatory elements, transitioning the tissue from a restrictive to a permissive environment. The beneficial role of inflammation in promoting remyelination has been substantiated by extensive research. For instance, immune cells facilitate the removal of myelin debris, which contains proteins that inhibit OPC differentiation[31–33]. Moreover, evidence suggests that remyelination occurs more frequently in active inflammatory MS lesions as compared to immunologically inactive plaques[34–37]. This is further supported by animal models of chronic demyelination that have shown that OPCs are capable of effective remyelination only when acute inflammation is

induced[38,39]. We demonstrated in a previous study that exposure to CSF from rMS patients, which is typically characterized by heightened immune activity, led to enhanced levels of OPC differentiation, as compared to exposure to CSF from pMS patients, wherein the inflammatory milieu is relatively limited[14]. Additionally, pro-inflammatory milieu has been observed to promote OPC differentiation via interactions with activated microglial cells and lymphocyte cytokine secretion[40]. Also, evidence indicates that experimental depletion of macrophages[41] and B or T cells[42,43] leads to remyelination impairment. Together with our results, these suggest that remyelination depends on an appropriate immune response controlled in time, space, and intensity[44,45].

Over recent years, significant progress has been made in elucidating the immune-modulatory function of OPCs. OPC have been demonstrated to play an active role in the inflammatory responses observed in a range of neurological disorders[10–12]. Following injury, OPC are activated by inflammatory signals, leading them to secrete IL1β and CCL2. This facilitates their mobilization to the site of injury, subsequently augmenting the post-injury inflammatory environment[10,13,43,46,47]. Further investigations have revealed that OPCs possess phagocytic abilities[12] and have the capability to present antigens through MHC-I and -II[12,14,15]. We demonstrated in a prior study that OPC exposed to CSF from rMS patients exhibited augmented immune-modulatory functions(such as MHC-II expression, NFκB activation, cytokines secretion, and T-Cell activation and proliferation) compared to CSF from pMS patients where these functions appeared compromised[14]. This suggests that an increased inflammatory response of OPC is inversely associated with the disease severity. Furthermore, in another study, we showed that merely pushing OPC to differentiate while suppressing their immune functions does not adequately address the remyelination failure in chronic MS[23]. These highlight the importance of the delicate balance between OPCs' immune and pro-myelinating functions and underscore the need for a pro-inflammatory supporting environment for OPCs to execute their pro-myelinating and immune-modulatory roles.

In line with these concepts, our current findings spotlighted the potential protective and reparative effects of TNFα on OPC in the inflamed CNS. Exposure to TNFα not only preserved the differentiation potential of OPCs but also sustained their immune-modulatory functions. Remarkably, TNFα appeared to counteract the detrimental effects of IFNγ on OPC differentiation, implying its potential beneficial role in remyelination within the context of MS. These results align with recent insights regarding the role of TNFα and TNFR2 in OPC homeostasis. A recent study showed that EAE mice lacking oligodendroglial TNFR2 exhibited peripheral immune cell infiltration, increased demyelination, and impaired remyelination

compared to wild-type[22]. Another research indicated that TNFR2 modulates OPCs' pro-inflammatory phenotype[21]. The absence of TNFR2 exacerbates the immune-modulatory and inflammatory response in OPCs post-inflammatory induction, consequently reducing their proliferation and differentiation potential[21]. In a recent study by Fiedler et al., the dual modulation of TNFR1 and TNFR2 in an EAE model yielded promising results, effectively mitigating EAE symptoms while decreasing demyelination and axonal degeneration[48]. It is important to note that the pivotal function of TNFα in maintaining a regenerative milieu within the CNS is further underscored by instances where patients treated with anti-TNF agents developed demyelinating syndromes[49]. These findings underscore the significance of immune signaling in determining OPC functions and indicate the promising therapeutic potential of targeting the TNFα and TNFR2 pathways in MS.

In conclusion, our study sheds lights on the nuanced interplay between pro- and anti-inflammatory environments and the subsequent impact on OPCs' immunological and differentiation functionalities. Our findings point towards the essentiality of a pro-regenerative inflammatory environment, emphasizing the pivotal role of TNFα and TNFR2 in harmonizing the balance between the regenerative and immunological tasks of OPCs in the inflamed CNS. These findings may pave the way for innovative therapies that encapsulate the full spectrum of OPC functions in MS.

References

1. Dendrou CA, Fugger L, Friese MA. Immunopathology of multiple sclerosis. *Nat Rev Immunol*. Published online 2015. doi:10.1038/nri3871

2. Torkildsen O, Myhr KM, Bø L. Disease-modifying treatments for multiple sclerosis - a review of approved medications. *Eur J Neurol*. Published online 2016. doi:10.1111/ene.12883

3. Lassmann H, Van Horssen J, Mahad D. Progressive multiple sclerosis: Pathology and pathogenesis. *Nat Rev Neurol*. Published online 2012. doi:10.1038/nrneurol.2012.168

4. Meinl E, Krumbholz M, Derfuss T, Junker A, Hohlfeld R. Compartmentalization of inflammation in the CNS: A major mechanism driving progressive multiple sclerosis. *J Neurol Sci*. Published online 2008. doi:10.1016/j.jns.2008.06.032

5. Lassmann H, Brück W, Lucchinetti C, Rodriguez M. Remyelination in multiple sclerosis. In: *Multiple Sclerosis*. ; 1997. doi:10.1177/135245859700300213

6. Bramow S, Frischer JM, Lassmann H, et al. Demyelination versus remyelination in progressive multiple sclerosis. *Brain*. Published online 2010. doi:10.1093/brain/awq250

7. Gruchot J, Weyers V, Göttle P, et al. The Molecular Basis for Remyelination Failure in Multiple Sclerosis. *Cells*. 2019;8(8):825. doi:10.3390/cells8080825

8. Hughes EG, Kang SH, Fukaya M, Bergles DE. Oligodendrocyte progenitors balance growth with self-repulsion to achieve homeostasis in the adult brain. *Nat Neurosci*. Published online 2013. doi:10.1038/nn.3390

9. Chang A, Nishiyama A, Peterson J, Prineas J, Trapp BD. NG2-positive oligodendrocyte progenitor cells in adult human brain and multiple sclerosis lesions. *Journal of Neuroscience*. Published online 2000. doi:10.1523/jneurosci.20-17-06404.2000

10. Akay LA, Effenberger AH, Tsai LH. Cell of all trades: oligodendrocyte precursor cells in synaptic, vascular, and immune function. *Genes Dev*. Published online 2021. doi:10.1101/gad.344218.120

11. Boccazzi M, Raffaele S, Fumagalli M. Not only myelination: the immune-inflammatory functions of oligodendrocytes. *Neural Regen Res*. 2022;17(12):2661. doi:10.4103/1673-5374.342678

12. Falcão AM, van Bruggen D, Marques S, et al. Disease-specific oligodendrocyte lineage cells arise in multiple sclerosis. *Nat Med*. Published online 2018. doi:10.1038/s41591-018-0236-y

13. Moyon S, Dubessy AL, Aigrot MS, et al. Demyelination causes adult CNS progenitors to revert to an immature state and express immune cues that support their migration. *Journal of Neuroscience*. Published online 2015. doi:10.1523/JNEUROSCI.0849-14.2015

14. Zveik O, Fainstein N, Rechtman A, et al. Cerebrospinal fluid of progressive multiple sclerosis patients reduces differentiation and immune functions of oligodendrocyte progenitor cells. *Glia*. 2022;70(6):1191-1209. doi:10.1002/glia.24165

15. Kirby L, Jin J, Cardona JG, et al. Oligodendrocyte precursor cells present antigen and are cytotoxic targets in inflammatory demyelination. *Nat Commun*. Published online 2019. doi:10.1038/s41467-019-11638-3

16. Nishri Y, Hampton D, Ben-Shushan E, et al. Continuous Immune-Modulatory Effects of Human Olig2+ Precursor Cells Attenuating a Chronic-Active Model of Multiple Sclerosis. *Mol Neurobiol*. 2020;57(2). doi:10.1007/s12035-019-01802-7

17.   Zhang Q, Zhu W, Xu F, et al. The interleukin-4/PPARγ signaling axis promotes oligodendrocyte differentiation and remyelination after brain injury. *PLoS Biol*. 2019;17(6):e3000330. doi:10.1371/journal.pbio.3000330

18.   Miron VE, Boyd A, Zhao JW, et al. M2 microglia and macrophages drive oligodendrocyte differentiation during CNS remyelination. *Nat Neurosci*. 2013;16(9):1211-1218. doi:10.1038/nn.3469

19.   Starost L, Lindner M, Herold M, et al. Extrinsic immune cell-derived, but not intrinsic oligodendroglial factors contribute to oligodendroglial differentiation block in multiple sclerosis. *Acta Neuropathol*. Published online 2020. doi:10.1007/s00401-020-02217-8

20.   Arnett HA, Mason J, Marino M, Suzuki K, Matsushima GK, Ting JPY. TNFα promotes proliferation of oligodendrocyte progenitors and remyelination. *Nat Neurosci*. Published online 2001. doi:10.1038/nn738

21.   Desu HL, Illiano P, Choi JS, et al. TNFR2 signaling regulates the immunomodulatory function of oligodendrocyte precursor cells. *Cells*. 2021;10(7). doi:10.3390/cells10071785

22.   Madsen PM, Desu HL, Vaccari JP de R, et al. Oligodendrocytes modulate the immune-inflammatory response in EAE via TNFR2 signaling. *Brain Behav Immun*. Published online 2020. doi:10.1016/j.bbi.2019.11.017

23.   Ganz T, Zveik O, Fainstein N, et al. Oligodendrocyte progenitor cells differentiation induction with MAPK/ERK inhibitor fails to support repair processes in the chronically demyelinated CNS. *Glia*. Published online 2023. doi:10.1002/glia.24453

24.   Zhang Y, Yang C, Ge S, Wang L, Zhang J, Yang P. EphB4/ TNFR2/ERK/MAPK signaling pathway comprises a signaling axis to mediate the positive effect of TNF-α on osteogenic differentiation. *BMC Mol Cell Biol*. 2020;21(1). doi:10.1186/s12860-020-00273-2

25.   Zveik O, Rechtman A, Haham N, et al. Sera of Neuromyelitis Optica Patients Increase BID-Mediated Apoptosis in Astrocytes. *Int J Mol Sci*. 2022;23(13):7117. doi:10.3390/ijms23137117

26.   Tosic M, Torch S, Comte V, Dolivo M, Honegger P, Matthieu J -M. Triiodothyronine Has Diverse and Multiple Stimulating Effects on Expression of the Major Myelin Protein Genes. *J Neurochem*. Published online 1992. doi:10.1111/j.1471-4159.1992.tb11009.x

27.   Barateiro A, Fernandes A. Temporal oligodendrocyte lineage progression: In vitro models of proliferation, differentiation and myelination. *Biochim Biophys Acta Mol Cell Res*. Published online 2014. doi:10.1016/j.bbamcr.2014.04.018

28.   Sharief MK, Hentges R. Association between Tumor Necrosis Factor-α and Disease Progression in Patients with Multiple Sclerosis. *New England Journal of Medicine*. 1991;325(7):467-472. doi:10.1056/NEJM199108153250704

29.   Maimone D, Gregory S, Arnason BGW, Reder AT. Cytokine levels in the cerebrospinal fluid and serum of patients with multiple sclerosis. *J Neuroimmunol*. 1991;32(1):67-74. doi:10.1016/0165-5728(91)90073-G

30.   Dostert C, Grusdat M, Letellier E, Brenner D. The TNF Family of Ligands and Receptors: Communication Modules in the Immune System and Beyond. *Physiol Rev*. 2019;99(1):115-160. doi:10.1152/physrev.00045.2017

31.   Kotter MR, Li WW, Zhao C, Franklin RJM. Myelin impairs CNS remyelination by inhibiting oligodendrocyte precursor cell differentiation. *Journal of Neuroscience*. Published online 2006. doi:10.1523/JNEUROSCI.2615-05.2006

32.   Rosenberg SS, Kelland EE, Tokar E, de La Torre AR, Chan JR. The geometric and spatial constraints of the microenvironment induce oligodendrocyte differentiation. *Proc Natl Acad Sci U S A*. Published online 2008. doi:10.1073/pnas.0805640105

33. Ruckh JM, Zhao JW, Shadrach JL, et al. Rejuvenation of regeneration in the aging central nervous system. *Cell Stem Cell*. Published online 2012. doi:10.1016/j.stem.2011.11.019

34. Confavreux C, Vukusic S, Moreau T, Adeleine P. Relapses and Progression of Disability in Multiple Sclerosis. *New England Journal of Medicine*. Published online 2000. doi:10.1056/nejm200011163432001

35. Khademi M, Dring AM, Gilthorpe JD, et al. Intense Inflammation and Nerve Damage in Early Multiple Sclerosis Subsides at Older Age: A Reflection by Cerebrospinal Fluid Biomarkers. *PLoS One*. Published online 2013. doi:10.1371/journal.pone.0063172

36. Patani R, Balaratnam M, Vora A, Reynolds R. Remyelination can be extensive in multiple sclerosis despite a long disease course. *Neuropathol Appl Neurobiol*. Published online 2007. doi:10.1111/j.1365-2990.2007.00805.x

37. Wolswijk G. Oligodendrocyte precursor cells in the demyelinated multiple sclerosis spinal cord. *Brain*. Published online 2002. doi:10.1093/brain/awf031

38. Foote AK, Blakemore WF. Inflammation stimulates remyelination in areas of chronic demyelination. *Brain*. Published online 2005. doi:10.1093/brain/awh417

39. Setzu A, Lathia JD, Zhao C, et al. Inflammation stimulates myelination by transplanted oligodendrocyte precursor cells. *Glia*. Published online 2006. doi:10.1002/glia.20371

40. El Behi M, Sanson C, Bachelin C, et al. Adaptive human immunity drives remyelination in a mouse model of demyelination. *Brain*. 2017;140(4):967-980. doi:10.1093/brain/awx008

41. Kotter MR, Setzu A, Sim FJ, Van Rooijen N, Franklin RJM. Macrophage depletion impairs oligodendrocyte remyelination following lysolecithin-induced demyelination. *Glia*. Published online 2001. doi:10.1002/glia.1085

42. Bieber AJ, Kerr S, Rodriguez M. Efficient central nervous system remyelination requires T cells. *Ann Neurol*. Published online 2003. doi:10.1002/ana.10578

43. Dombrowski Y, O'Hagan T, DIttmer M, et al. Regulatory T cells promote myelin regeneration in the central nervous system. *Nat Neurosci*. Published online 2017. doi:10.1038/nn.4528

44. Baaklini CS, Rawji KS, Duncan GJ, Ho MFS, Plemel JR. Central Nervous System Remyelination: Roles of Glia and Innate Immune Cells. *Front Mol Neurosci*. 2019;12. doi:10.3389/fnmol.2019.00225

45. Schwartz M, Kipnis J. Multiple sclerosis as a by-product of the failure to sustain protective autoimmunity: A paradigm shift. *Neuroscientist*. Published online 2002. doi:10.1177/107385802236966

46. Kang Z, Wang C, Zepp J, et al. Act1 mediates IL-17-induced EAE pathogenesis selectively in NG2 + glial cells. *Nat Neurosci*. Published online 2013. doi:10.1038/nn.3505

47. Kirby L, Castelo-Branco G. Crossing boundaries: Interplay between the immune system and oligodendrocyte lineage cells. *Semin Cell Dev Biol*. Published online 2020. doi:10.1016/j.semcdb.2020.10.013

48. Fiedler T, Fairless R, Pichi K, et al. Co-modulation of TNFR1 and TNFR2 in an animal model of multiple sclerosis. *J Neuroinflammation*. 2023;20(1):100. doi:10.1186/s12974-023-02784-z

49. Mohan N, Edwards ET, Cupps TR, et al. Demyelination occurring during anti-tumor necrosis factor α therapy for inflammatory arthritides. *Arthritis Rheum*. 2001;44(12). doi:10.1002/1529-0131(200112)44:12<2862::AID-ART474>3.0.CO;2-W

Figure legends

Figure 1. **OPC differentiation is impeded by anti-inflammatory cytokines and IFNγ, while preserved by TNFα**

(a) Time-course of differentiation experiments. OPCs were supplemented with PDGF and FGF2 for 48h. Then, each culture was exposed to simultaneously to thyroid hormone(T3) and either anti- or pro-inflammatory cytokines for 7 days and evaluated using OPC marker NG2 and mature oligodendrocyte marker O1,**(b)** Representative immunofluorescence analysis of mature oligodendrocyte marker O1 and progenitor marker NG2 on day 7(scale bar = 50 μm),**(c)** Exposure to IFNγ(25.74±3.37%) or anti-inflammatory cytokines(IL4: 26.39±3.32%; IL10: 24.15±3.48%) significantly decreased OPC differentiation into mature oligodendrocytes ac compared to T3 alone(63.1±4.64%), TNFα(60.9±4.63%), or TNFα and IFNγ together(55.32±4.19%); n=9 for each group,**(d)** OPCs exposed to IFNγ(65.63±3.96%) or anti-inflammatory cytokines(IL4: 64.24±4.63%; IL10: 65.07±5.38%) expressed higher levels of progenitor marker NG2 after 7 days in culture in comparison to T3 alone(28.55±6.09%), TNFα(30.13±3.1%), or TNFα and IFNγ together(35.43±6.18%); n=5 for each group,**(e-j)** Representative images of O1+ oligodendrocytes and cell outline with the appropriate fractal dimension(FD) value(scale bar = 30 μm),**(k)** Higher levels of morphological complexity among cells exposed to anti-inflammatory cytokines(IL4: 1.45±0.01; IL10: 1.56±0.08) or T3(1.51±0.07) compared with IFNγ(1.29±0.05), TNFα(1.39±0.04), or TNFα and IFNγ together(1.4±0.06),**(l)** Skeleton analysis resulted in higher endpoints/cell among cells exposed to anti-inflammatory cytokines(IL4: 31.32±4.21; IL10: 61.52±6.74) or T3(44.8±10.41) compared with IFNγ(15.6±2.8), TNFα(13.68±2.81), or TNFα and IFNγ together(22.5±3.88),**(m)** Skeleton analysis resulted in higher branches length/cell among cells exposed to anti-inflammatory cytokines(IL4: 1226.25±233.7; IL10: 1678.12±435.06) or T3(1382.12±401.8) compared with IFNγ(766.4±153.8), TNFα(383.9±98.9), or TNFα and IFNγ together(1023.3±149.46). Significance was determined by one-way ANOVA analysis followed by Tukey's multiple comparison analysis and by unpaired two-tailed student's t test(p * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001, ns = not significant). Data are presented as means±SD. Each assay was repeated independently at least three times.

Figure 2. Anti-inflammatory cytokines impede OPC phagocytic capability

(a) Time-course experiments of assessment of phagocytic capability. Primary OPCs were cultured under PDGF and FGF conditions(daily dose of 20 ng/mL) for 48 hours. After 48 hours, OPCs were stimulated with IFNγ(10ng/ml), then pro-inflammatory(TNFα) or anti-inflammatory cytokines(IL4 and IL10) were added for 24 hours. Afterward, latex beads were added to the cultures for 2 hours, followed by evaluation of phagocytosed latex beads out of NG2+ cells using immunofluorescence staining.,

(b) Simultaneous exposure of OPCs to TNFα and IFNγ significantly enhanced their phagocytic ability(36.29±2.81%) compared to IFNγ alone(28.88±2.86%), IL4(23.61±2.51%), IL10(17.37±2.13%), or both IL4 and IL10(19.12±3.78%); n=7 for each group,**(c)** Representative immunofluorescence analysis of NG2+ OPCs and phagocytosed latex beads upon culture with either anti- or pro-inflammatory cytokines for 24 hours(scale bar = 40 μm). Significance was determined by either student's unpaired *t* test or one-way ANOVA analysis followed by Tukey's multiple comparison analysis(p * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001, ns = not significant). Error bars in all graphs represent SD. Each assay was repeated independently at least three times.

Figure 3. Reduced MHC-II expression in OPC exposed to anti-inflammatory cytokines

(a) Time-course experiments of MHC-II expression. Primary OPCs were cultured under PDGF and FGF conditions(daily dose of 20 ng/mL) for 48 hours. After 48 hours, OPCs were stimulated with IFNγ(10ng/ml), then pro-inflammatory(TNFα) or anti-inflammatory cytokines(IL4 and IL10) were added for 72 hours, followed by evaluation of MHC-II expression out of NG2+ cells using immunofluorescence staining,**(b)** Exposure of OPCs to anti-inflammatory cytokines significantly reduced MHC-II expression(IL4: 25.8±4.66%; IL10: 18.47±2.65%; IL4 and IL10 together: 19.12±3.78%) compared to the exposure to IFNγ alone(33.63±3.6%) or TNFα and IFNγ together(55.23±3.22%); n=5 for each group,**(c)** Representative immunofluorescence analysis of MHC-II expression among OPCs upon culture with either anti- or pro-inflammatory cytokines for 72 hours(scale bar = 40 μm). Significance was determined by either student's unpaired *t* test or one-way ANOVA analysis followed by Tukey's multiple comparison analysis(p * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001, ns = not significant). Error bars in all graphs represent SD. Each assay was repeated independently at least three times.

Figure 4. **Increased secretion of CCL2, IL6, and IL10 upon exposure of OPCs to TNFα**

Using ELIAS analyses, we studied the secretion of CCL2, IL6, and IL10 in the supernatant of OPCs following culture with T3 and either anti- or pro-inflammatory cytokines for 7 days.**(a)** CCL2: T3 alone: 302.84±86.32pg/mL; IFNγ: 799.18±46.99pg/mL; TNFα: 953.75±87.46pg/mL; IL4: 477.93±115.14pg/mL; IL10: 610.37±154.64pg/mL; IFNγ and TNFα: 1164.05±137.99pg/mL; n=9 for each group,**(b)** IL6: T3 alone: 10.08±1.88pg/mL; IFNγ: 20.86±2.13pg/mL; TNFα: 13.63±1.76pg/mL; IL4: 5.02±1.13pg/mL; IL10: 3.74±2.07pg/mL; IFNγ and TNFα: 25.19±1.32pg/mL; n=9 for each group,**(c)** IL10: T3 alone: 54.5±6.14pg/mL; IFNγ: 14.81±2.07pg/mL; TNFα: 24.69±2.55pg/mL; IL4: 25.27±9.2pg/mL; IFNγ and TNFα: 39.85±4.15pg/mL; n=9 for each group. Significance was determined by either student's unpaired *t* test or one-way ANOVA analysis followed by Tukey's multiple comparison analysis($p * \leq 0.05$, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001, ns = not significant). Error bars in all graphs represent SD. Each assay was repeated independently at least three times.

Figure 5. **TNFα-mediated differentiation, but not MHC-II expression, is dependent on TNFR2**

(a) Representative immunofluorescence analysis of mature oligodendrocyte marker O1 and progenitor marker NG2 on day 7 for TNFR2 neutralizations differentiation experiments(scale bar = 50 μm),**(b)** Percentage of O1+ cells out of DAPI on day 7: T3 only: control 62.8±2.0%, a-TNFR2 62.4±0.5%; IFNγ: control 25.15±1.8%, a-TNFR2 24.32±0.9%; TNFα: control 63.02±1.92%, a-TNFR2 31.39±0.37%; IFNγ and TNFα: control 57.16±5.64%, a-TNFR2 29.79±2.64%; n=4 for each group,**(c)** Percentage of NG2+ cells out of DAPI on day 7: T3 only: control 28.55±1.67%, a-TNFR2 28.5±1.34%; IFNγ: control 65.36±1.49%, a-TNFR2 67.54±1.5%; TNFα: control 30.58±2.7%, a-TNFR2 59.1±1.0%; IFNγ and TNFα: control 35.9±5.57%, a-TNFR2 62.39±0.95%; n=4 for each group,**(d-g)** Representative images of O1+ oligodendrocytes and cell outline with the appropriate fractal dimension(FD) value for TNFR2 neutralizations differentiation experiments(scale bar = 30 μm),**(h)** FD index for TNFR2 neutralizations differentiation experiments: T3 only: control 1.51±0.06, a-TNFR2 1.5±0.06; IFNγ: control 1.34±0.04, a-TNFR2 1.31±0.07; TNFα: control 1.37±0.04, a-TNFR2 1.22±0.03; IFNγ and TNFα: control 1.43±0.04, a-TNFR2 1.28±0.02,**(i)** End points/cell index for TNFR2 neutralizations differentiation experiments: T3 only: control 43.8±5.5, a-TNFR2 42.96±5.72;

IFNγ: control 14.2±2.7, a-TNFR2 13.46±3.45; TNFα: control 13.2±2.07, a-TNFR2 8.2±2.02; IFNγ and TNFα: control 22.46±2.96, a-TNFR2 9.5±2.04,**(j)** Branch length/cell index for TNFR2 neutralizations differentiation experiments: T3 only: control 1369.693±225.33, a-TNFR2 1344.12±51.36; IFNγ: control 737.81±124.9, a-TNFR2 760.5±41.9; TNFα: control 364.68±72.17, a-TNFR2 293.76±48.47; IFNγ and TNFα: control 1101.54±80.36, a-TNFR2 322.01±37.78,**(k)** Percentage of MHC-II+ cells out of NG2+ cells on day 3: IFNγ: control 33.71±1.12%, a-TNFR2 32.89±0.61%; IFNγ and TNFα: control 55.42±0.46%, a-TNFR2 54.51±1.2%; n=4 for each group,**(l)** Representative immunofluorescence analysis of MHC-II expression among OPCs upon TNFR2 neutralization and culture with either anti- or pro-inflammatory cytokines for 72 hours(scale bar = 40 μm),**(m)** CCL2 levels for TNFR2 neutralizations differentiation experiments: T3 only: control 290.11±44.59pg/mL, a-TNFR2 286.15±40.32pg/mL; IFNγ: control 835.24±33.9pg/mL, a-TNFR2 847.69±6.02pg/mL; TNFα: control 946.68±29.89pg/mL, a-TNFR2 999.68±82.29pg/mL; IFNγ and TNFα: control 1088.49±55.63pg/mL, a-TNFR2 1223.11±43.07pg/mL, n=4 for each group,**(n)** IL6 levels for TNFR2 neutralizations differentiation experiments: T3 only: control 10.69±2.67pg/mL, a-TNFR2 10.26±1.97pg/mL; IFNγ: control 20.67±0.66pg/mL, a-TNFR2 20.28±1.05pg/mL; TNFα: control 12.82±0.66pg/mL, a-TNFR2 19.94±0.99pg/mL; IFNγ and TNFα: control 26.44±1.46pg/mL, a-TNFR2 32.08±2.22pg/mL, n=4 for each group,**(o)** IL10 levels for TNFR2 neutralizations differentiation experiments: T3 only: control 58.64±2.58pg/mL, a-TNFR2 58.04±3.91pg/mL; IFNγ: control 15.64±1.53pg/mL, a-TNFR2 16.52±1.66pg/mL; TNFα: control 24.23±1.56pg/mL, a-TNFR2 31.75±5.27pg/mL; IFNγ and TNFα: control 43.67±2.44pg/mL, a-TNFR2 50.6±2.97pg/mL, n=4 for each group. Significance was determined by one-way ANOVA analysis followed by Tukey's multiple comparison analysis and by unpaired two-tailed student's t test($p * \leq 0.05$, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001, ns = not significant). Data are presented as means±SD. Each assay was repeated independently at least three times.

Figure 1

Figure 2

a

Replaced media
and cytokines
were added

↓

-2   0   1

PFGF + FGF

↑

Latex beads were
added for 2h

b

c

IFN-γ

TNF-α + IFN-γ

IL4 + IFN-γ

IL10 + IFN-γ

IL10 + IL4 + IFN-γ

Enlargement of a phagocytic OPC

● DAPI ● NG2 ● Latex beads

Figure 3

Figure 4

Figure 5