Selective sweep probabilities in spatially expanding populations

Alexander Stein

Centre for Cancer Genomics and Computational Biology,

Barts Cancer Institute, Queen Mary University of London, London, UK and

Department of Physics, ETH Zurich, Zürich, Switzerland

Ramanarayanan Kizhuttil

Department of Physics, Indian Institute of Science Education and Research, Kolkata, India

Maciej Bak and Robert Noble *

Department of Mathematics, City, University of London, London, UK

Abstract

Evolution during range expansions shapes biological systems from microbial communities and tumours up to invasive species. A fundamental question is whether, when a beneficial mutation arises during a range expansion, it will evade clonal interference and sweep through the population to fixation. However, most theoretical investigations of range expansions have been confined to regimes in which selective sweeps are effectively impossible, while studies of selective sweeps have either assumed constant population size or have ignored spatial structure. Here we use mathematical modelling and analysis to investigate selective sweep probabilities in the alternative yet biologically relevant scenario in which mutants can outcompete and displace a slowly spreading wildtype. Assuming constant radial expansion speed, we derive probability distributions for the arrival time and location of the first surviving mutant and hence find surprisingly simple approximate and exact expressions for selective sweep probabilities in one, two and three dimensions, which are independent of mutation rate. Namely, the selective sweep probability is approximately $(1 - c_{\mathrm{wt}}/c_{\mathrm{m}})^d$, where c_{wt} and c_{m} are the wildtype and mutant radial expansion speeds, and d the spatial dimension. Using agent-based simulations, we show that our analytical results accurately predict selective sweep frequencies in the two-dimensional spatial Moran process. We further compare our results with those obtained for alternative growth laws. Parameterizing our model for human tumours, we find that selective sweeps are predicted to be rare except during very early solid tumour growth, thus providing a general, pan-cancer explanation for findings from recent sequencing studies.

INTRODUCTION

Range expansion – the spatial spread of populations into new regions – is ubiquitous across biological scales and alters the course of evolution in distinct, often profound ways that remain incompletely understood [1]. Among cell populations, evolution during range expansions determines the development and spatial heterogeneity of biofilms [2], tumours [3], mosaicism [4], and normal tissue [5]. At the species level, range expansions influenced human evolution [6] and are of growing importance as climate

---

* robert.noble@city.ac.uk

change forces organisms into new habitats [7, 8]. Prior theoretical and experimental investigations of evolution during range expansion have considered the case in which the wildtype population spreads into new territory much faster than any mutant can displace the wildtype (for example, [9–12]). In this scenario, which is typical of microbial colonies growing *in vitro*, mutants can expand only at the population boundary and selective sweeps are precluded. The alternative case of slow range expansion and strong selection has been unexplored but is widely plausible. Consider, for example, an invasive species that is rapidly adapting to its new environment while gradually displacing a resident competitor, or a bacterial colony whose growth is slowed by sub-inhibitory antibiotic treatment.

Somatic evolution provides especially strong motivation for investigating the likelihood of selective sweeps during slow range expansions. Cancer results from an accumulation of mutations that drive cells to proliferate uncontrollably and to invade surrounding tissue [13]. An early, highly influential model based on colorectal cancer genetics posited a linear mode of evolution in which tumours acquire mutations via sequential selective sweeps [14]. Yet recent advances in multi-region and single-cell sequencing have revealed pervasive genetic intratumour heterogeneity [15, 16]. Although fitter mutant clones arise throughout cancer progression, they apparently seldom achieve selective sweeps, except perhaps in very small tumours [17]. These observations have sparked an ongoing debate about the general nature of tumour evolution, the extent of clonal selection, and the causes of intratumour heterogeneity [18, 19]. Mathematical modelling offers a way to resolve the conflict but, with notable exceptions [20–22], theoretical investigations of cancer initiation and early evolution have either ignored spatial structure [23–25] or have relied on agent-based models [26–29]. The typical observed pattern of very early fixation events followed by branching or effectively neutral intratumour evolution [18, 19] – which has important clinical implications [30–32] – has eluded a general explanation.

Here we use mathematical analysis to explain why beneficial mutations typically fixate only – if at all – in the very early stages of range expansions, even when mutants can displace the wildtype faster than the wildtype expands its range. We focus on a simple model with short-range dispersal throughout the population. By solving our model in the canonical case of constant radial expansion speed we derive exact and sim-

ple approximate expressions for sweep probabilities in one, two and three dimensions. We confirm the accuracy and robustness of our analytical results using extensive agent-based simulations of a spatial Moran process, and we compare outcomes for alternative growth models. We discuss how these findings shed light on the nature of evolution in range expansions in general, and cancer development in particular.

RESULTS

Macroscopic model

We consider a wildtype population that starts expanding spherically at time $t = 0$, such that its radius x_{wt} grows at speed c_{wt}. Focusing on selective sweeps, we consider only advantageous mutations, which we assume spread within the wildtype at speed $c_m > c_{wt}$ (Figure 1). Mutations occur at per-capita rate $\tilde{\mu}$ with each surviving genetic drift with probability ρ. In our analytic model, it is enough to consider the compound parameter $\mu = \rho\tilde{\mu}$, which is the mutation rate conditioned on survival. For brevity, we will refer to μ as the mutation rate unless otherwise mentioned.

For mathematical and biological reasons (see Discussion), we focus on a model with constant radial expansion speeds; later we compare these results with those that pertain for alternative growth models. Various models link propagation speeds to fitness values and migration rates. The most prominent formula is obtained from a reaction-diffusion equation associated with Fisher [33] and Kolmogorov [34]: $c = 2\sqrt{aD}$, where D is the diffusion coefficient carrying information of migration and a is the difference in the local proliferation rate. More recent studies have sought to refine and generalise this result [35–38]. Our main results apply to any model that generates approximately constant speeds.

The first surviving mutation achieves a complete (or "hard" or "classic") selective sweep only if it reaches every part of the wildtype expansion front before a second mutant of equal or greater fitness arises within the wildtype (Figure 1). Otherwise, the outcome is clonal interference or possibly a multiple-origin soft selective sweep if the competing mutations are sufficiently similar [39]. For simplicity, we neglect mutants with fitness values between those of the wildtype and the first mutant, which would

Figure 1. Illustration of the macroscopic model (not to scale) showing the two possible fates of the first surviving mutant (blue) within the wildtype population (orange). In the first case, the mutant reaches every part of the wildtype boundary before further mutants arise in the wildtype, and so achieves a selective sweep. Otherwise, a second (green) mutant arises in the wildtype population and generates clonal interference.

slow the expansion of the first mutant and so reduce rather than negate the selective sweep probability. Neither do we investigate the case of a yet-fitter mutant that arises from the first mutant and achieves a selective sweep.

Sweep probability

The unconditional sweep probability is derived in four steps. First, we introduce the random variable X, the radius of the wildtype population when the first mutant arises, and we compute its probability density $f_X(x)$. Second, we introduce the random variable Y, the distance between the wildtype and mutant origins, and we calculate its probability density conditioned on X, namely $f_Y(y|X = x)$. Third, we derive an expression for the conditional sweep probability $\Pr(\text{sweep}|X = x, Y = y)$. Finally, we marginalize out X and Y to obtain the unconditional sweep probability $\Pr(\text{sweep})$. Here, we focus on the three-dimensional case; analogous results in one and two dimensions are presented in the SI Text.

 Probability that no mutations occur

The following result (proved in SI Text) will be useful in various parts of subsequent derivations.

Claim 1. *The probability that k mutations arise and survive during the time interval $[0, t]$ is Poisson distributed,*

$$P_k = \frac{e^{-\lambda} \lambda^k}{k!} \quad with \quad \lambda = \mu \int_0^t N(s) \, ds, \tag{1}$$

where $N(t)$ is the population size. In particular, the probability that no successful mutant arises is $P_0 = e^{-\lambda}$.

Although it is commonly assumed that mutations are coupled to divisions, it is straightforward to translate between per-capita and per-division mutation rates (see SI text).

Arrival time of the first mutant

In the absence of mutants, the wildtype population in three dimensions grows as $N_{\text{wt}} = \frac{4}{3}\pi x_{\text{wt}}^3 = \frac{4}{3}\pi (c_{\text{wt}} t)^3$. Applying Claim 1, we obtain the probability that no mutations arise in the time interval $[0, t]$,

$$P_0 = e^{-\mu \int_0^t \frac{4}{3}\pi (c_{\text{wt}} s)^3 \, ds} = e^{-\left(\frac{t}{\kappa}\right)^4}, \tag{2}$$

in terms of a characteristic duration $\kappa = \sqrt[4]{\frac{3}{\mu \pi c_{\text{wt}}^3}}$. We identify $1 - P_0$ as the cumulative distribution function for the arrival time T of the first surviving mutant. The probability density of T is then

$$f_T(t) = \frac{d(1 - P_0)}{dt} = \frac{4t^3}{\kappa^4} e^{-\left(\frac{t}{\kappa}\right)^4}, \tag{3}$$

which is the Weibull distribution with shape parameter 4 and scale parameter κ. Substituting $x = c_{\text{wt}} t$, we find that $f_X(x)$, the probability density of the radius of the wildtype population X at the time the first surviving mutant arises, is the Weibull distribution with shape parameter 4 and scale parameter $\theta = \sqrt[4]{\frac{3 c_{\text{wt}}}{\pi \mu}}$. It follows that $\mathbb{E}[X] \approx 0.91 \, \theta$ and $\text{Var}[X] \approx 0.065 \, \theta^2$. Analogous calculations in one and two dimensions yield similar Weibull distributions (SI text; Figure 2A-C).

 Location of the first mutant

Next, we compute the probability density for the distance Y between the first surviving mutant and the centre of the wildtype population. Since mutants arise in proportion to the number of wildtype cells, we have $f_Y(y|X = x)\,dy \propto D(y)$, where $D(y)$ is the number of cells at distance y. $D(y)$ corresponds to the infinitesimal shell, $D(y) = 4\pi y^2\,dy\,\mathbf{1}\{y \leq x\}$, where the last term is an indicator function that defines the boundary of the wildtype population. After normalisation, we obtain

$$f_Y(y|X = x) = \frac{3y^2}{x^3}\mathbf{1}\{y \leq x\}. \tag{4}$$

We calculate the unconditional probability density of Y by marginalizing out X,

$$f_Y(y) = \int_0^\infty f_Y(y|X = x)f_X(x)\,dx = \frac{3y^2}{\theta^3}\Gamma\left(\frac{1}{4}, \frac{y^4}{\theta^4}\right), \tag{5}$$

where $\Gamma(a, z) = \int_z^\infty t^{a-1}e^{-t}\,dt$ is the incomplete gamma function. We then find $\mathbb{E}[Y] \approx 0.68\,\theta$ and $\mathrm{Var}[Y] \approx 0.070\,\theta^2$. Similar results pertain in one and two dimensions (SI text; Figure 2A-C).

Conditional sweep probability

To compute the sweep probability, we first need an expression for the remaining wildtype population, N_{wt}. Therefore, we introduce time measure τ, with $\tau = 0$ when the first mutant arises. Recall that $t = 0$ at the origin of the wildtype population, so $t = \tau + x/c_{\mathrm{wt}}$. Once the mutant population starts expanding, we have $N_{\mathrm{wt}}(\tau) = \tilde{N}_{\mathrm{wt}}(\tau) - \Delta(\tau)$, where $\Delta(\tau)$ is the number of wildtype cells that the mutant has replaced, and $\tilde{N}_{\mathrm{wt}}(\tau) = \frac{4}{3}\pi x_{\mathrm{wt}}^3(\tau)$ is the wildtype population size had there been no mutant. While the mutant is surrounded by the wildtype, we have $\Delta(\tau) = \Delta_1(\tau) = \frac{4}{3}\pi x_{\mathrm{m}}^3(\tau)$. After the mutant breaches the wildtype boundary, $\Delta(\tau)$ is given by the intersection formula of two balls, which we denote $\Delta_2(\tau)$ (see SI text for the explicit formula). Together, we have

$$\begin{aligned} N_{\mathrm{wt}}(\tau) =&(\tilde{N}_{\mathrm{wt}}(\tau) - \Delta_1(\tau))\mathbf{1}\{[0, \tau_1]\} \\ &+ (\tilde{N}_{\mathrm{wt}}(\tau) - \Delta_2(\tau))\mathbf{1}\{[\tau_1, \tau_2]\} \end{aligned} \tag{6}$$

where τ_1 is the time at which the mutant reaches the wildtype boundary and τ_2 is the

Figure 2. Analytical and numerical solutions of the macroscopic model. **A.-C.** Probability density of the wildtype population radius at the time the first surviving mutant arises (dark curves) and of the distance between the origins of the mutant and wildtype expansions (light curves) in one dimension **(A)**, two dimensions **(B)** and three dimensions **(C)**. **D.** Approximate and exact solutions for the sweep probability conditioned on the wildtype population radius. **E.** Approximate and exact solutions for the unconditional sweep probability. **F.** Approximate and exact solutions for the probability density of the wildtype population radius at the time the first surviving mutant arises, conditioned on this mutant achieving a selective sweep. Except where parameter values are explicitly varied, we set $c_{\text{wt}} = 0.15, \tilde{\mu} = 10^{-5}, \rho = 0.23$ and $c_{\text{m}} = 0.31$. Formulas for all these curves are summarised in Table S2.

time at which the mutant has entirely replaced the wildtype. Noting that

$$\tau_1 c_{\text{m}} = (x - y) + \tau_1 c_{\text{wt}} \Rightarrow \tau_1 = \frac{x - y}{c_{\text{m}} - c_{\text{wt}}} \tag{7}$$

and

$$\tau_2 c_{\text{m}} = (x + y) + \tau_1 c_{\text{wt}} \Rightarrow \tau_2 = \frac{x + y}{c_{\text{m}} - c_{\text{wt}}}, \tag{8}$$

we express $N_{\text{wt}}(\tau)$ in terms of x and y (see SI text). We then apply Claim 1 to obtain the conditional sweep probability,

$$\Pr(\text{sweep}|X = x, Y = y) = e^{-\mu \int_0^\infty N_{\text{wt}}(\tau) \, d\tau}. \tag{9}$$

Although this integral can be solved analytically (SI text), the resulting expression is

complicated and not very enlightening, and we therefore seek simpler approximations. An especially fruitful approach is to assume $y = 0$, so that $f_Y(y|X = x) = \delta(y)$, where δ is the Dirac delta function. This yields an upper bound on the conditional sweep probability because, due to geometrical symmetry, the time required for a mutant to sweep (τ_2 in eqn. 8) is minimal when $y = 0$. With this approximation, eqn. 9 simplifies drastically as $\tau_1 = \tau_2 = \frac{x}{c_m - c_{wt}}$, and we do not need to integrate over $\Delta_2(\tau)$. The analytic solution is

$$\Pr(\text{sweep}|X = x, Y = 0) = e^{-\left(\frac{x}{\alpha}\right)^4}, \tag{10}$$

where $\alpha = \sqrt[4]{\frac{3(c_m - c_{wt})^3}{\pi \mu (c_{wt}^2 - 3c_{wt}c_m + 3c_m^2)}}$ is a characteristic length. Despite being based on a seemingly crude approximation, eqn. 10 is remarkably close to the exact conditional sweep probability (Figure 2D). The corresponding approximations in one and two dimensions (SI text) are likewise simple and useful.

Unconditional sweep probability

We now have all the necessary ingredients to compute the unconditional sweep probability by marginalizing out X and Y from the conditional sweep probability,

$$\Pr(\text{sweep}) =$$
$$\int_0^\infty \int_0^\infty \Pr(\text{sweep}|X = x, Y = y) f_Y(y|X = x) f_X(x) \, dy \, dx$$
$$= \int_0^\infty \int_0^x e^{-\mu \int_0^\infty N_{wt}(\tau) \, d\tau} \frac{3y^2}{x^3} \frac{4x^3 e^{-x^4/\theta^4}}{\theta^4} \, dy \, dx. \tag{11}$$

As before, we use the approximation $y = 0$ to obtain the strikingly simple result

$$\Pr(\text{sweep}) \leq \int_0^\infty \Pr(\text{sweep}|X = x, Y = 0) f_X(x) \, dx$$
$$= \int_0^\infty e^{-\left(\frac{x}{\alpha}\right)^4} \frac{4x^3 e^{-x^4/\theta^4}}{\theta^4} \, dx$$
$$= \left(\frac{c_m - c_{wt}}{c_m}\right)^3. \tag{12}$$

This result generalizes to analogous one- and two-dimensional models, with the exponent 3 replaced by the respective spatial dimension (SI text). The approximate expressions are close to the exact solution in one dimension, and to numerical evaluations of

the integral in two and three dimensions (Figure 2E). The sweep probability is independent of the mutation rate not only when $y = 0$ but also in the general case of eqn. 11. This follows from a more general result (SI text).

When the expanding wildtype must displace a resident competitor (as in our agent-based simulations, to follow), the sweep probability can be approximated in terms of evolutionary parameters via the FKPP solution as

$$\Pr(\text{sweep}) \approx \left(1 - \sqrt{\frac{r_{\mathrm{m}} - r_{\mathrm{wt}}}{r_{\mathrm{wt}} - r_{\mathrm{re}}}} \right)^d , \tag{13}$$

where r_{m}, r_{wt} and r_{re} are the proliferation rates of the mutant invader, wildtype invader and resident populations, respectively, and d is the spatial dimension.

Conditional arrival time of the first mutant

In biological systems where it is infeasible to track evolutionary dynamics, selective sweeps must be inferred from subsequent genetic data. For example, we might observe a public mutation in a tumour and ask when this mutation occurred. We can use our model to obtain the probability distribution of the radius X at the time the first mutant arose, given that we observe a selective sweep, simply by applying Bayes' theorem,

$$f_X(x|\text{sweep}) = \frac{\Pr(\text{sweep}|X = x) f_X(x)}{\Pr(\text{sweep})} . \tag{14}$$

Using the approximation $y = 0$, we obtain the Weibull distribution

$$f_X(x|\text{sweep}) \approx \frac{4x^3}{\beta^3 \theta^4} e^{-\frac{x^4}{\beta^3 \theta^4}} \tag{15}$$

where $\beta = \frac{c_{\mathrm{m}} - c_{\mathrm{wt}}}{c_{\mathrm{m}}}$. This approximation and the corresponding results in one and two dimensions are close to the exact solutions (Figure 2F).

Simulations

To gauge the robustness of our theoretical predictions against stochastic growth and discretization, we measured the frequency of selective sweeps in a two-dimensional agent-based model.

Figure 3. Agent-based simulation results versus macroscopic model predictions. **A.** Probability density of the wildtype population radius at the time the first surviving mutant arises. **B.** Probability density of the distance between the wildtype origin and the location at which the first successful mutant is born. **C.** Unconditional selective sweep probability versus the ratio of the proliferation rate differences $a_{\mathrm{wt}} = r_{\mathrm{n}} - r_{\mathrm{re}}$ and $a_{\mathrm{m}} = r_{\mathrm{m}} - r_{\mathrm{wt}}$. Each histogram in panels **A** and **B** and each data point in **C** is based on 1,000 replicates. Except where parameter values are explicitly varied, we set $m = 0.05, \tilde{\mu} = 10^{-5}, K = 16, r_{\mathrm{re}} = 0.91, r_{\mathrm{wt}} = 1$ and $r_{\mathrm{m}} = 1.3$, leading to speeds $c_{\mathrm{wt}} \approx 0.15$ and $c_{\mathrm{m}} \approx 0.31$ and mutant survival probability $\rho \approx 0.23$. Results for alternative parameter values and less stringent sweep definitions are shown in Figures S1-S3. The conversion from simulation parameters to macroscopic model parameters is explained in Materials and Methods.

An agent-based realisation of the two-dimensional spatial Moran process

We suppose that the wildtype population invades a habitat initially occupied by a resident competitor, which is a plausible biological scenario for both invasive species [40] and invasive tumours [28, 32]. Our agent-based model thus has three types of individuals: resident, wildtype invader, and mutant invader with local proliferation rates r_{re}, r_{wt} and r_{m}. Localised competition between wildype and resident slows the wildtype expansion and creates potential for selective sweeps. We implemented this model using the *demon* agent-based modelling framework [41] within the *warlock* computational workflow [42], which facilitates running large numbers of simulations on a high-performance computing cluster. We have previously applied the same framework to studying cancer evolution [28, 42]. Further model details are given in Materials and Methods.

The agent-based simulations provide useful insights additional to the macroscopic

model because, although the general setup is the same, they differ in several ways. Space in the simulations is divided into discrete patches; the times between birth and dispersal events are exponentially distributed random variables constituting another source of stochasticity; population boundaries are rough, not smooth; the expansion wave front is typically not sharp, and changes shape as the wave progresses; and the mutant will have increased propagation speed when competing with the resident population. Hence we would not expect perfect agreement between the results of the two models.

Linking the microscopic and macroscopic models

Our agent-based model approximately resembles a spatial death-birth Moran process (also known as the stepping stone model) [43, 44]. Expansion speeds in the spatial Moran process can in turn be approximated using the Fisher-Kolmogorov-Petrovsky-Piscounov (FKPP) equation [37, 38], which predicts that the mutant will expand within the wildtype with constant radial expansion speed c_m dependent on the difference in their local proliferation rates $a_m = r_m - r_{wt}$ [33, 34, 36]. Analogously, we have a constant expansion speed of the wildtype c_{wt} dependent on $a_{wt} = r_{wt} - r_{re}$. To compare the results of our discrete-space simulations to our continuous-space macroscopic model, we measured the propagation speeds of the wildtype within the resident, and of the mutant within the wildtype (see Materials and Methods). Further investigations of propagation speeds in this model will be the subject of a further study.

Simulation results

Given the considerable differences between the models, the probability density functions resulting from the macroscopic and microscopic models are reassuringly consistent. The radius at the time the first surviving mutant arises is slightly lower in the simulations than in our analytical model (Figure 3A). Similarly, the distribution for the location of the first surviving mutant coincides well except for a small offset of the mean (Figure 3B). Such offsets are expected due to discretization effects and the fact that, in the simulations, the propagation front needs to be established before the expansion can proceed. The sweep probabilities in simulations are nevertheless very close to our ana-

lytical predictions and change very little when varying the mutation rate over orders of magnitude, confirming our prediction that the sweep probability is independent of the mutation rate (Figure 3C; Figures S1-S4).

Comparison with other growth models

To further demonstrate the robustness of our findings, we compare them to results obtained from alternative models. Instead of permitting mutants to grow within the wildtype population, one might instead assume that dispersal is restricted to the wild-type population boundary. In this case, a complete sweep is impossible and we instead ask whether the first arising mutant envelopes the wildtype. Interestingly, this envel-opment probability is independent of the mutation rate and obtains values close to the sweep probabilities of our main model [21] (Figures 4 and S5; SI text). Alternatively, one might consider exponential growth with no competition, and ask whether a single mutant comes to dominate the population. Adapting our methods to this case (see SI text), we find that a single mutant is unlikely to become dominant unless its exponential growth rate is several times larger than that of the wildtype (Figure S6). Although our focus is on expanding populations, we also applied our methods to compute the sweep probability in constant populations (SI text), obtaining a result that depends on both mutation rate and population size (Figure 4), in agreement with prior analyses [45, 46].

Application to cancer

Our findings have especially interesting implications for understanding cancer evo-lution. Here we consider a wildtype tumour that evolves while invading a resident population of normal cells.

Arrival of the first driver mutation during tumour growth

Our model has only three parameters: the mutation rate conditional on survival, μ, and the wildtype and mutant propagation speeds, c_{wt} and c_{m}. To estimate c_{wt} for human tumours, we follow a similar procedure to prior studies [21, 47]. Consider a

Figure 4. Comparison of sweep probabilities between different growth models in 3D. We show the tuples (μ, c_{m}) that yield $\Pr(\text{sweep}) = 0.5$ for models with proliferation restricted to the boundary (solid purple line), population size fixed at radius x_0 (dashed red lines), and our main model in which mutants can expand within the expanding wildtype population (solid orange line). In the two range expansion models, the wildtype speed is set to $c_{\mathrm{wt}} = 0.15$ (dotted black line).

tumor that grows to volume V between 1 and 10 cm^3 in time T between 5 and 20 years. The propagation speed can then be estimated as

$$\tilde{c} = \frac{r}{T} = \frac{\sqrt[3]{\frac{3}{4\pi}V}}{T},$$

which equates to between 1 and 40 μm per day. Given that the diameter of a typical cancer cell is $l \approx 20\,\mu$m [48] and the generation time (cell cycle time) is $\tau_G \approx 4$ days [49], we can switch units to obtain $c = \tilde{c} \times \tau_G/l$, which is between 0.15 and 7 cell diameters per generation.

The rate of acquiring advantageous (driver) mutations is usually estimated at around $\tilde{\mu} = 10^{-5}$ per cell per generation [47, 50]. For the survival probability ρ, we assume values between 0.09 and 0.5, in agreement with inferred values in colorectal tumours [51].

Together, these parameter values lead to a typical length scale

$$\theta = \sqrt[4]{\frac{3c_{\mathrm{wt}}}{\pi\mu}} \approx 10 \text{ to } 50 \text{ cells.}$$

The expected values of X and Y are then $\mathbb{E}[X] = 9$ to 45 cell diameters and $\mathbb{E}[Y] = 7$
to 35 cell diameters. It follows that a tumour, having acquired sufficient mutations to
grow, will likely gain further driver mutations already during early development.

Ubiquity of clonal interference in cancer

During tumour growth, we predict very low sweep probabilities unless the mutant
grows much faster within the tumour than the tumour grows into surrounding tissue
(Figure 2E, 3C). When sweeps do occur, they will begin and end in the very early stages
of tumour growth. Assuming c_m is at most ten times c_wt, we find that the expected
tumour radius when a selective sweep occurred, given that a sweep did occur, is no
more than 50 cells, corresponding to a population size N of no more than 400,000 cells.
The time for the sweep to be completed is then $\tau_2 \approx 40$ generations, at a population
size of approximately 800,000 cells. These numerical estimates are highly insensitive to
varying parameter values because $\mathbb{E}[X]$, $\mathbb{E}[Y]$ and the expected value of $f_X(x|\text{sweep})$
are proportional to the characteristic length θ, which varies with the fourth root of μ
and c_wt. Hence, in three dimensions, changing μ or c_wt by a factor of 100 changes our
estimates by only a factor of $\sqrt[4]{100} \approx 3$.

DISCUSSION

Here we have used mathematical modelling and analysis to determine the expected
frequency of classic selective sweeps versus incomplete or soft sweeps during range ex-
pansions. We find that this frequency is generally expected to be low, even for mutations
with very high selection coefficients. Moreover, when the wildtype and mutant radial
expansion speeds are constant, the sweep probability can be expressed solely in terms of
those speeds, which can in turn be related, through the FKPP equation or other standard
models, to the selection coefficient, dispersal rates, and other basic parameters. Why is
the sweep probability independent of the mutation rate? An intuitive explanation is that
if the mutation rate is higher then, on the one hand, the first advantageous mutation is
likely to arise in a smaller population – meaning that it has less distance to travel to
achieve a sweep – but, on the other hand, competing mutations will also tend to arise

sooner than in the case of a lower mutation rate. These two effects exactly cancel out under the assumption of constant radial growth speeds.

We make three arguments to justify our focus on this particular model of range expansions. First, our model corresponds to the continuum approximation of standard mathematical models of range expansions and spatial population genetics: the spatial Moran process (or stepping stone model [44]) and the biassed voter model (which is equivalent to an Eden growth model extended to allow local dispersal and competition throughout the population) [36]. These models are well understood, intuitive, and easy to parameterise. Second, because our model is relatively permissive to selective sweeps, it provides useful upper bounds for selective sweep probabilities in more complex scenarios, as further explained below.

Our third justification is that, in much the same way as the Moran and Wright-Fisher processes are the most useful, tractable models of evolution in constant-sized, non-spatial populations, so the constant-radial-speed model yields the clearest results for range expansions. Haldane's famous rule of thumb is that the fixation probability of a weakly beneficial allele in a large, non-spatial population of constant size is approximately proportional to its relative advantage in terms of proliferation rate [52]. Here we have obtained the comparably simple result that the probability of a strongly beneficial allele achieving a selective sweep in an expanding population is approximately equal to its relative advantage in terms of radial expansion speed, raised to the power of the spatial dimension. For instance, if the radial expansion speed at which a mutant spreads within the wildtype population is twice the speed at which the wildtype expands then the probability of this mutant achieving a selective sweep can be approximated simply as $(1 - 1/2)^2 = 1/4$ in two dimensions and $1/8$ in three dimensions.

Some alternative models have been considered previously. Antal and colleagues [21] used a macroscopic model similar to ours to investigate the case in which mutations arise only at the boundary of a range expansion. Given that selective sweeps are then impossible, the interesting outcome is when the mutant envelops the wildtype. Ralph & Coop [45] and Martens and colleagues [20, 46] instead considered constant-sized populations, and found that selective sweeps are likely only if the population width is much smaller than a characteristic length scale, which depends on the mutation rate, the dispersal rate, the effective local population density, and the strength of selection. In SI

Text we compare our results to those of these prior studies and consider other alternative growth laws. The general conclusions are that selective sweeps are predicted to be rare except in small populations and that our model provides a useful upper bound on the sweep probability in range expansions. Selective sweeps will be even less frequent when the wildtype radial expansion speed increases over time (as is typical in the middle stage of tumour progression) or when mutant expansion is restricted to the tumour boundary (which may be the case in some tumours [28, 29, 53]).

A selective sweep can occur only if the rate of spread of an advantageous mutation exceeds the expansion speed of the wildtype population. This scenario is plausible, for example, in a biofilm growing slowly under antibiotic stress, in which case our findings predict the evolutionary dynamics of antibiotic resistance. Our results apply equally to an invasive species that is still adapting to its new conditions and whose range expansion is slowed by the need to modify its environment (niche construction) or by interspecific interactions [40]. If the invader must outcompete a resident then the sweep probability can be approximated via the FKPP solution in terms of proliferation rates as in eqn. 13. The dimensional exponent in this result implies that selective sweeps are most likely to occur in species invading essentially linear habitats, such as coastlines.

Our three-dimensional results are particularly relevant to understanding the nature of solid tumour evolution. There are two ways in which tumours might acquire the multiple clonal drivers (advantageous mutations) observed in sequencing studies. The first route is via sequential population bottlenecks. Suppose that early tumour development comprises not one but several successive range expansions. Growth repeatedly stalls due to constraints such as hypoxia, immune control, and physical barriers. Driver mutations enable subclones to escape these constraints and invade new territory, each time purging genetic diversity so that the final, prolonged expansion originates from a single highly transformed cell. This episodic model is conventional and has been particularly well characterised in recent studies of colorectal cancer [54, 55] and breast cancer [56]. The second way to acquire clonal drivers is via selective sweeps during the final range expansion. Our results suggest that the first process is the more important. Selective sweeps of even extremely strong drivers are highly unlikely to occur in tumours containing more than a few hundred thousand cells, equivalent to less than a cubic millimeter in volume. Pervasive genetic heterogeneity and parallel evolution [57] are

therefore revealed to be straightforward consequences of spatial expansion, expected to be ubiquitous across solid tumours, irrespective of mutation rate and degree of genomic instability.

METHODS

Numerical Integration

We performed numerical integration using the MATLAB function 'trapz'. For values of x close to 0 and beyond 2θ, $f_X(x)$ is relatively very small and its contribution to the integrals is negligible. Hence, to avoid numerical errors without compromising precision, we set the lower and upper bounds of integration to 0.001θ and 3θ when integrating over $f_X(x)$. Similarly, we set 0.001θ as the lower bound of the integration over y. We set interval widths to 0.001θ for $x \leq 0.5$, and 0.01θ for $x > 0.5$.

Agent-based simulations

Individuals in our agent-based model are subdivided into well-mixed demes on a regular two-dimensional grid. The demes have identical carrying capacities, K, and are initially filled with residents, except that a single wildtype invader is introduced at the centre of the grid. At each time step, an individual is chosen at random, with probability weighted by fitness, to be replaced by two offspring. Each offspring then either migrates, with probability m, to a neighbouring deme in a randomly chosen direction, or remains in its parent deme. Local density dependence is implemented by imposing a very high death rate whenever a deme is above carrying capacity. Mutation is coupled to wildtype reproduction. To improve computational efficiency, resident individuals are not permitted to disperse; we verified with additional simulations (to be published in a later study) that this asymmetry has only a minor effect on the wildtype expansion speed. Further model details have been published previously [28, 42]. All simulations were performed on City, University of London's Hyperion cluster.

For all simulations, we set the deme size to $K = 16$ and the migration probability $m = 0.05$. For such small m, the probability to survive drift can be estimated as the probability to become locally fixed in one deme. Since the within-deme dynamics approximate a

Moran process, this probability is $\rho = \frac{1 - r_\text{wt}/r_\text{m}}{1 - (r_\text{wt}/r_\text{m})^K}$, where r_wt and r_m are the proliferation rates [58]. We measured propagation speeds by simulating the expansion of a wildtype into a large resident population, or of a mutant into a large wildtype population, in the absence of mutation and then applying linear regression to the growth curve of the effective radius, defined as the square root of the population size divided by π. For each parameter set, we took the mean speed from ten replicates, for which the standard deviation was consistently below 2% of the mean. We set the proliferation rates to $r_\text{re} = 0.91$ and $r_\text{wt} = 1.0$ for the resident and wildtype, respectvively, leading to $a_\text{wt} = r_\text{wt} - r_\text{re} = 0.09$. The measured wildtype speed was then $c_\text{wt} \approx 0.15$. We varied the mutant proliferation rate r_m from 1.1 to 3.0, so that $a_\text{m} = r_\text{m} - r_\text{wt}$ ranged from 0.1 to 2.0. The measured mutant speeds are presented in Table I.

a_m	0.1	0.2	0.3	0.4	0.5	0.6	0.7
c_m	0.14	0.23	0.31	0.38	0.46	0.53	0.60

a_m	0.8	0.9	1.0	1.1	1.2	1.3	1.4
c_m	0.68	0.75	0.82	0.89	0.96	1.03	1.11

a_m	1.5	1.6	1.7	1.8	1.9	2.0
c_m	1.17	1.24	1.31	1.39	1.46	1.52

Table I. Correspondence between difference in proliferation rates a_m and mutant propagation speed c_m, measured in simulations.

AUTHOR CONTRIBUTIONS

RN conceived the research question and supervised the project. RN and AS designed the research. AS and RK carried out the mathematical analysis. MB performed agent-based simulations. All authors wrote and approved the manuscript.

ACKNOWLEDGMENTS

AS was supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie EvoGamesPlus grant agreement no.

955708. MB was supported by an award from the City, University of London Research Pump-priming Fund. RN was supported by the National Cancer Institute of the National Institutes of Health under Award Number U54CA217376. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

CODE AVAILABILITY

The code and data needed to reproduce our results can be found at https://zenodo.org/records/10775693.

---

[1] Laurent Excoffier, Matthieu Foll, and Rémy J Petit. Genetic consequences of range expansions. *Annual Review of Ecology, Evolution, and Systematics*, 40:481–501, 2009.

[2] Oskar Hallatschek, Pascal Hersen, Sharad Ramanathan, and David R Nelson. Genetic drift at expanding frontiers promotes gene segregation. *Proceedings of the National Academy of Sciences*, 104(50):19926–19930, 2007.

[3] Zaira Seferbekova, Artem Lomakin, Lucy R Yates, and Moritz Gerstung. Spatial biology of cancer evolution. *Nature Reviews Genetics*, 24(5):295–313, 2023.

[4] Leslie G Biesecker and Nancy B Spinner. A genomic view of mosaicism and human disease. *Nature Reviews Genetics*, 14(5):307–320, 2013.

[5] Iñigo Martincorena, Amit Roshan, Moritz Gerstung, Peter Ellis, Peter Van Loo, Stuart McLaren, David C Wedge, Anthony Fullam, Ludmil B Alexandrov, Jose M Tubio, et al. High burden and pervasive positive selection of somatic mutations in normal human skin. *Science*, 348(6237):880–886, 2015.

[6] Luigi L Cavalli-Sforza, Paolo Menozzi, and Alberto Piazza. Demic expansions and human evolution. *Science*, 259(5095):639–646, 1993.

[7] Gretta T Pecl, Miguel B Araújo, Johann D Bell, Julia Blanchard, Timothy C Bonebrake, I-Ching Chen, Timothy D Clark, Robert K Colwell, Finn Danielsen, Birgitta Evengård, et al. Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, 355(6332):eaai9214, 2017.

[8] Emily V Moran and Jake M Alexander. Evolutionary responses to global change: lessons from invasive species. *Ecology Letters*, 17(5):637–649, 2014.

[9] Oskar Hallatschek and David R Nelson. Life at the front of an expanding population. *Evolution*, 64(1):193–206, 2010.

[10] Kirill S Korolev, Melanie JI Müller, Nilay Karahan, Andrew W Murray, Oskar Hallatschek, and David R Nelson. Selective sweeps in growing microbial colonies. *Physical biology*, 9(2):026008, 2012.

[11] Diana Fusco, Matti Gralka, Jona Kayser, Alex Anderson, and Oskar Hallatschek. Excess of mutational jackpot events in expanding populations revealed by spatial luria–delbrück experiments. *Nature communications*, 7(1):12760, 2016.

[12] Serhii Aif, Nico Appold, Lucas Kampman, Oskar Hallatschek, and Jona Kayser. Evolutionary rescue of resistant mutants is governed by a balance between radial expansion and selection in compact populations. *Nature Communications*, 13(1):7916, 2022.

[13] Douglas Hanahan and Robert A Weinberg. The hallmarks of cancer. *cell*, 100(1):57–70, 2000.

[14] Eric R Fearon and Bert Vogelstein. A genetic model for colorectal tumorigenesis. *cell*, 61(5):759–767, 1990.

[15] Marco Gerlinger, Andrew J Rowan, Stuart Horswell, James Larkin, David Endesfelder, Eva Gronroos, Pierre Martinez, Nicholas Matthews, Aengus Stewart, Patrick Tarpey, et al. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *New England journal of medicine*, 366(10):883–892, 2012.

[16] Xiaodong Liu, Ke Zhang, Neslihan A Kaya, Zhe Jia, Dafei Wu, Tingting Chen, Zhiyuan Liu, Sinan Zhu, Axel M Hillmer, Torsten Wuestefeld, et al. Tumor phylogeography reveals block-shaped spatial heterogeneity and the mode of evolution in hepatocellular carcinoma. *Nature Communications*, 15(1):3169, 2024.

[17] Moritz Gerstung, Clemency Jolly, Ignaty Leshchiner, Stefan C Dentro, Santiago Gonzalez, Daniel Rosebrock, Thomas J Mitchell, Yulia Rubanova, Pavana Anur, Kaixian Yu, et al. The evolutionary history of 2,658 cancers. *Nature*, 578(7793):122–128, 2020.

[18] Alexander Davis, Ruli Gao, and Nicholas Navin. Tumor evolution: Linear, branching, neutral or punctuated? *Biochimica et Biophysica Acta (BBA)-Reviews on Cancer*, 1867(2):151–161, 2017.

[19] Samra Turajlic, Andrea Sottoriva, Trevor Graham, and Charles Swanton. Resolving genetic heterogeneity in cancer. *Nature Reviews Genetics*, 20(7):404–416, 2019.

[20] Erik A Martens, Rumen Kostadinov, Carlo C Maley, and Oskar Hallatschek. Spatial structure increases the waiting time for cancer. *New journal of physics*, 13(11):115014, 2011.

[21] Tibor Antal, PL Krapivsky, and MA Nowak. Spatial evolution of tumors with successive driver mutations. *Physical Review E*, 92(2):022705, 2015.

[22] Chay Paterson, Martin A Nowak, and Bartlomiej Waclaw. An exactly solvable, spatial model of mutation accumulation in cancer. *Scientific reports*, 6(1):39511, 2016.

[23] Peter Armitage and Richard Doll. The age distribution of cancer and a multi-stage theory of carcinogenesis. *British journal of cancer*, 91(12):1983–1989, 2004.

[24] Chay Paterson, Hans Clevers, and Ivana Bozic. Mathematical model of colorectal cancer initiation. *Proceedings of the National Academy of Sciences*, 117(34):20681–20688, 2020.

[25] Michael D Nicholson, David Cheek, and Tibor Antal. Sequential mutations in exponentially growing populations. *PLOS Computational Biology*, 19(7):e1011289, 2023.

[26] Ruping Sun, Zheng Hu, Andrea Sottoriva, Trevor A Graham, Arbel Harpak, Zhicheng Ma, Jared M Fischer, Darryl Shibata, and Christina Curtis. Between-region genetic divergence reflects the mode and tempo of tumor evolution. *Nature genetics*, 49(7):1015–1024, 2017.

[27] Jeffrey West, Ryan O Schenck, Chandler Gatenbee, Mark Robertson-Tessi, and Alexander RA Anderson. Normal tissue architecture determines the evolutionary course of cancer. *Nature communications*, 12(1):2060, 2021.

[28] Robert Noble, Dominik Burri, Cécile Le Sueur, Jeanne Lemant, Yannick Viossat, Jakob Nikolas Kather, and Niko Beerenwinkel. Spatial structure governs the mode of tumour evolution. *Nature ecology & evolution*, 6(2):207–217, 2022.

[29] Xiao Fu, Yue Zhao, Jose I Lopez, Andrew Rowan, Lewis Au, Annika Fendler, Steve Hazell, Hang Xu, Stuart Horswell, Scott TC Shepherd, et al. Spatial patterns of tumour growth impact clonal diversification in a computational model and the tracerx renal study. *Nature ecology & evolution*, 6(1):88–102, 2022.

[30] Noemi Andor, Trevor A Graham, Marnix Jansen, Li C Xia, C Athena Aktipis, Claudia Petritsch, Hanlee P Ji, and Carlo C Maley. Pan-cancer analysis of the extent and consequences of intratumor heterogeneity. *Nature medicine*, 22(1):105–113, 2016.

[31] Nicholas McGranahan and Charles Swanton. Clonal heterogeneity and tumor evolution: past, present, and the future. *Cell*, 168(4):613–628, 2017.

[32] Robert Noble, John T Burley, Cécile Le Sueur, and Michael E Hochberg. When, why and how tumour clonal diversity predicts survival. *Evolutionary applications*, 13(7):1558–1568, 2020.

[33] Ronald Aylmer Fisher. The wave of advance of advantageous genes. *Annals of eugenics*, 7(4):355–369, 1937.

[34] Andrey Nikolaevich Kolmogorov, Ivan Georgievich Petrovsky, and Nikolay Semenovich Piskunov. A study of the diffusion equation with increase in the amount of substance, and its application to a biological problem. *Bullentin of Moscow State University*, 1937.

[35] Eric Brunet and Bernard Derrida. Shift in the velocity of a front due to a cutoff. *Physical Review E*, 56(3):2597, 1997.

[36] James Dickson Murray. *Mathematical Biology: I: An introduction*, volume 17. Springer, 2002.

[37] Oskar Hallatschek and Kirill S Korolev. Fisher waves in the strong noise limit. *Physical review letters*, 103(10):108103, 2009.

[38] Bahram Houchmandzadeh and Marcel Vallade. Fisher waves: An individual-based stochastic model. *Physical Review E*, 96(1):012414, 2017.

[39] Joachim Hermisson and Pleuni S Pennings. Soft sweeps and beyond: understanding the patterns and probabilities of selection footprints under rapid adaptation. *Methods in Ecology and Evolution*, 8(6):700–716, 2017.

[40] Jens-Christian Svenning, Dominique Gravel, Robert D Holt, Frank M Schurr, Wilfried Thuiller, Tamara Münkemüller, Katja H Schiffers, Stefan Dullinger, Thomas C Edwards Jr, Thomas Hickler, et al. The influence of interspecific interactions on species range expansion rates. *Ecography*, 37(12):1198–1209, 2014.

[41] Robert Noble. demon: deme-based oncology model. https://github.com/robjohnnoble/demonmodel, 2023.

[42] Maciej Bak, Blair Colyer, Veselin Manojlović, and Robert Noble. Warlock: an automated computational workflow for simulating spatially structured tumour evolution. *arXiv preprint arXiv:2301.07808*, 2023.

[43] Richard Durrett, Jasmine Foo, and Kevin Leder. Spatial moran models, ii: cancer initiation in spatially structured tissue. *Journal of mathematical biology*, 72:1369–1400, 2016.

[44] Motoo Kimura and George H Weiss. The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics*, 49(4):561, 1964.

[45] Peter Ralph and Graham Coop. Parallel adaptation: one or many waves of advance of an advantageous allele? *Genetics*, 186(2):647–668, 2010.

[46] Erik A Martens and Oskar Hallatschek. Interfering waves of adaptation promote spatial mixing. *Genetics*, 189(3):1045–1060, 2011.

[47] Ivana Bozic, Tibor Antal, Hisashi Ohtsuki, Hannah Carter, Dewey Kim, Sining Chen, Rachel Karchin, Kenneth W Kinzler, Bert Vogelstein, and Martin A Nowak. Accumulation of driver and passenger mutations during tumor progression. *Proceedings of the National Academy of Sciences*, 107(43):18545–18550, 2010.

[48] Babita Shashni, Shinya Ariyasu, Reisa Takeda, Toshihiro Suzuki, Shota Shiina, Kazunori Akimoto, Takuto Maeda, Naoyuki Aikawa, Ryo Abe, Tomohiro Osaki, et al. Size-based differentiation of cancer and normal cells by a particle size analyzer assisted by a cell-recognition pc software. *Biological and Pharmaceutical Bulletin*, 41(4):487–503, 2018.

[49] Siân Jones, Wei-dong Chen, Giovanni Parmigiani, Frank Diehl, Niko Beerenwinkel, Tibor Antal, Arne Traulsen, Martin A Nowak, Christopher Siegel, Victor E Velculescu, et al. Comparative lesion sequencing provides insights into tumor evolution. *Proceedings of the National Academy of Sciences*, 105(11):4283–4288, 2008.

[50] Marc J Williams, Benjamin Werner, Timon Heide, Christina Curtis, Chris P Barnes, Andrea Sottoriva, and Trevor A Graham. Quantification of subclonal selection in cancer from bulk sequencing data. *Nature genetics*, 50(6):895–903, 2018.

[51] Benjamin Werner, Jack Case, Marc J Williams, Ketevan Chkhaidze, Daniel Temko, Javier Fernández-Mateos, George D Cresswell, Daniel Nichol, William Cross, Inmaculada Spiteri, et al. Measuring single cell divisions in human tissues from multi-region sequencing data. *Nature communications*, 11(1):1035, 2020.

[52] John Burdon Sanderson Haldane. A mathematical theory of natural and artificial selection. *Mathematical Proceedings of the Cambridge Philosophical Society*, 23(5):607–615, 1927.

[53] Maya A Lewinsohn, Trevor Bedford, Nicola F Müller, and Alison F Feder. State-dependent evolutionary models reveal modes of solid tumour growth. *Nature Ecology & Evolution*, 7(4):581–596, 2023.

[54] Marc D Ryser, Diego Mallo, Allison Hall, Timothy Hardman, Lorraine M King, Sergei Tatishchev, Inmaculada C Sorribes, Carlo C Maley, Jeffrey R Marks, E Shelley Hwang, et al. Minimal barriers to invasion during human colorectal tumor growth. *Nature communications*, 11(1):1280, 2020.

[55] William Cross, Michal Kovac, Ville Mustonen, Daniel Temko, Hayley Davis, Ann-Marie Baker, Sujata Biswas, Roland Arnold, Laura Chegwidden, Chandler Gatenbee, et al. The evolutionary landscape of colorectal tumorigenesis. *Nature ecology & evolution*, 2(10):1661–1672, 2018.

[56] Artem Lomakin, Jessica Svedlund, Carina Strell, Milana Gataric, Artem Shmatko, Gleb Rukhovich, Jun Sung Park, Young Seok Ju, Stefan Dentro, Vitalii Kleshchevnikov, et al. Spatial genomics maps the structure, nature and evolution of cancer clones. *Nature*, 611(7936):594–602, 2022.

[57] Alexander M Frankell, Michelle Dietzen, Maise Al Bakir, Emilia L Lim, Takahiro Karasaki, Sophia Ward, Selvaraju Veeriah, Emma Colliver, Ariana Huebner, Abigail Bunkum, et al. The evolution of lung cancer and impact of subclonal selection in tracerx. *Nature*, pages 1–9, 2023.

[58] Martin A Nowak. *Evolutionary dynamics: exploring the equations of life*. Harvard university press, 2006.