Paleolithic Gene Duplications Primed Adaptive Evolution of Human Amylase Locus Upon

Agriculture

Feyza Yılmaz[1]*, Charikleia Karageorgiou[2]*, Kwondo Kim[1]*, Petar Pajic[2], Kendra Scheer[2],

Human Genome Structural Variation Consortium, Christine R. Beck[1,3,4], Ann-Marie

Torregrossa[5,6], Charles Lee[1]**, Omer Gokcumen[2]**

Affiliations

1 - The Jackson Laboratory for Genomic Medicine, Farmington, CT, 06032, USA

2 - Department of Biological Sciences, University at Buffalo, Buffalo, NY, 14260, USA

3 - University of Connecticut, Institute for Systems Genomics, Storrs, CT, 06269, USA

4 - The University of Connecticut Health Center, Farmington, CT, 06032, USA

5 - Department of Psychology, University at Buffalo, Buffalo, NY 14260, USA

6 - University at Buffalo Center for Ingestive Behavior Research, University at Buffalo,

University at Buffalo, NY 14260, USA

* Contributed equally; ** Correspondence

Abstract

Starch digestion is a cornerstone of human nutrition. The amylase genes code for the starch-

digesting amylase enzyme. Previous studies suggested that the salivary amylase (*AMY1*) gene copy

number increased in response to agricultural diets. However, the lack of nucleotide resolution of

the amylase locus hindered detailed evolutionary analyses. Here, we have resolved this locus at

nucleotide resolution in 98 present-day humans and identified 30 distinct haplotypes, revealing

that the coding sequences of all amylase gene copies are evolving under negative selection. The

phylogenetic reconstruction suggested that haplotypes with three *AMY1* gene copies, prevalent

across all continents and constituting about 70% of observed haplotypes, originated before the out-

of-Africa migrations of ancestral modern humans. Using thousands of unique 25 base pair

sequences across the amylase locus, we showed that additional *AMY1* gene copies existed in the

genomes of four archaic hominin genomes, indicating that the initial duplication of this locus may

have occurred as far back 800,000 years ago. We similarly analyzed 73 ancient human genomes

dating from 300 - 45,000 years ago and found that the *AMY1* copy number variation observed

today existed long before the advent of agriculture (~10,000 years ago), predisposing this locus to

adaptive increase in the frequency of higher amylase copy number with the spread of agriculture.

Mechanistically, the common three-copy haplotypes seeded non-allelic homologous

recombination events that appear to be occurring at one of the fastest rates seen for tandem repeats

in the human genome. Our study provides a comprehensive population-level understanding of the

genomic structure of the amylase locus, identifying the mechanisms and evolutionary history

underlying its duplication and copy number variability in relation to the onset of agriculture.

Results

Copy number variation at the amylase locus is frequently attributed to human health and adaptation (*1*). As such, this structurally variable locus is a prime target for research on the fundamental biology of gene duplications. There are two types of amylase genes, *AMY1* and *AMY2,* which are reported to be expressed in the salivary glands and pancreas, respectively (*2*). Both genes encode for the amylase enzyme, which breaks down polymeric starch into simple sugar molecules, a crucial digestive process for starch-eating species (*3*). It has been shown that mammals that consume starch-rich diets underwent independent bursts of amylase gene duplications from the ancestral pancreatic *AMY2-like* gene (*4*). A great ape-specific duplication resulted in the formation of the salivary *AMY1* gene (*5*), which has since evolved to produce unusual copy number variations, ranging from 2 to 17 copies per diploid cell (*1*, *6*). This variation is especially prominent in human populations with high starch consumption, particularly those with a history of agriculture (*1*, *6*, *7*). These evolutionary insights indicate that copy number variation at the amylase locus may play an adaptive role in shaping the metabolic response to starchy diets, including the presence of microbes that break down amylase-resistant starch (*8*).

Given its adaptive and putative functional roles, the duplications of the *AMY1* gene was linked to the advent of agriculture approximately 10,000 years ago (*1*). However, the evolutionary analyses of the *AMY1* gene is hindered by the presence of highly similar DNA sequences (> 99%) within the amylase locus and makes it challenging to accurately resolve the haplotypes at this locus. The lack of precise nucleotide-level evolutionary analysis leads to disagreements about the timing and functional significance of *AMY1* gene duplications in relation to starch-rich diets and human evolution (*7, 9–12*). To address this issue, we have resolved this locus at nucleotide resolution at a population level across 98 individuals from different populations, using optical

genome mapping and long-read sequencing techniques. The nucleotide-resolved haplotypes of this

locus subsequently allowed us to conduct evolutionary genetic analyses on ancient human and

archaic hominin genomes to investigate the timing of *AMY1* gene duplications within the context

of agriculture.

Structural haplotypes at the human amylase locus

The amylase locus in the human genome is a ~212.5-kbp region on chromosome 1 (GRCh38;

chr1:103,554,220–103,766,732) which contains *AMY2B*, *AMY2A*, *AMY1A*, *AMY1B*, and *AMY1C*

genes (Fig. 1A). This locus is largely composed of segmental duplications with > 99% sequence

similarity, which complicates its accurate assembly using short-read sequencing (fig. S1). Using

the sequence similarity of segmental duplications and the labeling patterns from the optical

genome mapping data from GRCh38 *in silico* map, we defined six distinct amylase segments

overlapping the amylase genes, depicted by colored arrows (Fig. 1A and table S1). Using optical

genome mapping, which has been previously demonstrated to resolve similar complex regions

(*13–15*), we constructed haplotype-resolved diploid assemblies for 98 individuals (n = 196

sampled alleles (i.e., haploid sample size); table S2) and characterized this locus using the copy

number and orientation of the amylase segments (Fig. 1A). This approach allowed us to identify

51 distinct amylase haplotypes (fig. S2, A and B, and table S3), of which 7 were previously

reported (*7*) (fig. S3). These haplotypes were then classified based on the number of amylase gene

copies, adhering to the established nomenclature; HXAYBZ: HX: represents *AMY1*, AY:

represents *AMY2A*, and BZ: represents *AMY2B* copy numbers, with superscript "a" indicating

ancestral and superscript "r" indicating reference haplotypes (fig. S4) (*7*, *11*). We subsequently

defined 30 high-confidence haplotypes (from 117 observed alleles in 81 individuals) that were

  orthogonally supported by *de novo* assemblies based on long-read sequencing (Fig. 1B and Data

89 Files-S1). This represents the first nucleotide-level reconstruction of the amylase locus at a

90 population scale.

Fig. 1. Amylase structural haplotypes identified from present-day humans in this study. (**A**)
Segmental duplications (light to dark gray: 90-98% similarity, light to dark orange: > 99%
similarity), GENCODE V44 gene annotations, and long terminal repeats (LTRs) are represented
as tracks. The lower panel shows amylase segments (colored arrows), and haplotype structures of
GRCh38 and T2T-chm13 reference assemblies, represented as *in silico* maps with white
backgrounds and vertical blue lines displaying optical mapping labels. The AMY2B segment
overlaps the *AMY2B* gene, AMY2A.1 and AMY2A.2 segments overlap the *AMY2A* gene, and the
AMY1 segment overlaps the *AMY1* gene. (**B**) The high-confidence amylase structural haplotypes
resolved in our dataset (n = 30). The vertical black line in the second AMY1 segment of H3^r.3
represents the polymorphic label present in three alleles. The diagonal stripes in the second
AMY2B segment of H3B2.1 indicate that it is a partial copy of the first AMY2B segment.
Haplotype IDs describe: HX: X denotes the number of *AMY1* copies; AX: X denotes the number

of *AMY2A* copies; BX: X denotes the number of *AMY2B* copies. The superscript "a" denotes the
ancestral amylase haplotype structure, and the superscript "r" denotes the reference amylase
haplotype structure. The number in parentheses indicates the number of alleles. (**C**) The
distribution of common amylase haplotypes across 26 population samples. (**D**) The proportion of
singletons for tandem repeat loci (EnsembleTR) across the genome. For adequate comparison, we
used the same individuals (n = 33) for whom we were able to reconstruct amylase haplotypes in
our dataset. Additionally, we filtered the tandem repeat loci (719 loci, unit length 1-6 bp) that we
analyzed to match the number of distinct alleles (n = 21) observed in the amylase locus. The
asterisk (*) represents the proportion of singletons among all distinct haplotypes (~67%, 14 out of
21) detected at the amylase locus. (**E**) Rarefaction and extrapolation sampling curve based on 51
amylase haplotypes, displaying how the number of distinct haplotypes (blue line) is projected to
saturate with the increase in the number of alleles. The rate of change (red line, 0.13) indicates the
number of novel haplotypes discovered per unit increase in the number of analyzed alleles. The
dashed line shows the proportion of estimated number of samples (86.72%) captured in our study.

The length of amylase haplotypes ranged from 111-kbp ($H1^a.1$ and $H1^a.2$) to 402-kbp

(H7.1) (Fig. 1B), capturing those that are structurally identical to the GRCh38 ($H3^r.1$) and the T2T-

chm13 (H7.2) (*16*) assemblies (Fig. 1A). Four haplotypes, $H1^a.1$ (n = 20/117), $H3^r.1$ (n = 18/117),

$H3^r.2$ (n = 22/117), and $H3^r.4$ (n = 21/117), were categorized as common, each with an allele

frequency exceeding 5% across all populations. These four common haplotypes collectively

constitute approximately 70% of all amylase haplotypes (n = 81/117 identified in this study). We

also detected singletons, each detected only once (n = 19/117). Despite our limited sample size,

we found that all common haplotypes exist in all continental regions (Fig. 1C and fig. S5, A and

B). In addition, *AMY1* copy number variations do not exhibit a discernible geographic specificity

(p-value = 0.406, Kruskal-Wallis rank sum test).

Out of 30 distinct amylase haplotypes, we identified 19 (63%) as singletons. To infer the

relative mutation rate of the amylase locus, we compared this number to that of tandem repeats.

To avoid potential biases, we used the same 33 individuals that are present in both the tandem

repeat database (EnsembleTR) and our dataset. Only 21 of the 30 distinct amylase haplotypes were

present in these 33 individuals. To make sure the analysis was consistent across different genomic

regions, we only considered tandem repeat loci that had 21 detected alleles in the human

population. This yielded 719 tandem repeat loci (fig. S6). We found that the proportion of singleton

haplotypes was significantly higher for the amylase locus than the genome-wide average for the

719 tandem repeat loci (the observed empirical percentile = 0.017, Fig. 1D). This observation is

informative for understanding the mutation rate at this locus because the allele frequency spectrum

and proportion of singletons are determined by mutation rate and genetic drift (*17*). By using the

same individuals and matching the number of distinct haplotypes in our comparison, we control

for demographic biases and provide a relative estimation of the mutation rate for the amylase locus.

Considering that short tandem repeats have mutation rates as low as 10^{-8} (similar to single

nucleotide variant mutation rate) and, in some cases, can be as high as 10^{-2} mutations per locus

per generation (*18*), our analysis encompasses the entire range of mutation rates found in the

genome. We acknowledge that an ideal future comparison would involve other amylase-like loci,

which exhibit similar mutational mechanisms and levels of structural variation, and that are

resolved using comparable approaches once such databases become available. In the meantime,

when we repeated this analysis for the complex 3q29 locus, known for its segmental duplication-

rich nature and high levels of structural variation and resolved with similar approaches (*15*, *19*),

we found 11 singletons (fig. S7). Thus, the amylase locus is mutating faster than a typical structural

variation hotspot and 98.3% of analyzed tandem repeats.

To understand how much of all amylase haplotypes that exist in the human population has

been captured by our study, we conducted rarefaction analysis on the 98 samples and found that

we have identified all common haplotypes with a frequency of >= 5% (Fig. 1E and fig. S8, A and

B) and 86% of all haplotypes overall (fig. S9).

Strong negative selection limits functional variation among amylase gene copies

To systematically assess the selection pressure on the amylase coding sequences, we examined the

degree of protein-coding sequence variation associated with our high-confidence amylase

haplotypes (30 haplotypes from 117 alleles). Gene annotation predicted 582 distinct intact protein-

coding amylase gene copies in 117 alleles and we experimentally validated these copy number

predictions using digital droplet PCR on 18 randomly selected individuals ($R^2 = 0.94$; fig. S10A

and table S4, S5). The reconstruction of the coding sequence phylogeny revealed that all human

amylase gene copies can be robustly clustered into three distinct types: *AMY2B*, *AMY2A*, and

AMY1 (bootstrap value = 96%; Fig. 2A and fig. S11). We found that the *AMY2B*, *AMY2A*, and

AMY1 genes had 23, 23, and 36 fixed coding sequence variations unique to each type, resulting in

6, 11, and 19 gene-specific amino acid differences, respectively (table S6). Based on the coding

sequence alignment (Data Files-S2), we estimated the synonymous and nonsynonymous

substitution (dN/dS) ratios using codeml (*20*), where we found no evidence for lineage-specific

selection pressure acting on any of the amylase gene types (FDR adjusted p-value from χ^2 test >

0.05; table S7). In contrast, all amylase gene types show significant signatures of negative selection

(FDR adjusted p-value from χ^2 test < 0.05; Fig. 2A and table S7). These observations suggest that

negative selection (dN/dS ratio < 1) has acted to retain the amino acid sequence of amylase gene

copies, both within and between the three amylase gene types. It is of note that we identified two

amino acid variants at positions 211 and 366 (Accession: P0DUB6) that could contribute to

functional differences and may have biomedical significance. (Fig. 2B, fig. S12, and table S8).

Fig. 2. The variants in amylase coding sequences and negative selection on three amylase gene types. (**A**) The maximum likelihood phylogenetic tree of amylase coding sequences (left) and dN/dS estimate for each amylase gene type (right). The phylogenetic tree is rooted with a coding sequence from the sheep genome (Oar_rambouillet_v1.0). The number of nucleotide and resultant amino acid changes that are paralog-specific are indicated. The numbers in parenthesis indicate nucleotide and amino acid changes that are variable within the AMY1 branch. The numbers to the right of each bar represent the FDR-adjusted p-value for the likelihood ratio between $H0_2$ (dN/dS ratio is fixed to one on the foreground branches) and H1 (two dN/dS ratios are allowed on the foreground and background branches, respectively) for each gene type. (**B**) The positions of amino acid variants within and between *AMY2B*, *AMY2A* and *AMY1* protein sequences. The 211 and 366 positions are highlighted because they overlap with a conserved section of the amylase protein sequence and have a predicted functional impact (AlphaMissense (*21*). The coordinates are based on the residues of the amylase enzyme from UniProtKB (Accession: P0DUB6).

Our coding sequence data from gene annotations covering 582 intact amylase gene copies confirmed the explicit difference between *AMY1, AMY2A,* and *AMY2B* genes, which is crucial to distinguish the expression of these genes across tissues. Indeed, according to existing data portals, Genotype-Tissue Expression and The Human Protein Atlas, *AMY2A* and *AMY2B* are expressed in the pancreas and, to some level, in adipose and brain tissues, while the *AMY1* gene is expressed primarily in the parotid salivary gland (fig. S13). These observations are consistent with the idea

that *AMY1* first emerged in the ancestor of great apes, resulting in a gain of expression in the

parotid gland tissues (*5*). Furthermore, subsequent *AMY1* gene duplications in the human lineage

appear to affect the dosage of amylase in the parotid salivary glands (*1*). Among the haplotypes

we identified in this study, we found 110 amylase pseudogenes and showed they share a single

phylogenetic origin from an ancestral incomplete gene duplication of the *AMY2A* gene (*22*) (fig.

S14). Thus, the pseudogenization of *AMY2A* is more likely due to a single mutational event rather

than a loss of constraint and the repeated occurrence of novel loss-of-function variants.

Evolution of the copy number of AMY1 gene

We showed above that all human amylase gene copies can be robustly clustered into three distinct

types: *AMY2B*, *AMY2A*, and *AMY1*. However, to specifically study the evolution of the copy

number of the salivary *AMY1* gene, we needed to identify the sequences within the amylase locus

that were the most phylogenetically informative. To achieve this, we systematically evaluated the

haplotype variation in 117 alleles by aligning sequences and identified a consensus sequence for

each amylase segment (fig. S15, A and B). We identified an interval (from 22,850 to 26,730 bp;

Data Files-S2) within the AMY1 segment (fig. S15C), where all observed AMY1 segments (n =

337) can be phylogenetically structured into three distinct clusters; AMY1A (n = 124), AMY1B

(n = 99), and AMY1C (n = 114) (fig. S15D). These clusters correspond to the segments represented

in the GRCh38 assembly. Further, the chimpanzee (panTro6) and gorilla (gorGor6) reference

genomes each contain only one AMY1C-like segment diverged from human H1[a].1 haplotypes

(figs. S16 and S17). These findings suggest that the common ancestor of humans and chimpanzees

possessed a single AMY1C-like segment, and the AMY1A and AMY1B segments have evolved

only in the human lineage (fig. S17). Note that the bonobo reference genome (panPan3) contains

two AMY1 segments: one ancestral and one resulting from an independent, bonobo-specific

duplication, as determined through synteny and phylogenetic analysis (figs. S16 and S17). One of

these duplications is likely non-functional due to a previously reported a disrupted coding sequence

(*1*), a finding corroborated by the most recent annotation (RS_2024_02/NHGRI_mPanPan1-

v2.0_pri).

To further understand the evolution of the *AMY1* gene copy number, we aligned all AMY1

segments from the most common haplotypes, H3^r.1 and H3^r.2, that harbor all three AMY1 segment

types. Two independent Bayesian phylogenies based on these alignments indicate that the AMY1B

segment arose from the AMY1C approximately 140 to 270 kilo (thousand) years ago (KYA),

followed by the duplication of AMY1A from the AMY1B segment approximately 120 to 240

KYA (fig. S18 and table S9). Gene conversion between the GC-rich segmental duplications

complicates time estimation based on a molecular clock and is a known phenomenon at the

amylase locus (*6*, *23*). Considering gene conversion between AMY1 segments, the actual

duplication dates are expected to be older than the estimates above. Some studies have used single

nucleotide variants in the flanking regions to infer the phylogenetic history of this locus, positing

coalescence dates for human amylase locus at ~279 and ~450 KYA (*11*, *12*). However, as

described previously (*7*), we found that the linkage disequilibrium between the flanking single

nucleotide variants and structural haplotypes is low (e.g., H1^a.1 average: $R^2 = $ ~0.26 and median

$R^2 = $ ~0.03) (fig. S19), complicating the time estimation of *AMY1* duplications using flanking

regions. Therefore, our estimates avoid these complications and support the conclusion that the

initial *AMY1* gene duplications substantially predated out-of-Africa migrations, by at least 30 KYA

(table S9).

Fig. 3. Amylase gene duplication and the history of agriculture. (**A**) The read-depth of amylase locus spanning *RNPC3*, *AMY2B*, *AMY2A*, and *AMY1* genes (chr1:103494306-103668306) for GoyetQ56-1 (maroon line), a Neanderthal excavated in present-day Belgium, showing signatures of *AMY1* duplication, and for Denisova 11 (beige line), a hybrid hominin (Neanderthal and Denisovan) excavated from present-day Russia, showing signatures consistent with an ancestral single-copy AMY1 haplotype. The maroon and beige lines indicate average read-depths with 5-kbp window and 1-kbp step for each sample. The average read-depth of each 5-kbp window was normalized by the average read-depth of the *RNPC3* gene. Only uniquely mapped reads were used for this visualization. (**B**) A world map displaying the locations of ancient human samples. Sample locations are indicated with an 'X', with corresponding hexagons showing the estimated *AMY1* copy number. Carbon dating (Kilo Years Ago; KYA) estimated for each sample is indicated in blue. (**C**) Amylase copy number estimations from Europeans who were farmers (beige) and hunter-gatherers (maroon). Samples are binned on the x-axis according to three time periods; pre-agriculture (before the transition to agriculture, >9 KYA), during the transition (~ 9 - 3 KYA), and post-transition (complete transition to agriculture, ~ 3 KYA - 900 years ago). The shape of data points corresponds to sample dating estimates (Xs for more than 9 KYA, circles for 9 - 3 KYA, and triangles for 3 KYA - 900 years ago). The inset of the right panel shows nonparametric regression lines for *AMY1* copy number across time, for hunter-gatherers (maroon) and farmers (beige), with confidence intervals in lighter corresponding color, respectively. (**D**) Zoomed-in map showing the spread of agriculture into Europe from Asia. Major agricultural footholds are indicated by wheat pictograms. Tan arrows show general trends of human agricultural migration throughout Europe, with predicted time periods (*24*, *25*). Ancient human samples that were

266 analyzed for amylase copy number are annotated with shapes and colors for time period and
267 lifestyle, respectively.
268

AMY1 copy number variation in archaic hominin genomes

A complementary approach for estimating the relative timing of gene duplications involves

analyzing the read-depth of unique k-mers in ancient human and archaic hominin genomes. We

first tested a k-mer approach using the GeneToCN algorithm on short-read sequencing data (*26*)

to estimate the copy numbers of the *AMY1*, *AMY2A*, and *AMY2B* genes in 116 present-day human

genomes (table S5). Notably, the k-mer approach achieved an $R^2 > 0.99$ correlation for 32

individuals, each with both haplotypes of the amylase locus reconstructed (fig. S10B). We tested

our k-mer approach in a separate cohort of 101 samples with digital PCR validation data ($R^2 =$

0.95; fig. S10A). These results suggest that the GeneToCN algorithm may be more accurate than

digital PCR estimations and a viable option for estimating *AMY1* gene copy numbers in short-read

whole genome datasets.

We then aimed to estimate *AMY1* gene copy number in eight archaic hominin genomes

using two approaches: (1) the validated k-mer method described above, as well as (2) an

independent read-depth analysis (table S10). Given the varying genome-wide coverage in most of

these genomes, we needed to conduct a downsampling analysis to empirically determine that 1X

and 5X genome-wide coverage provides >85% and >95% accuracy in estimating *AMY1* copy

number, respectively (fig. S20 and table S11). Taking into account GC bias and coverage within

the amylase locus for each sample (see Methods), we were able to reliably estimate the *AMY1* copy

number for eight archaic hominin genomes (table S10). We found increased *AMY1* copy number

in two Eastern and one Western Neanderthals, as well as one Denisovan genome (Fig. 3A, fig. S21

and table S10). These include the Altai Neanderthal (2.6 copies), Denisova 2 (8 copies),

GoyetQ56-1 (5.0 copies), and Mezmaiskaya 2 (4.7 copies). Previous read-depth analysis of the

Altai Neanderthal genome found no evidence for an increase of *AMY1* gene copy number (*10*, *27*).

By incorporating eight archaic hominin genomes, we have now detected signatures of *AMY1* gene

duplication in some of the Neanderthals and Denisovan genomes, indicating that *AMY1* duplication

was actually polymorphic among archaic hominins.

The *AMY1* duplication in these archaic hominins could be postulated by four scenarios.

First, it is possible that because of the complexity of the locus and the complications inherent in

archaic hominin genomic sequencing, there may be a technical bias in our detection. However, the

observation of duplications using two different approaches and in multiple genomes provides

confidence in our results. Second, it is plausible that introgression into archaic hominins from

humans may explain the presence of duplications in the former. While still a possibility to be tested

when more archaic hominin genomes are available, we observed the duplication in western

Neanderthals (e.g., Goyet), which shows no signature for introgression from humans (*28*, *29*).

Third, it is plausible that the duplication evolved independently in the archaic hominin lineage.

However, we argue that two independent duplications (one in humans and another in archaic

hominins) in less than a million years is unlikely, given that initial duplications from single *AMY1*

copy haplotypes are rare in great apes (*4*) (fig. S16). The fourth and, in our opinion, the most

plausible scenario is that the *AMY1* gene might already have been copy number variable before the

human-Neanderthal/Denisovan divergence (~800 KYA (*30*)), albeit to a limited extent as

compared to what is observed in present-day humans. Overall, our results suggest a complex

history of amylase duplications, which will be further scrutinized as more high-coverage archaic

hominin genomes become available.

The frequency of higher copy number of AMY1 has increased in the last 10,000 years,

coinciding with the agricultural transition

To explore the changes in allele frequency of *AMY1* copy number since the migration out of Africa

60 - 90 KYA, we analyzed the genomes of 73 ancient human genomes (table S12). The oldest

genome analyzed was the Ust'Ishim sample from Siberia (~45 KYA), which has six *AMY1* gene

copies per diploid cell. The Peştera Muierii sample from Romania (~34 KYA) has eight *AMY1*

gene copies per diploid cell. This indicates that high *AMY1* gene copies (*defined here as >= 6*

copies per diploid cell) had already spread across Eurasia as far back as ~45 KYA (*31*)) (Fig. 3B).

We next analyzed the ancient human genomes in the context of agricultural development

and found a general trend where the *AMY1* gene copy number is significantly higher among

samples excavated from archaeological agricultural contexts compared to those from hunter-

gatherer contexts (p-value = 0.037; fig. S22). To further investigate the intricacies of this trend,

we examined ancient human genomes from Europe, where we have a clear timeline of the

Neolithic transition. Specifically, based on archaeological evidence, we categorized genomes

based on three distinct European time periods: the pre-agricultural period (> 9 KYA), where all

our samples are from hunter-gatherers; the agricultural transition period (9 to 4 KYA), representing

the long period of transition to agriculture in Europe; and the agricultural period, during which no

hunter-gatherers are recorded in our sample datasets (< 3 KYA). These period choices were made

with the assumption that the putative adaptive effects of agricultural diets would take some time

to become visible at the population level. We found that pre-agricultural genomes already harbored

four to eight *AMY1* copies per diploid cell (table S12). We also observed a consistent and

significant increase in *AMY1* copy numbers across these periods (*e.g.*, during vs. post p-value <

0.01; Fig. 3, C and D), and found similar trends for the *AMY2A* genes (Fig. 3C) and non-European

samples (figs. S23 and S24). These findings support the notion that haplotypes with higher

numbers of amylase gene copies have increased in frequency over the last 10 KYA. We found no

significant differences in *AMY1* copy number between samples excavated from agricultural versus

hunter-gatherer archaeological contexts during the "agricultural transition" period when farmers

and hunter-gatherers shared the same habitat.

Taken together, these findings are consistent with the idea that either neutral evolution or

weak-adaptive forces, perhaps due to preagricultural experimentation with food processing

techniques (*32*), such as flour production from wild cereals (*33*), have retained the wide range of

standing *AMY1* copy number variation in preagricultural Europe. The gradual increase in starch

availability as Europe transitioned to an agricultural lifestyle (*34*) likely underlies the observable

adaptive force acting on high copy number haplotypes, explaining the increase of *AMY1* copy

number in late post-agriculture European populations.

Multiple mutational origins that resulted in a burst of copy number variation since the initial

AMY1 duplication

We next investigated the likely mutational origins of present-day amylase haplotypes. To do this,

we first examined the relationship among the four most common haplotypes, H3^r.1, H3^r.2, H3^r.4,

and H1^a.1. Building upon the previously reported link between one-copy and three-copy

haplotypes (*35*) and the fact that we observe only three types of AMY1 segments, we propose an

evolutionary model linking the ancestral chimpanzee-like haplotype (H1^a-like) to the common

three-copy haplotypes (H3^r.1 and H3^r.2) (fig. S25). According to this model, the initial

duplications of *AMY1* starting from H1^a-like ancestral haplotype led to the emergence of H3^r

haplotypes. Given that multiple mutational steps are required to move from one-copy common

haplotypes to other common haplotypes and that we do not observe any intermediate haplotypes

in the present-day human genomes, the duplication from one-copy to three-copy haplotypes is

likely to have occurred only once in the human lineage. This is also supported by the absence of

duplications in the amylase segments within the H1^a.1 haplotype, which would impede non-allelic

homologous recombination (NAHR) events. In contrast, H3^r.1 and H3^r.4 haplotypes harbor copies

of identical and unidirectional sequences (AMY2A.2 segments), providing an ideal template for

recurrent NAHR events leading to the diverse haplotypes we see today.

To further investigate the mutational relationships between the H3^r haplotypes and other

extant haplotypes, we utilized dotplots and sequence alignments to delineate the breakpoints of

structural differences (Supplementary Results). In parallel, we conducted a scan for PRDM9-

binding motifs within the amylase locus to pinpoint possible recombination sites (Fig. 4A, fig. S18

and table S13). By integrating all these observations, we were able to construct a putative

evolutionary path with the fewest plausible mutational steps that can explain the origin of present-

day amylase haplotypes starting from NAHR-prone H3^r haplotypes (fig. S26, S27; Supplementary

Results). Our proposed evolutionary model of mutational events is consistent with our hypothesis

regarding the central role of the H3^r haplotypes in seeding extant haplotypic variation and offers

three major insights into the evolution of the amylase locus in humans.

Fig. 4. The evolutionary and mutational connections among common haplotypes. (**A**) The structural variation breakpoints and recombination hotspots in the amylase locus. The colored arrows represent amylase segments. The PRDM9 binding sites are represented with red dots. The non-allelic homologous recombination (NAHR) breakpoints are represented with purple, green and orange dots, and dashed lines. The microhomology-mediated break-induced replication breakpoints are represented with gray dots. (**B**) NAHR-mediated duplication and deletion of the *AMY1A-AMY1B* cluster. The AMY2.2 (orange) segment serves as the recombination substrate for the crossover, resulting in the duplication or deletion of the *AMY1A-AMY1B* cluster as illustrated in the middle panel. Chimeric AMY2.2 segments have been identified, utilizing parsimony informative sites within the AMY2.2 segment (right panel). (**C**) Microhomology-mediated break-induced replication based copy-number gain resulting in the formation of H2A2B2.1. The middle panel shows the mutational mechanism. Four nucleotides of microhomology internal to the breakends were identified at the breakpoint junction (right panel).

First, we found evidence for recurrent NAHR events among common haplotypes (H3^r.1, H3^r.2, and H3^r.4) harboring the AMY1A and AMY1B segments with breakpoints in the AMY2A.2 (orange) segment. These NAHR events, which may occur among different haplotype combinations, could concurrently result in the duplication and deletion of two *AMY1* gene copies (*e.g.*, Fig. 4B and fig. S28; Supplementary Results). Therefore, while other less likely scenarios

are possible, NAHR-based deletion and duplications underlie copy number variation of the AMY1

segments and thus *AMY1* genes. Specifically, as this proposed mechanism always adds or deletes

two copies of the *AMY1* genes, our finding explains how most human haploid genomes harbor

odd-numbered *AMY1* gene copies (*7*) (fig. S29). It would seem that the majority of H1^a.1

haplotypes among present-day humans have predominantly originated from H3^r haplotypes. If

true, this hypothesis explains the homoplastic occurrence of H1^a haplotypes across the amylase

phylogenetic tree (fig. S19C) and the lack of an out-of-Africa signal in H1^a nucleotide diversity,

which would be expected if the H1^a haplotypes arose before out-of-Africa migrations (table S14).

Therefore, we argue that although the H1^a.1 haplotype is structurally identical to the ancestral

human amylase haplotype, the current versions in our datasets have arisen recurrently from H3^r

haplotypes.

Second, we characterized three microhomology-mediated break-induced replication events

and identified the accompanying microhomologies at the breakpoint junctions (Fig. 4C;

Supplementary Results). Even though these three rearrangements constitute only five alleles

(H2A2B2.1, H3^r.6, and H3B2.1) (~ 4%), they hold substantial biological relevance since

H2A2B2.1 and H3B2.1 harbor duplications of the *AMY2* genes. Thus, different mechanisms drive

the copy number variation of the salivary *AMY1* genes and pancreatic *AMY2* genes, with the lower

copy number variation of *AMY2* genes explained by the slower rate of non-recurrent

microhomology-mediated break-induced replication events (*36*).

Third, recurrent NAHR-mediated inversion events at the amylase locus, similar to those

described previously (*37*), underpin the mutational connections between the common H3^r.1, H3^r.2,

and H3^r.4 haplotypes (fig. S30), as well as several other inversions among extant haplotypes

(Supplementary Results). Given that inversions underlie the structural differences between the

majority of the distinct haplotypes, their functional and adaptive relevance presents an intriguing

avenue for future research.

Discussion

In this study, we have dissected the evolution of the amylase locus. First, we hypothesize that the

initial duplications of the *AMY1* genes occurred once through multiple duplications, evolving from

a one-copy ancestral haplotype to the three-copy present-day haplotypes frequently observed in

our dataset. Analysis of archaic hominin genomes suggests that these initial duplications occurred

well before the split of the human-Neanderthal/Denisovan. This observation is concordant with

the recent evidence of Neanderthal starch consumption (*38*), and perhaps the availability of cooked

starch in archaic hominins made possible through the domestication of fire (*39*) (Fig. 5).

Second, we hypothesize that selection acted on abundant standing *AMY1* copy number

variation at this locus rather than on *de novo* variants. We observed a wide range of *AMY1* gene

copy number variation (3 - 9 copies) in samples that predate agriculture. We further found that late

agricultural populations consistently harbored amylase haplotypes with higher *AMY1* copy

numbers (Fig. 5). Yet, our data do not indicate strong, pervasive selection. Instead, we propose

that partial soft sweeps involving pre-existing amylase structural haplotypes may have influenced

AMY1 copy number variation in relation to historical starch consumption patterns in different

populations. The effects of amylase gene duplications on taste preferences and starch metabolism

may have also predisposed humans to prefer and tolerate consuming wild grains, as reported for

the Mesolithic groups in the Balkans (*40*), facilitating the adoption of starch-rich diets and eventual

domestication of these plants. Overall, our results support the complex narrative of the transition

to agriculture, which includes the replacement of western hunter-gatherers by Anatolian farmers

in Europe (*41*), potentially bringing with them amylase haplotypes that harbor higher *AMY1* gene

copies. Similarly, transient interactions between hunter-gatherer and agricultural groups (*42*) could

explain the similar copy numbers observed between these groups during the transition period.

Future comparative analyses focusing on partial soft sweeps in indigenous populations with

recently diverged dietary habits involving heavy starch consumption—such as the Andeans (*43*,

44) could provide further insights into these signatures of selection at the human amylase locus.

Third, we found that NAHR and microhomology-mediated break-induced replication

underlie the copy number variation of *AMY1* and *AMY2* genes, respectively, explaining the

different rates of their structural evolution. The extremely high rate of structural variation due to

NAHR led to remarkable *AMY1* copy number variation with distinct mutational propensities. For

example, our evolutionary model involving common H3^r haplotypes suggests recurrent NAHR

events mediated by highly similar sequences (> 99%), resulting in duplications or deletions of both

AMY1A and *AMY1B* genes. In contrast, the H1^a.1 haplotype harboring a single *AMY1* copy and

divergent *AMY2* genes rendering it less susceptible to NAHR events. Therefore, the mutation types

and rates of the amylase locus may differ depending on extant haplotype variation in a population,

especially in bottlenecked populations such as indigenous Americans (*45*). It is a distinct

possibility that a bottlenecked population ends up with a very high frequency of H1^a.1 due to drift.

In this case, the absence of segments with highly similar sequences within H1^a.1 haplotype would

mitigate recurrent NAHR events, resulting in a slower accumulation of variation in this population.

In contrast, if one of the larger amylase haplotypes were to become prevalent due to drift, the rate

of variation would increase exponentially. Within this general context, one interesting question for

future work is whether larger amylase haplotypes experience negative selection due to increased

genomic instability. Taken together, our study underscores how gene duplications in early human

history provided the genetic foundation for dietary flexibility during agricultural innovations,

contributing to modern human evolution.

Fig. 5. An evolutionary model of the human amylase genes and resulting hypotheses. Top: A timeline of human amylase locus evolution based on the results of this study, with relevant events indicated on top. Middle: A schematic view showing the increase in *AMY1* copy number variation and the mean number of *AMY1* copies in present-day human populations throughout history as starch consumption increases. Bottom: A phylogenetic representation of the hypothesized amylase duplication timeline.

Materials and Methods

Sample Collection and Datasets

Samples (n = 98) from diverse populations studied by the Human Genome Structural Variation

Consortium (HGSVC) (*19*), Human Pangenome Reference Consortium (HPRC) (*46*) and Genome

In a Bottle (GIAB) (HG002) datasets were included in this study. These datasets consist of African

(n = 43), American (n = 24), East Asian (n = 13), European (n = 9), and South Asian (n = 9) (table

S2) individuals. Bionano Genomics optical genome mapping and PacBio HiFi datasets of HGSVC

samples were downloaded from

http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/HGSVC2/working/ and

http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/HGSVC3/working/, and HPRC and

GIAB samples containing phased assemblies and optical genome mapping datasets were

downloaded from https://s3-us-west-2.amazonaws.com/human-

pangenomics/index.html?prefix=working/).

Amylase Segments

We characterized a ~212.5-kbp region on chromosome 1 (GRCh38; chr1:103,554,220–

103,766,732) overlapping paralogous copies of segmental duplications that have > 99% sequence

similarity. The amylase segments, depicted by colored arrows, were modified from a previous

study by Usher and colleagues (*7*), and determined based on the labeling pattern from the Bionano

Genomics optical genome mapping data and the sequence similarity between segments using

blastn (BLASTN 2.9.0+) (*47*). First, using optical genome mapping *in silico* map labeling patterns

of GRCh38 reference assembly, amylase segments were identified using pairwise alignments

between smaller duplicons within segmental duplications. Then, to obtain accurate coordinates of

segments, the GRCh38 fasta sequence of the locus was aligned against itself using blastn

(BLASTN 2.9.0+). The alignments with less than 99% sequence similarity and 60% coverage were

filtered out and coordinates of all reference segments were finalized. We identified one copy of

AMY2B (purple; 45.7-kbp), two copies of Intergenic1 (green; 5.3-kbp), one copy of AMY2A.1

(red; 12.4-kbp), two copies of AMY2A.2 (orange; 14.1-kbp), three copies of Intergenic2 (maroon;

4.3-kbp), and three copies of AMY1 (pink; 26.3-kbp) segments within the GRCh38 amylase locus

(table S1). Next, the GRCh38 segment sequences were used as a guide to detect amylase segments

from each sample included in this study using blastn. Same sequence similarity (> 99%) and

coverage thresholds (> 60%) were used to obtain the amylase segment coordinates from each

sample. Amylase segments in nonhuman primates, chimpanzee (Clint_PTRv2_panTro6), bonobo

(Mhudiblu_PPA_v0_panPan3) and gorilla (Kamilah_GGO_v0_gorGor6), were detected by using

the same approach described above with sequence similarity threshold of >= 90% using genome

reference assembly fasta files that were obtained from University of California Santa Cruz Genome

Browser.

De novo assembly of optical genome map dataset

Optical genome maps of HGSVC and HPRC samples were *de novo* assembled and using the

Bionano Solve v3.5 (https://bionanogenomics.com/support/software-downloads/) assembly

pipeline, with default settings as described previously (*48*) (fig. S31A).

```
517   python2.7 Solve3.5.1_01142020/Pipeline/1.0/pipelineCL.py -T 64 -U

518   -j 64 -jp 64 -N 6 -f 0.25 -i 5 -w -c 3 \

519   -y \

520   -b ${bionano_bnx} \
```

```
521  -l ${output_dir} \
522  -t Solve3.5.1_01142020/RefAligner/1.0/ \
523  -a
524  Solve3.5.1_01142020/RefAligner/1.0/optArguments_haplotype_DLE1_s
525  aphyr_human.xml \
526  -r ${reference_genome}
```

A pairwise comparison of DNA molecules (min. 250-kbp) was generated to produce the

initial consensus genome maps. During an extension step, molecules were aligned to genome

maps, and maps were extended based on the molecules aligning past the map ends. Overlapping

genome maps were then merged. Extension and merge steps were repeated five times before a

final refinement of the genome maps. Clusters of molecules aligned to genome maps with

unaligned ends > 30-kbp in the extension step were re-assembled to identify all alleles. To identify

alternate alleles with smaller size differences from the assembled allele, clusters of molecules

aligned to genome maps with internal alignment gaps of size < 50-kbp were detected, and the

genome maps were converted into two haplotype maps. The final genome maps were aligned to

the reference genome, GRCh38.p12.

Haplotype detection and single molecule support using optical genome maps

To resolve haplotype structures at the amylase locus, *de novo* assemblies of optical genome maps

of samples were first aligned to the GRCh38.p12 human genome reference assembly. Next,

alignment results were visualized using the Bionano Access™ software to resolve amylase

haplotypes. Haplotypes were identified from these visualizations using amylase segments and *in*

silico map labeling patterns as a guide. Next, molecule support for each haplotype was evaluated

using the following steps (fig. S31B): contigs aligning to the region of interest (ROI) (GRCh38;

chr1:103,554,220–103,766,732) were detected. Any samples with greater than or less than two

contigs aligning to the ROI were excluded. Next, ROIs in each contig were identified and extended

10-kbp on both proximal and distal regions. The number of molecules aligning to the ROI of the

amylase locus was collected, and the average molecule length was calculated for each contig. If

the average molecule length was less than the ROI, the number of molecules that were anchored

in the proximal and distal unique regions was counted. Next, the length of the ROI was divided by

the average molecule length, and new smaller ROIs were identified based on quotient, and the

number of molecules was counted for each new ROI. Only haplotypes supported by three or more

molecules were considered accurate haplotypes (Data File-S1).

De novo assembly and orthogonal support

PacBio HiFi long-read sequencing dataset from HGSVC (n = 60) was *de novo* assembled using

hifiasm (*49*) with the following command:

```
hifiasm -o ${outputdir}${sampleID}.asm -t 32 ${reads}
```

559   Publicly available phased assemblies from the HPRC were analyzed to validate the optical genome

560   mapping based haplotype structures, as described previously (*15*). Validation process for both

561   HPRC and HGSVC assemblies includes the following steps: First, fasta files of HPRC phased

562   assemblies and HGSVC hifiasm assemblies were converted to *in silico* maps using

563   fa2cmap_multi_color.pl (Bionano Solve™ v3.5.1). The resulting *in silico* maps were aligned to

564   the GRCh38p.12 reference assembly *in silico* map using the refAligner tool (Bionano Solve™

565   v3.5.1).

```
566   Solve3.5.1_01142020/RefAligner/1.0/RefAligner                -ref

567   ${referenceFile} \ -maxthreads ${threads} \

568   -i ${inputfile} \

569   -o ${outputdir}EXP_REFINEFINAL1
```

570   Next, a visual pairwise comparison of HPRC optical genome mapping and phased

571 assemblies for each sample was performed using Bionano Access™. Then, amylase segments in

572 each sample with a haplotype consistent with the optical genome mapping haplotype were detected

573 using blastn (BLASTN 2.9.0+), as described in the 'Amylase Segments' part. Next, all

574 orthogonally supported haplotype alignments were visualized using pygenomeviz (*50*) to confirm

575 haplotype structures. Finally, haplotypes were clustered using pgr-tk v0.5.1 (*51*) (fig. S4).

577 ***Significance of AMY1 copy number distribution across populations***

578 To evaluate whether the *AMY1* gene copy numbers exhibit population-specific patterns, we

579 performed Kruskal-Wallist rank sum test. This non-parametric test was used to determine if there

580 are statistically significant differences between the medians of three or more independent groups.

581 We chose this test due to its robustness in handling non-normally distributed data. We conducted

582 our analysis on R Studio (2022.07.2 Build 576). Below are the commands we used for the analysis:

```
583   counts <- c(174, 79, 36, 28, 23) #AMY1 copy number counts

584   populations <- factor(rep(c("AFR", "AMR", "EAS", "EUR", "SAS"),

585   times = c(1, 1, 1, 1, 1)))

586   kruskal_test_result <- kruskal.test(counts, populations).
```

588 ***Distribution of singleton proportions for genome-wide tandem repeat loci***

589   To quantitatively compare the fraction of singletons in the amylase locus with other fast-evolving

590   loci, we downloaded genotypes of tandem repeats (TR) loci from (https://github.com/gymrek-

591   lab/EnsembleTR), not taking into account the length of repetitive elements, and selected diploid

592   genotypes of individuals present in both the TR dataset and our dataset (33 individuals, 66 alleles).

593   Our study comprises 81 samples with 30 high-confidence haplotypes. Among these, 33 samples

594   are also present in the EnsembleTR dataset. By comparing data from the same individuals, we

595   eliminated the effect of genetic drift in our comparison. From the 33 samples, 21 out of the 30

596   amylase haplotypes were identified. We then selected 719 TR loci that have the same number of

597   distinct alleles (21 alleles from 33 individuals) as our comparative data. The remaining nine alleles

598   from the high-confidence haplotypes (n = 30) correspond to the alleles detected from samples that

599   are not present in the TR dataset. For the selected loci, the distribution of singleton proportions

600   was generated by dividing the number of singletons by the total number of alleles (n = 21) for each

601   locus. The empirical p-value is represented as the proportion of TR loci with a singleton proportion

602   greater than or equal to that observed at the amylase locus (n = 14). Note that the number of

603   singleton haplotypes (n = 19) that we detected as part of the high-confidence haplotypes were

604   reduced to 14 since the EnsembleTR didn't have samples that carry the remaining five singleton

605   haplotypes. For a comparison, we repeated this analysis for 3q29 locus, known for its segmental

606   duplication-rich nature and high levels of structural variation similar to the amylase locus. For

607   adequate comparison, we used the same individuals (n = 39) for whom we were able to reconstruct

608   3q29 haplotypes for both of the chromosomes. Haplotype information for the 3q29 locus was

609   obtained from a previous study (*15*). In addition, we filtered the tandem repeat loci (601 loci (unit

610   length 1-6 bp)) from EnsembleTR that we analyzed to match the number of distinct alleles (n =

611   22) that we observed in the 3q29 locus.

Rarefaction analysis

Rarefaction analysis was performed using the R package called "iNEXT" (*52*) to evaluate

haplotype diversity and determine whether the number of samples included sufficiently captures

majority of diversity at the amylase locus. The following command was used for both all amylase

haplotypes (n = 51) and four common amylase haplotypes:

```
Data <- read.table("rarefaction_allPopsCombined.txt", header =
TRUE, sep = "\t")[6,]

Obj <- iNEXT(as.numeric(Data), datatype="abundance", q=0,
knots=sum(Data)*2)

Temp_data <- Obj$iNextEst$size_based

Rate <- c()

for (i in 1:nrow(Temp_data)){

    Rate[i+1] <- Temp_data[i+1,5]-Temp_data[i,5]

}

Plot_data <- data.frame(NumChr=Temp_data$m, NumHap=Temp_data$qD,
Rate=Rate[-length(Rate)])

pdf("Sample-size-based_REcurve_with_rate_of_change.pdf", width=6,
height=4)

    twoord.plot(Plot_data$NumChr,Plot_data$NumHap,Plot_data$Num
Chr,Plot_data$Rate, type="l", xlab="Number of alleles",
```

```
ylab="Haplotype         count",        rylab="Rate        of        change",

lytickpos=seq(0,70,10), lcol="blue3",rcol="red2")

abline(v=sum(Data), lty="dashed")

dev.off()

pdf("Coverage-based_REcurve.pdf", width=6, height=4)

ggiNEXT(Obj, type=2)+

639           theme_bw()+

640           theme(legend.position="none")+

641           xlab("Number of alleles")

642   dev.off()

643
```

644   ***Amylase gene annotation using protein sequence homology***

645   To predict amylase gene coordinates and copy number on each haplotype, we used homology-

646   based gene prediction tool ("protein2genome") implemented in Exonerate v2.4.0 (*53*) with default

647   settings except for maximum intron length of 20-kbp (`--maxintron 20000`). For the

648   homology search, we used the amylase protein sequences (n = 3) with experimental evidence from

649   UNIPROT ("Evidence at protein level") (*54*) under the following accession numbers; *AMY2B*:

650   P19961, *AMY2A*: P04746, *AMY1A*: P0DUB6, *AMY1B*: P0DTE7, and *AMY1C*: P0DTE8 (The

651   protein sequences of *AMY1A*, *AMY1B*, and *AMY1C* are identical). We clustered any overlapping

652   hits from the prediction results and selected the best hit from each cluster. The hits that can be

653   translated into 511 full-length polypeptide sequences were then kept as a final annotation. In cases

654 where we found conflict between the homology-based annotation and the predicted copy number

655 of *AMY* genes from amylase haplotypes, the homology-based prediction was manually curated to

656 ensure it to be consistent with the optical mapping label. Additionally, genome annotations of *de*

657 *novo* assemblies were obtained by using the T2T-chm13 gff annotation file with liftoff (*55*).

658 `liftoff -db $dbfile -o $outputfile.gff -u $outputfile.unmapped -`

659 `dir $outputdir -p 8 -m $minimap2dir -sc 0.85 -copies $fastafile -`

660 `cds $refassembly`

661   The liftoff results were compared with homology-based annotation above and were used

662 when the manual curation was not able to produce annotation consistent with the optical mapping

663 label , where we found only one case (HG01175_H7.1; table S4). Lastly, pseudogene annotations

664 were obtained from the liftoff annotation files (table S4).

666 *Validation of AMY1 copy number using digital droplet PCR*

667 Custom primer and probe oligos were designed to target a conserved sequence within *AMY1* gene

668 in exon 5 across all annotated amylase gene copies (avoiding pseudogenes) in the T2T-chm13

669 human reference genome, with the following sequences: forward primer

670 (`CCTCAACAGGTCAGAGATTGTC`), reverse primer (`GGCAATCTTAGAACGCACATAATC`), and

671 probe (`TGGTCTTCTCGATCTTGCACTGGG`). The ddPCR protocol, reference primer, and reagents

672 used are as previously described (*4*). Human DNA samples from LCLs (Coriell Institute) of 101

673 individuals (table S5) were used to estimate total amylase gene copies in duplicates. Restriction

674 enzyme *HindIII* (New England Biolabs), was used in the suggested concentration to digest DNA

675 prior to ddPCR.

 Alignments of amylase segments and coding sequences

678   The alignments of each amylase segment and coding sequences (CDS) were constructed using

679   MAFFT v7.310 (*56*) and PAGAN v1.53 (*57*) with --codon option, respectively. As an outgroup,

680   chimpanzee sequences for each segment were prepared by searching a sequence that covers >=

681   30% of corresponding segment of the GRCh38 reference genome with >= 97% sequence similarity

682   in mPanTro3 genome from the Vertebrate Genome Project (https://www.genomeark.org/t2t-

683   all/Pan_troglodytes.html). Only one of the haplotypes ("haplotype 1") from mPanTro3 was used

684   to align segments. The raw alignments of each segment were summarized by averaging

685   frequencies of major alleles (including a gap) across a 200-bp window with a 10-bp step (fig.

686   S15A). We determined the phylogenetically informative interval (from 22,850 to 26,730 bp in the

687   AMY1 segment alignment, Data File-S2) in the AMY1 alignment based on the change point of

688   consensus determined by a statistical method, "AMOC" (*58*) implemented in the R package,

689   "changepoint", and transposable element information in the alignment. The CDSs of *AMY2B*,

690   *AMY2A*, and *AMY1* genes for chimpanzees were retrieved from the Ensembl annotation of panTro5

691   (release 109) (*59*). Additionally, as an outgroup to all amylase CDSs, we incorporated amylase

692   CDS from the sheep genome, Oar_rambouillet_v1.0 (release 109) (*59*) which is known to have

693   only one copy of the amylase gene (*4*). We also generated an alignment of the phylogenetically

694   informative interval of the AMY1 segment from humans and the non-human primates chimpanzee

695   (Clint_PTRv2_panTro6),   bonobo   (Mhudiblu_PPA_v0_panPan3),   and   gorilla

696   (Kamilah_GGO_v0_gorGor6) using MAFFT v7.310.

698   *Phylogenetic reconstruction of AMY1 segments and coding sequences*

699   We used the AMY1 segment, the phylogenetically informative interval of the AMY1 segment,

700   and CDS alignments (see "*Alignments of amylase segments and coding sequences*") to

701   phylogenetically cluster sequences. A maximum-likelihood tree for each alignment was

702   reconstructed using IQ-TREE v2.2.0 (*60*) with the following options: `-m MFP` (*61*), `-alrt`

703   `1000, -B 1000` (*62*). The chimpanzee (mPanTro3) AMY1 segment and CDSs from

704   chimpanzee (panTro5) and sheep (Oar_rambouillet_v1.0) amylase genes were used as outgroups

705   for the AMY1 segment and CDS, respectively. The tree for the phylogenetically informative

706   interval of the AMY1 segment was rooted by midpoint. All the reconstructed trees were visualized

707   using FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

709   *Estimation of dN/dS for amylase genes*

710   To estimate dN/dS ratios, we used codeml implemented in PAML v4.10.6 package (*20*). First, we

711   pruned the tree topology (fig. S11) and CDS alignment (Data File-S2) to keep only unique CDSs

712   that are not identical to each other. Based on the pruned tree and alignment, we employed a branch

713   model (H1), which allows the dN/dS ratio to vary among branches (model = 2, NSsites = 0,

714   fix_omega = 0 and omega = 0.4). We estimated dN/dS ratio for each gene type by labeling the

715   clade including all CDSs from the gene type of interest as foreground branches. To obtain

716   statistical significance for gene type-specific selection pressure, we calculated a likelihood for a

717   null branch model ($H0_1$) where dN/dS ratio did not vary among branches (model = 0, NSsites = 0,

718   fix_omega = 0 and omega = 0.4) and performed a χ^2 test with the likelihood ratio between $H0_1$ and

719   H1 models. To obtain statistical significance for negative selection pressure, we calculated a

720   likelihood for a null branch model ($H0_2$) where dN/dS ratio for the foreground branches (e.g.

721   human *AMY1* clade) is fixed to one (model = 2, NSsites = 0, fix_omega = 1 and omega = 1) and

722   performed a χ^2 test with the likelihood ratio between H0$_2$ and H1 models. The raw p-values were

723   adjusted by controlling false discovery rate (FDR) (*63*).

725   *Functional analysis for amylase protein sequences*

726   To examine the potential functional effects associated with differences between different amylase

727   gene types (*AMY1*, *AMY2A*, *AMY2B*), all predicted peptide amylase sequences, derived from

728   amylase haplotypes in this study, were aligned using CLUSTALO (*64*). Amino acid differences

729   that were present in six or more haplotypes within and between amylase sequences were then noted

730   (Fig. S12). Locations of site differences were subsequently intersected with previously identified

731   functional domains available through UniProtKB (Accession: P0DUB6). Further functional

732   analysis followed using data produced from recently published AlphaMissense (J. Cheng et al.

733   2023), examining all possible amino acid substitutions per site across the entire amylase peptide

734   sequence. Using this data, we examined the above noted amylase amino acid site differences and

735   mined previously assigned functional impact scores from AlphaMissense. Mutations from

736   AlphaMissense were described as having no effect ("likely benign"), undetermined effect

737   ("ambiguous"), or having a functional consequence ("likely pathogenic") (table S8).

739   *Detecting sequence origin and breakpoints in AMY1 segments*

740   To explore the origin of sequences within the AMY1 segment, we first generated consensus

741   sequences for each AMY1 segment cluster by selecting the major allele at each site in the

742   alignment. We then prepared a pseudo reference sequence including only a single copy of each

743   amylase segment (AMY1C segment was included as AMY1 segment) from the GRCh38 reference

744   genome. For each 200-bp sliding window with 10 step intervals in each cluster, we searched for

745 the best hit against the pseudo reference sequence using BLAT v. 37x1 (*65*), and determined it as

746 the origin of the 200-bp sequence. The windows where more than or equal to 50% of the sequence

747 is gap were excluded in this analysis. The breakpoints of possible NAHR events were manually

748 determined around the transition of origins by using segment-specific variants (table S15). To

749 visualize homology between the consensus sequences from each AMY1 segment cluster and the

750 other segments in the context of haplotype, we aligned the sequences using nucmer v3.1 and

751 generated dotplots using mummerplot v3.5 (*66*) (fig. S32).

753 ***Dating AMY1 duplication events***

754 To estimate the timing of *AMY1* duplication events, we first constructed a starting tree using the

755 whole AMY1 segment alignment above (see the "Alignments of segments and coding sequences"

756 section) with IQTREE v2.2.0 (*60*) and --alrt 1000 -B 1000 options. We then trimmed out the

757 phylogenetically informative interval (from 22,850 to 26,730 bp in the AMY1 segment alignment,

758 Data File-S2) that is potentially poorly aligned due to the presence of rearrangement breakpoints.

759 Exonerate v2.4.0 (*53*) with maximum intron length of 20-kbp (--maxintron 20000) was used to

760 determine coordinates of the coding region in the alignment and to ensure that all sequences in the

761 alignment have the same coordinates for the coding region. Based on the annotation, we partitioned

762 the alignment into four categories: non-coding region, 1st position of codon, 2nd position of codon,

763 and 3rd position of codon. This partitioned alignment was used as an input for the downstream

764 analyses. We extracted sequences and topology of $H3^r.1$ or $H3^r.2$ haplotypes and chimpanzee

765 genome (mPanTro3) from the partitioned alignment and the starting tree, respectively. We

766 restricted our analyses to $H3^r.1$ or $H3^r.2$ haplotype that is likely to be ancestral (due to their high

767 allele frequency) to other more complex haplotypes to reduce the genetic exchange effect of

768 inversion between different AMY1 segments.

769   For estimating the timing of *AMY1* gene duplication events (that is, AMY1 segment

770 duplication events), BEAST v2.7.5 (*67*), a Bayesian framework, was used with the alignments and

771 starting trees of H3^r.1 and H3^r.2 haplotypes, separately. First, we performed bModel Test

772 implemented in the BEAST to assess whether the four partitions are evolving differently or not

773 using unliked site models and calibrated Yule model. Based on the results of bModel Test, we

774 decided to use average substitution models using bModel Test in the real run to incorporate

775 heterogeneity across partitions. Then, a path sampling approach from BEAST's

776 MODEL_SELECTION package was used for all model combinations to determine best-fit clock

777 and tree models. We tested both strict and optimized relaxed clock models with constant and

778 Bayesian skyline tree models. For the path sampling, 40 path steps with 25,000,000 iterations were

779 used with an alpha parameter of 0.3, pre-burn-in of 75,000 iterations, and an 80% burn-in of the

780 complete chain. While the models with a strict clock model could not produce marginal likelihood

781 estimates, there was no significant difference among the models with an optimized relaxed clock

782 model (bayes factor = 1.32 and 0.14 for H3^r.1 and H3^r.2; table S9). Therefore, we estimated time

783 for both models with Markov chain Monte Carlos (MCMC) with 75,000,000 iterations, sampling

784 every 2,000 trees and using a burn-in of 10% iterations for each model. In all the time estimations

785 above, MRCA prior to assuming the human-chimpanzee divergence time to 6 million years ago

786 under normal distribution (a mean of 6 million years ago and variance of 0.5) was used to calibrate

787 time estimates. Also, we did not allow optimization of the topology of the starting tree during the

788 estimation to avoid shuffling between AMY1 segment clusters likely due to the observed gene

789 conversion effect. Tracer v1.7.2 was used to examine effective sample size and convergence of

790   estimates. Treeannotator v2.7.5 was used to merge and annotate the sampled trees into a single

791   tree. Figtree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/) was used for tree visualization.

793   *Amylase copy number detection in ancient humans*

794   We used the program GeneToCN (*26*) to determine amylase copy number in 73 ancient human

795   and 8 archaic hominin genomes (fig. S33 and, tables S10 and S12) (*27*, *42*, *68–92*). The raw

796   sequencing FASTQ formatted files were obtained through the European Nucleotide Archive

797   (https://www.ebi.ac.uk/ena/browser/). As GeneToCN is a read-depth based copy number caller,

798   we only utilized genomes that had at least 5X coverage both overall and in the amylase region to

799   minimize random error. Additionally, we did not use sequences whose FASTQ files were

800   generated by duplicate-removed bam files as we consistently found that this step generated no

801   output for the *AMY1* copy number analysis. We used the program provided databases

802   (https://github.com/bioinfo-ut/GeneToCN/tree/main/Kmer_db/Gene_kmers/AMY) for *AMY2A*,

803   *AMY2B*, and *AMY1* to calculate the copy number of each gene using 25 nucleotide sequences (25-

804   mers). We additionally used GeneToCN to calculate the amylase copy numbers from 101 present-

805   day humans (table S5). We downloaded mapped bam files for the individuals that are available in

806   both our diploid haplotype data and 1000 Genomes Project short-read sequencing dataset from

807   The International Genome Sample Resource (https://www.internationalgenome.org) and HPRC

808   (https://s3-us-west-2.amazonaws.com/human-pangenomics/index.html?prefix=working/).   We

809   validated these results using digital PCR (R^2 = 0.95 for *AMY1*) (table S5). We used the following

810   command for the GeneToCN:

```
811  python KmerToCN.py -db AMY_all_kmers.db -kp kmer_db_locations.txt
812  -s   ${file}   -d   AMY/${individual}   -o   ${individual}   -gm
813  ${direct}/GenomeTester4/src/ -r AMY.txt -i
```

814   Next, we compared the difference in *AMY1*, *AMY2A*, and *AMY2B* copy numbers between

815 all ancient human genomes practicing hunter-gatherer or agricultural lifestyles using a two-sided

816 Wilcoxon Rank Sum Test (P = 0.037 for *AMY1*) from the `stat_compare_means()` command

817 in the ggplot2 version 3.5.1 library (**fig. S22**). We then binned our time as "**Prior** to the major shift

818 in agriculture", "**During** the shift to agriculture", and "**Post** the shift to agriculture" to compare

819 the amylase copy number over time in both globally and in European populations only (Fig. 3C,

820 3D and fig. S23) using `wilcox.test(alternative = "less")` in R package "stats".

821 We calculated a one tailed Wilcoxon Rank test with the alternative hypothesis that the older time

822 period has a lower amylase copy number for *AMY1*, *AMY2A*, and *AMY2B*. Hungarian samples

823 from the site of Szólád were not included in this analysis as their large sample size in comparison

824 to all other archaeological sites and close kinship creates a bias in the "**Post** transitional" dataset.

826 *Downsampling ancient human genomes to estimate coverage bias*

827 To determine the effect of coverage on the amylase copy number estimation, we randomly

828 downsampled eight ancient human genomes (table S11) using seqtk (v1.4-r122)

829 (https://github.com/lh3/seqtk) for a range from 1X to 5X coverage. We calculated the *AMY1* gene

830 copy number five times and compared these results to the calculated copy number estimated when

831 using the full genome coverage prior to downsampling. To consider batch effects, we chose

832 samples from seven different studies with four different corresponding authors ranging in date

833 from 2014 to 2024 (table S12). Additionally, we chose samples that were a mix of hunter-gatherers

834 (n = 3) and farmers (n = 5) and had a wide range of both average genome coverage (8.2-30X) and

835 *AMY1* gene copy number (6-17 copies) to consider unknown systematic biases coming from

836 different individuals and sequencing settings. An example version of the script for the analysis is

837 provided below:

838 `random_seed=$((RANDOM % 1000 + 1))`

839 `Ratio=$( echo " scale=3; ${Downsample} / ${Coverage}" | bc -l )`

840 `seqtk sample -s seed=${random_Seed} ${Run_ID}_1.fastq.gz`

841 `${Ratio} | gzip >`

842 `${Individual}_1_downsampled_r${Round}_d${Downsample}.fastq.gz`

843 `seqtk sample -s seed=${random_Seed} ${Run_ID}_2.fastq.gz`

844 `${Ratio} | gzip >`

845 `${Individual}_2_downsampled_r${Round}_d${Downsample}.fastq.gz`

846 `coverage="high_coverage"`

847 `sample"${Individual}_downsampled_r${Round}_d${Downsample}"`

848 `GeneToCN.sh ${sample} ${coverage} ${directory}`

850 *Archaic hominin genome analyses*

851 We collected sequencing raw reads of archaic hominin genomes (n = 38) from publicly available

852 datasets (table S10), removed adaptor sequences, and merged paired-end reads in case of

853 overlapping using leeHom v1.2.17 (*93*). To increase sensitivity, we did not trim the reads by their

854 base quality scores. For Chagyrskaya Neanderthal genome, we used unmapped reads

855 (http://ftp.eva.mpg.de/neandertal/Chagyrskaya/rawBAM/) that were already merged, and adapter

856 trimmed. The clean reads were mapped to the GRCh38 reference genome

857 (http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/technical/reference/GRCh38_reference_genome/GRC

858   h38_full_analysis_set_plus_decoy_hla.fa), which includes decoy and HLA sequences, using "bwa

859   aln" v0.7.17 (*94*) with the following options: $-n$ 0.01 $-o$ 2 $-l$ 16500. Next, we employed

860   the "computeGCBias" command from the deepTools v3.5.1 (*95*) suite to assess GC bias in archaic

861   hominin genome dataset. Then, we used the "correctGCBias" command from deepTools v3.5.1 to

862   obtain corrected reads. This correction process is essential for minimizing potential errors due to

863   GC bias in downstream analyses. We evaluated the GC bias in GC-corrected reads one more time

864   and 29 out of 38 archaic hominin genomes were excluded from downstream analyses due to

865   moderate to significant GC bias (table S10). In addition, we calculated average read-depth for the

866   whole genome and the amylase locus using mosdepth v0.3.5 (*96*). Based on this result, we

867   excluded one more sample (Scladina I4-A) that has lower than 1X coverage both at the genome

868   wide level and at the amylase locus.

869   Following the completion of GC bias analysis, the GC-corrected reads were realigned to

870   the T2T-chm13 and GRCh38 assemblies to enhance the accuracy of the alignment and reduce the

871   reference bias (*97*), utilizing "bwa mem" v0.7.17-r1188 (*94*). A reference genome with a single

872   copy of the *AMY1* gene (*10*) was used in the final alignment stage to filter out reads that aligned

873   to non-target chromosomes. These sequential alignment and filtering steps were designed to

874   minimize background noise and enhance the specificity of read alignment to each amylase gene.

875   The average read depths were calculated and normalized by genome-wide depth using mosdepth

876   v0.3.5 (*96*) and copy number of amylase genes were calculated by using this read-depth

877   information. For visualization of read-depth around the *AMY1* gene, we extracted GC-corrected

878   reads mapped to the amylase locus and aligned them back to the reference genome containing a

879   single copy of the *AMY1* gene (Perry et al. 2015). The read-depth was normalized by the average

880   read-depth of *RNPC3* gene for comparable visualization in Fig. 3A.

Scanning PRDM9 binding sites across the H3^r.1 haplotype

To identify potential PRDM9 binding sites, FIMO v5.5.4 implemented in the MEME package (*98*) was employed using the degenerate 13-bp motif, "`CCNCCNTNNCCNC`" (*99*), and the non-redundant DNA database background frequency matrix in the H3^r.1 haplotype. We used a consensus sequence for the H3^r.1 haplotype to incorporate the standing nucleotide variation within the locus. Only hits with a FIMO score above 10 and a p-value below 0.00011 were considered (fig. S26 and table S13).

Investigating mechanisms underlying structural variation formation within haplotypes

To investigate putative mechanisms underlying structural variants between two haplotypes, we aligned the haplotypes using NUCmer v3.1 (*100*). The alignments were visualized as dotplots using mummerplot v3.5 and Miropeats-style plots using the R package SVbyEye (https://github.com/daewoooo/SVbyEye/tree/master). These sequence comparisons allowed us to identify the potential breakpoints of structural variants. Once we identified the potential breakpoints, we extracted 20-kbp long sequences upstream and downstream regions and then aligned them to each other, including the corresponding GRCh38 reference sequence using MAFFT v7.522 (*56*). Based on the analysis of these breakpoints, we evaluated replication and DNA recombination-based processes that underlie the mutational steps leading to the extant structural variants. We inferred NAHR when the variations between two structural haplotypes included (i) breakpoints within paralogous segments and (ii) absence of sequence motifs linked to replication-based mutational mechanisms.

References

906   1. G. H. Perry, N. J. Dominy, K. G. Claw, A. S. Lee, H. Fiegler, R. Redon, J. Werner, F. A.
907   Villanea, J. L. Mountain, R. Misra, N. P. Carter, C. Lee, A. C. Stone, Diet and the evolution
908   of human amylase gene copy number variation. *Nat. Genet.* **39**, 1256–1260 (2007).

909   2. Y. Nakamura, M. Ogawa, T. Nishide, M. Emi, G. Kosaki, S. Himeno, K. Matsubara,
910   Sequences of cDNAs for human salivary and pancreatic α-amylases. *Gene* **28**, 263–270
911   (1984).

912   3. C. Peyrot des Gachons, P. A. S. Breslin, Salivary Amylase: Digestion and Metabolic
913   Syndrome. *Curr. Diab. Rep.* **16**, 102 (2016).

914   4. P. Pajic, P. Pavlidis, K. Dean, L. Neznanova, R.-A. Romano, D. Garneau, E. Daugherity, A.
915   Globig, S. Ruhl, O. Gokcumen, Independent amylase gene copy number bursts correlate
916   with dietary preferences in mammals. *Elife* **8** (2019).

917   5. M. H. Meisler, C. N. Ting, The remarkable evolutionary history of the human amylase
918   genes. *Crit. Rev. Oral Biol. Med.* **4**, 503–509 (1993).

919   6. P. C. Groot, W. H. Mager, R. R. Frants, Interpretation of polymorphic DNA patterns in the
920   human alpha-amylase multigene family. *Genomics* **10**, 779–785 (1991).

921   7. C. L. Usher, R. E. Handsaker, T. Esko, M. A. Tuke, M. N. Weedon, A. R. Hastie, H. Cao, J.
922   E. Moon, S. Kashin, C. Fuchsberger, A. Metspalu, C. N. Pato, M. T. Pato, M. I. McCarthy,
923   M. Boehnke, D. M. Altshuler, T. M. Frayling, J. N. Hirschhorn, S. A. McCarroll, Structural
924   forms of the human amylase locus and their relationships to SNPs, haplotypes and obesity.
925   *Nat. Genet.* **47**, 921–925 (2015).

926   8. A. C. Poole, J. K. Goodrich, N. D. Youngblut, G. G. Luque, A. Ruaud, J. L. Sutter, J. L.
927   Waters, Q. Shi, M. El-Hadidi, L. M. Johnson, H. Y. Bar, D. H. Huson, J. G. Booth, R. E.
928   Ley, Human Salivary Amylase Gene Copy Number Impacts Oral and Gut Microbiomes.
929   *Cell Host Microbe* **25**, 553–564.e7 (2019).

930   9. S. Mathieson, I. Mathieson, FADS1 and the Timing of Human Adaptation to Agriculture.
931   *Mol. Biol. Evol.* **35**, 2957–2970 (2018).

932   10. G. H. Perry, L. Kistler, M. A. Kelaita, A. J. Sams, Insights into hominin phenotypic and
933   dietary evolution from ancient DNA sequence data. *J. Hum. Evol.* **79**, 55–63 (2015).

934   11. D. Bolognini, A. Halgren, R. N. Lou, A. Raveane, J. L. Rocha, A. Guarracino, N. Soranzo,
935   J. Chin, E. Garrison, P. H. Sudmant, Global diversity, recurrent evolution, and recent
936   selection on amylase structural haplotypes in humans. *bioRxiv*, doi:
937   10.1101/2024.02.07.579378 (2024).

938   12. C. E. Inchley, C. D. A. Larbey, N. A. A. Shwan, L. Pagani, L. Saag, T. Antão, G. Jacobs, G.

939   Hudjashov, E. Metspalu, M. Mitt, C. A. Eichstaedt, B. Malyarchuk, M. Derenko, J. Wee, S.
940   Abdullah, F.-X. Ricaut, M. Mormina, R. Mägi, R. Villems, M. Metspalu, M. K. Jones, J. A.
941   L. Armour, T. Kivisild, Selective sweep on human amylase genes postdates the split with
942   Neanderthals. *Sci. Rep.* **6**, 37198 (2016).

943   13. W. Demaerel, Y. Mostovoy, F. Yilmaz, L. Vervoort, S. Pastor, M. S. Hestand, A. Swillen,
944   E. Vergaelen, E. A. Geiger, C. R. Coughlin, S. K. Chow, D. McDonald-McGinn, B.
945   Morrow, P.-Y. Kwok, M. Xiao, B. S. Emanuel, T. H. Shaikh, J. R. Vermeesch, The 22q11
946   low copy repeats are characterized by unprecedented size and structural variability. *Genome*
947   *Res.* **29**, 1389–1401 (2019).

948   14. Y. Mostovoy, F. Yilmaz, S. K. Chow, C. Chu, C. Lin, E. A. Geiger, N. J. L. Meeks, K. C.
949   Chatfield, C. R. Coughlin, U. Surti, P.-Y. Kwok, T. H. Shaikh, Genomic regions associated
950   with microdeletion/microduplication syndromes exhibit extreme diversity of structural
951   variation. [Preprint] (2021). https://doi.org/10.1093/genetics/iyaa038.

952   15. F. Yilmaz, U. Gurusamy, T. J. Mosley, P. Hallast, K. Kim, Y. Mostovoy, R. H. Purcell, T.
953   H. Shaikh, M. E. Zwick, P.-Y. Kwok, C. Lee, J. G. Mulle, High level of complexity and
954   global diversity of the 3q29 locus revealed by optical mapping and long-read sequencing.
955   *Genome Med.* **15**, 35 (2023).

956   16. S. Nurk, S. Koren, A. Rhie, M. Rautiainen, A. V. Bzikadze, A. Mikheenko, M. R. Vollger,
957   N. Altemose, L. Uralsky, A. Gershman, S. Aganezov, S. J. Hoyt, M. Diekhans, G. A.
958   Logsdon, M. Alonge, S. E. Antonarakis, M. Borchers, G. G. Bouffard, S. Y. Brooks, G. V.
959   Caldas, N.-C. Chen, H. Cheng, C.-S. Chin, W. Chow, L. G. de Lima, P. C. Dishuck, R.
960   Durbin, T. Dvorkina, I. T. Fiddes, G. Formenti, R. S. Fulton, A. Fungtammasan, E.
961   Garrison, P. G. S. Grady, T. A. Graves-Lindsay, I. M. Hall, N. F. Hansen, G. A. Hartley, M.
962   Haukness, K. Howe, M. W. Hunkapiller, C. Jain, M. Jain, E. D. Jarvis, P. Kerpedjiev, M.
963   Kirsche, M. Kolmogorov, J. Korlach, M. Kremitzki, H. Li, V. V. Maduro, T. Marschall, A.
964   M. McCartney, J. McDaniel, D. E. Miller, J. C. Mullikin, E. W. Myers, N. D. Olson, B.
965   Paten, P. Peluso, P. A. Pevzner, D. Porubsky, T. Potapova, E. I. Rogaev, J. A. Rosenfeld, S.
966   L. Salzberg, V. A. Schneider, F. J. Sedlazeck, K. Shafin, C. J. Shew, A. Shumate, Y. Sims,
967   A. F. A. Smit, D. C. Soto, I. Sović, J. M. Storer, A. Streets, B. A. Sullivan, F. Thibaud-
968   Nissen, J. Torrance, J. Wagner, B. P. Walenz, A. Wenger, J. M. D. Wood, C. Xiao, S. M.
969   Yan, A. C. Young, S. Zarate, U. Surti, R. C. McCoy, M. Y. Dennis, I. A. Alexandrov, J. L.
970   Gerton, R. J. O'Neill, W. Timp, J. M. Zook, M. C. Schatz, E. E. Eichler, K. H. Miga, A. M.
971   Phillippy, The complete sequence of a human genome, *Science* (2021)p.
972   2021.05.26.445798.

973   17. A. Harpak, A. Bhaskar, J. K. Pritchard, Mutation Rate Variation is a Primary Determinant
974   of the Distribution of Allele Frequencies in Humans. *PLoS Genet.* **12**, e1006489 (2016).

975   18. M. Verbiest, M. Maksimov, Y. Jin, M. Anisimova, M. Gymrek, T. Bilgin Sonay, Mutation
976   and selection processes regulating short tandem repeats give rise to genetic and phenotypic
977   diversity across species. *J. Evol. Biol.* **36**, 321–336 (2023).

978   19. P. Ebert, P. A. Audano, Q. Zhu, B. Rodriguez-Martin, D. Porubsky, M. J. Bonder, A.

979   Sulovari, J. Ebler, W. Zhou, R. Serra Mari, F. Yilmaz, X. Zhao, P. Hsieh, J. Lee, S. Kumar,
980   J. Lin, T. Rausch, Y. Chen, J. Ren, M. Santamarina, W. Höps, H. Ashraf, N. T. Chuang, X.
981   Yang, K. M. Munson, A. P. Lewis, S. Fairley, L. J. Tallon, W. E. Clarke, A. O. Basile, M.
982   Byrska-Bishop, A. Corvelo, U. S. Evani, T.-Y. Lu, M. J. P. Chaisson, J. Chen, C. Li, H.
983   Brand, A. M. Wenger, M. Ghareghani, W. T. Harvey, B. Raeder, P. Hasenfeld, A. A.
984   Regier, H. J. Abel, I. M. Hall, P. Flicek, O. Stegle, M. B. Gerstein, J. M. C. Tubio, Z. Mu,
985   Y. I. Li, X. Shi, A. R. Hastie, K. Ye, Z. Chong, A. D. Sanders, M. C. Zody, M. E.
986   Talkowski, R. E. Mills, S. E. Devine, C. Lee, J. O. Korbel, T. Marschall, E. E. Eichler,
987   Haplotype-resolved diverse human genomes and integrated analysis of structural variation.
988   *Science* **372** (2021).

989   20.   Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**,
990   1586–1591 (2007).

991   21.   J. Cheng, G. Novati, J. Pan, C. Bycroft, A. Žemgulytė, T. Applebaum, A. Pritzel, L. H.
992   Wong, M. Zielinski, T. Sargeant, R. G. Schneider, A. W. Senior, J. Jumper, D. Hassabis, P.
993   Kohli, Ž. Avsec, Accurate proteome-wide missense variant effect prediction with
994   AlphaMissense. *Science* **381**, eadg7492 (2023).

995   22.   D. Carpenter, S. Dhar, L. M. Mitchell, B. Fu, J. Tyson, N. A. A. Shwan, F. Yang, M. G.
996   Thomas, J. A. L. Armour, Obesity, starch digestion and amylase: association between copy
997   number variants at human salivary (AMY1) and pancreatic (AMY2) amylase genes. *Hum.
998   Mol. Genet.* **24**, 3472–3480 (2015).

999   23.   M. R. Vollger, P. C. Dishuck, W. T. Harvey, W. S. DeWitt, X. Guitart, M. E. Goldberg, A.
1000   N. Rozanski, J. Lucas, M. Asri, Human Pangenome Reference Consortium, K. M. Munson,
1001   A. P. Lewis, K. Hoekzema, G. A. Logsdon, D. Porubsky, B. Paten, K. Harris, P. Hsieh, E.
1002   E. Eichler, Increased mutation and gene conversion within human segmental duplications.
1003   *Nature* **617**, 325–334 (2023).

1004   24.   M. Lipson, A. Szécsényi-Nagy, S. Mallick, A. Pósa, B. Stégmár, V. Keerl, N. Rohland, K.
1005   Stewardson, M. Ferry, M. Michel, J. Oppenheimer, N. Broomandkhoshbacht, E. Harney, S.
1006   Nordenfelt, B. Llamas, B. Gusztáv Mende, K. Köhler, K. Oross, M. Bondár, T. Marton, A.
1007   Osztás, J. Jakucs, T. Paluch, F. Horváth, P. Csengeri, J. Koós, K. Sebők, A. Anders, P.
1008   Raczky, J. Regenye, J. P. Barna, S. Fábián, G. Serlegi, Z. Toldi, E. Gyöngyvér Nagy, J.
1009   Dani, E. Molnár, G. Pálfi, L. Márk, B. Melegh, Z. Bánfai, L. Domboróczki, J. Fernández-
1010   Eraso, J. Antonio Mujika-Alustiza, C. Alonso Fernández, J. Jiménez Echevarría, R.
1011   Bollongino, J. Orschiedt, K. Schierhold, H. Meller, A. Cooper, J. Burger, E. Bánffy, K. W.
1012   Alt, C. Lalueza-Fox, W. Haak, D. Reich, Parallel palaeogenomic transects reveal complex
1013   genetic history of early European farmers. *Nature* **551**, 368–372 (2017).

1014   25.   N. E. Altınışık, D. D. Kazancı, A. Aydoğan, H. C. Gemici, Ö. D. Erdal, S. Sarıaltun, K. B.
1015   Vural, D. Koptekin, K. Gürün, E. Sağlıcan, D. Fernandes, G. Çakan, M. M. Koruyucu, V.
1016   K. Lagerholm, C. Karamurat, M. Özkan, G. M. Kılınç, A. Sevkar, E. Sürer, A. Götherström,
1017   Ç. Atakuman, Y. S. Erdal, F. Özer, A. Erim Özdoğan, M. Somel, A genomic snapshot of
1018   demographic and cultural dynamism in Upper Mesopotamia during the Neolithic
1019   Transition. *Sci Adv* **8**, eabo3609 (2022).

1020   26.   F.-D. Pajuste, M. Remm, GeneToCN: an alignment-free method for gene copy number
1021   estimation directly from next-generation sequencing reads. *Sci. Rep.* **13**, 17765 (2023).

1022   27.   K. Prüfer, F. Racimo, N. Patterson, F. Jay, S. Sankararaman, S. Sawyer, A. Heinze, G.
1023   Renaud, P. H. Sudmant, C. de Filippo, H. Li, S. Mallick, M. Dannemann, Q. Fu, M.
1024   Kircher, M. Kuhlwilm, M. Lachmann, M. Meyer, M. Ongyerth, M. Siebauer, C. Theunert,
1025   A. Tandon, P. Moorjani, J. Pickrell, J. C. Mullikin, S. H. Vohr, R. E. Green, I. Hellmann, P.
1026   L. F. Johnson, H. Blanche, H. Cann, J. O. Kitzman, J. Shendure, E. E. Eichler, E. S. Lein, T.
1027   E. Bakken, L. V. Golovanova, V. B. Doronichev, M. V. Shunkov, A. P. Derevianko, B.
1028   Viola, M. Slatkin, D. Reich, J. Kelso, S. Pääbo, The complete genome sequence of a
1029   Neanderthal from the Altai Mountains. *Nature* **505**, 43–49 (2014).

1030   28.   D. N. Harris, A. Platt, M. E. B. Hansen, S. Fan, M. A. McQuillan, T. Nyambo, S. W.
1031   Mpoloka, G. G. Mokone, G. Belay, C. Fokunang, A. K. Njamnshi, S. A. Tishkoff, Diverse
1032   African genomes reveal selection on ancient modern human introgressions in Neanderthals.
1033   *Curr. Biol.* **33**, 4905–4916.e5 (2023).

1034   29.   M. Hajdinjak, Q. Fu, A. Hübner, M. Petr, F. Mafessoni, S. Grote, P. Skoglund, V.
1035   Narasimham, H. Rougier, I. Crevecoeur, P. Semal, M. Soressi, S. Talamo, J.-J. Hublin, I.
1036   Gušić, Ž. Kućan, P. Rudan, L. V. Golovanova, V. B. Doronichev, C. Posth, J. Krause, P.
1037   Korlević, S. Nagel, B. Nickel, M. Slatkin, N. Patterson, D. Reich, K. Prüfer, M. Meyer, S.
1038   Pääbo, J. Kelso, Reconstructing the genetic history of late Neanderthals. *Nature* **555**, 652–
1039   656 (2018).

1040   30.   A. Gómez-Robles, Dental evolutionary rates and its implications for the Neanderthal-
1041   modern human divergence. *Sci Adv* **5**, eaaw1268 (2019).

1042   31.   L. Vallini, C. Zampieri, M. J. Shoaee, E. Bortolini, G. Marciani, S. Aneli, T. Pievani, S.
1043   Benazzi, A. Barausse, M. Mezzavilla, M. D. Petraglia, L. Pagani, The Persian plateau
1044   served as hub for Homo sapiens after the main out of Africa dispersal. *Nat. Commun.* **15**,
1045   1882 (2024).

1046   32.   A. Revedin, B. Aranguren, R. Becattini, L. Longo, E. Marconi, M. M. Lippi, N. Skakun, A.
1047   Sinitsyn, E. Spiridonova, J. Svoboda, Thirty thousand-year-old evidence of plant food
1048   processing. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 18815–18819 (2010).

1049   33.   D. R. Piperno, E. Weiss, I. Holst, D. Nadel, Processing of wild cereal grains in the Upper
1050   Palaeolithic revealed by starch grain analysis. *Nature* **430**, 670–673 (2004).

1051   34.   J. Jovanović, R. C. Power, C. de Becdelièvre, G. Goude, S. Stefanović, Microbotanical
1052   evidence for the spread of cereal use during the Mesolithic-Neolithic transition in the
1053   Southeastern Europe (Danube Gorges): Data from dental calculus analysis. *J. Archaeol. Sci.*
1054   **125**, 105288 (2021).

1055   35.   P. C. Groot, W. H. Mager, N. V. Henriquez, J. C. Pronk, F. Arwert, R. J. Planta, A. W.
1056   Eriksson, R. R. Frants, Evolution of the human alpha-amylase multigene family through
1057   unequal, homologous, and inter- and intrachromosomal crossovers. *Genomics* **8**, 97–105
1058   (1990).

36. F. Zhang, M. Khajavi, A. M. Connolly, C. F. Towne, S. D. Batish, J. R. Lupski, The DNA replication FoSTeS/MMBIR mechanism can generate genomic, genic and exonic complex rearrangements in humans. *Nat. Genet.* **41**, 849–853 (2009).

37. D. Porubsky, W. Höps, H. Ashraf, P. Hsieh, B. Rodriguez-Martin, F. Yilmaz, J. Ebler, P. Hallast, F. A. Maria Maggiolini, W. T. Harvey, B. Henning, P. A. Audano, D. S. Gordon, P. Ebert, P. Hasenfeld, E. Benito, Q. Zhu, Human Genome Structural Variation Consortium (HGSVC), C. Lee, F. Antonacci, M. Steinrücken, C. R. Beck, A. D. Sanders, T. Marschall, E. E. Eichler, J. O. Korbel, Recurrent inversion polymorphisms in humans associate with genetic instability and genomic disorders. *Cell* **185**, 1986–2005.e26 (2022).

38. J. A. F. Yates, I. M. Velsko, F. Aron, C. Posth, C. A. Hofman, R. M. Austin, C. E. Parker, A. E. Mann, K. Nägele, K. W. Arthur, J. W. Arthur, C. C. Bauer, I. Crevecoeur, C. Cupillard, M. C. Curtis, L. Dalén, M. D.-Z. Bonilla, J. C. D. Fernández-Lomana, D. G. Drucker, E. E. Escrivá, M. Francken, V. E. Gibbon, M. R. G. Morales, A. G. Mateu, K. Harvati, A. G. Henry, L. Humphrey, M. Menéndez, D. Mihailović, M. Peresani, S. R. Moroder, M. Roksandic, H. Rougier, S. Sázelová, J. T. Stock, L. G. Straus, J. Svoboda, B. Teßmann, M. J. Walker, R. C. Power, C. M. Lewis, K. Sankaranarayanan, K. Guschanski, R. W. Wrangham, F. E. Dewhirst, D. C. Salazar-García, J. Krause, A. Herbig, C. Warinner, The evolution and changing ecology of the African hominid oral microbiome. *Proceedings of the National Academy of Sciences* **118**, e2021655118 (2021).

39. C. Larbey, S. M. Mentzer, B. Ligouis, S. Wurz, M. K. Jones, Cooked starchy food in hearths ca. 120 kya and 65 kya (MIS 5e and MIS 4) from Klasies River Cave, South Africa. *J. Hum. Evol.* **131**, 210–227 (2019).

40. E. Cristiani, A. Radini, A. Zupancich, A. Gismondi, A. D'Agostino, C. Ottoni, M. Carra, S. Vukojičić, M. Constantinescu, D. Antonović, T. D. Price, D. Borić, Wild cereal grain consumption among Early Holocene foragers of the Balkans predates the arrival of agriculture. *Elife* **10** (2021).

41. G. M. Kılınç, A. Omrak, F. Özer, T. Günther, A. M. Büyükkarakaya, E. Bıçakçı, D. Baird, H. M. Dönertaş, A. Ghalichi, R. Yaka, D. Koptekin, S. C. Açan, P. Parvizi, M. Krzewińska, E. A. Daskalaki, E. Yüncü, N. D. Dağtaş, A. Fairbairn, J. Pearson, G. Mustafaoğlu, Y. S. Erdal, Y. G. Çakan, İ. Togan, M. Somel, J. Storå, M. Jakobsson, A. Götherström, The Demographic Development of the First Farmers in Anatolia. *Curr. Biol.* **26**, 2659–2666 (2016).

42. N. Marchi, L. Winkelbach, I. Schulz, M. Brami, Z. Hofmanová, J. Blöcher, C. S. Reyna-Blanco, Y. Diekmann, A. Thiéry, A. Kapopoulou, V. Link, V. Piuz, S. Kreutzer, S. M. Figarska, E. Ganiatsou, A. Pukaj, T. J. Struck, R. N. Gutenkunst, N. Karul, F. Gerritsen, J. Pechtl, J. Peters, A. Zeeb-Lanz, E. Lenneis, M. Teschler-Nicola, S. Triantaphyllou, S. Stefanović, C. Papageorgopoulou, D. Wegmann, J. Burger, L. Excoffier, The genomic origins of the world's first farmers. *Cell* **185**, 1842–1859.e18 (2022).

43. J. Lindo, R. Haas, C. Hofman, M. Apata, M. Moraga, R. A. Verdugo, J. T. Watson, C. Viviano Llave, D. Witonsky, C. Beall, C. Warinner, J. Novembre, M. Aldenderfer, A. Di

1099   Rienzo, The genetic prehistory of the Andean highlands 7000 years BP though European
1100   contact. *Sci Adv* **4**, eaau4921 (2018).

1101 44. K. Jorgensen, O. A. Garcia, M. Kiyamu, T. D. Brutsaert, A. W. Bigham, Genetic
1102   adaptations to potato starch digestion in the Peruvian Andes. *Am. J. Biol. Anthropol.* **180**,
1103   162–172 (2023).

1104 45. J. V. Moreno-Mayar, B. A. Potter, L. Vinner, M. Steinrücken, S. Rasmussen, J. Terhorst, J.
1105   A. Kamm, A. Albrechtsen, A.-S. Malaspinas, M. Sikora, J. D. Reuther, J. D. Irish, R. S.
1106   Malhi, L. Orlando, Y. S. Song, R. Nielsen, D. J. Meltzer, E. Willerslev, Terminal
1107   Pleistocene Alaskan genome reveals first founding population of Native Americans. *Nature*
1108   **553**, 203–207 (2018).

1109 46. W.-W. Liao, M. Asri, J. Ebler, D. Doerr, M. Haukness, G. Hickey, S. Lu, J. K. Lucas, J.
1110   Monlong, H. J. Abel, S. Buonaiuto, X. H. Chang, H. Cheng, J. Chu, V. Colonna, J. M.
1111   Eizenga, X. Feng, C. Fischer, R. S. Fulton, S. Garg, C. Groza, A. Guarracino, W. T.
1112   Harvey, S. Heumos, K. Howe, M. Jain, T.-Y. Lu, C. Markello, F. J. Martin, M. W. Mitchell,
1113   K. M. Munson, M. N. Mwaniki, A. M. Novak, H. E. Olsen, T. Pesout, D. Porubsky, P.
1114   Prins, J. A. Sibbesen, J. Sirén, C. Tomlinson, F. Villani, M. R. Vollger, L. L. Antonacci-
1115   Fulton, G. Baid, C. A. Baker, A. Belyaeva, K. Billis, A. Carroll, P.-C. Chang, S. Cody, D.
1116   E. Cook, R. M. Cook-Deegan, O. E. Cornejo, M. Diekhans, P. Ebert, S. Fairley, O. Fedrigo,
1117   A. L. Felsenfeld, G. Formenti, A. Frankish, Y. Gao, N. A. Garrison, C. G. Giron, R. E.
1118   Green, L. Haggerty, K. Hoekzema, T. Hourlier, H. P. Ji, E. E. Kenny, B. A. Koenig, A.
1119   Kolesnikov, J. O. Korbel, J. Kordosky, S. Koren, H. Lee, A. P. Lewis, H. Magalhães, S.
1120   Marco-Sola, P. Marijon, A. McCartney, J. McDaniel, J. Mountcastle, M. Nattestad, S. Nurk,
1121   N. D. Olson, A. B. Popejoy, D. Puiu, M. Rautiainen, A. A. Regier, A. Rhie, S. Sacco, A. D.
1122   Sanders, V. A. Schneider, B. I. Schultz, K. Shafin, M. W. Smith, H. J. Sofia, A. N. Abou
1123   Tayoun, F. Thibaud-Nissen, F. F. Tricomi, J. Wagner, B. Walenz, J. M. D. Wood, A. V.
1124   Zimin, G. Bourque, M. J. P. Chaisson, P. Flicek, A. M. Phillippy, J. M. Zook, E. E. Eichler,
1125   D. Haussler, T. Wang, E. D. Jarvis, K. H. Miga, E. Garrison, T. Marschall, I. M. Hall, H. Li,
1126   B. Paten, A draft human pangenome reference. *Nature* **617**, 312–324 (2023).

1127 47. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment
1128   search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

1129 48. H. Cao, A. R. Hastie, D. Cao, E. T. Lam, Y. Sun, H. Huang, X. Liu, L. Lin, W. Andrews, S.
1130   Chan, S. Huang, X. Tong, M. Requa, T. Anantharaman, A. Krogh, H. Yang, H. Cao, X. Xu,
1131   Rapid detection of structural variation in a human genome using nanochannel-based
1132   genome mapping technology. *Gigascience* **3**, 34 (2014).

1133 49. H. Cheng, G. T. Concepcion, X. Feng, H. Zhang, H. Li, Haplotype-resolved de novo
1134   assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175 (2021).

1135 50. Y. Shimoyama, pyGenomeViz: A genome visualization python package for comparative
1136   genomics. Jun [Preprint] (2022).

1137 51. C.-S. Chin, S. Behera, A. Khalak, F. J. Sedlazeck, P. H. Sudmant, J. Wagner, J. M. Zook,
1138   Multiscale analysis of pangenomes enables improved representation of genomic diversity

1139   for repetitive and clinically relevant genes. *Nat. Methods* **20**, 1213–1221 (2023).

1140 52. K. H. M. A. A. C. T. C. Hsieh, iNEXT: an R package for rarefaction and extrapolation of
1141   species diversity (Hill numbers). *Methods in Ecology and Evolution* **7**, 1451–1456 (2016).

1142 53. G. S. C. Slater, E. Birney, Automated generation of heuristics for biological sequence
1143   comparison. *BMC Bioinformatics* **6**, 31 (2005).

1144 54. The UniProt Consortium, UniProt: a worldwide hub of protein knowledge. *Nucleic Acids*
1145   *Res.* **47**, D506–D515 (2018).

1146 55. A. Shumate, S. L. Salzberg, Liftoff: accurate mapping of gene annotations. *Bioinformatics*,
1147   doi: 10.1093/bioinformatics/btaa1016 (2020).

1148 56. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:
1149   improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

1150 57. A. Löytynoja, A. J. Vilella, N. Goldman, Accurate extension of multiple sequence
1151   alignments using a phylogeny-aware graph algorithm. *Bioinformatics* **28**, 1684–1691
1152   (2012).

1153 58. D. V. Hinkley, Inference about the change-point in a sequence of random variables.
1154   *Biometrika* **57**, 1–17 (1970).

1155 59. F. J. Martin, M. R. Amode, A. Aneja, O. Austine-Orimoloye, A. G. Azov, I. Barnes, A.
1156   Becker, R. Bennett, A. Berry, J. Bhai, S. K. Bhurji, A. Bignell, S. Boddu, P. R. Branco Lins,
1157   L. Brooks, S. B. Ramaraju, M. Charkhchi, A. Cockburn, L. Da Rin Fiorretto, C. Davidson,
1158   K. Dodiya, S. Donaldson, B. El Houdaigui, T. El Naboulsi, R. Fatima, C. G. Giron, T.
1159   Genez, G. S. Ghattaoraya, J. G. Martinez, C. Guijarro, M. Hardy, Z. Hollis, T. Hourlier, T.
1160   Hunt, M. Kay, V. Kaykala, T. Le, D. Lemos, D. Marques-Coelho, J. C. Marugán, G. A.
1161   Merino, L. P. Mirabueno, A. Mushtaq, S. N. Hossain, D. N. Ogeh, M. P. Sakthivel, A.
1162   Parker, M. Perry, I. Piližota, I. Prosovetskaia, J. G. Pérez-Silva, A. I. A. Salam, N. Saraiva-
1163   Agostinho, H. Schuilenburg, D. Sheppard, S. Sinha, B. Sipos, W. Stark, E. Steed, R.
1164   Sukumaran, D. Sumathipala, M.-M. Suner, L. Surapaneni, K. Sutinen, M. Szpak, F. F.
1165   Tricomi, D. Urbina-Gómez, A. Veidenberg, T. A. Walsh, B. Walts, E. Wass, N. Willhoft, J.
1166   Allen, J. Alvarez-Jarreta, M. Chakiachvili, B. Flint, S. Giorgetti, L. Haggerty, G. R. Ilsley,
1167   J. E. Loveland, B. Moore, J. M. Mudge, J. Tate, D. Thybert, S. J. Trevanion, A.
1168   Winterbottom, A. Frankish, S. E. Hunt, M. Ruffier, F. Cunningham, S. Dyer, R. D. Finn, K.
1169   L. Howe, P. W. Harrison, A. D. Yates, P. Flicek, Ensembl 2023. *Nucleic Acids Res.* **51**,
1170   D933–D941 (2023).

1171 60. B. Q. Minh, H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von
1172   Haeseler, R. Lanfear, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
1173   Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

1174 61. S. Kalyaanamoorthy, B. Q. Minh, T. K. F. Wong, A. von Haeseler, L. S. Jermiin,
1175   ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,
1176   587–589 (2017).

1177   62.   D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2: Improving
1178   the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

1179   63.   Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: A practical and powerful
1180   approach to multiple testing. *J. R. Stat. Soc.* **57**, 289–300 (1995).

1181   64.   F. Sievers, D. G. Higgins, Clustal Omega for making accurate alignments of many protein
1182   sequences. *Protein Sci.* **27**, 135–145 (2018).

1183   65.   W. J. Kent, BLAT--the BLAST-like alignment tool. *Genome Res.* **12**, 656–664 (2002).

1184   66.   G. Marçais, A. L. Delcher, A. M. Phillippy, R. Coston, S. L. Salzberg, A. Zimin,
1185   MUMmer4: A fast and versatile genome alignment system. *PLoS Comput. Biol.* **14**,
1186   e1005944 (2018).

1187   67.   R. Bouckaert, J. Heled, D. Kühnert, T. Vaughan, C.-H. Wu, D. Xie, M. A. Suchard, A.
1188   Rambaut, A. J. Drummond, BEAST 2: a software platform for Bayesian evolutionary
1189   analysis. *PLoS Comput. Biol.* **10**, e1003537 (2014).

1190   68.   M. Rasmussen, Y. Li, S. Lindgreen, J. S. Pedersen, A. Albrechtsen, I. Moltke, M. Metspalu,
1191   E. Metspalu, T. Kivisild, R. Gupta, M. Bertalan, K. Nielsen, M. T. P. Gilbert, Y. Wang, M.
1192   Raghavan, P. F. Campos, H. M. Kamp, A. S. Wilson, A. Gledhill, S. Tridico, M. Bunce, E.
1193   D. Lorenzen, J. Binladen, X. Guo, J. Zhao, X. Zhang, H. Zhang, Z. Li, M. Chen, L.
1194   Orlando, K. Kristiansen, M. Bak, N. Tommerup, C. Bendixen, T. L. Pierre, B. Grønnow, M.
1195   Meldgaard, C. Andreasen, S. A. Fedorova, L. P. Osipova, T. F. G. Higham, C. B. Ramsey,
1196   T. V. O. Hansen, F. C. Nielsen, M. H. Crawford, S. Brunak, T. Sicheritz-Pontén, R.
1197   Villems, R. Nielsen, A. Krogh, J. Wang, E. Willerslev, Ancient human genome sequence of
1198   an extinct Palaeo-Eskimo. *Nature* **463**, 757–762 (2010).

1199   69.   D. Reich, R. E. Green, M. Kircher, J. Krause, N. Patterson, E. Y. Durand, B. Viola, A. W.
1200   Briggs, U. Stenzel, P. L. F. Johnson, T. Maricic, J. M. Good, T. Marques-Bonet, C. Alkan,
1201   Q. Fu, S. Mallick, H. Li, M. Meyer, E. E. Eichler, M. Stoneking, M. Richards, S. Talamo,
1202   M. V. Shunkov, A. P. Derevianko, J.-J. Hublin, J. Kelso, M. Slatkin, S. Pääbo, Genetic
1203   history of an archaic hominin group from Denisova Cave in Siberia. *Nature* **468**, 1053–
1204   1060 (2010).

1205   70.   Q. Fu, H. Li, P. Moorjani, F. Jay, S. M. Slepchenko, A. A. Bondarev, P. L. F. Johnson, A.
1206   Aximu-Petri, K. Prüfer, C. de Filippo, M. Meyer, N. Zwyns, D. C. Salazar-García, Y. V.
1207   Kuzmin, S. G. Keates, P. A. Kosintsev, D. I. Razhev, M. P. Richards, N. V. Peristov, M.
1208   Lachmann, K. Douka, T. F. G. Higham, M. Slatkin, J.-J. Hublin, D. Reich, J. Kelso, T. B.
1209   Viola, S. Pääbo, Genome sequence of a 45,000-year-old modern human from western
1210   Siberia. *Nature* **514**, 445–449 (2014).

1211   71.   C. Gamba, E. R. Jones, M. D. Teasdale, R. L. McLaughlin, G. Gonzalez-Fortes, V.
1212   Mattiangeli, L. Domboróczki, I. Kővári, I. Pap, A. Anders, A. Whittle, J. Dani, P. Raczky,
1213   T. F. G. Higham, M. Hofreiter, D. G. Bradley, R. Pinhasi, Genome flux and stasis in a five
1214   millennium transect of European prehistory. *Nat. Commun.* **5**, 5257 (2014).

1215 72. I. Lazaridis, N. Patterson, A. Mittnik, G. Renaud, S. Mallick, K. Kirsanow, P. H. Sudmant,
1216   J. G. Schraiber, S. Castellano, M. Lipson, B. Berger, C. Economou, R. Bollongino, Q. Fu,
1217   K. I. Bos, S. Nordenfelt, H. Li, C. de Filippo, K. Prüfer, S. Sawyer, C. Posth, W. Haak, F.
1218   Hallgren, E. Fornander, N. Rohland, D. Delsate, M. Francken, J.-M. Guinet, J. Wahl, G.
1219   Ayodo, H. A. Babiker, G. Bailliet, E. Balanovska, O. Balanovsky, R. Barrantes, G. Bedoya,
1220   H. Ben-Ami, J. Bene, F. Berrada, C. M. Bravi, F. Brisighelli, G. B. J. Busby, F. Cali, M.
1221   Churnosov, D. E. C. Cole, D. Corach, L. Damba, G. van Driem, S. Dryomov, J.-M.
1222   Dugoujon, S. A. Fedorova, I. Gallego Romero, M. Gubina, M. Hammer, B. M. Henn, T.
1223   Hervig, U. Hodoglugil, A. R. Jha, S. Karachanak-Yankova, R. Khusainova, E.
1224   Khusnutdinova, R. Kittles, T. Kivisild, W. Klitz, V. Kučinskas, A. Kushniarevich, L.
1225   Laredj, S. Litvinov, T. Loukidis, R. W. Mahley, B. Melegh, E. Metspalu, J. Molina, J.
1226   Mountain, K. Näkkäläjärvi, D. Nesheva, T. Nyambo, L. Osipova, J. Parik, F. Platonov, O.
1227   Posukh, V. Romano, F. Rothhammer, I. Rudan, R. Ruizbakiev, H. Sahakyan, A. Sajantila,
1228   A. Salas, E. B. Starikovskaya, A. Tarekegn, D. Toncheva, S. Turdikulova, I. Uktveryte, O.
1229   Utevska, R. Vasquez, M. Villena, M. Voevoda, C. A. Winkler, L. Yepiskoposyan, P.
1230   Zalloua, T. Zemunik, A. Cooper, C. Capelli, M. G. Thomas, A. Ruiz-Linares, S. A.
1231   Tishkoff, L. Singh, K. Thangaraj, R. Villems, D. Comas, R. Sukernik, M. Metspalu, M.
1232   Meyer, E. E. Eichler, J. Burger, M. Slatkin, S. Pääbo, J. Kelso, D. Reich, J. Krause, Ancient
1233   human genomes suggest three ancestral populations for present-day Europeans. *Nature* **513**,
1234   409–413 (2014).

1235 73. L. M. Cassidy, R. Martiniano, E. M. Murphy, M. D. Teasdale, J. Mallory, B. Hartwell, D.
1236   G. Bradley, Neolithic and Bronze Age migration to Ireland and establishment of the insular
1237   Atlantic genome. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 368–373 (2016).

1238 74. E. R. Jones, G. Gonzalez-Fortes, S. Connell, V. Siska, A. Eriksson, R. Martiniano, R. L.
1239   McLaughlin, M. Gallego Llorente, L. M. Cassidy, C. Gamba, T. Meshveliani, O. Bar-
1240   Yosef, W. Müller, A. Belfer-Cohen, Z. Matskevich, N. Jakeli, T. F. G. Higham, M. Currat,
1241   D. Lordkipanidze, M. Hofreiter, A. Manica, R. Pinhasi, D. G. Bradley, Upper Palaeolithic
1242   genomes reveal deep roots of modern Eurasians. *Nat. Commun.* **6**, 8912 (2015).

1243 75. M. G. Llorente, E. R. Jones, A. Eriksson, V. Siska, K. W. Arthur, J. W. Arthur, M. C.
1244   Curtis, J. T. Stock, M. Coltorti, P. Pieruccini, S. Stretton, F. Brock, T. Higham, Y. Park, M.
1245   Hofreiter, D. G. Bradley, J. Bhak, R. Pinhasi, A. Manica, Ancient Ethiopian genome reveals
1246   extensive Eurasian admixture in Eastern Africa. *Science* **350**, 820–822 (2015).

1247 76. C. Jeong, A. T. Ozga, D. B. Witonsky, H. Malmström, H. Edlund, C. A. Hofman, R. W.
1248   Hagan, M. Jakobsson, C. M. Lewis, M. S. Aldenderfer, A. Di Rienzo, C. Warinner, Long-
1249   term genetic stability and a high-altitude East Asian origin for the peoples of the high
1250   valleys of the Himalayan arc. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 7485–7490 (2016).

1251 77. S. Schiffels, W. Haak, P. Paajanen, B. Llamas, E. Popescu, L. Loe, R. Clarke, A. Lyons, R.
1252   Mortimer, D. Sayer, C. Tyler-Smith, A. Cooper, R. Durbin, Iron Age and Anglo-Saxon
1253   genomes from East England reveal British migration history. *Nat. Commun.* **7**, 10408
1254   (2016).

1255 78. C. M. Schlebusch, H. Malmström, T. Günther, P. Sjödin, A. Coutinho, H. Edlund, A. R.

1256   Munters, M. Vicente, M. Steyn, H. Soodyall, M. Lombard, M. Jakobsson, Southern African
1257   ancient genomes estimate modern human divergence to 350,000 to 260,000 years ago.
1258   *Science* **358**, 652–655 (2017).

1259   79.   C. E. G. Amorim, S. Vai, C. Posth, A. Modi, I. Koncz, S. Hakenbeck, M. C. La Rocca, B.
1260   Mende, D. Bobo, W. Pohl, L. P. Baricco, E. Bedini, P. Francalacci, C. Giostra, T. Vida, D.
1261   Winger, U. von Freeden, S. Ghirotto, M. Lari, G. Barbujani, J. Krause, D. Caramelli, P. J.
1262   Geary, K. R. Veeramah, Understanding 6th-century barbarian social organization and
1263   migration through paleogenomics. *Nat. Commun.* **9**, 3547 (2018).

1264   80.   C. de la Fuente, M. C. Ávila-Arcos, J. Galimany, M. L. Carpenter, J. R. Homburger, A.
1265   Blanco, P. Contreras, D. C. Dávalos, O. Reyes, M. S. Roman, A. Moreno-Estrada, P. F.
1266   Campos, C. Eng, S. Huntsman, E. G. Burchard, A.-S. Malaspinas, C. D. Bustamante, E.
1267   Willerslev, E. Llop, R. A. Verdugo, M. Moraga, Genomic insights into the origin and
1268   diversification of late maritime hunter-gatherers from the Chilean Patagonia. *Proceedings of*
1269   *the National Academy of Sciences* **115**, E4006–E4012 (2018).

1270   81.   S. S. Ebenesersdóttir, M. Sandoval-Velasco, E. D. Gunnarsdóttir, A. Jagadeesan, V. B.
1271   Guðmundsdóttir, E. L. Thordardóttir, M. S. Einarsdóttir, K. H. S. Moore, Á. Sigurðsson, D.
1272   N. Magnúsdóttir, H. Jónsson, S. Snorradóttir, E. Hovig, P. Møller, I. Kockum, T. Olsson, L.
1273   Alfredsson, T. F. Hansen, T. Werge, G. L. Cavalleri, E. Gilbert, C. Lalueza-Fox, J. W.
1274   Walser 3rd, S. Kristjánsdóttir, S. Gopalakrishnan, L. Árnadóttir, Ó. Þ. Magnússon, M. T. P.
1275   Gilbert, K. Stefánsson, A. Helgason, Ancient genomes from Iceland reveal the making of a
1276   human population. *Science* **360**, 1028–1032 (2018).

1277   82.   F. Sánchez-Quinto, H. Malmström, M. Fraser, L. Girdland-Flink, E. M. Svensson, L. G.
1278   Simões, R. George, N. Hollfelder, G. Burenhult, G. Noble, K. Britton, S. Talamo, N. Curtis,
1279   H. Brzobohata, R. Sumberova, A. Götherström, J. Storå, M. Jakobsson, Megalithic tombs in
1280   western and northern Neolithic Europe were linked to a kindred society. *Proc. Natl. Acad.*
1281   *Sci. U. S. A.* **116**, 9469–9474 (2019).

1282   83.   L. M. Cassidy, R. Ó. Maoldúin, T. Kador, A. Lynch, C. Jones, P. C. Woodman, E. Murphy,
1283   G. Ramsey, M. Dowd, A. Noonan, C. Campbell, E. R. Jones, V. Mattiangeli, D. G. Bradley,
1284   A dynastic elite in monumental Neolithic society. *Nature* **582**, 384–388 (2020).

1285   84.   C. Ning, T. Li, K. Wang, F. Zhang, T. Li, X. Wu, S. Gao, Q. Zhang, H. Zhang, M. J.
1286   Hudson, G. Dong, S. Wu, Y. Fang, C. Liu, C. Feng, W. Li, T. Han, R. Li, J. Wei, Y. Zhu,
1287   Y. Zhou, C.-C. Wang, S. Fan, Z. Xiong, Z. Sun, M. Ye, L. Sun, X. Wu, F. Liang, Y. Cao,
1288   X. Wei, H. Zhu, H. Zhou, J. Krause, M. Robbeets, C. Jeong, Y. Cui, Ancient genomes from
1289   northern China suggest links between subsistence changes and human migration. *Nat.*
1290   *Commun.* **11**, 2700 (2020).

1291   85.   P. Maisano Delser, E. R. Jones, A. Hovhannisyan, L. Cassidy, R. Pinhasi, A. Manica, A
1292   curated dataset of modern and ancient high-coverage shotgun human genomes. *Sci Data* **8**,
1293   202 (2021).

1294   86.   A. Seguin-Orlando, R. Donat, C. Der Sarkissian, J. Southon, C. Thèves, C. Manen, Y.
1295   Tchérémissinoff, E. Crubézy, B. Shapiro, J.-F. Deleuze, L. Dalén, J. Guilaine, L. Orlando,

Heterogeneous Hunter-Gatherer and Steppe-Related Ancestries in Late Neolithic and Bell Beaker Genomes from Present-Day France. *Curr. Biol.* **31**, 1072–1083.e10 (2021).

87. E. Svensson, T. Günther, A. Hoischen, M. Hervella, A. R. Munters, M. Ioana, F. Ridiche, H. Edlund, R. C. van Deuren, A. Soficaru, C. de-la-Rua, M. G. Netea, M. Jakobsson, Genome of Peştera Muierii skull shows high diversity and low mutational load in pre-glacial Europe. *Curr. Biol.* **31**, 2973–2983.e9 (2021).

88. B. Ariano, V. Mattiangeli, E. M. Breslin, E. W. Parkinson, T. R. McLaughlin, J. E. Thompson, R. K. Power, J. T. Stock, B. Mercieca-Spiteri, S. Stoddart, C. Malone, S. Gopalakrishnan, L. M. Cassidy, D. G. Bradley, Ancient Maltese genomes and the genetic geography of Neolithic Europe. *Curr. Biol.* **32**, 2668–2680.e6 (2022).

89. M. Srigyan, H. Bolívar, I. Ureña, J. Santana, A. Petersen, E. Iriarte, E. Kırdök, N. Bergfeldt, A. Mora, M. Jakobsson, K. Abdo, F. Braemer, C. Smith, J. J. Ibañez, A. Götherström, T. Günther, C. Valdiosera, Bioarchaeological evidence of one of the earliest Islamic burials in the Levant. *Commun Biol* **5**, 554 (2022).

90. K. Wang, K. Prüfer, B. Krause-Kyora, A. Childebayeva, V. J. Schuenemann, V. Coia, F. Maixner, A. Zink, S. Schiffels, J. Krause, High-coverage genome of the Tyrolean Iceman reveals unusually high Anatolian farmer ancestry. *Cell Genom* **3**, 100377 (2023).

91. L. G. Simões, R. Peyroteo-Stjerna, G. Marchand, C. Bernhardsson, A. Vialet, D. Chetty, E. Alaçamlı, H. Edlund, D. Bouquin, C. Dina, N. Garmond, T. Günther, M. Jakobsson, Genomic ancestry and social dynamics of the last hunter-gatherers of Atlantic France. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2310545121 (2024).

92. C. Valdiosera, T. Günther, J. C. Vera-Rodríguez, I. Ureña, E. Iriarte, R. Rodríguez-Varela, L. G. Simões, R. M. Martínez-Sánchez, E. M. Svensson, H. Malmström, L. Rodríguez, J.-M. Bermúdez de Castro, E. Carbonell, A. Alday, J. A. Hernández Vera, A. Götherström, J.-M. Carretero, J. L. Arsuaga, C. I. Smith, M. Jakobsson, Four millennia of Iberian biomolecular prehistory illustrate the impact of prehistoric migrations at the far end of Eurasia. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 3428–3433 (2018).

93. G. Renaud, U. Stenzel, J. Kelso, leeHom: adaptor trimming and merging for Illumina sequencing reads. *Nucleic Acids Res.* **42**, e141 (2014).

94. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

95. F. Ramírez, D. P. Ryan, B. Grüning, V. Bhardwaj, F. Kilpert, A. S. Richter, S. Heyne, F. Dündar, T. Manke, deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res.* **44**, W160–5 (2016).

96. B. S. Pedersen, A. R. Quinlan, Mosdepth: quick coverage calculation for genomes and exomes. *Bioinformatics* **34**, 867–868 (2018).

97. N.-C. Chen, B. Solomon, T. Mun, S. Iyer, B. Langmead, Reference flow: reducing

1333   reference bias using multiple population genomes. *Genome Biol.* **22**, 8 (2021).

1334   98.   T. L. Bailey, J. Johnson, C. E. Grant, W. S. Noble, The MEME Suite. *Nucleic Acids Res.*
1335   **43**, W39–49 (2015).

1336   99.   S. Myers, R. Bowden, A. Tumian, R. E. Bontrop, C. Freeman, T. S. MacFie, G. McVean, P.
1337   Donnelly, Drive against hotspot motifs in primates implicates the PRDM9 gene in meiotic
1338   recombination. *Science* **327**, 876–879 (2010).

1339   100.   S. Kurtz, A. Phillippy, A. L. Delcher, M. Smoot, M. Shumway, C. Antonescu, S. L.
1340   Salzberg, Versatile and open software for comparing large genomes. *Genome Biol.* **5**, R12
1341   (2004).

1342

1343   **Acknowledgements**

1344   We thank Sabriya Syed, Valerie Reyes-Ortiz, Nicholas Moskwa, Pille Hallast, Carmen Robinett,

1345   Stefan Ruhl, Victor Albert, Vincent Lynch, Derek Taylor, Leo Speidel, and John Novembre for

1346   technical help, discussions, suggestions, and feedback on the manuscript; the Human Genome

1347   Structural Variation Consortium and the Human Pangenome Reference Consortium for making

1348   their data publicly available; The Jackson Laboratory Scientific Services, including the Genome

1349   Technologies Service for expert assistance with the work described herein, and Research IT for

1350   computational infrastructure and support. We are grateful to the people who generously

1351   contributed their samples to the 1000 Genomes Project.

1352

1353   **Funding**

1354   C.L., and F.Y. were supported by the National Institute of Health (NIH) U24HG007497; K.K. was

1355   supported by The Jackson Laboratory Postdoctoral Scholar Award; O.G. was supported by the

1356   National Science Foundation (NSF) Awards # 2049947 and #2123284; C.R.B was supported by

1357   NIH GM133600.

1358

Authors Contributions

O.G. and C.L. conceived the study. F.Y. performed the analysis and interpretation of the Human Genome Structural Variation Consortium (HGSVC), and Human Pangenome Reference Consortium (HPRC) Bionano Genomics optical mapping, PacBio HiFi sequencing and phased assemblies, performed haplotype-resolved assemblies of HGSVC PacBio HiFi samples using hifiasm and amylase haplotype detection using HGSVC, and HPRC datasets. C.K. performed the mutational mechanisms analyses, breakpoint characterization, linkage disequilibrium, PCA and phylogenetic analyses. C.K. contributed to evolutionary and functional analysis. K.K. performed gene annotation, singleton analysis, selection/phylogenetic analyses of amylase coding sequences/segments, and interpretation/time estimation of initial *AMY1* gene duplication events. F.Y. and K.K. performed archaic hominin genome processing. P.P. performed ddPCR validation experiments, analysis of functional site differences in amylase amino acid sequences, and data visualization. K.S. calculated amylase gene copy numbers and performed downsampling in ancient human genomes. F.Y., C.K., K.K., P.P., C.R.B., A-M.T., C.L. and O.G. drafted and critically revised the article. Final approval of the version to be published was given by F.Y., C.K., K.K., P.P., K.S., C.R.B., A-M.T, C.L. and O.G. All authors read and approved the final manuscript.

Competing Interests

C.L. is a scientific advisory board member of Nabsys and Genome Insight.

Supplementary Materials
Supplementary Results

Supplementary Materials and Methods

1381   Figs. S1 to S40

1382   Tables S1 to S15

1383   References

1384

A

dN/dS estimate

23 SNV, 6 aa — **AMY2B** — p=0.0003

23 SNV, 11 aa — **AMY2A** — p=0.0451

36(1) SNV, 19(1) aa — **AMY1** — p=0.0226

Expectation for unconstrained sequence

B   Amino acid alignment across AMY2B, AMY2A, and AMY1

AMY2Bs, AMY2As, Chimp AMY1   I

98.2% AMY1s   L

1.8% AMY1s   I

DIDIGVAGFRIDASKHMWPGDI

LIDIGVAGFRLDASKHMWPGDIK

DIDIGVAGFRLDASKHMWPGDIA

position 211

AMY2Bs, AMY2As, Chimp AMY1   G

98% AMY1s   G

2% AMY1s   V

PRYFENGKDVNDW

PRYFENXKDVNDW

PRYFENVKDVNDW

position 366

A Archaic Hominin *AMY1* Copy Number

RNPC3 · AMY2B · AMY2A · AMY1

GoyetQ56-1
Denisova11

Normalized read depth(X)

GRCh38 chromosome 1(Mb)

103.50 · 103.55 · 103.60 · 103.65

B Ancient Pre-Agriculture *AMY1* Copy Number

③ Anzick ~12,500 YA

Peștera Muierii ~34,000YA
④ ~13,000YA
⑧
⑧ ~9,500YA
⑤ ~10,000YA

Ust'-Ishim ~45,000YA
⑥

C

AMY2A Copy Number

pre-agriculture · during transition · post-transition

0.29 · 0.07 · 0.07

AMY2B Copy Number

pre-agriculture · during transition · post-transition

0.09 · 0.08 · 0.56

AMY1 Copy Number

pre-agriculture · during transition · post-transition

0.33 · 0.01 · 0.01

D Spread of Agriculture through Europe

1KYA
5.5KYA
6KYA
7KYA
8.5KYA
10KYA

Hunter-Gatherer
Farmer
▲ 3,000-900YA (post)
● 9,000-3,000YA (during)
✕ >9,000YA (pre)

A

AMY2B   *AMY2A*   *AMY1A*   *AMY1B*   *AMYP1*   *AMY1C*

PRDM9 Binding sites

NAHR Breakpoints

MMBIR Breakpoints   • •Microhomology

B   NAHR - *AMY1* CNVs

H5.3

H3ʳ.1

H3ʳ.1

H1ᵃ.1

50kb   100kb   150kb   200kb   250kb

H5.3   A   B|A   B

H3ʳ.1   A   B   H3ʳ.1

H3ʳ.1   A   B

H1ᵃ.1   A|B

Parsimony
Informative sites

A   G   C

B   A   T

A|B   G   T

B|A   A   C

C   MMBIR - *AMY2* CNVs

H3ʳ.1

H2A2

B2.1

50kb   100kb   150kb   200kb

H3ʳ.1   B   C   A   D

A

C   B

D

H2A2B2.1   AB   CD

AB

Microhomology at
the junctions

...AATT**TGTT**
TGTTGTTA....

Human-Chimpanzee split

Fire allows for cooking of starch

Initial *AMY1* duplication

Human-Neanderthal split

Ust'-Ishim 6 *AMY1* genes

Pre-agriculture expansion

Post-agriculture expansion

Population-variable soft sweeps

Archaic
Ancient
Recent

~6MYA t_1 t_2 ~800KYA   ~45KYA t_3 t_4 t_5

AMY1 copies per diploid cell

Starch consumption

AMY1A
AMY1B
AMY1C

AMY1

Archaic Hominins

Modern Humans

Chimpanzee