1 **Epiregulon: Inference of single-cell transcription factor activity to dissect mechanisms of**
2 **lineage plasticity and drug response**
3

4 Tomasz Włodarczyk[1], Aaron Lun[1], Diana Wu[2], Shreya Menon[3,4,5], Shushan Toneyan[1], Kerstin
5 Seidel[2], Liang Wang[2], Jenille Tan[2], Shang-Yang Chen[1], Timothy Keyes[1], Aleksander
6 Chlebowski[1], Yu Guo[1], Ciara Metcalfe[2], Marc Hafner[1,2], Christian W. Siebel[2], M. Ryan
7 Corces[3,4,5], Robert Yauch[2]*, Shiqi Xie[2]*, Xiaosai Yao[1,2]*
8
[1]Department of Computational Sciences, Genentech Inc, South San Francisco, CA, USA
[2]Department of Discovery Oncology, Genentech Inc, South San Francisco, CA, USA
[3]Gladstone Institute of Neurological Disease, Gladstone Institutes, San Francisco CA, USA
[4]Gladstone Institute of Data Science and Biotechnology, Gladstone Institutes, San Francisco
CA, USA
[5]Department of Neurology, University of California San Francisco, San Francisco CA, USA
15

*correspondence

yauch.bob@gene.com; xie.shiqi@gene.com; yao.xiaosai@gene.com

Abstract

Transcription factors (TFs) and transcriptional coregulators represent an emerging and exciting
class of targets in oncology. Gene regulatory networks (GRNs) evaluate pharmacological
agents against these regulators through quantifying target gene modulation. However, methods
relying solely on gene expression cannot adequately model post-transcriptional interference of
TF functions. We introduce Epiregulon, a method to predict TF activity at the single-cell level
using GRNs constructed from RNA-seq and ATAC-seq data. Considering co-occurrence of
gene expression and chromatin accessibility addresses the decoupling of TF activity from gene
expression. The incorporation of ChIP-seq data extends inference to transcriptional coregulators
lacking defined motifs and facilitates the identification of interaction partners. We uncover
divergent cell fate transitions of prostate cancer cells driven by NKX2-1 and GATA6
overexpression. Epiregulon outperforms other methods in its consistency and accuracy in
predicting the effects of AR inhibition. Finally, degrader-treated networks recapitulate the
context-dependent activity of SMARCA4 consistent with the known etiologies of prostate
cancers. By mapping out gene regulation across a multitude of perturbations, Epiregulon can
accelerate the discovery of therapeutics targeting transcriptional regulators.

Introduction

Transcription factors and transcriptional coregulators shape cell fates and lineage commitment, and their dysregulation drives congenital diseases and tumor growth. Transcription factors bind to specific DNA sequences whereas coregulators interact with transcription factors in a context-specific manner and lack defined motifs. These transcriptional modulators represent an important class of therapeutic targets in oncology and beyond. Therapeutics targeting transcriptional regulators inhibit their activity by either blocking the ligand binding domains[1], degrading the protein[2,3] or disrupting protein-protein interactions[4]. Delineating the functions of transcriptional regulators can accelerate our understanding of disease biology and drug discovery.

Gene regulatory networks (GRNs) model the underlying circuitry of gene regulation and have been applied to understand lineage commitment and plasticity and drug response[5]. Early methods to construct gene regulatory networks relied exclusively on gene expression data and sought to identify association between expression of target genes and transcription factors[6,7]. Now, single-cell multiomics technologies provide chromatin accessibility information in addition to gene expression and recent GRN inference methods leverage both modalities to improve performance (Supplementary Table 1). Most of these methods, including FigR[8], Pando[9] and GRaNIE[10], rely on linear relationships between transcription factors and their target genes to model gene regulation. In addition, SCENIC+ utilizes random forest regression to model the regulatory relationships between TF, regulatory elements and target genes[11].

Several challenges still impede the broader utilization of these GRN inference methods in basic biology and drug discovery. First, none of these methods are specifically designed to predict changes in which the TF activity is decoupled from its gene expression. These include drug perturbations in which inhibitors disrupt protein complex formation or localization, and post-translational modifications that can impact TF activity and also CRISPR genome editing in which genes can be silenced by insertions or deletions at the DNA level but without causing appreciable changes in gene expression. Second, the use of motif sequences in GRN construction precludes prediction of transcriptional coregulators, an important class of regulators without defined motifs. Lastly, current studies rely on ChIP-seq and HiC data to benchmark GRNs but TF occupancy and chromatin conformation do not directly measure the true regulatory activity of TFs. Therefore, erroneous estimation of TF activity misled by gene expression, sole reliance on TF motifs and lack of appropriate benchmarking datasets are three limitations in current GRN inference methods.

Here we present Epiregulon, a method that constructs a GRN from single-cell multiomics data and TF occupancy data to infer activity of transcriptional regulators. Unlike most other GRN tools, Epiregulon utilizes accompanying ChIP-seq data, thus allowing inference of both transcription factors and transcriptional coregulators lacking defined motifs. Epiregulon utilizes weight estimation methods that account for co-occurrence of both TF gene expression and chromatin accessibility to handle decoupling of TF expression and activity. Functional interpretation of GRNs through Jaccard similarity of target genes and pathway annotations uncovers drivers and interaction partners of lineage transition and drug response in a cell-state

79 specific and context-aware manner. We performed simulations and generated ground truth data
80 to robustly benchmark its performance in a multitude of datasets including lineage development
81 and drug treatment. Epiregulon is implemented as a R package and interfaces with ArchR[12] or
82 any other preprocessing tools.
83
84 **Results**
85
Epiregulon constructs GRNs to infer TF activity at the single cell level
Epiregulon aims to map out target genes of transcriptional regulators and track changes in their
regulatory potential in each cell state, thereby identifying master regulators of cell fate changes.
It assumes that a transcription factor or a transcriptional coregulator (collectively referred to as
TF for simplicity) binds to regulatory elements (REs) and modulates the expression of nearby
target genes (TGs). Epiregulon assigns regulatory elements (REs) to their putative target genes
based on correlation between chromatin accessibility and gene expression (Figure 1). Each
regulatory element is then interrogated for TF occupancy based on user-provided or/and a
compilation of public TF ChIP-seq data from ChIP-Atlas and ENCODE. The use of ChIP-seq
data enables us to extend activity inference to chromatin modifiers and co-factors which do not
have well-defined binding motifs. Furthermore, accumulating evidence suggests that TF
occupancy is determined not just by motif sequences but also by other factors including
interactions with other TFs and co-factors, local DNA structure and genomic features[13]. A
substantial proportion of TF binding sites do not harbor motifs[14]. Therefore, ChIP-seq data is
likely to yield a more relevant set of target genes than motif annotations are.
101
We combine the RE-TG links with TF occupancy at REs to establish an initial unweighted
tripartite graph with TFs, REs and TGs being the nodes and their connections being edges. An
optional pruning step of the graph removes potentially spurious regulatory relationships and TFs
with undetectable gene expression. We then estimate the weights of the TF-RE-TG edges using
the Wilcoxon test statistic, computed by comparing the ranks of the target gene expression
between the group of cells that express both the corresponding TF and have chromatin
accessibility at that particular RE versus the remaining cells. This graph becomes the final gene
regulatory network returned by Epiregulon. Finally, the activity of a TF is defined as the
weighted sum of its target gene expression normalized by the number of target genes. The
exact algorithm is described in the Methods; see also Supplementary Note Section 1 for
comparison of parameter choices.
113
It is of great interest to understand the functions of a TF through its differential activity between
cell states or treatment conditions. Differential TF activity can result from at least two different
scenarios; 1) TFs can be turned on or off by changes in gene expression, or 2) they can
redistribute to different genomic regions due to mutations or hijacking by other factors. The first
scenario can be adequately captured by differences in total TF activity whereas the second
scenario requires the consideration of regulon membership. In addition to quantifying differential
total TF activity between cell states, Epiregulon employs a second method for differential
network analysis. We use the Wilcoxon method to derive cluster-specific weights, subtract the

edge weights between two networks and compute the degree centrality of the resulting edge-
subtracted network. See Supplementary Note Section 2 for simulation of differential networks.
124
125 Finally, it is desirable to interpret the functions of a TF through the pathway enrichment of its
126 regulons and through its interaction partners. Epiregulon provides pathway enrichment of the
127 regulons and further utilizes Jaccard similarity to estimate the overlap between regulons to
128 identify potential interaction partners. These functional annotations of GRNs serve to provide
129 biological interpretability.
130
Epiregulon identifies cell fate changes
To evaluate Epiregulon's performance, we generated data using Reprogram-Seq[15] to model
lineage transition of prostate adenocarcinomas by overexpressing defined transcription factors.
We separately introduced lentivirus encoding NKX2-1 and GATA6 into a prostate
adenocarcinoma cell line LNCaP and tracked changes in transcriptomic and epigenomic profiles
at the single cell level. NKX2-1 is a known driver of AR-independence and neuroendocrine
transition in prostate cancer[16,17] whereas GATA6 has unknown function in the prostate. These
factors are not expressed in parental LNCaP and were used for subsequent benchmarking
analysis. Reprogram-Seq data could serve as the ground truth to benchmark the inference of
TF activity because we expect to see increased TF activity in cells with exogenous expression
of NKX2-1 and GATA6.
142
Expression of these transcription factors altered the cell states of LNCaP, forming distinct
clusters with cluster 1 enriched in cells expressing GATA6 and cluster 3 enriched in cells
expressing NKX2-1 (Figure 2a). Due to the low expression of transcription factors and 3' bias of
scRNAseq, gene expression could not accurately map the identity of TFs being introduced
(Figure 2a). In contrast, Epiregulon enhanced the dynamic range of TF activity and retrieved the
correct cell identity (Figure 2a).
149
We performed a systematic benchmarking exercise to evaluate the performance of Epiregulon
against other multiomic GRN inference methods. Compared to FigR, Pando, GRaNIE and
SCENIC+, Epiregulon was the only method that was able to predict NKX2-1 activity (Figure 2b)
because the use of ChIP-seq data recovered the largest collection of regulators (Figure 2c). For
GATA6, Epiregulon had the second highest area under the receiving operating characteristic
(AUROC) value (Figure 2b). Epiregulon gave consistent predictions across multiple weight
methods (Figure S1a) and offers improved sensitivity in predicting activity of those factors with
low gene expression.
158
We further investigated the accuracy of the target genes inferred. We showed that both
unpruned and pruned GATA6 and NKX2-1 targets inferred by Epiregulon ranked highly by
differential expression (Figure S1b). Furthermore, cluster-specific GRN showed that NKX2-1
targets were exclusively expressed in cluster 3 while GATA6 targets were expressed in cluster 1
(Figure S1c). As another independent validation, we engineered a dox-inducible cell line
expressing NKX2-1 or eGFP as a control. The log changes of our inferred NXK2-1 targets upon
dox treatment correlated with the NKX2-1 target gene weights inferred by Epiregulon (Figure 2d,

S1d), suggesting that the GRN weights inferred by Epiregulon are informative of regulatory
strength.
168
Global analysis of TF activity identified other TFs whose activities were concomitantly altered by
introduction of these reprogramming factors. Jaccard similarity found FOSL2 (a paralog of
FOSL1), YAP1 and TEAD1 to be amongst the top interaction partners of GATA6 (Figure 2e,
S1e). These factors are known to reprogram prostate adenocarcinomas towards a stem cell-like
(SCL) state[18]. This led us to examine whether GATA6 itself could also reprogram LNCaP cells
towards the SCL subtype. We examined regions that are uniquely accessible in the SCL
tumors[18]. Strikingly, GATA6 overexpression led to increased chromatin accessibility at these
SCL-specific regions and GATA6 activity was highly correlated with the combined chromatin
accessibility score (Figure 2f, 2g). Lineage trajectory resulting from GATA6 expression differed
from the neuroendocrine cell state induced by NKX2-1 (Figure 2f, 2g, S1f). Taken together,
Epiregulon uncovered a previously unknown causal relationship between GATA6 expression
and lineage transition to SCL subtype of prostate cancer.
181
182
Epiregulon predicts responses towards TF targeting drugs
In the previous example, Epiregulon was able to predict changes in TF activity when they are
largely driven by changes in their expression. A more challenging task is to predict changes
when TF activity is decoupled from its gene expression. We generated a second dataset to
evaluate changes in AR activity following drug treatment. In total, 4 prostate cancer cell lines (2
AR-dependent and 2 AR-independent) were treated with 3 AR-modulating agents (Figure 3a).
The first agent is the clinically approved AR antagonist, enzalutamide, which interferes with AR
protein function by blocking its ligand binding domain[1]. The second agent is ARV-110, a
degrader of AR protein which acts by bringing an E3 ubiquitin ligase in close proximity to an AR
protein[3]. We also synthesized a third agent, SMARCA2.4_1, a degrader of the SWI/SNF
chromatin remodeling complex (Figure S2a-b); inhibition of the same protein complex is known
to decrease AR activity by evicting it from the chromatin[19]. These pharmacological agents do
not directly or consistently suppress the transcription of AR target genes, thus serving as an
example of the decoupling of the TF expression from its activity.
197
Epiregulon predicted decreased AR activity following enzalutamide and ARV110 treatment in
the known sensitive cell lines, LNCaP and VCaP and minimal changes in AR activity in the AR-
independent cell lines, DU145 and H660 (Figure 3b, Figure S2c). Other GRN inference methods
failed to predict decreased AR activity in either or both cell lines (Figure 3c). Gene expression of
AR was not informative of the effect on AR activity because enzalutamide increased AR gene
expression in VCaP cells despite its antagonistic function (Figure 3d). Like the Reprogram-Seq
data, Epiregulon recovered the greatest number of regulators (Figure S2d). By estimating
weights based on both TF expression and RE chromatin accessibility, Epiregulon outperformed
methods utilizing TF expression alone, and robustly predicted decreased AR activity despite
discordant AR gene expression (Figure 3c, Figure S2e). ChIP-seq data generated from merging
ChIP-atlas and ENCODE yielded comparable performance as cell line-matched ChIP-seq data

(Figure S2d, f). ChIP-seq data can outperform motif annotations using deep learning sequence-
to-function models or deep neural networks (DNNs) (see Supplementary note section 3).
211
Epiregulon could identify putative interaction partners to a regulator of choice. Using GRNs
derived from ARV110 and Enzalutamide treated cells, we retrieved FOXA1, a well-known
interaction of AR in both LNCaP and VCaP (Figure 3e). We also found ERG to be interacting
with AR in VCaP cells which harbor a TMPRSS2-ERG fusion, but not in LNCaP cells which lack
such a fusion. Top interaction partners also included lesser known but validated interaction
partners of AR, namely GATA2[20] and SUMO2[21,22](Figure 3e). DepMap shows dependency of
LNCaP and VCaP on SUMO2 and GATA2 (Figure 3f). Therefore, Epiregulon was able to
evaluate the efficacy of therapeutic agents targeting AR and identify additional interaction
partners including SUMO2 and GATA2 which are potential drug targets themselves.
221
Epiregulon predicts response of drug targeting transcriptional coregulator
Transcriptional coregulators cooperate with transcription factors to regulate gene expression
and are attractive targets in oncology. The challenge with predicting responses of transcriptional
coregulators is that these factors do not contain defined motifs as they can interact with different
transcription factors depending on the cellular context. As such, GRN inference methods which
rely on TF motifs cannot predict the activity of these factors. SMARCA2 and SMARCA4, two
mutually exclusive paralog proteins encoding the ATPase subunit of the SWI/SNF chromatin
remodeler are examples of transcriptional coregulators. We evaluated the performance of
Epiregulon based on its ability to predict decreased SMARCA2/4 activity upon treatment with a
degrader of SMARCA2/4 protein. We utilized the same AR dataset as in Figure 3 but focused
on the third agent, SMARCA2.4_1. We performed most of our analysis on SMARCA4 since it is
the predominant paralog that affects the proliferation of prostate cancer cell lines[23].
Pharmacological degradation of SMARCA4 allows benchmarking of activity inference for a
highly context-dependent transcriptional coregulator lacking a defined motif.
236
To circumvent the lack of motifs, Epiregulon utilizes public SMARCA4 ChIP-seq to infer the
most likely target genes in a cell-line aware manner. Epiregulon predicted decreased SMARCA4
activity across all 4 prostate cancer cell lines (Figure 4a). None of the other GRN inference
methods we benchmarked were able to predict the activity of SMARCA4. Within Epiregulon,
weight estimation methods that take into account both TF gene expression and RE chromatin
accessibility outperformed methods that only consider TF expression (Figure S3a).
Furthermore, SMARCA4 activities calculated from merged ChIP-seq and cell line-matched
ChIP-seq were consistent (Figure S3b). The SMARCA4 regulons recapitulated the unique
etiologies of prostate cancer, with AR-dependent cell lines LNCaP and VCaP enriched for
androgen response, stem-cell like cell line DU145 enriched for activation of YAP activity[18] and
the neuroendocrine cell line NCI-H660 enriched for the loss of TP53, activation of EZH2[24,25] and
MAPK signaling[26] (Figure 4b).
249
Epiregulon can inform mechanisms of action of the SMARCA4 degrader by discovering TFs co-
perturbed with SMARCA4 (Figure 4c). We identified differential TFs by subtracting DMSO
treated from SMARCA4 degrader treated networks. TFs were ranked by Jaccard similarity with

SMARCA4 after background correction using edge permuted networks. Common to all 4 cell
lines, highly ranked TFs included components of general transcriptional machinery including
BRD4 and EP300 and also regulators of chromatin structure including components of the
cohesin complex (RAD21, SMC3, SMC1A) and chromosome looping (YY1, CTCF and MED1).
These proteins have been previously implicated to associate with SMARCA4 (Table 1). Unique
to the individual cell lines, SMARCA4 interacts with AR in the AR-dependent cell lines (LNCaP
and VCaP). In the neuroendocrine cell line NCI-H660, SMARCA4 interacts with TCF12, a factor
known to be important for neuronal development[27] and EZH2, a factor known to mediate
neuroendocrine transition in prostate cancer[25]. Taken together, Epiregulon expands the
inference of activity to include transcriptional coregulators lacking defined motifs and is able to
capture their context specificity and infer the most relevant biological processes.
264
265 Table 1: Associated partners of SMARCA4 identified by differential network analysis

Function	Gene	References
Transcription	BRD4	25,28, 29, 30, 31
Transcription	EP300	32, 33, 34
Chromosome structure	Cohesin (RAD21, SMC1A, SMC3)	35, 36
Chromatin looping	YY1 and CTCF	37, 38

266
267
Epiregulon identifies lineage factors and predicts effects of perturbations
Finally, we ask whether regulons generated from single-cell multiomic data can be applied to
scRNA-seq dataset lacking chromatin information. We first used Epiregulon to compute the
activity of lineage factors in a scMultiomic dataset of human peripheral blood mononuclear cells
(PBMCs)[39] (Figure 5a). Epiregulon recapitulated the lineage-specific activity of known lineage
factors including GATA3 in T and NK cells, EBF1 and PAX5 in B cells and SPI1 in myeloid cells
(Figure 5b), and gene set enrichment of their target genes is consistent with their known
biological functions (Figure S4a). We used the Wilcoxon method to obtain cell-type specific
weights for all TF-TG pairs. As an example to illustrate the specificity of the GRN, the B cell
lineage factor EBF1 had its target genes largely confined to preB cells but not to CD4 memory T
cells (Figure 6c). We further utilized Epiregulon's similarity function to identify known interacting
partners (Figure 6d). The most similar TF to EBF1 was PAX5; EBF1 and PAX5 regulate B cell
development in a highly coordinated manner[40]. We further compared the global network
topology of monocytes and CD4 memory T cells; degree centrality of the differential graph
revealed known lineage factors of monocytes (SPI1, CEBPA, NR4A1)[41,42] and CD4 T cells
(TBX21, GATA3, LEF1, FOXO1)[43,44], suggesting that Epiregulon is able to prioritize lineage
factors from differential network analysis.
285

Next, we applied the GRN of the above PBMC dataset to a new scRNAseq dataset containing
K562, a lymphoblast cell line, engineered to express CRISPRa and guides targeting defined
factors[45]. The barcode information in each cell encodes the guide RNAs corresponding to the
TFs they activate and serves as the ground truth. GRN from Epiregulon better predicted the
effect of CRISPRa perturbations than gene expression, lineage-agnostic regulons (Dorothea[46]
and CollecTRI[47]) and lineage-specific regulons (Humanbase[48]) did, as reflected in the AUROC
values (Figure 5f, Figure S4b). Epiregulon also captured the highest number of TFs (Figure 5f).
The improvement in performance is largely due to the reduction of data sparsity by aggregating
the appropriate target genes, as shown by PRDM1 (Figure 5g). Taken together, based on this
ground truth dataset, GRNs reconstructed from Epiregulon could accurately identify known
lineage factors and predict effects of gene perturbation in datasets containing only scRNA-seq.
297
Discussion
We develop Epiregulon, a method to construct gene regulatory networks and infer activity of
both transcription factors and transcriptional coregulators lacking defined motifs, an important
class of regulators largely overlooked by previous GRN inference methods. When benchmarked
against other GRN inference methods, Epiregulon performs most robustly across a multitude of
datasets, especially when TF activity is decoupled from its gene expression such as in drug
perturbations. Epiregulon predicts the largest repertoire of regulators, constructing an extensive
regulatory network that is functionally annotated with pathway enrichment analysis and
interaction partners.
307
Unlike most other GRN inference methods, Epiregulon allows construction of GRNs from ChIP-
seq data but still preserves the option to filter ChIP-seq peaks based on the presence of motifs.
ChIP-seq data allows the prediction of chromatin co-activators and modifiers lacking defined
motifs and can often outperform motif annotations using deep-learning neural networks models.
Epiregulon was the only tool that was able to infer the activity of the chromatin remodeler and it
did so in a context-aware manner. Epiregulon predicted the suppression of SMARCA2/4 activity
after treatment by our newly synthesized degrader, and was able to identify the unique
interaction partners of SMARCA4 in both AR-dependent (AR, FOXA1, GATA2 and SUMO2) and
AR-independent prostate cancer cell lines (TCF12, EZH2).
317
Epiregulon's Wilcoxon weight estimation method, based on co-occurrence of TF gene
expression and chromatin accessibility, is especially well suited for applications in which TF
activity is discordant with its gene expression such as drug perturbations. The Wilcoxon method
also enables estimation of cell-cluster specific weights and subsequent differential network
topology analysis. Epiregulon outperformed other methods in its consistency and accuracy in
predicting the effects of multiple AR-targeting agents, and identifies both well-known (FOXA1)
and lesser-known interaction partners of AR (GATA2 and SUMO2). These interaction partners
hold the potential to become additional drug targets themselves.
326
Epiregulon enables us to make new discoveries from the ground truth data we generated. We
found an unexpected connection between GATA6, YAP1 and TEAD1 which led to the discovery
that GATA6 overexpression reprograms prostate adenocarcinoma cell line LNCaP towards a

stem-cell like subtype. This was possible because compared to other methods, Epiregulon
covers the most extensive regulators including coregulators lacking motifs (in this case YAP1),
enabling the identification of interaction partners based on shared target genes.

Several limitations of our study remain. While Epiregulon is robust in predicting the overall TF
activity, we have not formally evaluated the accuracy of individual target genes. Second,
Epiregulon does not explicitly model cooperativity between transcription factors and partners, so
direct interactions between functionally related factors may have been overlooked. Lastly, given
the use of ground truth data, our evaluation of performance is limited to a few transcription
factors and it remains unclear whether Epiregulon uniformly performs well across multiple TFs.
We speculate that the quality of available ChIP-seq data greatly influences the prediction of TF
activity, and wherever possible, ChIP-seq data of TF of interest should be generated in the
matching cell lines to ensure robustness of the biological findings.

We envision that Epiregulon can enable drug discovery by evaluating response to therapeutic
agents not limited to oncology and identifying new drivers of resistance. Epiregulon is available
at https://github.com/xiaosaiyao/epiregulon along with the data package scMultiome
(https://bioconductor.org/packages/release/data/experiment/html/scMultiome.html).

Methods

Epiregulon
The Epiregulon workflow consists of several components: GRN construction, pruning of
networks (optional), estimation of weights, calculation of activity, functional annotation of
regulons, differential network analysis and identification of interaction partners. Epiregulon is
designed to work seamlessly with ArchR but can accept any input as long as they are formatted
into SingleCellExperiment objects. It is available as three related R packages. Epiregulon is the
core   package   that   performs   GRN   construction   and   activity   inference
(https://github.com/xiaosaiyao/epiregulon/). Epiregulon.extra contains differential analysis and
plotting functions (https://github.com/xiaosaiyao/epiregulon.extra). Epiregulon.archr contains
functions interfacing with ArchR (https://github.com/xiaosaiyao/epiregulon.archr).

Data preprocessing
Epiregulon assumes that prior preprocessing of the data has been performed by users' methods
of choice, requiring paired gene expression matrix (scRNA) and peak matrix (scATAC) along
with dimensionality reduction matrix as the input. If gene expression and peak matrix are not
paired, integration of scATAC-seq and scRNA-seq data must be performed prior to Epiregulon
using methods such as ArchR's addGeneIntegrationMatrix function.

For the datasets included in this manuscript, reads were mapped by the Cell Ranger ARC 2.0
and were further processed with the use of ArchR[12]. Briefly, cells were filtered based on TSS
enrichment (>6) and number of ATAC-seq reads mapped to the nuclear genome (>2500 and
$<10^5$). Moreover, doublets were removed using dedicated ArchR functions. ATAC-seq data was
represented as a 500bp tile matrix on which iterative LSI was performed. Similarly, normalized

gene expression data was subject to iterative LSI. The dimensionality reductions from both
modalities were then combined into a single one that was used for cell clustering. The ATAC-
seq peaks were called separately for each cluster and after normalization were merged to
produce one peak set for all cells (peak x cell matrix). The cells were assigned to their sample
barcodes using demuxEM[49].
379
Network construction
Epiregulon provides two similar methods to establish peak to gene links using the calculateP2G
function. If there is an existing ArchR project, Epiregulon retrieves the peak to gene links that
have been previously assigned by ArchR. Briefly, ArchR creates 500 cell aggregates,
resampling cells if needed. ArchR computes the correlation between chromatin accessibility and
target genes within a size window (default ±250 kb) and retains peak-gene pairs that exceed a
correlation threshold (default Epiregulon cutoff is 0.5). In the absence of an ArchR project,
Epiregulon defines cell aggregates using k-means clustering based on the reduced
dimensionality matrix and performs correlation in the same manner as ArchR does. If cluster
labels are provided, cell aggregates are formed within each cluster and cluster-specific
correlations are reported in addition to overall correlations.
391
TF occupancy data
Each regulatory element is then interrogated for TF occupancy based on a compilation of public
TF ChIP-seq binding sites. ChIP-seq were downloaded from ChIP-Atlas (chip-atlas.org) and
ENCODE (encodeproject.org). TF ChIP-seq peaks with FDR < 1 x 10^{-5} were retained. ChIP-seq
datasets of the same TFs were merged, yielding 1557 unique factors (human) and 768 unique
factors (mouse). Data is provided in the scMultiome package as GRanges list objects and can
be accessed using the getTFMotifInfo function from the Epiregulon package or directly from the
scMultiome package using the tfBinding function.
400
If desired, ChIP-seq peaks can be further annotated for the presence of motifs using the
addMotifScore function. If motif annotation has been performed previously using ArchR's
addMotifAnnotations function, motif annotations can be easily retrieved and appended to the
peak matrix. In the ArchR independent workflow, Epiregulon can annotate peak matrix using
motifmatchr's motif matching function[50] and cisbp as the reference motif database[51].
Alternatively, users can start entirely from motif annotations and leverage deep learning motifs
to select motifs based on motif importance scores (see section on DNN models).
408
409 *Network pruning (optional)*
410 Epiregulon prunes the network by performing tests of independence on the observed number of
411 cells jointly expressing transcription factor (TF), regulatory element (RE) and target gene (TG)
412 vs the expected number of cells if TF/RE and TG are independently expressed using the
413 pruneRegulon function.
414
415 We define n as the total number of cells, k as the number of cells jointly expressing TF, TG and
416 RE above a set threshold, g as the number of cells jointly expressing TF and RE above a

417   threshold and h as the number of cells expressing TG above a threshold. p, the expected
418   probability of cells jointly expressing TF, TF and RE above a threshold is defined as follows:
419

$$P(cells\ expected\ to\ jointly\ express\ TF\ t, RE\ r\ and\ TG\ g) = p\ = \frac{g}{n} \times \frac{h}{n}$$

420
421   Two tests of independence are implemented, the binomial test and the chi-square test. In the
422   binomial test, the expected probability is p, the number of trials is the total number of cells n,
423   and the observed number of successes is k, the number of cells jointly expressing all three
424   elements.
425
426   In the chi-square test, the expected probability for having all 3 elements active is also p. The
427   observed cell count for the active category is k, and the cell count for the inactive category is
428      $n - k$. P-values are calculated from chi-squared distribution with degree of freedom equal to 1.
429   Cluster-specific p-values are calculated if users supply cluster labels. Finally, multiple
430   hypothesis testing was performed by the Holm method.
431
Estimation of weights
While network pruning provides statistics on the joint occurrence of TF-RE-TG, we would like to
further estimate the strength of regulation using the addWeights function. Biologically, this can
be interpreted as the magnitude of gene expression changes induced by changes in
transcription factor activity. Epiregulon provides 3 different methods to estimate weights. Two
measures (correlation and Wilcoxon statistics) give both the magnitude and directionality of
changes whereas weights computed by mutational information (MI) are always positive. Within
2 of the methods (correlation and MI), there is an option of modeling the TG expression based
only on TF expression, or on the product of TF expression and RE chromatin accessibility.
Consideration of both TF expression and RE chromatin accessibility is highly recommended,
especially for scenarios in which TF activity and TF gene expression are decoupled (as in the
case of drug perturbations or CRISPR genome editing). The correlation and mutual information
statistics are computed on pseudobulks by user-provided cluster labels and yield a single weight
across all clusters per each TF-RE-TG triplet. In contrast, the Wilcoxon method groups cells
based on the joint expression of TF, RE and TG in each single cell or in cell aggregates. Cell
aggregation uses a default value of 10 cells and can help overcome sparsity and speed up
computation. If cluster labels are provided, we can obtain cluster-specific weights using the
Wilcoxon method.
450
451   1) Wilcoxon test statistics
452   Cells were divided into two groups, with the first group jointly expressing TF gene expression
453   and chromatin accessibility at RE, and the remaining cells in the second group. The default
454   cutoff is 1 for normalized gene expression and 0 for normalized chromatin accessibility. There is
455   also an option to use the median of each feature as an adaptive cutoff.
456
2) Correlation

If we consider only TF expression (tf_re.merge set to FALSE), the weight is the correlation
coefficient between the TF gene expression and TG gene expression. If we consider both TF
expression and RE chromatin accessibility (tf_re.merge set to TRUE), the weight is the
correlation coefficient between the product of TF gene expression and RE chromatin
accessibility vs the TG gene expression.
463
464   3) Mutual information between the TF and target gene expression,
465
466   If we consider only TF expression (tf.re_merge is set to FALSE), the weight is the mutual
467   information between the TF gene expression and the TG gene expression. If we consider both
468   TF expression and RE chromatin accessibility (tf.re_merge is set to TRUE), the weight is the
469   mutual information between the product of TF gene expression and RE chromatin accessibility
470   vs the TG gene expression.
471
472   *Calculation of TF activity*
473   The activities for a specific TF in each cell are computed by averaging the weighted expressions
474   of target genes linked to the TF:
475
476

$$K_{t,c} = \frac{1}{q_t} \sum_{g \in R_t} \beta_{t,g,r} Y_{g,c}$$

477
478   where $K_{t,c}$ is the activity of a TF t for a cell c, q is the total number of target genes for a TF t, Y_g
479   is the normalized count of target gene g where g is regulated by TF t and $\beta_{t,g,r}$ is the regulatory
480   weight of TF t on target gene g at RE r. R_t is the regulon of TF t.
481
482   If cluster labels are provided, cluster-specific weights are used.
483
484   *Gene set enrichment of regulons*
485   Gene set enrichment of a regulon is performed by testing whether the target genes of a TF are
486   over-represented in known gene signatures such as those provided by MSigDB using a
487   hypergeometric test. Target genes can be refined by filtering the regulons on user-defined
488   weights.
489
Differential TF activity by total activity
This differential analysis compares the differences in the total activity of each transcription factor
between conditions. This analysis is well suited for identifying factors that have contrasting
levels of activities, for instance, lineage factors that are turned on or off during certain
developmental stages or cell types. TF activities are compared between groups using any
standard statistical methods. Here we use scran's findMarkers function to find differential activity
between user provided groups of cells (https://rdrr.io/bioc/scran/man/findMarkers.html).
497
Differential TF activity by network topology

A second approach to investigate differential TF activity is to compare and contrast target genes
or network topology. This is useful when a transcription factor differs in the target genes it
regulates, while maintaining a similar level of total activity. This can happen when a transcription
factor redistributes to a different set of genomic regions, due to mutations in the transcription
factors or changes in the interaction partners. Differences in network topology are calculated by
taking the degree centrality of the edge subtracted graphs between two conditions, with cluster-
specific regulon weights representing edge weights in each condition.
506

$G^{(1)}$ and $G^{(2)}$ are networks with identical node sets N and respective edge sets $E^{(1)}$ and $E^{(2)}$:

-    $E^{(1)} = \{e^{(1)}_1, e^{(1)}_2, e^{(1)}_3, ...e^{(1)}_m\}$
-    $E^{(2)} = \{e^{(2)}_1, e^{(2)}_2, e^{(2)}_3, ...e^{(2)}_m\}$

where $e^{(j)}_i$ represents the edge weight of the *i*th edge in network $G^{(j)}$.

The edge-subtracted network G' induced by $G^{(1)}$ and $G^{(2)}$ such that each edge e'_i in G' is given by
$e'_i = e^{(1)}_i - e^{(2)}_i$. We simplify the tripartite TF-RE-TG graph to a bipartite TF-TG graph by taking the
maximum of TF-RE-TG weights of the same TF-TG pairs. Degree centrality is the sum of the
weights of the TF-TG e'_i. Degree centrality is further normalized to account for differences in the
number of target genes of each transcription factor. The default normalization method is dividing
degree centrality by the square root of the number of target genes. This strikes a balance
between penalizing TFs with an abundance of target genes and prioritizing TFs with differential
target genes. Transcription factors are ranked by normalized degree centrality.
519

520 *Identification of interaction partners*
521 To identify potentially interacting TF or co-factors, we calculate Jaccard similarity between all
522 pairs of factors and rank factors by the degree of overlap in their regulons. The edges can be
523 filtered on a weight cutoff to retain the most confident set of target genes.
524

525 **Benchmarking**
526

Benchmarking using Reprogram-Seq data
We introduced 4 transcription factors into LNCaP cells - FOXA1, FOXA2, NKX2-1 and GATA6
and obtained paired gene expression and chromatin accessibility information. Because FOXA1
was already highly expressed in LNCaP cells at the basal level, its introduction did not have any
impact on lineage plasticity and therefore FOXA1 was excluded in subsequent analyses. We
focused on NKX2-1 and GATA6 because their expression resulted in distinct cell clusters. Each
cell is identified by the HTO tag corresponding to the well receiving virus encoding a particular
TF and this information serves as the ground truth cell labels. For each TF, a cell is classified
into either expressing this TF or not. AUROC was computed for all the packages being
benchmarked as well as for gene expression of the TF being evaluated. Note that because each
TF has different transduction efficacy and not all cells are able to take up the virus or express
the TF, the theoretical maximum of AUROC will not reach 1.
539

 Benchmarking using AR antagonist data

For the drug treatment dataset, we treated 4 prostate cancer cell lines with 3 different
therapeutic agents and obtained paired gene expression and chromatin accessibility
information. Only the 2 AR-dependent cell lines, LNCaP and VCaP, were included in the
benchmark since they are known to respond to AR-targeting agents. Similarly, only AR
antagonist, Enzalutamide[1] and AR degrader ARV-110[3] were used in the benchmark studies
since they are known to specifically inhibit AR activity. Cells treated with Enzalutamide and
ARV110 are supposed to show reduced AR activity compared to cells treated with DMSO
control. Each cell line was analyzed separately by retaining only peaks found in each cell line
and all the subsequent analyses were performed separately for both cell lines. Each cell is
identified by the HTO tag corresponding to the well receiving DMSO or one of the two AR
inhibitors and this information serves as the ground truth. For each of the two drugs
(enzalutamide or ARV110), a cell is classified into either treated with DMSO control or an AR
inhibitor. AUROC is computed for all the packages being benchmarked as well as for AR gene
expression. Because TF activity is decoupled from TF expression, we used the Wilcoxon
method to estimate weights with merged public AR ChIP-seq prefiltered for the presence of
motifs as the TF occupancy prior.

We used 5 GRN inference tools (FigR, SCENIC+, *GRaNIE* and *Pando)* to benchmark the
performance of Epiregulon. To ensure consistency, we applied the same gene expression and
chromatin accessibility matrices across all tools. We first used ArchR to determine the peaks i.e.
DNA regions with frequent Tn5 transposase insertion events from the fragment files output by
CellRanger. Then the peak x cell matrix was produced by counting the number of insertions per
peak per cell. From the same ArchR projects we also retrieved normalized gene expression to
be used by Epiregulon, GRaNIE, FigR. For the remaining tools, we used a subset of cells in the
gene expression data which matched the cells in the ArchR projects to make sure that all tools
work on the same cells. Below we describe the workflows used in each tool. We followed the
tutorial examples provided at official web sites with only minor changes to the default settings.

FigR

We performed benchmarking against FigR [8]. In the first step of GRN construction the correlation
between peak accessibility and target gene expression was determined using
runGenePeakcorr. We used a non-default search range around TSS (250 kb) to be consistent
across all the benchmarked tools. Correlation coefficients were also computed with background
peaks and the significance of gene-peak association was determined with one-tailed z-test.
Only gene-peak associations that show positive correlation and are statistically significant were
retained (p ≤ 0.05). DORaCs (High density domains of regulatory chromatin) correspond to
genes with ≥7 associated peaks and were calculated by summing the ATAC-seq reads in peaks
matched to each gene. Each TF is associated with a single motif by selecting the motifs most
highly correlated with other motifs of the same TF. Each DORC is evaluated for enrichment of
TFs by comparing the match frequency in its peakset vs the match frequency in a set of
background peaks; the p-values of the enrichment were obtained by z-test ($P_{Enrichment}$).
Smoothing of gene expression matrix and chromatin accessibility data summarized across
target genes was performed after determination of k nearest neighbor cells based on LSI space

retrieved from ArchR project and constructed using ATAC-seq data. The Spearman correlations
between the smoothed DORC accessibility and smoothed TF gene expression were computed
($Correlation$) and their significance were obtained using z-test ($P_{Correlation}$). From the output of
the main function (getFigRGRN) we retrieved scores indicating the strength of association
between transcription factors and target genes which were computed as below:
589

$$Regulation\ score\ =\ sign(Correlation)\ *\ -log10[1 - (1 - P_{Enrichment}) * (1 - P_{Correlation})]$$

590
591 The regulation scores were used as weights when calculating activity with Epiregulon.
592
593
SCENIC+
We performed benchmarking against SCENIC+ [11]. Briefly, we used pycisTopic which uses
Latent Dirichlet Allocation to group regulatory elements into the topics. The model evaluation
indicated 20, 25 and 20 as the number of topics for VCaP cells from AR dataset, LNCaP cells
from AR dataset, and Reprogram-Seq dataset respectively. The input peak matrix was retrieved
from the AchR project. We then used pycistarget to identify TF-region links. Pycistarget
identifies motif matches in the peak regions using HOMER, scores each region for motif
importance and identifies differentially enriched motifs above background regions. Only regions
showing NES > 3.0 and motifs with adjusted p-value <0.05 and log2FC >0.5 were retained. TF-
gene importance scores were calculated using gradient-boosting machine regression by
predicting TF expression from target gene expression. Region-gene importance scores were
calculated using gradient-boosting machine regression by predicting target gene expression
from region accessibility. Genes were ranked by TF-gene importance scores and only genes in
the leading edge of the gene set enrichment were used for the eRegulon. Gene set enrichment
was also performed for region-gene pairs. Peaks were ranked by imputed chromatin
accessibility and genes were ranked by gene expression counts per cell. Enrichment score was
defined as the AUC at 5% of the ranking and was calculated using AUCell. The enrichment
score was used as the activity score in the benchmark assessment.
612
GRaNIE
We performed benchmarking against GRaNIE [10]. Briefly, GRaNIE overlapped TF binding sites
obtained from HOCOMOCO-based TF motifs with ATAC-seq peaks. GRaNIE identified TF-peak
connections using Pearson correlation between TF expression and the peak accessibility across
samples. The cutoffs for the correlation were chosen based on an empirical FDR calculated
from the ratio of TF-peaks in the background peaks over the total number of TF-peaks in both
the background and foreground. Peak-gene connections were identified using correlation
between the gene expression and chromatin accessibility. All the GRN edges have a weight of
1. The default threshold is FDR < 0.2 for TF-peak links and FDR < 0.1 for peak-gene links.
Activities were computed using Epiregulon's calculateActivity.
623
Pando
We performed benchmarking against Pando[9]. Briefly, ATAC-seq peaks were intersected with
PhastCons conserved elements and cCREs derived from ENCODE. TFs present in the 4000

most variable genes were included in the downstream analysis. Motifs for the TFs were
obtained from JASPAR2020 and CISBP. Gene expression and chromatin accessibility counts
were smoothened by averaging cells within a neighborhood. Pando then models target gene
expression as the weighted sum of the product of TF expression and the chromatin accessibility
of the region where the TF binds to.
632

$$Y_g = \sum_t \beta_{t,g,r} X_t A_r + c$$

633   where Y_g is the expression of the target gene g, X_t is the expression of the transcription factor
634   t, A_r is the chromatin accessibility at region r, β is the fitted coefficient and c is the intersection.
635
636   Fitted coefficients were tested for significance using analysis of variance (ANOVA). Only edges
637   with FDR <0.05 were retained in the final GRN. Activities were calculated using Epiregulon's
638   calculateActivity function with the fitted coefficients $\beta_{t,g,r}$ as the weights.
639
640   *Benchmarking using precomputed regulons*
641   Gene expression and chromatin accessibility matrices were obtained from the unpaired PBMC
642   data[39] and integrated using ArchR's addGeneIntegrationMatrix function. Regulons were
643   constructed using Epiregulon and benchmarked against regulons obtained from Dorothea[46])
644   and Humanbase (hematopoiesis)[48].
645
These regulons were applied onto a single-cell RNA-seq dataset involving a CRISPRa screen
that overexpressed transcription factors using gRNA in K562 cells[45]. The identity of the gRNA
was used as the ground truth indicating which TF was activated in each cell. Regulon weights
were computed in the K562 dataset, using the correlation method with gRNA as the cluster
labels for all three regulons. ROC was computed using TF activity as the estimates for cluster
labels indicating whether a particular TF is active or not. TF gene expression was also used as
the estimates to generate a separate ROC.
653
654   **DNN training**
655   Steps for DNN-based TF-RE mapping involved dataset processing, model training, motif scoring
656   and thresholded GRN construction.
657
Dataset processing
The reference genome hg38 and the pseudo-bulk ATAC (taken from ArchR) of individual
clusters (C1 or C3) was collected. The whole genome excluding unmappable regions[52] was split
into 3072 bp regions. Using the coordinates of each region the corresponding DNA sequence
and bigwig coverage was obtained. The DNA sequence was converted to one-hot encoded
form. The DNA and coverage information were saved in tf records files for faster I/O during
training. Training, validation and test splits were done by chromosomes taking chr8 as test set
and chr9 as validation set.
666
667   *Model training*

668 We used Basenji2 architecture with 32 base-pair resolution (binned coverage at 32 bp). The
669 model was trained by randomly sampling 2048bp segment from the input DNA and taking the
670 corresponding coverage (one cluster per model). We used Poisson NLL as loss, used reverse
671 complement augmentation and trained for a maximum of 50 epochs with early stopping.
672
673 We additionally trained models with base-pair resolution using the ChromBPNet architecture.
674 We first trained a custom Tn5 sequence bias model for this dataset on non-peak regions. We
675 ensured that these models learned Tn5 bias by applying DeepSHAP to the model outputs.
676 Using this bias model to regress out Tn5 sequence bias, we then trained ChromBPNet TF
677 models. These models were trained on 2114bp sequences centered on ATAC-seq peaks.
678
Motif scoring
For each of the motif regions (taken from ArchR motif positions file) we obtained the DNA
sequence (denoted as wild type) by centering at the motif and extending to 2048bp. We
occluded the motif by replacing motif nucleotides with 'N' (or 0.25 in one-hot encoding) (denoted
as mutant). We computed the score - mean fraction change in predicted coverage by
subtracting mutant from wild type and normalizing by predicted wild type coverage.
685
686 *Thresholded GRN construction*
687 We computed several quartiles of scores as thresholds. We filtered for those motifs with
688 importance scores higher than the threshold corresponding to bigger differences in prediction.
689 Using the RE remaining we constructed the GRN as before.
690
Reprogram-Seq
Construction of Lentivirus Plasmids for TF Over-expression
All lentivirus plasmids were generated by GenScript. Briefly, the ORF of transcriptional factors
were codon optimized, synthesized and cloned after the hEF1a promoter. The puromycin
resistance gene is driven by a separate Cbh promoter to enable antibiotics selection. Maxi-prep
of each plasmid was performed to maximize the transfection efficiency.
697
Cell Culture and Virus Packaging
LNCaP Clone FGC cells were obtained from ATCC and cultured with RPMI-1640 media with
10% FBS and 2mM L-Glutamine. The cells were split every 4-5 days to maintain the appropriate
density. 293T cells were cultured in DMEM with 10% FBS, 100μM NEAA, 2mM Glutamine. They
were split every 2-3 days. One day before transfection, the cell culture dish was treated with
5mL 1% gelatin in PBS, incubated for 10 minutes, then aspirated. $3.5X10^6$ 293T cells were
seeded into each 10 cm dish to reach ~80% confluence. On the day of transfection, 20 μl
Lipofectamine 2000 was added to 480 μl OptiMEM. In a new tube, the plasmid was mixed in
500μl Opti-MEM at the following ratio: carrier plasmid, 5 μg; delta8.9, 16 μg; VSVG, 1 μg. Both
mixes were combined and incubated for 20 minutes before adding to the dish. The dish was
incubated at 37°C for 6 hours and then 6 ml of complete media was added. Virus was then
harvested 48 hours after transfection. Briefly, all supernatants were harvested and filtered
through a 0.45 μm filter bottle. The virus was concentrated by using the Lenti-X concentrator
(TAKARA Bio.) following the manufacturer's instructions and resuspended in 1 ml 1% BSA in

PBS per dish. The concentrated virus was divided into 200µl aliquots and stored in -80˚C until
infection.

Infection of LNCaP Cells by Lentivirus
Two days before infection, $4X10^5$ LNCaP cells were seeded into each well of a 6-well plate. On
the day of infection, aspirate the media and change to 0.5mL RPMI + 10% FBS + 1X Glutamax
with 8 µg/ml Polybrene. A total of 200 µl concentrated lentivirus was added to each well to
achieve high MOI, and then the plate was centrifuged at 1800 rpm for 45 minutes at room
temperature. After that the plate was put into the incubator for another 3 hours before 2 ml of
the full media was added to each well. Two days after infection, the media was refreshed with
1µg / ml puromycin. The cells were then grown for another 7 days before harvesting for the
single-cell analysis, during which the cells were split accordingly when the confluence reached
100%.

Engineering of dox-inducible NKX2-1 expressing cell lines
LNCAP Clone FGC cells were passaged two days before electroporation. 200k cells per
reaction were transduced with 375 ng PBIND-NKX2-1-IRES-eGFP-PGK-puro and 125ng pBO
(PiggyBac transposase) (3:1 vector:transposase ratio) according to manufacturer's instructions
for SF Cell Line 4D-Nucleofector™ X Kit S, using the EN-120 electroporation protocol in 20 µl
Nucleocuvette™ Strip. After electroporation, 100ul of warm media (RPMI + 10% FBS + 1X
Glutamax) was added to the cuvette and the cuvette was placed in the incubator for 1 hour to
recover. Cells were then transferred into a 6 well plate. Puromycin selection (RPMI + 10% FBS
+ 1X Glutamax, 1ug/ml Puromycin) was performed after 4-5 days, depending on cell recovery.

Drug treatment with AR inhibitors and SMARCA2/4 degrader
LNCap Clone FGC, VCAP and DU145 cells were obtained from ATCC and cultured with RPMI-
1640 media with 10% FBS and 2mM L-Glutamine; NCI-H660 cells were obtained from ATCC
and cultured with DMEM/F12 with 0.005 mg/ml insulin,0.01 mg/ml Transferrin, 30nM Sodium
selenite,10nM Hydrocortisone,10nM beta-estradiol, 4mM L-Glutamine. The cells were split
every 4-5 days to maintain the appropriate density. For the drug response assay, cells were
seeded in 6 well format. The cells were split every 4-5 days to maintain the appropriate density.

$1.5x10^5$ DU145 cells were seeded into each well of a 6 well plate and 5E3 DU145 cells were
seeded into each well of a 96 well plate; 1.5-2E6 LNCAP or VCAP cells were seeded into T75
flasks; $5x10^5$ cells NCI-H660 cells were seeded into each well of a 6 well plate. After 48 hours,
the media was removed and cells were treated with DMSO, 1uM Enzalutamide, 0.1 uM
SMARCA2.4_1 or 1 uM ARV-110 in complete media. After 24 hours from treatment, the cells
were harvested for the single-cell RNA-ATAC Co-assay

Single-cell RNA-seq and Single-cell RNA-ATAC Co-assay
The single-cell RNA-seq was performed using the Chromium Single Cell 3' kit (V3.1) from 10X
Genomics with cell hashing. Briefly, the cells were trypsinized into single-cell suspension from
the 6-well plates, and washed once with PBS with 1% BSA. The cells from different wells were
stained with human TotalSeq-A cell hashing antibodies (BioLegend) containing distinct

barcodes following the manufacturer's protocol. The cells were then washed twice with PBS with 1% BSA before combining and the live cells were sorted using the SONY SH800S FACS machine. For loading of the 10X chip G, we overloaded the channel with the aim of recovering 20K cells. The library construction was following the standard 10X protocol with the following exceptions: 1) the 1µl of 10µM HTO additive primer was spiked in during the cDNA PCR step; 2) during the SPRI beads cleanup step, the supernatant from the 0.6X cleanup was saved to recover the HTO fragment. Along with the transcriptome library, the HTO library was amplified from the supernatant using the HTO-specific primer. The detailed protocol can be found on the cell hashing website (https://cite-seq.com/cell-hashing/).

The single-cell RNA-ATAC co-assay was performed using the 10X Genomics Single Cell Multiome ATAC + Gene Expression kit. In a previous test, we found that the standard cell permeabilization protocol used for nuclei preparation does not completely remove the plasma membrane of the LNCap cells and robust B2M expression can still be detected with FACS (data not shown). Therefore, we speculated that the same cell hashing strategy would work for the 10X Multiome assay. Briefly, we followed the same hashing protocol as used in the scRNA-seq experiment. After the antibody staining and pooling, the cells were permeabilized by following the nuclei isolation protocol from 10X genomics (10X Nuclei Isolation Protocol). The cells were then counted and loaded on the Chip J with a recovery aim of 12K cells per lane. For the library construction, we followed the standard protocol with the following modifications: 1) 1µl of 10µM HTO additive primer was spiked in during the pre-AMP and cDNA PCR steps; 2) during the 0.6X cDNA cleanup, the supernatant was saved to amplify the HTO library, as described in the previous section.

All the libraries were sequenced on Illumina NextSeq 2000 or NovaSeq 4000. We were aiming for 20K, 40K and 2K raw reads per cell for the transcriptome, ATAC and HTO libraries, respectively.

Synthesis of SMARCA2.4_1

Scheme 1. Preparation of intermediate **E**.

tert-Butyl   4-((trans)-3-(2-(benzyloxy)ethoxy)cyclobutoxy)piperidine-1-carboxylate.   (**C**). To a solution of *tert*-butyl 4-((trans)-3-hydroxycyclobutoxy)piperidine-1-carboxylate (**A**, 500 mg, 1.84 mmol, 1.0 eq) in THF (7 mL) was added NaH (60% in mineral oil, 111 mg, 2.77 mmol, 1.5 eq) at 0 °C. The resulting solution was stirred at 0 °C for 0.5 h afterwhich ((2-bromoethoxy)methyl)benzene (**B**, 592 mg, 2.76 mmol, 1.5 eq) was added. The mixture was warmed to room temperature and was stirred for an additional 16 h then was quenched with saturated aqueous NH$_4$Cl (5 mL). The resulting solution was extracted with EtOAc (10 mL x 3),

and the combined organic layers were dried over sodium sulfate and concentrated under vacuum. The residue was purified by silica gel chromatography (EtOAc:petroleum ether, 0:100 to 24:76) to afford the title compound (330 mg, 44%) as a colorless oil. [1]H NMR (300 MHz, DMSO-d_6) δ 7.42-7.23 (m, 5H), 4.49 (s, 2H), 4.25-4.17 (m, 1H), 4.09-4.02 (m, 1H), 3.68-3.61 (m, 2H), 3.57-3.51 (m, 2H), 3.47-3.37 (m, 2H), 2.95 (t, J = 11.0 Hz, 2H), 2.23-1.98 (m, 4H), 1.79-1.67 (m, 2H), 1.39 (s, 9H), 1.35-1.17 (m, 3H). LC-MS: (ES+): m/z 406 [M+H]$^+$.

tert-Butyl 4-((trans)-3-(2-hydroxyethoxy)cyclobutoxy)piperidine-1-carboxylate. (**D**). A solution of intermediate **C** (330 mg, 0.815 mmol) and 10% Pd/C (66.0 mg) in ethanol (10 mL) and THF (10 mL) was stirred at room temperature for 1 h under a hydrogen atmosphere (3 atm). The resulting mixture was filtered, and the filter cake was washed with DCM (10 mL x 2). The filtrate was concentrated under reduced pressure to afford the title compound (245 mg, 95% yield) as a white oil. [1]H NMR (300 MHz, DMSO-d_6) δ 4.58 (t, J = 5.5 Hz, 1H), 4.26-4.18 (m, 1H), 4.08-4.01 (m, 1H), 3.69-3.61 (m, 2H), 3.53-3.34 (m, 3H), 3.28 (t, J = 5.3 Hz, 2H), 2.96 (t, J = 11.4 Hz, 2H), 2.24-1.98 (m, 4H), 1.76-1.71 (m, 2H), 1.39 (s, 9H), 1.36-1.18 (m, 2H). LC-MS: (ES+): m/z 316 [M+H]$^+$.

*tert-Butyl 4-((1r,3r)-3-(2-(tosyloxy)ethoxy)cyclobutoxy)piperidine-1-carboxylate (**E**).* To a solution of intermediate **D** (245 mg, 0.78 mmol, 1.0 eq) and pyridine (245 mg, 3.11 mmol, 4.0 eq) in DCM (2 mL) was added TsCl (223 mg, 1.16 mmol, 1.5 eq) at 0 °C. The mixture was then warmed to room temperature and stirred for 12 h. The solvent was subsequently removed under vacuum, and the residue was purified by silica gel chromatography (EtOAc:petroleum ether, 0:100 to 29:71) to afford the title compound (303 mg, 83%) as a colorless oil. [1]H NMR (300 MHz, DMSO-d_6) δ 7.82-7.72 (m, 2H), 7.52-7.41 (m, 2H), 4.22-4.05 (m, 3H), 4.01-3.88 (m, 1H), 3.67-3.59 (m, 2H), 3.45-3.34 (m, 3H), 2.94 (t, J = 11.4 Hz, 2H), 2.41 (s, 3H), 2.13-1.92 (m, 4H), 1.76-1.64 (m, 2H), 1.37 (s, 9H), 1.33-1.15 (m, 2H). LC-MS: (ES+): m/z 470 [M+H]$^+$.

Scheme 2. Preparation of SMARCA2.4_1 (compound **1)**.

6-Chloro-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridazin-3-amine **(H)**.   A
solution of 4-bromo-6-chloropyridazin-3-amine (**F**, 2.00 g, 9.60 mmol, 1.0 eq), (1-(tetrahydro-2H-
pyran-2-yl)-1H-pyrazol-4-yl)boronic acid (**G**, 2.46 g, 12.5 mmol, 1.3 eq), Pd(PPh$_3$)$_4$ (1.12 g,
0.969 mmol, 0.1 eq) and K$_2$CO$_3$ (2.67 g, 19.3 mmol, 2.0 eq) in 1,4-dioxane (20 mL) and water (2
mL) was stirred at 100 °C for 2 h. The reaction was then cooled to room temperature and was
quenched with water (20 mL). The resulting solution was extracted with EtOAc (30 mL x 3), and
the combined organic layers were dried over sodium sulfate and concentrated under vacuum.
The residue was purified by silica gel chromatography (EtOAc:petroleum ether, 0:100 to 58:42)
to afford the title compound (2.40 g, 89%) as a red oil. ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.52 (d,
J = 0.8 Hz, 1H), 8.11 (d, *J* = 0.8 Hz, 1H), 7.65 (s, 1H), 6.41 (s, 2H), 5.46 (dd, *J* = 9.9, 2.2 Hz,
1H), 4.07-3.90 (m, 1H), 3.76-3.57 (m, 1H), 2.24-2.05 (m, 1H), 2.00-1.91 (m, 2H), 1.79-1.64 (m,
1H), 1.60-1.53 (m, 2H). LC-MS: (ES+): m/z 280 [M+H]$^+$.

2-(6-Amino-5-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridazin-3-yl)phenol **(J)**.   A
solution of intermediate **H** (1.35 g, 4.83 mmol, 1.0 eq), (2-hydroxyphenyl)boronic acid (**I**, 870
mg, 6.31 mmol, 1.3 eq), XphosPdG3 (410 mg, 0.484 mmol, 0.1 eq) and K$_2$CO$_3$ (1.33 g, 9.64
mmol, 2.0 eq) in 1,4-dioxane (11.5 mL) and water (2.5 mL) was stirred at 95 °C for 5 h. The
mixture was cooled to room temperature, and the solvent was removed under vacuum. The
residue was purified by silica gel chromatography (MeOH:DCM, 0:100 to 5:95) to afford the title
compound (900 mg, 55%) as a yellow solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.77 (s, 1H),
8.60 (d, *J* = 0.8 Hz, 1H), 8.26-8.17 (m, 2H), 8.02 (dd, *J* = 8.3, 1.6 Hz, 1H), 7.28-7.23 (m, 1H),
6.97-6.85 (m, 2H), 6.51 (s, 2H), 5.48 (dd, *J* = 9.9, 2.2 Hz, 1H), 3.99-3.90 (m, 1H), 3.74-3.60 (m,
1H), 2.25-2.06 (m, 1H), 2.00-1.95 (m, 2H), 1.76-1.64 (m, 1H), 1.59-1.52 (m, 2H). LC-MS: (ES+):
m/z 338 [M+H]$^+$.

2-(6-Amino-5-(1H-pyrazol-4-yl)pyridazin-3-yl)phenol **(K)**.   A solution of intermediate **J**
(300 mg, 0.890 mmol) in HCl (4.0 M in MeOH, 1 mL) was stirred at room temperature for 2 h.
The solvent was then removed under vacuum, and the residue was purified by pre-packed C18
column (solvent gradient: 0-100% MeOH in water (0.1% TFA)) to afford the title compound (188
mg, 83%) as a brown solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.85 (s, 1H), 13.42 (s, 1H), 8.33
(s, 2H), 8.22 (s, 1H), 8.02 (dd, *J* = 8.3, 1.6 Hz, 1H), 7.30-7.24 (m, 1H), 6.95-6.90 (m, 2H), 6.50
(s, 2H). LC-MS: (ES+): m/z 254 [M+H]$^+$.

tert-Butyl   4-((trans)-3-(2-(4-(3-amino-6-(2-hydroxyphenyl)pyridazin-4-yl)-1H-pyrazol-1-
yl)ethoxy)cyclobutoxy)piperidine-1-carboxylate **(L)**. A solution of intermediate **K** (100 mg, 0.395
mmol, 1 eq), intermediate **E** (223 mg, 0.475 mmol, 1.2 eq) and K$_2$CO$_3$ (136 mg, 0.985 mmol, 2.5
eq) in DMF (1.3 mL) was stirred at 80 °C for 2 h. The reaction mixture was cooled to room
temperature, then was diluted with EtOAc (5 mL) and washed with brine (3 mL x 5). The
organic layer was dried over sodium sulfate and was concentrated under vacuum. The residue
was purified by silica gel chromatography (MeOH:DCM, 0:100 to 5:95) to afford the title
compound (60 mg, 27%) as a brown solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.79 (s, 1H), 8.45
(d, *J* = 0.8 Hz, 1H), 8.24-8.17 (m, 2H), 8.03-7.97 (m, 1H), 7.25 (t, *J* = 7.7 Hz, 1H), 6.97-6.87 (m,
2H), 6.48 (s, 2H), 4.30 (t, *J* = 5.4 Hz, 2H), 4.15-4.06 (m, 1H), 4.06-4.00 (m, 1H), 3.69 (t, *J* = 5.4
Hz, 2H), 3.61-3.53 (m, 2H), 3.24-3.13 (m, 1H), 2.84 (t, *J* = 11.3 Hz, 2H), 2.05-2.01 (m, 4H),
1.66-1.60 (m, 2H), 1.35 (s, 9H), 1.24-1.17 (m, 2H). LC-MS: (ES+): m/z 551 [M+H]$^+$.

2-(6-Amino-5-(1-(2-(trans)-3-(piperidin-4-yloxy)cyclobutoxy)ethyl)-1H-pyrazol-4-
*yl)pyridazin-3-yl)phenol, TFA salt (**M**).* A solution of intermediate **L** (60 mg, 0.11 mmol) in TFA
(0.3 mL) and DCM (0.3 mL) was stirred at room temperature for 0.5 h.　The solvent was
removed under vacuum to afford the title compound (66 mg, crude) as a yellow solid. ^{1}H NMR
(300 MHz, DMSO-d_6) δ 8.46 (s, 1H), 8.26 (s, 1H), 8.14 (s, 1H), 7.71-7.68 (m, 1H), 7.39-7.33 (m,
1H), 7.03-6.91 (m, 2H), 4.32 (t, J = 5.4 Hz, 2H), 4.20-4.00 (m, 2H), 3.70 (t, J = 5.4 Hz, 2H), 3.53-
3.45 (m, 1H), 3.12-3.08 (m, 2H), 2.90-2.84 (m, 2H), 2.09-2.05 (m, 4H), 1.91-1.78 (m, 2H), 1.59-
1.47 (m, 2H). LC-MS: (ES+): m/z 451 [M+H]$^{+}$.

(2S,4R)-1-((R)-2-(3-(2-(4-((trans)-3-(2-(4-(3-amino-6-(2-hydroxyphenyl)pyridazin-4-yl)-1H-
pyrazol-1-yl)ethoxy)cyclobutoxy)piperidin-1-yl)ethoxy)isoxazol-5-yl)-3-methylbutanoyl)-4-
*hydroxy-N-((S)-1-(4-(4-methylthiazol-5-yl)phenyl)ethyl)pyrrolidine-2-carboxamide (**1**).* A solution
of intermediate **M** (66 mg, crude), intermediate **N** (prepared as described in Berlin, M.; et al.
"PROTACs Targeting BRM (SMARCA2) Afford Selective In Vivo Degradation Over BRG1
(SMARCA4) and are Active in BRG1-Mutant Xenograft Tumor Models", *J. Med. Chem*.
accepted for publication; 85 mg, 0.157 mmol) and NaOAc (54 mg, 0.658 mmol) in MeOH (0.8
mL) was stirred at room temperature for 1 h.　Then NaBH$_3$CN (21.0 mg, 0.33 mmol) was added,
and the mixture was stirred at room temperature for an additional 0.5 h.　The solvent was
removed under vacuum, and the residue was purified by Prep-HPLC [column: Xselect CSH F-
Phenyl OBD column, 30*150 mm, 5 μm; mobile Phase A: water (10 mmol/L NH$_4$HCO$_3$), mobile
phase B: CH$_3$CN; flow rate: 60 mL/min; gradient: 43% to 63% B over 8 min; wavelength:
254/220nm; RT: 8.97 min] to afford the title compound (12.8 mg, 12% over two steps) as a
white solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.83 (s, 1H), 9.00-8.99 (m, 1H), 8.51-8.46 (m,
1H), 8.43 (d, J = 7.6 Hz, 1H), 8.26-8.19 (m, 2H), 8.05-8.01 (m, 1H), 7.47-7.34 (m, 4H), 7.29-7.24
(m, 1H), 6.98-6.87 (m, 2H), 6.50 (s, 2H), 6.09-5.92 (m, 1H), 5.11-5.02 (m, 1H), 4.96-4.86 (m,
1H), 4.44-4.24 (m, 4H), 4.15 (t, J = 5.6 Hz, 2H), 4.11-4.00 (m, 2H), 3.74-3.51 (m, 4H), 3.46-3.43
(m, 1H), 3.14-3.08 (m, 1H), 2.66-2.63 (m, 2H), 2.57-2.53 (m, 2H), 2.46 (s, 3H), 2.31-2.16 (m,
1H), 2.07-1.94 (m, 7H), 1.82-1.74 (m, 1H), 1.67-1.63 (m, 2H), 1.46-1.37 (m, 3H), 1.34-1.20 (m,
2H), 0.98 -0.95 (m, 3H), 0.84-0.79 (m, 3H).   LC-MS: (ES+): m/z 975 [M+H]$^{+}$.
896

Measurements of SMARCA2/4 degradation using In-cell Western

22Rv1 cells were seeded at 8000/well in 96-well black/clear-bottom plates with 180ul DMEM
containing 1% pen-strep, 1% HEPES and 10% FBS and incubated overnight at 37□°C to allow
adherence. The next morning cells were treated with 20□uL of 10× compound and incubated for
an additional 20□h. Cells were then washed with ice cold DPBS, then 50□uL of ice cold 4%
PFA (Electron Microscopy Sciences 15711)/DPBS was added, and plates were then incubated
at RT for 20□min. PFA was then removed and 200□uL of TBS-T containing 0.5% Triton X 100
(Sigma T8787) was added. Plates were incubated at RT for 30□min. TBS-T/Triton X was then
removed, 50□uL of Li-cor blocking solution (Li-Cor 927-50003) was added, and plates were
incubated at RT for 1□h. Blocking solution was removed, 50□uL of Li-Cor blocking solution
containing primary antibody to SMARCA2 (1:2000) and α-Tubulin (1:2000) or SMARCA4
(1:1000) and α-Tubulin (1:2000) was added, and plates were incubated at 4□°C overnight. The
next day, plates were washed 3× with TBS-T and 50uL of secondary antibody cocktail in Li-Cor
blocking solution was added (IRDye 800CW Goat anti-rabbit IgG, Li-Cor 926-32211 and IRDye
680RD Goat anti-mouse IgG, Li-Cor 926-68070; both secondary antibodies are diluted 1:5000).

Plates were incubated at RT for 1□h protected from light. Then plates were washed twice with
TBS-T and excess liquid removed. Plates were read on the Li-Cor Odyssey with default
intensity setting of 5.0 for both channels. Li-Cor images were analyzed with the in-cell Western
feature of Image Studio Lite. Assays were run with technical duplicates and multiple biological
replicates (>2) with error calculated with GraphPad prism using 95% confidence interval.
917

Acknowledgments
We would like to thank Xinpeng Wang, Fang Zhang, Mingtao He, Yunxing Cheng and Michael
Berlin (Arvinas) for the synthesis of SMARCA2.4_1 and Peter Dragovich for guiding the
synthesis. We would like to thank Jayaram Kancherla for his help and advice.

Figure legends
Figure 1. Figure constructs GRNs to infer TF activity at the single cell level
Schematics of Epiregulon. RNA and ATAC counts of similar cells are first aggregated. Based on
the correlation between ATAC and RNA, each target gene (TG) is assigned a regulatory
element (RE) within a defined distance (default ±250kb). Each RE is interrogated for possible
TF occupancy using ChIP-seq data from ENCODE and ChIP-Atlas. An optional network pruning
step retains TF-RE-TG triples that co-occur within the same cells. Co-occurrence is defined by
TF and TG expression and chromatin accessibility exceeding a threshold. The regulatory
potential of TF for each TG is estimated using 1 of the 3 methods (correlation, mutual
information and Wilcoxon) and this serves as the weight of each TF-TG pair. The weights are
computed for all the cells and/or for cells in each cluster to yield cluster-specific weights. The TF
activity is computed as the weighted sum of the target gene expression normalized by the
number of genes with non-zero edges. The regulon of each TF can be annotated with gene set
enrichment. Jaccard similarity between the target genes is used to identify transcription factors
that potentially interact. Differential network is performed on edge-subtracted networks of two
cell states.
939

Figure 2: Epiregulon identifies cell fate changes
a) UMAP representation of 3,903 LNCaP cells transduced with virus encoding GATA6 and
NKX2-1. The cells were infected in individual wells, hashtagged with HTO and then
pooled into a single run. Shown is the level of TF gene expression, TF activity computed
by Epiregulon for GATA6 and NKX2-1 or hash assignment.
b) Each cell was identified by the HTO tag corresponding to the well receiving virus
encoding a particular TF and this information serves as the ground truth. For each TF, a
cell was classified into either expressing this TF or not based on the TF activity. AUROC
was computed for all the packages being benchmarked as well as for gene expression
of the TF being evaluated. Note that because each TF had different transduction efficacy
and not all cells were able to take up the virus or express the TF, the theoretical
maximum of AUROC will not reach 1.
c) Number of TFs retrieved from the different packages being benchmarked
d) As a validation, LNCaP cells were transduced with dox-inducible constructs expressing
either NKX2-1 or eGFP control. Correlation between weights of NKX2-1 targets

(computed by the Wilcoxon method) against logFC of NKX2-1 targets between no dox
and dox treatments.
e) Genes most similar to GATA6 were ranked by Jaccard similarity between the GATA6
regulon and regulons of all the other TFs.
f) Chromatin accessibility at neuroendocrine (NE) - and stem cell like (SCL)- specific
regions defined by Tang et al was computed by ChromVAR
g) Shown are the Pearson's correlation between GATA6 activity and chromatin
accessibility at regions specific to stem cell like (SCL) subtype of prostate cancer, and
the Pearson's correlation between NKX2-1 activity and chromatin accessibility at
neuroendocrine (NE) subtype of prostate cancer.
966
Figure 3: Epiregulon predicts responses towards TF-targeting drugs
a) UMAP representation of scMultiome data of 4 prostate cancer cell lines treated with 3
drug compounds (5,035 VCaP cells, 5,935 LNCaP cells, 3,961 NCI-H660 cells and
3,639 DU145 cells). VCaP and LNCaP are AR-dependent whereas DU145 and H660
are AR-independent. The 3 drugs include an AR antagonist (enzalutamide), an AR
degrader (ARV110) and a degrader of SMARCA2/4, the ATPase subunit of the
SWI/SNF remodeling complex (SMARCA2.4_1). Enzalutamide and ARV110 are known
and specific inhibitors of AR activity and were used for benchmarking. Cells were
merged from 2 technical replicates.
b) AR activity computed by Epiregulon.
c) Each cell was identified by the HTO tag corresponding to the well receiving DMSO or
one of the two AR inhibitors and this information served as the ground truth. For each of
the two drugs (enzalutamide or ARV110), a cell was classified into either treated with
DMSO control or an AR inhibitor based on AR activity. AUROC was computed for all the
packages being benchmarked as well as for AR gene expression.
d) AR expression of LNCaP and VCaP cells treated with the 3 drugs.
e) Potential interaction partners of AR were nominated using Jaccard similarity between its
target genes and the target genes of all other transcription factors in differential networks
between DMSO-treated or Enzalutamide/ARV110-treated cells.
f) Chronos score of GATA2 and SUMO2 in all prostate cancer cells screened by DepMap.
987
988
989
Figure 4: Epiregulon predicts responses towards chromatin-modifying drugs
a) SMARCA4 activity computed by Epiregulon
b) Hypergeometric enrichment of gene sets (hallmark and oncogenic signature) of the
target genes of SMARCA4 inferred by Epiregulon for each of the 4 prostate cancer cell
lines. VCaP and LNCaP are AR-dependent and DU145 and H660 are AR-independent.
c) Potential interaction partners of SMARCA4 were nominated using Jaccard similarity
between its target genes and the target genes of all other transcription factors. The top
interacting genes are reflective of the unique etiologies of the different prostate cancer
subtypes.
999

Figure 5: Epiregulon-constructed GRNs predict effects of perturbation

    a) UMAP representation of PBMC data with unpaired and integrated scATAC-seq and scRNA-seq for which Epiregulon was used to compute the GRN.

    b) Representation of TF expression or activity inferred by Epiregulon of well-known lineage factors (EBF1, PAX5 - B cells, GATA3 - T cells and SPI1- myeloid cells). Weights were estimated using the Wilcoxon method.

    c) Cell-type specific GRN weights were computed for PreB and CD4 memory T cells (CD4.M) respectively. The GRN weights are cell type aware since only PreB cells, not CD4 memory T cells, have positive weights in the target genes of the B cell lineage factor, EBF1. Weight cutoff was >0.

    d) EBF1 was interrogated for potential interaction partners using Jaccard similarity between its target genes and the target genes of all other transcription factors. The top genes (PAX5, IRF4 and TCF3 (also known as E2A)) are known to interact with EBF1 in B cell development.

    e) Differential networks between monocytes and CD4 memory T cells were computed by subtracting the edges of the cell type specific GRNs. Genes were ranked by degree centrality of the edge-subtracted graph. SPI1 and CEBPA are well known lineage factors of the myeloid lineage whereas TBX21 and GATA3 are well known lineage factors of the T cell lineage.

    f) The GRN computed from the PBMC data was applied onto a CRISPRa dataset of K562 cells containing only gene expression data (no scATAC-seq). Each cell was identified by the guide RNA corresponding to a TF that was being activated and this information serves as the ground truth. For each TF, a cell was classified into containing either the TF activating guide or the non-targeting control based on its TF activity. Shown is the area under the receiver operating curves (AUROC) using either TF gene expression or TF activity computed from GRN constructed from the PBMC dataset as the classifier. AUROC is also calculated for activity inferred from other GRNs (Dorothea, CollecTRI or humanbase).

    g) TF activity inferred by epiregulon could better classify PRDM1-activated cells from control cells than gene expression or TF activity estimated from the Dorothea regulons. TF activity avoids zero inflation as seen with single cell gene expression.

Figure S1

    a) AUROC of the reprogram-seq dataset was computed with each of the 3 weight methods offered by Epiregulon. Correlation (corr) and mutual information (MI) weights are computed either using 1) transcription factor (TF) expression vs target gene (TG) expression or 2) TF expression multiplied with the chromatin accessibility at regulatory elements (RE) vs TG expression. Each cell is identified by the HTO tag corresponding to the well receiving virus encoding a particular TF and this information serves as the ground truth. For each TF, a cell is classified into either expressing this TF or not.

    b) Differences in rank between target and non-target genes with respect to GATA6 and NKX2-1 transcription factors. Gene rank was created from the product of two ranks: one based on the differences in the mean expression between the treatment and control

1044   groups, and the second based on the proportion of the total gene expression across all
1045   the cells contributed by the treated cells.
1046   c)   Representation of target genes of NKX2-1 and GATA6. C1 cluster contains cells infected
1047   with GATA6, C3 contains cells infected with NKX2-1 and C5 contains uninfected cells.
1048   Edges with weight > 0 were retained and target genes of NKX2-1 and GATA6 were
1049   shown for each cluster.
1050   d)   As a validation, LNCaP cells were transduced with dox-inducible constructs expressing
1051   either NKX2-1 or eGFP control. Shown are the putative target genes of NKX2-1 ranked
1052   by weights.
1053   e)   Representation of target genes of YAP1 and GATA6 with C1-specific weights.
1054   f)   Chromatin accessibility at adenocarcinoma- and neuroendocrine-specific regions
1055   defined by Baca et al was computed by ChromVAR
1056

1057 **Figure S2**
1058   a)   Chemical structure of SMARCA2.4_1, a degrader of SMARCA2 and SMARCA4
1059   b)   Levels of SMARCA2 and SMARCA4 after SMARCA2.4_1 treatment. SMARCA2 DC50 =
1060   0.22nM, SMARCA4 DC50 = 0.31 nM in 22Rv1 cells.
1061   c)   AUROC of the AR dataset was computed with the Wilcoxon method in 4 prostate cancer
1062   cell lines using AR activity as the classifier of treatment condition: DMSO or AR inhibitor
1063   (ARV110 or enzalutamide). The two AR-dependent cell lines (VcaP and LNCaP) are
1064   known to decrease AR activity upon ARV110 or enzalutamide treatment. The two AR-
1065   independent cell lines (DU145 and H660) do not respond to these AR inhibitors.
1066   d)   Same as c) but calculated with all 5 different weight methods in LNCaP.
1067   e)   Number of TFs retrieved by each of the packages
1068   f)   TF activity was computed with either AR ChIP-seq merged across ENCODE and ChIP-
1069   atlas or with LNCaP or VCaP AR and FOXA1 ChIP-seq (cell line-matched). There is a
1070   slight discrepancy between the cell culture condition used to obtain scMultiome data (in
1071   this manuscript) vs the condition used to obtain cell line-matched ChIP-seq. The cell
1072   line-matched ChIP-seq was obtained in cell lines cultured in charcoal stripped medium
1073   supplemented with R1881 whereas the scMultiome data was performed in cell lines
1074   cultured in 10% fetal bovine serum (FBS) but the identities of the cell lines are matched.
1075
1076

1077 **Figure S3**
1078   a)   AUROC of the AR dataset was computed with each of the 3 weight methods offered by
1079   Epiregulon. Correlation (corr) and mutual information (MI) weights are computed either
1080   using transcription factor (TF) expression vs target gene (TG) expression or TF
1081   expression multiplied with the chromatin accessibility at regulatory elements (RE) vs TG
1082   expression. Each cell is identified by the HTO tag corresponding to the well receiving
1083   DMSO or the SMARCA2/4 degrader SMARCA2.4_1 and this information serves as the
1084   ground truth. A cell is classified into either treated with DMSO control or SMARCA2.4_1
1085   based on the SMARCA2 and SMARCA4 activity inferred by Epiregulon.
1086   b)   TF activity was computed with either SMARCA4 ChIP-seq merged across ENCODE and
1087   ChIP-atlas or with VCaP SMARCA4 ChIP-seq (cell line-matched). There is a slight

1088   discrepancy between the cell culture condition used to obtain scMultiome data (in this
1089   manuscript) vs the condition used to obtain cell line-matched ChIP-seq. The cell line-
1090   matched ChIP-seq was obtained in cell lines cultured in charcoal stripped medium
1091   supplemented with R1881 whereas the scMultiome data was performed in cell lines
1092   cultured in 10% fetal bovine serum (FBS) but the identities of the cell lines are matched.
1093
Figure S4
a) Gene set enrichment of regulons inferred by Epiregulon (Wilcoxon weight cutoff > 0).
EBF1 and PAX5 are B cell lineage factors. GATA3 is a lymphoid lineage factor. SPI1 is
a myeloid lineage factor.
b) Shown is the area under the receiver operating curves (AUROC) using either TF gene
expression or TF activity computed from GRN obtained from humanbase, collecTRI or
Dorothea.
1101
Figure S5
a) We used the scMultiome package to simulate paired scRNA-seq and scATAC-seq data
(see Supplementary Note). Briefly, we provided a lineage tree that specified the
identities of the individual cells and a ground truth GRN that specified the regulatory
relationships between TF-TG and RE-TG. Gene expression and peak matrices were
generated taking into account both cell identities and gene regulatory relationships. We
obtained the ground truth activity of each TF by summing the differences in target gene
expression as defined by the GRN. For all assessments from b-e, we use the fixed cutoff
and tf_re.merge set to TRUE.
b) We varied the degree of sparsity by increasing the technical noise and evaluated the
correlation of activity against ground truth TF activity across all genes for each of the
weight methods. Each data point represents a TF. The ATAC effect was set at basic.
GRN had 100% true connections.
c) We added noise to GRN by adding false connections and removing true connections,
and correlated the activity against ground truth activity for each TF. The ATAC effect
was set at basic and data sparsity was set at medium.
d) Cell clusters were first defined using k-means clustering with a median cluster size of 10
cells for the Wilcoxon method. We correlated the activity against ground truth activity for
each TF. The ATAC effect was set at normal and data sparsity was set at medium. GRN
had 100% true connections. n indicates the number of TFs retrieved in the
benchmarking.
e) We varied the effect of ATAC on the gene expression, with the strong ATAC effect
indicating a greater importance of ATAC counts on shaping gene expression counts.
GRN had 100% true connections. True counts were generated without data sparsity and
observed counts were generated with medium data sparsity. n indicates the number of
TFs retrieved in the benchmarking.
f) We spiked in 100% false connections to the true GRN. We examined the recall and
precision of true GRN connections with varying degrees of z score cutoffs computed by
chi-square statistics

g) We spiked in 100% false connections to the true GRN. We show that true connections
have increased RE-TG correlations and this translated to higher weights in the true
connections.
h) We evaluated the effects of pruning on the accuracy of TF activity predictions. We
computed the ground truth TF activity from the predefined GRN with high data sparsity
and strong ATAC effects. We spiked in 100% false connections and compared the
correlation between the TF activity inferred from either the unpruned or pruned GRN vs
the ground truth GRN. Shown is the difference between the two correlations. A positive
difference indicates improved performance due to pruning.
1140

Figure S6
a) We added noise to the network by increasing proportions of randomly permuted edges
on top of true differential edges (15% of total edges). Centrality subtraction was
computed by subtracting the degree centralities of two networks whereas edge
subtraction was computed by centrality of the edge subtracted network. See methods for
details.
b) We varied the signal strength by increasing the proportions of altered edges
(enrichment, depletion or permutations) with noise fixed at 10% of total edges.
1149

Figure S7
a) Each cell was identified by the HTO tag corresponding to the well receiving virus
encoding a particular TF and this information served as the ground truth cell labels. For
each TF, a cell was classified into either expressing this TF or not. GATA6 ChIP-seq
was obtained by merging ChIP-seq from ChIP-atlas and ENCODE. Merged ChIP + motif
refers to merged ChIP-seq peaks that contain GATA6 motifs. We trained a DNN model
on the cluster of cells expressing GATA6 using Basenji2 and predicted an importance
score for each motif based on the difference between the original sequence and motif
occluded sequence. We filtered for those motifs with importance scores higher than
quartiles of scores.
b) Same as a) but computed for NKX2-1 using Nkx2.1.Homeobox_182 as the motif.
c) Same as a) but performed with ChromBPNet
d) Same as b) but performed with ChromBPNet
1163

Figure S8
a) To evaluate whether GRN weights improves performance of TF activity, weights of
target genes were randomly permuted and AUROC is computed with GRNs with
permuted weights
b) Same as a) but performed on NKX2-1
c) Each cell was identified by the HTO tag corresponding to the well receiving DMSO or the
AR antagonist enzalutamide and this information serves as the ground truth cell labels.
Public ChIP-seq was derived from merged AR ChIP-seq from ChIP-atlas and ENCODE.
Motif was derived from the AR motif in CisBP. Merged ChIP-seq + motif refers to merged
public ChIP-seq peaks that contain AR motifs. We trained a DNN model on cells treated
with DMSO using Basenji2 and predicted an importance score for each motif based on

the difference between the original sequence and motif occluded sequence. We filtered
for those motifs with importance scores higher than quartiles of scores.
1177

1178 **Code availability**

1179

Epiregulon: https://github.com/xiaosaiyao/epiregulon
Epiregulon.extra: https://github.com/xiaosaiyao/epiregulon.extra
Epiregulon.archr: https://github.com/xiaosaiyao/epiregulon.archr
ScMultiome: https://github.com/xiaosaiyao/scMultiome
Epiregulon benchmarking: https://github.com/xiaosaiyao/epiregulon.benchmark
Deep neural network model: https://github.com/xiaosaiyao/epiregulon.sequence.modeling
Differential network simulation: https://github.com/xiaosaiyao/epiregulon.diffnetwork.simulation
1187

1188 **Data availability**
1189 GEO (GSE252883)
1190 GSE251977:
1191 https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE251977
1192 token: snczwmievvwnbyl
1193

1194 GSE251978:
1195 https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE251978
1196 token: sdkpsuwutdoblep
1197

1198

1   Tran, C. *et al.* Development of a second-generation antiandrogen for treatment of
advanced prostate cancer. *Science* **324**, 787-790, doi:10.1126/science.1168175 (2009).
2   Sakamoto, K. M. *et al.* Protacs: chimeric molecules that target proteins to the Skp1-
Cullin-F box complex for ubiquitination and degradation. *Proc Natl Acad Sci U S A* **98**,
8554-8559, doi:10.1073/pnas.141230798 (2001).
3   Snyder, L. B. *et al.* Discovery of ARV-110, a first in class androgen receptor degrading
PROTAC for the treatment of men with metastatic castration resistant prostate cancer.
Cancer Res **81**, 43 (2021).
4   Hagenbeek, T. J. *et al.* An allosteric pan-TEAD inhibitor blocks oncogenic YAP/TAZ
signaling and overcomes KRAS G12C inhibitor resistance. *Nat Cancer* **4**, 812-828,
doi:10.1038/s43018-023-00577-0 (2023).
5   Badia, I. M. P. *et al.* Gene regulatory network inference in the era of single-cell multi-
omics. *Nat Rev Genet*, doi:10.1038/s41576-023-00618-5 (2023).
6   Alvarez, M. J. *et al.* Functional characterization of somatic mutations in cancer using
network-based inference of protein activity. *Nat Genet* **48**, 838-847, doi:10.1038/ng.3593
(2016).
7   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat
Methods* **14**, 1083-1086, doi:10.1038/nmeth.4463 (2017).
8   Kartha, V. K. *et al.* Functional inference of gene regulation using single-cell multi-omics.
Cell Genom **2**, doi:10.1016/j.xgen.2022.100166 (2022).
9   Fleck, J. S. *et al.* Inferring and perturbing cell fate regulomes in human brain organoids.
Nature **621**, 365-372, doi:10.1038/s41586-022-05279-8 (2023).

10 Kamal, A. *et al.* GRaNIE and GRaNPA: inference and evaluation of enhancer-mediated
gene regulatory networks. *Mol Syst Biol* **19**, e11627, doi:10.15252/msb.202311627
(2023).
11 Bravo Gonzalez-Blas, C. *et al.* SCENIC+: single-cell multiomic inference of enhancers
and gene regulatory networks. *Nat Methods* **20**, 1355-1367, doi:10.1038/s41592-023-
01938-4 (2023).
12 Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell
chromatin accessibility analysis. *Nat Genet* **53**, 403-411, doi:10.1038/s41588-021-
00790-6 (2021).
13 Inukai, S., Kock, K. H. & Bulyk, M. L. Transcription factor-DNA binding: beyond binding
site motifs. *Curr Opin Genet Dev* **43**, 110-119, doi:10.1016/j.gde.2017.02.007 (2017).
14 Worsley Hunt, R. & Wasserman, W. W. Non-targeted transcription factors motifs are a
systemic component of ChIP-seq datasets. *Genome Biol* **15**, 412, doi:10.1186/s13059-
014-0412-4 (2014).
15 Duan, J. *et al.* Rational Reprogramming of Cellular States by Combinatorial Perturbation.
Cell Rep **27**, 3486-3499 e3486, doi:10.1016/j.celrep.2019.05.079 (2019).
16 Baca, S. C. *et al.* Reprogramming of the FOXA1 cistrome in treatment-emergent
neuroendocrine prostate cancer. *Nat Commun* **12**, 1979, doi:10.1038/s41467-021-
22139-7 (2021).
17 Han, M. *et al.* FOXA2 drives lineage plasticity and KIT pathway activation in
neuroendocrine prostate cancer. *Cancer Cell* **40**, 1306-1323 e1308,
doi:10.1016/j.ccell.2022.10.011 (2022).
18 Tang, F. *et al.* Chromatin profiles classify castration-resistant prostate cancers
suggesting therapeutic targets. *Science* **376**, eabe1505, doi:10.1126/science.abe1505
(2022).
19 Xiao, L. *et al.* Targeting SWI/SNF ATPases in enhancer-addicted prostate cancer.
Nature **601**, 434-439, doi:10.1038/s41586-021-04246-z (2022).
20 He, B. *et al.* GATA2 facilitates steroid receptor coactivator recruitment to the androgen
receptor complex. *Proc Natl Acad Sci U S A* **111**, 18261-18266,
doi:10.1073/pnas.1421415111 (2014).
21 Rytinki, M. *et al.* Dynamic SUMOylation is linked to the activity cycles of androgen
receptor in the cell nucleus. *Mol Cell Biol* **32**, 4195-4205, doi:10.1128/MCB.00753-12
(2012).
22 Dufour, C. R. *et al.* The mTOR chromatin-bound interactome in prostate cancer. *Cell
Rep* **38**, 110534, doi:10.1016/j.celrep.2022.110534 (2022).
23 Cyrta, J. *et al.* Role of specialized composition of SWI/SNF complexes in prostate
cancer lineage plasticity. *Nat Commun* **11**, 5549, doi:10.1038/s41467-020-19328-1
(2020).
24 Beltran, H. *et al.* Divergent clonal evolution of castration-resistant neuroendocrine
prostate cancer. *Nat Med* **22**, 298-305, doi:10.1038/nm.4045 (2016).
25 Ku, S. Y. *et al.* Rb1 and Trp53 cooperate to suppress prostate cancer lineage plasticity,
metastasis, and antiandrogen resistance. *Science* **355**, 78-83,
doi:10.1126/science.aah4199 (2017).
26 Bluemn, E. G. *et al.* Androgen Receptor Pathway-Independent Prostate Cancer Is
Sustained through FGF Signaling. *Cancer Cell* **32**, 474-489 e476,
doi:10.1016/j.ccell.2017.09.003 (2017).
27 Mesman, S. & Smidt, M. P. Tcf12 Is Involved in Early Cell-Fate Determination and
Subset Specification of Midbrain Dopamine Neurons. *Front Mol Neurosci* **10**, 353,
doi:10.3389/fnmol.2017.00353 (2017).

28   Liu, W. *et al.* BRD4 regulates Nanog expression in mouse embryonic stem cells and
preimplantation embryos. *Cell Death Differ* **21**, 1950-1960, doi:10.1038/cdd.2014.124
(2014).
29   Choi, H. I. *et al.* BET inhibitor suppresses migration of human hepatocellular carcinoma
by inhibiting SMARCA4. *Sci Rep* **11**, 11799, doi:10.1038/s41598-021-91284-2 (2021).
30   Rahman, S. *et al.* The Brd4 extraterminal domain confers transcription activation
independent of pTEFb by recruiting multiple proteins, including NSD3. *Mol Cell Biol* **31**,
2641-2652, doi:10.1128/MCB.01341-10 (2011).
31   Xu, X. *et al.* Bromodomain Containing Protein 4 (BRD4) Regulates Expression of its
Interacting Coactivators in the Innate Response to Respiratory Syncytial Virus. *Front Mol*
Biosci **8**, 728661, doi:10.3389/fmolb.2021.728661 (2021).
32   Alver, B. H. *et al.* The SWI/SNF chromatin remodelling complex is required for
maintenance of lineage specific enhancers. *Nat Commun* **8**, 14648,
doi:10.1038/ncomms14648 (2017).
33   Blumli, S. *et al.* Acute depletion of the ARID1A subunit of SWI/SNF complexes reveals
distinct pathways for activation and repression of transcription. *Cell Rep* **37**, 109943,
doi:10.1016/j.celrep.2021.109943 (2021).
34   Sobczak, M., Pitt, A. R., Spickett, C. M. & Robaszkiewicz, A. PARP1 Co-Regulates
EP300-BRG1-Dependent Transcription of Genes Involved in Breast Cancer Cell
Proliferation and DNA Repair. *Cancers (Basel)* **11**, doi:10.3390/cancers11101539
(2019).
35   Shang, X. Y. *et al.* ARID1A deficiency weakens BRG1-RAD21 interaction that
jeopardizes chromatin compactness and drives liver cancer cell metastasis. *Cell Death*
Dis **12**, 990, doi:10.1038/s41419-021-04291-6 (2021).
36   Lazar, J. E. *et al.* Global Regulatory DNA Potentiation by SMARCA4 Propagates to
Selective Gene Expression Programs via Domain-Level Remodeling. *Cell Rep* **31**,
107788, doi:10.1016/j.celrep.2020.107788 (2020).
37   Marino, M. M. *et al.* Interactome mapping defines BRG1, a component of the SWI/SNF
chromatin remodeling complex, as a new partner of the transcriptional regulator CTCF. *J*
Biol Chem **294**, 861-873, doi:10.1074/jbc.RA118.004882 (2019).
38   Liao, L. *et al.* Multiple tumor suppressors regulate a HIF-dependent negative feedback
loop via ISGF3 in human clear cell renal cancer. *eLife* **7**, e37925,
doi:10.7554/eLife.37925 (2018).
39   Granja, J. M. *et al.* Single-cell multiomic analysis identifies regulatory programs in mixed-
phenotype acute leukemia. *Nat Biotechnol* **37**, 1458-1465, doi:10.1038/s41587-019-
0332-7 (2019).
40   Ungerback, J., Ahsberg, J., Strid, T., Somasundaram, R. & Sigvardsson, M. Combined
heterozygous loss of Ebf1 and Pax5 allows for T-lineage conversion of B cell
progenitors. *J Exp Med* **212**, 1109-1123, doi:10.1084/jem.20132100 (2015).
41   Yeamans, C. *et al.* C/EBPalpha binds and activates the PU.1 distal enhancer to induce
monocyte lineage commitment. *Blood* **110**, 3136-3142, doi:10.1182/blood-2007-03-
080291 (2007).
42   Thomas, G. D. *et al.* Deleting an Nr4a1 Super-Enhancer Subdomain Ablates Ly6C(low)
Monocytes while Preserving Macrophage Gene Function. *Immunity* **45**, 975-987,
doi:10.1016/j.immuni.2016.10.011 (2016).
43   Kanhere, A. *et al.* T-bet and GATA3 orchestrate Th1 and Th2 differentiation through
lineage-specific targeting of distal regulatory elements. *Nat Commun* **3**, 1268,
doi:10.1038/ncomms2260 (2012).
44   Newton, R. H. *et al.* Maintenance of CD4 T cell fitness through regulation of Foxo1. *Nat*
Immunol **19**, 838-848, doi:10.1038/s41590-018-0157-4 (2018).

45   Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and
Activation. *Cell* **159**, 647-661, doi:10.1016/j.cell.2014.09.029 (2014).
46   Garcia-Alonso, L., Holland, C. H., Ibrahim, M. M., Turei, D. & Saez-Rodriguez, J.
Benchmark and integration of resources for the estimation of human transcription factor
activities. *Genome Res* **29**, 1363-1375, doi:10.1101/gr.240663.118 (2019).
47   Müller-Dott, S. *et al.* Expanding the coverage of regulons from high-confidence prior
knowledge for accurate estimation of transcription factor activities. *bioRxiv*,
2023.2003.2030.534849, doi:10.1101/2023.03.30.534849 (2023).
48   Greene, C. S. *et al.* Understanding multicellular function and disease with human tissue-
specific networks. *Nat Genet* **47**, 569-576, doi:10.1038/ng.3259 (2015).
49   Gaublomme, J. T. *et al.* Nuclei multiplexing with barcoded antibodies for single-nucleus
genomics. *Nat Commun* **10**, 2907, doi:10.1038/s41467-019-10756-2 (2019).
50   Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat
Methods* **14**, 975-978, doi:10.1038/nmeth.4401 (2017).
51   Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor
sequence specificity. *Cell* **158**, 1431-1443, doi:10.1016/j.cell.2014.08.009 (2014).
52   Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of
Problematic Regions of the Genome. *Sci Rep* **9**, 9354, doi:10.1038/s41598-019-45839-z
(2019).
1340

Figure 1: Epiregulon constructs GRNs to infer TF activity at the single cell level

Figure 2: Epiregulon identifies cell fate changes

Figure 3: Epiregulon predicts responses towards TF-targeting drugs

Figure 4: Comparison of ChIP-seq and motif annotations

a

b

c

Figure 5: Epiregulon predicts responses towards chromatin-modifying drugs

Figure 6: Epiregulon-constructed GRNs predict effects of perturbations