1 **Epiregulon: Inference of single-cell transcription factor activity to dissect mechanisms of**
2 **lineage plasticity and drug response**
3

4 Tomasz Włodarczyk[1], Aaron Lun[1], Diana Wu[2], Shreya Menon[3,4,5], Shushan Toneyan[1], Kerstin
5 Seidel[2], Liang Wang[2], Jenille Tan[2], Shang-Yang Chen[1], Timothy Keyes[1], Aleksander
6 Chlebowski[1], Yu Guo[1], Ciara Metcalfe[2], Marc Hafner[1,2], Christian W. Siebel[2], M. Ryan Corces[3,4,5],
7 Robert Yauch[2]*, Shiqi Xie[2]*, Xiaosai Yao[1,2]*
8
[1]Department of Computational Sciences, Genentech Inc, South San Francisco, CA, USA
[2]Department of Discovery Oncology, Genentech Inc, South San Francisco, CA, USA
[3]Gladstone Institute of Neurological Disease, Gladstone Institutes, San Francisco CA, USA
[4]Gladstone Institute of Data Science and Biotechnology, Gladstone Institutes, San Francisco CA,
USA
[5]Department of Neurology, University of California San Francisco, San Francisco CA, USA
15

*correspondence

yauch.bob@gene.com; xie.shiqi@gene.com; yao.xiaosai@gene.com

Abstract

Transcription factors (TFs) and transcriptional coregulators represent an emerging and exciting
class of targets. By quantifying target gene modulation, gene regulatory networks (GRNs)
delineate disease biology and evaluate pharmacological agents targeting these regulators.
However, none of the existing methods are explicitly designed to measure the effects of
perturbations in which TF expression is decoupled from its activity. We present Epiregulon, a
method that constructs GRNs from single-cell ATAC-seq and RNA-seq data for accurate
prediction of TF activity. Our weight estimation, based on co-occurrence of TF expression and
chromatin accessibility, avoids erroneous inflation of TF activity as seen with TF expression only
approaches. Furthermore, our utilization of ChIP-seq data expands inference to transcriptional
coregulators lacking defined motifs. Our extensive network of regulators facilitates identification
of cell-state specific interaction partners. Using Epiregulon, we uncover divergent cell fate
transitions of prostate cancer cells driven by NKX2-1 and GATA6 overexpression. We accurately
predicted the effects of AR inhibition across various drug modalities. Finally, Epiregulon was able
to infer the context-dependent activity of a chromatin remodeler lacking a defined motif sequence
and recapitulate the unique etiologies of prostate cancer. By mapping out the network of key
regulators across a multitude of perturbations, Epiregulon can accelerate the discovery of new
therapeutics targeting transcription factors.

Introduction

Transcription factors and transcriptional coregulators shape cell fates and lineage commitment, and their dysregulation drives congenital diseases and tumor growth. Transcription factors bind to specific DNA sequences whereas coregulators interact with transcription factors in a context-specific manner and lack defined motifs. These transcriptional modulators represent an important class of therapeutic targets in oncology and beyond. Therapeutics targeting transcriptional regulators inhibit their activity by either blocking the ligand binding domains[1], degrading the protein[2,3] or disrupting protein-protein interactions[4]. Delineating the functions of transcriptional regulators can accelerate our understanding of disease biology and drug discovery.

Gene regulatory networks (GRNs) model the underlying circuitry of gene regulation and have been applied to understand lineage commitment and plasticity and drug response[5]. Early methods to construct gene regulatory networks relied exclusively on gene expression data and sought to identify association between expression of target genes and transcription factors[6,7]. Now, single-cell multiomics technologies provide chromatin accessibility information in addition to gene expression and recent GRN inference methods leverage both modalities to improve performance (Table 1). Most of these methods, including CellOracle[8], FigR[9], Pando[8] and GRaNIE[10], rely on linear relationships between transcription factors and their target genes to model gene regulation. In addition, SCENIC+ utilizes random forest regression to model the regulatory relationships between TF, regulatory elements and target genes[11].

Several challenges still impede the broader utilization of these GRN inference methods in basic biology and drug discovery. First, none of these methods are specifically designed to predict changes in which the TF activity is decoupled from its gene expression. These include drug perturbations in which inhibitors disrupt protein complex formation or localization, and post-translational modifications that can impact TF activity and also CRISPR genome editing in which genes can be silenced by insertions or deletions at the DNA level but without causing appreciable changes in gene expression. Second, the use of motif sequences in GRN construction precludes prediction of transcriptional coregulators, an important class of regulators without defined motifs. Lastly, current studies rely on ChIP-seq and HiC data to benchmark GRNs but TF occupancy and chromatin conformation do not directly measure the true regulatory activity of TFs. Therefore, erroneous estimation of TF activity misled by gene expression, sole reliance on TF motifs and lack of appropriate benchmarking datasets are three limitations in current GRN inference methods.

Here we present Epiregulon, a method that constructs a GRN from single-cell multiomics data and TF occupancy data to infer activity of transcriptional regulators. Unlike most other GRN tools, Epiregulon utilizes accompanying ChIP-seq data, thus allowing inference of both transcription factors and transcriptional coregulators lacking defined motifs. Epiregulon utilizes weight estimation methods that account for co-occurrence of both TF gene expression and chromatin accessibility to handle decoupling of TF expression and activity. Functional interpretation of GRNs through Jaccard similarity of target genes and pathway annotations uncovers drivers and interaction partners of lineage transition and drug response in a cell-state specific and context-aware manner. We performed simulations and generated ground truth data to robustly benchmark

80   its performance in a multitude of datasets including lineage development and drug treatment.

81   Epiregulon is implemented as a R package and interfaces with ArchR[12] or any other

82   preprocessing tools.

84   Table 1: Comparison of different GRN inference methods

Feature/package	**Epiregulon**	**GRaNIE**	**SCENIC+**	**FigR**	**Pando**	**cellOracle**
Prior information to predict TF occupancy	Granges list of ChIP-seq peaks or motif coordinates	BED file with motif coordinates	Pre-computed database with motif scores	Motif database compiled from cisBP	Motif database compiled from cisBP and JASPAR	Motif data from cisBC v. 2.0
TF-target gene model	Wilcoxon-based co-occurrence, correlation and mutual information	Expression correlation	Gradient Boosting Machine regression	TF-DORC correlation and TF enrichment	Regularized GLM, Bayesian regression models, gradient boosting regression, bagging and ridge Bayesian models	Regularized machine - learning linear model
Pruning method	TF-RE-TG co-occurrence (chi-sq)	Based on permuted data	Filtering TF-TG pairs using GSEApy	No pruning	Linear regression	Incorporated in the main model
Addressing sparsity	pseudobulks	no	imputation	KNN smoothing	no	KNN imputation, pseudocells
Data pre-processing compatibility	ArchR or SingleCellExperiment	Peak and gene expression matrices	scanpy, pycisTopic, pycisTarget	Summarized Experiment (peak), gene expression matrix	Signac	cicero, scanpy

Results

Epiregulon constructs GRNs to infer TF activity at the single cell level

Epiregulon aims to map out target genes of transcriptional regulators and track changes in their regulatory potential in each cell state, thereby identifying master regulators of cell fate changes. It assumes that a transcription factor or a transcriptional coregulator (collectively referred to as TF for simplicity) binds to regulatory elements (REs) and modulates the expression of nearby target genes (TGs). Epiregulon assigns regulatory elements (REs) to their putative target genes based on correlation between chromatin accessibility and gene expression (Figure 1). Each regulatory element is then interrogated for TF occupancy based on user-provided or/and a compilation of public TF ChIP-seq data from ChIP-Atlas and ENCODE. The use of ChIP-seq data enables us to extend activity inference to chromatin modifiers and co-factors which do not have well-defined binding motifs. Furthermore, accumulating evidence suggests that TF occupancy is determined not just by motif sequences but also by other factors including interactions with other TFs and co-factors, local DNA structure and genomic features[13]. A substantial proportion of TF binding sites do not harbor motifs[14]. Therefore, ChIP-seq data is likely to yield a more relevant set of target genes than motif annotations are.

We combine the RE-TG links with TF occupancy at REs to establish an initial unweighted tripartite graph with TFs, REs and TGs being the nodes and their connections being edges. An optional pruning step of the graph removes potentially spurious regulatory relationships and TFs with undetectable gene expression. We then estimate the weights of the TF-RE-TG edges using the Wilcoxon test statistic, computed by comparing the ranks of the target gene expression between the group of cells that express the corresponding TF and have chromatin accessibility at that particular RE versus the remaining cells. This graph becomes the final gene regulatory network returned by Epiregulon. Finally, the activity of a TF is defined as the weighted sum of its target gene expression normalized by the number of target genes. The exact algorithm is described in the Methods; see also Supplementary Note Section 1 for comparison of parameter choices.

It is of great interest to understand the functions of a TF through its differential activity between cell states or treatment conditions. Differential TF activity can result from at least two different scenarios; 1) TFs can be turned on or off by changes in gene expression, or 2) they can redistribute to different genomic regions due to mutations or hijacking by other factors. The first scenario can be adequately captured by differences in total TF activity whereas the second scenario requires the consideration of regulon membership. In addition to quantifying differential total TF activity between cell states, Epiregulon employs a second method for differential network analysis. We use the Wilcoxon method to derive cluster-specific weights, subtract the edge weights between two networks and compute the degree centrality of the resulting edge-subtracted network. See Supplementary Note Section 2 for simulation of differential networks.

Finally, it is desirable to interpret the functions of a TF through the pathway enrichment of its regulons and through its interaction partners. Epiregulon provides pathway enrichment of the regulons and further utilizes Jaccard similarity to estimate the overlap between regulons to identify

131   potential interaction partners. These functional annotations of GRNs serve to provide biological
132   interpretability.
133
Epiregulon identifies cell fate changes
To evaluate Epiregulon's performance, we generated our own data using Reprogram-Seq[15] to
model lineage transition of prostate adenocarcinomas by overexpressing defined transcription
factors. We separately introduced lentivirus encoding NKX2-1 and GATA6 into a prostate
adenocarcinoma cell line LNCaP and tracked changes in transcriptomic and epigenomic profiles
at the single cell level. NKX2-1 is a known driver of AR-independence and neuroendocrine
transition in prostate cancer[16,17] whereas GATA6 has unknown function in the prostate. These
factors are not expressed in parental LNCaP and were used for subsequent benchmarking
analysis. Reprogram-Seq data could serve as the ground truth to benchmark the inference of TF
activity because we expect to see increased TF activity in cells with exogenous expression of
NKX2-1 and GATA6.
145
Expression of these transcription factors altered the cell states of LNCaP, with cluster 1 enriched
in cells expressing GATA6 and cluster 3 enriched in cells expressing NKX2-1 (Figure 2a). Gene
expression of these factors could not accurately map the identity of TFs being introduced, largely
because transcription factors are lowly expressed and 3' bias of scRNAseq makes detection of
exogenous transcripts difficult (Figure 2b). In contrast, TF activity computed from Epiregulon
recapitulated the correct cell identity despite the low expression of these TFs (Figure 2b).
152
We performed a systematic benchmarking exercise to evaluate the performance of Epiregulon
against other multiomic GRN inference methods. Compared to FigR, Pando, cellOracle, GRaNIE
and SCENIC+, Epiregulon was the only method that was able to predict NKX2-1 activity (Figure
2c). This is because the use of ChIP-seq data allowed Epiregulon to predict the largest collection
of regulators (Figure 2d). We evaluated the GRN inference methods on whether TF activity is able
to classify cells expressing the matching TF vs the remaining cells. For GATA6, Epiregulon had
the second highest area under the receiving operating characteristic (AUROC) value (Figure 2c).
We next examined Epiregulon's performance across its different weight methods. Because TF
activity was driven by increased TF expression in reprogram-seq, weight estimation based on the
correlation between TF-TG expression was the best performing method (Figure S1a). This implies
that Epiregulon gives consistent predictions as other top-performing GRN inference methods and
offers improved sensitivity in predicting activity of those factors with low gene expression.
165
We further investigated the accuracy of the target genes inferred. We showed that both unpruned
and pruned GATA6 and NKX2-1 targets inferred by Epiregulon ranked highly by differential
expression (Figure S1b). Furthermore, cluster-specific GRN showed that NKX2-1 targets were
exclusively expressed in cluster 3 while GATA6 targets were exclusively expressed in cluster 1
(Figure S1c). As another independent validation, we engineered a dox-inducible cell line
expressing NKX2-1 or eGFP as a control. The log changes of our inferred NXK2-1 targets upon
dox treatment correlated with the NKX2-1 target gene weights inferred by Epiregulon (Figure 2e,
S1d), suggesting that the GRN weights inferred by Epiregulon are informative of regulatory
strength.

Global analysis of TF activity identified other TFs whose activities were concomitantly altered by
introduction of these reprogramming factors. Jaccard similarity found FOSL2 (a paralog of
FOSL1), YAP1 and TEAD1 to be amongst the top interaction partners of GATA6 (Figure 2f, S1e).
These factors are known to reprogram prostate adenocarcinomas towards a stem cell-like (SCL)
state[18]. This led us to examine whether GATA6 itself could also reprogram LNCaP cells towards
the SCL subtype. We examined regions that are uniquely accessible in the SCL tumors[18].
Strikingly, GATA6 overexpression led to increased chromatin accessibility at these SCL-specific
regions and GATA6 activity was highly correlated with the combined chromatin accessibility score
(Figure 2g, 2h). Lineage trajectory resulting from GATA6 expression differed from the
neuroendocrine cell state induced by NKX2-1 (Figure 2g, 2h, S1f). Taken together, Epiregulon
uncovered a previously unknown causal relationship between GATA6 expression and lineage
transition to SCL subtype of prostate cancer.

Epiregulon predicts responses towards TF targeting drugs
In the previous example, Epiregulon was able to predict changes in TF activity when they are
largely driven by changes in their expression. A more challenging task is to predict changes when
TF activity is decoupled from its gene expression. We generated a second dataset to evaluate
changes in AR activity following drug treatment. In total, 4 prostate cancer cell lines (2 AR-
dependent and 2 AR-independent) were treated with 3 AR-targeting agents (Figure 3a). The first
agent is the clinically approved AR antagonist, enzalutamide, which interferes with AR protein
function by blocking its ligand binding domain[1]. The second agent is ARV-110, a degrader of AR
protein which acts by bringing an E3 ubiquitin ligase in close proximity to an AR protein[3]. We also
synthesized a third agent, SMARCA2.4_1, a degrader of the SWI/SNF chromatin remodeling
complex (Figure S2a-b); inhibition of the same protein complex is known to decrease AR activity
by evicting it from the chromatin[19]. These pharmacological agents do not directly or consistently
suppress the transcription of AR target genes, thus serving as an example of the decoupling of
the TF expression from its activity.

Epiregulon predicted decreased AR activity following enzalutamide and ARV110 treatment in the
known sensitive cell lines, LNCaP and VCaP and minimal changes in AR activity in the AR-
independent cell lines, DU145 and H660 (Figure 3b). Other GRN inference methods failed to
predict decreased AR activity in either or both cell lines, with CellOracle even predicting an
increased AR activity in VCaP (Figure 3c). Gene expression of AR was not informative of the
effect on AR activity because enzalutamide increased AR gene expression in VCaP cells despite
its antagonistic function (Figure 3d). Like the Reprogram-Seq data, Epiregulon recovered the
greatest number of regulators (Figure 3e). By estimating weights based on both TF expression
and RE chromatin accessibility, Epiregulon outperformed methods utilizing TF expression alone,
and robustly predicted decreased AR activity despite discordant AR gene expression (Figure 3c,
Figure S2c).

We next asked whether ChIP-seq data generated from merging ChIP-atlas and ENCODE
(thereafter referred as merged ChIP-seq data) could yield comparable performance as cell line-

matched ChIP-seq data. TF activity inferred from AR and FOXA1 ChIP-seq data of LNCaP and
VCaP cells correlated well with that produced by merged ChIP-seq (Figure S2c, S2d). Taken
together, GRN constructed using merged ChIP-seq data could accurately recapitulate TF activity.
222
223   **GRN constructed from ChIP-seq data is a good predictor of TF activity**
224   Epiregulon uses ChIP-seq data as the default method to predict TF occupancy but can also
225   incorporate motif annotations. Given that the majority of the other GRN inference methods start
226   with motif sequences, we would like to systematically compare the accuracy of TF activity
227   prediction starting from ChIP-seq data or motif annotations.
228
Using the Reprogram-Seq dataset, we showed that ChIP-seq (obtained by merging public ChIP-
seq datasets from ENCODE and ChIP-Atlas) is superior to motif annotations in predicting TF
activity for both GATA6 and NKX2-1 (Figure 4a, 4b). We further asked whether selecting a set of
high-confidence motifs can improve TF activity prediction. We leveraged deep learning sequence-
to-function models or deep neural networks (DNNs) to prioritize motif patterns of importance
within the regulatory elements. DNNs are capable of learning complex patterns in the input data
associated with accessible regions to make predictions of ATAC-seq coverage[20-22]. Here in
addition to comparing ChIP-seq and motif annotations, we also applied thresholds to motif
importance scores assigned by two independent sequence deep learning models, Basenji and
ChromBPNet (see Methods). However, thresholded motifs did not improve the performance of TF
activity; in fact, overly stringent thresholding resulted in a reduced number of target genes and
worse performance (Figure 4a, 4b, Figure S3a, S3b). The weights of the target genes contributed
to the accuracy of TF prediction since shuffling of weights reduced AUROCs (Figure S3c, S3d).
242
Next, we leveraged the prostate dataset to evaluate the performance of GRN. We found that in
contrast to the Reprogram-Seq data, AR motifs outperformed merged ChIP-seq in terms of
differentiating drug treated vs DMSO treated cells (Figure 4c). We could however improve the
accuracy of ChIP-seq by utilizing cell line-matched ChIP-seq. In fact, combining motif annotation
with cell line-matched ChIP-seq binding was the best predictor of AR activity (Figure 4c). DNN-
based models did not further improve the performance of motif annotation (Figure 4c). Taken
together, we showed that target genes inferred by ChIP-seq data outperforms motif annotations
in the estimation of TF activity.
251
Epiregulon predicts response of drug targeting transcriptional coregulator
Transcriptional coregulators cooperate with transcription factors to regulate gene expression and
are attractive targets in oncology. The challenge with predicting responses of transcriptional
coregulators is that these factors do not contain defined motifs as they can interact with different
transcription factors depending on the cellular context. As such, GRN inference methods which
rely on TF motifs cannot predict the activity of these factors. SMARCA2 and SMARCA4, two
mutually exclusive paralog proteins encoding the ATPase subunit of the SWI/SNF chromatin
remodeler are examples of transcriptional coregulators. We evaluated the performance of
Epiregulon based on its ability to predict decreased SMARCA2/4 activity upon treatment with a
degrader of SMARCA2/4 protein. We utilized the same AR dataset as in Figure 3 but focused on
the third agent, SMARCA2.4_1. We performed most of our analysis on SMARCA4 since it is the

predominant paralog that affects the proliferation of prostate cancer cell lines[23]. Pharmacological
degradation of SMARCA4 allows benchmarking of activity inference for a highly context-
dependent transcriptional coregulator lacking a defined motif.
266
To circumvent the lack of motifs, Epiregulon utilizes public SMARCA4 ChIP-seq to infer the most
likely target genes in a cell-line aware manner. Epiregulon predicted decreased SMARCA4
activity across all 4 prostate cancer cell lines (Figure 5a). None of the other GRN inference
methods we benchmarked were able to predict the activity of SMARCA4. Within Epiregulon,
weight estimation methods that take into account both TF gene expression and RE chromatin
accessibility outperformed methods that only take into account TF expression (Figure S4a).
Furthermore, SMARCA4 activities calculated from merged ChIP-seq and cell line-matched ChIP-
seq were consistent (Figure S4b). The SMARCA4 regulons recapitulated the unique etiologies of
prostate cancer, with AR-dependent cell lines LNCaP and VCaP enriched for androgen response,
stem-cell like cell line DU145 enriched for activation of YAP activity[18] and the neuroendocrine cell
line NCI-H660 enriched for the loss of TP53, activation of EZH2[24,25] and MAPK signaling[26] (Figure
5b).
279
Epiregulon can inform mechanisms of action of the SMARCA4 degrader by discovering TFs co-
perturbed with SMARCA4 (Figure 5c). We identified differential TFs by subtracting DMSO treated
from SMARCA4 degrader treated networks. TFs were ranked by Jaccard similarity with
SMARCA4 after background correction using edge permuted networks. Common to all 4 cell
lines, highly ranked TFs included components of general transcriptional machinery including
BRD4 and EP300 and also regulators of chromatin structure including components of the cohesin
complex (RAD21, SMC3, SMC1A) and chromosome looping (YY1, CTCF and MED1). These
proteins have been previously implicated to associate with SMARCA4 (Table 2). Unique to the
individual cell lines, SMARCA4 interacts with AR in the AR-dependent cell lines (LNCaP and
VCaP). In the neuroendocrine cell line NCI-H660, SMARCA4 interacts with TCF12, a factor
known to be important for neuronal development[27] and EZH2, a factor known to mediate
neuroendocrine transition in prostate cancer[25]. Taken together, Epiregulon expands the inference
of activity to include transcriptional coregulators lacking defined motifs and is able to capture their
context specificity and infer the most relevant biological processes.
294
295   Table 2: Associated partners of SMARCA4 identified by differential network analysis

Function	Gene	References
Transcription	BRD4	25,28, 29, 30, 31
Transcription	EP300	32, 33, 34
Chromosome structure	Cohesin (RAD21, SMC1A, SMC3)	35, 36
Chromatin looping	YY1 and CTCF	37, 38

Epiregulon identifies lineage factors and predicts effects of perturbations

Finally, we ask whether regulons generated from single-cell multiomic data can be applied to
scRNA-seq dataset lacking chromatin information. We first used Epiregulon to compute the
activity of lineage factors in a scMultiomic dataset of human peripheral blood mononuclear cells
(PBMCs)[39] (Figure 6a). Epiregulon recapitulated the lineage-specific activity of known lineage
factors including GATA3 in T and NK cells, EBF1 and PAX5 in B cells and SPI1 in myeloid cells
(Figure 6b), and gene set enrichment of their target genes is consistent with their known biological
functions (Figure S5a). We used the Wilcoxon method to obtain cell-type specific weights for all
TF-TG pairs. As an example to illustrate the specificity of the GRN, the B cell lineage factor EBF1
had its target genes largely confined to preB cells but not to CD4 memory T cells (Figure 6c). We
further utilized Epiregulon's similarity function to identify known interacting partners (Figure 6d).
The most similar TF to EBF1 was PAX5; EBF1 and PAX5 regulate B cell development in a highly
coordinated manner[40]. We further compared the global network topology of monocytes and CD4
memory T cells; degree centrality of the differential graph revealed known lineage factors of
monocytes (SPI1, CEBPA, NR4A1)[41,42] and CD4 T cells (TBX21, GATA3, LEF1, FOXO1)[43,44],
suggesting that Epiregulon is able to prioritize lineage factors from differential network analysis.
314

Next, we applied the GRN of the above PBMC dataset to a new scRNAseq dataset containing
K562, a lymphoblast cell line, engineered to express CRISPRa and guides targeting defined
factors[45]. The barcode information in each cell encodes the guide RNAs corresponding to the TFs
they activate and serves as the ground truth. GRN from Epiregulon better predicted the effect of
CRISPRa perturbations than gene expression, lineage-agnostic regulons (Dorothea[46] and
CollecTRI[47]) and lineage-specific regulons (Humanbase[48]) did, as reflected in the AUROC values
(Figure 6f, Figure S5b). Epiregulon also captured the highest number of TFs (Figure 6f). The
improvement in performance is largely due to the reduction of data sparsity by aggregating the
appropriate target genes, as shown by PRDM1 (Figure 6g). Taken together, based on this ground
truth dataset, GRNs reconstructed from Epiregulon could accurately identify known lineage
factors and predict effects of gene perturbation in datasets containing only scRNA-seq.
326
327

Discussion

We develop Epiregulon, a method to construct gene regulatory networks and infer activity of both
transcription factors and transcriptional coregulators lacking defined motifs, an important class of
regulators largely overlooked by previous GRN inference methods. When benchmarked against
other GRN inference methods, Epiregulon performs most robustly across a multitude of datasets,
especially when TF activity is decoupled from its gene expression such as in drug perturbations.
Epiregulon predicts the largest repertoire of regulators, constructing an extensive regulatory
network that is functionally annotated with pathway enrichment analysis and interaction partners.
336

Unlike most other GRN inference methods, Epiregulon allows construction of GRNs from ChIP-
seq data but still preserves the option to filter ChIP-seq peaks based on the presence of motifs.
ChIP-seq data allows the prediction of chromatin co-activators and modifiers lacking defined

motifs. Importantly, we were able to infer the activity of the chromatin remodeler SMARCA4 in a
context-aware manner. We systematically compared the performance of TF prediction based on
GRNs constructed from ChIP-seq vs motif annotations, fine tuning motif annotations using deep-
learning neural networks models. We found that TF occupancy obtained from public or user-
provided ChIP-seq data can accurately predict TF activity.
345
Weight estimation by Epiregulon enhances the accuracy of TF activity by assigning the
importance scores to target genes. The Wilcoxon weight estimation method, based on co-
occurrence of TF gene expression and chromatin accessibility, is especially well suited for
applications in which TF activity is discordant with its gene expression such as drug perturbations.
The Wilcoxon method also enables estimation of cell-cluster specific weights and subsequent
differential network topology analysis. Epiregulon correctly recapitulates the response of multiple
AR-targeting agents in various prostate cancer cell line models, and is well positioned to predict
TF activity in a wide range of biological applications and systems.
354
Epiregulon enables us to make new discoveries from the ground truth data we generated. We
found an unexpected connection between GATA6, YAP1 and TEAD1 which led to the discovery
that GATA6 overexpression leads to reprogramming of the prostate adenocarcinoma cell line
LNCaP towards a stem-cell like subtype. This was possible because compared to other methods,
Epiregulon covers the most extensive regulators including coregulators lacking motifs (in this case
YAP1), enabling the identification of interaction partners based on shared target genes.
361
Several limitations of our study remain. While Epiregulon is robust in predicting the overall TF
activity, we have not formally evaluated the accuracy of individual target genes. Second,
Epiregulon does not explicitly model cooperativity between transcription factors and partners, so
direct interactions between functionally related factors may have been overlooked. Lastly, given
the use of ground truth data, our evaluation of performance is limited to a few transcription factors
and it remains unclear whether Epiregulon uniformly performs well across multiple TFs. We
speculate that the quality of available ChIP-seq data greatly influences the prediction of TF
activity, and wherever possible, ChIP-seq data of TF of interest should be generated in the
matching cell lines to ensure robustness of the biological findings.
371
372 We envision that Epiregulon can enable drug discovery by evaluating response to therapeutic
373 agents not limited to oncology and identifying new drivers of resistance. Epiregulon is available
374 at https://github.com/xiaosaiyao/epiregulon along with the data package scMultiome
375 (https://bioconductor.org/packages/release/data/experiment/html/scMultiome.html).
376
377 **Methods**
378
Epiregulon
The Epiregulon workflow consists of several components: GRN construction, pruning of networks
(optional), estimation of weights, calculation of activity, functional annotation of regulons,
differential network analysis and identification of interaction partners. Epiregulon is designed to
work seamlessly with ArchR but can accept any input as long as they are formatted into

SingleCellExperiment objects. It is available as three related R packages. Epiregulon is the core
package that performs GRN construction and activity inference
(https://github.com/xiaosaiyao/epiregulon/). Epiregulon.extra contains differential analysis and
plotting functions (https://github.com/xiaosaiyao/epiregulon.extra). Epiregulon.archr contains
functions interfacing with ArchR (https://github.com/xiaosaiyao/epiregulon.archr).
389
Data preprocessing
Epiregulon assumes that prior preprocessing of the data has been performed by users' methods
of choice, requiring paired gene expression matrix (scRNA) and peak matrix (scATAC) along with
dimensionality reduction matrix as the input. If gene expression and peak matrix are not paired,
integration of scATAC-seq and scRNA-seq data must be performed prior to Epiregulon using
methods such as ArchR's addGeneIntegrationMatrix function.
396
For the datasets included in this manuscript, reads were mapped by the Cell Ranger ARC 2.0
and were further processed with the use of ArchR[12]. Briefly, cells were filtered based on TSS
enrichment (>6) and number of ATAC-seq reads mapped to the nuclear genome (>2500 and
<10^5). Moreover, doublets were removed using dedicated ArchR functions. ATAC-seq data was
represented as a 500bp tile matrix on which iterative LSI was performed. Similarly, normalized
gene expression data was subject to iterative LSI. The dimensionality reductions from both
modalities were then combined into a single one that was used for cell clustering. The ATAC-seq
peaks were called separately for each cluster and after normalization were merged to produce
one peak set for all cells (peak x cell matrix). The cells were assigned to their sample barcodes
using demuxEM[49].
407
Network construction
Epiregulon provides two similar methods to establish peak to gene links using the calculateP2G
function. If there is an existing ArchR project, Epiregulon retrieves the peak to gene links that
have been previously assigned by ArchR. Briefly, ArchR creates 500 cell aggregates, resampling
cells if needed. ArchR computes the correlation between chromatin accessibility and target genes
within a size window (default ±250 kb) and retains peak-gene pairs that exceed a correlation
threshold (default Epiregulon cutoff is 0.5). In the absence of an ArchR project, Epiregulon defines
cell aggregates using k-means clustering based on the reduced dimensionality matrix and
performs correlation in the same manner as ArchR does. If cluster labels are provided, cell
aggregates are formed within each cluster and cluster-specific correlations are reported in
addition to overall correlations.
419
TF occupancy data
Each regulatory element is then interrogated for TF occupancy based on a compilation of public
TF ChIP-seq binding sites. ChIP-seq were downloaded from ChIP-Atlas (chip-atlas.org) and
ENCODE (encodeproject.org). TF ChIP-seq peaks with FDR < 1 x 10^{-5} were retained. ChIP-seq
datasets of the same TFs were merged, yielding 1557 unique factors (human) and 768 unique
factors (mouse). Data is provided in the scMultiome package as GRanges list objects and can be
accessed using the getTFMotifInfo function from the Epiregulon package or directly from the
scMultiome package using the tfBinding function.

If desired, ChIP-seq peaks can be further annotated for the presence of motifs using the
addMotifScore function. If motif annotation has been performed previously using ArchR's
addMotifAnnotations function, motif annotations can be easily retrieved and appended to the peak
matrix. In the ArchR independent workflow, Epiregulon can annotate peak matrix using
motifmatchr's motif matching function[50] and cisbp as the reference motif database[51]. Alternatively,
users can start entirely from motif annotations and leverage deep learning motifs to select motifs
based on motif importance scores (see section on DNN models).
436

437 *Network pruning (optional)*
438 Epiregulon prunes the network by performing tests of independence on the observed number of
439 cells jointly expressing transcription factor (TF), regulatory element (RE) and target gene (TG) vs
440 the expected number of cells if TF/RE and TG are independently expressed using the
441 pruneRegulon function.
442

443 We define n as the total number of cells, k as the number of cells jointly expressing TF, TG and
444 RE above a set threshold, g as the number of cells jointly expressing TF and RE above a threshold
445 and h as the number of cells expressing TG above a threshold. p, the expected probability of cells
446 jointly expressing TF, TF and RE above a threshold is defined as follows:
447

448 $$P(cells\ expected\ to\ jointly\ express\ TF\ t, RE\ r\ and\ TG\ g) = p\ =\ \frac{g}{n} \times \frac{h}{n}$$

450 Two tests of independence are implemented, the binomial test and the chi-square test. In the
451 binomial test, the expected probability is p, the number of trials is the total number of cells n, and
452 the observed number of successes is k, the number of cells jointly expressing all three elements.
453

454 In the chi-square test, the expected probability for having all 3 elements active is also p. The
455 observed cell count for the active category is k, and the cell count for the inactive category is $n -$
456 k. P-values are calculated from chi-squared distribution with degree of freedom equal to 1.
457 Cluster-specific p-values are calculated if users supply cluster labels.
458

Estimation of weights
While network pruning provides statistics on the joint occurrence of TF-RE-TG, we would like to
further estimate the strength of regulation using the addWeights function. Biologically, this can be
interpreted as the magnitude of gene expression changes induced by changes in transcription
factor activity. Epiregulon provides 3 different methods to estimate weights. Two measures
(correlation and Wilcoxon statistics) give both the magnitude and directionality of changes
whereas weights computed by mutational information (MI) are always positive. Within 2 of the
methods (correlation and MI), there is an option of modeling the TG expression based only on TF
expression, or on the product of TF expression and RE chromatin accessibility. Consideration of
both TF expression and RE chromatin accessibility is highly recommended, especially for
scenarios in which TF activity and TF gene expression are decoupled (as in the case of drug
perturbations or CRISPR genome editing). The correlation and mutual information statistics are

computed on pseudobulks by user-provided cluster labels and yield a single weight across all
clusters per each TF-RE-TG triplet. In contrast, the Wilcoxon method groups cells based on the
joint expression of TF, RE and TG in each single cell or in cell aggregates. Cell aggregation uses
a default value of 10 cells and can help overcome sparsity and speed up computation. If cluster
labels are provided, we can obtain cluster-specific weights using the Wilcoxon method.
476
477   1) Wilcoxon test statistics
478   Cells were divided into two groups, with the first group jointly expressing TF gene expression and
479   chromatin accessibility at RE, and the remaining cells in the second group. The default cutoff is 1
480   for normalized gene expression and 0 for normalized chromatin accessibility. There is also an
481   option to use the median of each feature as an adaptive cutoff.
482
2) Correlation
If we consider only TF expression (tf_re.merge set to FALSE), the weight is the correlation
coefficient between the TF gene expression and TG gene expression. If we consider both TF
expression and RE chromatin accessibility (tf_re.merge set to TRUE), the weight is the correlation
coefficient between the product of TF gene expression and RE chromatin accessibility vs the TG
gene expression.
489
490   3) Mutual information between the TF and target gene expression,
491
492   If we consider only TF expression (tf.re_merge is set to FALSE), the weight is the mutual
493   information between the TF gene expression and the TG gene expression. If we consider both
494   TF expression and RE chromatin accessibility (tf.re_merge is set to TRUE), the weight is the
495   mutual information between the product of TF gene expression and RE chromatin accessibility vs
496   the TG gene expression.
497
498   *Calculation of TF activity*
499   The activities for a specific TF in each cell are computed by averaging the weighted expressions
500   of target genes linked to the TF:
501
502

503
$$K_{t,c} = \frac{1}{q_t} \sum_{g \in R_t} \beta_{t,g,r} Y_{g,c}$$

504
505   where $K_{t,c}$ is the activity of a TF t for a cell c, q is the total number of target genes for a TF t, Y_g
506   is the normalized count of target gene g where g is regulated by TF t and $\beta_{t,g,r}$ is the regulatory
507   weight of TF t on target gene g at RE r. R_t is the regulon of TF t.
508
509   If cluster labels are provided, cluster-specific weights are used.
510
511   *Gene set enrichment of regulons*
512   Gene set enrichment of a regulon is performed by testing whether the target genes of a TF are
513   over-represented in known gene signatures such as those provided by MSigDB using a

514 hypergeometric test. Target genes can be refined by filtering the regulons on user-defined
515 weights.
516
Differential TF activity by total activity
This differential analysis compares the differences in the total activity of each transcription factor
between conditions. This analysis is well suited for identifying factors that have contrasting levels
of activities, for instance, lineage factors that are turned on or off during certain developmental
stages or cell types. TF activities are compared between groups using any standard statistical
methods. Here we use scran's findMarkers function to find differential activity between user
provided groups of cells (https://rdrr.io/bioc/scran/man/findMarkers.html).
524
Differential TF activity by network topology
A second approach to investigate differential TF activity is to compare and contrast target genes
or network topology. This is useful when a transcription factor differs in the target genes it
regulates, while maintaining a similar level of total activity. This can happen when a transcription
factor redistributes to a different set of genomic regions, due to mutations in the transcription
factors or changes in the interaction partners. Differences in network topology are calculated by
taking the degree centrality of the edge subtracted graphs between two conditions, with cluster-
specific regulon weights representing edge weights in each condition.
533
$G^{(1)}$ and $G^{(2)}$ are networks with identical node sets N and respective edge sets $E^{(1)}$ and $E^{(2)}$:

- $E^{(1)} = \{e^{(1)}_1, e^{(1)}_2, e^{(1)}_3, ...e^{(1)}_m\}$
- $E^{(2)} = \{e^{(2)}_1, e^{(2)}_2, e^{(2)}_3, ...e^{(2)}_m\}$

where $e^{(j)}_i$ represents the edge weight of the *i*th edge in network $G^{(j)}$.

The edge-subtracted network G' induced by $G^{(1)}$ and $G^{(2)}$ such that each edge e'_i in G' is given by
$e'_i = e^{(1)}_i - e^{(2)}_i$. We simplify the tripartite TF-RE-TG graph to a bipartite TF-TG graph by taking the
maximum of TF-RE-TG weights of the same TF-TG pairs. Degree centrality is the sum of the
weights of the TF-TG e'_i. Degree centrality is further normalized to account for differences in the
number of target genes of each transcription factor. The default normalization method is dividing
degree centrality by the square root of the number of target genes. This strikes a balance
between penalizing TFs with an abundance of target genes and prioritizing TFs with differential
target genes. Transcription factors are ranked by normalized degree centrality.
546
547 *Identification of interaction partners*
548 To identify potentially interacting TF or co-factors, we calculate Jaccard similarity between all
549 pairs of factors and rank factors by the degree of overlap in their regulons. The edges can be
550 filtered on a weight cutoff to retain the most confident set of target genes.
551
552 **Benchmarking**
553
Benchmarking using Reprogram-Seq data

We introduced 4 transcription factors into LNCaP cells - FOXA1, FOXA2, NKX2-1 and GATA6
and obtained paired gene expression and chromatin accessibility information. Because FOXA1
was already highly expressed in LNCaP cells at the basal level, its introduction did not have any
impact on lineage plasticity and therefore FOXA1 was excluded in subsequent analyses. We
focused on NKX2-1 and GATA6 because their expression resulted in distinct cell clusters. Each
cell is identified by the HTO tag corresponding to the well receiving virus encoding a particular TF
and this information serves as the ground truth cell labels. For each TF, a cell is classified into
either expressing this TF or not. AUROC was computed for all the packages being benchmarked
as well as for gene expression of the TF being evaluated. Note that because each TF has different
transduction efficacy and not all cells are able to take up the virus or express the TF, the
theoretical maximum of AUROC will not reach 1.

Benchmarking using AR antagonist data
For the drug treatment dataset, we treated 4 prostate cancer cell lines with 3 different therapeutic
agents and obtained paired gene expression and chromatin accessibility information. Only the 2
AR-dependent cell lines, LNCaP and VCaP, were included in the benchmark since they are
known to respond to AR-targeting agents. Similarly, only AR antagonist, Enzalutamide[1] and AR
degrader ARV-110[3] were used in the benchmark studies since they are known to specifically
inhibit AR activity. Cells treated with Enzalutamide and ARV110 are supposed to show reduced
AR activity compared to cells treated with DMSO control. Each cell line was analyzed separately
by retaining only peaks found in each cell line and all the subsequent analyses were performed
separately for both cell lines. Each cell is identified by the HTO tag corresponding to the well
receiving DMSO or one of the two AR inhibitors and this information serves as the ground truth.
For each of the two drugs (enzalutamide or ARV110), a cell is classified into either treated with
DMSO control or an AR inhibitor. AUROC is computed for all the packages being benchmarked
as well as for AR gene expression. Because TF activity is decoupled from TF expression, we
used the Wilcoxon method to estimate weights with merged public AR ChIP-seq prefiltered for
the presence of motifs as the TF occupancy prior.

We used 5 GRN inference tools (CellOracle, FigR, SCENIC+, *GRaNIE* and *Pando)* to benchmark
the performance of Epiregulon. To ensure consistency, we applied the same gene expression
and chromatin accessibility matrices across all tools. We first used ArchR to determine the peaks
i.e. DNA regions with frequent Tn5 transposase insertion events from the fragment files output by
CellRanger. Then the peak x cell matrix was produced by counting the number of insertions per
peak per cell. From the same ArchR projects we also retrieved normalized gene expression to be
used by Epiregulon, GRaNIE, FigR. For the remaining tools, we used a subset of cells in the gene
expression data which matched the cells in the ArchR projects to make sure that all tools work on
the same cells. Below we describe the workflows used in each tool. We followed the tutorial
examples provided at official web sites with only minor changes to the default settings.

CellOracle
We performed benchmarking against CellOracle[52]. The peak count matrices retrieved from AchR
projects were used to create a CellDataSet object which was subject to dimensionality reduction
and cell aggregation. The ATAC-peaks were first annotated by HOMER to identify transcription

start sites (TSSs). Co-accessibility regions were further identified by the cicero R package using
the run_cicero function. Pairs of peaks that show high co-accessibility (≥0.8) and overlap with a
TSS were used to link peaks to their target genes. Peaks were scanned for the presence of TF
binding motifs using TFinfo.scan. Cell neighborhoods were defined by diffusion maps on the
dimensionality reduced matrix (PCA) and cell clusters were identified using the K nearest
neighbor graph using the Louvain method.

CellOracle models the expression of the target gene as the weighted sum of gene expression of
all the transcription factors it is regulated by.

608
$$Y_g = \sum_t \beta_{t,g} X_t + c$$

where Y_g is the expression of the target gene g, X_t is the expression of the transcription factor t,
β is the fitted coefficient and c is the intersection.

The final activities were calculated using Epiregulon's calculateActivity function using $\beta_{t,g}$ as the
weights. Weights were estimated for each cluster individually and the activities were assembled
together.

FigR
We performed benchmarking against FigR [9]. In the first step of GRN construction the correlation
between peak accessibility and target gene expression was determined using runGenePeakcorr.
We used a non-default search range around TSS (250 kb) to be consistent across all the
benchmarked tools. Correlation coefficients were also computed with background peaks and the
significance of gene-peak association was determined with one-tailed z-test. Only gene-peak
associations that show positive correlation and are statistically significant were retained (p ≤ 0.05).
DORaCs (High density domains of regulatory chromatin) correspond to genes with ≥7 associated
peaks and were calculated by summing the ATAC-seq reads in peaks matched to each gene.
Each TF is associated with a single motif by selecting the motifs most highly correlated with other
motifs of the same TF. Each DORC is evaluated for enrichment of TFs by comparing the match
frequency in its peakset vs the match frequency in a set of background peaks; the p-values of the
enrichment were obtained by z-test ($P_{Enrichment}$). Smoothing of gene expression matrix and
chromatin accessibility data summarized across target genes was performed after determination
of k nearest neighbor cells based on LSI space retrieved from ArchR project and constructed
using ATAC-seq data. The Spearman correlations between the smoothed DORC accessibility
and smoothed TF gene expression were computed ($Correlation$) and their significance were
obtained using z-test ($P_{Correlation}$). From the output of the main function (getFigRGRN) we
retrieved scores indicating the strength of association between transcription factors and target
genes which were computed as below:

$$Regulation\ score = sign(Correlation) * -log10[1 - (1 - P_{Enrichment}) * (1 - P_{Correlation})]$$

The regulation scores were used as weights when calculating activity with Epiregulon.

SCENIC+

We performed benchmarking against SCENIC+ [11]. Briefly, we used pycisTopic which uses Latent Dirichlet Allocation to group regulatory elements into the topics. The model evaluation indicated 20, 25 and 20 as the number of topics for VCaP cells from AR dataset, LNCaP cells from AR dataset, and Reprogram-Seq dataset respectively. The input peak matrix was retrieved from the AchR project. We then used pycistarget to identify TF-region links. Pycistarget identifies motif matches in the peak regions using HOMER, scores each region for motif importance and identifies differentially enriched motifs above background regions. Only regions showing NES > 3.0 and motifs with adjusted p-value <0.05 and log2FC >0.5 were retained. TF-gene importance scores were calculated using gradient-boosting machine regression by predicting TF expression from target gene expression. Region-gene importance scores were calculated using gradient-boosting machine regression by predicting target gene expression from region accessibility. Genes were ranked by TF-gene importance scores and only genes in the leading edge of the gene set enrichment were used for the eRegulon. Gene set enrichment was also performed for region-gene pairs. Peaks were ranked by imputed chromatin accessibility and genes were ranked by gene expression counts per cell. Enrichment score was defined as the AUC at 5% of the ranking and was calculated using AUCell. The enrichment score was used as the activity score in the benchmark assessment.

GRaNIE

We performed benchmarking against GRaNIE [10]. Briefly, GRaNIE overlapped TF binding sites obtained from HOCOMOCO-based TF motifs with ATAC-seq peaks. GRaNIE identified TF-peak connections using Pearson correlation between TF expression and the peak accessibility across samples. The cutoffs for the correlation were chosen based on an empirical FDR calculated from the ratio of TF-peaks in the background peaks over the total number of TF-peaks in both the background and foreground. Peak-gene connections were identified using correlation between the gene expression and chromatin accessibility. All the GRN edges have a weight of 1. The default threshold is FDR < 0.2 for TF-peak links and FDR < 0.1 for peak-gene links. Activities were computed using Epiregulon's calculateActivity.

Pando

We performed benchmarking against Pando[8]. Briefly, ATAC-seq peaks were intersected with PhastCons conserved elements and cCREs derived from ENCODE. TFs present in the 4000 most variable genes were included in the downstream analysis. Motifs for the TFs were obtained from JASPAR2020 and CISBP. Gene expression and chromatin accessibility counts were smoothened by averaging cells within a neighborhood. Pando then models target gene expression as the weighted sum of the product of TF expression and the chromatin accessibility of the region where the TF binds to.

$$Y_g = \sum_t \beta_{t,g,r} X_t A_r + c$$

where Y_g is the expression of the target gene g, X_t is the expression of the transcription factor t, A_r is the chromatin accessibility at region r, β is the fitted coefficient and c is the intersection.

Fitted coefficients were tested for significance using analysis of variance (ANOVA). Only edges
with FDR <0.05 were retained in the final GRN. Activities were calculated using Epiregulon's
calculateActivity function with the fitted coefficients $\beta_{t,g,r}$ as the weights.

Benchmarking using precomputed regulons
Gene expression and chromatin accessibility matrices were obtained from the unpaired PBMC
data[39] and integrated using ArchR's addGeneIntegrationMatrix function. Regulons were
constructed using Epiregulon and benchmarked against regulons obtained from Dorothea[46]) and
Humanbase (hematopoiesis)[48].

These regulons were applied onto a single-cell RNA-seq dataset involving a CRISPRa screen
that overexpressed transcription factors using gRNA in K562 cells[45]. The identity of the gRNA
was used as the ground truth indicating which TF was activated in each cell. Regulon weights
were computed in the K562 dataset, using the correlation method with gRNA as the cluster labels
for all three regulons. ROC was computed using TF activity as the estimates for cluster labels
indicating whether a particular TF is active or not. TF gene expression was also used as the
estimates to generate a separate ROC.

DNN training
Steps for DNN-based TF-RE mapping involved dataset processing, model training, motif scoring
and thresholded GRN construction.

Dataset processing
The reference genome hg38 and the pseudo-bulk ATAC (taken from ArchR) of individual clusters
(C1 or C3) was collected. The whole genome excluding unmappable regions[53] was split into 3072
bp regions. Using the coordinates of each region the corresponding DNA sequence and bigwig
coverage was obtained. The DNA sequence was converted to one-hot encoded form. The DNA
and coverage information were saved in tf records files for faster I/O during training. Training,
validation and test splits were done by chromosomes taking chr8 as test set and chr9 as validation
set.

Model training
We used Basenji2 architecture with 32 base-pair resolution (binned coverage at 32 bp). The
model was trained by randomly sampling 2048bp segment from the input DNA and taking the
corresponding coverage (one cluster per model). We used Poisson NLL as loss, used reverse
complement augmentation and trained for a maximum of 50 epochs with early stopping.

We additionally trained models with base-pair resolution using the ChromBPNet architecture. We
first trained a custom Tn5 sequence bias model for this dataset on non-peak regions. We ensured
that these models learned Tn5 bias by applying DeepSHAP to the model outputs. Using this bias
model to regress out Tn5 sequence bias, we then trained ChromBPNet TF models. These models
were trained on 2114bp sequences centered on ATAC-seq peaks.

Motif scoring
For each of the motif regions (taken from ArchR motif positions file) we obtained the DNA
sequence (denoted as wild type) by centering at the motif and extending to 2048bp. We occluded
the motif by replacing motif nucleotides with 'N' (or 0.25 in one-hot encoding) (denoted as mutant).
We computed the score - mean fraction change in predicted coverage by subtracting mutant from
wild type and normalizing by predicted wild type coverage.
735
736 *Thresholded GRN construction*
737 We computed several quartiles of scores as thresholds. We filtered for those motifs with
738 importance scores higher than the threshold corresponding to bigger differences in prediction.
739 Using the RE remaining we constructed the GRN as before.
740
Reprogram-Seq
Construction of Lentivirus Plasmids for TF Over-expression
All lentivirus plasmids were generated by GenScript. Briefly, the ORF of transcriptional factors
were codon optimized, synthesized and cloned after the hEF1a promoter. The puromycin
resistance gene is driven by a separate Cbh promoter to enable antibiotics selection. Maxi-prep
of each plasmid was performed to maximize the transfection efficiency.
747
Cell Culture and Virus Packaging
LNCaP Clone FGC cells were obtained from ATCC and cultured with RPMI-1640 media with 10%
FBS and 2mM L-Glutamine. The cells were split every 4-5 days to maintain the appropriate
density. 293T cells were cultured in DMEM with 10% FBS, 100μM NEAA, 2mM Glutamine. They
were split every 2-3 days. One day before transfection, the cell culture dish was treated with 5mL
1% gelatin in PBS, incubated for 10 minutes, then aspirated. $3.5X10^6$ 293T cells were seeded
into each 10 cm dish to reach ~80% confluence. On the day of transfection, 20 μl Lipofectamine
2000 was added to 480 μl OptiMEM. In a new tube, the plasmid was mixed in 500μl Opti-MEM at
the following ratio: carrier plasmid, 5 μg; delta8.9, 16 μg; VSVG, 1 μg. Both mixes were combined
and incubated for 20 minutes before adding to the dish. The dish was incubated at 37°C for 6
hours and then 6 ml of complete media was added. Virus was then harvested 48 hours after
transfection. Briefly, all supernatants were harvested and filtered through a 0.45 μm filter bottle.
The virus was concentrated by using the Lenti-X concentrator (TAKARA Bio.) following the
manufacturer's instructions and resuspended in 1 ml 1% BSA in PBS per dish. The concentrated
virus was divided into 200μl aliquots and stored in -80°C until infection.
763
Infection of LNCaP Cells by Lentivirus
Two days before infection, $4X10^5$ LNCaP cells were seeded into each well of a 6-well plate. On
the day of infection, aspirate the media and change to 0.5mL RPMI + 10% FBS + 1X Glutamax
with 8 μg/ml Polybrene. A total of 200 μl concentrated lentivirus was added to each well to achieve
high MOI, and then the plate was centrifuged at 1800 rpm for 45 minutes at room temperature.
After that the plate was put into the incubator for another 3 hours before 2 ml of the full media was
added to each well. Two days after infection, the media was refreshed with 1μg / ml puromycin.
The cells were then grown for another 7 days before harvesting for the single-cell analysis, during
which the cells were split accordingly when the confluence reached 100%.

773
Engineering of dox-inducible NKX2-1 expressing cell lines
774
LNCAP Clone FGC cells were passaged two days before electroporation. 200k cells per reaction
were transduced with 375 ng PBIND-NKX2-1-IRES-eGFP-PGK-puro and 125ng pBO (PiggyBac
transposase) (3:1 vector:transposase ratio) according to manufacturer's instructions for SF Cell
Line 4D-Nucleofector™ X Kit S, using the EN-120 electroporation protocol in 20 µl
Nucleocuvette™ Strip. After electroporation, 100ul of warm media (RPMI + 10% FBS + 1X
Glutamax) was added to the cuvette and the cuvette was placed in the incubator for 1 hour to
recover. Cells were then transferred into a 6 well plate. Puromycin selection (RPMI + 10% FBS +
1X Glutamax, 1ug/ml Puromycin) was performed after 4-5 days, depending on cell recovery.
783

Single-cell RNA-seq and Single-cell RNA-ATAC Co-assay
The single-cell RNA-seq was performed using the Chromium Single Cell 3' kit (V3.1) from 10X
Genomics with cell hashing. Briefly, the cells were trypsinized into single-cell suspension from the
6-well plates, and washed once with PBS with 1% BSA. The cells from different wells were stained
with human TotalSeq-A cell hashing antibodies (BioLegend) containing distinct barcodes
following the manufacturer's protocol. The cells were then washed twice with PBS with 1% BSA
before combining and the live cells were sorted using the SONY SH800S FACS machine. For
loading of the 10X chip G, we overloaded the channel with the aim of recovering 20K cells. The
library construction was following the standard 10X protocol with the following exceptions: 1) the
1µl of 10µM HTO additive primer was spiked in during the cDNA PCR step; 2) during the SPRI
beads cleanup step, the supernatant from the 0.6X cleanup was saved to recover the HTO
fragment. Along with the transcriptome library, the HTO library was amplified from the supernatant
using the HTO-specific primer. The detailed protocol can be found on the cell hashing website
(https://cite-seq.com/cell-hashing/).
798

The single-cell RNA-ATAC co-assay was performed using the 10X Genomics Single Cell
Multiome ATAC + Gene Expression kit. In a previous test, we found that the standard cell
permeabilization protocol used for nuclei preparation does not completely remove the plasma
membrane of the LNCap cells and robust B2M expression can still be detected with FACS (data
not shown). Therefore, we speculated that the same cell hashing strategy would work for the 10X
Multiome assay. Briefly, we followed the same hashing protocol as used in the scRNA-seq
experiment. After the antibody staining and pooling, the cells were permeabilized by following the
nuclei isolation protocol from 10X genomics (10X Nuclei Isolation Protocol). The cells were then
counted and loaded on the Chip J with a recovery aim of 12K cells per lane. For the library
construction, we followed the standard protocol with the following modifications: 1) 1µl of 10µM
HTO additive primer was spiked in during the pre-AMP and cDNA PCR steps; 2) during the 0.6X
cDNA cleanup, the supernatant was saved to amplify the HTO library, as described in the previous
section.
812

813 All the libraries were sequenced on Illumina NextSeq 2000 or NovaSeq 4000. We were aiming
814 for 20K, 40K and 2K raw reads per cell for the transcriptome, ATAC and HTO libraries,
815 respectively.
816

 Synthesis of SMARCA2.4_1

Scheme 1. Preparation of intermediate **E**.

A   **C**   **D**   **E**

*tert-Butyl 4-((trans)-3-(2-(benzyloxy)ethoxy)cyclobutoxy)piperidine-1-carboxylate. (**C**)*. To
a solution of *tert*-butyl 4-((trans)-3-hydroxycyclobutoxy)piperidine-1-carboxylate (**A**, 500 mg, 1.84
mmol, 1.0 eq) in THF (7 mL) was added NaH (60% in mineral oil, 111 mg, 2.77 mmol, 1.5 eq) at
0 °C. The resulting solution was stirred at 0 °C for 0.5 h afterwhich ((2-
bromoethoxy)methyl)benzene (**B**, 592 mg, 2.76 mmol, 1.5 eq) was added. The mixture was
warmed to room temperature and was stirred for an additional 16 h then was quenched with
saturated aqueous NH_4Cl (5 mL). The resulting solution was extracted with EtOAc (10 mL x 3),
and the combined organic layers were dried over sodium sulfate and concentrated under vacuum.
The residue was purified by silica gel chromatography (EtOAc:petroleum ether, 0:100 to 24:76)
to afford the title compound (330 mg, 44%) as a colorless oil. [1]H NMR (300 MHz, DMSO-d_6) δ
7.42-7.23 (m, 5H), 4.49 (s, 2H), 4.25-4.17 (m, 1H), 4.09-4.02 (m, 1H), 3.68-3.61 (m, 2H), 3.57-
3.51 (m, 2H), 3.47-3.37 (m, 2H), 2.95 (t, *J* = 11.0 Hz, 2H), 2.23-1.98 (m, 4H), 1.79-1.67 (m, 2H),
1.39 (s, 9H), 1.35-1.17 (m, 3H). LC-MS: (ES+): m/z 406 [M+H][+].

*tert-Butyl 4-((trans)-3-(2-hydroxyethoxy)cyclobutoxy)piperidine-1-carboxylate. (**D**)*. A
solution of intermediate **C** (330 mg, 0.815 mmol) and 10% Pd/C (66.0 mg) in ethanol (10 mL) and
THF (10 mL) was stirred at room temperature for 1 h under a hydrogen atmosphere (3 atm). The
resulting mixture was filtered, and the filter cake was washed with DCM (10 mL x 2). The filtrate
was concentrated under reduced pressure to afford the title compound (245 mg, 95% yield) as a
white oil. [1]H NMR (300 MHz, DMSO-d_6) δ 4.58 (t, *J* = 5.5 Hz, 1H), 4.26-4.18 (m, 1H), 4.08-4.01
(m, 1H), 3.69-3.61 (m, 2H), 3.53-3.34 (m, 3H), 3.28 (t, *J* = 5.3 Hz, 2H), 2.96 (t, *J* = 11.4 Hz, 2H),
2.24-1.98 (m, 4H), 1.76-1.71 (m, 2H), 1.39 (s, 9H), 1.36-1.18 (m, 2H). LC-MS: (ES+): m/z 316
[M+H][+].

*tert-Butyl 4-((1r,3r)-3-(2-(tosyloxy)ethoxy)cyclobutoxy)piperidine-1-carboxylate (**E**)*. To a
solution of intermediate **D** (245 mg, 0.78 mmol, 1.0 eq) and pyridine (245 mg, 3.11 mmol, 4.0 eq)
in DCM (2 mL) was added TsCl (223 mg, 1.16 mmol, 1.5 eq) at 0 °C. The mixture was then
warmed to room temperature and stirred for 12 h. The solvent was subsequently removed under
vacuum, and the residue was purified by silica gel chromatography (EtOAc:petroleum ether, 0:100
to 29:71) to afford the title compound (303 mg, 83%) as a colorless oil. [1]H NMR (300 MHz,
DMSO-d_6) δ 7.82-7.72 (m, 2H), 7.52-7.41 (m, 2H), 4.22-4.05 (m, 3H), 4.01-3.88 (m, 1H), 3.67-
3.59 (m, 2H), 3.45-3.34 (m, 3H), 2.94 (t, *J* = 11.4 Hz, 2H), 2.41 (s, 3H), 2.13-1.92 (m, 4H), 1.76-
1.64 (m, 2H), 1.37 (s, 9H), 1.33-1.15 (m, 2H). LC-MS: (ES+): m/z 470 [M+H][+].

Scheme 2. Preparation of SMARCA2.4_1 (compound **1)**.

6-Chloro-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridazin-3-amine **(H)**. A solution of 4-bromo-6-chloropyridazin-3-amine (**F**, 2.00 g, 9.60 mmol, 1.0 eq), (1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)boronic acid (**G**, 2.46 g, 12.5 mmol, 1.3 eq), Pd(PPh$_3$)$_4$ (1.12 g, 0.969 mmol, 0.1 eq) and K$_2$CO$_3$ (2.67 g, 19.3 mmol, 2.0 eq) in 1,4-dioxane (20 mL) and water (2 mL) was stirred at 100 °C for 2 h. The reaction was then cooled to room temperature and was quenched with water (20 mL). The resulting solution was extracted with EtOAc (30 mL x 3), and the combined organic layers were dried over sodium sulfate and concentrated under vacuum. The residue was purified by silica gel chromatography (EtOAc:petroleum ether, 0:100 to 58:42) to afford the title compound (2.40 g, 89%) as a red oil. ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.52 (d, *J* = 0.8 Hz, 1H), 8.11 (d, *J* = 0.8 Hz, 1H), 7.65 (s, 1H), 6.41 (s, 2H), 5.46 (dd, *J* = 9.9, 2.2 Hz, 1H), 4.07-3.90 (m, 1H), 3.76-3.57 (m, 1H), 2.24-2.05 (m, 1H), 2.00-1.91 (m, 2H), 1.79-1.64 (m, 1H), 1.60-1.53 (m, 2H). LC-MS: (ES+): m/z 280 [M+H]$^+$.

2-(6-Amino-5-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridazin-3-yl)phenol **(J)**. A solution of intermediate **H** (1.35 g, 4.83 mmol, 1.0 eq), (2-hydroxyphenyl)boronic acid (**I**, 870 mg, 6.31 mmol, 1.3 eq), XphosPdG3 (410 mg, 0.484 mmol, 0.1 eq) and K$_2$CO$_3$ (1.33 g, 9.64 mmol, 2.0 eq) in 1,4-dioxane (11.5 mL) and water (2.5 mL) was stirred at 95 °C for 5 h. The mixture was cooled to room temperature, and the solvent was removed under vacuum. The residue was purified by silica gel chromatography (MeOH:DCM, 0:100 to 5:95) to afford the title compound (900 mg, 55%) as a yellow solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.77 (s, 1H), 8.60 (d, *J* = 0.8 Hz, 1H), 8.26-8.17 (m, 2H), 8.02 (dd, *J* = 8.3, 1.6 Hz, 1H), 7.28-7.23 (m, 1H), 6.97-6.85 (m, 2H), 6.51 (s, 2H), 5.48 (dd, *J* = 9.9, 2.2 Hz, 1H), 3.99-3.90 (m, 1H), 3.74-3.60 (m, 1H), 2.25-2.06 (m, 1H), 2.00-1.95 (m, 2H), 1.76-1.64 (m, 1H), 1.59-1.52 (m, 2H). LC-MS: (ES+): m/z 338 [M+H]$^+$.

2-(6-Amino-5-(1H-pyrazol-4-yl)pyridazin-3-yl)phenol **(K)**. A solution of intermediate **J** (300 mg, 0.890 mmol) in HCl (4.0 M in MeOH, 1 mL) was stirred at room temperature for 2 h. The solvent was then removed under vacuum, and the residue was purified by pre-packed C18 column (solvent gradient: 0-100% MeOH in water (0.1% TFA)) to afford the title compound (188 mg, 83%) as a brown solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.85 (s, 1H), 13.42 (s, 1H), 8.33 (s, 2H), 8.22 (s, 1H), 8.02 (dd, *J* = 8.3, 1.6 Hz, 1H), 7.30-7.24 (m, 1H), 6.95-6.90 (m, 2H), 6.50 (s, 2H). LC-MS: (ES+): m/z 254 [M+H]$^+$.

tert-Butyl   *4-((trans)-3-(2-(4-(3-amino-6-(2-hydroxyphenyl)pyridazin-4-yl)-1H-pyrazol-1-*
yl)ethoxy)cyclobutoxy)piperidine-1-carboxylate (L). A solution of intermediate **K** (100 mg, 0.395
mmol, 1 eq), intermediate **E** (223 mg, 0.475 mmol, 1.2 eq) and K_2CO_3 (136 mg, 0.985 mmol, 2.5
eq) in DMF (1.3 mL) was stirred at 80 °C for 2 h. The reaction mixture was cooled to room
temperature, then was diluted with EtOAc (5 mL) and washed with brine (3 mL x 5). The organic
layer was dried over sodium sulfate and was concentrated under vacuum. The residue was
purified by silica gel chromatography (MeOH:DCM, 0:100 to 5:95) to afford the title compound (60
mg, 27%) as a brown solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.79 (s, 1H), 8.45 (d, *J* = 0.8 Hz,
1H), 8.24-8.17 (m, 2H), 8.03-7.97 (m, 1H), 7.25 (t, *J* = 7.7 Hz, 1H), 6.97-6.87 (m, 2H), 6.48 (s,
2H), 4.30 (t, *J* = 5.4 Hz, 2H), 4.15-4.06 (m, 1H), 4.06-4.00 (m, 1H), 3.69 (t, *J* = 5.4 Hz, 2H), 3.61-
3.53 (m, 2H), 3.24-3.13 (m, 1H), 2.84 (t, *J* = 11.3 Hz, 2H), 2.05-2.01 (m, 4H), 1.66-1.60 (m, 2H),
1.35 (s, 9H), 1.24-1.17 (m, 2H). LC-MS: (ES+): m/z 551 [M+H]$^+$.

2-(6-Amino-5-(1-(2-(trans)-3-(piperidin-4-yloxy)cyclobutoxy)ethyl)-1H-pyrazol-4-
yl)pyridazin-3-yl)phenol, TFA salt (M). A solution of intermediate **L** (60 mg, 0.11 mmol) in TFA
(0.3 mL) and DCM (0.3 mL) was stirred at room temperature for 0.5 h. The solvent was removed
under vacuum to afford the title compound (66 mg, crude) as a yellow solid. ^{1}H NMR (300 MHz,
DMSO-d_6) δ 8.46 (s, 1H), 8.26 (s, 1H), 8.14 (s, 1H), 7.71-7.68 (m, 1H), 7.39-7.33 (m, 1H), 7.03-
6.91 (m, 2H), 4.32 (t, *J* = 5.4 Hz, 2H), 4.20-4.00 (m, 2H), 3.70 (t, *J* = 5.4 Hz, 2H), 3.53-3.45 (m,
1H), 3.12-3.08 (m, 2H), 2.90-2.84 (m, 2H), 2.09-2.05 (m, 4H), 1.91-1.78 (m, 2H), 1.59-1.47 (m,
2H). LC-MS: (ES+): m/z 451 [M+H]$^+$.

(2S,4R)-1-((R)-2-(3-(2-(4-((trans)-3-(2-(4-(3-amino-6-(2-hydroxyphenyl)pyridazin-4-yl)-1H-
pyrazol-1-yl)ethoxy)cyclobutoxy)piperidin-1-yl)ethoxy)isoxazol-5-yl)-3-methylbutanoyl)-4-
hydroxy-N-((S)-1-(4-(4-methylthiazol-5-yl)phenyl)ethyl)pyrrolidine-2-carboxamide (1). A solution
of intermediate **M** (66 mg, crude), intermediate **N** (prepared as described in Berlin, M.; et al.
"PROTACs Targeting BRM (SMARCA2) Afford Selective In Vivo Degradation Over BRG1
(SMARCA4) and are Active in BRG1-Mutant Xenograft Tumor Models", *J. Med. Chem*. accepted
for publication; 85 mg, 0.157 mmol) and NaOAc (54 mg, 0.658 mmol) in MeOH (0.8 mL) was
stirred at room temperature for 1 h. Then NaBH$_3$CN (21.0 mg, 0.33 mmol) was added, and the
mixture was stirred at room temperature for an additional 0.5 h. The solvent was removed under
vacuum, and the residue was purified by Prep-HPLC [column: Xselect CSH F-Phenyl OBD
column, 30*150 mm, 5 μm; mobile Phase A: water (10 mmol/L NH$_4$HCO$_3$), mobile phase B:
CH$_3$CN; flow rate: 60 mL/min; gradient: 43% to 63% B over 8 min; wavelength: 254/220nm; RT:
8.97 min] to afford the title compound (12.8 mg, 12% over two steps) as a white solid. ^{1}H NMR
(300 MHz, DMSO-d_6) δ 13.83 (s, 1H), 9.00-8.99 (m, 1H), 8.51-8.46 (m, 1H), 8.43 (d, *J* = 7.6 Hz,
1H), 8.26-8.19 (m, 2H), 8.05-8.01 (m, 1H), 7.47-7.34 (m, 4H), 7.29-7.24 (m, 1H), 6.98-6.87 (m,
2H), 6.50 (s, 2H), 6.09-5.92 (m, 1H), 5.11-5.02 (m, 1H), 4.96-4.86 (m, 1H), 4.44-4.24 (m, 4H),
4.15 (t, *J* = 5.6 Hz, 2H), 4.11-4.00 (m, 2H), 3.74-3.51 (m, 4H), 3.46-3.43 (m, 1H), 3.14-3.08 (m,
1H), 2.66-2.63 (m, 2H), 2.57-2.53 (m, 2H), 2.46 (s, 3H), 2.31-2.16 (m, 1H), 2.07-1.94 (m, 7H),
1.82-1.74 (m, 1H), 1.67-1.63 (m, 2H), 1.46-1.37 (m, 3H), 1.34-1.20 (m, 2H), 0.98 -0.95 (m, 3H),
0.84-0.79   (m,   3H).   LC-MS: (ES+):   m/z   975   [M+H]$^+$.
928

Measurements of SMARCA2/4 degradation using In-cell Western

22Rv1 cells were seeded at 8000/well in 96-well black/clear-bottom plates with 180ul DMEM
containing 1% pen-strep, 1% HEPES and 10% FBS and incubated overnight at 37 °C to allow
adherence. The next morning cells were treated with 20 uL of 10× compound and incubated for
an additional 20 h. Cells were then washed with ice cold DPBS, then 50 uL of ice cold 4% PFA
(Electron Microscopy Sciences 15711)/DPBS was added, and plates were then incubated at RT
for 20 min. PFA was then removed and 200 uL of TBS-T containing 0.5% Triton X 100 (Sigma
T8787) was added. Plates were incubated at RT for 30 min. TBS-T/Triton X was then removed,
50 uL of Li-cor blocking solution (Li-Cor 927-50003) was added, and plates were incubated at RT
for 1 h. Blocking solution was removed, 50 uL of Li-Cor blocking solution containing primary
antibody to SMARCA2 (1:2000) and α-Tubulin (1:2000) or SMARCA4 (1:1000) and α-Tubulin
(1:2000) was added, and plates were incubated at 4 °C overnight. The next day, plates were
washed 3× with TBS-T and 50uL of secondary antibody cocktail in Li-Cor blocking solution was
added (IRDye 800CW Goat anti-rabbit IgG, Li-Cor 926-32211 and IRDye 680RD Goat anti-mouse
IgG, Li-Cor 926-68070; both secondary antibodies are diluted 1:5000). Plates were incubated at
RT for 1 h protected from light. Then plates were washed twice with TBS-T and excess liquid
removed. Plates were read on the Li-Cor Odyssey with default intensity setting of 5.0 for both
channels. Li-Cor images were analyzed with the in-cell Western feature of Image Studio Lite.
Assays were run with technical duplicates and multiple biological replicates (>2) with error
calculated with GraphPad prism using 95% confidence interval.
949

Acknowledgments

We would like to thank Xinpeng Wang, Fang Zhang, Mingtao He, Yunxing Cheng and Michael
Berlin (Arvinas) for the synthesis of SMARCA2.4_1 and Peter Dragovich for guiding the synthesis.
We would like to thank Jayaram Kancherla for his help and advice.

Figure legends
Figure 1. Figure constructs GRNs to infer TF activity at the single cell level
Schematics of Epiregulon. RNA and ATAC counts of similar cells are first aggregated. Based on
the correlation between ATAC and RNA, each target gene (TG) is assigned a regulatory element
(RE) within a defined distance (default ±250kb). Each RE is interrogated for possible TF
occupancy using ChIP-seq data from ENCODE and ChIP-Atlas. An optional network pruning step
retains TF-RE-TG triples that co-occur within the same cells. Co-occurrence is defined by TF and
TG expression and chromatin accessibility exceeding a threshold. The regulatory potential of TF
for each TG is estimated using 1 of the 3 methods (correlation, mutual information and Wilcoxon)
and this serves as the weight of each TF-TG pair. The weights are computed for all the cells
and/or for cells in each cluster to yield cluster-specific weights. The TF activity is computed as the
weighted sum of the target gene expression normalized by the number of genes with non-zero
edges. The regulon of each TF can be annotated with gene set enrichment. Jaccard similarity
between the target genes is used to identify transcription factors that potentially interact.
Differential network is performed on edge-subtracted networks of two cell states.
970

972 **Figure 2: Epiregulon constructs GRNs to infer TF activity at the single cell level**

a) UMAP representation of LNCaP cells transduced with virus encoding GATA6 and NKX2-1. The cells were infected in individual wells, hashtagged with HTO and then pooled into a single run.
b) Shown is the level of TF gene expression or TF activity computed by Epiregulon for GATA6 and NKX2-1
c) Each cell was identified by the HTO tag corresponding to the well receiving virus encoding a particular TF and this information serves as the ground truth. For each TF, a cell was classified into either expressing this TF or not based on the TF activity. AUROC was computed for all the packages being benchmarked as well as for gene expression of the TF being evaluated. Note that because each TF had different transduction efficacy and not all cells were able to take up the virus or express the TF, the theoretical maximum of AUROC will not reach 1.
d) Number of TFs retrieved from the different packages being benchmarked
e) As a validation, LNCaP cells were transduced with dox-inducible constructs expressing either NKX2-1 or eGFP control. Correlation between weights of NKX2-1 targets (computed by the Wilcoxon method) against logFC of NKX2-1 targets between no dox and dox treatments.
f) Genes most similar to GATA6 were ranked by Jaccard similarity between the GATA6 regulon and regulons of all the other TFs.
g) Chromatin accessibility at neuroendocrine (NE) - and stem cell like (SCL)- specific regions defined by Tang et al was computed by ChromVAR
h) Shown are the correlation between GATA6 activity and chromatin accessibility at regions specific to stem cell like (SCL) subtype of prostate cancer, and the correlation between NKX2-1 activity and chromatin accessibility at neuroendocrine (NE) subtype of prostate cancer.

Figure 3: Epiregulon predicts responses towards TF-targeting drugs
a) UMAP representation of scMultiome data of 4 prostate cancer cell lines treated with 3 drug compounds. VCaP and LNCaP are AR-dependent whereas DU145 and H660 are AR-independent. The 3 drugs include an AR antagonist (enzalutamide), an AR degrader (ARV110) and a degrader of SMARCA2/4, the ATPase subunit of the SWI/SNF remodeling complex (SMARCA2.4_1). Enzalutamide and ARV110 are known and specific inhibitors of AR activity and were used for benchmarking.
b) AR activity computed by Epiregulon.
c) Each cell was identified by the HTO tag corresponding to the well receiving DMSO or one of the two AR inhibitors and this information served as the ground truth. For each of the two drugs (enzalutamide or ARV110), a cell was classified into either treated with DMSO control or an AR inhibitor based on AR activity. AUROC was computed for all the packages being benchmarked as well as for AR gene expression.
d) AR expression of LNCaP and VCaP cells treated with the 3 drugs
e) Number of TFs retrieved by each of the packages

Figure 4: Comparison of ChIP-seq and motif annotations

a) Each cell was identified by the HTO tag corresponding to the well receiving virus encoding a particular TF and this information served as the ground truth cell labels. For each TF, a cell was classified into either expressing this TF or not. GATA6 ChIP-seq was obtained by merging ChIP-seq from ChIP-atlas and ENCODE. Merged ChIP + motif refers to merged ChIP-seq peaks that contain GATA6 motifs. We trained a DNN model on the cluster of cells expressing GATA6 using Basenji2 and predicted an importance score for each motif based on the difference between the original sequence and motif occluded sequence. We filtered for those motifs with importance scores higher than quartiles of scores.

b) Same as a) but computed for NKX2-1 using Nkx2.1.Homeobox_182 as the motif.

c) Each cell was identified by the HTO tag corresponding to the well receiving DMSO or the AR degrader ARV110 and this information serves as the ground truth cell labels. Public ChIP-seq was derived from merged AR ChIP-seq from ChIP-atlas and ENCODE. Motif was derived from the AR motif from CisBP. Merged ChIP-seq + motif refers to merged public ChIP-seq peaks that contain AR motifs. We trained a DNN model on cells treated with DMSO using Basenji2 and predicted an importance score for each motif based on the difference between the original sequence and motif occluded sequence. We filtered for those motifs with importance scores higher than quartiles of scores.

Figure 5: Epiregulon predicts responses towards chromatin-modifying drugs
   a) SMARCA4 activity computed by Epiregulon
   b) Hypergeometric enrichment of gene sets (hallmark and oncogenic signature) of the target genes of SMARCA4 inferred by Epiregulon for each of the 4 prostate cancer cell lines. VCaP and LNCaP are AR-independent and DU145 and H660 are AR-independent.
   c) Potential interaction partners of SMARCA4 were nominated using Jaccard similarity between its target genes and the target genes of all other transcription factors. The top interacting genes are reflective of the unique etiologies of the different prostate cancer subtypes.

Figure 6: Epiregulon-constructed GRNs predict effects of perturbation
   a) UMAP representation of PBMC data with unpaired and integrated scATAC-seq and scRNA-seq for which Epiregulon was used to compute the GRN.
   b) Representation of TF expression or activity inferred by Epiregulon of well-known lineage factors (EBF1, PAX5 - B cells, GATA3 - T cells and SPI1- myeloid cells). Weights were estimated using the Wilcoxon method.
   c) Cell-type specific GRN weights were computed for PreB and CD4 memory T cells (CD4.M) respectively. The GRN weights are cell type aware since only PreB cells, not CD4 memory T cells, have positive weights in the target genes of the B cell lineage factor, EBF1. Weight cutoff was >0.
   d) EBF1 was interrogated for potential interaction partners using Jaccard similarity between its target genes and the target genes of all other transcription factors. The top genes (PAX5, IRF4 and TCF3 (also known as E2A)) are known to interact with EBF1 in B cell development.

e) Differential networks between monocytes and CD4 memory T cells were computed by
subtracting the edges of the cell type specific GRNs. Genes were ranked by degree
centrality of the edge-subtracted graph. SPI1 and CEBPA are well known lineage factors
of the myeloid lineage whereas TBX21 and GATA3 are well known lineage factors of the
T cell lineage.
f) The GRN computed from the PBMC data was applied onto a CRISPRa dataset of K562
cells containing only gene expression data (no scATAC-seq). Each cell was identified by
the guide RNA corresponding to a TF that was being activated and this information serves
as the ground truth. For each TF, a cell was classified into containing either the TF
activating guide or the non-targeting control based on its TF activity. Shown is the area
under the receiver operating curves (AUROC) using either TF gene expression or TF
activity computed from GRN constructed from the PBMC dataset as the classifier. AUROC
is also calculated for activity inferred from other GRNs (Dorothea, CollecTRI or
humanbase).
g) TF activity inferred by epiregulon could better classify PRDM1-activated cells from control
cells than gene expression or TF activity estimated from the Dorothea regulons. TF activity
avoids zero inflation as seen with single cell gene expression.
1077
1078
Figure S1
a) AUROC of the reprogram-seq dataset was computed with each of the 3 weight methods
offered by Epiregulon. Correlation (corr) and mutual information (MI) weights are
computed either using 1) transcription factor (TF) expression vs target gene (TG)
expression or 2) TF expression multiplied with the chromatin accessibility at regulatory
elements (RE) vs TG expression. Each cell is identified by the HTO tag corresponding to
the well receiving virus encoding a particular TF and this information serves as the ground
truth. For each TF, a cell is classified into either expressing this TF or not.
b) Differences in rank between target and non-target genes with respect to GATA6 and
NKX2-1 transcription factors. Gene rank was created from the product of two ranks: one
based on the differences in the mean expression between the treatment and control
groups, and the second based on the proportion of the total gene expression across all
the cells contributed by the treated cells.
c) Representation of target genes of NKX2-1 and GATA6. C1 cluster contains cells infected
with GATA6, C3 contains cells infected with NKX2-1 and C5 contains uninfected cells.
Edges with weight > 0 were retained and target genes of NKX2-1 and GATA6 were shown
for each cluster.
d) As a validation, LNCaP cells were transduced with dox-inducible constructs expressing
either NKX2-1 or eGFP control. Shown are the putative target genes of NKX2-1 ranked
by weights.
e) Representation of target genes of YAP1 and GATA6 with C1-specific weights.
f) Chromatin accessibility at adenocarcinoma- and neuroendocrine-specific regions defined
by Baca et al was computed by ChromVAR
1102
1103 **Figure S2**

1104   a) Chemical structure of SMARCA2.4_1, a degrader of SMARCA2 and SMARCA4

1105   b) Levels of SMARCA2 and SMARCA4 after SMARCA2.4_1 treatment. SMARCA2 DC50 =
1106   0.22nM, SMARCA4 DC50 = 0.31 nM in 22Rv1 cells.

1107   c) AUROC of the AR dataset was computed with each of the 3 weight methods offered by
1108   Epiregulon. Correlation (corr) and mutual information weights are computed using either
1109   1) transcription factor (TF) expression vs target gene (TG) expression or 2) TF expression
1110   multiplied with the chromatin accessibility at regulatory elements (RE) vs TG expression.

1111   d) TF activity was computed with either AR ChIP-seq merged across ENCODE and ChIP-
1112   atlas or with LNCaP or VCaP AR and FOXA1 ChIP-seq (cell line-matched). There is a
1113   slight discrepancy between the cell culture condition used to obtain scMultiome data (in
1114   this manuscript) vs the condition used to obtain cell line-matched ChIP-seq. The cell line-
1115   matched ChIP-seq was obtained in cell lines cultured in charcoal stripped medium
1116   supplemented with R1881 whereas the scMultiome data was performed in cell lines
1117   cultured in 10% fetal bovine serum (FBS) but the identities of the cell lines are matched.

1118

1119   **Figure S3**

1120   a) Same as Figure 4a but performed with ChromBPNet

1121   b) Same as Figure 4b but performed with ChromBPNet

1122   c) To evaluate whether GRN weights improves performance of TF activity, weights of target
1123   genes were randomly permuted and AUROC is computed with GRNs with permuted
1124   weights

1125   d) Same as S3c) but performed on NKX2-1

1126

1127   **Figure S4**

1128   a) AUROC of the AR dataset was computed with each of the 3 weight methods offered by
1129   Epiregulon. Correlation (corr) and mutual information (MI) weights are computed either
1130   using transcription factor (TF) expression vs target gene (TG) expression or TF expression
1131   multiplied with the chromatin accessibility at regulatory elements (RE) vs TG expression.
1132   Each cell is identified by the HTO tag corresponding to the well receiving DMSO or the
1133   SMARCA2/4 degrader SMARCA2.4_1 and this information serves as the ground truth. A
1134   cell is classified into either treated with DMSO control or SMARCA2.4_1 based on the
1135   SMARCA4 activity inferred by Epiregulon.

1136   b) TF activity was computed with either SMARCA4 ChIP-seq merged across ENCODE and
1137   ChIP-atlas or with VCaP SMARCA4 ChIP-seq (cell line-matched). There is a slight
1138   discrepancy between the cell culture condition used to obtain scMultiome data (in this
1139   manuscript) vs the condition used to obtain cell line-matched ChIP-seq. The cell line-
1140   matched ChIP-seq was obtained in cell lines cultured in charcoal stripped medium
1141   supplemented with R1881 whereas the scMultiome data was performed in cell lines
1142   cultured in 10% fetal bovine serum (FBS) but the identities of the cell lines are matched.

1143

1144   **Figure S5**

1145   a) Gene set enrichment of regulons inferred by Epiregulon (Wilcoxon weight cutoff > 0).
1146   EBF1 and PAX5 are B cell lineage factors. GATA3 is a lymphoid lineage factor. SPI1 is a
1147   myeloid lineage factor.

1148   b) Shown is the area under the receiver operating curves (AUROC) using either TF gene
1149     expression or TF activity computed from GRN obtained from humanase, collecTRI or
1150     Dorothea.
1151

1152 **Figure S6**
1153   a) We used the scMultiome package to simulate paired scRNA-seq and scATAC-seq data
1154     (see Supplementary Note). Briefly, we provided a lineage tree that specified the identities
1155     of the individual cells and a ground truth GRN that specified the regulatory relationships
1156     between TF-TG and RE-TG. Gene expression and peak matrices were generated taking
1157     into account both cell identities and gene regulatory relationships. We obtained the ground
1158     truth activity of each TF by summing the differences in target gene expression as defined
1159     by the GRN. For all assessments from b-e, we use the fixed cutoff and tf_re.merge set to
1160     TRUE.
1161   b) We varied the degree of sparsity by increasing the technical noise and evaluated the
1162     correlation of activity against ground truth TF activity across all genes for each of the
1163     weight methods. Each data point represents a TF. The ATAC effect was set at basic. GRN
1164     had 100% true connections.
1165   c) We added noise to GRN by adding false connections and removing true connections, and
1166     correlated the activity against ground truth activity for each TF. The ATAC effect was set
1167     at basic and data sparsity was set at medium.
1168   d) Cell clusters were first defined using k-means clustering with a median cluster size of 10
1169     cells for the Wilcoxon and logFC methods. We correlated the activity against ground truth
1170     activity for each TF. The ATAC effect was set at basic and data sparsity was set at
1171     medium. GRN had 100% true connections.
1172   e) We varied the effect of ATAC on the gene expression, with the strong ATAC effect
1173     indicating a greater importance of ATAC counts on shaping gene expression counts. GRN
1174     had 100% true connections. True counts were generated without data sparsity and
1175     observed counts were generated with medium data sparsity.
1176   f) We spiked in 100% false connections to the true GRN. We examined the recall and
1177     precision of true GRN connections with varying degrees of z score cutoffs computed by
1178     chi-square statistics
1179   g) We spiked in 100% false connections to the true GRN. We show that true connections
1180     have increased RE-TG correlations and this translated to higher weights in the true
1181     connections.
1182   h) We evaluated the effects of pruning on the accuracy of TF activity predictions. We
1183     computed the ground truth TF activity from the predefined GRN with high data sparsity
1184     and strong ATAC effects. We spiked in 100% false connections and compared the
1185     correlation between the TF activity inferred from either the unpruned or pruned GRN vs
1186     the ground truth GRN. Shown is the difference between the two correlations. A positive
1187     difference indicates improved performance due to pruning.
1188

1189 **Figure S7**
1190   a) We added noise to the network by increasing proportions of randomly permuted edges on
1191     top of true differential edges (15% of total edges). Centrality subtraction was computed by

1192   subtracting the degree centralities of two networks whereas edge subtraction was
1193   computed by centrality of the edge subtracted network. See methods for details.
1194   b) We varied the signal strength by increasing the proportions of altered edges (enrichment,
1195   depletion or permutations) with noise fixed at 10% of total edges.

1196

Code availability

1198

Epiregulon: https://github.com/xiaosaiyao/epiregulon
Epiregulon.extra: https://github.com/xiaosaiyao/epiregulon.extra
Epiregulon.archr: https://github.com/xiaosaiyao/epiregulon.archr
ScMultiome: https://github.com/xiaosaiyao/scMultiome
Epiregulon benchmarking: https://github.com/xiaosaiyao/epiregulon.benchmark
Deep neural network model: https://github.com/xiaosaiyao/epiregulon.sequence.modeling
Differential network simulation: https://github.com/xiaosaiyao/epiregulon.diffnetwork.simulation

1   Tran, C. *et al.* Development of a second-generation antiandrogen for treatment of
advanced prostate cancer. *Science* **324**, 787-790, doi:10.1126/science.1168175 (2009).
2   Sakamoto, K. M. *et al.* Protacs: chimeric molecules that target proteins to the Skp1-Cullin-
F box complex for ubiquitination and degradation. *Proc Natl Acad Sci U S A* **98**, 8554-
8559, doi:10.1073/pnas.141230798 (2001).
3   Snyder, L. B. *et al.* Discovery of ARV-110, a first in class androgen receptor degrading
PROTAC for the treatment of men with metastatic castration resistant prostate cancer.
Cancer Res **81**, 43 (2021).
4   Hagenbeek, T. J. *et al.* An allosteric pan-TEAD inhibitor blocks oncogenic YAP/TAZ
signaling and overcomes KRAS G12C inhibitor resistance. *Nat Cancer* **4**, 812-828,
doi:10.1038/s43018-023-00577-0 (2023).
5   Badia, I. M. P. *et al.* Gene regulatory network inference in the era of single-cell multi-
omics. *Nat Rev Genet*, doi:10.1038/s41576-023-00618-5 (2023).
6   Alvarez, M. J. *et al.* Functional characterization of somatic mutations in cancer using
network-based inference of protein activity. *Nat Genet* **48**, 838-847, doi:10.1038/ng.3593
(2016).
7   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat
Methods* **14**, 1083-1086, doi:10.1038/nmeth.4463 (2017).
8   Fleck, J. S. *et al.* Inferring and perturbing cell fate regulomes in human brain organoids.
Nature **621**, 365-372, doi:10.1038/s41586-022-05279-8 (2023).
9   Kartha, V. K. *et al.* Functional inference of gene regulation using single-cell multi-omics.
Cell Genom **2**, doi:10.1016/j.xgen.2022.100166 (2022).
10   Kamal, A. *et al.* GRaNIE and GRaNPA: inference and evaluation of enhancer-mediated
gene regulatory networks. *Mol Syst Biol* **19**, e11627, doi:10.15252/msb.202311627
(2023).
11   Bravo Gonzalez-Blas, C. *et al.* SCENIC+: single-cell multiomic inference of enhancers and
gene regulatory networks. *Nat Methods* **20**, 1355-1367, doi:10.1038/s41592-023-01938-
4 (2023).

12   Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell
chromatin accessibility analysis. *Nat Genet* **53**, 403-411, doi:10.1038/s41588-021-00790-
6 (2021).
13   Inukai, S., Kock, K. H. & Bulyk, M. L. Transcription factor-DNA binding: beyond binding
site motifs. *Curr Opin Genet Dev* **43**, 110-119, doi:10.1016/j.gde.2017.02.007 (2017).
14   Worsley Hunt, R. & Wasserman, W. W. Non-targeted transcription factors motifs are a
systemic component of ChIP-seq datasets. *Genome Biol* **15**, 412, doi:10.1186/s13059-
014-0412-4 (2014).
15   Duan, J. *et al.* Rational Reprogramming of Cellular States by Combinatorial Perturbation.
Cell Rep **27**, 3486-3499 e3486, doi:10.1016/j.celrep.2019.05.079 (2019).
16   Baca, S. C. *et al.* Reprogramming of the FOXA1 cistrome in treatment-emergent
neuroendocrine prostate cancer. *Nat Commun* **12**, 1979, doi:10.1038/s41467-021-22139-
7 (2021).
17   Han, M. *et al.* FOXA2 drives lineage plasticity and KIT pathway activation in
neuroendocrine prostate cancer. *Cancer Cell* **40**, 1306-1323 e1308,
doi:10.1016/j.ccell.2022.10.011 (2022).
18   Tang, F. *et al.* Chromatin profiles classify castration-resistant prostate cancers suggesting
therapeutic targets. *Science* **376**, eabe1505, doi:10.1126/science.abe1505 (2022).
19   Xiao, L. *et al.* Targeting SWI/SNF ATPases in enhancer-addicted prostate cancer. *Nature*
601, 434-439, doi:10.1038/s41586-021-04246-z (2022).
20   Kelley, D. R. Cross-species regulatory sequence activity prediction. *PLoS Comput Biol* **16**,
e1008050, doi:10.1371/journal.pcbi.1008050 (2020).
21   Avsec, Z. *et al.* Base-resolution models of transcription-factor binding reveal soft motif
syntax. *Nat Genet* **53**, 354-366, doi:10.1038/s41588-021-00782-6 (2021).
22   Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-
range interactions. *Nat Methods* **18**, 1196-1203, doi:10.1038/s41592-021-01252-x (2021).
23   Cyrta, J. *et al.* Role of specialized composition of SWI/SNF complexes in prostate cancer
lineage plasticity. *Nat Commun* **11**, 5549, doi:10.1038/s41467-020-19328-1 (2020).
24   Beltran, H. *et al.* Divergent clonal evolution of castration-resistant neuroendocrine prostate
cancer. *Nat Med* **22**, 298-305, doi:10.1038/nm.4045 (2016).
25   Ku, S. Y. *et al.* Rb1 and Trp53 cooperate to suppress prostate cancer lineage plasticity,
metastasis, and antiandrogen resistance. *Science* **355**, 78-83,
doi:10.1126/science.aah4199 (2017).
26   Bluemn, E. G. *et al.* Androgen Receptor Pathway-Independent Prostate Cancer Is
Sustained through FGF Signaling. *Cancer Cell* **32**, 474-489 e476,
doi:10.1016/j.ccell.2017.09.003 (2017).
27   Mesman, S. & Smidt, M. P. Tcf12 Is Involved in Early Cell-Fate Determination and Subset
Specification of Midbrain Dopamine Neurons. *Front Mol Neurosci* **10**, 353,
doi:10.3389/fnmol.2017.00353 (2017).
28   Liu, W. *et al.* BRD4 regulates Nanog expression in mouse embryonic stem cells and
preimplantation embryos. *Cell Death Differ* **21**, 1950-1960, doi:10.1038/cdd.2014.124
(2014).
29   Choi, H. I. *et al.* BET inhibitor suppresses migration of human hepatocellular carcinoma
by inhibiting SMARCA4. *Sci Rep* **11**, 11799, doi:10.1038/s41598-021-91284-2 (2021).
30   Rahman, S. *et al.* The Brd4 extraterminal domain confers transcription activation
independent of pTEFb by recruiting multiple proteins, including NSD3. *Mol Cell Biol* **31**,
2641-2652, doi:10.1128/MCB.01341-10 (2011).
31   Xu, X. *et al.* Bromodomain Containing Protein 4 (BRD4) Regulates Expression of its
Interacting Coactivators in the Innate Response to Respiratory Syncytial Virus. *Front Mol*
Biosci **8**, 728661, doi:10.3389/fmolb.2021.728661 (2021).

32 Alver, B. H. *et al.* The SWI/SNF chromatin remodelling complex is required for
maintenance of lineage specific enhancers. *Nat Commun* **8**, 14648,
doi:10.1038/ncomms14648 (2017).
33 Blumli, S. *et al.* Acute depletion of the ARID1A subunit of SWI/SNF complexes reveals
distinct pathways for activation and repression of transcription. *Cell Rep* **37**, 109943,
doi:10.1016/j.celrep.2021.109943 (2021).
34 Sobczak, M., Pitt, A. R., Spickett, C. M. & Robaszkiewicz, A. PARP1 Co-Regulates
EP300-BRG1-Dependent Transcription of Genes Involved in Breast Cancer Cell
Proliferation and DNA Repair. *Cancers (Basel)* **11**, doi:10.3390/cancers11101539 (2019).
35 Shang, X. Y. *et al.* ARID1A deficiency weakens BRG1-RAD21 interaction that jeopardizes
chromatin compactness and drives liver cancer cell metastasis. *Cell Death Dis* **12**, 990,
doi:10.1038/s41419-021-04291-6 (2021).
36 Lazar, J. E. *et al.* Global Regulatory DNA Potentiation by SMARCA4 Propagates to
Selective Gene Expression Programs via Domain-Level Remodeling. *Cell Rep* **31**,
107788, doi:10.1016/j.celrep.2020.107788 (2020).
37 Marino, M. M. *et al.* Interactome mapping defines BRG1, a component of the SWI/SNF
chromatin remodeling complex, as a new partner of the transcriptional regulator CTCF. *J*
Biol Chem **294**, 861-873, doi:10.1074/jbc.RA118.004882 (2019).
38 Liao, L. *et al.* Multiple tumor suppressors regulate a HIF-dependent negative feedback
loop via ISGF3 in human clear cell renal cancer. *eLife* **7**, e37925, doi:10.7554/eLife.37925
(2018).
39 Granja, J. M. *et al.* Single-cell multiomic analysis identifies regulatory programs in mixed-
phenotype acute leukemia. *Nat Biotechnol* **37**, 1458-1465, doi:10.1038/s41587-019-0332-
7 (2019).
40 Ungerback, J., Ahsberg, J., Strid, T., Somasundaram, R. & Sigvardsson, M. Combined
heterozygous loss of Ebf1 and Pax5 allows for T-lineage conversion of B cell progenitors.
J Exp Med **212**, 1109-1123, doi:10.1084/jem.20132100 (2015).
41 Yeamans, C. *et al.* C/EBPalpha binds and activates the PU.1 distal enhancer to induce
monocyte lineage commitment. *Blood* **110**, 3136-3142, doi:10.1182/blood-2007-03-
080291 (2007).
42 Thomas, G. D. *et al.* Deleting an Nr4a1 Super-Enhancer Subdomain Ablates Ly6C(low)
Monocytes while Preserving Macrophage Gene Function. *Immunity* **45**, 975-987,
doi:10.1016/j.immuni.2016.10.011 (2016).
43 Kanhere, A. *et al.* T-bet and GATA3 orchestrate Th1 and Th2 differentiation through
lineage-specific targeting of distal regulatory elements. *Nat Commun* **3**, 1268,
doi:10.1038/ncomms2260 (2012).
44 Newton, R. H. *et al.* Maintenance of CD4 T cell fitness through regulation of Foxo1. *Nat*
Immunol **19**, 838-848, doi:10.1038/s41590-018-0157-4 (2018).
45 Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and
Activation. *Cell* **159**, 647-661, doi:10.1016/j.cell.2014.09.029 (2014).
46 Garcia-Alonso, L., Holland, C. H., Ibrahim, M. M., Turei, D. & Saez-Rodriguez, J.
Benchmark and integration of resources for the estimation of human transcription factor
activities. *Genome Res* **29**, 1363-1375, doi:10.1101/gr.240663.118 (2019).
47 Müller-Dott, S. *et al.* Expanding the coverage of regulons from high-confidence prior
knowledge for accurate estimation of transcription factor activities. *bioRxiv*,
2023.2003.2030.534849, doi:10.1101/2023.03.30.534849 (2023).
48 Greene, C. S. *et al.* Understanding multicellular function and disease with human tissue-
specific networks. *Nat Genet* **47**, 569-576, doi:10.1038/ng.3259 (2015).
49 Gaublomme, J. T. *et al.* Nuclei multiplexing with barcoded antibodies for single-nucleus
genomics. *Nat Commun* **10**, 2907, doi:10.1038/s41467-019-10756-2 (2019).

50   Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
1340   transcription-factor-associated accessibility from single-cell epigenomic data. *Nat*
1341   *Methods* **14**, 975-978, doi:10.1038/nmeth.4401 (2017).
1342   51   Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor
1343   sequence specificity. *Cell* **158**, 1431-1443, doi:10.1016/j.cell.2014.08.009 (2014).
1344   52   Kamimoto, K. *et al.* Dissecting cell identity via network inference and in silico gene
1345   perturbation. *Nature* **614**, 742-751, doi:10.1038/s41586-022-05688-9 (2023).
1346   53   Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of
1347   Problematic Regions of the Genome. *Sci Rep* **9**, 9354, doi:10.1038/s41598-019-45839-z
1348   (2019).
1349

Figure 1: Epiregulon constructs GRNs to infer TF activity at the single cell level

Figure 2: Epiregulon constructs GRNs to infer TF activity at the single cell level

Figure 3: Epiregulon predicts responses towards TF-targeting drugs

Figure 4: Comparison of ChIP-seq and motif annotations

Figure 5: Epiregulon predicts responses towards chromatin-modifying drugs

Figure 6: Epiregulon-constructed GRNs predict effects of perturbations