Title: Pathogenic IgE-fated B cell memory retains functional plasticity

Authors: Kelly Bruton,[1†‡*], Allyssa Phelps,[1‡] Atai Ariaz[1], Tina D. Walker[1], Jianping Wen[1],

Sharon S. Khavkine-Binstock[1], Allison Fang[1], Olivia Mann-Delany[1], Niels Peter H. Knudsen[2],

Siyon Gadkar[1], Emily Grydziuszko[1], Joshua F. E. Koenig[1], Aidan Gagnon[1], Susan Waserman[1],

Peter S. Andersen[2], Manel Jordana[1]

Affiliations:

[1]Schroeder Allergy & Immunology Research Institute and Department of Medicine (Division of

Clinical Immunology & Allergy), McMaster University; Hamilton, Ontario, CA.

[2]ALK-Abelló A/S; Hørsholm, DK

[†]Current Address: Department of Pathology, Stanford University School of Medicine; Stanford,

California, USA.

[‡]These authors contributed equally to this work.

[*]Corresponding author. Email: brutonk@stanford.edu

Abstract: Long-lived immunoglobulin (Ig) E responses against innocuous environmental and

dietary antigens (Ags) are maintained by an IgG1-dominant memory B cell (MBC) compartment

primed for IL-4 responsiveness. In this work, we investigated the plasticity of the MBC

compartment fated for IgE class switch recombination upon Ag re-exposure. Antibody-mediated

IL-4Rα blockade augmented the germinal center response and uncovered an IL-4/IL-13

dependency within the type 2 memory B cell (MBC2) skewed phenotype. In the absence of IL-

4/IL-13 signaling during the recall response, a long-lived Ag-specific IgG2c MBC population

emerged, shifting the MBC response away from a type 2 phenotype and towards type 1. The

emergence of this IgG2c response was dependent on IFN-γ signaling and arose from both

unswitched and class-switched Ag-specific B cells *in vivo*. We further demonstrated that a type

1-polarizing adjuvant can augment the redirection of type 2-polarized class-switched MBCs,

suggesting with the correct cues pathogenic MBCs can be functionally reprogrammed.

One Sentence Summary: B cell responses to allergens can be reprogrammed away from a

pathogenic fate through IL-4/IL-13 signaling blockade.

Main Text:

INTRODUCTION

Classical immunological memory is marked by the propensity for rapid and optimized responses

upon re-exposure to the target antigen (Ag). Within the B cell compartment, immunological

memory is distinguished by high-affinity binding to Ag, a class-switched constant region

conferring specialized effector functions, and altered cellular processes (*e.g.,* metabolism).

Following vaccination, for example, viral exposure rapidly reactivates memory B cells (MBCs)

facilitating plasma cell differentiation and boosting of neutralizing IgG titers, lessening the

opportunity for infection (*1–3*). This same process, however, is pathogenic when perpetuating

inflammatory responses to innocuous Ag, as occurs in allergic disease.

Allergies categorized as type I hypersensitivities are mediated by IgE. Humoral IgE responses

are relatively short-lasting, yet reactivity to some allergens (*e.g.,* peanuts) can persist for a

lifetime (*4–8*). Central to this persistence are $IgG1^+$ MBCs which undergo sequential class-

switching to replenish the transient IgE plasma cell pool (*5, 9–11*). Likewise, it has been

identified that IgE-fated MBCs, termed MBC2s, exhibit an IL-4-responsive phenotype marked

by high expression of CD23 (FcεRII) and IL-4 receptor α (IL-4Rα) and low CD32 (FcγRII)

expression (*12–14*). Indeed, seminal work on IgE established IL-4 as a critical molecule for IgE

class switch recombination (CSR) upon primary sensitization (*15, 16*). Despite the formation of a

more specialized MBC compartment, IL-4 continues to be required for secondary IgE responses

(*17–19*). Beyond inhibition of IgE CSR, loss of IL-4 signaling upregulates IFN-γ production

suggestive of plasticity within the T cell compartment (*17*). Whether a parallel degree of

plasticity exists in Ag-specific MBCs, or if the commitment is irreversible, remains to be

elucidated.

In this study, we identified a previously unrecognized plasticity of MBCs within type 2

immunity. By inhibiting IL-4 signaling in a recall response, the pathogenic MBC2 phenotype

was diminished and upregulation of IFN-γ supported the emergence of long-lived IgG2

responses. Even beyond transient IL-4Rα blockade, IgG2 reemerged upon Ag exposure,

suggesting some degree of stability in the type I-like reprogrammed Ab response despite

discontinued IL-4Rα blockade. Ag-specific IgG2 responses appeared to derive from both

unswitched B cells and class-switched MBCs, where the addition of a type 1 adjuvant could

augment the redirection of IgG1 memory to IgG2. Collectively, our findings elucidate a

malleable fate of pathogenic IgE-fated MBCs and highlights the potential to therapeutically

reprogram refractory lifelong allergies.

RESULTS

IL-4Rα signaling augments secondary germinal center activity

To assess the impact of *in vivo* anti-IL-4α (A4RA) treatment on the B cell compartment in type 2

immunity, we employed a well characterized model of epicutaneous sensitization to ovalbumin

(OVA). This strategy establishes a humoral OVA-IgG1 response and IgG1$^+$ MBCs which, upon

subsequent non-sensitizing subcutaneous (s.c.) re-exposures, facilitate IgE production and IgE-

mediated clinical reactivity (*20*). With temporal control over the IgE response, we treated

sensitized mice with A4RA or an isotype control (IC) prior to s.c. re-exposures, as previously

described (*17*). Draining lymph nodes (dLNs) and spleens were collected for single-cell RNA-

sequencing (scRNA-seq), single-cell VDJ-sequencing (scVDJ-seq), and flow cytometry, and

blood was collected for immunoglobulin (Ig) analysis (**Fig. 1A**). For scRNA- and scVDJ-seq

experiments an additional group of mice were administered cholera toxin (CT) and OVA (s.c.) to

model a classical type 2 immunization strategy that yields robust humoral IgE responses and

IgG1$^+$ MBCs.

Four weeks following immunization and s.c. re-exposures, paired scRNA-seq and scVDJ-seq

were performed on 7,369 B cells. Clustering analysis revealed subsets of naïve B cells (*Ighm$^+$*

Ighd$^+$), class-switched MBCs (*Ighm$^-$ Ighd$^-$*), and an *Ighd$^{+/-}$* subset defined by high *Fcrl5* and

Apoe expression (**Fig. 1B and 1C**). Naïve-1 was the dominant naïve subset in CT and IC-treated

mice, though was almost completely replaced by the Naïve-4 cluster in A4RA-treated mice (**Fig.**

1D). Naïve-4 cells exhibited marked transcriptional downregulation, particularly across several

metabolic pathways and genes relating to Ag presentation and cellular localization (**fig. S1A and**

S1B). Indeed, it has been shown that *in vitro* B cell stimulation with IL-4 initiates metabolic

reprogramming and MHC-II upregulation (*21–23*). The Naïve-3 subset, which was almost

exclusively found in epicutaneously sensitized mice, had upregulated *Il31ra* expression, the

ligand (IL-31) of which is implicated in promoting dermatitis (**fig. S1C and S1D**) (*24*). Other

naïve and MBC clusters were similarly represented between the three groups. Across all core

naïve clusters (1-5) and MBCs, A4RA treatment resulted in downregulated oxidative

phosphorylation and B cell receptor (BCR) signaling gene modules – two imperative processes

for germinal center (GC) activity (**Fig. 1E**) (*25, 26*).

Previous reports have demonstrated a critical requirement of IL-4 signaling for primary GC

activity in models of type 2 immunity, though it remains unclear whether this requirement is

maintained for secondary responses where primary responses can prime MBCs for a GC or PC

fate (*27–29*). The requirement of secondary GC activity to mount an IgE response in

epicutaneously sensitized mice was confirmed using CD40L blockade before and during s.c. re-

exposures, whereby anti-CD40L prevented OVA-specific IgE production and clinical reactivity

(**fig. S2A-C**). To investigate how an IL-4/IL-13 signaling deficiency might impact secondary

GCs, we utilized the paired scRNA- and scVDJ-seq data to determine the extent of somatic

hypermutation (SHM) within the MBC cluster. Despite an absence of highly mutated sequences

with A4RA treatment, the overall extent of somatic hypermutation was not significantly different

between groups (**Fig. 1F**).

As the above findings provided only transcriptional insight within the bulk (non-Ag-specific) B

cell repertoire, we assessed the impact of A4RA on OVA-specific B cells by flow cytometry

using fluorochrome-conjugated OVA tetramers (*30*). In agreement with the altered B cell

metabolic profile in the transcriptomic data suggestive of compromised B cell survival and/or

proliferative capacity, the frequency of OVA^+ B cells was 3-fold lower in A4RA-treated mice,

with an average of 406 versus 1269 cells per dLN pair in A4RA and IC groups, respectively

(**Fig. 1G and 1H**). Expression of CD38 and GL7 identifies three populations of B cells differing

in their activation states: $CD38^+$ $GL7^-$ resting naïve/memory (1), $CD38^+$ $GL7^+$ transitional or

multipotent precursor cells (2), and $CD38^-$ $GL7^+$ GC B cells (3) (**Fig. 1I**) (*31*). At three days

post-re-exposures, the A4RA group had significantly impaired OVA-specific B cell activation,

evidenced by an increased frequency of CD38$^+$ GL7$^-$ OVA-specific B cells and a concomitant

decrease in the CD38$^-$ GL7$^+$ population (**Fig. 1J**). However, 23% of OVA-specific B cells in

A4RA-treated mice were in a transitional state compared to 2.6% in IC-treated mice, which

suggested that loss of IL-4/IL-13 signaling altered the kinetics of MBC reactivation. Indeed,

assessment of GC activity at 7-, 14-, and 28-days post-re-exposure demonstrated that the

magnitude of GC activity is no different between groups and A4RA treatment merely stalls early

GC formation (**Fig. 1K**). Taken together, these data suggest that loss of Il-4/IL-13 signaling

elicits profound transcriptional changes to naïve B cells and MBCs, but GC-dependent recall

responses can ultimately proceed.

MBC2 phenotype is ameliorated with IL-4Rα blockade in mice and humans

MBC2s have been identified as an important reservoir for pathological IgE responses.

Previously, IL-4 was shown to be necessary for MBC2 development and, likewise, blockade of

IL-4/IL-13 signaling issued a decline in MBC2 frequency (*12*, *13*, *17*). Despite the observed

decrease in MBC2s a small number persist, though whether expression of the pathogenic IL-4-

responsive phenotype is maintained to the same extent has not been explored.

To address this, we leveraged publicly reposited B cell scRNA-seq datasets involving

perturbation of IL-4/IL-13 signaling. Including our own data generated above, our analysis

involved four data sets comprising both mouse and human, various models of type 2 immunity,

enhanced and abrogated IL-4/IL-13 signaling, and with or without Ag re-exposure (**Fig. 2A**) (*17*,

32, *33*). For each dataset, unsupervised clustering was performed on B cells (*Cd79a*$^+$), which

resolved clusters of MBCs classified by absence of *Ighd*, *Tcl1a* (human), and *Bcl6* (mouse) and

the presence of *Hhex* (mouse) expression (**Fig. 2B and S3A-D**). *Fcer2*, *Il4r*, and *Ighg1*

expression was compared between each control and interventional group to evaluate the impact

of enhanced or inhibited IL-4/IL-13 signaling on MBC2-related gene expression (**Fig. 2C**).

Fcer2 expression in MBCs was consistently downregulated across datasets involving IL-4/IL-13

signaling inhibition. IL-4Rα blockade also elicited a decline in *Il4r* expression, apart from

epicutaneously sensitized mice where *Il4r* transcript abundance was also low in IC mice.

Conversely, with increased IL-4/IL-13 bioavailability to B cells through elimination of its uptake

by lymphoid stromal cells (*Ccl19$^{Cre/+}$ Il4ra$^{fl/fl}$* bone marrow chimeras), *Fcer2* and *Il4r* expression

was enhanced within the MBC cluster (*32*). Intriguingly, *Ighg1* expression was downregulated

only in experimental systems involving Ag re-exposure during IL-4Rα blockade.

To corroborate the diminished MBC2 phenotype within Ag-specific cells, we assessed co-

expression of surface IL-4Rα and CD23 on tetramer[+] MBCs by flow cytometry four weeks

following *s.c.* re-exposures in epicutaneously sensitized mice (**Fig. 2D**). A4RA-treated mice

exhibited a 61% reduction in the frequency of IL-4Rα[+] CD23[+] cells in the OVA-specific MBC

compartment (**Fig. 2E**). Loss of IL-4Rα expression was not an artifact of impaired detection due

to the *in vivo* blocking antibody as there was no increase in the frequency of CD23 single-

positive cells (**Fig. 2E**). Consistent with a role of IL-4 in inducing CD23 expression, A4RA

treatment decreased surface expression of CD23 (**Fig. 2F**) (*34*). CD23 downregulation was not,

however, unique to Ag-specific cells, but instead was broadly observed throughout non-specific

MBCs (**Fig. 2F**).

To also validate the human scRNA-seq observations, we stimulated peripheral blood

mononuclear cells (PBMCs) from peanut-allergic donors (*n*=3) with crude peanut extract for 8

days in the presence or absence of A4RA. CD23[+] IL-4Rα[+] class-switched B cells were observed

in stimulated cells but were virtually undetectable with A4RA treatment (**Fig. 2G**). Consistent

with previous reports, CD23[+] IL-4Rα[+] cells had variable CD27 expression, though both

populations were similarly impacted by A4RA (**Fig. 2H**) (*12*, *13*). Moreover, as was observed in

mice, A4RA broadly decreased CD23 mean fluorescence intensity (MFI) within class-switched

B cells (**Fig. 2I**). Collectively, these data demonstrate a conserved requirement for IL-4/IL-13

signaling to maintain heightened type 2 polarization within established MBC compartments,

which contests a previous suggestion regarding MBC2 phenotypic resiliency (*12*). These data

also raise the issue of how Ag re-exposure during IL-4Rα blockade may skew the isotype

distribution away from IgG1 – the dominant intermediary for IgE CSR.

Long-lived Ag-specific IgG2 responses emerge in the absence of IL-4/IL-13 signaling

As we observed that recall responses occurred in the absence of IL-4/IL-13 signaling, alongside

dampened MBC2 polarization (including *Ighg1*) and prevented IgE CSR, we sought to define the

Ig product of recall responses. Evaluating isotypes downstream to IgG1 – the dominant memory

reservoir in epicutaneously sensitized mice – an increase in *Ighg2b* and *Ighg2c*, but not *Igha*,

transcripts was observed in the MBC cluster of A4RA-treated mice (**Fig. 3A**). From the paired

scVDJ-seq analysis we plotted the assigned *Ighc* gene for each MBC with at least three

nucleotide mutations (**Fig. 3B**). Consistent with the notion that IgE$^+$ MBCs are non-existent in

mice, *Ighe* sequences were undetected (*35*, *36*). The distribution of isotype usage was

comparable between CT and IC mice, except for *Ighg2b*, which had a substantially greater

presence in CT in agreement with the reported CT-induced secretion of TGF-β, a driver of

IgG2b CSR (*37*, *38*). By contrast, A4RA induced a pronounced increase in *Ighg2c* above both

CT and IC – an isotype that is not associated with type 2 immunity.

To assess whether the skewed isotype usage occurred within the Ag-specific B cell repertoire

and the longevity of such response, OVA$^+$ MBCs and the OVA-specific serum Ig profile were

interrogated up to 15 weeks after the recall response. OVA$^+$ IgG1$^+$ MBCs were present in both

IC and A4RA-treated mice, though with a significantly decreased frequency with A4RA treatment (**Fig. 3C and 3D**). By comparison, OVA[+] IgG2c[+] MBCs emerged with A4RA treatment and were sustained beyond three months. The emergence of OVA[+] IgG2c[+] MBCs was halted with concurrent IFN-γ receptor 1 blockade (**Fig. 3E**). This coincides with IgG2c CSR being driven by IFN-γ and our previous work demonstrating an increase in IFN-γ production with A4RA treatment (*17, 39, 40*). Skewed isotype usage was also reflected in the analysis of serum OVA-specific Igs (**Fig. 3F**). As per our previous observations, serum OVA-IgE responses were short-lived in IC mice with a completely abrogated response in A4RA-treated mice (*17, 20*). Consistent with the induction of IgG2c[+] MBCs, serum OVA-IgG2c was observed with A4RA treatment though was not long-lived and declined at a similar rate to IgE in IC mice. Disparate to the reduction in OVA[+] IgG1[+] MBCs, Ag re-exposure induced a comparable resurgence of serum OVA-IgG1 in both groups, suggesting that A4RA may have limited impact on MBCs fated for PC differentiation. Serum OVA-IgG2b titers were comparable between both groups and OVA-IgA was undetectable.

We reasoned that if the skewed isotype response by A4RA was stably maintained then a resurgence of IgG2c should be facilitated by seeded IgG2c[+] MBCs. To this end, mice were provided tertiary OVA re-exposures nine weeks post-A4RA/IC and s.c. OVA administration. At this point there is clearance of both serum OVA-IgG2c and the blocking A4RA antibody (*17*). Indeed, tertiary Ag re-exposure provided a significant regeneration of OVA-IgG2b and OVA-IgG2c though abrogation of OVA-IgE production remained (**Fig. 3G**). Collectively, these data demonstrate that induction of a recall response with IL-4Rα blockade permit the emergence of a non-type 2 antibody response and propose that this reprogrammed B cell response may be maintained even following A4RA clearance.

LNs at distinct anatomical sites have been shown to differ in their predisposition to promote

inflammatory or tolerogenic responses towards Ags (*41*). In this regard, we also investigated

A4RA-induced isotype skewing in the context of oral immunization, which preferentially issues

tolerogenic IgA-dominant responses. As previously shown, one intragastric administration of Ag

+ CT does not issue a humoral response (*42*, *43*), though Ag-specific Igs arise after subsequent

oral administration of Ag alone (*42*). We adopted this framework to provide systemic

administration of IC or A4RA prior to each oral Ag re-exposure (**fig. S4A**). Expectedly, IC-

treated mice exhibited a classical type 2 response marked by OVA$^+$ IgG1$^+$ MBCs in gut dLNs

and serum OVA-IgG1 and -IgE (**fig. S4B and C**). Mirroring observations from epicutaneous

sensitization, A4RA elicited the emergence of OVA$^+$ IgG2c$^+$ MBCs and serum OVA-IgG2b and

-IgG2c, while preventing OVA-IgE synthesis. Conversely, OVA re-exposures did initiate an IgA

response which was abrogated with A4RA, reconciling the requirement of IL-4 signaling for

mucosal IgA responses (*44*) (**fig. S4C**). Thus, loss of IL-4/IL-13 signaling during Ag re-exposure

preferentially promotes IgG2 responses irrespective of tissue site.

Class switch origins of IgG2$^+$ clones are comprised of multiple isotypes

The observed loss of MBC2 polarization and divergence towards non-IgE humoral response

indicate the potential to reprogram pathogenic IgE responses; however, it remains unclear

whether skewed isotype usage arises *de novo* (i.e., from naïve or unswitched B cells) or from

direct reprogramming of existing class-switched MBCs. To address this, we performed bulk VDJ

sequencing on the OVA-enriched B cell fraction from pooled LNs and spleens. Parallel analyses

with Ig sequences derived in the 10X dataset were carried out to gain insight as to whether the

effects of A4RA are occurring broadly throughout all B cells or are contained within the Ag-

specific repertoire. We found that IL-4/IL-13 signaling did not influence CDR3 length or SHM

in unswitched and class-switched B cell repertoires (**fig. S5A-D**); Apparent biases in V gene

usage were observed within the class-switched repertoire of OVA-enriched B cells (**fig. S5E**);

however, this was also observed within polyclonal B cells and thus is unlikely related to the

OVA-specific recall response (**fig. S5F**).

Broadly probing whether unswitched or class-switched cells were putative precursors to IgG2

clones, we first measured the correlation of paired IGHV and IGHJ gene usage between isotypes.

Expectedly, paired IGHV-IGHJ gene usage was most highly correlated between *Ighm* and *Ighd*

clones in both groups (**Fig. 4A**). IGHV-IGHJ gene usage in *Ighg2* clones also correlated most

strongly with *Ighm* and *Ighd* clones, though overlap with *Ighg3* and *Ighg1* clones was increased

in A4RA (correlation coefficient = 0.31 *Ighg3* and 0.34 *Ighg1*) compared to IC (correlation

coefficient = 0.14 *Ighg3* and 0.22 *Ighg1*). Next, we partitioned sequences into clonal lineages

through hierarchical clustering, as described previously (*45*, *46*). Consistent with the correlation

of V-J gene and isotype usage, the majority of clones related to IgG1 and IgG2 sequences were

IgD or IgM. The isotype distribution of other shared IgG1 and IgG2 clones was similar between

IC and A4RA (**Fig. 4B**), suggesting that loss of IL-4/IL-13 signaling did not influence the class

switch origins of IgG2. Accordingly, mutated IgG2 sequences were present in B cell lineage

trees containing expanded IgM, IgG3, and IgG1 clones, demonstrating that both naïve and class-

switched cells can give rise to IgG2 (**Fig. 4C**).

While IL-4/IL-13 signaling did not appear to influence isotype origins of IgG2, we reasoned that

an increased propensity for IgG2 class switching would be reflected in the transcriptome of Ag-

specific B cells. To this end, we performed Smart-Seq2 on OVA-specific cells yielding the

reconstruction of full length BCRs for 40 and 41 OVA[+] B cells from IC- and A4RA-treated

mice, respectively. Of the sequenced cells, 22 were class-switched, where *Ighg2b* ($n = 2$) and

Ighg2c (*n* =1) sequences were only present in the A4RA group, consistent with upregulated

Ighg2b and *Ighg2c* expression in the 10X dataset (**table. S2 and Fig. 3A and B**). The limited

throughput of this approach did not permit the detection of clonally related sequences within

either group. Comparing gene expression of unswitched cells, A4RA upregulated genes relating

to IFN-γ signaling, including *Ifitm2, Cd63, Ifi203,* and *Mars*; conversely, *Ptpn2*, a negative

regulator of IFN-γ signaling was downregulated, collectively supporting an altered disposition of

unswitched B cells to undergo IgG2 CSR (**Fig. 4D**). Likewise, A4RA *Ighg1*[+] OVA-specific B

cells upregulated some IFN-γ-related genes (*Ifitm2, Cd63, Ets1,* and *Jak1*), which were similarly

expressed in *Ighg2b/c*[+] cells (**Fig. 4E**). Taken together, blockade of IL-4/IL-13 signaling can

influence the transcriptional profile of both unswitched and class-switched OVA-specific B cells

to give rise to IgG2 responses *in vivo.*

Type 1-polarizing adjuvant can redirect isotypic fate of type 2-polarized class-switched

MBCs

Having demonstrated evidence that IgG3 and IgG1 – though infrequently – can be an

intermediate to an IgG2 isotype we next assessed whether the addition of a type 1-polarizing

adjuvant could augment the redirection of class-switched MBCs to IgG2b/c. We utilized an

adjuvanted re-exposure system to assess whether the type 1 adjuvant, CpG, can divert the Ag-

specific response of class-switched MBCs from type 2 immunized mice (**Fig. 5A**). Class-

switched OVA-enriched MBCs were sorted from OVA + Alum-immunized CD45.1[+] donor mice

and were transferred to B cell-deficient (muMT) CD45.2[+] naïve mice. Recipient mice were then

immunized with either OVA + CT or OVA + CpG. Of note, OVA + Alum immunization yields

an IgG1 and IgE dominant response, where over 85% of our transferred class-switched MBC

pool was IgG1[+] and less than 4% were IgG2b/c[+] (*47*) (**Fig. 5B**). Thus, any emergence of OVA-

IgG2 responses in recipient mice is not likely to represent an expansion of donor IgG2 clones.

Moreover, OVA + CT immunization produces more OVA$^+$ IgG2b/c$^+$ MBCs than Alum;

therefore, if any IgG2 clones in the donor pool do contribute to the OVA-IgG2 response in

recipients, we would expect to observe this with both CT and CpG immunization (**Fig. 5B**).

One week following adoptive transfer and immunization, donor (CD45.1$^+$) OVA$^+$ B cells and

OVA-specific Igs were detected in recipient mice. As anticipated, immunization of recipient

mice with OVA + CT drove emergence of OVA$^+$ IgG1$^+$ MBCs and serum OVA-IgG1 and -IgE.

In contrast, OVA$^+$ IgG2c$^+$ MBCs and serum OVA-IgG2b and -IgG2c became present in recipient

mice immunized with OVA + CpG, with a sustained absence of OVA-IgE (**Fig. 5C and 5D**).

Collectively, these data support that Ag re-exposure in the presence of a type 1 adjuvant can

redirect the fate of class-switched MBCs established under type 2-polarizing conditions.

DISCUSSION

MBCs, compared to naïve counterparts, acquire enhanced functional and metabolic properties

increasing fitness for recall responses to the same Ag. The MBCs established upon allergic

sensitization gain a unique propensity to switch to IgE upon subsequent allergen exposures,

under the influence of IL-4. Whether this specialized B cell memory is functionally plastic was

not yet established. Assessing transcriptional profile, BCR repertoire, Ag-specific cellular

memory, and humoral Ab responses, our work identifies the capacity to reprogram the

pathogenic IgE-fated response towards a type 1-like, non-pathogenic IgG2 program.

Numerous studies in mice and humans have now established that IgG1 is the dominant MBC

reservoir for IgE (*5*, *10*, *11*, *20*, *48*). Recent work by us and others have extended upon this

concept illustrating that IgE-fated MBCs ("MBC2s") express ε germline transcripts and

upregulate surface expression of CD23 and IL-4Rα, seemingly remaining prepared for IL-4

responsiveness (*12–14*). In the absence of IL-4/IL-13 signaling in a recall response, we have

previously shown that IgE responses are indeed averted, and cytokine production skews toward a

Th1 profile (*17*). In the present work investigating how IL-4Rα blockade impacts the B cell

repertoire beyond avoidance of IgE production, we found that loss of IL-4/IL-13 signaling *in*

vivo broadly suppresses gene expression, particularly relating to metabolic processes and BCR

signaling. Intriguingly, Ag re-exposure could still elicit Ag-specific GC activity in the absence of

IL-4/IL-13, an issue which has remained contentious (*27, 49, 50*).

As a primary product of a GC response is PC production, we were particularly intrigued to see a

sustained absence of serum IgE without a compensatory increase in IgG1. Coinciding with our

previous observations of an increased IFN-γ to IL-4 ratio, we found that A4RA treatment

induced an IFN-γ-dependent Ag-specific IgG2c response, which was otherwise absent with IC.

IgG2 antibodies, unlike IgG1, only weakly bind Fcγ receptors and thus, can regulate the pro-

inflammatory effector functions of IgG1 through competition for Ag binding (*51, 52*). IgG2 is

postulated to also regulate IgE effector functions directly through engagement with an inhibitory

receptor, CD32b, on mast cells and basophils (*53*). Indeed, allergen desensitization is associated

with increases in allergen-specific IgG2, paralleling the well documented increases in IgG4 (*54–*

56). In our studies we found that serum titers of Ag-specific IgG2c were short-lived, however

tertiary Ag exposure could re-boost serum IgG2 titers without any observed increase in IgE.

Notably, the boosted IgG2 response occurred in the absence of continued IL-4Rα blockade

illustrating a sustained reprogramming of the Ag-specific B cell response.

The molecular cues which sustain the MBC2 phenotype have not yet been described. In the T

cell compartment, a subset of T helper cells, termed "Th2A" is uniquely associated with allergic

disease, much like the novel MBC2 subset (*57*). Th2A cells exhibit heightened Th2-like

properties and are terminally differentiated, persisting even with IL-4Rα blockade (*17, 57, 58*).

Here, we demonstrate that unlike Th2A cells, MBC2s do not harbor a refractory pathogenic

phenotype. This requirement of IL-4 to sustain heightened CD23 and IL-4Rα expression was

consistent amongst mouse models and humans and reaffirms early observations of IL-4-

regulated expression in bulk B cell populations (*34, 59*). Interestingly, CD23 purportedly

augments allergen-specific responses through IgE-facilitated Ag-presentation (*60, 61*). Thus, it is

plausible that depressed CD23 expression observed not only on MBC2s, but throughout the B

cell compartment, may curtail reactivation of allergen-specific CD4 T cells, though this remains

an area for future investigation.

Our analysis of BCR sequences revealed clonal relationships between IgG2 and all upstream

isotypes, including IgM/D, IgG3, and IgG1. These data, therefore, suggest that both naïve and

Ag-experienced B cells can contribute to the A4RA-induced IgG2 response. Indeed, we showed

functionally that the adoptive transfer of class-switched MBCs (composed of over 85% IgG1$^+$

cells) from type 2 immunized mice could issue IgG2 responses following a boost with a type 1-

polarizing adjuvant. Sequential CSR has been most extensively studied in the IgM to IgE

trajectory, where IgG1 is a dominant intermediary. Interestingly, over 50% of the bulk IgG2

repertoire in humans also arises through sequential CSR with an IgG1 intermediary (*62*). This is

somewhat surprising given that sequential rounds of V(D)J recombination are likely more

metabolically taxing to the host. One explanation for this could be that sequential, rather than

direct, CSR provides a means to regulate pro-inflammatory antibody-mediated effector

functions. In agreement, both vaccination and allergy yield IgG1 dominant responses, though

with chronic Ag exposure through repeated vaccination or allergen immunotherapy, a shift to the

"anti-inflammatory" isotype, IgG4, is observed (*55, 63–66*). Here, we demonstrated that Ag

exposure in the absence in IL-4/IL-13 signaling could produce a similar effect – a deviation from

the IgG1-IgE axis to an IgG2 response. As IgG2 is uninvolved in allergic inflammation in both

mice and humans, this shift to allergen-specific IgG2 production could be considered "anti-

inflammatory". Nevertheless, clonal relationships between IgG2 and downstream IgE have been

reported (*9*); whether the induced IgG2$^+$ MBCs can later revert to a pathogenic IgE fate in

humans is yet to be studied.

In summary, our research demonstrates that IL-4/IL-13 signaling is required to sustain the IgE

fate of B cell responses in established allergic disease and that loss of IL-4/IL-13 signaling can

transform the allergen response to an innocuous type 1 phenotype. This not only bolsters the

applicability of anti-IL-4Rα as a novel therapeutic in food allergy, but also broadly illuminates

the potential to reprogram deviant B cell responses. Thus, further research is warranted to

establish the intrinsic determinants that govern MBC fate, which will be imperative in the

development of therapeutic strategies to correct deviant B cell responses in settings like allergy,

autoimmunity, and transplantation.

MATERIALS AND METHODS

Mice

Female C57BL/6 mice (6-8 weeks old) were purchased from Charles River Laboratories

(Wilmington, MA). Female B6.129S2-*Ighm*^{tmCgn} and B6.SJL-*Ptprc*^{a} *Pepc*^{b}/Boy mice (6-8 weeks

old) were purchased from Jackson Laboratories (Bar Harbor, ME). All experimentation was in

compliance with the McMaster University Research Ethics Board.

Immunizations

Epicutaneous sensitization: Epicutaneous sensitization to ovalbumin (Sigma-Aldrich, St. Louis,

MO) was conducted as previously described (*20*). For recall responses, three low-back

subcutaneous injections of 100 μg OVA were given over the course of a week. *Gastric*

sensitization: Mice received one intragastric gavage of 1 mg OVA with 5 μg cholera toxin (CT)

in PBS. Four weeks later, mice received 1 mg OVA in PBS weekly for three weeks via gavage.

Subcutaneous sensitization: Mice were immunized subcutaneously with 250 μg OVA and 5 μg

CT. *Intraperitoneal sensitization:* Mice were immunized intraperitoneally with 250 μg OVA and

250 μL Alhydrogel 2% (aluminum hydroxide [alum], InvivoGen, San Diego, CA), 40 μg CpG,

or 5 μg CT.

Serum for analysis of OVA-specific antibodies was collected through retro-orbital bleeding. For

allergen challenges, 5 mg OVA was intraperitoneally injected. Core body temperature and

clinical signs were recorded, as previously described, and hematocrit was measured at 40

minutes (*67*).

Antibody Administration

Anti-CD40L (MR-1), anti-IFNγ-R1 (GR-20), and anti-trinitrophenol IgG2a isotype control

(2A3) were purchased from Bio X Cell (Lebanon, NH). Anti-IL-4Rα (M1) was produced in-

house, as previously described (*17*). Original hybridomas were generated by Immunex

Corporation (Amgen, Thousand Oaks, CA) and were generously provided by Dr. Fred

Finkelman (University of Cincinnati).

One mg of anti-IL-4Rα, anti-IFNγ-R1, or isotype control was intraperitoneally injected the day

prior to s.c. re-exposures. Anti-CD40L (0.35 mg) or isotype control was intraperitoneally

injected on days -1, 0, and 2 of s.c. re-exposures.

Tetramer Staining and Enrichment

Purified OVA (Sigma-Aldrich, St. Louis, MO) was biotinylated (ThermoFisher Scientific, Waltham, MA) and tetramerized with streptavidin-PE or streptavidin-APC (Agilent, Santa Clara, CA), as previously described (*30*). For decoy tetramer, β-lactoglobulin (Sigma-Aldrich, St. Louis, MO) was biotinylated. Streptavidin-PE or -APC was conjugated with DyLight 594 or DyLight 755 esters (ThermoFisher Scientific) and was used to tetramerize the biotinylated- β-lactoglobulin.

Cell pellets were resuspended and stained with 5 nM decoy tetramer in 20 μL FACS buffer (10% FBS and 1 μM EDTA in PBS) and 2% rat serum for five minutes on ice. Five nM OVA tetramer was added and incubation continued for 30 minutes. Cells were washed to remove unbound tetramer and were incubated with anti-PE or -APC microbeads (Miltenyi Biotec, Germany) for 15 minutes on ice. Cells were passed through LS columns affixed to MACS magnetic separator (Miltenyi Biotec). Bound fractions were eluted with FACS buffer.

For CD45.1 enrichments, cell pellets were resuspended in 20 μL FACS buffer with 0.25 μg FcBlock (BioLegend) for five minutes on ice, then stained with anti-CD45.1-APC (A20; BioLegend) for 30 minutes. Enrichment with anti-APC microbeads proceeded as described above.

Patient Samples

A cohort of three peanut allergic donors, aged 9-23, were recruited to donate blood from McMaster Children's Hospital and Hamilton Health Sciences (Hamilton, Ontario, Canada). Allergic status was confirmed by peanut-specific IgE (higher than 0.35 kU/L) ImmunoCap at Laboratory Reference Centre Hamilton (McMaster Children's Hospital). Donors were excluded

from the study for previous or current allergen immunotherapy, omalizumab, dupilumab or other

systemic immunomodulatory treatment, as well as diagnosed autoimmune or immunodeficiency

diseases. All donors were recruited with written consent and ethical approval from the Hamilton

Integrated Research Ethics Board.

Peripheral blood was collected into heparinized tubes (BD). PBMCs were isolated via Ficoll-

Paque (GE Healthcare) density centrifugation and were cryopreserved in FBS + 10% DMSO for

long-term storage in liquid nitrogen. Before culture, viability of thawed PBMCs was confirmed

to be ≥ 70%. PBMCs were cultured for 8 days and stimulated with 2.5 ng/mL crude peanut

extract (Stallergenes Greer) with or without 50 ug/mL anti-human IL-4Ra (R&D Systems), as

previously described (*17*).

Flow Cytometry and Cell Sorting

Single-cell suspensions were plated in U-bottom 96-well plates and incubated with 0.25 µg

FcBlock for 10 minutes on ice. Cells were stained extracellularly for 30 minutes on ice with

antibody cocktails prepared in FACS buffer. For intracellular staining, cells were fixed and

permeabilized with Cytofix/Cytoperm (BD Biosciences), as per manufacturer's instructions.

Intracellular antibody cocktails were prepared in 1x perm wash and cells were stained for 30

minutes on ice.

The following anti-mouse fluorochrome-conjugated antibodies were used: B220-AF700 (RA3-

6B2), CD138-PE/Dazzle 594 (281-2), CD3-BV711 (17A2), CD3-BV510 (17A2), F4/80-BV711

(BM8), F4/80-BV510 (BM8), CD38-PE/Cy7 (90), GL7-PerCP.Cy5.5 (GL7), GL7-BV421 (GL7;

BD, Franklin Lakes, NJ), IgD-Pacific Blue (11-26c.2a), IgD-BV605 (11-26c.2a), IgM-BV786

(II/41; BD), IgM-FITC (II/41), IgG1-BV650 (RMG1-1), IgG2b-PECy7 (RMG2b-1), IgG2c-

FITC (polyclonal; SouthernBiotech, Birmingham, AL), IL-4Rα-PE (REA235; Miltenyi Biotec),

and CD23-BV421 (B3B4). The following anti-human antibodies were used: IL-4Rα-PE

(G077F6), CD23-BV421 (EBVCS-5), IgD-AF700 (IA6-2), CD27-BV605 (O323), IgM-BV650

(MHM-88), CD38-PECy7 (HB-7), CD19-PerCP.Cy-5.5 (SJ25C1), CD3-BV711 (OKT3), CD14-

BV711 (M5E2), and CD16-BV711 (3G8). All antibodies were purchased from BioLegend (San

Diego, CA), unless otherwise noted. Viability staining was executed with Fixable Viability Dye

eFluor780 (eBioscience, San Diego, CA) or Fixable Viability Stain 510 (BD Biosciences).

Data were acquired with an LSRFortessa flow cytometer (BD). Bulk cell sorting was performed

on a MoFlo XDP cell sorter (Beckman Coulter, Brea, CA). All data were analyzed with FlowJo,

version 10 software (Tree Star, Ashland, OR).

Adoptive Transfers

Class-switched MBCs (B220$^+$ CD3$^-$ F4/80$^-$ CD38$^+$ GL7$^-$ IgD$^-$ IgM$^-$) from tetramer enriched

fractions were isolated by flow sorting four weeks after donor immunization. 10^4 cells were

intravenously injected in the tail-vein of naïve recipient (muMT) mice. The day after transfer

recipients were intraperitoneally injected with 250 μg OVA with either 40 μg CpG or 5 μg CT.

One week later, spleens and mesenteric LNs were collected for flow cytometric analysis of donor

B cells, and serum was collected for OVA-specific antibody analysis by ELISA.

ELISAs

OVA-specific immunoglobulins were measured by sandwich ELISA, described previously, with

some modifications (*68*). Briefly, MaxiSorp plates (ThermoFisher Scientifc) were coated with

100 ng anti-IgG2b, -IgG2c, -IgE, or -IgA overnight at 4°C. Plates were blocked in 5% skim milk

at 37°C for one hour, then diluted samples were added and incubated overnight at 4°C.

Digoxigenated OVA (15 ng) prepared in-house was added for two hours at room temperature,

followed by anti-digoxigenin Fab fragments (Roche, Basel, Switzerland) for one hour. Plates

were developed with 3,3',5,5'-tetramethylbenzidinie (Sigma-Aldrich) and stopped with sulfuric

acid. Optical densities were acquired with a Multiskan FC photometer (ThermoFisher Scientific).

For OVA-IgG1, plates were coated with 400 ng OVA in carbonate-bicarbonate buffer overnight

at 4°C. Plates were blocked in 5% skim milk at 37°C for two hours, then diluted samples were

added and incubated overnight at 4°C. Biotinylated anti-mouse IgG1 (SouthernBiotech;1.25 ng)

was added for two hours at room temperature, followed by alkaline-phosphatase streptavidin

(ThermoFisher Scientific) for one hour. Plates were developed with 4-nitrophenyl phosphate

(Sigma-Aldrich) and stopped with sodium hydroxide. Optical densities were acquired as above.

10x Chromium Sequencing & Analysis

Sequencing: Spleens and inguinal LNs were collected one month post-recall response from

epicutaneously or subcutaneously sensitized mice and B cells were sorted for sequencing.

Single-cell 5' gene expression and V(D)J libraries were prepared with Chromium Single-Cell V3

Reagent Kit and Controller (10x Genomics; Pleasanton, CA). Libraries were sequenced with

NextSeq2000 (Illumina; San Diego, CA). Data were processed with the Cell Ranger Count and

V(D)J pipelines (version 7.0.1).

Gene Expression Analysis: Sequencing reads were aligned to the mouse transcriptome

(GRCm38), followed by filtering, and correction of cell barcodes and Unique Molecular

Identifiers (UMIs). Reads associated with retained barcodes were quantified and used to build

expression matrices for further analysis with Seurat (version 4.1.3). Cells expressing less than

200 genes, genes that were detected in less than 3 cells, and cells expressing over 10%

mitochondrial genes were filtered out. Normalization, variable feature selection, and scaling

were performed with Seurat default parameters. Principal component analysis (PCA) was

performed and the top 15 components were used for UMAP dimensionality reduction and

clustering at resolution 0.4. Contaminating non-B cells were filtered out and PCA and clustering

were repeated on the subsetted Seurat object at resolution 0.3. Differentially expressed genes or

gene modules were identified with the Wilcoxon Rank Sum test.

BCR Analysis: BCR analysis was performed with pipelines in the Immcantation suite (*46*). The

"filtered_contig.fasta" and "filtered_contig_annotations.csv" outputs from Cell Ranger V(D)J

were used to assign V, D, and J genes with IgBLAST and cells assigned with 0 or >1 heavy

chains were removed. Clonal groups were defined with the hierarchialClones function using

applying a Hamming distance of 0.1. Germline sequences were reconstructed with

createGermlines and IMGT germline references (version 3.1.39). Somatic hypermutation

frequency within the IGHV gene was calculated with observedMutations.

Bulk VDJ Sequencing & Analysis

Sequencing: Spleens and inguinal LNs were collected one month post-recall response from

epicutaneously sensitized mice. ~12,000-15,000 B cells from the tetramer-enriched fraction were

sorted directly into Buffer RLT (Qiagen; Hilden, DE) prior to shipment on dry ice to iRepertoire.

RNA extraction was performed prior to 3-chain long read library preparation (Mouse BCR

Heavy, Kappa, Lambda). Amplified PCR products were run on MiSeq System with MiSeq

Reagent Nano Kit v2 (iRepertoire, Huntsville, AL).

Analysis: Nucleotide sequences and UMI count were extracted from "joinedSeq" and "copy"

columns of the "pep.csv" output from iRweb (iRepertoire) and were aligned with IgBLAST

(version 1.18.0) using IMGT reference sequences. Processing of IgBLAST output was

performed with the Immcantation suite to assign V, D, and J genes, cluster into clonal groups,

assign unmutated germlines, and calculate IGHV somatic hypermutation frequency, as described

in the 10x Chromium section. Lineage tree topologies were predicted by maximum parsimony in

IgPhyML with constraints for unidirectional isotype switching and trees were subsequently

visualized with dowser (*69, 70*).

Smart-Seq2 Sequencing & Analysis

Sequencing: Spleens and inguinal LNs were collected one month post-recall response from

epicutaneously sensitized mice. OVA-specific B cells were-single cell sorted into 96 U-bottom

well PCR-grade RNase and DNase free plates before being flash frozen at -80C. Full-length

cDNA was generated with Smart-Seq Single Cell Kit (Takarabio; Mountain View, CA).

Libraries were then prepared for sequencing with NovaSeq 600 Reagent Kits (Illumina;

20028400) and sequenced on Illumina NovaSeq 6000 (Ilumina; San Diego, CA).

Analysis: Sequencing reads were aligned to the mouse transcriptome (GRCm38) with STARsolo

(*71*). Count matrix output was normalized and scaled with Seurat (version 4.1.3). BCR

sequences were reconstructed with the TRUST4 algorithm, followed by integration with the gene

expression data using Platypus (*72, 73*)

Statistical Analysis

Statistical analyses not pertaining to RNA sequencing were performed with Prism 9 (GraphPad

Software; San Diego, CA). Comparisons were drawn between groups (CT, IC, A4RA) or across

time points using two-sided Student's *t*-test or ANOVA. Data were considered significant with a

p value < 0.05.

List of Supplementary Materials

Fig. S1. Gene expression profiles of naïve B cell clusters.

Fig. S2. Requirement of B-T interactions for IgE recall responses.

Fig. S3. Analysis of publicly reposited scRNA-seq datasets.

Fig. S4. B cell reprogramming in orally-sensitized mice.

Fig. S5. VDJ repertoire analysis in OVA-enriched bulk sequencing and 10X datasets.

References and Notes

1.   R. R. Goel, S. A. Apostolidis, M. M. Painter, D. Mathew, A. Pattekar, O. Kuthuru, S. Gouma, P. Hicks, W.
Meng, A. M. Rosenfeld, S. Dysinger, K. A. Lundgreen, L. Kuri-Cervantes, S. Adamski, A. Hicks, S. Korte,
D. A. Oldridge, A. E. Baxter, J. R. Giles, M. E. Weirick, C. M. McAllister, J. Dougherty, S. Long, K.
D'Andrea, J. T. Hamilton, M. R. Betts, E. T. Luning Prak, P. Bates, S. E. Hensley, A. R. Greenplate, E. J.
Wherry, Distinct antibody and memory B cell responses in SARSCoV-2 naïve and recovered individuals
following mRNA vaccination. *Sci Immunol* **6**, 1–19 (2021).
2.   J. Wrammert, K. Smith, J. Miller, W. A. Langley, K. Kokko, C. Larsen, N. Y. Zheng, I. Mays, L. Garman,
C. Helms, J. James, G. M. Air, J. D. Capra, R. Ahmed, P. C. Wilson, Rapid cloning of high-affinity human
monoclonal antibodies against influenza virus. *Nature 2008 453:7195* **453**, 667–671 (2008).
3.   T. Onodera, Y. Takahashi, Y. Yokoi, M. Ato, Y. Kodama, S. Hachimura, T. Kurosaki, K. Kobayashi,
Memory B cells in the lung participate in protective humoral immune responses to pulmonary influenza
virus reinfection. *Proc Natl Acad Sci U S A* **109**, 2485–2490 (2012).
4.   Z. Yang, B. M. Sullivan, C. D. C. Allen, Fluorescent in vivo detection reveals that IgE(+) B cells are
restrained by an intrinsic cell fate predisposition. *Immunity* **36**, 857–72 (2012).
5.   R. Jiménez-Saiz, D. K. Chu, T. S. Mandur, T. D. Walker, M. E. Gordon, R. Chaudhary, J. Koenig, S. Saliba,
H. J. Galipeau, A. Utley, I. L. King, K. Lee, R. Ettinger, S. Waserman, R. Kolbeck, M. Jordana, Lifelong
Memory Responses Perpetuate Humoral Th2 Immunity and Anaphylaxis in Food Allergy. *Journal of*
Allergy and Clinical Immunology **140**, 1604–1615 (2017).
6.   N. Carballeda-Sangiao, A. I. Rodríguez-Mahillo, M. Careche, A. Navas, I. Moneo, M. González-Muñoz,
Changes over Time in IgE Sensitization to Allergens of the Fish Parasite Anisakis spp. *PLoS Negl Trop Dis*
10, e0004864 (2016).
7.   M. S. Kim, M. D. Straesser, B. Keshavarz, L. Workman, E. C. McGowan, T. A. E. Platts-Mills, J. M.
Wilson, IgE to galactose-α-1,3-galactose wanes over time in patients who avoid tick bites. *J Allergy Clin*
Immunol Pract **8**, 364-367.e2 (2020).
8.   D. L. Neuman-Sunshine, J. A. Eckman, C. A. Keet, E. C. Matsui, R. D. Peng, P. J. Lenehan, R. A. Wood,
The natural history of persistent peanut allergy. *Annals of Allergy, Asthma & Immunology* **108**, 326-331.e3
(2012).
9.   T. J. Looney, J.-Y. Lee, K. M. Roskin, R. A. Hoh, J. King, J. Glanville, Y. Liu, T. D. Pham, C. L. Dekker,
M. M. Davis, S. D. Boyd, Human B-cell isotype switching origins of IgE. *J Allergy Clin Immunol* **137**, 579-
586.e7 (2016).
10.   J.-S. He, S. Subramaniam, V. Narang, K. Srinivasan, S. P. Saunders, D. Carbajo, T. Wen-Shan, N. Hidayah
Hamadee, J. Lum, A. Lee, J. Chen, M. Poidinger, F. Zolezzi, J. J. Lafaille, M. A. Curotto de Lafaille, IgG1
memory B cells keep the memory of IgE responses. *Nat Commun* **8**, 641 (2017).
11.   I. Hoof, V. Schulten, J. A. Layhadi, T. Stranzl, L. H. Christensen, S. Herrera de la Mata, G. Seumois, P.
Vijayanand, C. Lundegaard, K. Niss, A. Lund, J. Ahrenfeldt, J. Holm, E. Steveling, H. Sharif, S. R. Durham,

B. Peters, M. H. Shamji, P. S. Andersen, Allergen-specific IgG+ memory B cells are temporally linked to
IgE memory responses. *J Allergy Clin Immunol* **146**, 180–191 (2020).

12. C. J. Aranda, E. Gonzalez-Kozlova, S. P. Saunders, W. Fernandes-Braga, M. Ota, S. Narayanan, J. S. He, E.
Del Duca, B. Swaroop, S. Gnjatic, G. Shattner, J. Reibman, N. A. Soter, E. Guttman-Yassky, M. A. Curotto
de Lafaille, IgG memory B cells expressing IL4R and FCER2 are associated with atopic diseases. *Allergy*
00, 1–15 (2022).

13. J. F. E. Koenig, N. P. H. Knudsen, A. Phelps, K. Bruton, I. Hoof, G. Lund, D. Della Libera, A. Lund, L. H.
Christensen, D. R. Glass, T. Walker, A. Fang, S. Waserman, M. Jordana, P. S. Andersen, A Distinct
Phenotype of Polarized Memory B cell holds IgE Memory. *bioRxiv*, 2023.01.25.525495 (2023).

14. M. Ota, K. B. Hoehn, T. Ota, C. J. Aranda, S. Friedman, W. F. Braga, A. Malbari, S. H. Kleinstein, S. H.
Sicherer, M. A. C. de Lafaille, The memory of pathogenic IgE is contained within CD23+IgG1+ memory B
cells poised to switch to IgE in food allergy. *bioRxiv*, 2023.01.25.525506 (2023).

15. F. D. Finkelman, J. F. Urban, M. P. Beckmann, K. A. Schooley, J. M. Holmes, I. M. Katona, Regulation of
murine in vivo IgG and IgE responses by a monoclonal anti-IL-4 receptor antibody. *Int Immunol* **3**, 599–607
(1991).

16. F. D. Finkelman, I. M. Katona, J. F. Urban, J. Holmes, J. Ohara, A. S. Tung, J. Sample vG., W. E. Paul, IL-4
is required to generate and sustain in vivo IgE responses. *J Immunol* (1988).

17. K. Bruton, P. Spill, S. Vohra, O. Baribeau, S. Manzoor, S. Gadkar, M. Davidson, T. D. Walker, J. F. E.
Koenig, Y. Ellenbogen, A. Florescu, J. Wen, D. K. Chu, S. Waserman, R. Jiménez-Saiz, S. Epelman, C.
Robbins, M. Jordana, Interrupting reactivation of immunologic memory diverts the allergic response and
prevents anaphylaxis. *J Allergy Clin Immunol* **147**, 1381–1392 (2021).

18. J. K. Nono, H. Ndlovu, N. Abdel Aziz, T. Mpotje, L. Hlaka, F. Brombacher, Interleukin-4 receptor alpha is
still required after Th2 polarization for the maintenance and the recall of protective immunity to Nematode
infection. *PLoS Negl Trop Dis* **11** (2017).

19. I. M. Katona, J. F. Urban, S. S. Kang, W. E. Paul, F. D. Finkelman, IL-4 requirements for the generation of
secondary in vivo IgE responses. *J Immunol* **146**, 4215–21 (1991).

20. R. Jiménez-Saiz, Y. Ellenbogen, J. Koenig, M. E. Gordon, T. D. Walker, D. Rosace, P. Spill, K. Bruton, J.
Kong, K. Monteiro, J. Wen, E. I. Tuomanen, R. Kolbeck, D. K. Chu, S. Waserman, M. Jordana, IgG1 $^+$ B
cell immunity predates IgE responses in epicutaneous sensitization to foods. *Allergy*, doi: 10.1111/all.13481
(2018).

21. K. Haniuda, S. Fukao, D. Kitamura, Metabolic Reprogramming Induces Germinal Center B Cell
Differentiation through Bcl6 Locus Remodeling. *Cell Rep* **33** (2020).

22. L. R. Waters, F. M. Ahsan, D. M. Wolf, O. Shirihai, M. A. Teitell, Initial B Cell Activation Induces
Metabolic Reprogramming and Mitochondrial Remodeling. *iScience* **5**, 99 (2018).

23. R. Noelle, P. H. Krammer, J. Ohara, J. W. Uhr, E. S. Vitetta, Increased expression of Ia antigens on resting
B cells: an additional role for B-cell growth factor. *Proceedings of the National Academy of Sciences* **81**,
6149–6153 (1984).

24. S. R. Dillon, C. Sprecher, A. Hammond, J. Bilsborough, M. Rosenfeld-Franklin, S. R. Presnell, H. S.
Haugen, M. Maurer, B. Harder, J. Johnston, S. Bort, S. Mudri, J. L. Kuijper, T. Bukowski, P. Shea, D. L.
Dong, M. Dasovich, F. J. Grant, L. Lockwood, S. D. Levin, C. LeCiel, K. Waggie, H. Day, S. Topouzis, J.
Kramer, R. Kuestner, Z. Chen, D. Foster, J. Parrish-Novak, J. A. Gross, Interleukin 31, a cytokine produced
by activated T cells, induces dermatitis in mice. *Nature Immunology 2004 5:7* **5**, 752–760 (2004).

25. D. Chen, Y. Wang, G. K. Manakkat Vijay, S. Fu, C. W. Nash, D. Xu, D. He, N. Salomonis, H. Singh, H.
Xu, Coupled analysis of transcriptome and BCR mutations reveals role of OXPHOS in affinity maturation.
Nature Immunology 2021 22:7 **22**, 904–913 (2021).

26. S. C. Choi, L. Morel, Immune metabolism regulation of the germinal center response. *Experimental &*
Molecular Medicine 2020 52:3 **52**, 348–355 (2020).

27. A. Turqueti-Neves, M. Otte, O. P. da Costa, U. E. Höpken, M. Lipp, T. Buch, D. Voehringer, B-cell-
intrinsic STAT6 signaling controls germinal center formation. *Eur J Immunol* **44**, 2130–2138 (2014).

28. C. AF, S. K, T. KM, K. M, A. J, B. F, M. ICM, Pinpointing IL-4-independent acquisition and IL-4-
influenced maintenance of Th2 activity by CD4 T cells. *Eur J Immunol* **34** (2004).

29. G. v Zuccarino-Catania, S. Sadanand, F. J. Weisel, M. M. Tomayko, H. Meng, S. H. Kleinstein, K. L. Good-
Jacobson, M. J. Shlomchik, CD80 and PD-L2 define functionally distinct memory B cell subsets that are
independent of antibody isotype. *Nat Immunol* **15**, 631–637 (2014).

30. J. J. Taylor, R. J. Martinez, P. J. Titcombe, L. O. Barsness, S. R. Thomas, N. Zhang, S. D. Katzman, M. K.
Jenkins, D. L. Mueller, Deletion and anergy of polyclonal B cells specific for ubiquitous membrane-bound
self-antigen. *J Exp Med* **209**, 2065–77 (2012).

31. J. J. Taylor, K. A. Pape, M. K. Jenkins, A germinal center-independent pathway generates unswitched
memory B cells early in the primary response. *J Exp Med* **209**, 597–606 (2012).

32. L. Duan, D. Liu, H. Chen, M. A. Mintz, M. Y. Chou, D. I. Kotov, Y. Xu, J. An, B. J. Laidlaw, J. G. Cyster,
Follicular dendritic cells restrict interleukin-4 availability in germinal centers and foster memory B cell
generation. *Immunity* **54**, 2256-2272.e6 (2021).

33. J. D. Mountz, M. Gao, D. M. Ponder, S. Liu, C. W. Sun, F. Alduraibi, K. Sullivan, B. Pat, L. J. Dell'Italia,
H. C. Hsu, IL-4 receptor blockade is a global repressor of naïve B cell development and responses in a
dupilumab-treated patient. *Clin Immunol* **244** (2022).

34. M. Sarfati, S. Fournier, M. Christoffersen, G. Biron, Expression of CD23 antigen and its regulation by IL-4
in chronic lymphocytic leukemia. *Leuk Res* **14**, 47–55 (1990).

35. J.-S. He, M. Meyer-Hermann, D. Xiangying, L. Y. Zuan, L. A. Jones, L. Ramakrishna, V. C. de Vries, J.
Dolpady, H. Aina, S. Joseph, S. Narayanan, S. Subramaniam, M. Puthia, G. Wong, H. Xiong, M. Poidinger,
J. F. Urban, J. J. Lafaille, M. A. Curotto de Lafaille, The distinctive germinal center phase of IgE+ B
lymphocytes limits their contribution to the classical memory response. *Journal of Experimental Medicine*
210, 2755–2771 (2013).

36. Z. Yang, M. J. Robinson, C. D. C. Allen, Regulatory constraints in the generation and differentiation of IgE-
expressing B cells. *Curr Opin Immunol* **28**, 64–70 (2014).

37. A. K. Gloudemans, M. Plantinga, M. Guilliams, M. A. Willart, A. Ozir-Fazalalikhan, A. van der Ham, L.
Boon, N. L. Harris, H. Hammad, H. C. Hoogsteden, M. Yazdanbakhsh, R. W. Hendriks, B. N. Lambrecht,
H. H. Smits, The Mucosal Adjuvant Cholera Toxin B Instructs Non-Mucosal Dendritic Cells to Promote
IgA Production Via Retinoic Acid and TGF-β. *PLoS One* **8**, e59822 (2013).

38. T. M. McIntyre, D. R. Klinma, P. Rohman, M. Lugo, J. R. Dasch, J. J. Mond, C. M. Snapper, Transforming
growth factor beta 1 selectivity stimulates immunoglobulin G2b secretion by lipopolysaccharide-activated
murine B cells. *Journal of Experimental Medicine* **177**, 1031–1037 (1993).

39. C. M. Snapper, W. E. Paul, Interferon-γ and B Cell Stimulatory Factor-1 Reciprocally Regulate Ig Isotype
Production. *Science (1979)* **236**, 944–947 (1987).

40. A. Bossie, E. S. Vitetta, IFN-gamma enhances secretion of IgG2a from IgG2a-committed LPS-stimulated
murine B cells: implications for the role of IFN-gamma in class switching. *Cell Immunol* **135**, 95–104
(1991).

41. D. Esterházy, M. C. C. Canesso, L. Mesin, P. A. Muller, T. B. R. de Castro, A. Lockhart, M. ElJalby, A. M.
C. Faria, D. Mucida, Compartmentalized gut lymph node drainage dictates adaptive immune responses.
Nature 2019 569:7754 **569**, 126–130 (2019).

42. B. Zhang, E. Liu, J. A. Gertie, J. Joseph, L. Xu, E. Y. Pinker, D. A. Waizman, J. Catanzaro, K. H. Hamza,
K. Lahl, U. Gowthaman, S. C. Eisenbarth, Divergent T follicular helper cell requirement for IgA and IgE
production to peanut during allergic sensitization. *Sci Immunol* **5** (2020).

43. Q. Chen, M. Xie, H. Liu, A. L. Dent, Development of allergen-specific IgE in a food-allergy model requires
precisely timed B cell stimulation and is inhibited by Fgl2. *Cell Rep* **39**, 110990 (2022).

44. M. Vajdy, M. H. Kosco-Vilbois, M. Kopf, G. Köhler, N. Lycke, Impaired mucosal immune responses in
interleukin 4-targeted mice. *J Exp Med* **181**, 41–53 (1995).

45. Z. Chen, A. M. Collins, Y. Wang, B. A. Gata, Clustering-based identification of clonally-related
immunoglobulin gene sequence sets. *Immunome Res* **6**, 1–7 (2010).

46. N. T. Gupta, J. A. Vander Heiden, M. Uduman, D. Gadala-Maria, G. Yaari, S. H. Kleinstein, Change-O: a
toolkit for analyzing large-scale B cell immunoglobulin repertoire sequencing data. *Bioinformatics* **31**,
3356–3358 (2015).

47. Z. Yang, C.-A. M. Wu, S. Targ, C. D. C. Allen, IL-21 is a broad negative regulator of IgE class switch
recombination in mouse and human B cells. *Journal of Experimental Medicine* **217** (2020).

48. R. Jiménez-Saiz, Y. Ellenbogen, K. Bruton, P. Spill, D. D. Sommer, H. Lima, S. Waserman, S. U. Patil, W.
G. Shreffler, M. Jordana, Human BCR analysis of single-sorted, putative IgE+ memory B cells in food
allergy. *J Allergy Clin Immunol* **144**, 336-339.e6 (2019).

49. L. Shehata, C. D. Thouvenel, B. D. Hondowicz, L. A. Pew, D. J. Rawlings, J. Choi, M. Pepper, IL-4
downregulates BCL6 to promote memory B cell selection in germinal centers. *bioRxiv*, 2023.01.26.525749
(2023).

50.   D. G. Gonzalez, C. M. Cote, J. R. Patel, C. B. Smith, Y. Zhang, K. M. Nickerson, T. Zhang, S. M. Kerfoot,
A. M. Haberman, Nonredundant Roles of IL-21 and IL-4 in the Phased Initiation of Germinal Center B
Cells and Subsequent Self-Renewal Transitions. *J Immunol* **201**, 3569–3579 (2018).

51.   B. G. De Jong, H. Ijspeert, L. Marques, M. Van Der Burg, J. J. M. Van Dongen, B. G. Loos, M. C. Van
Zelm, Human IgG2- and IgG4-expressing memory B cells display enhanced molecular and phenotypic signs
of maturity and accumulate with age. *Immunol Cell Biol* **95**, 744 (2017).

52.   A. M. Collins, K. J. L. Jackson, A Temporal Model of Human IgE and IgG Antibody Function. *Front
Immunol* **4** (2013).

53.   D. MacGlashan, R. G. Hamilton, Parameters Determining the Efficacy of CD32 to Inhibit Activation of
FcεRI in Human Basophils. *J Allergy Clin Immunol* **137**, 1256 (2016).

54.   M. Sugimoto, N. Kamemura, M. Nagao, M. Irahara, S. Kagami, T. Fujisawa, H. Kido, Differential response
in allergen-specific IgE, IgGs, and IgA levels for predicting outcome of oral immunotherapy. *Pediatr
Allergy Immunol* **27**, 276–282 (2016).

55.   J. J. Heeringa, C. I. McKenzie, N. Varese, M. Hew, A. T. C. M. Bakx, P. M. Aui, J. M. Rolland, R. E.
O'Hehir, M. C. van Zelm, Induction of IgG2 and IgG4 B-cell memory following sublingual immunotherapy
for ryegrass pollen allergy. *Allergy* **75**, 1121 (2020).

56.   O. T. Burton, S. L. Logsdon, J. S. Zhou, J. Medina-Tamayo, A. Abdel-Gadir, M. Noval Rivas, K. J.
Koleoglou, T. A. Chatila, L. C. Schneider, R. Rachid, D. T. Umetsu, H. C. Oettgen, Oral immunotherapy
induces IgG antibodies that act via FcγRIIb to suppress IgE-mediated hypersensitivity. *J Allergy Clin
Immunol* **134**, 1310 (2014).

57.   E. Wambre, V. Bajzik, J. H. DeLong, K. O'Brien, Q.-A. Nguyen, C. Speake, V. H. Gersuk, H. A. DeBerg,
E. Whalen, C. Ni, M. Farrington, D. Jeong, D. Robinson, P. S. Linsley, B. P. Vickery, W. W. Kwok, A
phenotypically and functionally distinct human TH2 cell subpopulation is associated with allergic disorders.
Sci Transl Med **9**, eaam9171 (2017).

58.   C. Bangert, K. Rindler, T. Krausgruber, N. Alkon, F. M. Thaler, H. Kurz, T. Ayub, D. Demirtas, N.
Fortelny, V. Vorstandlechner, W. M. Bauer, T. Quint, M. Mildner, C. Jonak, A. Elbe-Bürger, J. Griss, C.
Bock, P. M. Brunner, Persistence of mature dendritic cells, T H 2A, and Tc2 cells characterize clinically
resolved atopic dermatitis under IL-4Rα blockade. *Sci Immunol* **6** (2021).

59.   H. Renz, J. Domenico, E. W. Gelfand, IL-4-dependent up-regulation of IL-4 receptor expression in murine T
and B cells. *Journal of Immunology* **146**, 3049–3055 (1991).

60.   J. Holm, N. Willumsen, P. A. Würtzen, L. H. Christensen, K. Lund, Facilitated antigen presentation and its
inhibition by blocking IgG antibodies depends on IgE repertoire complexity. *J Allergy Clin Immunol* **127**,
1029–37 (2011).

61.   C. Liu, K. Richard, M. Wiggins, X. Zhu, D. H. Conrad, W. Song, CD23 can negatively regulate B-cell
receptor signaling. *Sci Rep* **6**, 25629 (2016).

62.   F. Horns, C. Vollmers, D. Croote, S. F. Mackey, G. E. Swan, C. L. Dekker, M. M. Davis, S. R. Quake,
Lineage tracing of human B cells reveals the in vivo landscape of human antibody class switching. *Elife* **5**
(2016).

63.   P. Irrgang, J. Gerling, K. Kocher, D. Lapuente, P. Steininger, K. Habenicht, M. Wytopil, S. Beileke, S.
Schäfer, J. Zhong, G. Ssebyatika, T. Krey, V. Falcone, C. Schülein, A. S. Peter, K. Nganou-Makamdop, H.
Hengel, J. Held, C. Bogdan, K. Überla, K. Schober, T. H. Winkler, M. Tenbusch, Class switch toward
noninflammatory, spike-specific IgG4 antibodies after repeated SARS-CoV-2 mRNA vaccination. *Sci
Immunol* **8** (2023).

64.   B. P. Vickery, J. Lin, M. Kulis, Z. Fu, P. H. Steele, S. M. Jones, A. M. Scurlock, G. Gimenez, L. Bardina, H.
A. Sampson, A. W. Burks, Peanut oral immunotherapy modifies IgE and IgG4 responses to major peanut
allergens. *J Allergy Clin Immunol* **131**, 128-34.e1–3 (2013).

65.   P. Varshney, S. M. Jones, A. M. Scurlock, T. T. Perry, A. Kemper, P. Steele, A. Hiegel, J. Kamilaris, S.
Carlisle, X. Yue, M. Kulis, L. Pons, B. Vickery, A. W. Burks, A randomized controlled study of peanut oral
immunotherapy: clinical desensitization and modulation of the allergic response. *J Allergy Clin Immunol*
127, 654–60 (2011).

66.   T. Rispens, M. G. Huijbers, The unique properties of IgG4 and its roles in health and disease. *Nature
Reviews Immunology 2023 23:11* **23**, 763–778 (2023).

67.   K. Arias, D. K. Chu, K. Flader, F. Botelho, T. D. Walker, N. Arias, A. A. Humbles, A. J. Coyle, H. C.
Oettgen, H.-D. Chang, N. van Rooijen, S. Waserman, M. Jordana, Distinct immune effector pathways
contribute to the full expression of peanut-induced anaphylactic reactions in mice. *J Allergy Clin Immunol*
127, 1552–61.e1 (2011).

68. J. Sun, K. Arias, D. Alvarez, R. Fattouh, T. Walker, S. Goncharova, B. Kim, S. Waserman, J. Reed, A. J.
Coyle, M. Jordana, Impact of CD40 Ligand, B Cells, and Mast Cells in Peanut-Induced Anaphylactic
Responses. *The Journal of Immunology* **179** (2007).
69. K. B. Hoehn, J. A. Vander Heiden, J. Q. Zhou, G. Lunter, O. G. Pybus, S. H. Kleinstein, Repertoire-wide
phylogenetic models of B cell molecular evolution reveal evolutionary signatures of aging and vaccination.
Proc Natl Acad Sci U S A **116**, 22664–22672 (2019).
70. K. B. Hoehn, O. G. Pybus, S. H. Kleinstein, Phylogenetic analysis of migration, differentiation, and class
switching in B cells. *PLoS Comput Biol* **18** (2022).
71. B. Kaminow, D. Yunusov, A. Dobin, STARsolo: accurate, fast and versatile mapping/quantification of
single-cell and single-nucleus RNA-seq data. *bioRxiv*, 2021.05.05.442755 (2021).
72. L. Song, D. Cohen, Z. Ouyang, Y. Cao, X. Hu, X. S. Liu, TRUST4: immune repertoire reconstruction from
bulk and single-cell RNA-seq data. *Nature Methods 2021 18:6* **18**, 627–630 (2021).
73. A. Yermanos, A. Agrafiotis, R. Kuhn, D. Robbiani, J. Yates, C. Papadopoulou, J. Han, I. Sandu, C. Weber,
F. Bieberich, R. Vazquez-Lombardi, A. Dounas, D. Neumeier, A. Oxenius, S. T. Reddy, Platypus: an open-
access software for integrating lymphocyte single-cell immune repertoires with transcriptomes. *NAR Genom
Bioinform* **3** (2021).

Acknowledgments: We kindly thank S. Eisenbarth for thoughtful review of the manuscript. We

also thank M. Subapanditha and for assistance with cell sorting and C. Mader, the

McMaster Genomics Facility, and G. Kongsgaard Koed, and J. Skovsgaard for

sequencing support. Experimental schematics were created with BioRender

(www.biorender.com). KB was supported by a CIHR CGS-Doctoral award and CIHR

Postdoctoral Fellowship. AP was supported by the Schroder Allergy & Immunology

Research Institute Scholarship and an Ontario Graduate Scholarship.

Funding:

Schroeder Foundation (SW, MJ)

Food Allergy Canada (SW, MJ)

ALK Abelló (SW, MJ)

Canadian Allergy, Asthma, and Immunology Foundation (MJ)

Michael Zych Family (SW, MJ)

Author contributions:

Conceptualization: KB, AP, MJ

Investigation: KB, AP, AA, TDW, JW, SSK-B, AF, OM-D, SG, EG, JFEK, AG

Visualization: KB, AP

Funding acquisition: SW, PSA, MJ

Supervision: MJ

Writing – original draft: KB, AP

Writing – review & editing: KB, AP

Competing interests: JFEK, SW, and MJ receive funding from ALK Abelló A/S. All other authors declare that they have no competing interests.

Data and materials availability: This manuscript produced no new code. Single-cell and bulk sequencing data will be deposited to the GEO database prior to publication. All other data are available within the main text or the supplementary materials.

Fig. 1. **IL-4Rα signaling augments secondary germinal center activity. (A)** Study schematic.

(B) UMAP of B cell clusters from scRNA-seq analysis of B cells isolated from spleens and

inguinal LNs of CT-immunized mice and A4RA/IC-treated epicutaneously sensitized mice. $n = 5$

CT, $n = 10$ IC, $n = 10$ A4RA. **(C)** UMAP plots of genes defining naïve and MBC clusters. **(D)**

Percent of cells in each cluster by treatment group. **(E)** Expression level of gene modules in

naïve B cell clusters 1-5 (grey) and MBCs (green) by treatment group. Point indicates median.

(F) VDJ analysis of MBCs identifying the number of nucleotide mutations compared to the

germline sequence. Shading in circle plots depict the fraction of cells with each number of

mutations. Mean (SEM) number of mutations are displayed within the circles. **(G)**

Representative contour plots of enriched OVA^+ B cell population from inguinal LNs of

epicutaneously sensitized mice. Pre-gated as live > singlet > $F4/80^-$ $CD3^-$ > $B220^+$. **(H)** Summary

plots of OVA^+ B cell total count and frequency. **(I)** Representative contour plot of resting

($CD38^+$ $GL7^-$), transitional ($CD38^+$ $GL7^+$) and germinal center ($CD38^-$ $GL7^+$) B cells. Pre-gated

from OVA^+ B cells > IgM^- IgD^-. **(J)** Summary plots of OVA^+ B cell frequency within the

populations from (I) at three days post-boost. **(K)** Summary plot of OVA^+ B cell frequency

within transitional (empty) and germinal center (filled) populations in post-boost time course.

Data represent 2 experiments, each with 4 mice per group. Data are presented as median ±

min/max (H, J) or mean ± SEM (K). ** $p < 0.01$, *** $p < 0.001$.

Fig. 2. MBC2 phenotype is ameliorated with IL-4Rα blockade in mice and humans. (A) Overview of analyzed datasets. **(B)** UMAP of B cell clusters from scRNA-seq data sets. Colored points indicate class-switched cluster used for downstream analyses. **(C)** Expression level of MBC2-related genes in MBC clusters. **(D)** Representative contour plots of OVA$^+$ MBC2s from inguinal LNs of epicutaneously sensitized mice four weeks after re-exposure and antibody treatment. Pre-gated as OVA$^+$ B cells > CD38$^+$ GL7$^-$ > IgM$^-$ IgD$^-$. **(E)** Summary plots of MBC2 and CD23$^+$ IL-4R^{low} MBC frequency from IC and A4RA-treated epicutaneously sensitized mice. **(F)** MFI of CD23 within OVA$^+$ (left) and OVA$^-$ (right) MBCs from IC and A4RA-treated epicutaneously sensitized mice. Pre-gated as OVA$^{+/-}$ B cells > CD38$^+$ GL7$^-$ > IgM$^-$ IgD$^-$. **(G)** Representative contour plots of MBC2s from human PBMCs 8 days following *in vitro* allergen stimulation and treatment with IC or A4RA. Pre-gated as live > singlet > CD3$^-$ CD14$^-$ CD16$^-$ CD19$^+$ > IgD$^-$ IgM$^-$ > CD38$^{low-med}$. **(H)** Summary plots of human MBC2 frequency in **G**, separated by CD27 expression. **(I)** MFI of CD23 within CD27$^{+/-}$ class-switched B cells from human PBMCs 8 days following *in vitro* allergen stimulation and treatment with IC or A4RA. Data represents 2-3 experiments with 5 mice per group and are presented as median ± min/max (E, F) or 2 experiments (n = 3 blood donors) with each line representing a single donor (H, I). * p < 0.05, ** p < 0.01, *** p < 0.001. Ab, antibody; Ag, antigen; LSC, lymphoid stromal cell.

Fig. 3. Longitudinal analysis of Ag-specific B cell responses with absent IL-4 signaling. (A)
Expression level of *Ighc* genes downstream of *Ighg1* from scRNA-seq analysis of B cells
isolated from spleens and inguinal LNs of CT-immunized mice and A4RA/IC-treated
epicutaneously sensitized mice. Gene expression is plotted for cells within the MBC cluster (Fig.
1B). **(B)** VDJ analysis of MBC cluster. Plot depicts the assigned *Ighc* gene in MBCs with ≥3
nucleotide mutations compared to the germline sequence. **(C)** Representative contour plots of
IgG1$^+$ and IgG2c$^+$ populations from inguinal LNs of IC/A4RA-treated epicutaneously sensitized
mice four weeks post-boost and antibody treatment. **(D)** IC/A4RA-treated mice were rested up to
15 weeks after re-exposures. IgG1$^+$ (left) and IgG2c$^+$ (right) MBC frequency are plotted at four
and 15 weeks post-boost. Pre-gated as OVA$^+$ B cells > CD38$^+$ GL7$^-$ > IgM$^-$ IgD$^-$. **(E)**
Epicutaneously sensitized mice were treated with A4RA, AIFNG, or both at the time of
subcutaneous re-exposure. IgG1$^+$ (left) and IgG2c$^+$ (right) MBC frequency are plotted at four
weeks post-boost. Pre-gated as OVA$^+$ B cells > CD38$^+$ GL7$^-$ > IgM$^-$ IgD$^-$. **(F)** OVA-specific
serum immunoglobulin profiles of IC/A4RA-treated mice at zero to 15 weeks post-boost. **(G)**
IC/A4RA-treated mice were rested for nine weeks to allow for clearance of OVA-specific IgG2c
and clearance of IC/A4RA blocking antibodies. Subsequently, mice received tertiary OVA
exposures and serum was collected at week 12. OVA-specific serum immunoglobulin profiles
are depicted before and after tertiary exposures. (C-G) Data represents 2 experiments with at
least 4 mice per group. Data are presented as mean ± SEM (F, G) or median ± min/max (D, E). *
p < 0.05, ** *p* < 0.01, *** *p* < 0.001. Seq, sequences; swIg, class-switched; AIFNG, anti-IFN-
γR1; OD, optical density.

A Paired V-J gene usage

B Isotype Distribution of Shared Clones

IC — *IGHG1⁺*: 82.4, 5.1, 11.6, 0.9
A4RA — *IGHG1⁺*: 84.7, 3.4, 10.4, 1.5
IC — *IGHG2⁺*: 88.3, 6.6, 3.5, 1.6
A4RA — *IGHG2⁺*: 84.5, 9.1, 3.3, 3.1

Legend: *IGHM/D*, *IGHG3*, *IGHG1*, *IGHG2*, *IGHA*

C
IGHV1-26 / IGHJ1
IGHV1-50 / IGHJ4
IGHV1-76 / IGHJ2

D IC A4RA

Expression Level

Ifitm2, *Cd63*, *Ifi203*, *Mars*, *Ets1*, *Jak1*, *Ptpn2*

Ighm/d, *Ighm/d*

E IC A4RA

Ifitm2, *Cd63*, *Ifi203*, *Mars*, *Ets1*, *Jak1*, *Ptpn2*

Ighg1, *Ighg1*, *Ighg2b/c*

Fig. 4. Clonal analysis of OVA-enriched B cells. WT mice were epicutaneously sensitized with

OVA, then re-exposed s.c. to OVA in the presence of A4RA or IC. Four weeks post-boost,

spleens and inguinal LNs were collected. B cells from the OVA tetramer-enriched fraction were

sorted for bulk VDJ sequencing. **(A)** Correlation matrix of paired IGHV-IGHJ gene usage

between isotypes in IC and A4RA. Black boxes highlight the relationships between IgG2 and

upstream isotypes. **(B)** Circle plots showing the isotype distribution within clonal families

containing IGHG1 (top) and IGHG2 (bottom). **(C)** Representative clonal lineage trees containing

IGHG2 sequences and upstream isotypes (IGHM/D, IGHG3, and IGHG1) from A4RA-treated

mice. Scale indicates IGHV SHM frequency. **(D, E)** OVA-specific B cells were single-cell

sorted from epicutaneously sensitized mice at four weeks post-boost for Smart-Seq. $n = 40$ IC, n

$= 41$ A4RA. Expression level of select DEG are shown for unswitched (D) and IGHG1/2$^+$ (E)

OVA-specific B cells. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

A

Donors: WT (CD45.1)

OVA-Enriched
swIg⁺ MBCs

Recip: muMT (CD45.2)

or

Ag-Specific
Response

○ OVA + Alum   ● OVA + CT   ● OVA + CpG   ⊢—⊣ 1 Week

B

IgG1

Alum   CT   CpG

87.70   67.43   16.72

2.93   4.05   61

IgG2b

0.64   1.99   10.11

IgG2c

C

% of OVA⁺ swIg MBCs

IgG1⁺   IgG2c⁺

*

D

OD

OVA-IgG2b   OVA-IgG2c   OVA-IgE

*   *

Fig. 5. Boost with CpG produces IgG2 response from type 2-primed class-switched MBCs.

(A) Adoptive transfer schematic. **(B)** WT mice were immunized with OVA + Alum, CT, or CpG. Representative contour plots class-switched OVA$^+$ MBCs. Pre-gated as OVA$^+$ B cells > CD38$^+$ GL7$^-$ > IgM$^-$ IgD$^-$. Numbers indicate the proportion of IgG1$^+$, IgG2b$^+$, and IgG2c$^+$ cells from the parent class-switched MBC population. **(C)** OVA-enriched B cells were adoptively transferred from WT OVA-Alum donor mice to naïve muMT mice and recipient were boosted with either OVA + CT or OVA + CpG. Summary plots show OVA$^+$ IgG1$^+$ and OVA$^+$ IgG2c$^+$ B cell frequency in recipient mice one week post-transfer and boost. **(D)** OVA-specific serum immunoglobulin profiles in recipient mice one week post-transfer and boost. Data represents 2 experiments, each with 2-4 mice per group. Data are presented as median ± min/max (B) or mean ± SEM (D). * $p < 0.05$.