When the host's away, the pathogen will play:
the protective role of the skin microbiome during hibernation

Troitsky TS[1], Laine VN[1], Lilley TM[1]

1. BatLab Finland, Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland
Corresponding author: Thomas Lilley (thomas.lilley@helsinki.fi)

ABSTRACT

The skin of animals is enveloped by a symbiotic microscopic ecosystem known as the microbiome. The host and microbiome exhibit a mutualistic relationship, collectively forming a single evolutionary unit sometimes referred to as a holobiont. Although the holobiome theory highlights the importance of the microbiome, little is known about how the skin microbiome contributes to protecting the host. Existing studies focus on humans or captive animals, but research in wild animals is in its infancy. Specifically, the protective role of the skin microbiome in hibernating animals remains almost entirely overlooked. This is surprising, considering the massive population declines in hibernating North American bats caused by the fungal pathogen *Pseudogymnoascus destructans*, which causes white-nose syndrome. Hibernation offers a unique setting in which to study the function of the microbiome because, during torpor, the host's immune system becomes suppressed, making it susceptible to infection. We conducted a systematic review of peer-reviewed literature on the protective role of the skin microbiome in non-human animals. We selected 230 publications that mentioned pathogen inhibition by microbes residing on the skin of the host animal. We found that the majority of studies were conducted in North America and focused on the bacterial microbiome of amphibians infected by the chytrid fungus. Despite mentioning pathogen inhibition by the skin microbiome, only 30,4 % of studies experimentally tested the actual antimicrobial activity of symbionts. Additionally, only 7,8 % of all publications studied defensive cutaneous symbionts during hibernation. With this review, we want to highlight the knowledge gap surrounding skin microbiome research in hibernating animals. For instance, research looking to mitigate the effects of white-nose syndrome in bats should focus on the antifungal microbiome of Palearctic bats, as they survive exposure to the *Pseudogymnoascus destructans* -pathogen during hibernation. We also recommend future studies prioritize lesser-known microbial symbionts, such as fungi, and investigate the effects of a combination of anti-pathogen microbes, as both areas of research show promise as probiotic treatments. By incorporating the protective skin microbiome into disease mitigation strategies, conservation efforts can be made more effective.

KEY WORDS: skin microbiome, hibernation, symbiosis, wildlife disease, probiotics, holobiont, chiroptera, amphibian, white-nose syndrome, chytridiomycosis

INTRODUCTION

Animals are constantly under attack from a plethora of microorganisms that have the potential to cause disease and even mortality. However, to infect an animal, these microbes have to first permeate the skin, which is the primary barrier between the host and the environment [1, 2]. The skin is a cool, acidic environment that is covered by sebaceous glands that secrete an antimicrobial substance, sebum. Sebum lubricates the skin and facilitates the growth of commensal microbes such as archaea, bacteria, viruses, and fungi. Together these microbes form a mutualistic community referred to as the skin microbiome [1, 3]. The host provides the symbionts with a favorable environment to propagate in and the microbes contribute by helping the host heal wounds, educating the immune system and preventing colonization of new microbes, which may have pathogenic properties [1, 2, 4, 5].

Microbiome research has increased in popularity in recent years with studies focusing mainly on the beneficial role of the gut, oral and skin microbiome of organisms, notably in humans [6, 7]. These studies give support to the holobiome theory, which suggests the host and its microbiome can be viewed together as a single evolutionary unit instead of separate entities [8, 9]. This perspective changes the definition of an individual to include the microorganisms living in and on the host. In many regards, the host cannot survive without its microbial symbionts, which also outnumber the cells of the host [10, 11]. This obligatory symbiosis also exists in animals, plants, and various other organisms [8]. Since the genomes of the microbes contributing to the microbiome evolve faster than the genome of the host, it can play a fundamental role in the host's ability to rapidly adapt to environmental disturbances and new potentially pathogenic microbes [12]. This may be an important adaptation as climate change exposes species to novel pathogens.

The skin microbiome in particular is very sensitive to changes both in the environment and the host [1], which affects the holobiont's ability to respond to changes. *Dysbiosis* or disruption in the composition of the skin microbiome can cause an imbalance that has a negative effect on host survival [2]. Dysbiosis often occurs when the amount of commensal microbes is reduced due to factors like immune deficiencies or exposure to pathogens, resulting in the microbiome losing its ability to protect the host [13]. For example, the diversity of the sheep (*Ovis aries*) skin microbiome is known to decrease preceding the onset of foot rot [14]. In addition, the artificial reduction of skin microbiome richness in salamanders before exposure to the deadly fungal pathogen that causes chytridiomycosis (*Batrachochytrium dendrobatidis,* hereafter *Bd*) leads to higher mortality [15]. In general, tropical amphibian species threatened by chytridiomycosis have lower skin bacterial diversity than non-threatened species [16].

On the other hand, the enrichment of certain microbes can be beneficial to the holobiont. Antimicrobial bacteria that inhibit pathogen growth *in vitro* have been found on the skin of amphibians, reptiles, fish, and mammals [17–20]. These bacteria, along with other protective microbes, are often referred to as probiotics. Testing the inhibition ability of these bacteria is becoming exceedingly popular in amphibians [21–23], because chytridiomycosis has

caused major population declines in both the Americas and Eastern Australia [24, 25]. Mutualistic bacteria living on amphibian skin are known to produce antimicrobial agents, such as violacein and prodigiosin, that can inhibit the growth of *Bd* and suppress inflammation [13, 26, 27]. Thus, both positive and negative changes in skin microbiome composition seem to have a direct effect on the fitness of the host organism.

Due to its warm and moist nature, the skin provides an ideal environment for fungi to grow on, simultaneously making the skin more susceptible to fungal infections [28]. Over the past three decades, wildlife populations have experienced unprecedented, high-profile declines due to emerging infectious fungal diseases such as chytridiomycosis [29]. Another example of a deadly, skin-infecting mycosis that could potentially be treated with probiotics is white-nose syndrome (WNS) in insectivorous, hibernatory bats. WNS is caused by the psychrophilic fungus *Pseudogymnoascus destructans* (hereafter *Pd*), which invades and infects the skin causing a distinct fungal growth on the wings and muzzle of hibernating bats during winter [30, 31]. The fungal propagation arouses bats from torpor depleting their fat reserves, and eventually leading to starvation during a period when minimal insect-food is available. The disease was first discovered in the winter of 2006-2007 in New York and it has devastated Nearctic bat populations ever since, endangering once abundant species, such as the little brown bat (*Myotis lucifugus)* [32, 33].

The reason WNS has had such a calamitous effect on Nearctic bat populations can be attributed to the pathogen infecting bats when they are most vulnerable, during hibernation. The body temperature of hibernating bats drops drastically to resemble that of the ambient temperature in the hibernacula (2-14 °C) [30, 34]. Bats are, therefore, heterothermic, meaning they switch between an endothermic active state to an exothermic torpor state [35]. This radical change in thermoregulation is comparable to the ectothermic strategy of amphibians since both bats and their skin microbiome must tolerate substantial temperature fluctuations. This poses an added burden to both the host and its skin microbiome.

In addition, the metabolism and immune system of a bat become suppressed during hibernation, because they are energetically costly [36, 37]. This is exemplified by a significant decrease in the number of circulating leukocytes in the bloodstream during torpor [37]. Hibernation is an optimal strategy for insectivorous bats to save energy when food is scarce, and bats can remain torpid from days to months without eating [35]. However, the ability of the bat to defend itself against pathogens during this time becomes reduced due to its down-regulated immune system. Although most microscopic pathogens do not propagate well in cold temperatures [37], *Pd* thrives in the approximate temperature bats hibernate in, posing a significant threat [34].

However, not all bats get infected when exposed to *Pd*. In the Palearctic, where the fungus originates, bats tolerate exposure to the pathogen without infection or mortality [38, 39]. Species such as the greater mouse-eared bat (*Myotis myotis*) can tolerate high pathogen loads without apparent negative consequences [39, 40], suggesting the parasitic relationship has evolved into something that more resembles commensalism [41, 42]. One hypothesis to

explain this phenomenon is that Palearctic bats have evolved a tolerance due to their longer history of exposure to the pathogen [43]. Molecular evidence implies Palearctic bats have been exposed to *Pd* for an extensive period of time, while Nearctic bats have had a mere 20-year bout with the pathogen since it was introduced from Europe [44, 45]. The protective skin microbiome could have enabled bat populations in the Palearctic to endure *Pd* exposure until the host develops tolerance.

Hibernation offers a unique setting in which to study the protective role of the skin microbiome because as the bat is in a torpid state, the microbiome may remain active. The symbiotic bacteria living on the skin of bats benefit from host survival, thus, it is not surprising that several of these bacterial strains have been found to have antifungal properties that may inhibit the growth of *Pd* [19, 46, 47]. For example, the bacterial genus *Pseudomonas* that is commonly found on bat skin has been shown to inhibit the growth of *Pd* both *in vitro* [46–49] and *in vivo* [50, 51]. Viewing Palearctic bats as holobionts that have coevolved together with *Pd* can help explain how selection might have favored bats harboring these antifungal bacteria in abundance on their skin. It is also noteworthy to mention that many other animals, such as some frogs, snakes, bears, rodents, birds, and fish possess the ability to hibernate, exposing them to similar risks as bats [52–57]. Therefore, studying the composition and antifungal potential of the skin microbiome during hibernation is an exclusive opportunity to better understand disease dynamics and the protective role of the skin microbiome in animals.

The aims of this review are to determine: *(i)* whether the protective skin microbiome of hibernating animals has been studied; *(ii)* whether experimental research studying pathogen inhibition of the skin microbiome has increased in the past years; and *(iii)* which antifungal microbes have been identified and studied the most? We emphasize the importance of experimental research because without inhibition assays and probiotic trials, the protective capacity of the microbiome remains speculative at best. To address these questions, we conducted a systematic review encompassing a range of publications examining the protective function of the skin microbiome in animals (Fig 1).

MATERIALS AND METHODS

1) Systematic literature search

We performed a comprehensive keyword search on the *Web of Science* on the 13th of June 2023. Before defining the final search terms, we did multiple exploratory trials using different search words to determine which string of words would maximize the number of relevant references without adding an excessive number of irrelevant ones. For example, adding the words "probiotic" and "bioaugmentation" as a separate and obligatory search clause captured only 81 publications. Therefore, we added the words to the previous clause, making them facultative. We conducted the final optimized search using the following terms:

ALL=("microbiota" OR "microbiome") AND ALL=("skin" OR "cutaneous" OR "epidermis" OR "dermal") AND
ALL=("resistance" OR "inhibit" OR "antifungal" OR "pathogen" OR "fungal" OR "bioaugmentation" OR
"probiotic") AND ALL=("vertebrate" OR "invertebrate" OR "animal" OR "mammal" OR "reptile" OR
"amphibian" OR "fish" OR "bird" OR "bat")

This yielded 483 publications ranging from the years 2005-2023 that were screened by TST according to the
PRISMA diagram (Fig 1.). Articles were found suitable for this review based on the following inclusion criteria:
(i) they studied the skin microbiome (as opposed to just gut or oral microbiome); (ii) they mentioned skin-
infecting pathogens and antimicrobial symbionts living on the skin of the host; (iii) they studied non-human
animals. Reviews and publications that did not meet these criteria were excluded, including one publication that
was not accessible. In addition to this, we added 21 publications found elsewhere that fit the search criteria and
seven publications that we were notified about by *Web of Science* alert, resulting in the final data set (N=230).

142
143 *Figure 1.* PRISMA diagram [99] explaining screening process of publications.
144
2) Metadata extraction

We extracted metadata from all relevant references for the final database. We documented the geographical and
taxonomic range of the studies, the host's captivity status, whether the study solely examined microbiome
composition (descriptive) or also assessed the microbiome's response to pathogens (experimental), as well as how
the microbiome's response was tested and whether pathogens were known to infect hosts during hibernation. This
was done by cross referencing literature and/or checking pathogen propagation temperatures (if optimal pathogen
propagation temperature was not similar to the temperature in hibernacula, the pathogen was not considered a
threat during hibernation). For studies that experimentally tested microbes against pathogens we also determined
the type of pathogen and antimicrobial genera detected on skin and whether the microbes were successful in
inhibiting the pathogens.

3) Data visualization and statistical analyses

We performed data analysis and visualizations using R version 4.2.2 [58] using packages 'ggplot2' version 3.4.2

[59] and 'bipartite' version 2.18 [60]. Additionally, we used Inkscape version 1.3 [61] to edit the visualizations.

We used a binomial generalized linear model to analyze how many of the publications actually experimentally

tested the pathogen inhibition ability of the microbiome, in proportion to all published studies over the past years

(glm(formula = cbind(experimental, descriptive) ~ year, family = "binomial")). We excluded the year 2023 from

the analysis, since the year is not over, and more studies are likely to be published before the end of the year.

RESULTS

General summary of literature

In our initial *Web of Science* search, we identified 483 publications. Following the screening of titles and

abstracts, 318 were considered relevant and underwent full text inspection. Among these, 202 met our inclusion

criteria and seven publications were added after a Web of Science alert. An additional 21 studies were added from

other sources, resulting in 230 publications (Supplementary file 1).

The majority of skin microbiome studies were conducted in the Western Hemisphere, with 59,1 % of studies

taking place in North America and 8,3 % in South America (Fig 2). The remaining 32,6 % of studies were spread

between Europe (13,5 %), Asia (13,0 %), Africa (2,2 %), and Oceania (3,0 %). Additionally, two studies (0,9 %)

sampled animals from multiple continents.

Predictably, the most studied animal class was amphibians (59,6 %), followed by ray-finned fishes (19,6 %), and

mammals (14,8 %). The remaining 6,0 % of studied species were divided into reptiles (2,6 %), cartilaginous

fishes (1,7 %), and birds (0,4 %) with three studies investigating multiple classes (1,3 %). About half (52,0 %) of

the publications studied the skin microbiome of wild animals, while 42,3 % focused on captive animals, and 5,7%

considered both.

Despite the search set to contain at least one word regarding pathogen inhibition ("antifungal", "pathogen",

"resistance", "inhibit", "fungal", "bioaugmentation" or "probiotic"), most of the publications (69,6 %) only

described the skin microbiome composition of the host, without testing the inhibition ability of potentially

antifungal bacteria found on the skin. Altogether, only 30,4 % of publications tested inhibition ability by either

conducting inhibition assays *in vitro* or testing probiotic treatments *in vivo*. Out of these studies 65,2 % tested

inhibitory ability using inhibition assays, 24,2 % used probiotic treatment, and 10,6 % used both. Among these

studies 60,6 % experimented on antifungal amphibian symbionts, 21,2 % on mammalian symbionts, 16,7 % on

fish symbionts, and 1,5 % on reptile symbionts. The majority (54,7 %) of experimentally studied host species

were captive, while 37,5 % of publications studied wild animals, and 7,8 % studied both.

Figure 2. *Summary of literature. The proportion of skin microbiome studies conducted on different animal classes on all continents. Number of studies in parentheses.*

i) Has the protective skin microbiome of hibernating animals been studied?

Only 18 publications (7,8 % of all articles) studied the protective microbiome during hibernation. Six of these studies were solely descriptive and 12 were experimental. Although many amphibian and reptilian species are capable of hibernation, no studies were conducted on the protective role of their skin microbiome during hibernation. In fact, all studies that sampled the skin microbiomes of hibernating animals involved bats and *Pd*.

Among the 12 experimental publications that focused specifically on a pathogen that infects hosts during hibernation, nine studies tested antifungal microbes against *Pd* using inhibition assays *in vitro* and three studies tested probiotic treatments *in vivo*. Only two of these studies were conducted in the Palearctic (Germany and China), where bats survive exposure to *Pd* without infection [49, 62]. Both studies used bacteria in inhibition assays to successfully suppress the growth of *Pd*. The remaining ten studies were conducted in North America.

Altogether, only seven studies have been published about the skin microbiome of Palearctic bats. Among these, five were descriptive [63–67] and two experimental [49, 62]. Four took place in China [49, 63, 64, 66], while the remaining three were conducted in Germany [62], Poland/Armenia [65], and Belgium [67]. Out of these, only five publications sampled hibernating bats [49, 63–66], while one study sampled active bats and the wall of the hibernacula [67], and one acquired the symbiont tested against *Pd* from the environment (not bat skin) [62].

ii) Has experimental skin microbiome research studying pathogen inhibition increased in recent years?

The overall amount of research on the protective role of the skin microbiome in non-human vertebrates has increased dramatically over the past 18 years, with the first study conducted in 2005 (Fig 3). Regardless of the growing interest in this field, the proportion of experimental studies investigating inhibitory ability of antimicrobial microbes residing on the skin has decreased significantly ($p < 0,001$, -0.20187 ± 0.04862) in

210 proportion to the number of studies published. This might be explained by the fact that the topic is vastly
211 unexplored, and most studies focus on solely describing the skin microbiome composition of animals and whether
212 known antimicrobial taxa are found on the skin. It is, however, noteworthy to mention that the number of all
213 studies published (both descriptive and experimental) has been lower in the year 2023 compared to previous
214 years. It remains to be determined whether more studies will be published by the end of the year.

215
216 ***Figure 3.*** *Temporal trends in protective skin microbiome studies in non-human animals. The year 2023 was excluded from*
217 *the GLM and scatter plot since the year is not over.*
218

iii) Which antifungal microbes have been identified and studied the most?

A total of 105 microbial genera were found to show weak to strong pathogen inhibition in the experimental
studies. The majority of tested microbes were bacteria (84,8 %), but fungi (13,3 %), and archaea (1,9 %) were
also tested successfully. Out of these, 51 genera were experimentally tested more than once, and 27 genera were
tested on two or more classes of animals (Fig. 4).

The most popular bacterial genus tested against pathogens was *Pseudomonas*. It was tested against multiple
pathogens (*Pd, Bd,* and others) that infect several classes of animals including mammals, amphibians, and ray-
finned fishes. *Pseudomonas* species showed strong to moderate inhibition of pathogens and they are considered
one of the primary candidates for *Pd*-inhibition in bats [46, 68]. Altogether, 31 studies testing *Pseudomonas*
showed successful pathogen inhibition, however, two studies using *Pseudomonas* as a probiotic were
unsuccessful [22, 69]. The authors of one failed trial noted that the skin microbiome still retained a defensive role
against *Bd*, but that the antifungal isolates were unable to colonize the skin of the amphibian host. This issue may
be addressed in future experiments by reapplying treatment or prolonging exposure to treatment [22].

Other microbes successfully tested against pathogens in multiple studies included the genera *Janthinobacterium, Bacillus, Chryseobacterium, Stenotrophomonas, Serratia, Acinetobacter, Rhodococcus,* and *Enterobacter*, indicating these bacteria show promise as probiotics and should be investigated in more detail. However, it is important to mention that *Janthinobacterium*, *Chryseobacterium*, *Stenotrophomonas,* and *Rhodococcus* also failed some trials. For example, while the genus *Rhodococcus* showed strong inhibition of *Pd in vitro* [70], a recent *in vivo* experiment on bat skin was not successful [71]. These results highlight the need for more trials to determine whether these microbes can, in fact, be used in wildlife disease mitigation. Further experiments will also help assess the microbial mechanisms of inhibition, which provide important information about the conditions that best facilitate pathogen inhibition.

Not all experimental publications studied the inhibitory effects of a single microbial strain, six publications (9,1 % of experimental studies) researched the collective effect of a group of inhibiting microbes. Bacterial genera used in the consortium studies were *Pseudomonas, Janthinobacterium, Bacillus, Chryseobacterium, Stenotrophomonas, Serratia, Acinetobacter, Enterobacter, Microbacterium, Staphylococcus, Citrobacter, Comamonas, Pedobacter, Chitinophaga, Iodobacter*, *Collimonas, Curvibacter,* and *Sanguibacter.* Out of these, five publications were successful in inhibiting pathogens [20, 72–75], and one was not [69].

Figure 4. *Antifungal microbes found on the skin of two or more classes of animals that successfully inhibited pathogens. Size of block indicates number of studies conducted (larger block = more studies).*

DISCUSSION

In this review, we illustrate that the protective role of the skin microbiome is becoming an increasingly popular topic of research. However, most publications focus on describing the composition of the skin microbiome and

identifying known antifungal microbes without testing their pathogen inhibition ability. This phenomenon may be
explained by the novelty of the topic, as the majority of publications aim to simply describe the microbial
diversity on the skin of hosts, before experimentally testing them against pathogens. Additionally, the knowledge
attained so far centers mainly around amphibians in the Nearctic, leaving other animal classes and continents
overwhelmingly unexplored. The popularity of protective skin microbiome studies in North American amphibians
can be explained by the disastrous emergence of chytridiomycosis in 1993 [24] and the uneven distribution of
research funding opportunities that are overly represented in the Nearctic [76].

The main research gap we want to highlight with this review is the lack of publications about the protective role
of the skin microbiome in hibernating animals, specifically studies that experimentally test the inhibition ability of
cutaneous microbiota against pathogens. The suppression of the host's immune system during hibernation
amplifies the importance of the skin microbiome since it may remain active when the host is not. It would be
especially beneficial to study the protective role of the skin microbiome in species of animals that survive
exposure to pathogens during hibernation, such as Palearctic bats. So far, the skin microbiomes of only 13 of over
100 bat species in the Palearctic have been studied. Most of these studies focused on fungal symbionts and just
one study was conducted on the protective mycobiome of *M. myotis,* the flagship species known to tolerate high
Pd loads in Europe. To our knowledge, there are no published data on the mutualistic bacteria living on the
epidermis of *M. myotis*. In fact, the bacterial composition of most Palearctic bat species and possible temporal
changes in their microbiome composition (for example during hibernation) remain unknown.

Are probiotics the solution to lethal skin disease in wildlife?

Describing and experimentally testing microbial species found on host skin are the first steps to developing a non-
toxic disease mitigation strategy for lethal skin infections in wild animals. As mentioned earlier, using
antimicrobial bacteria as a preventative probiotic treatment on the skin to help mitigate disease has already been
explored in some organisms [26, 50, 77]. Results from these studies have varied, however, most studies have
found encouraging findings in several classes of animals. For example, the bioaugmentation of a known
antifungal bacterium (*Janthinobacterium lividum)* on frog skin successfully prevented mortality due to
chytridiomycosis [17]. Moreover, probiotic treatments tested on walleye fish (*Sander vitreus*) were found to have
a significant antagonistic effect against a common pathogen (*Flavobacterium columnare*) and increase the
survival of fish exposed to the pathogen [78]. In addition, a probiotic bacterium isolated from feline skin
successfully reduced the colonization of a pathogen when added to the epidermis of mice, indicating certain
probiotics could be effective across multiple species [79]. While this provides compelling evidence for the
justification of probiotic use, other publications have reported contradicting results [22, 69, 80], suggesting more
information is needed before probiotic treatments can be successfully applied to wildlife disease mitigation.

Our results indicate that several microbial species, mostly bacteria, have been shown to exhibit potential as
probiotics. Notably, the bacterial genus *Pseudomonas* has demonstrated the inhibition ability of several pathogens
infecting multiple classes of animals, including bats and *Pd* [46, 81–83]. However, there are various other
microbial genera that have shown inhibition ability but are still overlooked. Fungi are among the often
disregarded species that have also shown promise in pathogen inhibition [84–86]. For example, North American
bat species resistant to WNS exhibit a more diverse cutaneous mycobiome compared to WNS-susceptible species
[86]. Some common fungal genera identified on bat skin, such as *Cutaneotrichosporon*, *Aureobasidium*, and
Holtermanniella, have also been found to inhibit the growth of *Pd in vitro*, albeit weakly [86, 87]. Additionally,
gram-positive bacteria may be overlooked in these studies since DNA extraction methods do not always
successfully permeate the thick outer layer of the bacteria [88, 89]. These bacteria may also possess the ability to
inhibit pathogens, but could be underrepresented in these datasets and, therefore, not tested for inhibition.

It is also important to acknowledge that certain mutualistic microbial genera, such as *Pseudomonas*, are known
pathogens for certain organisms [20, 90, 91], meaning the effect of the microbial genus is highly dependent on
context [21]. For example, *Pseudomonas fluorescens*, a commensal on bat skin [51], can be lethal to fruit flies
(*Drosophila melanogaster*) and ladybird beetles (*Henosepilachna vigintioctopunctata*) [92]. Timing of treatment
is also of importance since the addition of *P. fluorescens* to bat skin before exposure to *Pd* increased disease
severity, while simultaneous treatment and exposure reduced *Pd* invasion [50]. When utilizing probiotics, there is
always the risk that the symbionts could spread to and infect non-target species causing more harm than good.
Hence, it is advisable that the probiotic is indigenous to the local environment and has been studied adequately
before adding treatment to an ecosystem or species [21, 93].

In addition to testing the pathogen inhibiting ability of just one microbial strain, there seems to be an emerging
trend of testing a consortium of bacteria against pathogens. Multiple studies have found that more diverse
communities of bacteria can outperform single strains in inhibiting pathogen growth [73–75]. Bacterial growth
rate *in vitro* has also been found to be higher, when bacterial strains were grown together, instead of individually
[72]. This is understandable given that a diverse community of organisms is known to be more resistant to
invasions on both a macro- [94] and micro-scale [95]. For instance, as an analogous example, grassland plots with
higher species diversity are more resistant to colonization by invasive plants than homogenous plots [94]. The
interactions of microbial species within the microbiome mirror those of organisms in a macro-level ecosystem
(for example a forest), which is why diversity means better pathogen resistance in the skin microbiome as well [1,
86].

Future threats and conservation

316   As climate change progresses and humans encroach further into wildlife habitats, people and wildlife alike will be
317   more regularly exposed to new potential pathogens [96, 97]. Fungi, in particular, should be treated with concern

as fungal infections are notoriously difficult to treat due to their resilient nature. Over 600 species of fungi are known to infect vertebrates and many species have been identified as the causal agents of potential emerging infectious diseases (EIDs) in recent years [96, 98]. In fact, fungi are more closely related to animals than bacteria, and therefore, do not respond well to common antimicrobial treatments that work on bacterial infections [98]. WNS and chytridiomycosis have demonstrated how rapidly fungal disease outbreaks can devastate wildlife populations and highlight the need for preventative disease mitigation strategies.

It is often difficult to manage disease outbreaks in endangered wildlife populations, so captive breeding and reintroduction are occasionally used to attempt to restore declining populations [99]. These attempts are often costly and have varying success rates. In this review, the majority (54,7 %) of experimental skin microbiome studies were conducted on captive animals. However, since the skin microbiome is heavily influenced by the environment [1, 100], the results from these studies may not always be applicable to wild animals. For example, the skin microbiome of captive amphibians is known to be less diverse than that of their wild counterparts, which may become an issue when reintroducing captive animals back into the wild during conservation efforts [101]. Considering reduced diversity in the skin microbiome affects the host's ability to resist infection, the holobiont perspective could be beneficial when planning and upgrading conservation methods [99].

CONCLUSIONS

While the skin microbiome holds tremendous potential for disease mitigation, its protective role during hibernation is highly understudied. Not only is there a scarcity of publications describing the microbial diversity inhabiting the skin, but there is also a notable absence of experimental studies determining which microbes effectively inhibit pathogens. Hibernatory bats and WNS provide an exceptional study system for addressing this knowledge gap and we encourage researchers to tackle this subject by exploring the microbial species living on bat skin and their potential as probiotics in WNS mitigation. Specifically, the skin microbiome of Palearctic bats should be studied to determine how they survive exposure to *Pd*, as this information could be beneficial for solving the WNS crisis in North America. In particular, we recommend future research concentrate on testing the anti-pathogen activity of lesser-known symbionts, such as fungi, in addition to testing a consortium of known antifungal bacteria. We emphasize the importance of adopting a holistic approach which incorporates the holobiont perspective into conservation planning for more efficient results in disease mitigation.

LIST OF ABBREVIATIONS

Bd – Batrachochytrium dendrobatidis

WNS – White-nose syndrome

Pd - Pseudogymnoascus destructans

DECLARATIONS

Ethical Approval

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

TML, VNL and TST conceived the idea and designed the methodology. TST extracted data from the literature, analyzed the data, prepared the figures, and led the writing of the manuscript. All authors read, contributed to, and approved the final manuscript.

Acknowledgements

The authors would like to thank Dr. Melissa Meierhofer for her insightful and constructive input in the selection of figures and models used in this review.

Funding

This work was supported by funding from the Maj & Tor Nessling Foundation and the Academy of Finland.

Availability of data and materials

All data presented in the supplementary material.

Supplementary file 1: Data set (Review_Data_Troitsky.xlsx)

Contains list of chosen publications and data extracted from them that was used in model and figures.

REFERENCES

1. Grice EA, Segre JA. The skin microbiome. Nat Rev Microbiol. 2011;9:244–53.

2. Byrd AL, Belkaid Y, Segre JA. The human skin microbiome. Nat Rev Microbiol. 2018;16:143–55.

3. Belkaid Y, Segre JA. Dialogue between skin microbiota and immunity. Science. 2014;346:954–9.

4. Findley K, Grice EA. The skin microbiome: a focus on pathogens and their association with skin disease. PLoS Pathog. 2014;10:e1004436.

5. Smythe P, Wilkinson HN. The skin microbiome: current landscape and future opportunities. Int J Mol Sci. 2023;24:3950.

6. Ross AA, Rodrigues Hoffmann A, Neufeld JD. The skin microbiome of vertebrates. Microbiome. 2019;7:79.

7. Cullen CM, Aneja KK, Beyhan S, Cho CE, Woloszynek S, Convertino M, et al. Emerging priorities for microbiome research. Front Microbiol. 2020;11.

8. Zilber-Rosenberg I, Rosenberg E. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. FEMS Microbiol Rev. 2008;32:723–35.

9. Bordenstein SR, Theis KR. Host biology in light of the microbiome: ten principles of holobionts and hologenomes. PLOS Biol. 2015;13:e1002226.

10. Gilbert SF, Sapp J, Tauber AI. A symbiotic view of life: we have never been individuals. Q Rev
Biol. 2012;87:325–41.

11. Sender R, Fuchs S, Milo R. Revised estimates for the number of human and bacteria cells in the
body. PLOS Biol. 2016;14:e1002533.

12. Rosenberg E, Zilber-Rosenberg I. The hologenome concept of evolution after 10 years. Microbiome.
2018;6:78.

13. Nørreslet LB, Agner T, Clausen M-L. The skin microbiome in inflammatory skin diseases. Curr
Dermatol Rep. 2020;9:141–51.

14. Clifton R, Monaghan EM, Green MJ, Purdy KJ, Green LE. Differences in composition of
interdigital skin microbiota predict sheep and feet that develop footrot. Sci Rep. 2022;12:8931.

15. Becker MH, Harris RN. Cutaneous bacteria of the redback salamander prevent morbidity associated
with a lethal disease. PLOS ONE. 2010;5:e10957.

16. Greenspan SE, Peloso P, Fuentes-González JA, Bletz M, Lyra ML, Machado IF, et al. Low
microbiome diversity in threatened amphibians from two biodiversity hotspots. Anim Microbiome.
2022;4:69.

17. Harris RN, Brucker RM, Walke JB, Becker MH, Schwantes CR, Flaherty DC, et al. Skin microbes
on frogs prevent morbidity and mortality caused by a lethal skin fungus. ISME J. 2009;3:818–24.

18. Muñoz-Atienza E, Araújo C, Magadán S, Hernández PE, Herranz C, Santos Y, et al. In vitro and
in vivo evaluation of lactic acid bacteria of aquatic origin as probiotics for turbot (Scophthalmus
maximus L.) farming. Fish Shellfish Immunol. 2014;41:570–80.

19. Lemieux-Labonté V, Simard A, Willis CKR, Lapointe F-J. Enrichment of beneficial bacteria in the
skin microbiota of bats persisting with white-nose syndrome. Microbiome. 2017;5:115.

20. Xiong Y, Wu Q, Qin X, Yang C, Luo S, He J, et al. Identification of Pseudomonas aeruginosa from
the skin ulcer disease of crocodile lizards (Shinisaurus crocodilurus) and probiotics as the control
measure. Front Vet Sci. 2022;9.

21. Woodhams DC, Brandt H, Baumgartner S, Kielgast J, Küpfer E, Tobler U, et al. Interacting
symbionts and immunity in the amphibian skin mucosome predict disease risk and probiotic
effectiveness. PLOS ONE. 2014;9:e96375.

22. Becker MH, Walke JB, Cikanek S, Savage AE, Mattheus N, Santiago CN, et al. Composition of
symbiotic bacteria predicts survival in Panamanian golden frogs infected with a lethal fungus. Proc R
Soc B Biol Sci. 2015;282:20142881.

23. Madison JD, Berg EA, Abarca JG, Whitfield SM, Gorbatenko O, Pinto A, et al. Characterization of
Batrachochytrium dendrobatidis inhibiting bacteria from amphibian populations in Costa Rica. Front
Microbiol. 2017;8.

24. Berger L, Speare R, Daszak P, Green DE, Cunningham AA, Goggin CL, et al. Chytridiomycosis
causes amphibian mortality associated with population declines in the rain forests of Australia and
Central America. Proc Natl Acad Sci. 1998;95:9031–6.

25. Voyles J, Young S, Berger L, Campbell C, Voyles WF, Dinudom A, et al. Pathogenesis of
chytridiomycosis, a cause of catastrophic amphibian declines. Science. 2009;326:582–5.

26. Becker MH, Brucker RM, Schwantes CR, Harris RN, Minbiole KPC. The bacterially produced
metabolite violacein is associated with survival of amphibians infected with a lethal fungus. Appl
Environ Microbiol. 2009;75:6635–8.

27. Woodhams DC, LaBumbard BC, Barnhart KL, Becker MH, Bletz MC, Escobar LA, et al.
Prodigiosin, violacein, and volatile organic compounds produced by widespread cutaneous bacteria of
amphibians can inhibit two Batrachochytrium fungal pathogens. Microb Ecol. 2018;75:1049–62.

28. Garber G. An overview of fungal infections. Drugs. 2001;61:1–12.

29. Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, et al. Emerging fungal
threats to animal, plant and ecosystem health. Nature. 2012;484:186–94.

30. Blehert DS, Hicks AC, Behr M, Meteyer CU, Berlowski-Zier BM, Buckles EL, et al. Bat white-nose
syndrome: an emerging fungal pathogen? Science. 2009;323:227–227.

31. Frick WF, Pollock JF, Hicks AC, Langwig KE, Reynolds DS, Turner GG, et al. An emerging
disease causes regional population collapse of a common North American bat species. Science.
2010;329:679–82.

32. Warnecke L, Turner JM, Bollinger TK, Lorch JM, Misra V, Cryan PM, et al. Inoculation of bats
with European Geomyces destructans supports the novel pathogen hypothesis for the origin of white-
nose syndrome. Proc Natl Acad Sci U S A. 2012;109:6999–7003.

33. Frick WF, Puechmaille SJ, Willis CKR. White-nose syndrome in bats. In: Voigt CC, Kingston T,
editors. Bats in the Anthropocene: conservation of bats in a changing world. Cham: Springer
International Publishing; 2016. p. 245–62.

34. Meteyer CU, Buckles EL, Blehert DS, Hicks AC, Green DE, Shearn-Bochsler V, et al.
Histopathologic criteria to confirm white-nose syndrome in bats. J Vet Diagn Investig Off Publ Am
Assoc Vet Lab Diagn Inc. 2009;21:411–4.

35. Altringham JD. Bats: from evolution to conservation. Oxford University Press; 2011.

36. Geiser F. Metabolic rate and body temperature reduction during hibernation and daily torpor. Annu
Rev Physiol. 2004;66:239–74.

37. Bouma HR, Carey HV, Kroese FGM. Hibernation: the immune system at rest? J Leukoc Biol.
2010;88:619–24.

38. Leopardi S, Blake D, Puechmaille SJ. White-nose syndrome fungus introduced from Europe to
North America. Curr Biol. 2015;25:R217–9.

39. Zukal J, Bandouchova H, Brichta J, Cmokova A, Jaron KS, Kolarik M, et al. White-nose syndrome
without borders: Pseudogymnoascus destructans infection tolerated in Europe and Palearctic Asia but
not in North America. Sci Rep. 2016;6:19829.

40. Fritze M, Puechmaille SJ. Identifying unusual mortality events in bats: a baseline for bat hibernation
monitoring and white-nose syndrome research. Mammal Rev. 2018;48:224–8.

41. Iliev ID, Underhill DM. Striking a balance: fungal commensalism versus pathogenesis. Curr Opin
Microbiol. 2013;16:366–73.

42. Harazim M, Horáček I, Jakešová L, Luermann K, Moravec JC, Morgan S, et al. Natural selection in
bats with historical exposure to white-nose syndrome. BMC Zool. 2018;3:8.

43. Wibbelt G, Kurth A, Hellmann D, Weishaar M, Barlow A, Veith M, et al. White-nose syndrome
fungus (Geomyces destructans) in bats, Europe. Emerg Infect Dis. 2010;16:1237–43.

44. Drees KP, Lorch JM, Puechmaille SJ, Parise KL, Wibbelt G, Hoyt JR, et al. Phylogenetics of a
fungal invasion: origins and widespread dispersal of white-nose syndrome. mBio.
2017;8:10.1128/mbio.01941-17.

45. Hoyt JR, Kilpatrick AM, Langwig KE. Ecology and impacts of white-nose syndrome on bats. Nat
Rev Microbiol. 2021;19:196–210.

46. Hoyt JR, Cheng TL, Langwig KE, Hee MM, Frick WF, Kilpatrick AM. Bacteria isolated from bats
inhibit the growth of Pseudogymnoascus destructans, the causative agent of white-nose syndrome.
PLOS ONE. 2015;10:e0121329.

47. Forsythe A, Fontaine N, Bissonnette J, Hayashi B, Insuk C, Ghosh S, et al. Microbial isolates with
Anti-Pseudogymnoascus destructans activities from Western Canadian bat wings. Sci Rep.
2022;12:9895.

48. Grisnik M, Bowers O, Moore AJ, Jones BF, Campbell JR, Walker DM. The cutaneous microbiota of
bats has in vitro antifungal activity against the white nose pathogen. FEMS Microbiol Ecol.
2020;96:fiz193.

49. Li Z, Li A, Hoyt JR, Dai W, Leng H, Li Y, et al. Activity of bacteria isolated from bats against
Pseudogymnoascus destructans in China. Microb Biotechnol. 2021;15:469–81.

50. Cheng TL, Mayberry H, McGuire LP, Hoyt JR, Langwig KE, Nguyen H, et al. Efficacy of a
probiotic bacterium to treat bats affected by the disease white-nose syndrome. J Appl Ecol.
2017;54:701–8.

51. Hoyt JR, Langwig KE, White JP, Kaarakka HM, Redell JA, Parise KL, et al. Field trial of a
probiotic bacteria to protect bats from white-nose syndrome. Sci Rep. 2019;9:9158.

52. Gillingham JC, Carpenter CC. Snake hibernation: construction of and observations on a man-made
hibernaculum (Reptilia, Serpentes). J Herpetol. 1978;12:495–8.

53. Buck CL, Barnes BM. Annual cycle of body composition and hibernation in free-living arctic
ground squirrels. J Mammal. 1999;80:430–42.

54. McKechnie AE, Ashdown RAM, Christian MB, Brigham RM. Torpor in an African caprimulgid, the
freckled nightjar Caprimulgus tristigma. J Avian Biol. 2007;38:261–6.

55. Campbell HA, Fraser KPP, Bishop CM, Peck LS, Egginton S. Hibernation in an Antarctic fish: on
ice for winter. PLoS ONE. 2008;3:e1743.

56. Costanzo JP, do Amaral MCF, Rosendale AJ, Lee RE Jr. Hibernation physiology, freezing
adaptation and extreme freeze tolerance in a northern population of the wood frog. J Exp Biol.
2013;216:3461–73.

57. Evans AL, Singh NJ, Friebe A, Arnemo JM, Laske TG, Fröbert O, et al. Drivers of hibernation in
the brown bear. Front Zool. 2016;13:1–14.

58. R Core Team. R: A language and environment for statistical computing. R Found Stat Comput
Vienna Austria. 2022.

59. Wickham H. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York; 2016.

60. Dormann C, Gruber B, Fründ J. Introducing the bipartite package: analysing ecological networks. R
News. 2008;8:8–11.

61. Harrington B. Inkscape. 2004.

62. Fritze M, Pham TLH. Effekt des bodenbakteriums Pseudomonas veronii-like PAZ1 auf das
wachstum des white-nose erregers Geomyces destructans in antagonisten-tests. Nyctalus. 2012;17:104–
7.

63. Li Z, Li A, Dai W, Leng H, Liu S, Jin L, et al. Skin microbiota variation among bat species in china
and their potential defense against pathogens. Front Microbiol. 2022;13.

64. Li A, Li Z, Leng H, Jin L, Xiao Y, Sun K, et al. Seasonal assembly of skin microbiota driven by
neutral and selective processes in the greater horseshoe bat. Mol Ecol. 2023;n/a n/a.

65. Seidlova V, Pikula J, Kolarik M, Nováková A, Cmokova A, Ghazaryan A, et al. Higher white-nose
syndrome fungal isolate yields from UV-guided wing biopsies compared with skin swabs and optimal
culture media. BMC Vet Res. 2023;19:40.

66. Li A, Li Z, Dai W, Parise KL, Leng H, Jin L, et al. Bacterial community dynamics on bats and the
implications for pathogen resistance. Environ Microbiol. 2022;24:1484–98.

67. Becker P, van den Eynde C, Baert F, D'hooge E, De Pauw R, Normand A-C, et al. Remarkable
fungal biodiversity on northern Belgium bats and hibernacula. Mycologia. 2023;115:484–98.

68. Cheng TL, Mayberry H, McGuire LP, Hoyt JR, Langwig KE, Nguyen H, et al. Efficacy of a probiotic bacterium to treat bats affected by the disease white☐nose syndrome. J Appl Ecol. 2017;54:701–8.

69. Woodhams DC, Rollins-Smith LA, Reinert LK, Lam BA, Harris RN, Briggs CJ, et al. Probiotics modulate a novel amphibian skin defense peptide that is antifungal and facilitates growth of antifungal bacteria. Microb Ecol. 2020;79:192–202.

70. Cornelison CT, Keel MK, Gabriel KT, Barlament CK, Tucker TA, Pierce GE, et al. A preliminary report on the contact-independent antagonism of Pseudogymnoascus destructans by Rhodococcus rhodochrousstrain DAP96253. BMC Microbiol. 2014;14:246.

71. Hooper S, Amelon S. Contact-independent exposure to Rhodococcus rhodochrous DAP96253 volatiles does not improve the survival rate of Myotis lucifugus (little brown bats) affected by White-nose Syndrome. PeerJ. 2023;11:e15782.

72. Alexiev A, Chen MY, Korpita T, Weier AM, McKenzie VJ. Together or alone: evaluating the pathogen inhibition potential of bacterial cocktails against an amphibian pathogen. Microbiol Spectr. 2023;11:e01518-22.

73. Antwis RE, Harrison XA. Probiotic consortia are not uniformly effective against different amphibian chytrid pathogen isolates. Mol Ecol. 2018;27:577–89.

74. Loudon AH, Holland JA, Umile TP, Burzynski EA, Minbiole KPC, Harris RN. Interactions between amphibians' symbiotic bacteria cause the production of emergent anti-fungal metabolites. Front Microbiol. 2014;5.

75. Piovia-Scott J, Rejmanek D, Woodhams DC, Worth SJ, Kenny H, McKenzie V, et al. Greater species richness of bacterial skin symbionts better suppresses the amphibian fungal pathogen Batrachochytrium dendrobatidis. Microb Ecol. 2017;74:217–26.

76. Petersen OH. Inequality of research funding between different countries and regions is a serious problem for global science. Function. 2021;2:zqab060.

77. Boutin S, Audet C, Derome N. Probiotic treatment by indigenous bacteria decreases mortality without disturbing the natural microbiota of Salvelinus fontinalis. Can J Microbiol. 2013;59:662–70.

78. Seghouani H, Garcia-Rangel C-E, Füller J, Gauthier J, Derome N. Walleye autochthonous bacteria as promising probiotic candidates against Flavobacterium columnare. Front Microbiol. 2017;8.

79. O'Neill AM, Worthing KA, Kulkarni N, Li F, Nakatsuji T, McGrosso D, et al. Antimicrobials from a feline commensal bacterium inhibit skin infection by drug-resistant S. pseudintermedius. eLife. 2021;10:e66793.

80. Knapp RA, Joseph MB, Smith TC, Hegeman EE, Vredenburg VT, Jr JEE, et al. Effectiveness of antifungal treatments during chytridiomycosis epizootics in populations of an endangered frog. PeerJ. 2022;10:e12712.

81. Lauer A, Simon MA, Banning JL, Lam BA, Harris RN. Diversity of cutaneous bacteria with
antifungal activity isolated from female four-toed salamanders. ISME J. 2008;2:145–57.

82. Harris RN, Lauer A, Simon MA, Banning JL, Alford RA. Addition of antifungal skin bacteria to
salamanders ameliorates the effects of chytridiomycosis. Dis Aquat Organ. 2009;83:11–6.

83. Rosado D, Xavier R, Severino R, Tavares F, Cable J, Pérez-Losada M. Effects of disease, antibiotic
treatment and recovery trajectory on the microbiome of farmed seabass (Dicentrarchus labrax). Sci Rep.
2019;9:18946.

84. Kearns PJ, Fischer S, Fernández-Beaskoetxea S, Gabor CR, Bosch J, Bowen JL, et al. Fight fungi
with fungi: antifungal properties of the amphibian mycobiome. Front Microbiol. 2017;8.

85. Kearns PJ, Winter AS, Woodhams DC, Northup DE. The mycobiome of bats in the American
Southwest is structured by geography, bat species, and behavior. Microb Ecol. 2023.
https://doi.org/10.1007/s00248-023-02230-w.

86. Vanderwolf KJ, Campbell LJ, Goldberg TL, Blehert DS, Lorch JM. Skin fungal assemblages of bats
vary based on susceptibility to white-nose syndrome. ISME J. 2021;15:909–20.

87. Vanderwolf KJ, Campbell LJ, Taylor DR, Goldberg TL, Blehert DS, Lorch JM. Mycobiome traits
associated with disease tolerance predict many western North American bat species will be susceptible
to white-nose syndrome. Microbiol Spectr. 2021;9:e0025421.

88. Auer GK, Weibel D. Bacterial cell mechanics. Biochemistry. 2017;56:3710–24.

89. Li X, Bosch-Tijhof CJ, Wei X, de Soet JJ, Crielaard W, Loveren C van, et al. Efficiency of chemical
versus mechanical disruption methods of DNA extraction for the identification of oral gram-positive and
gram-negative bacteria. J Int Med Res. 2020;48:0300060520925594.

90. Eissa NME, El-Ghiet ENA, Shaheen AA, Abbass A. Characterization of Pseudomonas species
isolated from tilapia "Oreochromis niloticus" in Qaroun and Wadi-El-Rayan Lakes, Egypt. Glob Vet.
2010;5:116–21.

91. Wood SJ, Kuzel TM, Shafikhani SH. Pseudomonas aeruginosa: infections, animal modeling, and
therapeutics. Cells. 2023;12:199.

92. Teoh M-C, Furusawa G, Veera Singham G. Multifaceted interactions between the pseudomonads
and insects: mechanisms and prospects. Arch Microbiol. 2021;203:1891–915.

93. Myers JM, Ramsey JP, Blackman AL, Nichols AE, Minbiole KPC, Harris RN. Synergistic inhibition
of the lethal fungal pathogen Batrachochytrium dendrobatidis: the combined effect of symbiotic
bacterial metabolites and antimicrobial peptides of the frog Rana muscosa. J Chem Ecol. 2012;38:958–
65.

94. Kennedy TA, Naeem S, Howe KM, Knops JMH, Tilman D, Reich P. Biodiversity as a barrier to
ecological invasion. Nature. 2002;417:636–8.

95. Mallon CA, Elsas JD van, Salles JF. Microbial invasions: the process, patterns, and mechanisms.
Trends Microbiol. 2015;23:719–29.

96. Daszak P, Cunningham AA, Hyatt AD. Emerging infectious diseases of wildlife--threats to
biodiversity and human health. Science. 2000;287:443–9.

97. Garcia-Solache MA, Casadevall A. Global warming will bring new fungal diseases for mammals.
mBio. 2010;1:10.1128/mbio.00061-10.

98. Fisher MC, Gurr SJ, Cuomo CA, Blehert DS, Jin H, Stukenbrock EH, et al. Threats posed by the
fungal kingdom to humans, wildlife, and agriculture. mBio. 2020;11:e00449-20.

99. Carthey AJR, Blumstein DT, Gallagher RV, Tetu SG, Gillings MR. Conserving the holobiont. Funct
Ecol. 2020;34:764–76.

100. Colston TJ, Jackson CR. Microbiome evolution along divergent branches of the vertebrate tree of
life: what is known and unknown. Mol Ecol. 2016;25:3776–800.

101. Kueneman JG, Bletz MC, Becker M, Gratwicke B, Garcés OA, Hertz A, et al. Effects of captivity
and rewilding on amphibian skin microbiomes. Biol Conserv. 2022;271:109576.