Effects of transient, persistent, and resurgent sodium currents on excitability and spike regularity in vestibular ganglion neurons

Selina Baeza-Loya[1], Ruth Anne Eatock[2*]

[1]Virginia Merrill Bloedel Hearing Research Center, Department of Otolaryngology-HNS, University
of Washington, Seattle, WA, United States

[2]Department of Neurobiology, University of Chicago, Chicago, IL, United States

* Correspondence:
Ruth Anne Eatock
eatock@uchicago.edu

Keywords: Vestibular ganglion neurons, spiking, regularity, excitability, Na$_V$ currents,
persistent current, resurgent current

Abstract

Vestibular afferent neurons occur as two populations with differences in spike timing regularity that
are independent of rate. The more excitable regular afferents have lower current thresholds and
sustained spiking responses to injected currents, while irregular afferent neurons have higher
thresholds and transient responses. Differences in expression of low-voltage-activated potassium
(K$_{LV}$) channels are emphasized in models of spiking regularity and excitability in these neurons,
leaving open the potential contributions of the voltage-gated sodium (Na$_V$) channels responsible for
the spike upstroke. We investigated the impact of different Na$_V$ current modes (transient, persistent,
and resurgent) with whole-cell patch clamp experiments in mouse vestibular ganglion neurons
(VGNs), the cultured and dissociated cell bodies of afferents. All VGNs had transient Na$_V$ current,
many had a small persistent (non-inactivating) Na$_V$ current, and a few had resurgent current, which
flows after the spike peak when Na$_V$ channels that were blocked are unblocked. Na$_V$1.6 channels
conducted most or all of each Na$_V$ current mode, and a Na$_V$1.6-selective blocker decreased spike rate
and altered spike waveforms in both sustained and transient VGNs. A Na$_V$ channel agonist enhanced
persistent current and increased spike rate and regularity. We hypothesized that persistent and
resurgent currents have different effects on sustained (regular) VGNs vs. transient (irregular) VGNs.
Lacking blockers specific for the different current modes, we used modeling to isolate their effects on
spiking of simulated transient and sustained VGNs, driven by simulated current steps and noisy trains
of simulated EPSCs. In all simulated neurons, increasing transient Na$_V$ current increased spike rate
and rate-independent regularity. In simulated sustained VGNs, adding persistent current increased
both rate and rate-independent regularity, while adding resurgent current had limited impact. In
transient VGNs, adding persistent current had little impact, while adding resurgent current increased
both rate and rate-independent irregularity by enhancing sensitivity to synaptic noise. These
experiments show that the small Na$_V$ current modes may enhance the differentiation of afferent
populations, with persistent currents selectively making regular afferents more regular and resurgent
currents selectively making irregular afferents less regular.

1   Introduction

Vestibular hair cells transmit information about head motions and tilt to the peripheral terminals of
bipolar vestibular ganglion neurons (VGNs, Fig. 1A). VGNs in amniotes are well-known for their
differences in regularity of spike timing: regular and irregular VGNs synapse on hair cells in
peripheral and central zones of vestibular inner ear epithelia and have tonic and phasic response
dynamics, respectively (reviewed in Goldberg, 2000; Eatock and Songer, 2011). The difference in
regularity is rate-independent, such that it holds even when afferents have similar average rates.
Evidence from *in vivo* neuronal responses to head motions links regularity to encoding strategy: more
"rate-based" in regular afferents and more "temporal" in irregular afferents, with each strategy
favoring different kinds of sensory information (reviewed in Cullen, 2019). The cellular mechanisms
that give rise to the difference in regularity are therefore significant for vestibular information
processing. Here we examine whether the existence of different modes – transient, persistent, and
resurgent – of voltage-gated sodium current (I-Na$_V$) contributes to differences in afferent excitability
(tendency to spike) and rate-independent regularity.

Depolarizing current steps evoke a spectrum of firing patterns in isolated VGN cell bodies, ranging
from sustained spiking to transient. Several lines of evidence indicate that sustained and transient
firing patterns correspond, respectively, to regular and irregular spike timing *in vivo* (Iwasaki et al.,
2008; Kalluri et al., 2010; Ono et al., 2020)*. In vivo*, spikes initiate in the peripheral neurite at the
heminode near its hair cell synapse and travel through the cell body into the central neurite (Fig. 1A).
Over the first postnatal week, the afferent nerve fibers, including the cell body, become myelinated.
We investigated the Na$_V$ currents that give rise to firing patterns in dissociated VGN cell bodies.
Their compact form allows the high-quality voltage clamp necessary for recording very fast Na$_V$
currents. To gain access for the patch electrode to the VGN membrane beyond the first postnatal
week, we cultured dissociated cell bodies overnight, which loosens or eliminates the myelin
covering.

With this approach, previous work on rat VGNs in the first two postnatal weeks established that,
relative to sustained VGNs, transient VGNs are less excitable, with higher current thresholds for
spiking, reflecting their more negative resting potentials and lower input resistances (Iwasaki et al.,
2008; Kalluri et al., 2010). A major factor in these differences is the greater expression of low-
voltage-activated potassium (K$_{LV}$) channels from the K$_V$1 and K$_V$7 channel families in transient
VGNs. By stimulating VGNs with trains of simulated excitatory postsynaptic currents (EPSCs) with
pseudo-random timing, Kalluri et al. (2010) showed that transient VGNs have larger K$_{LV}$ currents
that drive irregular spike timing. Whether diversity in Na$_V$ currents contributes to afferent differences
in regularity and excitability has not been established.

Na$_V$ currents drive the rising phase of most neuronal action potentials (reviewed in Bean, 2007). In
rat vestibular ganglia (Liu et al. 2016), we showed diverse expression of Na$_V$ channel subunits and
characterized multiple transient Na$_V$ currents (I-Na$_V$T) in immature VGNs. Here we investigated a
different aspect of Na$_V$ current expression in VGNs that becomes a factor as the inner ear matures:
the expression of persistent (Na$_V$P) or resurgent (Na$_V$R) Na$_V$ modes of current, which flow through
the same channel subunits that produce larger Na$_V$T currents (Brown et al., 1994; Raman and Bean,
1997; Meredith and Rennie, 2020).

Most Na$_V$ channels strongly inactivate within milliseconds after depolarization, producing I-Na$_V$T. In
some cases, however, a small fraction do not inactivate even after seconds, creating I-Na$_V$P. I-Na$_V$P
activates more negatively than I-Na$_V$T and therefore enhances Na$_V$ channel availability near resting

potential, contributing to excitability and repetitive firing (Crill, 1996; Raman & Bean, 1997; Do and Bean, 2003). In another type of Na_V current, Na_V channels opened by a strong depolarization (like a large spike) are blocked by an intracellular molecule before they inactivate; upon repolarization, the channels rapidly unblock and carry I-Na_VR until they deactivate (Raman & Bean, 1997; Aman & Raman, 2024). These channels enhance depolarizing current on the downstroke of a spike and during the interspike interval.

Na_VP and Na_VR current modes have been recorded from gerbil calyces, the large terminals formed by vestibular afferents on type I hair cells and are attributed to tetrodotoxin (TTX)-sensitive Na_V1.6 channels (Meredith and Rennie, 2020). We speculated that diversity in Na_V current modes contributes to differences in excitability and/or regularity in VGNs, as suggested for certain brain neurons (Lewis and Raman, 2014). Our experiments and simulations together suggest that increasing I-Na_VT drives increases in spike rate and regularity in all VGNs, but that persistent and resurgent currents may enhance differentiation of spike regularity between sustained (regular) and transient (irregular) vestibular afferent neurons.

2   Methods

2.1   Electrophysiology

Animals: For all electrophysiology experiments, wildtype CD-1 mice were obtained from Charles River Laboratories (Wilmington, MA). Mice were housed at the University of Chicago Animal Care Facility and were handled in accordance with the National Research Council Guide for the Care and Use of Laboratory Animals. All procedures were approved by the animal care committee at the University of Chicago.

Preparation: Whole-cell currents and voltages were recorded from VGNs isolated from mice on postnatal day 3 (P3) to P25 (11 ± 0.5 (SEM) days old, median = 8, n = 146). VGNs from mice of both sexes were pooled into cell cultures; in the few cases where cells were from cultures made of a single sex (n = 7 female, 8 male cells), there were no significant differences in Na_V current voltage dependence of activation (p = 0.99, power = 0.05) or conductance density (p = 0.18, power = 0.27) between sexes.

Mice were anesthetized via isoflurane inhalation and decapitated. Temporal bones were dissected in chilled Liebovitz-15 (L-15) medium supplemented with 10 mM HEPES to pH 7.4, ~320 mmol/kg. Chemicals were purchased from ThermoFisher (Waltham, MA) unless otherwise stated.

Each otic capsule was exposed, and the superior compartment of the vestibular ganglion was dissected out. The superior ganglion houses the cell bodies of VGNs that synapse on hair cells in the utricle, part of the saccule, and the lateral and anterior semicircular canals, and project centrally to the vestibular nuclei in brain stem and cerebellum. We followed a dissociation protocol as described in Limón et al., 2005, with small changes. The tissue was placed in L-15 supplemented with 0.12% collagenase and 0.12% trypsin for 15-20 minutes at 37°C. The ganglion was then dissociated by gentle trituration into Minimal Essential Medium with Glutamax supplemented with 10 mM HEPES, 5% fetal bovine serum, and 1% penicillin (Sigma-Aldrich, St. Louis, MO). Cells were allowed to settle in glass-bottomed culture dishes (MatTek, Ashland, MA) pre-coated with poly-D-lysine. In most experiments, recordings were made after cells were incubated 10-20 hours in 5% CO_2 - 95% air at 37°C. Overnight incubation reduced myelin and satellite cell coverage from cell bodies. Age of cells does not include time in culture: e.g., data from a P17 VGN indicates the pup was 17 days old

when neurons were harvested, and the cells were P17 + ~1 day *in vitro*. The number of surviving
cells decreased with age of animal.

Recording solutions: During experiments, cells were bathed in either one of two external solutions,
summarized in Table 1. For voltage clamp experiments, we used a "Na^+-reduced" external solution
and a "Cs^+" -based internal solution, designed to minimize K^+ currents, Ca^{2+} currents, and reduce Na^+
currents. K^+ currents were minimized and Na^+ currents reduced by substituting Cs^+ for K^+ (in both
external and internal solutions, Table 1) and replacing about half of the external Na+ with equivalent
tetraethylammonium chloride (TEACl, a K channel blocker). By reducing K^+ and Na_V current, these
solutions improved voltage clamp quality by reducing voltage errors and the impact of the clamp
time constant. To minimize Ca^{2+} current, only trace Ca^{2+} was present and Mg^{2+} was added to block
voltage-gated calcium (Ca_V) channels. For current clamp experiments to gather spiking data, we used
more physiological external and internal solutions ("standard" solutions, Table 1).

Whole cell recordings: Cells were visualized at 600X on an inverted microscope equipped with
Nomarski optics (IMT-2; Olympus, Lake Success, NY). We chose round cells with diameters >5 µm
(range 8 – 25 µm). Mean membrane capacitance measured by the amplifier was 15 ± 0.5 (median =
16, n = 135). No correlation between age and cell size was observed. Neurons were distinguished
from glia by their shape and the presence of Na_V currents and/or spikes.

Stimuli were delivered, and responses recorded and amplified, with a Multiclamp 700B amplifier,
Digidata 1440A digitizer, and pClamp 11 software (Axon Instruments, Molecular Devices,
Sunnyvale, CA), with low-pass Bessel filtering set at 10 kHz and sampling interval set at 5 µs.
Stimuli include voltage steps and ramps, current steps, and synthetically generated frozen trains of
excitatory postsynaptic currents (EPSCs) based on recordings from vestibular terminals, as in Kalluri
et al., 2010. Electrodes were pulled from soda glass (King Precision Glass, Claremont, CA) to
resistances of 3 – 4 MΩ in our solutions and wrapped with parafilm to reduce pipette capacitance. All
membrane voltages were corrected offline for a liquid junction potential of either 4.7 (standard
solution) or 5.2 mV (reduced solution), calculated with JPCalc software (Barry 1994) in pClamp 11.

The series resistance (R_S) ranged between 3 and 10 MΩ and was compensated by $73 \pm 0.9\%$ (n =
134). Recordings were analyzed only when the seal exceeded 1 GΩ and, if standard solutions were
used, the cell maintained a stable resting membrane potential more negative than −60 mV. After
compensation for junction potentials, holding potential in voltage-clamp mode was −65 mV (−64.7
mV in standard solutions, −65.2 mV in Na^+ Reduced/Cs^+ solutions). Input resistance measured at
holding potential was 712 ± 56 MΩ. Fast Na_V currents are difficult to voltage clamp at body
temperature, so we recorded at room temperature (23 – 25°C). For our average cell capacitance (15
pF) and after R_S compensation, the voltage clamp time constant ranged from ~15-50 µs, adequate for
recording fast Na_V currents at room temperature, at which time-to-peak and inactivation time
constants are ~1 ms, and activation and deactivation time constants are shorter (Patel et al. 2015;
Alexander et al., 2021).

Pharmacology: On the day of experiments, we thawed stock solutions of pharmacological agents:
Na_V channel blockers tetrodotoxin (TTX; Enzo Life Sciences, Farmingdale, NY; 2 mM in distilled
water) and 4,9-anhydro-tetrodotoxin (4,9-ah-TTX; Alomone Labs, Jerusalem, Israel; 500 µM in
methanol), or Na_V channel agonist *Anemonia viridis* toxin 2 (ATX-II; Alomone Labs, Jerusalem,
Israel; 100 µM in distilled water). Aliquots were added to 5 mL of external solution for final
concentrations of 1 µM TTX, 100 nM 4,9-ah-TTX, and 100 nM ATX-II, chosen to maximize
blocking or agonizing effects (Oliveira et al., 2004; Rosker et al., 2007). 1 mg/ml of bovine serum

albumin was added to the ATX-II solution to reduce adhesion to the plastic delivery tubing. Toxin-
containing solutions were applied via local perfusion (Perfusion Fast-Step, Warner Instruments,
Holliston, MA) delivered with a Bee Hive Controller (BASI, West Lafayette, IN). Control and drug
solutions flowed through adjacent delivery tubes and a stepper mechanism selected the tube directed
at the patched cell. This system allows for no dead volume. Perfusion of control solution at the
beginning of each drug series provided additional control for flow effects.

Analysis and statistical treatments Analysis was performed in Clampfit (pClamp 10 or 11,
Molecular Devices) and MATLAB 2023b (The MathWorks, Natick, MA). Statistical analyses, curve
fitting, and figures were done in Origin Pro 2018 or 2024 (OriginLab, Northampton, MA). Means ±
SEM are presented. In box plots, box indicates SEM, middle line indicates median, open squares
indicate the mean, and whiskers mark 5-95% confidence intervals. For comparisons between 2
groups, we tested for normal distributions and homogeneity of variance (Levene's test), and
estimated statistical significance with Student's t-test or, if variances were unequal, Welch's t-test.
We used paired t-tests for drug effects on individual cells and an alpha level of 0.05 for all statistical
tests. For significant results, we calculated effect size with bias-corrected Hedges' g (small effect =
0.2, medium = 0.5, large = 0.8; Durlak, 2009). To compare >2 groups, we used one-way ANOVA
followed by the Bonferroni test for multiple comparisons.

The voltage dependencies of current activation and inactivation were analyzed for currents collected
with R_S voltage error <10 mV at peak current. Activation curves of conductance (G) *vs.* voltage (V)
were generated by dividing peak current (I) by the driving force ($V - E_{rev}$) to obtain G. E_{rev} was
approximated by the equilibrium potential for Na^+ (E_{Na}) for the specified Na^+ concentrations (Table
1). The resulting G-V curves were generally well described by fitting a simple Boltzmann function
using the Levenberg-Marquardt fitting algorithm as implemented in OriginPro:

$$G(V) = \frac{G_{max}}{1 + \exp\left(\frac{V_{act} - V}{s}\right)} \qquad \text{Eq 1}$$

where G_{max} is the maximum conductance, V_{act} is the voltage of half-maximal activation, and s is the
slope factor.

Inactivation G-V curves were generated by measuring how iterated prepulse voltages affect the
conductance evoked by a test pulse (step) in the activation range (–35 or –15 mV). Peak current
evoked by the test pulse was divided by driving force, plotted against the iterated prepulse voltage,
and fit by a Boltzmann function:

$$G(V) = \frac{G_{max}}{1 + \exp\left(\frac{V - V_{inact}}{s}\right)} \qquad \text{Eq 2}$$

195 with V_{inact} equaling the voltage of half-maximal inactivation.

196 Coefficient of variation (CV) was used to measure regularity of spike trains and was calculated for a
197 given spike train from the mean and standard deviation of interspike interval (ISI):

$$CV = \frac{\sigma(ISI)}{\mu(ISI)} \qquad \text{Eq 3}$$

To compare intrinsic, rate-independent, regularity, we wished to compare CV at similar rates.
Because CV varies inversely with spike rate, we manipulated the input current amplitude to hold
spike rate approximately constant between control and experimental conditions. Events were counted
as spikes when they crossed a voltage threshold, between –20 and 0 mV, which was set for each
neuron to exclude subthreshold excitatory postsynaptic potentials (EPSPs). We confirmed that APs
and not EPSPs were counted by comparing the waveforms: APs had higher peak values, faster rise
times and decays, and, often, afterhyperpolarizations (AHPs, measured as the difference between the
resting potential and the most negative potential following a spike).

206 2.2 Modeling

207 A Hodgkin-Huxley model of neuronal firing was used to assess the individual impacts of I-Na$_V$T, I-
208 Na$_V$R and I-Na$_V$P on spiking. The single-compartment model was implemented in MATLAB 2023b
209 as a differential equation in which the net current across the neuronal membrane was taken as the
210 sum of various individual currents (Hodgkin and Huxley, 1952):

$$I_{inj} = C_m\, S\, \frac{dV}{dt} + I_{Na} + I_{KLV} + I_{KH} + I_H + I_{leak} + I_{EPSC} \qquad \text{Eq 4}$$

This model is an extension of the single-compartment VGN model developed by Hight and Kalluri
(2016) and Ventura and Kalluri (2019) to fit rat VGN data. Membrane voltage $V(t)$ was solved
numerically using a backwards difference method. The specific membrane capacitance (C_m) was
fixed at 0.9 µF/cm^2 (Gentet et al., 2000). Cell surface area (S) was fixed to yield a net capacitance of
15 pF, the average for our recorded mouse VGNs. The 5 ionic currents represent key current types in
vestibular ganglion neurons: voltage-gated sodium (I_{Na}), low-voltage-activated potassium (I_{KLV}),
high-voltage-activated potassium (I_{KH}), hyperpolarization-activated cyclic nucleotide-gated (I_H), and
leak (I_{leak}). The model VGN was stimulated either by injected current steps (I_{inj}) with I_{EPSC} set to 0,
or with trains of simulated vestibular excitatory postsynaptic currents (pseudo-EPSCs, I_{EPSC})
with I_{inj} set to 0. A holding period of 500 ms preceded any current injection to allow the resting
membrane potential to equilibrate.

To ensure that the combinations of parameters for currents reproduced APs and firing patterns of
different VGNs observed in whole-cell recordings (see Fig. 5A and C), we fit the model output to
recorded data using a local search optimization algorithm. This algorithm compared the phase plane
plots of model APs, produced by different combinations of ionic, leak, and injected currents, against
the phase plane plots of averaged recorded APs (Fig. S1, Table 2). We constrained the Na$_V$
conductance parameter range to the specific range recorded for each neuron type in this study (e.g.,
Fig 5B) and, for the other ionic conductances, determined the ranges based on previously published
work (see Table 2 for citations). For each firing pattern, the combination of parameters yielding a
simulated AP with the lowest mean squared error relative to the average recorded AP was accepted;
the resulting simulations are shown in Fig. S1: panel D (AP waveforms), E (phase plane plots) and F
(firing patterns). The model parameter combinations for the 4 identified VGN firing patterns (Fig.
S1C, F) are summarized in Table 2.

In previous versions of the model (Hight and Kalluri, 2016; Ventura and Kalluri, 2019), I_{Na} was
entirely transient (I-Na$_V$T) and based on the formulation in Rothman and Manis (2003). Here, we
adapted I_{Na} to include persistent (I-Na$_V$P) and resurgent (I-Na$_V$R) Na$_V$ currents. We used I-Na$_V$T
equations from Hight and Kalluri (2016) and I-Na$_V$R and I-Na$_V$P equations from Venugopal et al.
(2019) and Wu et al. (2005), substituting in our mouse VGN values for conductance density and
voltage dependence. Figure 1E shows the *I-V* relations of both recorded (continuous lines) and
modeled (dashed lines) Na$_V$ current components.

The equation for total Na$_V$ current was based on the computational model by Venugopal et al. (2019)
and can be written as:

$$I_{Na} = I_{NaVT} + I_{NaVP} + I_{NaVR} \qquad \text{Eq 5}$$

243 where I_{NaVT} , I_{NaVP}, and I_{NaVR} are modeled as:

$$I_{NaVT} = \bar{g}_{NaVT}(m_T{}^3 h_T)(V - E_{Na}) \qquad \text{Eq 6}$$

$$I_{NaVP} = \bar{g}_{NaVP}(m_{P_\infty} h_P)(V - E_{Na}) \qquad \text{Eq 7}$$

$$I_{NaVR} = \bar{g}_{NaVT}((1 - b_R)^3 h_R{}^5)(V - E_{Na}) \qquad \text{Eq 8}$$

Model parameters for Na$_V$ current modes are summarized in Table 3. Conductance densities ($\bar{g}$)
cover the range of experimentally derived values from this study on VGN cell bodies and the data of
Meredith and Rennie (2020) on Na$_V$ currents in VGN calyx terminals on hair cells: maximal
persistent conductance ($\bar{g}_{NaVP}$) was set to 3% of the transient conductance ($\bar{g}_{NaVT}$) and the maximum
resurgent conductance ($\bar{g}_{NaVR}$) was set to 10% of $\bar{g}_{NaVT}$, as recorded from cell bodies in this study and
as recorded in vestibular afferent (calyceal) terminals (Meredith and Rennie, 2020).

Transient sodium current: The steady-state voltage-dependent activation (m_T) and inactivation (h_T)
of Na$_V$T currents are modeled as in Rothman and Manis (2003), with voltage of half-activation ($V_{1/2}$)
and slope factor (*s*) set equal to the mean values from this study.

$$m'_T = \frac{m_{T_\infty} - m_T}{\tau_{m_T}} \qquad\qquad \text{Eq 9}$$

$$h'_T = \frac{h_{T_\infty} - h_T}{\tau_{h_T}} \qquad\qquad \text{Eq 10}$$

$$m_{T_\infty} = \left[1 + exp\left(\frac{-(V - V_{1/2})}{s}\right)\right]^{-1} \qquad\qquad \text{Eq 11}$$

$$h_{T_\infty} = \left[1 + exp\left(\frac{V - V_{1/2}}{s}\right)\right]^{-1} \qquad\qquad \text{Eq 12}$$

253 Steady state voltage-dependent time constants of activation and inactivation functions for $Na_V T$
254 currents, originally from Rothman and Manis (2003), remained mostly unchanged:

$$\tau_{m_T} = 10\left\{5\ exp\left[\frac{V + 60}{18}\right] + 36\ exp\left[-\frac{V + 60}{25}\right]\right\}^{-1} + 0.04 \qquad\qquad \text{Eq 13}$$

$$\tau_{h_T} = 100\left\{7\ exp\left[\frac{V + 60}{11}\right] + 10\ exp\left[-\frac{V + 60}{25}\right]\right\}^{-1} + 0.6 \qquad\qquad \text{Eq 14}$$

255 ***Persistent and resurgent sodium currents***: Steady-state activation (h_P), inactivation (m_P), and
256 voltage-dependent time constant of inactivation (τ_{hP}) for $Na_V P$ current are based on Venugopal et al.
257 (2019) and Wu et al. (2005) and are modeled as:

$$m_{P_\infty} = \left[1 + exp\left(\frac{-(V - V_{1/2})}{s}\right)\right]^{-10} \qquad\qquad \text{Eq 15}$$

$$h'_P = \frac{h_{P_\infty} - h_P}{\tau_{h_P}} \qquad\qquad \text{Eq 16}$$

$$h_{P_\infty} = \left[1 + exp\left(\frac{V - V_{1/2}}{s}\right)\right]^{-1} \qquad\qquad \text{Eq 17}$$

$$\tau_{h_P} = 100 + \frac{10000}{1 + exp\left(\frac{V + 60}{10}\right)} \qquad\qquad \text{Eq 18}$$

258 The formulation for I-$Na_V R$ is from Venugopal et al. (2019). It alters the Hodgkin-Huxley
259 conductance-based formulation to incorporate state-dependent Na^+ resurgence due to unblocking of a

260 channel that was blocked upon opening. The equations that govern voltage-dependent
261 blocking/unblocking (b_R) kinetics are as follows:

$$b'_R = \alpha_b(1 - b_R)b_{R_\infty} - k_b\beta_{b_R}b_R \qquad \text{Eq 19}$$

$$b_{R_\infty} = \left(1 + exp\left(\frac{V - V_{1/2}}{s}\right)\right)^{-1} \qquad \text{Eq 20}$$

$$\beta_{b_R} = \left(1 + exp\left(\frac{-(V - 40)}{8}\right)\right)^{-2} \qquad \text{Eq 21}$$

262 where constants α_b (0.08) and k_b (0.9) control the rate of unblocking. The voltage-dependent
263 inactivation of I-Na$_V$R (h_R) functions include:

$$h'_R = \alpha_{h_R}h_{R_\infty} - 0.8\beta_{h_R}h_R \qquad \text{Eq 22}$$

$$h_{R_\infty} = \left(1 + exp\left(\frac{V - V_{1/2}}{s}\right)\right)^{-1} \qquad \text{Eq 23}$$

$$\alpha_{h_R} = \frac{1}{1 + exp\left(\frac{-(V + 45)}{8}\right)} \qquad \text{Eq 24}$$

$$\beta_{h_R} = \frac{0.5}{1 + exp\left(\frac{-(V + 45)}{15}\right)} \qquad \text{Eq 25}$$

Synaptic conductance and EPSC shape: Synaptic input was generated and modeled as described in
Hight and Kalluri (2016). Briefly, modeled synaptic events were randomly drawn from Gaussian
distributions of size and timing based on EPSC amplitudes and rates. For our model simulations, we
simulated EPSCs with a shape based on EPSCs of vestibular afferent calyces in the excised P8 CD-1
mouse utricular epithelium (lateral extrastriolar zone) at room temperature. These have longer onset
and decay times than the S1 EPSC shape used in Hight and Kalluri (2016) (Fig. S2). An exponential
function was fitted to an averaged synaptic event:

$$s(t) = 3.112 * exp(-0.4545t) - 3.112 * exp(-1.121t) \qquad \text{Eq 26}$$

3   Results

Whole-cell ruptured-patch recordings were taken from somata of vestibular ganglion neurons
(VGNs; n = 146) dissociated from CD1 mice of both sexes between P3 and P25 (median = P8) and
cultured overnight. VGN cell bodies allow better space clamp than afferents recorded _in vivo_.

Because Na$_V$P and Na$_V$R current modes upregulate with development (Browne et al., 2017; Hong et
al., 2018; Meredith and Rennie, 2020), we needed to record beyond the first postnatal week, by
which time myelin is extensive. With overnight culture, VGN cell bodies lose their myelin wrapping,
allowing access for recording, but have not yet acquired long neurites that occur with long term
culture.

First, we show Na$_V$T, Na$_V$P, and Na$_V$R current modes recorded from VGNs, and their contributions
to voltage dependence and time course of the overall Na$_V$ current (I-Na$_V$). We describe experiments
investigating contributions of a channel subunit, Na$_V$1.6, that carries all three current modes. We then
assess the roles of Na$_V$ currents in AP waveforms and spiking during current clamp. Finally, we
examine the roles of each current mode in a computational model of VGN spiking.

3.1   Properties of Na$_V$ Currents in VGNs

To collect and characterize Na$_V$ currents, we used
solutions that lowered R$_S$ voltage errors by
decreasing Na$_V$ currents: "Cs$^+$" internal solution
and "Na$^+$ Reduced" external solution (Table 1),
which had Cs$^+$ instead of K$^+$ and no added Ca^{2+}
to minimized contamination with K$^+$ or Ca^{2+}
current. We recorded at room temperature
(~24°C) to slow activation speed of fast Na$_V$
currents into the range for which our voltage
clamp time constant is adequate (~15-50 μs, see
Methods).

All VGNs had I-Na$_V$T, some had I-Na$_V$P, and few
had I-Na$_V$R.

Figure 1 shows exemplar Na$_V$ currents recorded
from VGNs in response to voltage protocols
previously designed to reveal Na$_V$T, Na$_V$P, and
Na$_V$R current components (Stafstrom et al., 1982;
Raman and Bean, 1997). Depolarizing steps
following a hyperpolarizing prepulse revealed
fast-inactivating I-Na$_V$T in all VGNs (Fig. 1B),
as previously reported (Chabbert et al., 1997;
Risner and Holt, 2006; Liu et al., 2016). Table 4
summarizes I-Na$_V$T properties.

Figure 1

To reveal Na$_V$P current, we eliminated rapidly inactivating I-Na$_V$T by applying a slow depolarizing
ramp (0.1 mV/ms from −80 to +60 mV) and obtained the TTX-sensitive non-inactivating component
by subtracting the ramp current remaining in 1 μM TTX (Fig. 1C). This method often revealed I-
Na$_V$P as a small TTX-sensitive inward current activating above –70 mV. I-Na$_V$P was evident in 42 of
78 (54%) VGNs, P3-25, always in combination with I-Na$_V$T and in 4 cases with I-Na$_V$R as well.

I-Na$_V$R was revealed by stepping briefly from –125 mV to +25 mV to open Na$_V$ channels, then
repolarizing to iterated negative voltages (Fig. 1D) (Raman and Bean, 1997). It is thought that a

blocking particle enters the channel as it opens at the activating voltage and prevents inactivation;
with repolarization the channel unblocks, yielding I-Na$_V$R, which then inactivates (Bant and Raman,
2010; White et al., 2019). Overall, I-Na$_V$R was much less common than I-Na$_V$P, occurring in just 6 of
78 VGNs (8%) tested in voltage clamp. All the cells with I-Na$_V$R were older than P10; for this age
group, the incidence was 6/49 or 12%. Two of the 6 VGNs had I-Na$_V$T and I-Na$_V$R but no I-Na$_V$P.
Developmental upregulation of I-Na$_V$R has been described in vestibular afferent endings, where I-
Na$_V$R was found in 35/67 (54%) calyces >P30 (Meredith and Rennie, 2020).

Relative to I-Na$_V$T, which for iterated voltage steps peaked at −20 mV, I-Na$_V$R and I-Na$_V$P reached
maximal amplitude at –45 mV repolarization voltage and −25 mV ramp voltage, respectively (Fig.
1E). Although, on average, peak I-Na$_V$P and peak I-Na$_V$R were just 1% and 3% of peak total I-Na$_V$,
the small Na$_V$P and Na$_V$R currents can be relatively more substantial at subthreshold voltages; for
example, from −60 to −50 mV, average I-Na$_V$P is 13% and I-Na$_V$R is 124% of the peak I-Na$_V$T over
the same voltage range. Note that I-Na$_V$R is present only following a depolarizing prepulse or, *in*
vivo, an AP.

I-Na$_V$T, I-Na$_V$P, and I-Na$_V$R were all completely blocked by 1 μM TTX. In contrast, in "acute"
recordings from immature rat VGNs (P < 8) on the day of dissociation, Liu et al. (2016) recorded
multiple kinds of I-Na$_V$T with different TTX sensitivities and kinetics: TTX-insensitive Na$_V$1.5
current (IC$_{50}$ ~300 nM TTX) and TTX-resistant Na$_V$1.8 current (no block at 5 μM TTX), in addition
to TTX-sensitive current. As in other studies of overnight-cultured VGNs (Chabbert et al., 1997;
Risner and Holt, 2006; Liu et al. 2016), we did not detect TTX-insensitive or -resistant currents, even
with 300 nM TTX to block the large TTX-sensitive currents (n = 5, not shown).

The addition of I-Na$_V$P and I-Na$_V$R affects overall activation voltage.

Cells with I-Na$_V$R had larger average peak current density relative to VGNs with both Na$_V$T and
Na$_V$P currents (I-Na$_V$(T+P)) or just I-Na$_V$T (Fig. 2A; collected with protocol shown in Fig. 1A). The
2 VGNs with I-Na$_V$(T+R) are shown separately in Figure 2A and B and had unusually large current
densities (Fig. 2C, green) and negative midpoints of activation and inactivation (Fig. 2B).

To measure how Na$_V$P and Na$_V$R currents influenced the voltage dependence of the total Na$_V$
current, we fit activation and inactivation (peak G-V) curves with Boltzmann functions (Eqs. 1 and 2;
Fig. 2B) and compared fit parameters (Table 4; statistical tests in Table S1). There was no difference
detected in half-maximal activation potential (V$_{1/2, Act}$) values between VGNs with I-Na$_V$T and I-
Na$_V$(T+P) (Fig. 2B); although the statistical power is low, the lack of clear difference is not
surprising given the small size of I-Na$_V$P. Cells with I-Na$_V$R (n = 6), however, had V$_{1/2, Act}$ values
shifted significantly negative to cells with I-Na$_V$(T+P) or just I-Na$_V$T (Fig. 2B). The negative shift of

Figure 2

activation voltage suggests that I-Na$_V$R may decrease the current threshold for spiking, possibly by
reducing overall rates of inactivation. Half-maximal inactivation potential (V$_{1/2, Inact}$) did not differ
significantly across groups (Fig. 2B).

In Figure 2C, peak current density values have been converted to maximum Na$_V$ conductance density
(Na$_V$ G$_{Max}$ density). To assess the effect of I-Na$_V$R, we pooled all VGNs with I-Na$_V$R [(T+P+R) and
(T+R)]. VGNs with I-Na$_V$(T+P+R) had greater total Na$_V$ G$_{Max}$ density relative to I-Na$_V$T and I-
Na$_V$(T + P) (Table 4). This indicates that VGNs with resurgent current have a greater I-Na$_V$
conductance. Later (see Fig. 8), we use the computational model to compare the effects of increasing
total conductance with just I-Na$_V$T current vs. I-Na$_V$T plus P and/or R modes.

Major fractions of Na$_V$T, Na$_V$P, and Na$_V$R currents flow through Na$_V$1.6 channels.

In neurons with resurgent currents, such as cerebellar Purkinje cells, Na$_V$1.6 channels can carry all
three current modes (Raman et al., 1997). Purkinje neurons in Na$_V$1.6-null mice have reduced Na$_V$T
and Na$_V$P current, and almost no Na$_V$R current (Raman and Bean, 1997; Raman et al., 1997; Khaliq
et al., 2003; Do and Bean, 2004). We tested for Na$_V$1.6 contributions to the Na$_V$T, Na$_V$P, and Na$_V$R
current components using the Na$_V$1.6 blocker, 4,9-ah-TTX, at a dose (100 nM) that is ~10-fold
higher than the IC50 (IC50 = 8 nM; Rosker et al., 2007) and still highly selective for Na$_V$1.6 relative
to other subunits.

Figure 3

In Na$^+$-reduced external
solution, 100 nM 4,9-ah-
TTX blocked approximately
70% of I-Na$_V$T in VGNs
(Fig. 3A, B). Control and
blocked (i.e., 4,9-ah-TTX
sensitive currents) currents
had similar V$_{1/2, Act}$ and V$_{1/2,}$
$_{Inact}$ values (Table 4). This is
not surprising, given that
the blocked current makes
up most of the total current.
V$_{1/2,Inact}$ was, however, more
negative for the unblocked
(residual, 4,9-ah-TTX
insensitive) current than for
control current; V$_{1/2,Act}$ was
not significantly different
(Table S1). This suggests
the possibility of a second

TTX-sensitive current that is not carried by Na$_V$1.6 channels and has a more negative inactivation
voltage dependence. The voltage dependencies of inactivation and activation of the two TTX-
sensitive, transient conductances (blocked putative Na$_V$1.6 and residual current) are consistent with
observations on isolated rat VGNs (Liu et al., 2016).

Na$_V$P and Na$_V$R currents were also blocked by 100 nM 4,9-ah-TTX. Block of I-Na$_V$P was seen in
responses to voltage steps (Fig. 3A inset) or to the slow voltage ramp (Fig. 3C) (n = 13). We were
able to test 4,9-ah-TTX on just 1 of the 6 VGNs with Na$_V$R; it produced a strong block of I-Na$_V$R

(Fig. 3D). These results with a $Na_V1.6$-selective blocker suggest that $Na_V1.6$ channels carry the
majority of I-Na_V in cultured VGNs, including ~50-70% I-Na_VT and >90% of I-Na_VP and most of I-
Na_VR. Similar results were obtained in calyx terminals (Meredith and Rennie, 2020).

I-Na_VT and I-Na_VP were enhanced by Na_V channel agonist ATX-II.

The sea anemone toxin ATX-II interacts with Na_V channel gating, slowing down or preventing
inactivation and thereby increasing Na_V current (Oliveira et al., 2004). ATX-II enhances Na_VP
current in vestibular afferent calyces (Meredith and Rennie, 2020) and Na_VR and Na_VP currents in
spiral ganglion neurons (Browne et al., 2017) and dorsal root ganglion neurons (Klinger et al., 2012).
We tested the impact of ATX-II on I-Na_V modes in mouse VGN cell bodies.

100 nM ATX-II increased maximum I-Na_VT in 3 of 7 VGNs tested (Fig. 4A). In all VGNs,
inactivation of I-Na_VT was slowed, resulting in increased I-Na_VP at the end of depolarizing steps
(Fig. 4A) and during voltage ramps (Fig. 4B). On average, I-Na_VP increased more than 5-fold (Fig.
4C); the example in Figure 4B was the largest effect seen. We detected no significant difference with
ATX-II in $V_{1/2, Act}$ and $V_{1/2, Inact}$ values for I-Na_VP and I-Na_VT (Table 4, Table S1). For I-Na_VT, ATX-
II increased the slope factor (decreased the steepness of voltage dependence) of inactivation (p =
0.009, Hedges' g = 0.85, large effect) but not activation (p = 0.85, power = 0.05) (not shown). We
were not able to test ATX-II on resurgent current due to its low incidence.

Figure 4

3.2   Roles of Na_V currents during action potentials and spike trains

To characterize AP waveforms and evoked firing patterns, we recorded in current clamp mode from
62 VGNs in K^+ based standard solutions (Table 1). Without reduced external Na^+, voltage clamp was
less satisfactory: the large fast I-Na_V often escaped voltage clamp, obscuring the small, non-
inactivating Na_VP and Na_VR currents. Although we could not identify I-Na_VP or I-Na_VR in whole-
cell voltage clamp recordings, we characterized some features of I-Na_VT and tested for effects of the
Na_V channel blocker and agonist on spike waveform and firing pattern.

Na_V conductance correlated with features of the action potential waveform.

We classified VGNs into four groups based on firing patterns evoked by depolarizing current steps
just above current threshold ($\pm$ 50 pA), following the scheme of Ventura and Kalluri (2019) (Fig.
5A). Transient neurons fired 1 or 2 spikes at step onset independent of step size. Sustained-firing
neurons ranged from sustained-A type, with spike trains lasting the 500-ms depolarizing step, to
sustained-B type, with shorter trains, to sustained-C type, with 2 or more small spikes that devolve

into voltage oscillations. Accommodation could increase as more current was injected (Kalluri et al.,
2010).

Figure 5

Transient VGNs had a significantly higher current threshold for spiking relative to sustained-A and
sustained-B VGNs but we detected no significant difference in resting potential (V_{rest}), input
resistance (R_{in}), or membrane capacitance (C_m) across firing patterns (Table S2). While the incidence
of the transient firing pattern was stable with age, at ~50%, the distribution of sustained firing
patterns changed from an approximate balance across sustained-A, -B, and -C subtypes for ages
below P10 (n = 21) to mostly sustained-B above P10 (6/7) (Fig. S3). Similar changes with age were
reported in rat VGNs (Ventura and Kalluri, 2019).

We assessed the AP waveform associated with each firing pattern (Fig. 5C, summarized in Table 5,
and detailed in Table S3). APs from transient VGNs were smaller than APs from sustained-A VGNs
(ΔV_{AP}, Fig. 5C) and had slower peak rates of depolarization (peak dV/dt) than APs from sustained-A
and sustained-B VGNs (Fig. 5F). Time-to-peak and voltage threshold of APs did not differ
significantly across firing patterns (Fig. 5C, D).

For 56 cells, we also collected Na_V currents in voltage clamp and fit activation (peak G-V) curves
(Eq. 1) to measure maximum Na_V conductance density (Na_V G_{Max} density) (Fig. 5B). We had

hypothesized that Na_V G_{Max} density would be highest for sustained-A VGNs and lowest for transient
VGNs but actually did not detect a significant difference in Na_V G_{Max} density across firing pattern
populations, although this was a low-powered observation (Fig. 5B, Table 5). Spike height (Fig. 5C)
and peak dV/dt from phase plane plots (Fig. 5F; Bean, 2007) both correlated with Na_V G_{Max} density
(Fig. 5D, G), as expected given that Na_V currents drive the rising phase of the AP. We saw no
relationship between Na_V G_{Max} density and age (Fig. 5H).

The spike afterhyperpolarization (AHP) was significantly greater in sustained-A neurons than any
other firing groups (Fig. 5E, Table 5, Table S3). The AHP preserves sodium channel availability by
relieving inactivation, shortening the refractory period, and allowing sustained and regular firing at
high rates (Gittis et al., 2010). Differences in AHP have previously been attributed to differences in
K_{LV} conductance that affect resting potential, current threshold, and membrane recovery time (Hight
and Kalluri, 2016; Kalluri et al., 2010; Iwasaki et al., 2008). In particular, the low K_{LV} conductance
of immature sustained-A VGNs can contribute by making V_{rest} less negative.

In summary, maximum Na_V conductance density did not clearly vary with firing pattern but
correlated with features of the spike waveform. We had hypothesized that sustained-A VGNs would
have the highest Na_V G_{Max} densities and transient VGNs the lowest densities but did not detect a
significant difference in average Na_V G_{Max} density across firing pattern populations, except that the
range of G_{Max} densities was much higher for the transient VGNs. Key features of the AP waveform –
spike amplitude, AHP, and peak rate of depolarization – did positively correlate with Na_V G_{Max}
density (Fig. 5).

Blocking Na_V1.6 currents reduced excitability and altered AP waveform.

We probed the effects of the Na_V1.6 blocker 4,9-ah-TTX on spiking (Fig. 6), which greatly reduced
Na_VT, Na_VP, and Na_VR currents recorded in Na^+ Reduced/Cs^+ solutions (Fig. 3). In standard
recording conditions, 100 nM 4,9-ah-TTX showed a similar percent block of I-Na_VT for transient
VGNs ($50.1 \pm 11.3\%$) and sustained VGNs ($52.0 \pm 5.2\%$). Given that Na_V1.6 current expressed in
HEK cells has similar 4,9-ah-TTX sensitivity (Denomme et al., 2020), Na_V1.6 channels may carry
most of the Na_V current in our isolated and cultured VGNs.

100 nM 4,9-ah-TTX affected step-evoked firing quantitatively, as shown in Figure 6. For all 13
VGNs tested, the Na_V1.6 blocker increased current threshold for 500-ms steps. For sustained VGNs,
4,9-ah-TTX reduced the number of APs (spike rate) at spiking threshold and throughout the family of
current steps (Fig. 6A). For transient VGNs, 100 nM 4,9-ah-TTX increased current threshold for
spiking and decreased spike amplitude (Fig 6B).

To assess changes in the AP waveform, we temporally aligned the peaks of the first APs evoked by
long current steps (Fig. 6C, detailed in Table S4). 4,9-ah-TTX reduced spike height by ~20 mV
(~25%) on average (Fig. 6D). AHP depth was also reduced, possibly because hyperpolarizing K^+
currents were less activated during the smaller spike (Fig. 6E) and AP time-to-peak increased (Fig.
6F). No change was detected in spike width at half-height, although control spikes are narrower at
their peaks, reflecting their higher rates of depolarization and repolarization (Fig. 6G, H).

4,9-ah-TTX substantially hyperpolarized V_{rest} in sustained VGNs (Fig. 6I) but not transient VGNs
(Fig. 6J). These data suggest that in sustained VGNs, 4,9-ah-TTX blocks depolarizing channels that
are open at rest, such as I-Na_VT and I-Na_VP. The larger impact of the blocker on sustained VGNs
may reflect a different balance of channels open at rest: they have smaller K_{LV} conductances (Kalluri

et al., 2010) and may also have larger
Na$_V$ conductances open at rest,
leading to a relatively more
depolarized V$_{rest}$.

To test the impact of Na$_V$1.6 current
on spike regularity (CV), we
stimulated firing with frozen trains of
synthetic ("pseudo") EPSCs. The
pseudo-EPSCs had pseudo-random
timing to represent noisy quantal
input from hair cells to afferent
terminals where spiking normally
initiates (Kalluri et al., 2010). In 4 of
14 VGNs tested, block of Na$_V$1.6
current with 4,9-ah-TTX eliminated
EPSC-induced spiking entirely. In
the remaining 10 VGNs (6 transient
and 4 sustained), we measured
regularity as coefficient of variation
(CV). To avoid rate effects on CV,
we held spike rates to control levels
(15.1 ± 1.3 vs 15.3 ± 1.7 spikes/s) by
controlling EPSC size (Fig. 6K; see

Figure 6

Kalluri et al., 2010). However, blocking Na$_V$1.6 current did not significantly affect CV (Fig. 6L).

In summary, in all firing pattern groups, blocking Na$_V$1.6 current with 4,9-ah-TTX altered spike
wave form, decreasing spike amplitude, AHP, time-to-peak, and rate of depolarization. In sustained
VGNs alone, 4,9-ah-TTX also made resting potential appreciably more negative, showing that
Na$_V$1.6 conductance is a significant factor in setting resting potential. Later (see Fig. S4), we use the
computational model to assess the effects of individual current modes on V$_{rest}$. In a sample of
transient and sustained neurons, blocking Na$_V$1.6 current had no consistent effect on spike regularity
when overall rate was held constant by increasing EPSC size.

ATX-II increased excitability and spiking regularity.

In voltage clamp, 100 nM ATX-II, which reduces Na$_V$ channel inactivation, increased Na$_V$P and
Na$_V$T currents (Fig. 4). We tested its impact on step-evoked firing patterns and pseudo-EPSC-evoked
spike trains (Fig. 7).

In a sample of 9 VGNs, ATX-II reduced current threshold to zero and changed step-evoked firing
patterns toward more sustained categories in 7 VGNs (3 sustained-A, 2 sustained-C, 2 transient); e.g.,
sustained-C VGNs became sustained-A (Fig. 7A). ATX-II also increased spike rate and decreased
CV in responses to trains of pseudo-EPSCs. To control for rate effects on CV, we decreased EPSC
amplitude in ATX-II to match spike rate (35 ± 3.4 sp/s) to the control value (31 ± 2.8, Fig. 7C) and
still found a consistent, if modest, decrease in CV in ATX-II (Fig. 7C, D). Figure 7B shows one of 2
of 9 VGNs, both transient, that remained transient, irregular, and relatively inexcitable in ATX-II.
There was no significant effect on AP waveform (Table S4).

Figure 7

In summary, reducing Na$_V$ channel inactivation with ATX-II lowered current thresholds for both
step-and pseudo-EPSC-evoked firing, in most cases to zero current (spontaneous firing). ATX-II also
slightly increased spike regularity independent of rate. These results are consistent with increased
Na$_V$ channel availability near resting potential, presumably through reduced proportion inactivated
channels, effectively increasing the persistent current.

3.3   Modeling the effects of transient, persistent, and resurgent Na$_V$ currents on VGN firing

We hypothesized that I-Na$_V$P and I-Na$_V$R increase excitability in VGNs by increasing channel
availability near spike threshold, and thus enhance the likelihood of firing. Because excitability has
been associated with regularity in VGNs, we further hypothesized that I-Na$_V$P and I-Na$_V$R enhance
spike timing regularity. Lacking pharmacological tools to disentangle the impacts of Na$_V$T, Na$_V$P and
Na$_V$R currents *in vitro*, we adapted existing models of neuronal firing to create model VGNs
("mVGNs") for which each current mode could be adjusted. Below we show that, while I-Na$_V$T
affected rate and regularity in all mVGNs, I-Na$_V$P significantly increased rate and rate-independent
regularity in the sustained mVGNs and I-Na$_V$R increased rate but decreased rate-independent
regularity in the transient mVGNs.

We combined a single-compartment conductance-based VGN spiking model (Hight and Kalluri,
2016; Ventura and Kalluri, 2019) with equations for I-Na$_V$P and I-Na$_V$R (Wu et al., 2005; Venupogal
et al., 2019). Our recorded range of Na$_V$ conductance density (g$_{NaVT}$) values was 0.2 – 18.4 nS/pF or
0.3 – 20.5 mS/cm^2. We refined the spiking parameters for each mVGN by fitting the phase plane plot
(PPP) of each mVGN to the PPP of the average recorded AP (Fig. 5F, see Methods). Based on the
fitting results, we used 16 mS/cm^2 for sustained-A, 13 mS/cm^2 for sustained-B, 11 mS/cm^2 for
sustained-C and, and 7 mS/cm^2 for transient model neurons (Table 3). These values neatly replicated
AP waveforms, firing patterns, spike rates, and regularity ranges of each VGN type.

To test the impact of Na$_V$ current components, we separately simulated four combinations of Na$_V$T,
Na$_V$P and Na$_V$R current modes of I-Na$_V$ to assess differential effects on firing: 1) "T", 2) "T + P", 3)
"T + R", and 4) "T + P + R". I-Na$_V$P and I-Na$_V$R were simulated with conductance density values as
a percentage of I-Na$_V$T; I-Na$_V$P at 3% I-Na$_V$T and for I-Na$_V$R at 10% I-Na$_V$T. These proportions are
based on our mouse VGN data and values from calyx afferent terminal recordings in semi-intact
vestibular organs (Meredith & Rennie, 2020) (Methods and Table 3). Current-voltage relations of the
simulated currents reproduce the voltage dependence of experimental data (dashed curves in Fig.
1D).

Another set of simulations explored whether any changes with adding I- Na_VP and I-Na_VR were
simply the effect of increasing Na_VT conductance (g_{NaVT}). We increased g_{NaVT} ("T+") to match total
conductances achieved in other simulations by adding Na_VP and/or Na_VR conductances, and to
explore the range of g_{NaVT} recorded (Fig. 5B).

Adding I-Na_VR or I-Na_VP increased instantaneous firing rates and altered spike waveforms evoked
by current steps.

We tested responses of mVGNs to
500-ms current steps for
comparison with step-evoked
spike patterns (illustrated for real
VGNs in Figure 5A). In Figure
8A, we show, for the 5 I-Na_V
combinations, the first 50 ms of
the response to a current injected
after a 500-ms holding period (0
injected current). Individual Na_V
current modes during spiking are
shown in Figure 8B – D. To
highlight any effects on spike
waveform, Figure 8 (E, F) shows
the waveforms and phase-plane
plots of aligned first spikes of
each mVGN firing pattern shown
above. These simulations show
that adding I-Na_VR and I-Na_VP
increased excitability in sustained
mVGNs.

Figure 8

Adding I-Na_VP produced a small
inward current at 0 injected
current, before spiking onset (Fig.
8D), in all mVGNs. This was
largest in the sustained-A mVGN
(Fig. 8A, left), where adding I-
Na_VP depolarized V_{rest} (blue
arrow), reduced spike height (Fig.
8E, blue arrow) and rate of rise
(Fig. 8F), thus altering the AP
waveform (Fig. 8F). In both T+P
and T+P+R conditions, the added
I-Na_VP affected the first spike in the train by decreasing time-to-peak by 11%, spike height by 19%,
spike width by 20% and peak dV/dt by 30% (Table S5); adding I-Na_VR (i.e., T+P+R) had no
additional effect. I-Na_VP also increased AHP by 36%, reducing I-Na_VT inactivation and so leading to
larger Na_VT and P currents and driving larger spikes later in the train. I-Na_VP had much smaller
effects on AP waveforms of other mVGNs. In sustained-B and -C mVGNs, I-Na_VP enhanced the
number/rate and size of spikes and oscillations after the first spike (1 – 4% decrease in ISI; blue
arrows, middle columns, Fig. 8A).

Effects of I-Na$_V$R on spike waveform were first evident after a spike had occurred, as expected from
Na$_V$R current's requirement for a large activating depolarization (AP upstroke) to block the channels,
followed by relief of block by the AP downstroke (Fig. 8C; also Fig. 1D). In sustained-A mVGNs,
adding I-Na$_V$R to I-Na$_V$T (i.e., T+R) decreased ISI, shortening AHP duration by 52% and reducing
ISI by 16%, therefore changing instantaneous spike rate and changing the timing of individual spikes
in the train (green arrow; Fig. 8A). I-Na$_V$R had smaller or negligible effects in sustained-B, -C and
transient mVGNs, likely because their smaller APs fail to reach the voltage range of "blocking" for I-
Na$_V$R (> +25 mV; Raman and Bean, 2001). This may also account for the dominance of I-Na$_V$P
current, such that T+P+R simulations resembled T+P simulations: depolarization of V$_{rest}$ by I-Na$_V$P
yielded smaller APs, effectively eliminating the impact of I-Na$_V$R (Fig. 8C, pink traces).

We compared the effects of adding Na$_V$R and Na$_V$P currents to the effect of increasing the Na$_V$T
conductance alone (i.e., T+; Fig. 8, orange traces). The effects on step-evoked firing of sustained
mVGNs were not the same. Increasing I-Na$_V$T decreased ISI after the first spike (by <1 ms, Table
S5) less than either I-Na$_V$P or I-Na$_V$R (~3%) (Fig. 8A, B). Increasing I-Na$_V$T affected first-spike
waveforms more than I-Na$_V$R or I-Na$_V$P, substantially increasing spike height and peak rate of rise of
first-spike AP waveforms for all firing patterns (compare orange curves with all others, Fig. 8E, F).

In summary, adding I-Na$_V$R and/or I-Na$_V$P slightly increased instantaneous spike rate in step-evoked
firing of sustained mVGNs. Increasing I-Na$_V$T to the same total conductance affected spike height
and rate of rise more than I-Na$_V$R and/or I-Na$_V$P.

Na$_V$P current affected resting potential more than Na$_V$T or Na$_V$R currents.

Adding I-Na$_V$P depolarized the resting membrane potential in sustained-A mVGNs (Fig. 8A, blue
arrow). To simulate the 4,9-ah-TTX block in real VGNs (Fig. 6), we simulated step-evoked firing in
the sustained-A mVGN and transient mVGN with I-Na$_V$T reduced by 70% and I-Na$_V$P and I-Na$_V$R
by 90% (Fig. S4). Again, we set I-Na$_V$P at 3% and I-Na$_V$R at 10% I-Na$_V$T. The simulated block
replicated observations in real VGNs. Current thresholds increased and spike heights fell in both
mVGNs. The diminution of the spike was most pronounced in the sustained-A mVGN, which also
showed reduced spike rate and clearly hyperpolarized V$_{rest}$. V$_{rest}$ was negligibly affected in the
transient mVGN.

We analyzed the influence of each Na$_V$ current mode on the change in V$_{rest}$ in sustained mVGN by
varying P, R, and T combinations (Fig. S4C). Blocking I-Na$_V$R had no effect on V$_{rest}$ in any mVGN
because resurgent current requires activity to activate. Blocking I-Na$_V$T had at most a small effect on
V$_{rest}$ and then only in sustained mVGN (~1 mV). Virtually all the effect on V$_{rest}$ was through I-Na$_V$P:
6 mV for sustained mVGN and 0.5 mV for the transient mVGN.

Therefore, reduction in Na$_V$T, Na$_V$P, and Na$_V$R currents in mVGNs reproduced key effects of a
Na$_V$1.6 channel blocker and show I-Na$_V$P driving 5-10 mV depolarization of V$_{rest}$ in sustained-A
mVGNs. This simulation indicates that even if Na$_V$P currents are equally present in sustained and
transient VGNs (as we have modeled them), the effect of Na$_V$P current is substantially stronger in
sustained VGNs.

Adding I-Na$_V$T, I-Na$_V$P, and I-Na$_V$R increased spike rate evoked by pseudo-EPSC trains and
differentially affected regularity.

To examine the roles of Na$_V$ current modes in spike rate and regularity, we drove sustained-A and
transient mVGNs with 1-second trains of simulated synaptic events (pseudo-EPSCs, first 500 ms

shown in Fig. 9A, B), modeled
after EPSCs recorded from
vestibular afferent calyx events
and quasi-randomly distributed in
time (see Methods). We generated
a set of 5 different pseudo-EPSC
trains and applied the same set to
each simulation condition shown
in Figure 9. Therefore, the input
noise is similar across each g_{NaVT}
value in Figure 9. Each
measurement in panels C-H is the
mean and SEM for the 5 pseudo-
EPSC trains at the specified g_{NaVT}
and combinations of NavR and
NavP conductances. Figure 9B
(bottom trace) shows one of the
pseudo-EPSC trains, with an inset
highlighting a short onset segment
at high gain. Again, we held NavP
and NavR conductance at 3% and
10% of g_{NaVT}, respectively. Larger
% values of NavP are explored in
Supplemental Figure 5.

Figure 9

First, we assessed the effects of I-
NavT alone on spike rate and spike
regularity at 12 values of g_{NaVT}
density chosen to cover the range
we encountered in VGN
recordings (6 to 22 mS/cm^2 in
intervals of 2; orange traces in Fig.
9A and B). To study spike rate

effects, we generated EPSC trains with average inter-event interval fixed at 5 ms (200 EPSCs/s) and
an amplitude that produced ~20 spikes/s at 14 mS/cm^2, then varied g_{NaVT} density. This firing rate
corresponds approximately to the measured background rate for mouse vestibular afferents (~50
spikes/s, Lasker and Minor 2006) after compensation for the ~15°C lower temperature of our room
temperature VGN recordings. For all mVGNs, spike rate increased and CV decreased with increasing
g_{NaVT} density (Fig. 9C – F), as previously demonstrated by Hight and Kalluri, 2016. For transient
neurons, spike rate increased with g_{NaVT} density, again especially at the low end (Fig. 9D). But CV
showed much less variation with g_{NaVT} density (Fig. 9F). Note that the sustained mVGN in the 'T'
condition is no more regular than the transient mVGN for g_{NaVT} densities <12 mS/cm^2 or more
(compare panels E and F).

In a second set of simulations, we assessed CV at a constant spike rate of 20 spikes/s (CV$_{20sp/s}$) to
eliminate the confounding effect of spike rate on regularity. Rate was controlled by titrating the size
of pseudo-EPSCs in each train, while not interfering with the pseudo-randomly distributed inter-
event intervals. For the sustained-A mVGN (Fig. 9G), increasing g_{NaVT} density decreased CV$_{20sp/s}$ up
to about 12 mS/cms^2, above which CV plateaued. For the transient mVGN (Fig. 9H), in contrast,

there was no obvious change in CV over the entire range of g_{NaVT} tested. The addition of I-Na$_V$R
and/or I-Na$_V$P to I-Na$_V$T affected rate, CV and CV$_{20sp/s}$ differently for the sustained-A mVGN vs. the
transient mVGN (Fig. 9C-H).

Na$_V$P and Na$_V$R currents increase spike rate in sustained and transient mVGNs, respectively.

For the sustained-A mVGN, rate increased substantially when Na$_V$P current was added (p < 0.0001;
Fig. 9C; Table S6) but was not affected by adding Na$_V$R current except for the (T+P+R) condition at
the highest level of g_{NaVT} density tested. For the transient mVGN, adding both I-Na$_V$P and I-Na$_V$R
increased rate for g_{NaVT} density > 12 mS/cm^2, reaching significance at the highest level. The effects
of the two components appeared to be additive (Fig. 9D). In simulations of all three components
(T+P+R), the increase in spike rate is compounded: sustained mVGNs showed a medium to large
effect (Hedge's g = 0.6) and transient mVGNs a small effect (Hedge's g = 0.2) (Fig. 9C, D, Table
S6).

In Supplemental Figure 5, we show the effect of increasing Na$_V$P current to the highest values
recorded at afferent terminals on hair cells, near the spike initiation zone: 5 to 10% of g_{NaVT}
(Meredith and Rennie, 2020). Spike rate increased further as I-Na$_V$P increased. For the sustained-A
mVGN, spike rate doubled as I-Na$_V$P increased from 0-10% (p < 0.0001, Hedges' g = 1.5). For the
transient mVGN, the impact on rate was smaller (~25% increase; p < 0.001, Hedges' g = 0.7).

I-Na$_V$P decreases regularity in sustained mVGNs whereas I-Na$_V$R increases regularity in transient
mVGNs.

Adding Na$_V$P and Na$_V$R conductances had complementary effects on the regularity (CV and CV$_{20sp/s}$)
of sustained-A and transient mVGNs. For the sustained-A mVGN (Fig. 9G), adding Na$_V$P (at 3%)
halved CV$_{20sp/s}$ for all g_{NaVT} density > 8 mS/cm^2, whether Na$_V$R conductance was present or not. For
the transient mVGN, in contrast, Na$_V$P current had no detectable effect, but Na$_V$R current
increasingly enhanced irregularity (both CV and CV$_{20sp/s}$) for g_{NaVT} density > 12 mS/cm^2 (Fig. 9F, H).
One factor in the enhanced irregularity is likely to be depolarization block at high total Na$_V$
conductance (e.g., green arrow, Fig. 9B): instead of returning to V$_{rest}$, the increased proportion of
unblocked channels held V$_m$ depolarized, leading to profound inactivation of Na$_V$T current.

In summary, simulations suggest that I-Na$_V$P and I-Na$_V$R differentially affect spike rate and reinforce
the difference in regularity at constant rate of sustained-A and transient VGNs. Adding P current
made the sustained-A mVGN more regular, and adding R current made the transient mVGN more
irregular, in both cases by factors of ~2.

4   Discussion

4.1   VGN as a model preparation for firing patterns in the vestibular nerve

VGNs are cell bodies that, when mature and *in vivo*, are bipolar, inter-nodal and myelinated.
Research from multiple groups (Chabbert et al., 1997; Limon et al., 2005; Iwasaki et al., 2008) has
shown that isolated VGNs express voltage-gated currents and firing patterns that resemble currents
and firing at spike initiation zones just below the hair cell-afferent terminal synapses in the vestibular
epithelium, as illustrated by immunohistochemistry and direct recordings from calyx terminals (e.g.,
Lysakowski et al., 2011; Songer and Eatock, 2013; Meredith and Rennie, 2020). It is not clear
whether the presence of these channels in the dissociated cell bodies represents true somatic
expression, or whether nodal Na$_V$ channels and paranodal K$_V$ channels on either side of the soma

come along during dissociation. In most cases, investigators have cultured the dissociated VGNs
overnight, causing the VGNs to shed their myelin and expose the neuronal membrane for whole-cell
patch recording.

In rat VGNs, there is some evidence that overnight culturing eliminates expression of some types of
pore-forming Nav channels. Dissociated young rat VGNs that were studied acutely had multiple
transient Nav currents with distinct voltage dependence and TTX sensitivity (Liu et al., 2016),
including a $Na_V1.5$ current (negatively shifted voltage range, TTX-insensitive) and a $Na_V1.8$ current
(positively shifted voltage range, TTX-resistant). These non-TTX-sensitive forms were not detected
in overnight-cultured rat VGNs (Liu et al., 2016). Age differences across studies may also matter.
$Na_V1.5$ expression in ganglion cell bodies may be an immature feature, as suggested by its time
course of expression in small rat DRG neurons (Ranganathan et al. 2002) and some decline in current
expression over the first postnatal week in rat VGNs (Liu et al. 2016). Meredith and Rennie (2020)
recorded TTX-insensitive currents (consistent with $Na_V1.5$) from gerbil afferent calyceal terminals,
and but only in afferents younger than P12.

In summary, isolated and cultured VGNs are compact cell bodies that allow high-quality voltage
clamp recordings and reproduce some but not all naturally occurring ion channel expression.
Importantly for our purposes, they show a range of firing patterns consistent with a role for intrinsic
neuronal properties in setting up the different encoding strategies (temporal vs. rate) characteristic of
irregular and regular afferent populations (Jamali et al., 2016; Cullen, 2019). Previous work has
documented the difference in firing patterns of VGNs and probed their relationship to specific ion
channels: low-voltage-activated K_V1 (Iwasaki et al., 2008) and K_V7 channels (Kalluri et al., 2010),
Ca^{2+}-dependent K channels (Limon et al. 2005), HCN channels (Horwitz et al., 2014) and Nav
channels (Liu et al. 2016). By going beyond correlation and developing a method to interrogate
regularity in these synapse-less somata, a particularly strong case was made for K_{LV} channels (Kalluri
et al., 2010). We took this approach with Nav channel modes and provide support for a substantial
role of relatively small persistent and resurgent currents in excitability and related properties such as
resting potential.

4.2 $Na_V1.6$ is the major Nav subunit in VGNs, contributing to resting potential and excitability

All VGNs in our sample expressed I-NavT and about half also expressed I-NavP with no clear
change in incidence with age (range P3-28), in contrast to mouse cochlear ganglion neurons where
both persistent and resurgent forms increased with maturation (Browne et al., 2017). We saw no I-
NavR before P10. Beyond P10, the incidence was 12% (6/49 tested). VGNs also have voltage-
dependent Ca^{2+} (Cav) currents (L-, N-, P/Q-, R-, and T-type) (Desmadryl et al. 1997; Chambard et al.
1999) which drive Ca^{2+}-dependent K^+ currents that reduce sustained firing (Limón et al., 2005). Here
we suppressed both Cav and Ca^{2+}-dependent K^+ currents by eliminating all but trace Ca^{2+} from the
external medium. Our experiments do not indicate whether sustained VGNs have more NavP or
NavR current than transient neurons, because the conditions for studying firing pattern did not allow
isolation of the small P and R currents. However, our modeling suggests that for transient neurons,
NavP and NavR currents, if present, would have little effect on regularity and resting potential.

RT-PCR of whole rat vestibular ganglia indicated expression of most Nav pore-forming (α) subunits
and all auxiliary (β) subunits (Liu et al. 2016). Block by the $Na_V1.6$-selective blocker, 4,9-ah-TTX,
however, indicates that $Na_V1.6$ channels make up much or all of the current we recorded, for all
modes, again consistent with results from calyx terminals in gerbil cristae (Meredith and Rennie,
2020). In our study, 100 nM 4,9-ah-TTX blocked NavT current by a similar amount (~50%) in

sustained and transiently firing VGNs, suggesting that they express similar proportions of $Na_V1.6$
channels. Persistent and resurgent I-Na_V were almost fully blocked. Cochlear spiral ganglion neurons
also express 4,9-ah-TTX-sensitive transient, persistent, and resurgent currents (Browne et al. 2017),
and are immunoreactive for $Na_V1.6$ at spike initiation zones next to their terminals on hair cells (Kim
and Rutherford, 2016). I-Na_VR is theorized to arise from the internal block of Na_V channels,
canonically $Na_V1.6$, by a positively charged molecule such as Na_V auxiliary subunit β4 (present in
RT-PCR data of the ganglia; Liu et al., 2016) or fibroblast growth factor homologous factor 14
(Raman and Lewis, 2014; White et al., 2019; Aman and Raman, 2024).

The residual Na_V current in 4,9-ah-TTX had significantly more negative inactivation than the control
and blocked currents, but not nearly as negative as the TTX-insensitive current carried by $Na_V1.5$ in
acutely cultured, immature rat VGNs (Liu et al., 2016). RT-PCR data from rat vestibular ganglia (Liu
et al., 2016) suggest $Na_V1.1$, 1.2, 1.3, and 1.7 as candidates.

As expected for a major Na_V current, the $Na_V1.6$ currents are critical to excitability in both firing
types. In both sustained and transient neurons, spikes were substantially smaller in 4,9-ah-TTX, as
expected from the strong block of the dominant transient mode of $Na_V1.6$ current. A large decrease in
AHP size presumably reflected both the small spike height (reducing activation of K^+ currents that
cause AHPs) and the more negative V_{rest} (reducing the driving force, i.e., voltage difference from
E_K). 4,9-ah-TTX blocking experiments showed that $Na_V1.6$ channels contribute to resting potential in
sustained neurons (Fig. 6I; Table S4).

Our modeling also suggests that Na_VP and, to a lesser extent, Na_VT current modes may significantly
contribute to resting conductance in sustained VGNs (Fig. S4). Although I-Na_VP contributed just 3%
of maximum Na_V current density in VGNs, Na_VP current at resting potential may be closer to 10% of
total current, based on its relative voltage dependence and the substantial (~30%) steady-state
inactivation of Na_VT current even at resting potential (Fig. 2B). The small to negligible effect of
blocking $Na_V1.6$ channels on resting potential in transient VGNs (Fig. 6J vs. 6I) is consistent with
resting conductance being dominated by their greater density of K_{LV} channels. HCN conductances
may also contribute to resting membrane potential in VGNs, depending on the balance of resting
conductances (Ventura and Kalluri, 2019).

4.3   Effects of Na_V current modes on AP waveforms, firing patterns, rate, and regularity
AP height and rate of rise varied across firing pattern (Fig. 5C, F) and correlated modestly with
maximum Na_V conductance density (Fig. 5D), both for all VGNs and for transient VGNs alone,
which showed the greatest variance in maximum Na_V conductance density. Firing pattern, in
contrast, did not correlate closely with Na_V G_{max} density. G_{max} density is dominated by the Na_VT
current mode, so these correlations do not reveal roles for I-Na_VP, which has a more significant effect
near resting potential, and I-Na_VR, which acts during spike repolarization. To isolate their effects, we
used simulations.

Our simulations support previous work on isolated VGNs indicating a dominant role for K_{LV} currents
in differentiating spike rate and regularity of sustained (presumed regular) and transient (presumed
irregular) VGNs (Kalluri et al., 2010; Ventura and Kalluri, 2019). K_{LV} currents increase irregularity
independent of rate by enhancing the afferent's sensitivity to high-frequency noise (Kalluri et al.,
2010). We found that the large K_{LV} currents of transient VGNs limited the influence of Na_V current
modes on rate and regularity, but when K_{LV} currents are small, as in sustained VGNs, augmenting
transient and/or persistent Na_V currents increased excitability, spike rate, and rate-dependent
regularity.

In sustained model VGNs, I-Na$_V$P had a stronger effect than other modes on resting potential,
depolarizing V$_{rest}$ by ~6 mV. The effect on V$_{rest}$, in turn, shapes many AP metrics by affecting resting
(input) conductance, inactivation state of I-Na$_V$T, and current and voltage thresholds for spikes (Fig.
8F). I-Na$_V$P also changed AP waveform by reducing time-to-peak and spike height in sustained-A
mVGN APs. Primarily, however, I-Na$_V$P strongly increased firing rate in sustained-A mVGNs.
Regularly firing neurons, like sustained-A VGNs, use firing rate to preferentially encode low
frequency information (Jamali et al., 2016; Cullen, 2019). A mechanism that increases firing at or
near V$_{rest}$ without inactivating will enhance ability to encode small excitatory and inhibitory inputs,
increasing response linearity.

In sustained mVGNs current step simulations (Fig. 8A, B), adding I-Na$_V$R altered instantaneous
spike rate by truncating the AHP, decreasing the current threshold for spiking, and shortening ISI
between the first two spikes. However, when tested with trains of pseudo-EPSCs, I-Na$_V$R had no
effect on average spike rate across multiple spike trains (Table S6), suggesting that long term effects
of resurgent current reduce the impact of short-term changes in rate. In transient mVGNs, resurgent
current affected spike rate and regularity (CV) during pseudo-EPSC trains, independent of spike rate
and not. We attribute these effects to sensitization during the inter-spike interval of transient mVGNs
to the pseudo-randomly times EPSCs.

The isolated VGN patch preparation and model provide an opportunity to isolate the impact of the
afferents' diverse voltage-gated channels, but incorporation into a fuller model of the inner ear may
yield more insight into their impact on shaping the vestibular afferent signal. *In vivo,* many features
differ across the peripheral and central epithelial zones that give rise to regular and irregular
afferents, including the macromechanical input to hair cells, numbers and types of synaptic inputs,
relative contributions of non-quantal and quantal transmission, size of dendritic arbors, and
complements of expressed ion channels, all of which might shape voltage noise at spike initiation
zones. Our analyses suggest that diverse Na$_V$ currents flowing through one kind of pore-forming
subunit can play significant roles in differentiating regular and irregular firing in the vestibular nerve.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6   Author Contributions

SBL conceived the project, planned, and performed the electrophysiological recordings and
computational modeling, and wrote the first draft of the manuscript. RAE participated in planning,
design, analysis, and writing the manuscript.

7   Funding

This study was supported by the National Institutes of Health (R01 DC012347) awarded to RAE, and
the Howard Hughes Medical Institute Gilliam Fellowship for Advanced Study, awarded to SBL.

8   Acknowledgments

We are grateful to Dr. Omar López-Ramírez for sharing excitatory postsynaptic current data from
vestibular afferent terminals and to Dr. Radha Kalluri for access to the original VGN spiking model.

We thank Dr. Hannah Martin, Dr. Aravind Govindaraju, Dr. Kalluri, Dr. Daniel Bronson, Dr.
Nathaniel Nowak, and Katherine Regalado for comments on early versions of the manuscript.

9   References

Alexander, S. P. H., Mathie, A., Peters, J. A., Veale, E. L., Striessnig, J., Kelly, E., et al. (2021). THE
CONCISE GUIDE TO PHARMACOLOGY 2021/22: Ion channels. British Journal of Pharmacology
178, S157–S245. doi: 10.1111/bph.15539

Aman, T. K., and Raman, I. M. (2023). Resurgent current in context: Insights from the structure and
function of Na and K channels. Biophysical Journal, S0006349523041541. doi:
10.1016/j.bpj.2023.12.016

Bant, J. S., and Raman, I. M. (2010). Control of transient, resurgent, and persistent current by open-
channel block by Na channel β4 in cultured cerebellar granule neurons. Proceedings of the National
Academy of Sciences 107, 12357–12362. doi: 10.1073/pnas.1005633107

Barry, P. H. (1994). JPCalc, a software package for calculating liquid junction potential corrections
in patch-clamp, intracellular, epithelial and bilayer measurements and for correcting junction
potential measurements. Journal of Neuroscience Methods 51, 107–116. doi: 10.1016/0165-
0270(94)90031-0

Bean, B. P. (2007). The action potential in mammalian central neurons. Nature Reviews
Neuroscience 8, 451–465. doi: 10.1038/nrn2148

Brown, A. M., Schwindt, P. C., and Crill, W. E. (1994). Different voltage dependence of transient
and persistent Na+ currents is compatible with modal-gating hypothesis for sodium channels. Journal
of Neurophysiology 71, 2562–2565. doi: 10.1152/jn.1994.71.6.2562

Browne, L., Smith, K. E., and Jagger, D. J. (2017). Identification of Persistent and Resurgent Sodium
Currents in Spiral Ganglion Neurons Cultured from the Mouse Cochlea. eNeuro, ENEURO.0303-
17.2017. doi: 10.1523/ENEURO.0303-17.2017

Chabbert, C., Chambard, J.-M., Valmier, J., Sans, A., and Desmadryl, G. (1997). Voltage-activated
sodium currents in acutely isolated mouse vestibular ganglion neurones. NeuroReport 8, 1253.

Chambard, J. M., Chabbert, C., Sans, A., and Desmadryl, G. (1999). Developmental changes in low
and high voltage-activated calcium currents in acutely isolated mouse vestibular neurons. The Journal
of Physiology 518, 141–149. doi: 10.1111/j.1469-7793.1999.0141r.x

Crill, W. E. (1996). Persistent Sodium Current in Mammalian Central Neurons. Annual Review of
Physiology 58, 349–362. doi: 10.1146/annurev.ph.58.030196.002025

Cullen, K. E. (2019). Vestibular processing during natural self-motion: implications for perception
and action. Nat Rev Neurosci 20, 346–363. doi: 10.1038/s41583-019-0153-1

Denomme, N., Lukowski, A. L., Hull, J. M., Jameson, M. B., Bouza, A. A., Narayan, A. R. H., et al.
(2020). The voltage-gated sodium channel inhibitor, 4,9-anhydrotetrodotoxin, blocks human Nav1.1
in addition to Nav1.6. Neuroscience Letters 724, 134853. doi: 10.1016/j.neulet.2020.134853

Desmadryl, G., Chambard, J.-M., Valmier, J., and Sans, A. (1997). Multiple voltage-dependent
calcium currents in acutely isolated mouse vestibular neurons. Neuroscience 78, 511–522. doi:
10.1016/S0306-4522(96)00595-7

Do, M. T. H., and Bean, B. P. (2004). Sodium Currents in Subthalamic Nucleus Neurons From
Nav1.6-Null Mice. Journal of Neurophysiology 92, 726–733. doi: 10.1152/jn.00186.2004

Durlak, J. A. (2009). How to Select, Calculate, and Interpret Effect Sizes. Journal of Pediatric
Psychology 34, 917–928. doi: 10.1093/jpepsy/jsp004

Eatock, R. A., Xue, J., and Kalluri, R. (2008). Ion channels in mammalian vestibular afferents may
set regularity of firing. Journal of Experimental Biology 211, 1764–1774. doi: 10.1242/jeb.017350

Gentet, L. J., Stuart, G. J., and Clements, J. D. (2000). Direct Measurement of Specific Membrane
Capacitance in Neurons. Biophysical Journal 79, 314–320. doi: 10.1016/S0006-3495(00)76293-X

Gittis, A. H., Moghadam, S. H., and du Lac, S. (2010). Mechanisms of Sustained High Firing Rates
in Two Classes of Vestibular Nucleus Neurons: Differential Contributions of Resurgent Na, Kv3, and
BK Currents. Journal of Neurophysiology 104, 1625–1634. doi: 10.1152/jn.00378.2010

Goldberg, J. M. (2000). Afferent diversity and the organization of central vestibular pathways. Exp
Brain Res 130, 277–297. doi: 10.1007/s002210050033

Hight, A. E., and Kalluri, R. (2016). A biophysical model examining the role of low-voltage-
activated potassium currents in shaping the responses of vestibular ganglion neurons. Journal of
Neurophysiology 116, 503–521. doi: 10.1152/jn.00107.2016

Hodgkin, A. L., and Huxley, A. F. (1952). A quantitative description of membrane current and its
application to conduction and excitation in nerve. The Journal of Physiology 117, 500–544. doi:
https://doi.org/10.1113/jphysiol.1952.sp004764

Holmes, W. R., Huwe, J. A., Rowe, M. H., and Peterson, E. H. (2014). Afferent-hair cell connectivity
as a possible source of spike train irregularity in turtle vestibular bouton afferents. BMC
Neuroscience 15, P69. doi: 10.1186/1471-2202-15-S1-P69

Hong, H., Lu, T., Wang, X., Wang, Y., and Sanchez, J. T. (2018). Resurgent sodium current
promotes action potential firing in the avian auditory brainstem. The Journal of Physiology 596, 423–
443. doi: 10.1113/JP275083

Horwitz, G. C., Risner-Janiczek, J. R., and Holt, J. R. (2014). Mechanotransduction and
hyperpolarization-activated currents contribute to spontaneous activity in mouse vestibular ganglion
neurons. Journal of General Physiology 143, 481–497. doi: 10.1085/jgp.201311126

Iwasaki, S., Chihara, Y., Komuta, Y., Ito, K., and Sahara, Y. (2008). Low-Voltage-Activated
Potassium Channels Underlie the Regulation of Intrinsic Firing Properties of Rat Vestibular Ganglion
Cells. Journal of Neurophysiology 100, 2192–2204. doi: 10.1152/jn.01240.2007

Kalluri, R., Xue, J., and Eatock, R. A. (2010). Ion Channels Set Spike Timing Regularity of
Mammalian Vestibular Afferent Neurons. Journal of Neurophysiology 104, 2034–2051. doi:
10.1152/jn.00396.2010

Kay, A. R., Sugimori, M., and Llinás, R. (1998). Kinetic and Stochastic Properties of a Persistent
Sodium Current in Mature Guinea Pig Cerebellar Purkinje Cells. Journal of Neurophysiology 80,
1167–1179. doi: 10.1152/jn.1998.80.3.1167

Khaliq, Z. M., Gouwens, N. W., and Raman, I. M. (2003). The Contribution of Resurgent Sodium
Current to High-Frequency Firing in Purkinje Neurons: An Experimental and Modeling Study. J.
Neurosci. 23, 4899–4912. doi: 10.1523/JNEUROSCI.23-12-04899.2003

Kim, K. X., and Rutherford, M. A. (2016). Maturation of NaV and KV Channel Topographies in the
Auditory Nerve Spike Initiator before and after Developmental Onset of Hearing Function. J.
Neurosci. 36, 2111–2118. doi: 10.1523/JNEUROSCI.3437-15.2016

Klinger, A. B., Eberhardt, M., Link, A. S., Namer, B., Kutsche, L. K., Schuy, E. T., et al. (2012).
Sea-Anemone Toxin ATX-II Elicits A-Fiber-Dependent Pain and Enhances Resurgent and Persistent
Sodium Currents in Large Sensory Neurons. Mol Pain 8, 1744-8069-8–69. doi: 10.1186/1744-8069-
8-69

Lewis, A. H., and Raman, I. M. (2014). Resurgent current of voltage-gated Na+ channels. The
Journal of Physiology 592, 4825–4838. doi: 10.1113/jphysiol.2014.277582

Limón, A., Pérez, C., Vega, R., and Soto, E. (2005). Ca2+-Activated K+-Current Density Is
Correlated With Soma Size in Rat Vestibular-Afferent Neurons in Culture. Journal of
Neurophysiology 94, 3751–3761. doi: 10.1152/jn.00177.2005

Liu, X.-P., Wooltorton, J. R. A., Gaboyard-Niay, S., Yang, F.-C., Lysakowski, A., and Eatock, R. A.
(2016). Sodium channel diversity in the vestibular ganglion: NaV1.5, NaV1.8, and tetrodotoxin-
sensitive currents. Journal of Neurophysiology 115, 2536–2555. doi: 10.1152/jn.00902.2015

Lysakowski, A., Gaboyard-Niay, S., Calin-Jageman, I., Chatlani, S., Price, S. D., and Eatock, R. A.
(2011). Molecular Microdomains in a Sensory Terminal, the Vestibular Calyx Ending. J. Neurosci.
31, 10101–10114. doi: 10.1523/JNEUROSCI.0521-11.2011

Meredith, F. L., and Rennie, K. J. (2015). Zonal variations in K+ currents in vestibular crista calyx
terminals. Journal of Neurophysiology 113, 264–276. doi: 10.1152/jn.00399.2014

Meredith, F. L., and Rennie, K. J. (2018). Regional and Developmental Differences in Na+ Currents
in Vestibular Primary Afferent Neurons. Front. Cell. Neurosci. 12. doi: 10.3389/fncel.2018.00423

Meredith, F. L., and Rennie, K. J. (2020). Persistent and resurgent Na+ currents in vestibular calyx
afferents. Journal of Neurophysiology 124, 510–524. doi: 10.1152/jn.00124.2020

Oliveira, J. S., Redaelli, E., Zaharenko, A. J., Cassulini, R. R., Konno, K., Pimenta, D. C., et al.
(2004). Binding Specificity of Sea Anemone Toxins to Nav 1.1-1.6 Sodium Channels:
UNEXPECTED CONTRIBUTIONS FROM DIFFERENCES IN THE IV/S3-S4 OUTER LOOP *.
Journal of Biological Chemistry 279, 33323–33335. doi: 10.1074/jbc.M404344200

Ono, K., Keller, J., López Ramírez, O., González Garrido, A., Zobeiri, O. A., Chang, H. H. V., et al.
(2020). Retinoic acid degradation shapes zonal development of vestibular organs and sensitivity to
transient linear accelerations. Nat Commun 11, 63. doi: 10.1038/s41467-019-13710-4

Patel, R. R., Barbosa, C., Xiao, Y., and Cummins, T. R. (2015). Human Nav1.6 Channels Generate
Larger Resurgent Currents than Human Nav1.1 Channels, but the Navβ4 Peptide Does Not Protect
Either Isoform from Use-Dependent Reduction. PLOS ONE 10, e0133485. doi:
10.1371/journal.pone.0133485

Raman, I. M., and Bean, B. P. (1997). Resurgent Sodium Current and Action Potential Formation in
Dissociated Cerebellar Purkinje Neurons. J. Neurosci. 17, 4517–4526. doi: 10.1523/JNEUROSCI.17-
12-04517.1997

Raman, I. M., and Bean, B. P. (2001). Inactivation and Recovery of Sodium Currents in Cerebellar
Purkinje Neurons: Evidence for Two Mechanisms. Biophysical Journal 80, 729–737. doi:
10.1016/S0006-3495(01)76052-3

Raman, I. M., Sprunger, L. K., Meisler, M. H., and Bean, B. P. (1997). Altered subthreshold sodium
currents and disrupted firing patterns in Purkinje neurons of Scn8a mutant mice. Neuron 19, 881–
891. doi: 10.1016/s0896-6273(00)80969-1

Renganathan, M., Dib-Hajj, S., and Waxman, S. G. (2002). Nav1.5 underlies the 'third TTX-R
sodium current' in rat small DRG neurons. Molecular Brain Research 106, 70–82. doi:
10.1016/S0169-328X(02)00411-4

Risner, J. R., and Holt, J. R. (2006). Heterogeneous Potassium Conductances Contribute to the
Diverse Firing Properties of Postnatal Mouse Vestibular Ganglion Neurons. J Neurophysiol 96,
2364–2376. doi: 10.1152/jn.00523.2006

Rosker, C., Lohberger, B., Hofer, D., Steinecker, B., Quasthoff, S., and Schreibmayer, W. (2007).
The TTX metabolite 4,9-anhydro-TTX is a highly specific blocker of the Nav1.6 voltage-dependent
sodium channel. American Journal of Physiology-Cell Physiology 293, C783–C789. doi:
10.1152/ajpcell.00070.2007

Rothman, J. S., and Manis, P. B. (2003). The Roles Potassium Currents Play in Regulating the
Electrical Activity of Ventral Cochlear Nucleus Neurons. Journal of Neurophysiology 89, 3097–
3113. doi: 10.1152/jn.00127.2002

Songer, J. E., and Eatock, R. A. (2013). Tuning and Timing in Mammalian Type I Hair Cells and
Calyceal Synapses. J. Neurosci. 33, 3706–3724. doi: 10.1523/JNEUROSCI.4067-12.2013

Stafstrom, C. E., Schwindt, P. C., and Crill, W. E. (1982). Negative slope conductance due to a
persistent subthreshold sodium current in cat neocortical neurons in vitro. Brain Research 236, 221–
226. doi: 10.1016/0006-8993(82)90050-6

Taddese, A., and Bean, B. P. (2002). Subthreshold Sodium Current from Rapidly Inactivating
Sodium Channels Drives Spontaneous Firing of Tuberomammillary Neurons. Neuron 33, 587–600.
doi: 10.1016/S0896-6273(02)00574-3

Ventura, C. M., and Kalluri, R. (2019). Enhanced Activation of HCN Channels Reduces Excitability
and Spike-Timing Regularity in Maturing Vestibular Afferent Neurons. J. Neurosci. 39, 2860–2876.
doi: 10.1523/JNEUROSCI.1811-18.2019

Venugopal, S., Seki, S., Terman, D. H., Pantazis, A., Olcese, R., Wiedau-Pazos, M., et al. (2019).
Resurgent Na+ Current Offers Noise Modulation in Bursting Neurons. PLOS Computational Biology
15, e1007154. doi: 10.1371/journal.pcbi.1007154

Wanke, E., Zaharenko, A. J., Redaelli, E., and Schiavon, E. (2009). Actions of sea anemone type 1
neurotoxins on voltage-gated sodium channel isoforms. Toxicon 54, 1102–1111. doi:
10.1016/j.toxicon.2009.04.018

White, H. V., Brown, S. T., Bozza, T. C., and Raman, I. M. (2019). Effects of FGF14 and NaVβ4
deletion on transient and resurgent Na current in cerebellar Purkinje neurons. Journal of General
Physiology 151, 1300–1318. doi: 10.1085/jgp.201912390

Wu, N., Enomoto, A., Tanaka, S., Hsiao, C.-F., Nykamp, D. Q., Izhikevich, E., et al. (2005).
Persistent Sodium Currents in Mesencephalic V Neurons Participate in Burst Generation and Control
of Membrane Excitability. Journal of Neurophysiology 93, 2710–2722. doi: 10.1152/jn.00636.2004

10   Data Availability Statement

The associated code is accessible in the following repository: https://github.com/eatocklab/NaV-
currents-in-VGN-spiking.

The datasets generated and analyzed for both electrophysiology and modeling experiments can be
found in our Dryad repository:
https://datadryad.org/stash/share/sbukzLlDkLPlsuTAISqONfWVmLOuSwvnNNXr5LKdD6c
(Reviewer URL).

11   Figure Legends

Figure 1 *VGNs expressed NaVT current, many expressed NaVP current, and a few expressed NaVR*
currents.

(A) VGNs are the isolated cell bodies of bipolar vestibular afferents (orange) synapsing on a hair cell
(blue). *In vivo*, APs initiate at a heminode close to the synapse and adjacent to the first myelinated
internode and propagate through the myelinated cell body, toward the brain. **(B)** Transient sodium
current (I-NaVT) in a P13 VGN, evoked by stepping up from a prepulse of –125 mV in 5 mV
increments. I-NaVT was always fully blocked by 1 μM TTX. *Inset:* Diagram of patching an isolated,
demyelinated VGN. **(C)** A small, non-inactivating (persistent) current (I-NaVP) was isolated from a
P3 VGN by applying a 0.1 mV/ms voltage ramp (–80 mV to +60 mV) and subtracting current in 1
μM TTX from control current. **(D)** Resurgent sodium current (I-NaVR) in a P18 VGN, evoked by
applying a +25 mV prepulse followed by repolarizing steps to –25 to –80 mV; and subtracting
current in 1 μM TTX from control current. **(E)** I-V curves compared for NaVT, NaVP and NaVR
currents. Solid lines (means) plus shading (± SEM) are averaged from recordings; dashed lines show
modeling of NaVT, NaVP and NaVR currents (see Methods).

Figure 2 *Voltage dependence of total NaV current varied with different combinations of NaVT, NaVP*
and/or NaVR current modes.

(A) I-V curves averaged for cells with different current modes. 2 VGNs with I-NaVT plus I-NaVR
(green) had large current densities and are shown individually. **(B)** G-V activation curves show that
voltage dependence differed in cells with different combinations of NaV current modes. $V_{1/2}$ of

activation and inactivation are marked with square symbols. Voltage dependence of activation was
more negative in VGNs with I-Na$_V$R (I-NaV(T+R) and I-NaV(T+P+R); green and pink lines) than I-
Na$_V$(T+P) (p = 0.02, blue) and I-Na$_V$T (p = 0.007, black). V$_{1/2}$ of inactivation did not differ (one-way
3-factor ANOVA, p = 0.7). **(C)** VGNs with I-Na$_V$R, I-Na$_V$(T+P+R) (pink) and I-Na$_V$(T+R) (green),
had larger (pooled) conductance densities than VGNs without I-Na$_V$R: I-Na$_V$(T+P) (blue; one-way 3-
factor ANOVA, p = 0.02) and I-Na$_V$T (black; p = 0.002).

Figure 3 *Na$_V$1.6-selective channel blocker (4,9-ah-TTX) reveals strong Na$_V$1.6 contribution to Na$_V$T,*
Na$_V$P, and Na$_V$R current modes.

(A) I-Na$_V$T (P6 VGN) by 100 nM 4,9-ah-TTX is blocked by ~70% (n = 13). Inset highlights block of
Na$_V$P current during voltage step. Control current is in black, red is the current measured during the
drug (i.e., the "residual" current), and the blue is the blocked current (red current subtracted from the
black). **(B)** Boltzmann fits of G-V activation and inactivation curves for data from (A). Square
symbols indicate V$_{1/2}$ values. 4,9-ah-TTX-insensitive current (red) had more negative inactivation
V$_{1/2}$ (n = 12, p = 0.02, Table 4). **(C & D)** Na$_V$P (C, P17 VGN) and Na$_V$R (D, P18 VGN) currents are
also blocked (~90%) by 100 nM 4,9-ah-TTX.

Figure 4 *Na$_V$ channel agonist (ATX-II) strongly enhanced Na$_V$ T and Na$_V$ P currents.*

(A) An example of enhanced Na$_V$T and Na$_V$P current by 100 nM ATX-II (red trace) (top panel) as
elicited by depolarizing voltage steps (bottom panel). **(B)** I-V relation of Na$_V$P current in a P7 VGN,
before (grey) and in 100 nM ATX-II (red). Voltage dependence of I-Na$_V$P was not significantly
altered by ATX-II: V$_{1/2}$ of G-V curves –38.8 ± 4.2 mV (n = 7) vs. –30.6 ± 2.9 mV, p = 0.12, power =
0.33 (not shown). **(C)** Na$_V$P currents grew in ATX-II from 27.4 ± 14.4 pA to 163.5 ± 45.9 pA (n = 7,
paired t-test; p = 0.01; Hedges' g = 0.93, large effect). Na$_V$R currents were not tested.

Figure 5 *Correlating VGN firing patterns with spike waveforms and maximum Na$_V$ conductance*
density.

(A) Exemplar firing patterns in VGNs evoked by a 500-ms current steps of different size reflecting
different current thresholds. **(B)** Variation in maximum Na$_V$ conductance density with firing pattern
is not significant (homogeneous variance, Levene's Test). **C:** Average spikes for each firing pattern,
aligned to peak; spike height (ΔV$_{AP}$) and afterhyperpolarization (AHP) are measured from V$_{rest}$. **(D)**
AP height correlated weakly with Na$_V$ G$_{max}$ density. **(E)** AHP depth varied with firing pattern (****,
p = 0.00007; ***, p = 0.0005; **, p = 0.009; *, p = 0.04). **(F)** Phase plane plots of averaged APs from
C. Squares denote peak dV/dt (rate of spike rise) values. **(G)** Peak dV/dt (see squares in F) correlated
weakly with Na$_V$ G$_{Max}$ density. **(H)** No relationship between Na G$_{max}$ Density and age.

Figure 6 *4,9-ah-TTX reduced step-evoked spiking excitability and altered the AP waveform.*

In 100 nM 4,9-ah-TTX, the 50% block of I-Na$_V$T was similar in transient VGNs (n = 8) and
sustained VGNs (n = 5: 1 sustained-A, 1 sustained-B, 3 sustained-C) (Welch's t-test, p = 0.94, power
= 0.05). **(A & B)** In 100 nm 4,9-ah-TTX (red), current threshold increased in a sustained VGN (A)
and a transient VGN (B). In 7 cells, threshold increased from 92.3 ± 14.8 pA to 165.4 ± 23.6 pA
(paired t-test, p = 0.002, Hedges' g = 0.63; medium effect). **(C)** First AP from firing patterns such as
(A) and (B), averaged (n = 7). **(D)** In 4,9-ah-TTX, for sustained (squares) and transient (circles)
neurons, spike height decreased (p = 0.01), **(E)** afterhyperpolarization depth increased (p = 0.006),
(F) and spike latency (i.e., time-to-peak) increased (p = 0.05). In some cases, AHP was depolarized
relative to (V$_{rest}$) by the current step. **(G)** Phase plane plots of mean AP waveforms in (C) highlight

the change in peak dV/dt and V_{AP}. (**H**) Peak dV/dt, from (E), was decreased by 4,9-ah-TTX (p = 0.03, Hedges' g = 0.14, small effect). (**I & J**) Sustained VGNs (I) were significantly hyperpolarized (p = 0.03, Hedges' g = 0.67, medium-large effect) but not transient VGNs (J) (p = 0.16). (**K**) 100 nm 4,9-ah-TTX (*red middle panel*) did not significantly reduce CV in spike trains evoked by pseudo-EPSCs (bottom panel) relative to control conditions (*black top panel*) in a P5 transient VGN. Dashed lines at –10 mV indicate event threshold for spike count; dashed lines at –60 mV indicate V_{rest}. *Inset:* first 50 ms of EPSC train. (**L**) CV was not significantly altered in 4,9-ah-TX (paired t-test, 0.85 ± 0.09 vs 0.94 ± 0.07, p = 0.15, power = 0.29). Sustained VGNs (n = 4, *squares*); transient VGNs (n = 6, *circles*).

Figure 7 *ATX-II increased spike rate and rate-independent regularity for current steps and EPSC-train stimuli.*

(**A**) 100 nm ATX-II (*red*) increased excitability relative to control (*black*) in a P12 sustained-C VGN: increasing the number of APs per step and inducing spontaneous spiking at rest (step current threshold = 0). (**B**) In a P12 transient VGN, current threshold was reduced, but the number of APs remained unchanged. (**C**) Pseudo-EPSC trains (*bottom panel*) evoked spikes in the sustained-C VGN from A. ATX-II increased spike timing regularity in all VGNs. Inset shows the first 50 ms of EPSC train. To keep spike rate constant between conditions at ~38 spikes/s, smaller EPSCs were applied during ATX-II (*bottom panel*). (**D**) ATX-II decreased CV in every VGN tested (transient = open circles, sustained = filled squares) (0.70 ± 0.06 to 0.34 ± 0.08, paired t-test; n = 7; p = 0.003).

Figure 8 *Adding I-Na$_V$P altered AP waveform and ISI in current-step responses of model VGNs.*

Step-evoked firing in model VGNs with only I-Na$_V$T (T); added I-Na$_V$R (T+R) (10% of I-Na$_V$T); added I-Na$_V$P (T+P) (3% of I-Na$_V$T); both added (T+P+R); and increased I-Na$_V$T (T+). 1st 50 ms of responses to current step. Current steps, below, were increased (left to right) to account for increased current thresholds for spiking as firing patterns progressed from sustained-A to transient. Rows B-D show currents corresponding to the above firing pattern and isolated for each current mode (T, R, P). (**A**) In sustained-A, -B, and –C models, adding I-Na$_V$P had variable effects on refractory periods and therefore AP time-to-peak in sustained-A, -B, -C mVGNs, and increased the size of post-spike voltage oscillations in sustained-B and sustained-C mVGNs (blue arrows). Adding I-Na$_V$P depolarized Vrest in sustained-A mVGN (blue arrow), and slightly decreased refractory periods (time-to-peak) in all sustained mVGNs. (**B**) I-Na$_V$T current flow during AP train. To control for increased Na$_V$ conductance with added P and/or R conductance, we also ran a simulation with Na$_V$T conductance density increased by the same amount (18.1, 14.7, 12.4, 7.9 mS/cm^2 in sustained-A, -B, -C, transient simulations, respectively; T+ (orange) traces). (**C**) I-Na$_V$R current flow during APs; note that I-Na$_V$R flows during repolarization of each AP and decreases in amplitude with each successive AP in sustained-A model neuron. (**D**) I-Na$_V$P current flow during AP train; note small variations in peak I-Na$_V$P and the increase in amplitude with each AP in sustained mVGNs. (**E**) Temporally aligning the first APs from trains in (A) shows I-Na$_V$P decreased spike height and I-Na$_V$R decreases AHP (green arrows) in sustained-A, and increased Na$_V$T increases spike height (orange arrows) in all mVGNs. (**F**) Phase plane plots show that increasing Na$_V$T also increases peak dV/dt in all mVGNs and I-Na$_V$P decreases peak dV/dt in sustained-S mVGN.

Figure 9 *Adding I-Na$_V$P and/or I-Na$_V$R to I-Na$_V$T differentially affected spike rate and regularity in model VGNs.*

(A) Sustained-A mVGN under different Na_V current combinations: adding I-Na_VP increased spike rate, whether I-Na_VR was also added or not. (B) Transient mVGN under different Na_V current combinations: adding I-Na_VR increased spike rate, but sometimes caused depolarization block (green arrow). *Bottom panel,* pseudo-EPSC train used to evoke spike trains shown. (C & D) Summary of spike rates for each combination of Na_V current modes at each level of g_{NaVT}. Increasing Na_VT increased spike rate in both sustained-A and transient mVGNs. *C,* Adding I-Na_VP (T+P and T+P+R) increased spike rate relative to Na_VT in sustained-A mVGN, while adding R has little effect on spike rate except at the highest value of (T+P+R). *D*, I-Na_VR (T+R and T+P+R) increased spike rate in transient VGN, while I-Na_VP had no effect. (E & F) Spike train regularity (CV) for spike trains in C & D. I-Na_VP had little effect on CV in either mVGN. I-Na_VR increased CV especially in the transient mVGN, but also in sustained-A mVGN for $g_{NaVT} > 14$ mS/cm^2. Two-way ANOVAs confirmed interactions between effects of g_{NaVT} and Na_V current modes on spike rate and CV in sustained-A and on CV in transient mVGNs (Spike rate: $F_{(8,24)} = 3$ and 0.8, p <0.0001 and 0.67. CV: $F_{(8,24)} = 2$ and 3.6 and p = 0.009 and <0.001). (G & H) CVs with spike rate held at ~20 spikes/s ($CV_{20sp/s}$) across conditions, used to assess the effects of Na_V currents on CV independent of spike rate. *G,* Adding I-Na_VP to sustained-A mVGN significantly decreased $CV_{20sp/s}$ in sustained-A mVGN, with or without I-Na_VR. *H*, Adding I-Na_VR to transient mVGN significantly increased $CV_{20sp/s}$ in sustained-A mVGN for $g_{NaVT} > 12$ mS/cm^2, with or without I-Na_VP.

12   Tables

Table 1 *Whole-cell recording solutions.* pH was adjusted with CsOH for Na$^+$ reduced external and Cs internal solutions, NaOH for standard external solution, and KOH for standard internal solution.

Table 1: Whole-cell recording solutions

External

in mM	NaCl	KCl	CsCl	CaCl$_2$	MgCl$_2$	TEACl	NaH$_2$PO$_4$	D-Glucose		HEPES	pH	Osm (mmol/kg)
Na$^+$ Reduced	75	0	5.4	0	2.5	75	0	10		5	7.4	310
Standard	144	5.1	0	1.3	0.9	0	0.7	5.6		10	7.4	310

Internal

in mM	KCl	CsCl	CaCl$_2$	MgCl$_2$	Na$_2$ creatine phosphate	Mg-ATP	Li-GTP	Na-cAMP	EGTA	HEPES	pH	Osm (mmol/kg)
Cs$^+$	0	148	0.8	0	3.5	3.5	0.1	0.1	5	5	7.4	300
Standard	135	0	0.1	0.5	5	3	0.1	0.1	5	5	7.25	285

Table 2 *Conductance parameters used for spiking by model VGNs.* Sources: [a]Hight and Kalluri (2016), [b]Kalluri et al. (2010), [c]Iwasaki et al. (2008), [d]Ventura and Kalluri (2019). g_{NaVT} and V_{rest} from this study.

Table 2: Conductance parameters used for spiking

Firing Pattern	$\bar{g}_{NaVT}$, (mS/cm^2)	$\bar{g}_{KLV}$ [a,b,d]	$\bar{g}_{KH}$ [a,c,d]	$\bar{g}_H$ [d]	$\bar{g}_{leak}$ [a,d]	V_{rest} (mV)
Sustained-A	16	0	4.5	0.2	0.02	-60.1
Sustained-B	13	0.2	4	0.5	0.05	-63.5
Sustained-C	11	0.5	4	0.1	0.05	-64.1
Transient	7	1.2	2.5	0.9	0.1	-65.7

1160 **Table 3** *Steady-state activation and inactivation parameters used in modeling Na$_V$ current modes.*

Table 3: Steady-state parameters for Na$_V$ current modes		Activation		Inactivation	
	$\bar{g}$ (mS/cm^2)	V$_{1/2}$ (mV)	S (mV)	V$_{1/2}$, (mV)	S (mV)
I-Na$_V$T	6 – 22	-36	6	-68	8
I-Na$_V$P	0.18 – 0.44	-27	10	-52	14
I-Na$_V$R	0.6 – 2.2	-40	22	-40	28

1161

1162 **Table 4** *Electrophysiological properties of Na$_V$ currents in VGNs.* Asterisks indicate significance in
1163 one or two comparisons. See Table S1 for full summary of statistical analyses.

Table 4: Electrophysiological properties of Na$_V$ currents in VGNs		Activation		Inactivation		
		V$_{1/2}$ (mV)	S (mV)	V$_{1/2}$ (mV)	S (mV)	G density (nS/pF)
Na$_V$ current	I-Na$_V$ (Na+ Red Ext Sol'n)	-35.8 ± 0.7 (n = 75)	5.7 ± 0.2 (76)	-68.1 ± 1.1 (73)	8.4 ± 0.5 (74)	3.8 ± 0.3 (74)
	I-Na$_V$ (Standard Ext Sol'n)	-38.9 ± 1.2 (29)	3.2 ± 0.3 (29)	-59.1 ± 1.6 (23)	4.4 ± 0.8 (23)	6.8 ± 0.7 (29)
	I-Na$_V$ *(Liu et al., 2016, rat VGN)*	*-36.5 ±1.6 (11)*	*5.7 ±0.4 (11)*	*-76.3 ±0.2 (12)*	*7.6 ±0.1 (12)*	*8.7 ±1.0 (12)*
I-Na$_V$ modes (Figure 2)	I-Na$_V$ (T)	-35.6 ± 1.0 (10)	6.6 ± 0.3 (10)	-70.1 ± 2.1 (10)	12.3 ± 1.3 (10)	2.9 ± 0.3 (31)
	I-Na$_V$ (T+P)	-37.7 ± 1.6 (12)	7.1 ± 0.4 (12)	-66.4 ± 2.3 (12)	12.8 ± 0.8 (12)	3.6 ± 0.4 (34)
	I-Na$_V$ (T+P+R)	-48.6 ± 5.7* (6)	6.0 ± 0.4 (6)	-67.5 ± 6.3 (6)	11.5 ±1.0 (6)	5.9 ± 0.9* (6)
4,9-ah-TTX (Figure 3)	Control (I-Na$_V$)	-38.7 ± 1.3 (12)	5.7 ± 0.5 (12)	-68.3 ± 2.6 (12)	9.1 ± 1.4 (12)	3.1 ± 0.4 (13)
	Subtraction	-39.4 ± 1.6	4.7 ± 0.6	-68.4 ± 2.4	8.8 ± 1.3	2.4 ± 0.3
	Residual	-42.3 ± 2.3	9.2 ± 2.1	-80.6 ± 3.8*	14.5 ± 4.9	0.9 ± 0.3*
ATX-II (Figure 4)	Control (I-Na$_V$)	-37.7 ± 1.2 (7)	5.7 ± 0.3 (7)	-76.7 ± 3.6 (7)	11.2 ± 0.8 (7)	3.7 ± 1.7 (7)
	ATX-II	-38.7 ± 1.3	5.8 ± 0.1	-80.9 ± 2.9	16.3 ± .3*	1.9 ± 0.4

1164

1165 **Table 5** *AP waveform differences between firing patterns from Figure 5.*

Table 5: AP waveform difference between firing patterns (one-way 4-factor ANOVA)						
Firing pattern (n)	Time-to-peak (ms)	AHP (mV)	V$_{AP}$ (mV)	Voltage threshold (mV)	Peak dV/dt (mV/ms)	Na$_V$ G$_{max}$ Den (nS/pF)
Sustained-A (6)	4.5 ± 0.4	11.0 ± 2	110 ± 6.2	-41.0 ± 3.3	208.3 ± 52.2	9.6 ± 0.7
Sustained-B (13)	4.7 ± 0.3	6.0 ± 0.8	83.5 ± 6.4	-44.3 ± 1.1	166.3 ± 34.2	8.0 ± 1.4
Sustained-C (10)	4.5 ± 0.3	4.1 ± 0.9	80.0 ± 7.7	-44.0 ± 1.3	123.7 ± 28.3	7.0 ± 1.2
Transient (28)	4.2 ± 0.2	2.9 ± 0.6	75.1 ± 4.0	-42.9 ± 1.3	78.6 ± 12.5	7.5 ± 0.9
p	0.45	0.00001	0.009	0.16	0.005	0.68
power	0.23	0.99	0.83	0.16	0.88	0.14

13   Supplementary Material

13.1 Supplementary Figures

Figure S1 *Comparison of VGN responses to model VGN simulations.*

(A – C) (reproduced from Fig. 5). Averaged recorded APs (A), corresponding mean phase plane
plots (B), and exemplar step-evoked firing patterns (C). **(D)** Model-generated APs based on APs
from (A). **(E)** Phase-plane plots for simulated APs from (D)**.** Simulated APs have smaller peak rates
of depolarization. **(F)** Step-evoked firing patterns of model VGNs capture key features of exemplar
data in (C).

1174

1175 **Figure S2** *Shape of simulated EPSCs.*

1176 Simulated excitatory postsynaptic currents (EPSCs) used to evoke spiking in our cell-autonomous
1177 model were modeled after spontaneous vestibular synaptic potentials (s_V) recorded at room
1178 temperature in mouse calyces. EPSC shapes previously tested in Hight and Kalluri (2016) (s1, s2,
1179 and s3) are shown for comparison.

1180

1181 **Figure S3** *Changes in firing patterns and Na$_V$ conductance density with age.*

1182 Change in distribution of firing pattern in younger and older VGNs.

1183

Figure S4 *Reducing Na$_V$ current modes replicates 4,9-ah-TTX block in model sustained and*
transient VGNs.

(A & B) Firing patterns for a sustained-A (A) and transient (B) mVGNs. Reducing (red) I-Na$_V$T by
70% and I-Na$_V$R and I-Na$_V$P by 90% replicates results observed with application of 4,9-ah-TTX in
sustained-A mVGNs: increase in current threshold, reduction in spike height and spike rate, and
slight hyperpolarization in V$_{rest}$. **(B)** Transient mVGNs had reduced spike height and (not shown)
greater current threshold. **(C)** Na$_V$P and Na$_V$T drive hyperpolarization in sustained mVGNs. Each
component was reduced by the estimated 4,9-ah-TTX block: Na$_V$R and Na$_V$P by 90% and Na$_V$T by
70%. Reducing ("blocking") components in various combinations shows that blocking Na$_V$T and
Na$_V$P hyperpolarizes resting membrane potential.

1194

Figure S5 *Increasing I-Na$_V$P further increases spike rate*

(A & B) Firing patterns for a sustained-A (A) and transient (B) mVGNs with I-Na$_V$T density of 16
mS/cm^2 with different amounts of I-Na$_V$P: 0% (control, black), 3% (red), 5% (blue), 8% (green) and
10% (purple). **(C & D)** Summary of spike rates for sustained-A and transient mVGN under each
level of I-Na$_V$P. Increasing I-Na$_V$P increased spike rate in both mVGNs. **(E & F)** A summary of
spike train regularity as measured by CV. Increased I-Na$_V$P had no effect on CV in either mVGN.

1201

1202

1203

1204 ### 13.2 Supplementary Tables

Table S1: Summary of one-way ANOVA tests for voltage clamp experiments

		F(2,70) = 6.7, p = 0.002, power = 0.90	
		p	effect size, interpretation
Na$_V$ conductance density (Fig 2B)	Na$_V$T vs Na$_V$T+P	0.33	
	Na$_V$T vs Na$_V$T+P+R	0.002	0.27, small
	Na$_V$T+P vs Na$_V$T+P+R	0.02	0.41, med
		Activation, V$_{1/2}$		**Inactivation, V$_{1/2}$**	
		F(2,27) = 5.9, p = 0.008, power = 0.83		F(2,26) = 0.38, p = 0.68, power = 0.10	
Na$_V$ current modes (Fig 2C)	Bonferroni test:	p	effect size	p	
	Na$_V$T vs Na$_V$T+P	1		0.66	
	Na$_V$T vs Na$_V$T+P+R	0.009	0.75, big	0.86	
	Na$_V$T+P vs Na$_V$T+P+R	0.02	0.59, med	0.97	
4,9-ah-TTX (Fig 3B)	(n = 12)	F(2, 35) = 1.1, p = 0.35, power = 0.22		F(2, 35) = 5.5, p = 0.009, power = 0.82	
	Bonferroni test:	p		p	effect size

		Activation, $V_{1/2}$		Inactivation, $V_{1/2}$	
	Control vs Residual	0.52	0.02	0.69, med	
	Control vs Blocked	1	1		
	Residual vs Blocked	0.83	0.02	0.66, med	
ATX-II (Fig 4)	(n = 7)	**Activation, $V_{1/2}$**		**Inactivation, $V_{1/2}$**	
	Paired t-test:	p	power	p	power
	Na_VT: Control vs ATX-II	0.34	0.13	0.06	0.48
	Na_VP: Control vs ATX-II	0.12	0.33	-	-

1205

Table S2: Firing pattern passive properties (one-way 4-factor ANOVA)				
Firing pattern	Current threshold (pA) (n)	V_{rest} (mV)	R_{in} (MΩ)	Cm (pF)
Sustained-A	133.3 ± 16.7 (3)	-60.0 ± 2.2 (6)	1020.0 ± 247.8 (5)	21.9 ± 2.3 (6)
Sustained-B	261.1 ± 26.1 (9)	-64.1 ± 1.8 (13)	852.2 ± 280.6 (9)	16.9 ±1.8 (12)
Sustained-C	295.5 ± 34.7 (11)	-63.5 ± 1.3 (10)	475.3 ± 128.8 (7)	15.1 ± 2.1 (9)
Transient	426.8 ± 33.9 (28)	-65.7 ± 0.9 (33)	464.9 ± 73.8 (20)	16.1 ± 1.1 (31)
p	0.001	0.11	0.08	0.18
power	0.94	0.51	0.56	0.41

1206

Table S3: Direct comparisons of AP waveform differences (one-way 4-factor ANOVA)								
	Current threshold (pA)		Spike height (V_{AP}) (mV)		Peak dV/dt (mV/ms)		AHP (mV)	
	p	Effect size	p	Effect size	p	Effect size	p	Effect size
Sustained-B vs Sustained-A	0.35		0.08		0.79		**0.009**	0.6
Sustained-C vs Sustained-A	0.58		0.05		0.31		**0.0005**	0.9
Sustained-C vs Sustained-B	0.95		0.98		0.71		0.46	
Transient vs Sustained-A	**0.01**	0.4	**0.004**	**0.6**	**0.01**	0.5	**0.000006**	1.0
Transient vs Sustained-B	**0.03**	0.4	0.66		**0.03**	0.3	**0.04**	0.5
Transient vs Sustained-C	0.08		0.92		0.58		0.77	

1207   c

Table S4: Pharmacological effects on AP waveform (paired t-test)								
4,9-ah-TTX	V_{rest} (mV) Transient	V_{rest} (mV) Sustained	Time-to-peak (ms)	Spike Width (ms)	AHP (mV)	V_{AP} (mV)	Voltage threshold (mV)	Peak dV/dt (mV/ms)
Control (n = 12)	-59.5 ± 1.5	-57.6 ± 3.3	4.1 ± 0.3	2.9 ± 0.5	0.0 ± 1.9	74.4 ± 6.4	-42.3 ± 2.3	102 ± 20
Drug	-63.6 ± 2.9	-66.8 ± 3.4	4.8 ± 0.4	3.9 ± 0.6	6.8 ± 1.2	55.1 ± 7.0	-43.3 ± 3.2	64.4 ± 20

p	0.16	**0.03**	**0.05**	0.13	**0.006**	**0.01**	0.90	**0.03**
Effect size		0.67	0.42	0.19	0.72	0.49	0.05	0.41
Power	0.27	0.67	0.53	0.31	0.86	0.76	0.05	0.62
ATX-II	V_{rest} (mV)							
Control (n = 7)	-59.2 ± 0.9		3.8 ± 0.3	1.2 ± 0.1	4.8 ± 0.8	92.7 ± 5.9	-49.7 ± 2.1	229 ± 30
Drug	-64.0 ± 2.2		5.5 ± 0.8	2.4 ± 0.6	4.1 ± 1.5	92.8 ± 6.3	-52.2 ± 3.5	209 ± 50
p	0.07		0.07	0.09	0.58	0.97	0.60	0.41
Effect size	0.83		0.45	0.35	0.10	0.01	0.23	0.14
Power	0.45		0.45	0.41	0.07	0.05	0.07	0.11

1208

Table S6: Effects of I-Na$_V$P and I-Na$_V$R on spike rate and CV in model VGNs (two-way 4-factor ANOVA)									
		Spike Rate (sp/s)			**CV**			**CV$_{20sp/s}$**	
			vs. Na$_V$T			vs. Na$_V$T			vs. Na$_V$T
		μ ± SEM	p	g	μ ± SEM	p	μ ± SEM	p
Sustained-A	Na$_V$T	16 ± 0.9	-		0.42 ± 0.03	-	0.45 ± 0.03	-
	Na$_V$T+R	16 ± 1.1	1		0.45 ± 0.03	1	0.45 ± 0.03	1
	Na$_V$T+P	23 ± 1.2	**<0.0001**	**0.6**	0.36 ± 0.03	0.08	0.35 ± 0.03	<0.0001
	Na$_V$T+R+P	24 ± 1.5	**<0.0001**	**0.6**	0.45 ± 0.03	0.96	0.32 ± 0.03	<0.0001
Transient	Na$_V$T	15 ± 0.7	-		0.60 ± 0.02	-	0.60 ± 0.02	-
	Na$_V$T+R	16 ± 0.9	0.06		0.79 ± 0.03	<0.0001	0.81 ± 0.03	<0.0001
	Na$_V$T+P	16 ± 0.8	0.2		0.57 ± 0.02	1	0.59 ± 0.02	1
	Na$_V$T+R+P	18 ± 1.0	<0.0001	0.2	0.79 ± 0.03	<0.0001	0.80 ± 0.03	<0.0001

1209

1210

Table S5: AP waveform differences between model VGNs											
mVGN (Figure 8)			1st AP:					ISI (ms)	2nd AP:		
	Nav modes	Vrest (mV)	Time-to-peak (ms)	V_{AP} (mV)	Spike width (ms)	Peak dV/dt (mV/ms)	AHP (mV)		Time-to-peak (ms)	V_{AP} (mV)	Spike width (ms)
Sustained-A	T	-59.6	2.6	96	1	143.4	-6.9	8.6	6.4	69.2	1.4
	T+P	-54.7	2.3	77.9	1.2	100.6	-9.6	8.3	6	61.8	1.4
	T+R	-59.6	2.6	96	1	143.4	-3.3	7.2	5.1	63.7	1.6
	T+P+R	-54.7	2.3	77.9	1.2	100.6	-9.4	8.2	5.9	61.8	1.5
	T+	-59.6	2.5	101.6	0.9	162.4	-7.3	8.6	6.2	74	1.3
Sustained-B	T	-61.7	3.9	85.9	1.2	105.1	-5.6	12.9	10.3	57	2.3
	T+P	-61.1	3.8	84.7	1.2	106	-6.1	12.4	9.8	58.8	1.8
	T+R	-61.7	3.9	85.9	1.2	105.1	-5.4	12.8	10.2	57.1	2.3
	T+P+R	-61.1	3.8	84.7	1.1	106	-6	12.3	9.7	58.9	2
	T+	-62.7	3.8	92.7	1.1	123.1	-5.3	12.7	10.2	63.7	1.9
Sustained-C	T	-65.2	5.1	86.6	1.4	92.8	-0.8	16.6	13.5	48.5	3.4
	T+P	-64.7	4.9	86.5	1.5	94.2	-1.2	15.9	12.7	51.2	3.2
	T+R	-65.2	5.1	86.6	1.4	92.8	-0.7	16.5	13.4	48.8	3.8
	T+P+R	-64.7	4.9	86.4	1.5	94.2	-1.2	15.7	12.1	51.5	3.2
	T+	-65.2	5	91.9	1.5	108.7	-1.5	16.1	12.9	53.7	2.8
Transient	T	-64.3	2.5	77.6	1.5	77.9	5.4				
	T+P	-64.2	2.5	77.9	1.5	79.7	5.3				
	T+R	-64.2	2.5	77.9	1.5	77.9	5.3				
	T+P+R	-64.2	2.5	77.9	1.5	79.7	5.3				
	T+	-64.3	2.5	82	1.4	90.9	4.7				

1211

A

B I-Na$_V$T

+ 1 µM TTX

2 nA
1 ms

-25

-125 mV

C I-Na$_V$P

+ 1 µM TTX

Control

50 pA

400 ms

-80 mV   60   0.1 mV/ms

D I-Na$_V$R

100 pA
10 ms

25   -45

-135 mV

E I-V Curves

0   0

-2

-4

-80   -40   0   40
V (mV)

I-Na$_V$T (nA)

I-Na$_V$P & I-Na$_V$R (pA)

-80

-160

Model
Ephys