1

Comparing the predictors of mutability among healthy human tissues inferred from mutations in single cell genome data

Madeleine Oman[1,2,*], and Rob W. Ness[1,2,3]

[1] Dept of Ecology and Evolutionary Biology, University of Toronto, Toronto, Canada
[2] Dept of Biology, University of Toronto, Mississauga, Canada
[3] Dept of Cell and Systems Biology, University of Toronto, Toronto, Canada
[*] Corresponding Author email: madeleine.oman@mail.utoronto.ca

Article type: Research

Running Title: Mutability across the human body

Competing Interest Statement: The authors declare no conflict of interest.

Abstract

Studying mutation in healthy somatic tissues is key for understanding the genesis of cancer and other genetic diseases. Mutation rate varies from site to site in the human genome by up to 100-fold and is influenced by numerous epigenetic and genetic factors including GC content, trinucleotide sequence context, DNAse accessibility, and histone modifications. These factors influence mutation at both local and regional scales and are often interrelated with one another, meaning that predicting mutability or uncovering its drivers requires modelling multiple factors and scales simultaneously. Historically, most investigations have focused either on analyzing the local sequence scale through triplet signatures or on examining the impact of epigenetic processes at larger scales, but not both concurrently. Additionally, sequencing technology limitations have restricted analyses to mutations only in coding regions (RNA-seq) or to those that have been influenced by selection (bulk tissue). Here we present a comprehensive analysis of epigenetic and genetic factors at multiple scales in the germline and three somatic tissues. We leverage publicly available data for 21 genomic predictors, and somatic mutations from single cell whole genome sequencing. We create models that accurately predict mutability in each tissue, and compare how the genomic predictors of mutability vary across the human body. Our analysis reveals that triplets emerge as robust predictors of mutability in comparison to epigenetic factors. Importantly, we observe both universal and tissue-specific mutagenic processes in healthy tissues, with implications for understanding the maintenance of germline versus soma and the mechanisms underlying early tumorigenesis.

Introduction

Mutation is a continual process that occurs in both somatic and germline tissues. The consequences of germline mutation are widely appreciated as they underpin both heritable genetic disease and adaptive evolution. The consequences of somatic mutation are similarly profound including the origin of cancers (García-Nieto et al. 2019), neurodegeneration (Lodato et al. 2015) and ageing (Finkel et al. 2007). Although mutation is a random process, the likelihood of mutation (mutability) at any given site is not uniform across the genome. Variation can be seen in large genomic windows but also can vary between adjacent sites, and the rate at which mutations occur can vary from site to site more than 100-fold (Ellegren et al. 2003).

Mutability variation is driven by both genetic and epigenetic processes. A well known genetic effect on mutability is CpG dinucleotides which are 10 fold more likely to mutate than non CpG sites (Hwang and Green 2004; Hodgkinson and Eyre-Walker 2011; Oman et al. 2022), when methylated cytosines are converted to thymine through deamination (Coulondre et al. 1978; Razin and Riggs 1980). Numerous epigenetic processes have also been associated with mutability variation. For instance,

regions close to the origin of replication are associated with lower mutation rates, due to the increased
activity of DNA repair (Supek and Lehner 2015). Many other processes have been found to affect
mutation rate variation, including transcription level, chromatin conformation, various histone marks,
and recombination (Michaelson et al. 2012; Schuster-Böckler and Lehner 2012; Blokzijl et al. 2016;
Carlson et al. 2018; García-Nieto et al. 2019; Moore et al. 2021). It is important to note that many of
these genetic and epigenetic processes that affect mutability are interdependent (Roadmap
Epigenomics Consortium et al. 2015), so if we hope to determine the relative contribution of these
different processes on mutation rate variation we need to consider them together in models that can
account for correlations among these forces (Chen et al. 2017; Supek and Lehner 2019). For
example, though DNAse accessibility correlates negatively with mutation at the megabase scale
(Polak et al. 2013), this effect was found to be minimal after controlling for other confounds (Supek
and Lehner 2017). Specifically, DNAse accessibility was identified as a predictor for replication timing
(Gindin et al. 2014). The few studies that have analysed multiple epigenetic and genetic factors
simultaneously (Chen et al. 2017; García-Nieto et al. 2019) have been used at larger genomic scales
such as on individual genes. Consequently, these studies do not capture the significant variations
observed at smaller scales, including per base variation. This limitation is critical given that small-scale
sequence context has been shown to explain 81% of the variance in human substitution probabilities
(Aggarwala and Voight 2016), emphasizing the necessity for analyses that include finer resolutions
86
Cellular environments and epigenetic landscapes impose substantially different mutagenic pressures
across tissues and cell types (Roadmap Epigenomics Consortium et al. 2015; García-Nieto et al.
2019). This is in evidence from a large variation in mutation load among tissues, ranging 23-fold from
the highest in intestinal crypts to the lowest in germline (Ren et al. 2022). However, the forces
responsible for this variation and its full extent across the body remains unclear. A critical limitation to
exploration of somatic mutation has been the inherent difficulty of sequencing rare, mosaic somatic
mutations. Until recently, our knowledge of somatic mutation has relied primarily on bulk sequencing
of tumours (Schuster-Böckler and Lehner 2012; Alexandrov et al. 2013), which can only extract
mutations that occur at relatively high frequency and therefore have been influenced by clonal
selection (Malikic et al. 2019) and may be biassed due to changes in the cellular and mutagenic
environment when tissue becomes cancerous. Other studies have used RNA (García-Nieto et al.
2019) or exome sequencing (Chen et al. 2017), and thus can detect patterns present in the coding
regions but can not analyse the majority of genome space that is non-coding. Single cell sequencing
(SCS) is a technology that may offer a method to identify low frequency mutations in healthy tissue.
SCS has the power to greatly increase the quantity and variety of discoverable mutations, facilitating
the analysis of the complete spectrum of somatic mutations and providing an unbiased view of the
mutations and their drivers across tissues. Until recently SCS has been prohibitively costly (Wang and

Navin 2015), and there has therefore been no studies that use SCS mutations to analyse multiple

epigenetic and genetic predictors of mutation at fine resolution across germline and soma tissues

(Supek and Lehner 2019).

To address this gap, we built multivariate regression models for four healthy tissues that predict the

relative probability of mutation at individual sites from 17 epigenetic and genetic predictors. We

leveraged pre existing datasets, including high quality SCS datasets for blood, liver and skin tissues,

and parent-offspring trios mutations in germline. We explored how somatic mutational processes vary

across healthy human tissue, and specifically addressed the questions:

1. How accurately can genetic and epigenetic processes predict mutation rate variation in different

114   healthy tissues?

115   2. What genetic and epigenetic factors predict mutation in healthy tissue?

116   3. How do the factors that predict mutation rate vary between tissues?

117   4. How much does the mutability of a site vary among tissues?

Results

How accurately can genetic and epigenetic processes predict mutation rate variation in different healthy tissues?

Our mutability models accurately predicted mutability in all four tissues analysed. In blood, germline,

liver and skin we report low mean absolute error (average MAE = 0.0205) based on the fit of predicted

mutability with the fraction of mutations in each group of sites (figure 1). The fact that mutability trained

on one set of sites (training set) was able to predict the proportion of mutations in each bin on

independent test datasets demonstrated that the genetic and epigenetic properties can predict the

main mutagenic processes occurring in blood, liver, germline and skin. We see a wide spread of

predicted mutability, ranging from 0.2 to 0.9, implying a wide range of actual mutability among sites.

The wide range of mutability is another indicator that the model is making meaningful predictions

because in the absence of a strong signal the null estimate for each site would be 0.5 mutability since

our dataset contained 50% mutations. Extreme mutability values that deviate from this null

demonstrate that the model is able to differentiate high and low mutability sites; indeed, when we

randomize the mutation status of the training data, mutability values cluster strongly around 0.5 and

73% of sites have mutability between 0.45 and 0.55, in contrast to only 18% of sites in this range in

the real data (see figure S1).

Interestingly, in the germline model we see a considerable gap between the highest mutability sites

and the rest, which appears to be driven by the extreme difference in mutability of germline CpG sites.

CpG sites are extremely mutable and so to test if these sites alone are driving model accuracy across

tissues, we created models removing these sites. Without CpGs, the spread of germline mutability

predictions reduces 40% from 0.24-0.93 to 0.27-0.67, however, the model maintains the same low

error on the remaining sites (MAE=0.02). MAE remains low in blood, liver and skin when CpGs are

removed (mean MAE = 0.022). No other tissue exhibits the same reduction in the spread of mutability

values as in the germline, with the skin only experiencing a 2.5% decrease from 0.17-0.97 to 0.17-

0.95 (see figure S2). Thus we find that, while CpG conversion accounts for the main mutagenic

process in highly mutability sites in all tissues but skin, there are numerous other critical mutagenic

processes that we have identified to accurately predict the ratio of mutant and non-mutant sites at

lower mutability.

Figure 1: Tissue specific mutability model fit and accuracy in healthy tissues. Here accuracy is estimated from how well the estimated mutability predicts the proportion of the sites in a given bin that are actually mutations as mean absolute error (MAE). Each data point represents a bin of 400 mutated or unmutated sites within a given mutability range. The proportion of sites that are true mutations in that bin is plotted against the average predicted mutability, which represents the model's estimation of the probability that a site would be a mutation in the test set. The dashed grey line is the 1:1 where the predicted mutability and proportion of mutations are equal.

What genetic and epigenetic factors predict mutation in healthy tissue?

The impact of each epigenetic and genetic factor on mutability can be extracted from each tissue model by analysing t-statistics. The t-statistic is a standardised parameter that indicates the strength and direction of effect a given predictor has on mutability, controlling for the confidence in that value. Across tissues, sequence triplets consistently emerge as strong predictors of mutation, with both mutagenic and protective effects seen for different triplets (figure 2). Uniformly across tissues we find that CpG sites (XCG triplets) are positively associated with mutation, however this effect is strongest in the germline, where CpG have two fold higher mutability than non CpG triplets. Furthermore, t-statistics for CpG triplets are 40% higher in the germline than other tissues. Another significant pattern is that all triplets with a central T (or A in the complementary strand) are significant and negatively associated with mutation (denoted here as "XTX"). This effect varies among XTX triplets, with some triplets exhibiting extremely protective effects against mutability (e.g. mean t-statistic for TTT across tissues= -7.0±1.6), while other triplets have an effect close to zero (e.g. mean t-statistic for ATC across tissues = -1.1±1.1). This trend is seen in all tissues, however the protective effect of XTX triplets is much stronger in the blood: t-statistics for XTX triplets are 170% more negative in blood compared to germline, skin and liver. A third triplet signature we find is that [T/C]CX triplets are extremely strong positive predictors of mutation in skin (220% higher t-statistic values than all other triplets), but this effect is not seen in other tissues, likely due to the fact that mutations at these sites are typical of UV light damage.

175
176
Figure 2: Comparison of the importance of genomic and epigenetic predictors of mutability among tissues. Every
predictor is represented by four points corresponding to each of the tissues, where each point represents the t-
statistic for a given tissue. The larger deviations from zero (grey dotted line) indicate a stronger effect on
mutability, with points above the line increasing mutability and those below reduce mutability. Here we only plot
predictors that have significant effect (p <0.005) in at least one tissue; tissues in which a given genomic property
has no significant effect are indicated with translucent data points. Predictors are ordered by the the most
extreme t-statistic value of the four tissues from most positive to most negative. A predictor with the same
mutagenic importance in two tissues would exhibit a larger t-statistic in the tissue with the larger dataset. Thus,
to ensure discrepancies in t-statistics reflect biologically meaningful differences, here the germline, liver and skin
datasets are downsampled to be equivalent to the smallest dataset, blood (n mutations = 9,082). See figure S8
for t-statistics values from the full models without downsampling.

188
189
We see that a diversity of epigenetic and genetic factors are important to predicting mutation beyond
triplets. Being in repetitive DNA is strongly indicative of mutation in the blood and skin, with t-statistics
of 8.9 and 15.9 respectively. Interestingly, while the effect of repeats in skin is the most extreme seen
across all tissue models, repeats have close to no effect on the liver or germline mutability (t-statistics
= -0.07 and -0.26 respectively). The GC content of the surrounding 10Kbp protects against mutation,

with a strong negative influence on the predicted mutability of a central site. However, this trend is only
seen in the liver (t-statistic = -3.6), with either no effect in blood or germline (t-statistic = -0.91 and -
0.28 respectively) or only slightly mutagenic in skin (t-statistic = 2.04). Epigenetic processes are also
important to predicting mutation. Here we find that sites at the nuclear periphery (indicated by high
laminB1 values) are more likely to mutate, though this effect is only significant in the blood and skin
tissue (t-statistic = 5.89 and 4.66 respectively). Histones marks are also important, though typically
only significant in one tissue; H3k27me3 marks (indicating facultatively repressed genes) are
mutagenic in the liver (t-statistic = 3.0), while H3k36me3 marks (actively transcribed genes) are
protective in the blood (t-statistic = -3.3). The variation seen in the importance and direction of effect of
epigenetic and genetic factors included in the model suggests that mutagenic processes are specific
to the cellular environments within each tissue.
206
Triplets play an important role in the predictive ability of these models, however we wanted to test
whether model accuracy relies solely on these predictors. Thus we created models without triplet
predictors and evaluated accuracy to see if other genetic and epigenetic factors contain enough
biologically relevant signal to accurately predict mutability. Models without triplet predictors still exhibit
high accuracy, with MAE only increasing on average 7.5% from 0.0205 to 0.022 across tissues.
Interestingly, in the liver model we see a decrease in error when triplets are removed, with MAE
reduced 10% from 0.021 to 0.019 (see figure S3 and table S3). Recall that another method to analyse
the fit is the range of mutability values: accurate extreme predictions (close to either 1 or 0) indicate
that the model has the power to leverage the epigenetic and genetic factors to assign a site high or
low mutability, while models that do not find associations between the predictors and mutation will
default to predicting 0.5 mutability as there are 50% mutations in the dataset. Without these predictors,
we find that the spread of predicted mutability drops 43% across tissues. In the absence of triplets, we
find the emergence of other predictors that can capture mutagenic signals (see figure S4). Across
tissues we find that GC-content at the 100 scale is mutagenic (average t-statistic = 7.7±3.0), possibly
reflecting some of the signal from high mutability CpG triplets. Similarly, sites at the nuclear periphery
are universally mutagenic in the absence of triplets (average t-statistic= 3.8±0.6), as opposed to the
tissue specific effect seen in the full model (see figure 2 above). Models without triplets also reveal
robust tissue-specific signals: for example, repetitive sequences exhibit pronounced mutagenicity in
skin and blood (t-statistic = 13.7 and 7.2 respectively), yet confer a protective effect in the germline (t-
statistic = -5.9). Similarly, many histone marks become important for predicting mutation but only in
one or two tissues; for example sites enriched for H3k27me3 is strongly mutagenic in the germline and
liver (t-statistic = 8.3 and 5.9 respectively), but has close to no effect in the blood or skin (t-statistic =
0.6 and -0.02 respectively). The fact that models without triplets maintain accuracy demonstrates that,

while triplets are important indicators of mutagenic processes, these mutagenic signals are also
encompassed within diverse genetic and epigenetic factors.
232
Notably, we included many genomic properties and genetic processes in our model that have been
found to drive mutation in other studies, but found no significant predictive power for mutability here.
For example, recombination rate has been shown to correlate with mutation (Lercher and Hurst 2002)
due to imperfect repair at recombination sites. However here we find no importance of recombination
in the germline tissue, with a t-statistic of 2.07 and p value of 0.038; though traditionally this would be
considered significant, here we consider over 300 predictors across all tissue models and so stricter
significance cutoffs are required to protect against type 1 error. Similarly, distance to replication origin
has been correlated to human SNP density (Wolfe et al. 1989; Stamatoyannopoulos et al. 2009; Chen
et al. 2010) and mutations in cancer genomes (Woo and Li 2012; Liu et al. 2013; Supek and Lehner
2017), in part due to the fact that DNA repair machinery is recruited to regions that replicate early
(Supek and Lehner 2015). Again we find no significant effect of replication timing in mutation in blood,
germine or skin models (p values range from 0.027 to 0.208).
245

How do the factors that predict mutation rate vary between tissues?
We can use model performance to assess the similarity of mutagenesis across tissues; if the
epigenetic and genetic factors have similar mutagenic roles in both tissues, a model trained on one
tissue will accurately predict mutability in the other. To assess accuracy, we analysed MAE and the
shape of the relationship between predicted and observed mutability. As expected, we see that
models have lower accuracy when predicting on other tissues than on datasets from the same tissue
on which they were trained, however the degree of decrease varies depending on the two tissues
being compared (see table 1). A strong pattern emerged that the skin model poorly predicts mutability
outcomes in other tissues, with MAE ranging from 5 to 5.7 times the value observed when predicting
on the skin mutations (for blood and germline tissue, respectively, see table 2). In these comparisons
we see that when mutability is 0 - 0.6 the skin model underestimates mutability but with a general
linear relationship (figure 3C). However, in bins with mutability > 0.6 we see the skin model is not
linear, with some sites having a negative relationship between observed and predicted mutability - a
pattern that remains consistent across all tissues, see figure S5. The loss of accuracy in higher
mutability bins is driven by [T/C]CX triplets: the skin model predicts these triplets to have high
mutability, but in the absence of UV damage in non-skin tissue, [T/C]CX triplets do not have a
markedly higher mutation rate.

Figure 3. Tissue specific mutability model fit and accuracy in healthy tissues, where the model predicts mutations

from tissues other than the one on which the model was trained. Three of six possible combinations of tissues

were chosen to illustrate the varying degrees of accuracy with which tissue-specific models can predict the

mutability of sites on other tissues, see figure S5 for all combinations. Each data point represents a bin of sites

which includes both mutated and unmutated sites (n=400). The proportion of sites that are mutations in that bin

is plotted against predicted mutability, which represents the model's estimation of the probability that a site would

be a mutation in the test set, averaged for all sites in the bin. The dashed grey line is the one-to-one line, where

the predicted mutability and proportion of mutations are equal

Table 1: Comparison of mean absolute error (MAE) of the predicted mutability and the estimated mutability when

mutability models are used to predict mutability on test mutation data sets from the same or different tissue

types. When models are tested on a different tissue, the fold change in error (MAE) is displayed in brackets.

		Mutation dataset			
		blood	**germline**	**liver**	**skin**
Mutability model	**blood**	0.028	0.064 (2.3x)	0.058 (2.1x)	0.035 (1.3x)
	germline	0.038 (2.0x)	0.019	0.036 (1.9x)	0.087 (4.6x)
	liver	0.024 (1.2x)	0.047 (2.4x)	0.02	0.04 (2.0x)
	skin	0.11 (5.0x)	0.125 (5.7x)	0.113 (5.1x)	0.022

The blood, liver, and germline tissues exhibit greater similarities, as each tissue's mutability model is

able to accurately predict mutations in other tissues. The most similar models are blood and liver, with

an increase in MAE of 20% when the liver model predicts blood mutations as outlined in table 1.

Furthermore, the data points closely align with the 1:1 relationship, as illustrated in figure 3A. While

the blood model displays higher error when predicting on liver mutations, it is noteworthy that the

observed trend maintains linearity and does not diverge greatly from the 1:1 line (figure S5). Germline

tissue diverges slightly from the blood and liver, as comparisons between these tissues display

moderate increases in error and deviations from linearity. The germline model exhibits a linear

relationship and modest accuracy when predicting on the blood and liver tissues (increase 100% and

90% respectively, see table 1 and figure S5). However, when the blood and liver models predict on

germline data we see greater increases in MAE (140% increase in the liver model, 130% in blood

model, table 1), and a lack of linearity: the model overestimates the mutability for bins with mutability

from 0.50-0.75 and underestimating some high mutability bins (>0.75) (see figure 3B). These results

suggest that there are tissue specific mutagenic processes present in the germline and not other

tissues. Additionally, the unique characteristics of the germline may stem from the augmented sample

size, as training the model on a larger dataset enables the detection of finer scale effects on

mutability.

We can analyse directly how the mutagenic effects of epigenetic and genetic factors vary between

tissues by comparing the t-statistic between models in a pairwise fashion: if predictors have similar

mutagenic effects in the two tissues being compared, then the data will center around the 1:1 line with

little dispersion. Again we find that the skin is distinct from other tissues with an average MAE of 0.081

compared to the other three tissues, which is a 1.5x larger difference than observed between other

pairs (see table 2). We can see the extent to which mutability differs in skin in figure 4B where many

genomic and epigenetic properties that are negatively associated with mutation in blood, are positively

associated with mutation in skin (MAE=0.091). In contrast, the greater similarity in the strength and

direction of mutagenic processes in other tissues is exemplified in figure 4A. When comparing the

germline and liver, we find a linear relationship between the t-statistics, though they do not always lie

along the 1:1 line. We see low error (MAE=0.044) with only GC content switching in its direction of

effect on mutation between the tissues. Interestingly, the pairwise comparison reveals that predictors

in the blood are generally lower than in other tissues (t-statistics are 3.5 times lower than in other

tissues, ±1.2). Comparisons between all tissues can be found in the supplemental figures S6-7 and

while they show varying degrees of similarity between the blood, liver and germline, it is clear that the

skin tissue is the most unique among those compared here.

Figure 4: Pairwise comparisons of the strength and direction of different genetic and epigenetic properties and
processes on mutability across tissues. Only predictors that are significant in at least one of the tissues after
correcting for multiple tests are displayed (p<0.01). These pairwise comparisons were chosen to illustrate how
the trends differ between similar models (A. germline vs. liver) and markedly different models (B. blood vs. skin).
Predictors that have a similar effect in both tissues are near 1:1 line (grey dotted line), while those that deviate
from equality have an effect unique to that tissue. Predictors in the top left or bottom right quadrant change the
direction of effect between tissues, indicating a process that is mutagenic in one tissue and protective in another.
Select informative points are labelled. Because the magnitude of t-statistics are proportional to the size of the
data set we downsampled each tissue's training set to be equivalent to the smallest dataset for this analysis
(blood $n_{mutations}$ = 9,082). For comparison of other tissue pairs, see figures S6-S7.

Table 2: Summary of the overall similarity of t-statistic values between tissues. Below we report the mean
absolute difference of the corresponding t-statistic for a given pair of mutability models. Larger numbers indicate
bigger discrepancies between the mutagenic effect of different genetic and epigenetic factors across tissues.

	germline	liver	skin
blood	0.068	0.051	0.091
germline		0.044	0.076
liver			0.077

How much does the mutability of a site vary among tissues?

We compared the predicted mutability of the same 15000 random sites estimated from each of the tissue specific models to determine the consistency of mutation rates across the genome in different cellular contexts. Overall the mutability of sites was clearly correlated across tissues with a mean MAE of 0.10 (figure 5) but the errors varied from 0.141 between skin and germline mutability to 0.064 in germline versus liver. Differences in the strength of correlation were driven by strong patterns of clustering, especially in skin and germline tissues. In all tissues CpG sites are enriched for high mutability, but the effect is exaggerated in germline where the mutability of CpG sites is separated from the rest of sites (figure 5, purple). In skin the [T/C]CX triplets, likely the result of UV damage, are distinct in that they have high mutability in skin and relatively average mutability in other tissues (figure 5, blue). There are also a group of sites from repetitive DNA that have higher mutability in blood (mutability 0.6-1) and very high in skin (mutability >0.9), but are not very mutable in germline or liver. Therefore, while there are correlations in the mutation rate across tissues, strong patterns are apparent that suggest the same site may have very different mutation rates depending on the cell it finds itself in.

Figure 5: Pairwise comparisons of per site mutability in different tissues. Blood, germline, liver and skin models were all used to predict mutability on a set of 15000 random sites. Sites were coloured to highlight the key factors that result in mutability variation between tissues, including CpG triplets, [T/C]CX triplets, and DNA repeat context. Any site that was a combination of these categories was labelled 'combined' and all other sites are shown as translucent black points. The 1:1 line is shown in grey and the mean absolute error (MAE) from that line is reported in each panel.

Discussion

Mutability models are able to predict variation in mutation rate with high accuracy

Our goal was to explore the effects of genetic and epigenetic factors together on mutability at the single base resolution in healthy human tissues. We compiled the main genomic factors known to affect mutation and created mutability models for the germline and soma (blood, liver and skin) that all demonstrate impressive precision and low error (average MAE = 0.0205). In all tissues we successfully discriminated between highly mutable and less mutable sites. For example, 90% of sites classified to have a mutability score of 0.9 are indeed true mutations, even though the model was not trained on these sites and only 50% of test sites were mutations. The converse is also true, as these models can successfully classify non-mutant sites into lower mutability bins where there are very few true mutations (as low as 17.3%). Note that non-mutant sites in high mutability bins and mutant sites in low mutability bins are not necessarily misclassified; instead it is expected that most high mutability sites in the genome do not mutate regularly and some low mutability sites do mutate despite this being relatively less probable. This fact is what necessitates binning sites by predicted mutability to evaluate the performance of the models. The fact that predicted mutability correlates strongly with the observed fraction of mutations suggests the mean mutability of sites in each bin is well estimated, even with narrow ranges of mutability in bins and relatively few sites (400 sites). This does not negate the possibility of some inaccuracy and unexplained variance in the predictions of mutation rates of individual sites (Eyre-Walker and Eyre-Walker 2014). Very similar approaches have been used to predict mutability in model organisms (Ness et al. 2015; Saxena et al. 2019; López-Cortegano et al. 2021) and the human germline (Michaelson et al. 2012) but our work represents the first time this approach has been used in human soma. The accuracy of these models reinforces the utility of this statistical approach, and that mutability can be predicted from models trained with as few as 9,082 mutations. Importantly, the fit of the models demonstrates that the epigenetic and genetic predictors included here capture important mutagenic signals, which is important for understanding the mutational processes in healthy tissues.

DNA sequence-based properties are the strongest predictors of mutation rate variation across all tissues.

In our analysis, the DNA sequence itself emerges as the strongest predictor of mutation. Historically, studies have analysed the impact of genetic and epigenetic changes on mutation rate variation in large genomic regions (Chen et al. 2017; García-Nieto et al. 2019). In contrast we find that the local scale (<10bp) is the most important for predicting mutation. For example, a site being in a repeat (1bp scale) is the strongest indicator of mutability seen in all tissues, but there is no effect of being in a 10kbp region that is enriched for repeats. In line with previous observations, both the specific nucleotide at a site and its two flanking sites (triplet) strongly predict human mutability (Michaelson et al. 2012; Aggarwala and Voight 2016). Among triplets, mutagenic effect varies 180 standard deviations (t-statistics, ACG =29.4 , TTT = -25.0). As expected, deamination of methylated cytosines causes CpG sites to be highly and universally mutable, however the strength of this effect across the four different CpG triplets (XCG) varies up to 2.4 fold within a tissue. Conversely, we find that triplets with a central A/T (denoted X[A/T]X) are protective against mutation across all tissues, but with up to 7-fold variation between the most and least protective X[A/T]X triplet within a tissue. Our findings suggest that local sequence context is one of if not the best predictor of mutability and can also drive changes in the mutation rate of the same site in different cell types. This argues that models of sequence evolution should be using the sequence context to determine the null expectation of mutation rate variation, but each species or cell type requires its own dataset to completely capture the variation in mutation rate in different contexts.

One of the reasons that triplets are such a powerful predictor of mutation is because they capture many different mutagenic processes that affect DNA in a biassed way (Alexandrov et al. 2013; Moore et al. 2021). These effects can be due to different repair mechanisms or variable agents that cause specific types of DNA damage. This means that although the sequence is the same across tissues, the triplets can have different mutation rates depending on the specific cellular environment. For example, the protective effect of triplets with an A/T at the central base is 170% stronger in blood than all other tissues, suggesting that transient strand denaturation at A/T sites may play a lesser role in mutagenesis in blood (Leroy et al. 1988). We also find that CpG sites are 40% more mutable in the germline than soma (see below). Variable triplet mutation rates have been shown to reflect numerous external mutagens (Alexandrov et al. 2013). A clear example is UV light that causes mutations in triplets with a central C and either a C or T upstream (denoted [T/C]CX). Consequently, we find [T/C]CX triplets are highly mutable in skin, while having a protective or inconsequential effect in other tissues. Interestingly, this effect is seen in the skin regardless of chromatin accessibility, suggesting that mutations from UV damage accumulate independent of condensation state (Pich et al. 2018).

414   Thus, consistent with other work (Alexandrov et al. 2013; Moore et al. 2021), we find that triplets are

415   important for capturing the mutagenic processes that are endogenous to cells, as well as those

416   incurred via external mutagens.

Epigenetic factors that capture mutagenic processes vary among tissues

Though triplets are powerful predictors of mutation, other genetic and epigenetic factors also account

for a significant fraction of the mutagenic signal. We created models without triplets and found that

models lose the ability to predict extreme mutability, with a mean 43% reduction in the range of

predicted mutability (see table S3). However, models still retain accuracy, as MAE increases only

7.5% within this range. This is in line with other studies that find that epigenetic elements can explain

between 55-80% of mutation variation in cancer (Schuster-Böckler and Lehner 2012; Polak et al.

2015). Many internal processes of cells are interdependent and involve interactions of DNA with

epigenetic mechanisms, thus both the sequence-based and epigenetic predictors may capture the

same mutagenic processes. When all predictors are combined triplets emerge as superior predictors,

likely owing to the fact they can capture the notable local-scale variability in human mutation rates.

However, the variation in triplet mutability among the limited set of tissues analyzed here argues for

much more comprehensive sampling of different cell types, individuals, ages and even species if we

hope to produce accurate predictive models of mutation.

Despite this, the effect of epigenetic processes on mutation provides both insight into the mechanism

of mutation and captures information not reflected by triplet sequence. In the full models that include

triplets, many epigenetic factors significantly predict mutation. Though trends are more nuanced, we

find that open chromatin regions are associated with lower mutability (e.g. H3k36me3). Regions at the

nuclear periphery (laminB1) are silenced as heterochromatin (Filion et al. 2010), and being in these

regions is a strong indicator of mutation in blood and skin. In both germline and liver tissue, sites in

facultatively repressed chromatin (H3k27me3) are associated with higher mutability (t-statistics =5.4

and 5.7 respectively) while in the blood, regions that are annotated (t-statistic = -2.8) and marked

(H3k36me3, t-statistic = -4.0) as genes have lower mutability. Our models therefore support the

hypothesis that transcription coupled repair outweighs the damage caused by transcription (Moore et

al. 2021). However, the fact that we do not see as strong and consistent a signal across tissues

suggests that the relationship between chromatin state and mutability may be more nuanced than

previously thought or not particularly strong. In fact, other previously established trends are weaker or

have variable effects across tissues in our analyses. For example, we do not see an effect of

replication timing and find that H3k36me3 protects against mutation only in blood, whereas these

predictors have previously been strongly linked with mutability (García-Nieto et al. 2019; Supek and

Lehner 2019). These discrepancies amongst previous literature highlight the importance of including
many factors simultaneously to understand independent effects and control for multicollinearity. The
fact that the same predictors have variable effects across tissues suggests that there are important
tissue specific effects that influence mutability and contribute to the overall mutation rate and spectra
in each tissue.
454
Healthy tissues exhibit varying degrees of similarity
We also explored the similarity of mutability across healthy human tissues by comparing how the
effect of each epigenetic and genetic predictor and how similar model predictions were between
tissues and how this leads to variation in mutability of sites across tissues. The first major finding is
that the mutation profile of skin is the most distinct, driven primarily by the preponderance of mutations
with the triplet signature of UV damage, exemplifying how external mutagens can dominate mutation.
Interestingly, the liver and blood models can predict mutations in skin with only moderately reduced
accuracy. This suggests that, besides the external mutagen UV light signature, the mutagenic
processes that occur in these tissues are similar, and that the liver and blood models are accurately
predicting mutations not caused by UV damage in skin. This is supported by work that extracted the
triplet signatures in healthy tissue and found that the two endogenous triplet signatures that dominate
many tissues are also present in skin (Moore et al. 2021). The similarity between skin and liver may be
further explained by oxidative damage in the liver, as [T/C]CX triplets are often mutated in the
signature associated with APOBEC (Alexandrov et al. 2013).
469
We also find that the germline exhibits important differences from blood and liver tissue. The most
distinctive feature of germline mutability is the very well characterized mutagenic effect of CpG sites.
This effect is three times stronger in germline than in other tissues. Interestingly, in an earlier study
investigating the effect of triplet sequence on mutation they found a similar contribution of the CpG
signature in germline and soma (Moore et al. 2021). Despite the fact that the major mutagenic process
(CpG deamination) varies between these tissues, we find that the germline, liver and blood exhibit
moderate to high predictive accuracy on each other (see table 1 and 2). This is consistent with (Moore
et al. 2021) who reported that the major triplet signatures were shared across most tissues. In the two
studies that have examined the effects of epigenetic processes on mutation in human germline and
soma, differences between the soma and germline were identified (Chen et al. 2017; Moore et al.
2021). Specifically, transcription played a protective role in soma (over ten tissues including skin and
liver) but not in germline, and the correlation between late replicative timing and increased mutation
was stronger in soma. Notably these associations were based on mutations in coding sequence only
that were modelled with one variable at a time (Moore et al. 2021) or from mutations from bulk cancer

tumours (Chen et al. 2017), and these differences could explain why we did not observe either pattern

in our data. The fact that our results differ highlights the importance of including local scales, analysing

intergenic regions, and modelling multiple variables simultaneously to control for correlations among

genomic properties. These differences between germline and soma are worth noting because

analyzing patterns of genetic variation in a pedigree, a population or through evolutionary time reflects

the process of mutation in the germline, and those findings may bias our assumptions about mutability

across the body and how it influences somatic mutation and disease.

Differences in mutation profiles across various cancers have been well established (Alexandrov et al.

2013; Chen et al. 2017), but the time when these changes arise remains unclear. It raises a critical

question: to what extent do mutation differences among tumours reflect mutability in healthy tissues

versus changes in mutability acquired during tumour evolution? Comprehensive analyses of healthy

somatic tissues are scarce. (García-Nieto et al. 2019) detected differences between tissues using

epigenetic and genetic factors, however, their analysis was limited to RNA-seq data, capturing only

the 1.5% (Lander 2011) of the genome in coding regions. (Moore et al. 2021) found that genome-wide,

two triplet signatures accounted for the majority of mutations in most tissues, arguing that healthy

tissues generally share mutation processes. However, in a limited analysis of the impact of gene

expression and replication timing, Moore et al. did find differences between germline and soma

tissues. By collating data from four tissues and seventeen predictors we have shown that somatic

mutability is both complex and differs among tissues. Our comparison of mutability of the same sites in

different cellular contexts (figure 5) clearly demonstrates that sites mutate at very different rates in

different cells. Therefore, the probability of a site contributing to ageing and disease across tissues is

highly variable. If this variation results in some genes being more prone to mutation, it could mean that

their association with disease could be driven by higher mutation rate, by causing the disease, or both.

We are currently limited by somatic sequencing technology: many tissues are not amenable to the

current standard practices, because they cannot be clonally expanded (e.g. post mitotic tissues like

the brain) or because they do not have local monoclonal structures available for for laser-

microdissection (e.g. the blood). Error corrected sequencing technologies have reported error rates as

low as 5 in a billion bases (Abascal et al. 2021) and will offer a new way to identify somatic mutations.

Analysing larger datasets and more tissues than ever before will help deconvolute the mutagenic

processes in healthy tissue.

Methods

Mutation data sets

To do a comprehensive analysis of how epigenetic and genomic processes impact mutability in four healthy tissues we leveraged publicly available mutation datasets. We used 104,941 single nucleotide mutations in the germline, identified from whole genome sequencing of 1.5k parent-offspring trios (Jónsson et al. 2017). For somatic tissues, mutations from SCS were downloaded from the Database for Somatic Mutations in Normal Cells (Miao et al. 2019) or the database of somatic mutations in normal tissues (Sun et al. 2022), see supplemental table 1 for a summary. We also used a mappability filter that describes whether sites can be readily aligned to the genome and removed both mutations and non-mutations at non-mappable sites as in previous work (Carlson et al. 2018).

Genomic predictor data

We used publicly available tissue-specific data for genomic predictors of mutation to train each of our mutability models. Each model contains the same predictor categories to facilitate comparison across tissues, including cell-type specific processes (e.g. histone modification, transcription) and genome sequence-based properties (e.g. GC content) that have been used previously to predict mutability (Michaelson et al. 2012; Chen et al. 2017; Carlson et al. 2018; García-Nieto et al. 2019).

It has been shown that some predictors affect mutation at different scales, for example G/C bases tend to mutate at a higher rate than A/T but high GC at the scale of 1000bp tends to reduce spontaneous mutation (Ellegren et al. 2003; Supek and Lehner 2019). We therefore calculated the value of these properties across various spatial contexts. Predictors at multiple different scales were only included in the model where intercorrelation was low enough to permit (see supplemental tables S13-16 for VIF summaries). Supplemental table 2 contains a summary of the predictors used in our model.

Measurements of some tissue-specific predictors were vulnerable to technological inconsistencies that could bias our analysis. For example, transcription data measured with RNASeq across different tissues could have different depth of coverage, read length or read architecture. Thus we strictly standardised tissue specific predictors so that for a given predictor we chose datasets that were as similar as possible based on the molecular method, source lab, fragment length, read length, date submitted, file type, cell type, age, and sex. In-depth details regarding these predictors including the data source and lab technique can be found in table S2.

Training mutability models

To explore which genomic or epigenetic properties drive mutation, we trained models that used these properties to predict whether a given site was a mutant or non-mutant in our training data. The relative probability (0 - 1.0) that a site would be in the mutation category is referred to as 'mutability'. Extending previous work (Michaelson et al. 2012; Ness et al. 2015; Carlson et al. 2018), we used logistic regression to train a predictive model that used the genetic and epigenetic properties to distinguish between mutated and non-mutated sites. Non-mutant sites were randomly selected from the mappable genome without overlap with the mutation dataset. The full dataset was split evenly into a training and testing set, with equal proportions of mutant and non-mutant sites in each half. Model accuracy was assessed on the independent "testing" dataset. All code used for these analyses is available at: https://github.com/MadeleineOman/mutability/tree/master

Model predictions and performance

To assess how well our models predicted mutability, we used mean absolute error (MAE). MAE is generally calculated as the mean absolute difference between the predicted and actual response value of a given data point. In our case this would be whether a mutant site (1) is properly predicted to have a mutability >0.5 or a non-mutant site (0) with a mutability <0.5. However, mutation rate is a continuous property with some sites being more likely to mutate than others. Therefore we need to test whether sites that are predicted to have higher mutability are proportionally more likely to be mutations and vice versa. To do so we can aggregate all the sites in the test set with a given mutability value and determine if the fraction of mutations in that bin is proportional to mutability. Using this method means that we may not specifically know the accuracy of any one site prediction nor how much variance there truly is in a bin (Eyre-Walker and Eyre-Walker 2014), but instead we can test if the mean mutability of the bin is accurate which allows us to interpret the fit and relative effect of predictors. Bin size was set at 400 sites per bin, as it is the maximum size that allows at least 20 bins in the smallest training dataset (blood n_{sites}=9082). To test the validity of this binning method we compare accuracy to a null model where the model predicts on data where the mutation status of the sites are randomized so that there should be no genomic properties that predict mutability (see figure S1).

To explore how mutational processes vary between tissues, we also used trained models from each tissue to predict mutability of sites from other tissues. If a model presents high error when predicting on another tissue, this indicates differences in the strength and direction of predictors in the fitted model in each respective tissue.

Genetic and epigenetic predictors of mutability

To understand which genomic properties and epigenetic processes are the best predictors of mutation

we extracted the effect of each predictor from each tissue model. In regression, each predictor is

assigned a β coefficient (slope) that describes the direction and strength of that predictor in

determining mutability. The t-statistic of each predictor represents a normalized indicator of the

importance of that predictor and more is suitable for comparison across models. The t-statistic for a

given predictor is calculated as follows,

$$t_{\widehat{\beta}} = \frac{\widehat{\beta}}{SE(\widehat{\beta})}$$

592   where β(hat) is the estimator for the true β value in the model and SE(β(hat)) is its standard error. To

593   compare how the same genomic properties affected mut differently across tissues, we normalized all

594   predictors in all tissues to a mean of zero and standard deviation of one. This allowed us to directly

595   compare the varying importance of each predictor in each tissue across models.

Variation of tissue-specific mutability of individual sites

We also wanted to compare how much the mutability of any given site in the genome might vary

depending on the tissue. To achieve this we chose 15000 random sites from the genome. For each

site we used the previously trained tissue-specific mutability model and predictor data to estimate the

site's predicted mutability in each of the four tissues. We then compared these sites across tissues in

a pairwise manner and analyzed the similarity of mutability with a linear model and reported the R^2 as

a measure of similarity.

Data access

606   The full analysis is available online at GitHub at https://github.com/MadeleineOman/mutability and all

607   data used in this study are from public databases, details of which can be found in supplemental table

608   S1 and corresponding code in the GitHub repository.

Competing interest statement

610   The authors have no competing interests to declare.

Acknowledgements

This work was supported by a Natural Sciences and Engineering Research Council (NSERC)
Discovery grant (RGPIN/06331-2016) and Canadian Foundation for Innovation John R. Evans
Leaders fund (35591) to R.W.N. and a Canada Graduate Scholarship-Doctoral (CGS D) to M.O.

References

Abascal F, Harvey LMR, Mitchell E, Lawson ARJ, Lensing SV, Ellis P, Russell AJC, Alcantara RE, Baez-Ortega A, Wang Y, et al. 2021. Somatic mutation landscapes at single-molecule resolution. Nature. 593(7859):405–410.

Aggarwala V, Voight BF. 2016. An expanded sequence context model broadly explains variability in polymorphism levels across the human genome. Nat Genet. 48(4):349–355.

Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, Bignell GR, Bolli N, Borg A, Børresen-Dale AL, et al. 2013. Signatures of mutational processes in human cancer. Nature. 500(7463):415–421.

Blokzijl F, De Ligt J, Jager M, Sasselli V, Roerink S, Sasaki N, Huch M, Boymans S, Kuijk E, Prins P, et al. 2016. Tissue-specific mutation accumulation in human adult stem cells during life. Nature. 538(7624):260–264.

Carlson J, Locke AE, Flickinger M, Zawistowski M, Levy S, Consortium, Bridges, Myers RM, Boehnke M, Kang HM, Scott LJ, et al. 2018. Extremly rare variants reveal patterns of germline mutation rate heterogeneity in humans. Nat Commun.(2018). doi:10.1038/s41467-018-05936-5. https://www.nature.com/articles/s41467-018-05936-5.

Chen C-L, Rappailles A, Duquenne L, Huvet M, Guilbaud G, Farinelli L, Audit B, d'Aubenton-Carafa Y, Arneodo A, Hyrien O, et al. 2010. Impact of replication timing on non-CpG and CpG substitution rates in mammalian genomes. Genome Res. 20(4):447–457.

Chen C, Qi H, Shen Y, Pickrell J, Przeworski M. 2017. Contrasting determinants of mutation rates in germline and soma. PLoS Genet. 207(September):255–267.

Coulondre C, Miller JH, Farabaugh PJ, Gilbert W. 1978. Molecular basis of base substitution hotspots in Escherichia coli. Nature. 274(5673):775–780.

Ellegren H, Smith NGC, Webster MT. 2003. Mutation rate variation in the mammalian genome. Curr Opin Genet Dev. 13(6):562–568.

Eyre-Walker A, Eyre-Walker YC. 2014. How Much of the Variation in the Mutation Rate Along the Human Genome Can Be Explained? G3 Genes|Genomes|Genetics. 4(9):1667–1670.

Filion GJ, van Bemmel JG, Braunschweig U, Talhout W, Kind J, Ward LD, Brugman W, de Castro IJ, Kerkhoven RM, Bussemaker HJ, et al. 2010. Systematic Protein Location Mapping Reveals Five Principal Chromatin Types in Drosophila Cells. Cell. 143(2):212–224.

Finkel T, Serrano M, Blasco MA. 2007. The common biology of cancer and ageing. Nature. 448(7155):767–774.

García-Nieto PE, Morrison AJ, Fraser HB. 2019. The somatic mutation landscape of the human body.

Genome Biol. 20(298). doi:10.1186/S13059-019-1919-5.
https://genomebiology.biomedcentral.com/articles/10.1186/s13059-019-1919-5.

Gindin Y, Valenzuela MS, Aladjem MI, Meltzer PS, Bilke S. 2014. A chromatin structure-based model
accurately predicts DNA replication timing in human cells. Mol Syst Biol. 10(3):722.

Hodgkinson A, Eyre-Walker A. 2011. Variation in the mutation rate across mammalian genomes. Nat
Rev Genet. 12(11):756–766.

Hwang DG, Green P. 2004. Bayesian Markov chain Monte Carlo sequence analysis reveals varying
neutral substitution patterns in mammalian evolution. Proc Natl Acad Sci U S A. 101(39):13994–
14001.

Jónsson H, Sulem P, Kehr B, Kristmundsdottir S, Zink F, Hjartarson E, Hardarson MT, Hjorleifsson
KE, Eggertsson HP, Gudjonsson SA, et al. 2017. Parental influence on human germline de novo
mutations in 1,548 trios from Iceland. Nature. 549(7673):519–522.

Lander ES. 2011. Initial impact of the sequencing of the human genome. Nature. 470(7333):187–197.

Leroy JL, Kochoyan M, Huynh-Dinh T, Guéron M. 1988. Characterization of base-pair opening in
deoxynucleotide duplexes using catalyzed exchange of the imino proton. J Mol Biol. 200(2):223–238.

Liu L, De S, Michor F. 2013. DNA replication timing and higher-order nuclear organization determine
single-nucleotide substitution patterns in cancer genomes. Nat Commun. 4(1):1–9.

Lodato MA, Woodworth MB, Lee S, Evrony GD, Mehta BK, Karger A, Lee S, Chittenden TW, D'Gama
AM, Cai X, et al. 2015. Somatic mutation in single human neurons tracks developmental and
transcriptional history. Science. 350(6256):94–98.

López-Cortegano E, Craig RJ, Chebib J, Samuels T, Morgan AD, Kraemer SA, Böndel KB, Ness RW,
Colegrave N, Keightley PD. 2021. De Novo Mutation Rate Variation and Its Determinants in
Chlamydomonas. Mol Biol Evol. 38(9):3709–3723.

Malikic S, Jahn K, Kuipers J, Sahinalp SC, Beerenwinkel N. 2019. Integrative inference of subclonal
tumour evolution from single-cell and bulk sequencing data. Nature Communications 2019 10:1.
10(1):1–12.

Miao X, Li X, Wang L, Zheng C, Cai J. 2019. DSMNC: a database of somatic mutations in normal
cells. Nucleic Acids Res. 47(D1):D971–D975.

Michaelson JJ, Shi Y, Gujral M, Zheng H, Malhotra D, Jin X, Jian M, Liu G, Greer D, Bhandari A, et al.
2012. Whole-genome sequencing in autism identifies hot spots for de novo germline mutation. Cell.
151(7):1431–1442.

Moore L, Cagan A, Coorens THH, Neville MDC, Sanghvi R, Sanders MA, Oliver TRW,
Leongamornlert D, Ellis P, Noorani A, et al. 2021. The mutational landscape of human somatic and
germline cells. Nature. 597(7876):381–386.

Ness RW, Morgan AD, Vasanthakrishnan RB, Colegrave N, Keightley PD. 2015. Extensive de novo
mutation rate variation between individuals and across the genome of Chlamydomonas reinhardtii.
Genome Res. 25(11):1739–1749.

Oman M, Alam A, Ness RW. 2022. How Sequence Context-Dependent Mutability Drives Mutation
Rate Variation in the Genome. Genome Biol Evol. 14(3):evac032.

Pich O, Muiños F, Sabarinathan R, Reyes-Salazar I, Gonzalez-Perez A, Lopez-Bigas N. 2018.
Somatic and germline mutation periodicity follow the orientation of the DNA minor groove around
nucleosomes. Cell. 175(4):1074–1087.e18.

Polak P, Karlic R, Koren A, Thurman R, Sandstrom R, Lawrence MS, Reynolds A, Rynes E,
Vlahovicek K, Stamatoyannopoulos JA, et al. 2015. Cell-of-origin chromatin organization shapes the
mutational landscape of cancer. Nature. 518(7539):360–364.

Polak P, Lawrence MS, Haugen E, Stoletzki N, Stojanov P, Thurman RE, Garraway LA, Mirkin S, Getz
G, Stamatoyannopoulos JA, et al. 2013. Reduced local mutation density in regulatory DNA of cancer
genomes is linked to DNA repair. Nature Biotechnology 2013 32:1. 32(1):71–75.

Razin A, Riggs AD. 1980. DNA methylation and gene function. Science. 210(4470):604 LP–610.

Ren P, Dong X, Vijg J. 2022. Age-related somatic mutation burden in human tissues. Frontiers in
Aging. 3. doi:10.3389/FRAGI.2022.1018119. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9534562/.

Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-
Moussavi A, Kheradpour P, Zhang Z, Wang J, et al. 2015. Integrative analysis of 111 reference
human epigenomes. Nature. 518(7539):317–330.

Saxena AS, Salomon MP, Matsuba C, Yeh SD, Baer CF. 2019. Evolution of the mutational process
under relaxed selection in Caenorhabditis elegans. Mol Biol Evol. 36(2):239–251.

Schuster-Böckler B, Lehner B. 2012. Chromatin organization is a major influence on regional mutation
rates in human cancer cells. Nature. 488(7412):504–507.

Stamatoyannopoulos JA, Adzhubei I, Thurman RE, Kryukov GV, Mirkin SM, Sunyaev SR. 2009.
Human mutation rate associated with DNA replication timing. Nat Genet. 41(4):393–395.

Sun S, Wang Y, Maslov AY, Dong X, Vijg J. 2022. SomaMutDB: a database of somatic mutations in
normal human tissues. Nucleic Acids Res. 50(D1):D1100–D1108.

Supek F, Lehner B. 2015. Differential DNA mismatch repair underlies mutation rate variation across
the human genome. Nature. 521(7550):81–84.

Supek F, Lehner B. 2017. Clustered Mutation Signatures Reveal that Error-Prone DNA Repair Targets
Mutations to Active Genes. Cell. 170(3):534–547.e23.

Supek F, Lehner B. 2019. Scales and mechanisms of somatic mutation rate variation across the
human genome. DNA Repair . 81(July):102647–102647.

Wang Y, Navin NE. 2015. Advances and Applications of Single-Cell Sequencing Technologies. Mol
Cell. 58(4):598–609.

Wolfe KH, Sharp PM, Li W-H. 1989. Mutation rates differ among regions of the mammalian genome.
Nature. 337(6204):283–285.

Woo YH, Li W-H. 2012. DNA replication timing and selection shape the landscape of nucleotide
variation in cancer genomes. Nat Commun. 3(1):1–8.