1

The Small Tumor Antigen of Merkel Cell Polyomavirus Accomplishes Cellular

Transformation by Uniquely Localizing to the Nucleus Despite the Absence of a Known

Nuclear Localization Signal

Kaira R. Thevenin [a], Isabella S. Tieche [a], Cody E. Di Benedetto [a], Matt Schrager [a], Kristine N. Dye [a b]

[a] Department of Health Sciences, Stetson University, DeLand, FL 32723, USA

[b] Department of Biology, Stetson University, DeLand, FL 32723, USA

Corresponding Author: Kristine N. Dye (kdye1@stetson.edu); 421 N Woodland Blvd,

DeLand, FL, 32723

ABSTRACT

Background: Merkel Cell Carcinoma (MCC) is an aggressive skin cancer that is three times deadlier than melanoma. In 2008, it was found that 80% of MCC cases are caused by the genomic integration of a novel polyomavirus, Merkel Cell Polyomavirus (MCPyV), and the expression of its small and truncated large tumor antigens (ST and LT-t, respectively). MCPyV belongs to a family of human polyomaviruses; however, it is the only one with a clear association to cancer.

Methods: To investigate the role and mechanisms of various polyomavirus tumor antigens in cellular transformation, Rat-2, 293A, and human foreskin fibroblasts were transduced with pLENTI MCPyV LT-t, MCPyV ST, TSPyV ST, HPyV7 ST, or empty pLENTI and assessed through multiple transformation assays, mass spectrometry, and subcellular fractionations. One-way ANOVA tests were used to assess statistical significance.

Results: Soft agar, proliferation, doubling time, glucose uptake, and serum dependence assays confirmed ST to be the dominant transforming protein of MCPyV. Furthermore, it was found that MCPyV ST is uniquely transforming, as the ST antigens of other non-oncogenic human polyomaviruses such as Trichodysplasia Spinulosa Polyomavirus (TSPyV) and Human Polyomavirus 7 (HPyV7) were not transforming when similarly assessed. Identification of structural dissimilarities between transforming and non-transforming tumor antigens revealed that the uniquely transforming domain(s) of MCPyV ST are likely located within the structurally dissimilar loops of the MCPyV ST unique region. Of all potential MCPyV ST cellular interactors, 62% are exclusively or transiently nuclear, suggesting that MCPyV ST localizes to the nucleus despite the absence of a canonical nuclear localization signal. Indeed, subcellular fractionations confirmed that MCPyV ST could achieve nuclear localization through a currently unknown, regulated mechanism independent of its small size, as HPyV7 and TSPyV ST proteins were incapable of nuclear translocation. Although nuclear localization was found to be important for several

transforming properties of MCPyV ST, some properties were also performed by a cytoplasmic

sequestered MCPyV ST, suggesting that MCPyV ST may perform different transforming functions in

individual subcellular compartments.

Conclusions: Together, these data further elucidate the unique differences between MCPyV ST and other

polyomavirus ST proteins necessary to understand MCPyV as the only known human oncogenic

polyomavirus.

66   **Keywords:** Merkel Cell Polyomavirus, Merkel Cell Carcinoma, Cellular Transformation,

67   Trichodysplasia Spinulosa Polyomavirus, Human Polyomavirus 7, Nuclear Localization

BACKGROUND

Merkel Cell Carcinoma (MCC) was first described in the 1970s as a rare, aggressive skin cancer, likely a consequence of ultraviolet radiation-induced mutations in cell cycle-regulation genes (1–3). However, in 2008, a novel human polyomavirus, subsequently termed Merkel Cell Polyomavirus (MCPyV), was found to be integrated within, and the etiologic agent of, ~80% of MCC tumors (4). Further investigation identified MCPyV as a ubiquitous, asymptomatic human virus that, in rare cases of immunosuppressed individuals, could integrate into the human genome and lead to the development of cancer (5–9).

The small, 5.3kb dsDNA genome of MCPyV is divided into an early region and a late region (ER and LR, respectively) by a bidirectional promoter (4,10). Due to its constrained genomic size, the ER utilizes alternative splicing and overprinting to generate 4 viral proteins, including the large tumor antigen (LT), small tumor antigen (ST), alternate large T open reading frame (ALTO), and the 57kT protein, while the LR encodes the VP1 and VP2 structural proteins (10–12). Although all of these viral proteins are expressed for successful viral replication in the normal MCPyV viral lifecycle, in MCC, only a truncated LT (LT-t) and ST are expressed and necessary for the viability of MCC tumors, suggesting a role for these proteins in tumorigenesis and maintenance (4,13,14). Several researchers have identified transforming capabilities of both ST and LT-t, including, but not limited to, perturbation of MYCL and Rb, respectively (10,15–19).

MCPyV belongs to a family of 14 known human polyomaviruses but is currently the only one with a clear association to cancer (20,21). However, polyomaviruses of other species, such as Simian Virus 40 (SV40), which infects simian species, have been found to be oncogenic in rodents (22–25). As one of the most intensively studied oncogenic viruses, the study of SV40 oncogenesis has led to the development of several transformation assays, which have subsequently been used to direct the study of MCPyV transformation and tumorigenesis.

Transformation is defined as the acquisition of expanded proliferation and/or survival potential of a cell (26). Many *in vitro* transformation assays exist, including soft agar assays, focus formation,

doubling time, proliferation rate, serum dependence, and metabolic assays, each of which assesses

different aspects and pathways involved in cellular transformation that may collectively contribute to

tumorigenesis *in vivo.* In order to accomplish cellular transformation, viral proteins bind and perturb

cellular pathways that allow for expanded proliferation and survival potential (27,28). Cellular proteins

and pathways operate in specific subcellular compartments, including the cytoplasm, nucleus, and

membrane (29,30). As localization influences the regulation and activity of specific proteins and

pathways, it is tightly regulated (30,31). Therefore, protein localization to the nucleus is tightly regulated

and restricted to proteins containing a nuclear localization signal (NLS), as this is where the DNA is

contained, and gene expression is initiated (32).

Herein, we employed an alternative approach to elucidate the mechanism of transformation of

MCPyV T antigens. As opposed to previous approaches of comparing similarities between SV40 and

MCPyV T antigens, several transformation assays on various oncogenic and non-oncogenic human

polyomavirus tumor antigens were performed to identify the unique domains and mechanisms of MCPyV

T-antigen mediated oncogenesis. Together, we identified MCPyV ST to be predominantly and uniquely

transforming when compared to MCPyV LT-t and the ST of other, skin-tropic human polyomaviruses.

Furthermore, MCPyV ST was found to uniquely localize and interact with several nuclear proteins

despite the absence of a canonical NLS, in contrast to non-oncogenic human polyomavirus ST proteins.

The nuclear localization of MCPyV ST was found to be responsible for many, but not all, MCPyV ST

transforming properties, suggesting MCPyV ST binds to and perturbs many pathways in both the

cytoplasm and nucleus to accomplish cellular transformation. Together, these data further increase our

understanding of MCPyV ST-mediated cellular transformation and may prove influential in further

discoveries of MCPyV ST-mediated oncogenesis and the consequent development of MCPyV-targeted

MCC therapies.

METHODS

Plasmids and Mutagenesis

MCPyV ST and GFP were subcloned into pLENTI-puro (Addgene #39481) for lentiviral

transductions in human foreskin fibroblasts (HFF). Furthermore, codon-optimized constructs

with a Kozak sequence were subcloned into pLENTI-puro to create MCPyV ST ko/co, TSPyV

ST ko/co, and HPyV7 ST ko/co, which were also transduced into Rat-2 cells. pLENTI-puro

empty and pLENTI-puro MCPyV LT-t were also transduced into Rat-2 cells. The Q5 NEB

Mutagenesis Kit (New England Biolabs) was used to add an NES (LQKKLEELEL) or NLS

(SPKKKRKVE) followed by a $(GGGGS)_2$ flexible linker to the 5' end (N-terminus) of pLENTI-

puro MCPyV ST ko/co to create pLENTI-puro NES-MCPyV ST ko/co and pLENTI-puro NLS-

MCPyV ST ko/co. pMTBS (donated by Dr. Christopher Buck, NIH) was used to transfect 293A

cells with MCPyV ST. TSPyV ST and HPyV 7 ST and were subcloned into pCS2 to transfect

293A cells. Empty pLENTI and pCS2 were used as negative controls.

Tissue Culture, Transductions, and Transfections

Adenovirus-transformed human embryonic kidney cells (HEK293A), human foreskin

fibroblasts (HFF), 293TNs, and Rat-2 cells were cultured in 1x DMEM (ThermoFisher

Scientific) supplemented with 10% fetal bovine serum (FBS) (Neuromics), GlutaMax

(ThermoFisher Scientific), Non-Essential Amino Acids (NEAA) (ThermoFisher Scientific), and

100 units/mL penicillin-streptomycin (ThermoFisher Scientific). MCC virus-positive MKL-1

cells were cultured in 1x RPMI (ThermoFisher Scientific) supplemented similarly as stated

above. All cell lines were incubated at 37°C in 5% CO_2.

293TN cells were transfected with the lentiviral envelope and packaging plasmids

psPAX2 (Addgene #12260) and pMD2.G (Addgene #12259) and lentiviral transfer plasmid

pLENTI-puro using TransIT-293 Transfection reagent (Mirus). Viral harvests were conducted at

24 and 48 hours post-transfection. Rat-2 or HFFs were transduced with the pseudotyped

lentiviruses supplemented with 4μg/mL polybrene (SantaCruz Biotechnology) and selected with

puromycin (ThermoFisher Scientific) 48 hours post-transduction at 1.5μg/mL or 4μg/mL for

HFFs and Rat-2 cells, respectively.

HEK293 were transfected with pMTBS and pCS2-ST at ~80% confluence using

TransIT-293 transfection reagent (Mirus). Cells were harvested for downstream applications 48

hours post-transfection.

Immunoblotting and Antibodies

Cell lysates were prepared in RIPA lysis buffer (ThermoFisher Scientific) supplemented

with a protease inhibitor cocktail (ThermoFisher Scientific) on ice, followed by sonication at

20% amplitude and centrifugation at 16,000 x g for 15 minutes. A BCA assay (ThermoFisher

Scientific) was performed to determine protein concentrations for normalization in 2x SDS

sample buffer (ThermoFisher Scientific) at 1μg/μL followed by boiling at 95°C. 30μg of protein

was loaded in an 8-16% SDS-PAGE gel (ThermoFisher Scientific) and transferred to an

Immobilon-P PVDF membrane, followed by a 1 hour block in 4% milk and overnight primary

antibody incubation with Ab5 (donated by James Decaprio, Harvard University) diluted at

1:1000 for MCPyV ST, HPyV7 ST, and TSPyV ST, and MCPyV LT, HSP90 (Cell Signaling

Technologies, 4874S) (1:5000), HDAC2 (Cell Signaling Technologies, 2540S) (1:5000),

NA$^+$/K$^+$-ATPase (Cell Signaling Technologies, 3010S) (1:2500), or actin (Cell Signaling

Technologies, 5125S) (1:10,000). After TBST washes, membranes were incubated in 4% milk

supplemented with either anti-mouse IgG HRP-conjugated secondary antibody (Cell Signaling

Technologies, 7076P2) (1:10,000) or anti-rabbit IgG HRP-conjugated secondary antibody (Cell

Signaling Technologies, 7074P2) (1:10,000) at room temperature for 1 hour. Following

secondary washing, proteins were detected using Femto (ThermoFisher Scientific) and a

ChemiDoc Imaging System (Bio-Rad).

Transformation Assays

Soft agar assays were performed in triplicate according to the JOVE online protocol at

50,000 cells/well (33). The top layer of 1x DMEM was supplemented with puromycin. The cells

were monitored for three weeks, followed by imaging with a Nikon Ti2 Eclipse microscope

(Nikon).

Metabolism experiments were performed in triplicate using a Glucose Uptake-Glo Assay

Kit (Promega) according to manufacturer instructions at 10,000 cells/96-well.

Serum dependence experiments were performed by plating 50,000 cells/6-well in

triplicate in complete DMEM supplemented with either 10%, 5%, or 1% FBS (Neuromics). Cells

were quantified at days 2, 5, and 10. Puromycin-supplemented media was changed every 3 days

or as needed in later time points.

Proliferation rate experiments were conducted by plating 50,000 cells/6-well in triplicate.

Cells were quantified at days 2, 4, 8, 14, 20, and 28. Doubling time was evaluated between days

0 and 28 to be representative of the entire experiment. Puromycin-supplemented media was

changed every 3 days or as needed in later time points.

Subcellular Fractionation

Subcellular fractionations were conducted using a Subcellular Fractionation Kit

(ThermoFisher Scientific) following a manufacturer-modified protocol. In short, cells were

plated at 1×10^6 in 4 10cm plates 24 hours prior to harvest through trypsinization (ThermoFisher

Scientific) for adherent cells. Provided fractionation extraction buffers were supplemented with a

1x protease inhibitor cocktail and used to fractionate the cell sample as instructed with the

addition of extra wash steps to lessen fraction contamination. Following protein extraction,

lysates were quantified and run in a western blot as described previously. Controls included Heat

Shock Protein 90 (HSP90) for the cytoplasmic fraction, Sodium Potassium ATPase (NA^+/K^+-

ATPase) for the membrane fraction, and Histone Deacetylase 2 (HDAC2) for the nuclear soluble

fraction.

Mass Spectrometry

The mutant BirA biotin ligase was fused in frame on the N-terminus of MCPyV ST with

a 4xGS linker separating the proteins and a V5 epitope tag (TID-ST). TurboID was performed on

HFFs expressing TID-ST as well as HFFs expressing negative controls (i.e., a biotin ligase fused

GFP with an (NLS) to mimic the subcellular compartment of TID-ST (TID-GFP-NLS); and the

biotin ligase expressed alone (TID)). Streptavidin agarose slurry was reductively methylated

using a reductive alkylation kit (Hampton Research HR2-434). For each set of triplicates, 16

15cm HFF MCPyV ST or control plates at 90% confluency were incubated in DMEM with

500μM biotin for 30 minutes. After 30 minutes, cells were placed on ice and washed five times

with ice-cold PBS to terminate the biotinylation reaction. Cells were then scrapped and spun

down into cellular pellets that were resuspended in ice-cold lysis buffer and benzonase (Sigma),

and rotated for 30 minutes at room temperature. Lysates were then spun at 10,800rpm for 15

minutes in an ultracentrifuge using an SW41 rotor, before collecting the supernatant. Protein

concentrations were normalized to 6mg/ml. For each 1ml replicate, 125µl of equilibrated

reductively methylated streptavidin beads were added and rotated for 2 hours at room

temperature, followed by a series of 25 washes. Finally, the beads were resuspended in a

digestion buffer and submitted to proteomics for mass spectrometry. A more detailed protocol

will be shared upon request.

Statistical Methods

Statistical analyses were conducted using the Jamovi statistical computer software, Version 2.3.

All experiments were repeated in triplicate. One-way ANOVA analyses were used, and a *p-value*

<0.05 was considered to indicate a statistically significant difference.

RESULTS

The ST unique region is responsible for the dominant transforming properties of MCPyV ST

In order to investigate the mechanism(s) of MCPyV cellular transformation, we first sought to

assess various transforming capabilities of the MCPyV tumor antigens (TAgs) expressed in virus positive

MCC (VP-MCC): ST and LT-t. As done previously by others, Rat-2 cells were transduced with

pseudotyped lentiviruses containing plasmids encoding MCPyV ST, LT-t, or empty pLENTI control (34).

After confirmation of protein expression (Fig. 1A), soft agar assays were performed to assess the ability

of the MCPyV TAgs to induce anchorage-independent cellular proliferation, a common property of

cancerous cells that promotes tumor formation and metastasis (35). Consistent with the work of others,

MCPyV ST was capable of inducing anchorage-independent growth, whereas MCPyV LT-t was

incapable of colony formation (Fig. 1B and C) (34,36).

Because cellular transformation can be defined by properties other than anchorage-independent

growth, additional transformation assays were performed between the MCPyV TAgs to further define the

specific transforming properties of both MCPyV ST and LT-t. It has been frequently described that

cancerous cells have an increased metabolic rate and, therefore, glucose uptake than healthy cells to

support their robust cellular division (37,38). Consistently, MCPyV ST expression significantly increased

the glucose uptake of Rat-2 cells, whereas MCPyV LT-t did not (Fig. 1D). Similarly, MCPyV ST was

also capable of significantly increasing the proliferation rate and decreasing the doubling time of Rat-2

cells (Fig. 1E and F); however, these assays demonstrated that MCPyV LT-t was also capable of

significantly increasing the cellular proliferation rate and decreasing the doubling time compared to

control cells, albeit to a lesser degree than MCPyV ST. These data support the transforming properties of

both MCPyV ST and LT-t, consistent with the necessity of both ST and LT-t to be expressed for the

viability of VP-MCC cells.

Finally, the serum dependence of Rat-2 cells was compared between control, ST, and LT-t

expressing cells, as cancerous cells are frequently described as capable of maintaining cellular

proliferation in the absence of the growth factors and nutrients found in serum (39). Consistent with

proliferation rate experiments containing the normal 10% serum supplementation (Fig. 1E), both ST and

LT-t were capable of significantly increasing proliferation in 10% serum; however, MCPyV LT-t

exhibited less proliferation than MCPyV ST with reduced serum concentrations of either 5% or 1% (Fig.

1G). To illustrate the effect of reducing serum concentrations on the proliferation of the control, ST, and

LT-t expressing cells, the percent change in doubling time was calculated going from 10% to 1%, 10% to

5%, and 5% to 1% serum. Although the percent change in doubling time between these different serum

concentrations was not significantly different between control, ST, and LT-t expressing cells when going

from 10% to 1% serum, normalization by glucose uptake (Fig. 1D) found MCPyV ST to significantly

decrease the percent change in doubling time upon serum reductions to 1% (Fig. 1H). Together, these

data found serum concentrations to affect the proliferation of ST-expressing cells the least, despite these

cells having the highest metabolic rate as inferred by glucose uptake.

Upon confirmation of ST as the dominant transforming protein of MCPyV, we sought to

elucidate the domain(s) responsible for this phenotype. Both MCPyV ST and LT-t share the same start

codon and are created through the alternative splicing of the same mRNA (11). Therefore, both ST and

LT-t share the same N-terminal amino acid sequence, referred to as "common-T", whereas they differ in

their C-terminal amino acid sequence, which is referred to as the "ST- and LT – unique regions" (Fig. 1I).

The finding of ST to be the dominant transforming protein of MCPyV led to the hypothesis that the

domain(s) responsible for these unique transforming properties are likely located in the ST-unique region,

as location within the common-T region would suggest MCPyV LT-t to have similar transforming

properties. Together, these data confirm ST as the dominant transforming protein of MCPyV by

assessment of various transformation properties such as anchorage-independent growth, metabolism,

proliferation rate, doubling time, and serum dependence, and that the domain(s) responsible for this

phenotype are likely found in the ST unique region.

The structurally dissimilar loops in the ST unique region of MCPyV ST may be responsible

for the uniquely transforming capabilities of MCPyV ST when compared to other skin-tropic

human polyomaviruses

Upon confirmation of ST being the dominant transforming protein of MCPyV, likely a

result of activities in the ST unique region, we sought to further elucidate the region(s) of

MCPyV ST responsible for various transformation properties. As noted, MCPyV ST belongs to a

family of 14 known human polyomaviruses, but is the only polyomavirus found to be directly

responsible for the development of cancer (20,21). Therefore, we sought to determine whether

the ST antigens of other, skin tropic, human polyomaviruses, such as Human Polyomavirus 7

(HPyV7) and Trichodysplasia Spinulosa Polyomavirus (TSPyV), were also capable of any of the

mechanisms of cellular transformation similar to MCPyV ST. Rat-2 cells were transduced to

express MCPyV, TSPyV, or HPyV7 ST (Fig. 2A). Although it appeared that both HPyV7 and

TSPyV ST proteins were expressed in lower quantities than MCPyV ST, it is likely that the

antibody used to detect these proteins has higher affinity to MCPyV ST, as it was developed

through its recognition of the MCPyV ST antigen; however, each T antigen was similarly codon

optimized and under the control of the same CMV promoter.

Consistent with TSPyV and HPyV7 not being found to be associated with cancer,

expression of HPyV7 and TSPyV ST proteins in Rat-2 cells failed to significantly induce

anchorage-independent growth above control cells (Fig. 2B and C). Furthermore, the metabolic

rate, as measured by glucose uptake, of HPyV7 and TSPyV ST were lower than that of MCPyV

ST (Fig. 2D). Interestingly, although the proliferation rate, doubling time, and serum dependence

of HPyV7 ST were more similar to control cells, TSPyV ST induced a cellular proliferation rate,

doubling time, and serum dependence similar to that of MCPyV ST, further demonstrating that

transformation is a combination of many cellular abnormalities that are not necessarily connected

(Fig. 2E, F, G, and H).

Together, our results indicate that MCPyV ST can uniquely induce anchorage-

independent growth, increase glucose uptake and proliferation rate, and reduce doubling time

and serum dependence when compared to HPyV7 ST; however, although TSPyV ST was

incapable of inducing anchorage-independent growth and increased glucose uptake, it was able

to increase cellular proliferation and reduce doubling time and serum dependence. In an effort to

determine the domain(s) responsible for the unique transforming capabilities of MCPyV ST, an

amino acid alignment of MCPyV, TSPyV, and HPyV7 ST was performed (Supplementary

Figure 1). Unfortunately, although all evolutionarily related ST proteins, they contained

exceptionally different amino acid sequences that made it extremely difficult to identify MCPyV

ST unique domains possibly responsible for transformation. Despite dissimilar amino acid

sequences, MCPyV, TSPyV, and HPyV7 ST proteins were found to have very similar protein

structures upon protein structure alignment (Fig. 2I). Consistent with the common T region being

hypothesized to not contain the domain(s) necessary for MCPyV ST transformation (Fig. 1I), the

common-T structure of the transforming MCPyV ST, and non-transforming TSPyV and HPyV7

ST were predicted to be very similar. However, despite their similarity, MCPyV, TSPyV, and

HPyV7 ST are predicted to contain structurally dissimilar loops in the ST-unique region,

consistent with the previous hypothesis of the transforming domains being found in the ST-

unique region (Fig. 1I). Together, these data suggest that the unique transforming capabilities of

MCPyV ST may be a result of the activities of the structurally dissimilar loops within the unique

region.

MCPyV ST binds to many nuclear cellular proteins

To identify the cellular proteins and pathways that MCPyV ST perturbs to accomplish its

dominant and unique cellular transformation, Turbo-ID Mass Spectrometry was performed on

human foreskin fibroblasts (HFF) expressing MCPyV ST or GFP control (Fig. 3A). Out of the

104 significant potential MCPyV ST cellular interactors, 39 (37.5%) were exclusively

cytoplasmic proteins, 30 (28.8%) were exclusively nuclear, and 35 (33.6%) were proteins with

both cytoplasmic and nuclear localization (Fig. 3A and B). These findings were unexpected, as

the amino acid sequence of MCPyV ST does not contain a known, canonical NLS, nor do many

cellular localization programs such as PSORTII, seqNLS, and BaCelLO predict MCPyV ST to

exhibit nuclear localization (Fig. 3C and D). Although MCPyV LT has been found to localize to

the nucleus as a result of its NLS, the NLS of LT is found in the LT unique region, and mutation

of the common-T region of LT did not affect its nuclear localization (Fig. 3E) (40). Therefore, in

addition to transformation, the potential nuclear localization of MCPyV ST may also be due to

the activity of a domain found in the ST unique region.

MCPyV ST uniquely localizes to the nucleus independent of size

In order to determine whether MCPyV ST can localize to the nucleus in the absence of a

canonical NLS, subcellular fractionations (SCF) were performed. Rat-2 cells transduced with

pLENTI MCPyV ST were fractionated into cytoplasmic, nuclear, and membrane fractions, with

MCPyV ST being found to localize to both the cytoplasmic and nuclear subcellular

compartments (Fig. 4A). Although a small amount of MCPyV ST protein was detected in the

membrane fraction, it is consistent with the level of HSP90 cytoplasmic contamination within the

membrane fraction, and is likely a false signal. To confirm this finding in another cell line,

293As transfected with pMTBS also identified MCPyV ST to localize to the cytoplasmic and

nuclear fractions, with membrane localization again likely a result of cytoplasmic contamination

(Fig. 4B). Of note, it was consistently found that SCF experiments in 293A cells frequently led

to more contamination than Rat-2 cells; however, the results from both cell lines confirmed both

MCPyV ST cytoplasmic and nuclear localization.

To confirm these findings in a physiologically relevant cell line, an SCF was performed

on MKL-1 cells, a virus-positive MCC cell line. Interestingly, in a VP-MCC cell line,

endogenous MCPyV ST localized exclusively to the nucleus (Fig. 4C). This finding was

consistent with exogenous MCPyV ST nuclear localization in Rat-2 and 293A cells, but

inconsistent with shared cytoplasmic localization.

As shown previously, MCPyV ST is uniquely transforming when compared to other

human, skin-tropic polyomaviruses such as TSPyV and HPyV7 ST (Fig. 2). To determine

whether MCPyV ST is also unique in its ability to localize to the nucleus despite the absence of a

canonical NLS, SCFs were performed on Rat-2 and 293A cells transfected with pCS2 TSPyV ST

or pCS2 HPyV7 ST. Interestingly, both TSPyV ST (Fig. 4D and E) and HPyV7 ST (Fig. 4F and

G) exclusively localized to the cytoplasm in both Rat-2 and 293A cells. These data suggest that

MCPyV ST is unique in both its transforming capacity and nuclear localization. Furthermore, the

domain(s) responsible for the unique transformation and nuclear localization of MCPyV ST may

be found in the structurally dissimilar loops of the ST unique region (Fig. 2I).

MCPyV ST nuclear localization is important for anchorage-independent growth, increased

metabolism, increased proliferation, and decreased serum dependence of MCPyV ST, but is

expendable for doubling time

In order to determine whether nuclear localization of MCPyV ST is necessary for cellular

transformation, both a cytoplasmic and nuclear sequestered MCPyV ST were created by adding

the NLS from SV40, or the nuclear export signal (NES) from Mitogen-Activated Protein Kinase

Kinase (MAPKK), to MCPyV ST (NLS-ST and NES-ST, respectively). Importantly, localization

domains were added to the N-terminus of MCPyV ST to ensure the furthest distance from the

likely transforming domains within the structurally dissimilar loops of the MCPyV ST unique

region in the C-terminus (Fig. 1I and 2I). Furthermore, localization domains were added with the

addition of a flexible linker to decrease the likelihood of the added domain interfering with the

structure and/or functions of MCPyV ST. To ensure the expression and stability of the MCPyV

ST localization mutants, a confirmation western blot was performed and found the localization

mutants and wild-type MCPyV (WT-ST) to all be expressed; however, protein densitometry

revealed NLS-ST and NES-ST to be expressed at 41% and 59% of WT-ST, respectively. SCF of

Rat-2 cells confirmed the appropriate localization of WT-ST and the localization mutants (Fig.

5B). Similar to WT-ST, NLS-ST retained localization to both the nucleus and the cytoplasm,

with a higher prevalence of nuclear localization. More importantly, NES-ST localized

exclusively to the cytoplasm, necessary to assess the importance of nuclear localization in

MCPyV ST-mediated cellular transformation.

To assess the effect of localization on MCPyV ST-mediated cellular transformation,

various transformation assays were performed comparing WT-ST, NLS-ST, and NES-ST,

followed by normalization based on protein densitometry (Fig. 5A). NLS-ST was capable of

significantly inducing colony formation above WT-ST post-normalization, whereas NES-ST also

led to colony formation, but to a lesser extent (Fig. 5 C and D). Normalized glucose uptake of

NLS-ST was significantly higher than that of WT-ST, whereas the glucose uptake of NES-ST

was similar to that of WT-ST, suggesting nuclear localization of MCPyV ST may influence the

ability of MCPyV ST to increase cellular metabolism (Fig. 5E). The proliferation curves of WT-

ST and both localization mutants were significantly increased when compared to the empty

pLENTI transduced controls; however NLS-ST was not significantly different from WT-ST,

whereas NES-ST had a proliferation rater significantly less than WT-ST (Fig. 5F). However, the

protein-level normalized doubling times of NLS-ST and NES-ST were both significantly faster

than WT-ST, with NLS-ST having the fastest doubling time (Fig. 5G).

No significant differences were observed when comparing the proliferation rate curves of

WT-ST compared to the localization mutants at differing serum concentrations, likely as a result

of the differing protein levels and metabolic rates of each ST (Fig. 5H). Therefore, the percent

change in doubling time between differing serum concentrations was calculated and normalized

by both the protein levels (Fig. 5A) and glucose uptake rates (Fig. 5E) of each ST (Fig. 5I).

Together, WT-ST and NES-ST were each found to have similar changes in the percent doubling

time between 10% and 1% serum containing media, whereas NLS-ST was found to have a

significantly lower percent change in the doubling time between 10% and 1% serum-containing

media. Together, these data suggest nuclear localization of ST is largely responsible for the

ability of MCPyV ST to proliferate in the absence of serum despite having an increased

metabolic rate, as inferred by an increased glucose uptake. In conclusion, these data suggest that

doubling time may be a combined result of both nuclear and cytoplasmic localization, whereas

increased anchorage-independent growth, metabolism, proliferation, and decreased serum

dependence may be influenced by the nuclear localization of MCPyV ST.

DISCUSSION

The mechanism(s) of MCPyV-mediated cellular transformation and tumorigenesis have

been investigated since the discovery of MCPyV in 2008; however, many of these early analyses

were hindered by using the approach of comparing the similarities between the human oncogenic

polyomavirus MCPyV, and the simian oncogenic virus SV40. The findings of the current study

and of others that ST is the dominant transforming protein of MCPyV is opposite to SV40, in

which ST plays only a supportive or accessory role, with LT being the dominant transforming

protein (41). Similarity comparisons between the dissimilar SV40 and MCPyV are responsible

for many years spent investigating MCPyV ST binding to PP2A and Fbw7, which have been

found to be irrelevant for transformation by MCPyV ST (42,43). Furthermore, we found MCPyV

ST to be uniquely transforming among skin-tropic human polyomavirus ST antigens, consistent

with MCPyV being the only human polyomavirus clearly associated with cancer. Together, these

data refute the method of identifying mechanisms of MCPyV transformation through similarity

comparisons to SV40, and alternatively support the dissimilarity approach of investigating the

differences between MCPyV ST and the ST of other non-oncogenic human polyomaviruses to

identify the unique mechanisms of MCPyV ST mediated transformation.

The dissimilarity approach can also be applied when comparing MCPyV ST and LT-t, as

they share a common T region, have their own unique regions, and accomplish dissimilar

patterns of transformation. Through this approach, we were able to narrow down the location of

the domain(s) responsible for MCPyV ST transformation to the ST unique region. Although

transformation assays performed on Rat-2 cells expressing the ST unique region alone may be

able to confirm this finding, it is likely that the absence of common-T may influence the

structure of the ST unique region, and it is possible that domain(s) within the common-T region

may be necessary to facilitate the activities of the ST unique region.

Although the discovery that MCPyV is uniquely oncogenic among human

polyomaviruses was discouraging at the genesis of MCPyV oncogenesis research, the

dissimilarity approach has exploited it for our benefit. Comprehensive transformation assay

analysis of TSPyV ST and HPyV7 ST identified that although the ST of HPyV7 and TSPyV

were not as robustly transforming as MCPyV ST, TSPyV ST was capable of achieving some

properties of cellular transformation, further supporting the fact that cellular transformation is a

collection of properties rather than a single entity, and that the ability to induce several

transforming properties may translate to being able to accomplish tumorigenesis *in vivo.*

Furthermore, structural alignments allowed us to further narrow down the unique transforming

domains of MCPyV ST to the structurally dissimilar loops consistent with the unique region of

MCPyV ST. It is currently hypothesized that the similar transforming capabilities of MCPyV and

TSPyV ST may reside in domains shared by MCPyV and TSPyV ST, but not found in HPyV7

ST. Alternatively, TSPyV ST and MCPyV ST may accomplish similar methods of cellular

transformation through distinct mechanisms.

The identification of MCPyV ST potentially binding to several nuclear proteins

suggested nuclear localization, despite the absence of a canonical NLS in MCPyV ST.

Surprisingly, subcellular fractionation confirmed nuclear localization of MCPyV ST in various

cell lines, whereas the non-oncogenic TSPyV and HPyV7 ST proteins were confined to the

cytoplasm, further proving the usefulness of the dissimilarity hypothesis. Although MCPyV ST

is under the 30kDa size necessary for passive diffusion through the nuclear pore, the finding that

the 30kDa TSPyV and HPyV7 ST proteins are cytoplasmic confirmed that MCPyV ST

localization to the nucleus is regulated (44). Interestingly, MCPyV ST exhibited both nuclear and

cytoplasmic localization in transduced Rat-2 and 293A cells, yet it was exclusively nuclear in

VP-MCC cells. Although it is possible that the differences in MCPyV ST localization between

the cell lines may be an artifact of differing cell lines, hypotheses to explain these differences

include the following: First, it is possible that the mechanism of MCPyV ST nuclear

translocation is saturated in Rat-2 and 293A cells overexpressing exogenous ST, leading to

spillover into the cytoplasm. Second, the presence of MCPyV LT-t in MKL-1 cells, but not Rat-2

and 293A cells, may enhance the efficiency of MCPyV ST nuclear localization. Although both

hypotheses require further experimentation, it is clear from these experiments that MCPyV ST is

capable of uniquely localizing to the nucleus despite the absence of a canonical NLS.

It can be hypothesized that the noncanonical NLS or domain responsible for MCPyV ST

nuclear localization is found within the MCPyV ST unique region, as previous investigation into

479 MCPyV LT-t nuclear localization found the common T region to not be responsible for nuclear

480 localization (40). As the function of an NLS is dependent on both structure and function, it is

481 possible that the MCPyV ST NLS is located in the highly dissimilar amino acid sequence or the

482 structurally dissimilar loops between MCPyV, TSPyV, and HPyV7 ST.

483   Although most of the attention regarding MCPyV-mediated oncogenesis has been

484 directed towards ST, it is clear that the large T antigen is also important due to its necessity for

485 the viability of MCC cells and its ability to influence cellular proliferation and doubling time,

486 despite not inducing anchorage-independent growth, increased metabolism, and decreased serum

487 dependence. Due to the sharing of the common-T region between LT and ST, the domain(s)

488 responsible for increasing proliferation rate and reducing doubling time may be found in the

489 common T region; however, it is more likely that ST and LT-t accomplish this through different

490 mechanisms attributed to their common-T region, as they affected proliferation and doubling

491 time of Rat-2 cells dissimilarly, despite similar protein expression levels. It is likely that

492 perturbation of Rb by the Rb binding domain of MCPyV LT-t contributes to these phenotypes;

493 however, this has not yet been directly assessed in these assays.

494   Finally, the export of MCPyV ST to the cytoplasm did not ablate all properties of cellular

495 transformation, suggesting a potential role of MCPyV in both the cytoplasm and nucleus.

496 However, it is not currently understood how this finding translates to the observation of MCPyV

497 ST being exclusively nuclear in VP-MCC cells, and therefore warrants further investigation.

498 Furthermore, although normalized NES-ST was still capable of anchorage-independent growth,

499 increased glucose uptake, and increased proliferation rate, NLS-ST normalization led to a

500 significant difference in anchorage-independent growth, glucose uptake, proliferation rate, and

501 serum dependence when compared to both WT-ST and NES-ST. Although protein levels of WT-

ST and NLS-ST differed, the ratio of WT-ST and NLS-ST in the cytoplasmic and nuclear

compartments were similar. Therefore, it is of great interest why NLS-ST was capable of

increasing transformation above WT-ST in some assays but not others.

CONCLUSIONS

Together, it has been found that elucidating mechanisms of transformation guided by a

dissimilarity approach may prove useful in identifying otherwise hidden, unique mechanisms of

MCPyV oncogenesis, such as the unique nuclear localization of MCPyV ST. Cellular

transformation was found to be multi-variable and not simply a positive or negative

characteristic of viral proteins, consistent with our understanding of the role of viral proteins in

viral replication. Further investigation into the mechanisms and role of MCPyV nuclear

localization will further enhance our understanding of the natural viral life cycle and MCPyV-

mediated oncogenesis necessary for basic virology and the treatment of MCC.

525 **LIST OF ABBREVIATIONS**

527	MCC	Merkel Cell Carcinoma
528	MCPyV	Merkel Cell Polyomavirus
529	HPyV7	Human Polyomavirus 7
530	TSPyV	Trichodysplasia Spinulosa Polyomavirus
531	HPyV	Human Polyomavirus
532	NES	Nuclear Export Signal
533	NLS	Nuclear Localization Signal
534	TAg	Tumor Antigens
535	ST	Small Tumor Antigen
536	LT-t	Truncated Large Tumor Antigen
537	ALTO	Alternate Large T Open Reading Frame
538	ER	Early Region
539	LR	Late Region
540	VP-MCC	Virus Positive Merkel Cell Carcinoma
541	FBS	Fetal Bovine Serum
542	SCF	Subcellular Fractionation
543	SAA	Soft Agar Assay
544	SV40	Simian Virus 40

DECLARATIONS

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article and

supplementary materials.

Competing interests

The authors declare no conflict of interest.

Funding

Financial support was provided by the Stetson University Health Sciences Department and Cell

Biology Education Consortium.

Author Contributions

Kaira R. Thevenin, Isabella S. Tieche, and Cody E. Di Benedetto facilitated conceptualization,

investigation, data curation, writing of the original draft, review and editing, and visualization.

Matt Schrager performed formal analysis and review and editing. Kristine N. Dye performed

conceptualization, methodology, validation, investigation, writing of original draft and review

and editing, visualization, supervision, and funding acquisition.

Acknowledgments

We would like to thank Dr. Denise Galloway (University of Washington) for project

conceptualization and Dr. John Lee (University of Washington) for facilitating Turbo-ID mass

spectrometry experiments. Furthermore, we would also like to thank Richard Wang (University

of Texas) for sharing Rat-2 cells, Dr. Patrick Moore and Dr. Yuan Chang (University of

Pittsburgh) for sharing MKL-1 cells, and Dr. James DeCaprio (Harvard) for donating Ab5.

FIGURE LEGENDS

Figure 1. The ST unique region is responsible for the dominant transforming properties of

MCPyV ST. *Rat-2 cells were transduced with MCPyV LT-t, MCPyV ST, or pLENTI control and*

protein expression was confirmed (A). Various transformation assays were performed including

soft agar assays (B and C), glucose uptake (D), proliferation rate (E), doubling time (F), and

serum dependence (G) (grey – pLENTI, coral – MCPyV ST, blue – MCPyV LT-t). The percent

change in serum dependence doubling time (solid bars) was normalized (dashed bars) by

glucose uptake compared to pLENTI control (H). MCPyV ST protein structure was determined

by PyMol and colored to show the common-T (blue) and ST unique (pink) regions (I). Each point

*represents mean ± the standard error of independent triplicates. One-way ANOVA * p < 0.05; ***

< 0.01; *** p < 0.001, significantly different from pLENTI.

Figure 2. The structurally dissimilar loops in the ST unique region of MCPyV ST may be

responsible for the uniquely transforming capabilities of MCPyV ST when compared to other

skin-tropic human polyomaviruses. *Rat-2 cells were transduced with MCPyV ST, HPyV7 ST,*

TSPyV ST, or pLENTI control, and protein expression was confirmed (A). Various

transformation assays were performed, including soft agar assays (B and C), glucose uptake (D),

proliferation rate (E), doubling time (F), and serum dependence (G) (grey – LENTI, coral –

MCPyV ST, purple – HPyV7 ST, yellow – TSPyV ST). The percent change in serum dependence

doubling time (solid bars) was normalized (dashed bars) by glucose uptake compared to pLENTI

control (H). MCPyV ST, HPyV7 ST, and TSPyV ST protein structures were determined and

aligned using PyMol and black arrows denote the structurally dissimilar regions in protein

structure (I). Each point represents mean ± the standard error of independent triplicates. One-

way ANOVA * p < 0.05; ** < 0.01; *** p < 0.001, significantly different from pLENTI.

***Figure 3. MCPyV ST binds to many nuclear cellular proteins.** Human foreskin fibroblasts*

(HFFs) were transduced with MCPyV ST or GFP, followed by TurboID Mass Spectrometry

analysis. A total of 121 proteins were enriched in ST expressing HFFs defined as a False

Discovery Rate (FDR) <0.01 and Fold Change (FC) > 2. These 121 proteins were colored based

on their subcellular localization, with teal representing nuclear proteins, pink representing

cytoplasmic proteins, and green representing proteins with both nuclear and cytoplasmic

localization (A). A Venn diagram was also included to visualize the number of MCPyV ST

potential interactors that are nuclear, cytoplasmic, or both nuclear and cytoplasmic (B). The

amino acid sequence of MCPyV ST contains no known canonical NLS (C). Online amino acid

and protein subcellular localization predictor tools were utilized to predict the localization of

MCPyV ST (D). The location of the MCPyV LT-t NLS is found within the MCPyV LT-t unique

region (orange), but no known NLS is found within the common-T region (red) or ST unique

region (blue) (E).

***Figure 4. MCPyV ST uniquely localizes to the nucleus independent of size.** Rat-2 (A) and 293A*

(B) cells were transduced and transfected, respectively, with MCPyV ST and a subcellular

fractionation was performed to isolate proteins for the cytoplasmic (C), membrane (M), and

nuclear (N) fractions. A subcellular fractionation for MCPyV ST was also performed on the VP-

MCC cells, MKL-1 (C). Rat-2 and 293A cells were also transduced and transfected, respectively,

with TSPyV ST (D and E) and HPyV7 ST (F and G), followed by subcellular fractionation.

HSP90 was used as a cytoplasmic control, NA⁺/K⁺-ATPase was used as a membrane control,

and HDAC2 was used as a nuclear control.

Figure 5. MCPyV ST nuclear localization is important for anchorage-independent growth,

increased metabolism, increased proliferation, and decreased serum dependence of MCPyV

ST, but is expendable for doubling time. *Rat-2 cells were transduced with MCPyV ST (ST-WT),*

a cytoplasmic sequestered MCPyV ST (NES-ST), a nuclear sequestered MCPyV ST (NLS-ST), or

pLENTI control, and protein expression was confirmed and quantified by protein densitometry

(A). A subcellular fractionation was performed to verify the localization of the wildtype ST and

localization mutants. Subcellular fractionations included cytoplasmic (C), membrane (M), and

nuclear (N) fractions, with HSP90 acting as a cytoplasmic control, NA⁺/K⁺-ATPase as a

membrane control, and HDAC2 as a nuclear control (B). Various transformation assays were

performed, including soft agar assays (C and D), glucose uptake (E), proliferation rate (F),

doubling time (G), and serum dependence (H) (grey – pLENTI control, coral – MCPyV ST,

maroon – NES-ST, teal – NLS-ST). The number of soft agar colonies (D), glucose uptake (E),

and doubling time (G) were normalized based on protein levels determined by densitometry

(solid bars – unnormalized, dashed bars – normalized) (A). The percent change in serum

dependence doubling time (solid bars) was normalized (dashed bars) by densitometry

determined protein levels and glucose uptake compared to wildtype MCPyV ST (I). Each point

*represents mean ± the standard error of independent triplicates. One-way ANOVA * $p < 0.05$; ***

*< 0.01; *** $p < 0.001$, significantly different from MCPyV ST.*

662
663
664
665

666 **SUPPLEMENTARY FIGURE LEGENDS**
667
668 ***Supplementary Figure 1. Amino acid alignment of MCPyV ST, HPyV7 ST, and TSPyV ST.*** *An*
669 *amino acid alignment was created using Clustal Omega for MCPyV ST (Sequence ID:*
670 *YP_009111422.1), HPyV7 ST (Sequence ID: ADE45476.1), and TSPyV ST (Sequence ID:*
671 *AID54936.1).*

REFERENCES

1. Toker C. Trabecular carcinoma of the skin. Arch Dermatol. 1972;105(1):107–10.

2. Becker JC. Merkel cell carcinoma. Ann Oncol. 2010;21(suppl_7):vii81–5.

3. Goh G, Walradt T, Markarov V, Blom A, Riaz N, Doumani R, et al. Mutational landscape of
MCPyV-positive and MCPyV-negative Merkel cell carcinomas with implications for
immunotherapy. Oncotarget. 2015;7(3):3403–15.

4. Feng H, Shuda M, Chang Y, Moore PS. Clonal Integration of a Polyomavirus in Human
Merkel Cell Carcinoma. Science. 2008;319(5866):1096–100.

5. Martel-Jantin C, Pedergnana V, Nicol JTJ, Leblond V, Trégouët DA, Tortevoye P, et al.
Merkel cell polyomavirus infection occurs during early childhood and is transmitted between
siblings. J Clin Virol. 2013;58(1):288–91.

6. Gossai A, Waterboer T, Nelson HH, Michel A, Willhauck-Fleckenstein M, Farzan SF, et al.
Seroepidemiology of Human Polyomaviruses in a US Population. Am J Epidemiology.
2016;183(1):61–9.

7. Zhang C, Liu F, He Z, Deng Q, Pan Y, Liu Y, et al. Seroprevalence of Merkel Cell
Polyomavirus in the General Rural Population of Anyang, China. PLoS ONE.
2014;9(9):e106430.

8. Šroller V, Hamšíková E, Ludvíková V, Vochozková P, Kojzarová M, Fraiberk M, et al.
Seroprevalence rates of BKV, JCV, and MCPyV polyomaviruses in the general Czech Republic
population. J Méd Virol. 2014;86(9):1560–8.

9. Tolstov YL, Knauer A, Chen JG, Kensler TW, Kingsley LA, Moore PS, et al. Asymptomatic
Primary Merkel Cell Polyomavirus Infection among Adults - Volume 17, Number 8—August
2011 - Emerging Infectious Diseases journal - CDC. Emerg Infect Dis. 2011;17(8):1371–80.

10. Shuda M, Feng H, Kwun HJ, Rosen ST, Gjoerup O, Moore PS, et al. T antigen mutations are
a human tumor-specific signature for Merkel cell polyomavirus. Proc Natl Acad Sci.
2008;105(42):16272–7.

11. DeCaprio JA. Merkel cell polyomavirus and Merkel cell carcinoma. Philos Trans R Soc B:
Biol Sci. 2017;372(1732):20160276.

12. Carter JJ, Daugherty MD, Qi X, Bheda-Malge A, Wipf GC, Robinson K, et al. Identification
of an overprinting gene in Merkel cell polyomavirus provides evolutionary insight into the birth
of viral genes. Proc Natl Acad Sci. 2013;110(31):12744–9.

13. Shuda M, Kwun HJ, Feng H, Chang Y, Moore PS. Human Merkel cell polyomavirus small T
antigen is an oncoprotein targeting the 4E-BP1 translation regulator. J Clin Investig.
2011;121(9):3623–34.

14. Sunshine JC, Jahchan NS, Sage J, Choi J. Are there multiple cells of origin of Merkel cell
carcinoma? Oncogene. 2018;37(11):1409–16.

15. Park DE, Cheng J, Berrios C, Montero J, Cortés-Cros M, Ferretti S, et al. Dual inhibition of
MDM2 and MDM4 in virus-positive Merkel cell carcinoma enhances the p53 response. Proc
Natl Acad Sci. 2019;116(3):1027–32.

16. Cheng J, Park DE, Berrios C, White EA, Arora R, Yoon R, et al. Merkel cell polyomavirus
recruits MYCL to the EP400 complex to promote oncogenesis. PLoS Pathog.
2017;13(10):e1006668.

17. Hesbacher S, Pfitzer L, Wiedorfer K, Angermeyer S, Borst A, Haferkamp S, et al. RB1 is the
crucial target of the Merkel cell polyomavirus Large T antigen in Merkel cell carcinoma cells.
Oncotarget. 2016;7(22):32956–68.

18. Houben R, Adam C, Baeurle A, Hesbacher S, Grimm J, Angermeyer S, et al. An intact
retinoblastoma protein-binding site in Merkel cell polyomavirus large T antigen is required for
promoting growth of Merkel cell carcinoma cells. Int J Cancer. 2012;130(4):847–56.

19. Schrama D, Hesbacher S, Angermeyer S, Schlosser A, Haferkamp S, Aue A, et al. Serine
220 phosphorylation of the Merkel cell polyomavirus large T antigen crucially supports growth
of Merkel cell carcinoma cells. Int J Cancer. 2016;138(5):1153–62.

20. Prado JCM, Monezi TA, Amorim AT, Lino V, Paladino A, Boccardo E. Human
polyomaviruses and cancer: an overview. Clinics. 2018;73(suppl 1):e558s.

21. DeCaprio JA, Garcea RL. A cornucopia of human polyomaviruses. Nat Rev Microbiol.
2013;11(4):264–76.

22. Eddy BE, Borman GS, Berkeley WH, Young RD. Tumors Induced in Hamsters by Injection
of Rhesus Monkey Kidney Cell Extracts. Proc Soc Exp Biol Med. 1961;107(1):191–7.

23. Eddy BE, Borman GS, Grubbs GE, Young RD. Identification of the oncogenic substance in
rhesus monkey kidney cell cultures as simian virus 40. Virology. 1962;17(1):65–75.

24. Shein HM, Enders JF. TRANSFORMATION INDUCED BY SIMIAN VIRUS 40 IN
HUMAN RENAL CELL CULTURES, I. MORPHOLOGY AND GROWTH
CHARACTERISTICS*. Proc Natl Acad Sci. 1962;48(7):1164–72.

25. Rabson AS, Kirschstein RL. Induction of Malignancy in vitro in Newborn Hamster Kidney
Tissue Infected with Simian Vacuolating Virus (SV40). Proc Soc Exp Biol Med.
1962;111(2):323–8.

26. Ahuja D, Sáenz-Robles MT, Pipas JM. SV40 large T antigen targets multiple cellular
pathways to elicit cellular transformation. Oncogene. 2005;24(52):7729–45.

27. Moore PS, Chang Y. Why do viruses cause cancer? Highlights of the first century of human
tumour virology. Nat Rev Cancer. 2010;10(12):878–89.

28. Butel JS. Viral carcinogenesis: revelation of molecular mechanisms and etiology of human
disease. Carcinogenesis. 2000;21(3):405–26.

29. Scott MS, Calafell SJ, Thomas DY, Hallett MT. Refining Protein Subcellular Localization.
PLoS Comput Biol. 2005;1(6):e66.

30. Hung MC, Link W. Protein localization in disease and therapy. J Cell Sci.
2011;124(20):3381–92.

31. Laurila K, Vihinen M. Prediction of disease-related mutations affecting protein localization.
BMC Genom. 2009;10(1):122.

32. Lu J, Wu T, Zhang B, Liu S, Song W, Qiao J, et al. Types of nuclear localization signals and
mechanisms of protein import into the nucleus. Cell Commun Signal. 2021;19(1):60.

33. Borowicz S, Scoyk MV, Avasarala S, Rathinam MKK, Tauler J, Bikkavilli RK, et al. The
Soft Agar Colony Formation Assay. J Vis Exp. 2014;(92):e51998.

34. Zhao J, Jia Y, Shen S, Kim J, Wang X, Lee E, et al. Merkel Cell Polyomavirus Small T
Antigen Activates Noncanonical NF-κB Signaling to Promote Tumorigenesis. Mol Cancer Res.
2020;18(11):1623–37.

35. Mori S, Chang JT, Andrechek ER, Matsumura N, Baba T, Yao G, et al. Anchorage-
independent cell growth signature identifies tumors with metastatic potential. Oncogene.
2009;28(31):2796–805.

36. Kwun HJ, Shuda M, Feng H, Camacho CJ, Moore PS, Chang Y. Merkel Cell Polyomavirus
Small T Antigen Controls Viral Replication and Oncoprotein Expression by Targeting the
Cellular Ubiquitin Ligase SCFFbw7. Cell Host Microbe. 2013;14(2):125–35.

37. Lin X, Xiao Z, Chen T, Liang SH, Guo H. Glucose Metabolism on Tumor Plasticity,
Diagnosis, and Treatment. Front Oncol. 2020;10:317.

38. Fadaka A, Ajiboye B, Ojo O, Adewale O, Olayide I, Emuowhochere R. Biology of glucose
metabolization in cancer cells. J Oncol Sci. 2017;3(2):45–51.

39. White EZ, Pennant NM, Carter JR, Hawsawi O, Odero-Marah V, Hinton CV. Serum
deprivation initiates adaptation and survival to oxidative stress in prostate cancer cells. Sci Rep.
2020;10(1):12505.

40. Nakamura T, Sato Y, Watanabe D, Ito H, Shimonohara N, Tsuji T, et al. Nuclear localization
of Merkel cell polyomavirus large T antigen in Merkel cell carcinoma. Virology.
2010;398(2):273–9.

41. Cheng J, DeCaprio JA, Fluck MM, Schaffhausen BS. Cellular transformation by Simian
Virus 40 and Murine Polyoma Virus T antigens. Semin Cancer Biol. 2009;19(4):218–28.

42. Kwun HJ, Shuda M, Camacho CJ, Gamper AM, Thant M, Chang Y, et al. Restricted Protein
Phosphatase 2A Targeting by Merkel Cell Polyomavirus Small T Antigen. J Virol.
2015;89(8):4191–200.

43. Dye KN, Welcker M, Clurman BE, Roman A, Galloway DA. Merkel cell polyomavirus
Tumor antigens expressed in Merkel cell carcinoma function independently of the ubiquitin
ligases Fbw7 and β-TrCP. PLoS Pathog. 2019;15(1):e1007543.

44. Timney BL, Raveh B, Mironska R, Trivedi JM, Kim SJ, Russel D, et al. Simple rules for
passive diffusion through the nuclear pore complex. J Cell Biol. 2016;215(1):57–76.

A)

B) Cytoplasmic — Nuclear

39 | 35 | 30

C) MDLVLNRKEREALCKLLEIAPNCYGNI
PLMKAAFKRSCLKHHPDKGGNPVIM
MELNTLWSKFQQNIHKLRSDFSMFD
EVSTKFPWEEYGTLKDYMQSGYNAR
FCRGPGCMLKQLRDSKCACISCKLSR
QHCSLKTLKQKNCLTWGECFCYQCFI
LWFGFPPTWESFDWWQKTLEETDY
CLLHLHLF

D)

	BaCelLO	Euk-mPloc 2.0	ESLPred	DeepLoc	PSORTII	seqNLS
Nucleus			✓	✓		
Cytosol	✓	✓			✓	✓

E)

NLS
277-280
(RKRK)

LT

ST

Common T 🟥
ST Unique Region 🟦
LT Unique Region 🟧