1

The Small Tumor Antigen of Merkel Cell Polyomavirus Accomplishes Cellular

Transformation by Uniquely Localizing to the Nucleus Despite the Absence of a Known

Nuclear Localization Signal

10 Kaira R. Thevenin [a], Isabella S. Tieche [a], Cody E. Di Benedetto [a], Matt Schrager [a], Kristine N. Dye [a b]

[a] Department of Health Sciences, Stetson University, DeLand, FL 32723, USA

[b] Department of Biology, Stetson University, DeLand, FL 32723, USA

Corresponding Author: Kristine N. Dye (kdye1@stetson.edu); 421 N Woodland Blvd,

DeLand, FL, 32723

ABSTRACT

Background: Merkel Cell Carcinoma (MCC) is an aggressive skin cancer that is three times deadlier than melanoma. In 2008, it was found that 80% of MCC cases are caused by the genomic integration of a novel polyomavirus, Merkel Cell Polyomavirus (MCPyV), and the expression of its small and truncated large tumor antigens (ST and LT-t, respectively). MCPyV belongs to a family of human polyomaviruses; however, it is the only one with a clear association to cancer.

Methods: To investigate the role and mechanisms of various polyomavirus tumor antigens in cellular transformation, Rat-2, 293A, and human foreskin fibroblasts were transduced with pLENTI MCPyV LT-t, MCPyV ST, TSPyV ST, HPyV7 ST, or empty pLENTI and assessed through multiple transformation assays, and subcellular fractionations. One-way ANOVA tests were used to assess statistical significance.

Results: Soft agar, proliferation, doubling time, glucose uptake, and serum dependence assays confirmed ST to be the dominant transforming protein of MCPyV. Furthermore, it was found that MCPyV ST is uniquely transforming, as the ST antigens of other non-oncogenic human polyomaviruses such as Trichodysplasia Spinulosa Polyomavirus (TSPyV) and Human Polyomavirus 7 (HPyV7) were not transforming when similarly assessed. Identification of structural dissimilarities between transforming and non-transforming tumor antigens revealed that the uniquely transforming domain(s) of MCPyV ST are likely located within the structurally dissimilar loops of the MCPyV ST unique region. Of all known MCPyV ST cellular interactors, 62% are exclusively or transiently nuclear, suggesting that MCPyV ST localizes to the nucleus despite the absence of a canonical nuclear localization signal. Indeed, subcellular fractionations confirmed that MCPyV ST could achieve nuclear localization through a currently unknown, regulated mechanism independent of its small size, as HPyV7 and TSPyV ST proteins were incapable of nuclear translocation. Although nuclear localization was found to be important for several transforming properties of MCPyV ST, some properties were also performed by a cytoplasmic

51   sequestered MCPyV ST, suggesting that MCPyV ST may perform different transforming functions in

52   individual subcellular compartments.

54   **Conclusions:** Together, these data further elucidate the unique differences between MCPyV ST and other

55   polyomavirus ST proteins necessary to understand MCPyV as the only known human oncogenic

56   polyomavirus.

65   **Keywords:** Merkel Cell Polyomavirus, Merkel Cell Carcinoma, Cellular Transformation,

66   Trichodysplasia Spinulosa Polyomavirus, Human Polyomavirus 7, Nuclear Localization

BACKGROUND

Merkel Cell Carcinoma (MCC) was first described in the 1970s as a rare, aggressive skin cancer, likely a consequence of ultraviolet radiation-induced mutations in cell cycle-regulation genes (1–3). However, in 2008, a novel human polyomavirus, subsequently termed Merkel Cell Polyomavirus (MCPyV), was found to be integrated within, and the etiologic agent of, ~80% of MCC tumors (4). Further investigation identified MCPyV as a ubiquitous, asymptomatic human virus that, in rare cases of immunosuppressed individuals, could integrate into the human genome and lead to the development of cancer (5–9).

The small, 5.3kb dsDNA genome of MCPyV is divided into an early region and a late region (ER and LR, respectively) by a bidirectional promoter (4,10). Due to its constrained genomic size, the ER utilizes alternative splicing and overprinting to generate 4 viral proteins, including the large tumor antigen (LT), small tumor antigen (ST), alternate large T open reading frame (ALTO), and the 57kT protein, while the LR encodes the VP1 and VP2 structural proteins (10–12). Although all of these viral proteins are expressed for successful viral replication in the normal MCPyV viral lifecycle, in MCC, only a truncated LT (LT-t) and ST are expressed and necessary for the viability of MCC tumors, suggesting a role for these proteins in tumorigenesis and maintenance (4,13,14). Several researchers have identified transforming capabilities of both ST and LT-t, including, but not limited to, perturbation of MYCL and Rb, respectively (10,15–19).

MCPyV belongs to a family of 14 known human polyomaviruses but is currently the only one with a clear association to cancer (20,21). However, polyomaviruses of other species, such as Simian Virus 40 (SV40), which infects simian species, has been found to be oncogenic in rodents (22–25). As one of the most intensively studied oncogenic viruses, the study of SV40 oncogenesis has led to the development of several transformation assays, which have subsequently been used to direct the study of MCPyV transformation and tumorigenesis.

Transformation is defined as the acquisition of expanded proliferation and/or survival potential of a cell (26). Many *in vitro* transformation assays exist, including soft agar assays, focus formation,

doubling time, proliferation rate, serum dependence, and metabolic assays, each of which assesses

different aspects and pathways involved in cellular transformation that may collectively contribute to

tumorigenesis *in vivo.* In order to accomplish cellular transformation, viral proteins bind and perturb

cellular pathways that allow for expanded proliferation and survival potential (27,28). Cellular proteins

and pathways operate in specific subcellular compartments, including the cytoplasm, nucleus, and

membrane (29,30). As localization influences the regulation and activity of specific proteins and

pathways, it is tightly regulated (30,31). Therefore, protein localization to the nucleus is tightly regulated

and restricted to proteins containing a nuclear localization signal (NLS), as this is where the DNA is

contained, and gene expression is initiated (32).

Herein, we employed an alternative approach to elucidate the mechanism of transformation of

MCPyV T antigens. As opposed to previous approaches of comparing similarities between SV40 and

MCPyV T antigens, several transformation assays on various oncogenic and non-oncogenic human

polyomavirus tumor antigens were performed to identify the unique domains and mechanisms of MCPyV

T-antigen mediated oncogenesis. Together, we identified MCPyV ST to be predominantly and uniquely

transforming when compared to MCPyV LT-t and the ST of other, skin-tropic human polyomaviruses.

Furthermore, MCPyV ST was found to uniquely localize to the nucleus, consistent with its known

interaction with cellular nuclear proteins, despite the absence of a canonical NLS. This is in contrast to

non-oncogenic human polyomavirus ST proteins, which localize exclusively to the cytoplasm. The

nuclear localization of MCPyV ST was found to be responsible for many, but not all, MCPyV ST

transforming properties, suggesting MCPyV ST binds to and perturbs many pathways in both the

cytoplasm and nucleus to accomplish cellular transformation. Together, these data further increase our

understanding of MCPyV ST-mediated cellular transformation and may prove influential in further

discoveries of MCPyV ST-mediated oncogenesis and the consequent development of MCPyV-targeted

MCC therapies.

METHODS

Plasmids and Mutagenesis

Codon-optimized constructs with a Kozak sequence were subcloned into pLENTI-puro (Addgene #39481) to create MCPyV ST ko/co, TSPyV ST ko/co, and HPyV7 ST ko/co, which were transduced into Rat-2 cells. pLENTI-puro empty and pLENTI-puro MCPyV LT-t were also transduced into Rat-2 cells. The Q5 NEB Mutagenesis Kit (New England Biolabs) was used to add an NES (LQKKLEELEL) or NLS (SPKKKRKVE) followed by a $(GGGGS)_2$ flexible linker to the 5' end (N-terminus) of pLENTI-puro MCPyV ST ko/co to create pLENTI-puro NES-MCPyV ST ko/co and pLENTI-puro NLS-MCPyV ST ko/co. pMTBS (donated by Dr. Christopher Buck, NIH) was used to transfect 293A cells with MCPyV ST. TSPyV ST and HPyV 7 ST were subcloned into pCS2 to transfect 293A cells. Empty pLENTI and pCS2 were used as negative controls.

Tissue Culture, Transductions, and Transfections

Adenovirus-transformed human embryonic kidney cells (HEK293A), 293TNs, and Rat-2 cells were cultured in 1x DMEM (ThermoFisher Scientific) supplemented with 10% fetal bovine serum (FBS) (Neuromics), GlutaMax (ThermoFisher Scientific), Non-Essential Amino Acids (NEAA) (ThermoFisher Scientific), and 100 units/mL penicillin-streptomycin (ThermoFisher Scientific). MCC virus-positive MKL-1 cells were cultured in 1x RPMI (ThermoFisher Scientific) supplemented similarly as stated above. All cell lines were incubated at 37°C in 5% CO_2.

293TN cells were transfected with the lentiviral envelope and packaging plasmids

psPAX2 (Addgene #12260) and pMD2.G (Addgene #12259) and lentiviral transfer plasmid

pLENTI-puro using TransIT-293 Transfection reagent (Mirus). Viral harvests were conducted at

24 and 48 hours post-transfection. Rat-2 cells were transduced with the pseudotyped lentiviruses

supplemented with 4μg/mL polybrene (SantaCruz Biotechnology) and selected with 4μg/mL

puromycin (ThermoFisher Scientific) 48 hours post-transduction.

HEK293 cells were transfected with pMTBS and pCS2-ST at ~80% confluence using

TransIT-293 transfection reagent (Mirus). Cells were harvested for downstream applications 48

hours post-transfection.

Immunoblotting and Antibodies

Cell lysates were prepared in RIPA lysis buffer (ThermoFisher Scientific) supplemented

with a protease inhibitor cocktail (ThermoFisher Scientific) on ice, followed by sonication at

20% amplitude and centrifugation at 16,000 x g for 15 minutes. A BCA assay (ThermoFisher

Scientific) was performed to determine protein concentrations for normalization in 2x SDS

sample buffer (ThermoFisher Scientific) at 1μg/μL followed by boiling at 95°C. 30μg of protein

was loaded in an 8-16% SDS-PAGE gel (ThermoFisher Scientific) and transferred to an

Immobilon-P PVDF membrane, followed by a 1 hour block in 4% milk and overnight primary

antibody incubation with Ab5 (donated by James Decaprio, Harvard University) diluted at

1:1000 for MCPyV ST, HPyV7 ST, and TSPyV ST, and MCPyV LT, HSP90 (Cell Signaling

Technologies, 4874S) (1:5000), HDAC2 (Cell Signaling Technologies, 2540S) (1:5000), Na/K-

ATPase (Cell Signaling Technologies, 3010S) (1:2500), or actin (Cell Signaling Technologies,

5125S) (1:10,000). After TBST washes, membranes were incubated in 4% milk supplemented

with either anti-mouse IgG HRP-conjugated secondary antibody (Cell Signaling Technologies,

7076P2) (1:10,000) or anti-rabbit IgG HRP-conjugated secondary antibody (Cell Signaling

Technologies, 7074P2) (1:10,000) at room temperature for 1 hour. Following secondary

washing, proteins were detected using Femto (ThermoFisher Scientific) and a ChemiDoc

Imaging System (Bio-Rad).

Transformation Assays

Soft agar assays were performed in triplicate according to the JOVE online protocol at

50,000 cells/well (33). The top layer of 1x DMEM was supplemented with puromycin. The cells

were monitored for three weeks, followed by imaging with a Nikon Ti2 Eclipse microscope

(Nikon).

Metabolism experiments were performed in triplicate using a Glucose Uptake-Glo Assay

Kit (Promega) according to manufacturer instructions at 10,000 cells/96-well.

Serum dependence experiments were performed by plating 50,000 cells/6-well in

triplicate in complete DMEM supplemented with either 10%, 5%, or 1% FBS (Neuromics). Cells

were quantified at days 2, 5, and 10. Puromycin-supplemented media was changed every 3 days

or as needed in later time points.

Proliferation rate experiments were conducted by plating 50,000 cells/6-well in triplicate.

Cells were quantified at days 2, 4, 8, 14, 20, and 28. Doubling time was evaluated between days

0 and 28 to be representative of the entire experiment. Puromycin-supplemented media was

changed every 3 days or as needed in later time points.

Subcellular Fractionation

Subcellular fractionations were conducted using a Subcellular Fractionation Kit

(ThermoFisher Scientific) following a manufacturer-modified protocol. In short, cells were

plated at 1×10^6 in 4 10cm plates 24 hours prior to harvest through trypsinization (ThermoFisher

Scientific) for adherent cells. Provided fractionation extraction buffers were supplemented with a

1x protease inhibitor cocktail and used to fractionate the cell sample as instructed with the

addition of extra wash steps to lessen fraction contamination. Following protein extraction,

lysates were quantified and run in a western blot as described previously. Controls included Heat

Shock Protein 90 (HSP90) for the cytoplasmic fraction, Sodium Potassium ATPase (NA/K

ATPase) for the membrane fraction, and Histone Deacetylase 2 (HDAC2) for the nuclear soluble

fraction.

Statistical Methods

Statistical analyses were conducted using the Jamovi statistical computer software, Version 2.3.

All experiments were repeated in triplicate. One-way ANOVA analyses were used, and a *p-value*

<0.05 was considered to indicate a statistically significant difference.

RESULTS

The ST unique region is responsible for the dominant transforming properties of MCPyV ST

In order to investigate the mechanism(s) of MCPyV cellular transformation, we first

sought to assess various transforming capabilities of the MCPyV tumor antigens (TAgs)

expressed in virus positive MCC (VP-MCC): ST and LT-t. As done previously by others, Rat-2

cells were transduced with pseudotyped lentiviruses containing plasmids encoding MCPyV ST,

LT-t, or empty pLENTI control (34). After confirmation of protein expression (Fig. 1A), soft

agar assays were performed to assess the ability of the MCPyV TAgs to induce anchorage-

independent cellular proliferation, a common property of cancerous cells that promotes tumor

formation and metastasis (35). Consistent with the work of others, MCPyV ST was capable of

inducing anchorage-independent growth, whereas MCPyV LT-t was incapable of colony

formation (Fig. 1B and C) (34,36).

Because cellular transformation can be defined by properties other than anchorage-

independent growth, additional transformation assays were performed between the MCPyV

TAgs to further define the specific transforming properties of both MCPyV ST and LT-t. It has

been frequently described that cancerous cells have an increased metabolic rate and, therefore,

glucose uptake than healthy cells to support their robust cellular division (37,38). Consistently,

MCPyV ST expression significantly increased the glucose uptake of Rat-2 cells, whereas

MCPyV LT-t did not (Fig. 1D). Similarly, MCPyV ST was also capable of significantly

increasing the proliferation rate and decreasing the doubling time of Rat-2 cells (Fig. 1E and F);

however, these assays demonstrated that MCPyV LT-t was also capable of significantly

increasing the cellular proliferation rate and decreasing the doubling time compared to control

cells, albeit to a lesser degree than MCPyV ST. These data support the transforming properties of

both MCPyV ST and LT-t, consistent with the necessity of both ST and LT-t to be expressed for

the viability of VP-MCC cells.

Finally, the serum dependence of Rat-2 cells was compared between control, ST, and LT-

t expressing cells, as cancerous cells are frequently described as capable of maintaining cellular

proliferation in the absence of the growth factors and nutrients found in serum (39). Consistent

with proliferation rate experiments containing the normal 10% serum supplementation (Fig. 1E),

both ST and LT-t were capable of significantly increasing proliferation in 10% serum; however,

MCPyV LT-t exhibited less proliferation than MCPyV ST with reduced serum concentrations of

either 5% or 1% (Fig. 1G). To illustrate the effect of reducing serum concentrations on the

proliferation of the control, ST, and LT-t expressing cells, the percent change in doubling time

was calculated going from 10% to 1%, 10% to 5%, and 5% to 1% serum. Although the percent

change in doubling time between these different serum concentrations was not significantly

different between control, ST, and LT-t expressing cells when going from 10% to 1% serum,

normalization by glucose uptake (Fig. 1D) found MCPyV ST to significantly decrease the

percent change in doubling time upon serum reductions to 1% (Fig. 1H). Together, these data

found serum concentrations to affect the proliferation of ST-expressing cells the least, despite

these cells having the highest metabolic rate as inferred by glucose uptake.

Upon confirmation of ST as the dominant transforming protein of MCPyV, we sought to

elucidate the domain(s) responsible for this phenotype. Both MCPyV ST and LT-t share the

same start codon and are created through the alternative splicing of the same mRNA (11).

Therefore, both ST and LT-t share the same N-terminal amino acid sequence, referred to as

"common-T", whereas they differ in their C-terminal amino acid sequence, which is referred to

as the "ST- and LT – unique regions" (Fig. 1I). The finding of ST to be the dominant

transforming protein of MCPyV led to the hypothesis that the domain(s) responsible for these

unique transforming properties are likely located in the ST-unique region, as location within the

common-T region would suggest MCPyV LT-t to have similar transforming properties.

Together, these data confirm ST as the dominant transforming protein of MCPyV by assessment

of various transformation properties such as anchorage-independent growth, metabolism,

proliferation rate, doubling time, and serum dependence, and that the domain(s) responsible for

this phenotype are likely found in the ST unique region.

The structurally dissimilar loops in the ST unique region of MCPyV ST may be responsible

for the uniquely transforming capabilities of MCPyV ST when compared to other skin-tropic

human polyomaviruses

Upon confirmation of ST being the dominant transforming protein of MCPyV, likely a

result of activities in the ST unique region, we sought to further elucidate the region(s) of

MCPyV ST responsible for various transformation properties. As noted, MCPyV ST belongs to a

family of 14 known human polyomaviruses, but is the only polyomavirus found to be directly

responsible for the development of cancer (20,21). Therefore, we sought to determine whether

the ST antigens of other, skin tropic, human polyomaviruses, such as Human Polyomavirus 7

(HPyV7) and Trichodysplasia Spinulosa Polyomavirus (TSPyV), were also capable of any of the

mechanisms of cellular transformation similar to MCPyV ST. Rat-2 cells were transduced to

express MCPyV, TSPyV, or HPyV7 ST (Fig. 2A). Although it appeared that both HPyV7 and

TSPyV ST proteins were expressed in lower quantities than MCPyV ST, it is likely that the

antibody used to detect these proteins has higher affinity to MCPyV ST, as it was developed

through its recognition of the MCPyV ST antigen; however, each T antigen was similarly codon

optimized and under the control of the same CMV promoter.

Consistent with TSPyV and HPyV7 not being found to be associated with cancer,

expression of HPyV7 and TSPyV ST proteins in Rat-2 cells failed to significantly induce

anchorage-independent growth above control cells (Fig. 2B and C). Furthermore, the metabolic

rate, as measured by glucose uptake, of HPyV7 and TSPyV ST were lower than that of MCPyV

ST (Fig. 2D). Interestingly, although the proliferation rate, doubling time, and serum dependence

of HPyV7 ST were more similar to control cells, TSPyV ST induced a cellular proliferation rate,

doubling time, and serum dependence similar to that of MCPyV ST, further demonstrating that

transformation is a combination of many cellular abnormalities that are not necessarily connected

(Fig. 2E, F, G, and H).

Together, our results indicate that MCPyV ST can uniquely induce anchorage-

independent growth, increase glucose uptake and proliferation rate, and reduce doubling time

and serum dependence when compared to HPyV7 ST; however, although TSPyV ST was

incapable of inducing anchorage-independent growth and increased glucose uptake, it was able

to increase cellular proliferation and reduce doubling time and serum dependence. In an effort to

determine the domain(s) responsible for the unique transforming capabilities of MCPyV ST, an

amino acid alignment of MCPyV, TSPyV, and HPyV7 ST was performed (Supplementary

Figure 1). Unfortunately, although all evolutionarily related ST proteins, they contained

exceptionally different amino acid sequences that made it extremely difficult to identify MCPyV

ST unique domains possibly responsible for transformation. Despite dissimilar amino acid

sequences, MCPyV, TSPyV, and HPyV7 ST proteins were found to have very similar protein

structures upon protein structure alignment (Fig. 2I). Consistent with the common T region being

hypothesized to not contain the domain(s) necessary for MCPyV ST transformation (Fig. 1I), the

common-T structure of the transforming MCPyV ST, and non-transforming TSPyV and HPyV7

ST were very similar. However, despite their similarity, MCPyV, TSPyV, and HPyV7 ST

contained structurally dissimilar loops in the ST-unique region, consistent with the previous

hypothesis of the transforming domains being found in the ST-unique region (Fig. 1I). Together,

these data suggest that the unique transforming capabilities of MCPyV ST may be a result of the

activities of the structurally dissimilar loops in the unique region.

MCPyV ST binds to many nuclear cellular proteins

Previous studies of the MCPyV viral life cycle and MCPyV ST-mediated cellular

transformation have identified many MCPyV ST cellular interactors (13,15,16,34,36,40–45).

Interestingly, when reviewing these findings, we found that 74% of the previously identified

MCPyV ST cellular interactors are nuclear proteins (Fig. 3A). These findings were unexpected,

as the amino acid sequence of MCPyV ST does not contain a known, canonical NLS, nor do

many cellular localization programs such as PSORTII, seqNLS, and BaCelLO predict MCPyV

ST to exhibit nuclear localization (Fig. 3B and C). Although MCPyV LT has been found to

localize to the nucleus as a result of its NLS, the NLS of LT is found in the LT unique region,

and mutation of the common-T region of LT did not affect its nuclear localization (Fig. 3D) (46).

Therefore, in addition to transformation, the potential nuclear localization of MCPyV ST may

also be due to the activity of a domain found in the ST unique region.

MCPyV ST uniquely localizes to the nucleus independent of size

In order to determine whether MCPyV ST can localize to the nucleus in the absence of a

canonical NLS, subcellular fractionations (SCF) were performed. Rat-2 cells transduced with

pLENTI MCPyV ST were fractionated into cytoplasmic, nuclear, and membrane fractions, with

MCPyV ST being found to localize to both the cytoplasmic and nuclear subcellular

compartments (Fig. 4A). Although a small amount of MCPyV ST protein was detected in the

membrane fraction, it is consistent with the level of HSP90 cytoplasmic contamination within the

membrane fraction, and is likely a false signal. To confirm this finding in another cell line,

293As transfected with pMTBS also identified MCPyV ST to localize to the cytoplasmic and

nuclear fractions, with membrane localization again likely a result of cytoplasmic contamination

(Fig. 4B). Of note, it was consistently found that SCF experiments in 293A cells frequently led

to more contamination than Rat-2 cells; however, the results from both cell lines confirmed both

MCPyV ST cytoplasmic and nuclear localization.

To confirm these findings in a physiologically relevant cell line, an SCF was performed

in MKL-1 cells, a virus-positive MCC cell line. Interestingly, in a VP-MCC cell line,

endogenous MCPyV ST localized exclusively to the nucleus (Fig. 4C). This finding was

consistent with exogenous MCPyV ST nuclear localization in Rat-2 and 293A cells, but

inconsistent with shared cytoplasmic localization.

As shown previously, MCPyV ST is uniquely transforming when compared to other

human, skin-tropic polyomaviruses such as TSPyV and HPyV7 ST (Fig. 2). To determine

whether MCPyV ST is also unique in its ability to localize to the nucleus despite the absence of a

canonical NLS, SCFs were performed on Rat-2 and 293A cells transfected with pCS2 TSPyV ST

or pCS2 HPyV7 ST. Interestingly, both TSPyV ST (Fig. 4D and E) and HPyV7 ST (Fig. 4F and

G) exclusively localized to the cytoplasm in both Rat-2 and 293A cells. These data suggest that

MCPyV ST is unique in both its transforming capacity and nuclear localization. Furthermore, the

domain(s) responsible for the unique transformation and nuclear localization of MCPyV ST may

be found in the structurally dissimilar loops of the ST unique region (Fig. 2I).

MCPyV ST nuclear localization is important for anchorage-independent growth, increased

metabolism, and decreased serum dependence of MCPyV ST, but is expendable for

proliferation rate and doubling time

In order to determine whether nuclear localization of MCPyV ST is necessary for cellular

transformation, both a cytoplasmic and nuclear sequestered MCPyV ST were created by adding

the NLS from SV40, or the nuclear export signal (NES) from Mitogen-Activated Protein Kinase

Kinase (MAPKK), to MCPyV ST (NLS-ST and NES-ST, respectively). Importantly, localization

domains were added to the N-terminus of MCPyV ST to ensure the furthest distance from the

likely transforming domains within the structurally dissimilar loops of the MCPyV ST unique

region in the C-terminus (Fig. 1I and 2I). Furthermore, localization domains were added with the

addition of a flexible linker to decrease the likelihood of the added domain interfering with the

structure and/or functions of MCPyV ST. To ensure the expression and stability of the MCPyV

ST localization mutants, a confirmation western blot was performed and found the localization

mutants and wild-type MCPyV (WT-ST) to all be expressed; however, protein densitometry

revealed NLS-ST and NES-ST to be expressed at 41% and 59% of WT-ST, respectively. SCF

experiments in Rat-2 cells confirmed the appropriate localization of WT-ST and the localization

mutants (Fig. 5B). Similar to WT-ST, NLS-ST retained localization to both the nucleus and the

cytoplasm, with a higher prevalence of nuclear localization. More importantly, NES-ST localized

exclusively to the cytoplasm, necessary to assess the importance of nuclear localization in

MCPyV ST-mediated cellular transformation.

To assess the effect of localization on MCPyV ST-mediated cellular transformation,

various transformation assays were performed comparing WT-ST, NLS-ST, and NES-ST,

followed by normalization based on protein densitometry. NLS-ST was capable of forming

colonies above WT-ST post-normalization, whereas NES-ST also led to colony formation, but in

fewer numbers (Fig. 5 C and D). Normalized glucose uptake of NLS-ST was significantly higher

than that of WT-ST, whereas the glucose uptake of NES-ST was similar to that of WT-ST,

suggesting nuclear localization of MCPyV ST may influence the ability of MCPyV ST to

increase cellular metabolism (Fig. 5E). The proliferation curves of WT-ST and both localization

mutants were significantly increased when compared to the empty pLENTI transduced controls,

but were not significantly different from each other (Fig. 5F). However, the protein-level

normalized doubling times of NLS-ST and NES-ST were both significantly faster than WT-ST,

with NLS-ST having the fastest doubling time (Fig. 5G).

No significant differences were observed when comparing the proliferation rate curves of

WT-ST compared to the localization mutants at differing serum concentrations, likely as a result

of the differing protein levels and metabolic rates of each ST (Fig. 5H). Therefore, the percent

change in doubling time between differing serum concentrations was calculated and normalized

by both the protein levels (Fig. 5A) and glucose uptake rates (Fig. 5E) of each ST (Fig. 5I).

Together, WT-ST and NES-ST were each found to have similar changes in the percent doubling

time between 10% and 1% serum containing media, whereas NLS-ST was found to have a

significantly lower percent change in the doubling time between 10% and 1% serum-containing

media. Together, these data suggest nuclear localization of ST is largely responsible for the

ability of MCPyV ST to proliferate in the absence of serum despite having an increased

metabolic rate, as inferred by an increased glucose uptake. In conclusion, these data suggest that

cellular proliferation and doubling time may be a combined result of both nuclear and

cytoplasmic localization, whereas increased anchorage independent growth, metabolism, and

decreased serum dependence may be influenced by the nuclear localization of MCPyV ST.

Discussion

The mechanism(s) of MCPyV-mediated cellular transformation and tumorigenesis have

been investigated since the discovery of MCPyV in 2008; however, many of these early analyses

were hindered by using the approach of comparing the similarities between the human oncogenic

polyomavirus MCPyV, and the simian oncogenic virus SV40. The findings of the current study

and of others that ST is the dominant transforming protein of MCPyV is opposite to SV40, in

which ST plays only a supportive or accessory role, with LT being the dominant transforming

protein (47). Similarity comparisons between the dissimilar SV40 and MCPyV are responsible

for many years spent investigating MCPyV ST binding to PP2A and Fbw7, which have been

found to be irrelevant for transformation by MCPyV ST (40,41). Furthermore, we found MCPyV

ST to be uniquely transforming among skin-tropic human polyomaviruses ST antigens,

consistent with MCPyV being the only human polyomavirus clearly associated with cancer.

Together, these data refute the method of identifying mechanisms of MCPyV transformation

through similarity comparisons to SV40, and support the alternative dissimilarity approach of

investigating the differences between MCPyV ST and the ST of other non-oncogenic human

polyomaviruses to identify the unique mechanisms of MCPyV ST mediated transformation.

The dissimilarity approach can also be applied when comparing MCPyV ST and LT-t, as

they share a common T region, have their own unique regions, and accomplish dissimilar

patterns of transformation. Through this approach, we were able to narrow down the location of

the domain(s) responsible for MCPyV ST transformation to the ST unique region. Although

transformation assays performed on Rat-2 cells expressing the ST unique region alone may be

able to confirm this finding, it is likely that the absence of common-T may influence the

structure of the ST unique region, and it is possible that domain(s) within the common-T region

may be necessary to facilitate the activities of the ST unique region.

Although the discovery that MCPyV is uniquely oncogenic among human

polyomaviruses was discouraging at the genesis of MCPyV oncogenesis research, the

dissimilarity approach exploited it for our benefit. Comprehensive transformation assay analysis

of TSPyV ST and HPyV7 ST identified that although the ST of HPyV7 and TSPyV were not as

robustly transforming as MCPyV ST, TSPyV ST was capable of achieving some properties of

cellular transformation, further supporting the fact that cellular transformation is a collection of

properties rather than a single entity, and that the ability to induce several transforming

properties may translate to being able to accomplish tumorigenesis *in vivo*. Furthermore,

structural alignments allowed us to further narrow down the unique transforming domains of

MCPYV ST to the structurally dissimilar loops consistent with the unique region of MCPyV ST.

It is currently hypothesized that the similar transforming capabilities of MCPyV and TSPyV ST

may reside in domains shared by MCPyV and TSPyV ST, but not found in HPyV7

ST. Alternatively, TSPyV ST and MCPyV ST may accomplish similar methods of cellular

transformation through distinct mechanisms.

The identification of MCPyV ST binding to several nuclear proteins suggested nuclear

localization despite the absence of a canonical NLS in MCPyV ST. Surprisingly, subcellular

fractionation confirmed nuclear localization of MCPyV ST in various cell lines, whereas the

non-oncogenic TSPyV and HPyV7 ST proteins were confined to the cytoplasm, further proving

the usefulness of the dissimilarity hypothesis. Although MCPyV ST is under the 30kDa size

necessary for passive diffusion through the nuclear pore, the finding that 30kDa TSPyV and

HPyV7 ST proteins are cytoplasmic confirms that MCPyV ST localization to the nucleus is

regulated (48). Interestingly, MCPyV ST exhibited both nuclear and cytoplasmic localization in

transduced Rat-2 and 293A cells, yet it was exclusively nuclear in VP-MCC cells. Although it is

possible that the differences in MCPyV ST localization between the cell lines may be an artifact

of differing cell lines, hypotheses to explain these differences include the following: First, it is

possible that the mechanism of MCPyV ST nuclear translocation is saturated in Rat-2 and 293A

cells, overexpressing exogenous ST, leading to spillover into the cytoplasm. Second, the

presence of MCPyV LT-t in MKL-1 cells, but not Rat-2 and 293A cells, may enhance the

efficiency of MCPyV ST nuclear localization. Although both hypotheses require further

experimentation, it is clear from these experiments that MCPyV ST is capable of uniquely

localizing to the nucleus despite the absence of a canonical NLS.

It can be hypothesized that the noncanonical NLS or domain responsible for MCPyV ST

nuclear localization is found in the MCPyV ST unique region, as previous investigation into

MCPyV LT-t nuclear localization identified the common T region to not be responsible for

nuclear localization (46). As the function of an NLS is dependent on both structure and function,

it is possible that the MCPyV ST NLS is located in the highly dissimilar amino acid sequence or

the structurally dissimilar loops between MCPyV, TSPyV, and HPyV7 ST.

Although most of the attention regarding MCPyV-mediated oncogenesis has been

directed towards ST, it is clear that the large T antigen is also important due to its necessity for

the viability of MCC cells and its ability to influence cellular proliferation and doubling time,

despite not inducing anchorage-independent growth, metabolism, and serum dependence. Due to

the sharing of the common-T region between LT and ST, the domain(s) responsible for inducing

proliferation rate and doubling time may be found in the common T region; however, it is more

likely that ST and LT-t accomplish this through different mechanisms attributed to their

common-T region, as they affected Rat-2 cells dissimilarly, despite similar protein expression

levels. It is likely that perturbation of Rb by the Rb binding domain of MCPyV LT-t contributes

to these phenotypes; however, this has not yet been directly assessed.

Finally, the export of MCPyV ST to the cytoplasm did not ablate all properties of cellular

transformation, suggesting a potential role of MCPyV in both the cytoplasm and nucleus.

However, it is not currently understood how this finding translates to the observation of MCPyV

ST being exclusively nuclear in VP-MCC cells, and therefore warrants further investigation.

Furthermore, although normalized NES-ST was still capable of anchorage-independent growth,

increased glucose uptake, and reduced serum dependence, NLS-ST normalization led to

anchorage-independent growth, glucose uptake, and serum dependence significantly higher than

WT-ST and NES-ST. Although protein levels of WT-ST and NLS-ST differed, the ratio of WT-

ST and NLS-ST in the cytoplasmic and nuclear compartments were similar. Therefore, it is of

great interest why NLS-ST was capable of increasing transformation above WT-ST in some

assays but not others.

Conclusions

Together, it has been found that elucidating mechanisms of transformation guided by a dissimilarity approach may prove useful in identifying otherwise hidden, unique mechanisms of MCPyV oncogenesis, such as the unique nuclear localization of MCPyV ST. Cellular transformation was found to be multi-variable and not simply a positive or negative characteristic of viral proteins, consistent with our understanding of the role of viral proteins in viral replication. Further investigation into the mechanisms and role of MCPyV nuclear localization in transformed and MCC cells will further enhance our understanding of the natural viral life cycle and MCPyV-mediated oncogenesis necessary for basic virology and the treatment of MCC.

List of Abbreviations

MCC   Merkel Cell Carcinoma

MCPyV   Merkel Cell Polyomavirus

HPyV7   Human Polyomavirus 7

TSPyV   Trichodysplasia Spinulosa Polyomavirus

HPyV   Human Polyomavirus

NES   Nuclear Export Signal

NLS   Nuclear Localization Signal

TAg   Tumor Antigens

ST   Small Tumor Antigen

LT-t   Truncated Large Tumor Antigen

ALTO   Alternate Large T Open Reading Frame

ER   Early Region

LR   Late Region

VP-MCC   Virus Positive Merkel Cell Carcinoma

FBS   Fetal Bovine Serum

SCF   Subcellular Fractionation

SAA   Soft Agar Assay

SV40   Simian Virus 40

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article and

supplementary materials.

Competing interests

The authors declare no conflict of interest.

Funding

Financial support was provided by the Stetson University Health Sciences Department and Cell

Biology Education Consortium.

Author Contributions

K.T., I.T., and C.D. facilitated conceptualization, investigation, data curation, writing of the

original draft, review and editing, and visualization. M.S. performed formal analysis and review

and editing. K.D. performed conceptualization, methodology, validation, investigation, writing

of original draft and review and editing, visualization, supervision, and funding acquisition.

Acknowledgments

The plasmids and lentiviruses used in these studies, and the modeling of ST structures were

made in the laboratory of Dr. Denise A. Galloway at the Fred Hutchinson Cancer Center when

Dr. Dye was a graduate student. This work was supported by R35-CA209979 to DAG and by

T32-AIO83203 to KND. Furthermore, we would also like to thank Richard Wang (University of

Texas) for sharing Rat-2 cells, and Dr. Patrick Moore and Dr. Yuan Chang (University of

Pittsburgh) for sharing MKL-1 cells, and Dr. James DeCaprio (Harvard) for donating Ab5.

 <u>Figure Legends</u>

Figure 1. The ST unique region is responsible for the dominant transforming properties of

MCPyV ST. *Rat-2 cells were transduced with MCPyV LT-t, MCPyV ST, or pLENTI control and*

protein expression was confirmed (A). Various transformation assays were performed including

soft agar assays (B and C), glucose uptake (D), proliferation rate (E), doubling time (F), and

serum dependence (G) (grey – pLENTI, coral – MCPyV ST, blue – MCPyV LT-t). The percent

change in serum dependence doubling time (solid bars) was normalized (dashed bars) by

glucose uptake compared to pLENTI control (H). MCPyV ST protein structure was determined

by PyMol and colored to show the common-T (blue) and ST unique (pink) regions (I). Each point

*represents mean ± the standard error of independent triplicates. One-way ANOVA * $p < 0.05$; ***

*< 0.01; *** $p < 0.001$, significantly different from pLENTI.*

Figure 2. The structurally dissimilar loops in the ST unique region of MCPyV ST may be

responsible for the uniquely transforming capabilities of MCPyV ST when compared to other

skin-tropic human polyomaviruses. *Rat-2 cells were transduced with MCPyV ST, HPyV7 ST,*

TSPyV ST, or pLENTI control, and protein expression was confirmed (A). Various

transformation assays were performed, including soft agar assays (B and C), glucose uptake (D),

proliferation rate (E), doubling time (F), and serum dependence (grey – LENTI, coral – MCPyV

ST, purple – HPyV7 ST, yellow – TSPyV ST). The percent change in serum dependence doubling

time (solid bars) was normalized (dashed bars) by glucose uptake compared to pLENTI control

(H). MCPyV ST, HPyV7 ST, and TSPyV ST protein structures were determined and aligned

using PyMol and black arrows denote the structurally dissimilar regions in protein structure (I).

*Each point represents mean ± the standard error of independent triplicates. One-way ANOVA * p*

*< 0.05; ** < 0.01; *** p < 0.001, significantly different from pLENTI.*

***Figure 3. MCPyV ST binds to many nuclear cellular proteins.** The localization of known*

MCPyV ST cellular interactors were recorded, with 31% being nuclear (orange), 26% being

cytoplasmic (blue), and 43% that are both cytoplasmic and nuclear (A). The amino acid

sequence of MCPyV ST contains no known canonical NLS (B). Online amino acid and protein

subcellular localization predictor tools were utilized to predict the localization of MCPyV ST

(C). The location of the MCPyV LT-t NLS is found within the MCPyV LT-t unique region

(orange), but no known NLS is found within the common-T region (red) or ST unique region

(blue) (D).

***Figure 4. MCPyV ST uniquely localizes to the nucleus independent of size.** Rat-2 (A) and 293A*

(B) cells were transduced and transfected, respectively, with MCPyV ST and a subcellular

fractionation was performed to isolate proteins for the cytoplasmic (C), membrane (M), and

nuclear (N) fractions. A subcellular fractionation for MCPyV ST was also performed on the VP-

MMCC cells, MKL-1 (C). Rat-2 and 293A cells were also transduced and transfected,

respectively, with TSPyV ST (D and E) and HPyV7 ST (F and G), followed by subcellular

fractionation. HSP90 was used as a cytoplasmic control, NA^+/K^+-ATPase was used as a

membrane control, and HDAC2 was used as a nuclear control.

Figure 5. MCPyV ST nuclear localization is important for anchorage-independent growth,

increased metabolism, and decreased serum dependence of MCPyV ST, but is expendable for

proliferation rate and doubling time. *Rat-2 cells were transduced with MCPyV ST (ST-WT), a*

cytoplasmic sequestered MCPyV ST (NES-ST), a nuclear sequestered MCPyV ST (NLS-ST), or

pLENTI control, and protein expression was confirmed with protein densitometry (A). A

subcellular fractionation was performed to verify localization of the wildtype ST and localization

mutants. Subcellular fractionations included cytoplasmic (C), membrane (M), and nuclear (N)

fractions, with HSP90 acting as a cytoplasmic control, NA^+/K^+-ATPase as a membrane control,

and HDAC2 as a nuclear control (B). Various transformation assays were performed, including

soft agar assays (C and D), glucose uptake (E), proliferation rate (F), doubling time (G), and

serum dependence (H) (grey – pLENTI control, coral – MCPyV ST, maroon – NES-ST, teal –

NLS-ST). The number of soft agar colonies (D), glucose uptake (E), and doubling time (G) were

normalized based on protein levels determined by densitometry (solid bars – unnormalized,

dashed bars – normalized) (A). The percent change in serum dependence doubling time (solid

bars) was normalized (dashed bars) by densitometry determined protein levels and glucose

uptake compared to wildtype MCPyV ST (I). Each point represents mean ± the standard error of

*independent triplicates. One-way ANOVA * $p < 0.05$; ** < 0.01; *** $p < 0.001$, significantly*

different from MCPyV ST.

Supplementary Figure 1: *An amino acid alignment of MCPyV ST, HPyV7 ST, and TSPyV ST*

created using ClustalOmega.com.

1. Toker C. Trabecular carcinoma of the skin. Arch Dermatol. 1972;105(1):107–10.

2. Becker JC. Merkel cell carcinoma. Ann Oncol. 2010;21(suppl_7):vii81–5.

3. Goh G, Walradt T, Markarov V, Blom A, Riaz N, Doumani R, et al. Mutational landscape of
MCPyV-positive and MCPyV-negative Merkel cell carcinomas with implications for
immunotherapy. Oncotarget. 2015;7(3):3403–15.

4. Feng H, Shuda M, Chang Y, Moore PS. Clonal Integration of a Polyomavirus in Human
Merkel Cell Carcinoma. Science. 2008;319(5866):1096–100.

5. Martel-Jantin C, Pedergnana V, Nicol JTJ, Leblond V, Trégouët DA, Tortevoye P, et al.
Merkel cell polyomavirus infection occurs during early childhood and is transmitted between
siblings. J Clin Virol. 2013;58(1):288–91.

6. Gossai A, Waterboer T, Nelson HH, Michel A, Willhauck-Fleckenstein M, Farzan SF, et al.
Seroepidemiology of Human Polyomaviruses in a US Population. Am J Epidemiology.
2016;183(1):61–9.

7. Zhang C, Liu F, He Z, Deng Q, Pan Y, Liu Y, et al. Seroprevalence of Merkel Cell
Polyomavirus in the General Rural Population of Anyang, China. PLoS ONE.
2014;9(9):e106430.

8. Šroller V, Hamšíková E, Ludvíková V, Vochozková P, Kojzarová M, Fraiberk M, et al.
Seroprevalence rates of BKV, JCV, and MCPyV polyomaviruses in the general Czech Republic
population. J Méd Virol. 2014;86(9):1560–8.

9. Tolstov YL, Knauer A, Chen JG, Kensler TW, Kingsley LA, Moore PS, et al. Asymptomatic
Primary Merkel Cell Polyomavirus Infection among Adults - Volume 17, Number 8—August
2011 - Emerging Infectious Diseases journal - CDC. Emerg Infect Dis. 2011;17(8):1371–80.

10. Shuda M, Feng H, Kwun HJ, Rosen ST, Gjoerup O, Moore PS, et al. T antigen mutations are
a human tumor-specific signature for Merkel cell polyomavirus. Proc Natl Acad Sci.
2008;105(42):16272–7.

11. DeCaprio JA. Merkel cell polyomavirus and Merkel cell carcinoma. Philos Trans R Soc B:
Biol Sci. 2017;372(1732):20160276.

12. Carter JJ, Daugherty MD, Qi X, Bheda-Malge A, Wipf GC, Robinson K, et al. Identification
of an overprinting gene in Merkel cell polyomavirus provides evolutionary insight into the birth
of viral genes. Proc Natl Acad Sci. 2013;110(31):12744–9.

13. Shuda M, Kwun HJ, Feng H, Chang Y, Moore PS. Human Merkel cell polyomavirus small T
antigen is an oncoprotein targeting the 4E-BP1 translation regulator. J Clin Investig.
2011;121(9):3623–34.

14. Sunshine JC, Jahchan NS, Sage J, Choi J. Are there multiple cells of origin of Merkel cell
carcinoma? Oncogene. 2018;37(11):1409–16.

15. Park DE, Cheng J, Berrios C, Montero J, Cortés-Cros M, Ferretti S, et al. Dual inhibition of
MDM2 and MDM4 in virus-positive Merkel cell carcinoma enhances the p53 response. Proc
Natl Acad Sci. 2019;116(3):1027–32.

16. Cheng J, Park DE, Berrios C, White EA, Arora R, Yoon R, et al. Merkel cell polyomavirus
recruits MYCL to the EP400 complex to promote oncogenesis. PLoS Pathog.
2017;13(10):e1006668.

17. Hesbacher S, Pfitzer L, Wiedorfer K, Angermeyer S, Borst A, Haferkamp S, et al. RB1 is the
crucial target of the Merkel cell polyomavirus Large T antigen in Merkel cell carcinoma cells.
Oncotarget. 2016;7(22):32956–68.

18. Houben R, Adam C, Baeurle A, Hesbacher S, Grimm J, Angermeyer S, et al. An intact
retinoblastoma protein□binding site in Merkel cell polyomavirus large T antigen is required for
promoting growth of Merkel cell carcinoma cells. Int J Cancer. 2012;130(4):847–56.

19. Schrama D, Hesbacher S, Angermeyer S, Schlosser A, Haferkamp S, Aue A, et al. Serine
220 phosphorylation of the Merkel cell polyomavirus large T antigen crucially supports growth
of Merkel cell carcinoma cells. Int J Cancer. 2016;138(5):1153–62.

20. Prado JCM, Monezi TA, Amorim AT, Lino V, Paladino A, Boccardo E. Human
polyomaviruses and cancer: an overview. Clinics. 2018;73(suppl 1):e558s.

21. DeCaprio JA, Garcea RL. A cornucopia of human polyomaviruses. Nat Rev Microbiol.
2013;11(4):264–76.

22. Eddy BE, Borman GS, Berkeley WH, Young RD. Tumors Induced in Hamsters by Injection
of Rhesus Monkey Kidney Cell Extracts. Proc Soc Exp Biol Med. 1961;107(1):191–7.

23. Eddy BE, Borman GS, Grubbs GE, Young RD. Identification of the oncogenic substance in
rhesus monkey kidney cell cultures as simian virus 40. Virology. 1962;17(1):65–75.

24. Shein HM, Enders JF. TRANSFORMATION INDUCED BY SIMIAN VIRUS 40 IN
HUMAN RENAL CELL CULTURES, I. MORPHOLOGY AND GROWTH
CHARACTERISTICS*. Proc Natl Acad Sci. 1962;48(7):1164–72.

25. Rabson AS, Kirschstein RL. Induction of Malignancy in vitro in Newborn Hamster Kidney
Tissue Infected with Simian Vacuolating Virus (SV40). Proc Soc Exp Biol Med.
1962;111(2):323–8.

26. Ahuja D, Sáenz-Robles MT, Pipas JM. SV40 large T antigen targets multiple cellular
pathways to elicit cellular transformation. Oncogene. 2005;24(52):7729–45.

27. Moore PS, Chang Y. Why do viruses cause cancer? Highlights of the first century of human
tumour virology. Nat Rev Cancer. 2010;10(12):878–89.

28. Butel JS. Viral carcinogenesis: revelation of molecular mechanisms and etiology of human
disease. Carcinogenesis. 2000;21(3):405–26.

29. Scott MS, Calafell SJ, Thomas DY, Hallett MT. Refining Protein Subcellular Localization.
PLoS Comput Biol. 2005;1(6):e66.

30. Hung MC, Link W. Protein localization in disease and therapy. J Cell Sci.
2011;124(20):3381–92.

31. Laurila K, Vihinen M. Prediction of disease-related mutations affecting protein localization.
BMC Genom. 2009;10(1):122.

32. Lu J, Wu T, Zhang B, Liu S, Song W, Qiao J, et al. Types of nuclear localization signals and
mechanisms of protein import into the nucleus. Cell Commun Signal. 2021;19(1):60.

33. Borowicz S, Scoyk MV, Avasarala S, Rathinam MKK, Tauler J, Bikkavilli RK, et al. The
Soft Agar Colony Formation Assay. J Vis Exp. 2014;(92):e51998.

34. Zhao J, Jia Y, Shen S, Kim J, Wang X, Lee E, et al. Merkel Cell Polyomavirus Small T
Antigen Activates Noncanonical NF-κB Signaling to Promote Tumorigenesis. Mol Cancer Res.
2020;18(11):1623–37.

35. Mori S, Chang JT, Andrechek ER, Matsumura N, Baba T, Yao G, et al. Anchorage-
independent cell growth signature identifies tumors with metastatic potential. Oncogene.
2009;28(31):2796–805.

36. Kwun HJ, Shuda M, Feng H, Camacho CJ, Moore PS, Chang Y. Merkel Cell Polyomavirus
Small T Antigen Controls Viral Replication and Oncoprotein Expression by Targeting the
Cellular Ubiquitin Ligase SCFFbw7. Cell Host Microbe. 2013;14(2):125–35.

37. Lin X, Xiao Z, Chen T, Liang SH, Guo H. Glucose Metabolism on Tumor Plasticity,
Diagnosis, and Treatment. Front Oncol. 2020;10:317.

38. Fadaka A, Ajiboye B, Ojo O, Adewale O, Olayide I, Emuowhochere R. Biology of glucose
metabolization in cancer cells. J Oncol Sci. 2017;3(2):45–51.

39. White EZ, Pennant NM, Carter JR, Hawsawi O, Odero-Marah V, Hinton CV. Serum
deprivation initiates adaptation and survival to oxidative stress in prostate cancer cells. Sci Rep.
2020;10(1):12505.

40. Kwun HJ, Shuda M, Camacho CJ, Gamper AM, Thant M, Chang Y, et al. Restricted Protein
Phosphatase 2A Targeting by Merkel Cell Polyomavirus Small T Antigen. J Virol.
2015;89(8):4191–200.

41. Dye KN, Welcker M, Clurman BE, Roman A, Galloway DA. Merkel cell polyomavirus
Tumor antigens expressed in Merkel cell carcinoma function independently of the ubiquitin
ligases Fbw7 and β-TrCP. PLoS Pathog. 2019;15(1):e1007543.

42. Shuda M, Velásquez C, Cheng E, Cordek DG, Kwun HJ, Chang Y, et al. CDK1 substitutes
for mTOR kinase to activate mitotic cap-dependent protein translation. Proc Natl Acad Sci.
2015;112(19):5875–82.

43. Pietropaolo V, Prezioso C, Moens U. Merkel Cell Polyomavirus and Merkel Cell Carcinoma.
Cancers. 2020;12(7):1774.

44. Wu JH, Narayanan D, Limmer AL, Simonette RA, Rady PL, Tyring SK. Merkel Cell
Polyomavirus Small T Antigen Induces DNA Damage Response. Intervirology. 2019;62(2):96–
100.

45. Abdul-Sada H, Müller M, Mehta R, Toth R, Arthur JSC, Whitehouse A, et al. The PP4R1
sub-unit of protein phosphatase PP4 is essential for inhibition of NF-κB by merkel polyomavirus
small tumour antigen. Oncotarget. 2017;8(15):25418–32.

46. Nakamura T, Sato Y, Watanabe D, Ito H, Shimonohara N, Tsuji T, et al. Nuclear localization
of Merkel cell polyomavirus large T antigen in Merkel cell carcinoma. Virology.
2010;398(2):273–9.

47. Cheng J, DeCaprio JA, Fluck MM, Schaffhausen BS. Cellular transformation by Simian
Virus 40 and Murine Polyoma Virus T antigens. Semin Cancer Biol. 2009;19(4):218–28.

48. Timney BL, Raveh B, Mironska R, Trivedi JM, Kim SJ, Russel D, et al. Simple rules for
passive diffusion through the nuclear pore complex. J Cell Biol. 2016;215(1):57–76.

A)

- Nuclear
- Cytoplasmic
- Both nuclear and cytoplasmic

B)

MDLVLNRKEREALCKLLEIAPNCYGNI
PLMKAAFKRSCLKHHPDKGGNPVIM
MELNTLWSKFQQNIHKLRSDFSMFD
EVSTKFPWEEYGTLKDYMQSGYNAR
FCRGPGCMLKQLRDSKCACISCKLSR
QHCSLKTLKQKNCLTWGECFCYQCFI
LWFGFPPTWESFDWWQKTLEETDY
CLLHLHLF

C)

	BaCelLO	Euk-mPloc 2.0	ESLPred	DeepLoc	PSORTII	seqNLS
Nucleus			✓	✓		
Cytosol	✓	✓			✓	✓

D)

```
HPyV7_ST    MDKLLGRDEVKELMELIGLNMACWGNLPLIQHKVRLASKKYHPDKGGDPQKMQRLNVLKD      60
MCPyV_ST    MDLVLNRKEREALCKLLEIAPNCYGNIPLMKAAFKRSCLKHHPDKGGNPVIMMELNTLWS      60
TSPyV_ST    MDKFLSREESLELMDLLQIPRHCYGNFALMKINHKKMSLKYHPDKGGDPEKMSRLNQLWQ      60
            ** .*.*.*    * .*: :   *:**: *::    :  . *:******:*  *  .** * .

HPyV7_ST    KLEATLRDQRSGSPMWHYSSDEVSFWDIE-LTVGEFLGP-EFNRKKVWNYNLCVVQGL--     116
MCPyV_ST    KFQQNIHKLRSDFSMF-DEVSTKFPWEEY-GTLKDYMQS-GYNA-RFCRGPGCMLKQLRD     116
TSPyV_ST    KLQEGIYNARQEFPTSFSSQVGSWYWEANLISLKEYFGKKKYDENVIKHWPQCAEKAL--     118
            *::   :  . *.          .       *:      :: :::      ::   . .    *  : *

HPyV7_ST    RACCCIHCILKRKHKKKAKEYAKDQRGPLSWGKCWCFHCYLNWFGVERSEES-FMWWSHI     175
MCPyV_ST    SKCACISCKLSRQHCSLK---TLKQKNCLTWGECFCYQCFILWFGFPPTWES-FDWWQKT     172
TSPyV_ST    KECKCLTCKIGLQHFVYK---QMHQKKCVVWGECFCYKCYCAWFGEDLYCLDSLWAWSCI     175
             * *: * :  :*            .*:  : **:*:*::*:  ***        . :  *.

HPyV7_ST    IFQTPMDVLNLWGQLNLL------     193
MCPyV_ST    LEETDYCLLHLHLF----------     186
TSPyV_ST    VGEVDFHLVNLYLRVNQGFNWGK     198
            : :.     :::*
```