Human dynein-dynactin is a fast processive motor in living cells

Vikash Verma[1], Patricia Wadsworth[1,2], and Thomas J. Maresca[1,2,]

[1] Biology Department, University of Massachusetts, Amherst

[2] Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst

Correspondence to:

Thomas J. Maresca
Biology Department
University of Massachusetts, Amherst
Amherst, MA 01003
Phone: 413-545-0957
Fax: 413-545-3243
E-mail: tmaresca@umass.edu

Patricia Wadsworth
Biology Department
University of Massachusetts, Amherst
Amherst, MA 01003
Phone: 413-545-4877
Fax: 413-545-3243
E-mail: patw@umass.edu

Running title: Direct visualization of dynein-dynactin motility in human cells

ABSTRACT

Minus-end directed transport along microtubules in eukaryotes is primarily mediated by

cytoplasmic dynein and its cofactor dynactin. Significant advances have been made in recent

years characterizing human dynein-dynactin structure and function using in vitro assays,

however, there is limited knowledge about the motile properties and functional organization of

dynein-dynactin in living human cells. Total internal reflection fluorescence microscopy (TIRFM)

of CRISPR-engineered human cells is employed here to visualize fluorescently tagged dynein

heavy chain (DHC) and p50 with high spatio-temporal resolution. We find that p50 and DHC

exhibit indistinguishable motility properties in their velocities, run lengths, and run times. The

dynein-dynactin complexes are fast (~1.2 μm/s) and typically run for several microns (~2.7 μm).

Quantification of the fluorescence intensities of motile puncta reveals that dynein-dynactin runs

are mediated by at least one DHC dimer while the velocity is consistent with that measured for

double dynein (two DHC dimers) complexes in vitro.

INTRODUCTION

The eukaryotic microtubule cytoskeleton plays a critical role in the organization, positioning and motility of organelles, mRNA, and proteins. Intracellular motility is mediated by molecular motor proteins that move cargoes along polarized microtubule tracks (for reviews see (Cason and Holzbaur, 2022; Reck-Peterson et al., 2018). The kinesin superfamily includes motors that are responsible for plus-end directed transport, and others that contribute to minus-end motility and regulation of microtubule dynamics (Hirokawa et al., 2009). In contrast, cytoplasmic dynein 1 motor complexes (hereafter dynein) (Canty et al., 2021; Pfister et al., 2005) are exclusively minus-end directed. The dynein complex is composed of two catalytic heavy chains, and additional light, light intermediate and intermediate chains (Carter et al., 2016; Cason and Holzbaur, 2022). The isolated dynein motor complex displays predominantly diffuse motility along microtubules in vitro, an observation that led to the identification of the dynactin complex, which is important for dynein motility (Gill et al., 1991; McKenney et al., 2014; Schlager et al., 2014; Schroer and Sheetz, 1991). In addition, a number of adaptor complexes that link dynein and dynactin to specific cargoes and activate the motor have been identified and characterized (Canty et al., 2021; Cason and Holzbaur, 2022; Reck-Peterson et al., 2018; Splinter et al., 2012).

Vesicular transport has been directly visualized by live-cell imaging in diverse cells including highly polarized axons (Canty et al., 2021). Motility of vesicles is bidirectional, both plus- and minus-end directed motors co-purify with cargoes, and adaptor complexes have been shown to associate with both kinesin and dynein motors (Ali et al., 2023; Hancock, 2014; Hendricks et al., 2010; Maeder et al., 2014; Vale, 1987). Together these observations indicate that the number and activity of cargo associated motors must be regulated to ensure efficient delivery to appropriate cellular destinations (Cason and Holzbaur, 2022). In vitro studies in which the number of motors bound to artificial cargoes can be precisely controlled demonstrate that cargoes with both plus and minus motors often stall or pause, or show directed motion that is distinct from either individual motor acting alone; however, once directed motion is initiated, reversals are

infrequent (Belyy et al., 2016; Derr et al., 2012). Cryo-EM of dynein-dynactin complexes revealed

that the stoichiometry of dynein relative to dynactin is determined by the activating adaptors that

link them. Specifically, the adaptor BICD2 recruited a single dynein to dynactin while BICDR1 and

HOOK3 supported assembly of a "double dynein" complex (Chaaban and Carter, 2022; Grotjahn

et al., 2018; Urnavicius et al., 2018; Urnavicius et al., 2015). Interestingly, the processivity and

velocity of dynein-dynactin differs depending on the number of dyneins such that the double

dynein complexes assembled with BICDR1 and HOOK3 had a higher frequency of processive

motility events and moved faster than the single dynein complexes assembled with BICD2

(Urnavicius et al., 2018).

In contrast to the relative ease of visualizing motor proteins in in vitro motility assays, the

crowded three-dimensional intracellular environment poses a challenge for imaging and

quantifying motility events in living cells. In human cells constitutively over-expressing a GFP-

tagged intermediate chain of dynein both dot-like, bidirectional motility and comet-like motile

events, which co-localized with EB1, were observed, confirming a role for dynein in both tip-

tracking and vesicle motility (Kobayashi and Murayama, 2009). Quantification of organelle motility

in the highly polarized filamentous fungus *Ustilago maydis* expressing endogenously tagged

fluorescent dynein revealed that binding of a single dynein to an anterograde-directed early

endosome was sufficient to trigger directional reversal of its transport (Schuster et al., 2011).

In this study, HeLa cells expressing CRISPR/Cas9 modified dynein heavy chain (DHC) or

the p50 subunit of dynactin were visualized using high-resolution fluorescence microscopy

approaches to gain insights into the behaviors of dynein and dynactin in vivo. As a result, we were

able to directly observe and quantify core motility parameters of dynein-dynactin while analyses

of their fluorescence intensities relative to a known standard was informative of the stoichiometry

of dynein in motile dynein-dynactin complexes in proliferating human cells.

RESULTS AND DISCUSSION

Human HeLa cells were engineered using CRISPR/Cas9 to insert a cassette encoding FKBP and EGFP tags in frame at the 3' end of the dynein heavy chain (DYNC1H1) gene (SF1). A clonal DHC-EGFP-expressing HeLa cell line was generated and subjected to live-cell fluorescence imaging to directly visualize DHC activities in interphase cells. The most striking dynein behavior was tip-tracking on polymerizing MTs as an abundance of DHC comets traveling at the speed of MT polymerization (Cassimeris et al., 1988; Rusan et al., 2001; Walker et al., 1988) were readily observed by both spinning disc confocal microscopy and total internal reflection fluorescence microscopy (TIRFM) (**Fig. 1 A, B, Videos 1 and 2**). SiR-Tubulin was next introduced to visualize DHC localization and behaviors relative to MTs (**Fig. 1 C**). Interestingly, because SiR-Tubulin is a docetaxol derivative, its addition suppressed plus-end MT polymerization resulting in a significant reduction in the DHC tip-tracking population and a much clearer view of a different population of MT-associated DHC puncta (**Fig. 1 C**). The SiR-Tubulin-treated cells were subjected to two-color TIRFM to determine if the DHC puncta exhibited motility and; indeed, puncta were observed streaming along MTs, which was especially evident via TIRFM on MTs that were pinned between the nucleus and the plasma membrane (**Video 3**). DHC puncta were next visualized with higher temporal resolution by acquiring single color TIRFM time-lapses of the EGFP channel at a rate of 5 frames per second (fps) (**Fig. 1 D, Video 4**). The TIRFM time-lapses were assessed by eye and several parameters of the motile puncta were measured using kymographs (**Fig. 1 E**). The average DHC puncta moved at 1.2 ± 0.05 (mean ± SEM) μm/sec over a distance of 2.8 ± 0.2 μm for 2.6 ± 0.2 seconds (**Fig. 1 F-H**). While some motile puncta appeared to switch "tracks" at MT intersections, which was inferred from observations of sudden high angle turns, directional switches on the same MT were infrequent (~5-10% of runs). It should also be noted that the run lengths and run times for DHC are likely underestimates as the puncta could only be tracked while they remained in the TIRFM evanescent field and prior to their photobleaching.

Activation of dynein motility requires its interaction with the dynactin complex (Chowdhury

et al., 2015; Grotjahn et al., 2018; Urnavicius et al., 2018; Zhang et al., 2017). CRISPR/Cas9-

based genomic engineering was used to insert the cassette encoding FKBP and EGFP tags at

the 3' end of the p50/dynamitin gene and a clonal p50-EGFP-expressing HeLa cell line was

subjected to live-cell fluorescence imaging to visualize dynactin in interphase cells. Consistent

with prior observations of dynactin localization (Vaughan et al., 1999; Vaughan et al., 2002) and

like dynein, dynactin exhibited robust tip-tracking activity on growing MT plus-ends that was

clearly visualized via both spinning disc confocal imaging and TIRFM (**Fig. 2 A, B; Videos 5 and**

6). Suppression of plus-end polymerization dynamics upon introduction of SiR-Tubulin caused a

loss of the tip-tracking pool of dynactin, which made a second pool of motile, MT-associated

puncta of p50 more pronounced especially when visualized with TIRFM (**Fig. 2 C, Video 7**). The

p50-EGFP-expressing cells were then subjected to single color TIRFM at 5 fps and kymograph

analysis to measure the motility parameters of the dynactin complex (**Fig. 2 D**). The mean velocity

of the p50 puncta was 1.2 ± 0.07 μm/sec (**Fig. 2 E**) while the mean run length was 2.6 ± 0.2 μm

(**Fig. 2 F**) and the mean run time was 2.2 ± 0.2 seconds (**Fig. 2 G**). Similar to dynein, dynactin

was observed to switch MT "tracks" but infrequently switched direction.

In comparing the motility parameters of the motile dynein and dynactin puncta, their

velocities (**Fig. 3 A**), run lengths (**Fig. 3 B**), and run times (**Fig. 3 C**) were statistically

indistinguishable. Thus, we inferred that the motile population of dynein was associated with the

dynactin complex. The fluorescence intensities of motile DHC and p50 puncta were next

compared to EGFP-tagged kinesin-1 dimer as a standard to assess the stoichiometries of the

motile dynein and dynactin complexes (**Fig. 3 D-F**). Interestingly, the intensity of motile DHC-

EGFP puncta was not statistically significantly different from the intensity of motile kinesin-1-

EGFP dimers; however, the intensity of motile p50 puncta had a mean fluorescence intensity that

was about half that of the motile kinesin-1-EGFP puncta (**Fig. 3 G**). Thus, the motile dynein and

dynactin complexes visualized here via high-speed TIRFM were comprised, on average, of two

EGFP-tagged DHC molecules and a single p50-EGFP molecule. When considering how these

values relate to physiological stoichiometries of the motile dynein-dynactin complexes, it is

important to note that both the DHC-EGFP and p50-EGFP HeLa cell line clones are

heterozygotes (**Fig. 3 H, I**). Dynein motility requires DHC dimerization and if there is an equal

likelihood of EGFP-tagged and untagged DHC incorporation into a functional dimer then the

motile dynein puncta we visualized would actually contain two DHC dimers. While we were unable

to assess the relative expression levels of tagged versus untagged DHC for technical reasons,

we can confidently conclude from our data that there is at least one DHC dimer in the motile

puncta. Interestingly, western blotting of cell extract prepared from the p50-EGFP clone revealed

that the cells expressed a ~5-6-fold molar excess of the untagged p50 compared to p50-EGFP

(**Fig. 3 I**). Differential allele regulation has been observed for endogenously tagged proteins

(Mann and Wadsworth, 2018; Roberts et al., 2017) suggesting that regulation of gene expression

may help avoid deleterious effects when cells can tolerate a fraction of the total protein pool being

modified. The relative levels of EGFP-tagged versus untagged p50 most likely explains why motile

p50 puncta only had a single EGFP molecule despite the fact that the dynactin complex is known

to have four copies of p50 (Eckley et al., 1999; Urnavicius et al., 2015). Thus, we conclude that

the motile dynactin complexes visualized here are typically comprised of one p50-EGFP molecule

and 3 untagged copies of p50.

While several groups have recently imaged endogenously tagged dynein in mitotic HeLa

cells and iNeurons (Fellows et al., 2023; Ide et al., 2023), to our knowledge - this is the first direct

visualization of motile endogenously-tagged human dynein-dynactin complexes in proliferating,

non-neuronal cells. Two distinct motile populations of dynein-dynactin exist in human interphase

cells: one pool that tip-tracks on polymerizing MT plus-ends and a second pool that moves

processively at high velocities (~7X faster than the tip-tracking population) along MTs. We

propose that the ~3 μm run lengths measured here are an underestimate of what the dynein-

dynactin complex is capable of due to the relatively short and dynamic MT tracks available to it in

interphase HeLa cells and because motile puncta could only be visualized while they were in the

TIRF field and prior to their photobleaching. Indeed, dynein was recently shown to be incredibly

processive in iNeurons where its mean run length was ~ 35 μm and runs >100 μm were observed

(Fellows et al., 2023). It is possible that the tip-tracking pool of dynein-dynactin is regulated in a

cell-cycle dependent manner because dynein-dynactin tracking events were less robust in mitotic

cells than in interphase cells. The dynein-dynactin pool that exhibited directional motility on MTs

was capable of switching MT tracks but rarely switched directions suggesting that there is not a

constant tug-of-war between dynein-dynactin and kinesins in proliferating cells as compared to

observations of bidirectional vesicular transport in neurons (Hancock, 2014; Hendricks et al.,

2010; Maeder et al., 2014). The low directional switching frequency observed here is consistent

with in vitro studies reconstituting the tug-of-war phenomenon showing that one type of motor

dominates once directional movement begins and that reversal events are rare (Ali et al., 2023;

Belyy et al., 2016; Derr et al., 2012). The velocities measured here in HeLa cells were comparable

to those measured for unopposed dynein and; therefore, did not suggest that there was significant

resistive drag from associated kinesin motors as has been observed in vitro. This suggests that

the cargos, which we presume motile dynein-dynactin puncta are bound to, any kinesins

associated with the cargos are inactive and/or cannot effectively bind the MT lattice during dynein-

dynactin-mediated transport in interphase HeLa cells. Interestingly, directional switching was also

uncommon during retrograde dynein-dynactin movements in iNeuron axons (Fellows et al., 2023)

and in retrograde trafficking of early endosomes in *Ustilago maydis* mediated by dynein-dynactin

containing the HOOK3 adaptor (Bielska et al., 2014; Schuster et al., 2011).

Dynein-dynactin motility is also affected by tubulin modifications (Barisic and Maiato,

2016; Roll-Mecak, 2020; Sirajuddin et al., 2014) most notably by the tyrosination state of α-tubulin

such that dynein-dynactin preferentially associates with tyrosinated MTs to promote the initiation

of motility (McKenney et al., 2016; Nirschl et al., 2016; Peris et al., 2006). Since all of the motility

parameters reported here were measured for dynein-dynactin walking on docetaxol-stabilized

MTs it is likely that the MT lattice was more detyrosinated than normal in our experimental

conditions (Gundersen et al., 1987; Webster et al., 1990). Thus, it would be informative to apply

our high resolution TIRFM imaging of dynein-dynactin in cells where tubulin modifications –

starting with the tyrosination state – can be experimentally manipulated (Barisic et al., 2015;

Ferreira et al., 2020).

Finally, our cell-based data are consistent with recent in vitro characterizations of dynein-

dynactin structure, function, and regulation (Belyy et al., 2016; Chaaban and Carter, 2022;

Urnavicius et al., 2018). Based on our fluorescence intensity measurements, we favor the

interpretation that the motile dynein-dynactin complexes visualized here are typically comprised

of a single dynactin complex bound to a tandem array of two dyneins. Our measured dynein-

dynactin velocity of 1.2 μm/sec further supports this conclusion since this speed is consistent with

in vitro velocities of dynein-dynactin complexes containing activating adaptors that recruit two

dyneins although we do not exclude that the range of DHC velocities and intensities measured

here may include sub-populations of dynein-dynactin containing a single dynein dimer. It would

be valuable to test whether dynein-dynactin regulation mechanisms that have been characterized

in vitro also apply to the physiological regulation of dynein-dynactin activity in living cells by

measuring DHC motility parameters and intensities in cells depleted of different cargo adaptors.

Ultimately, we hope that the direct visualization of human dynein-dynactin and quantification of

its motility parameters and stoichiometries in living cells will be an important physiological

complement to in vitro assays, which altogether will better inform mechanistic models of the many

cellular processes that rely on dynein-dynactin.

MATERIALS AND METHODS

Cell culture

Parental HeLa and CRISPR-edited HeLa clones expressing DHC-EGFP or p50-EGFP were grown in standard DMEM medium (Gibco, USA) supplemented with 10% non-heat-inactivated FBS (Gibco, USA) and 0.5× antibiotic/antimycotic cocktail (Sigma-Aldrich) and maintained at 37°C with 5% CO2. Human-Kinesin-1-EGFP-expressing *Drosophila* S2 cells were grown in Schneider's medium (Life Technologies) supplemented with 10% heat inactivated fetal bovine serum (FBS) and 0.5x antibiotic/antimycotic cocktail (Sigma), and maintained at 25°C. Kinesin-1-EGFP expression was induced with 500µM $CuSO_4$ for 16-18 hours.

CRISPR-engineered cell line production

CRISPR gene editing.

FKBP-EGFP tags were added to the C-terminus of human dynein heavy chain (DHC) and p50 using methods described previously (Sheridan and Bentley, 2016; Stewart-Ornstein and Lahav, 2016). In brief, repair cassettes comprised of FKBP-EGFP linked to a cleavable peptide (T2A) followed by a selectable marker (Neomycin) were cloned into pCMV and used for PCR reactions. Glycine-Alanine linkers were included between proteins in the construct. Guide sequences were selected using the CRISPR design tool (http://crispr.mit.edu/) from the Zhang lab at MIT (Ran et al., 2013) using "other regions" and the human target genome (hg19). The search tool was used to select guides close to the C-terminus of the protein of interest (~100nt surrounding and including the stop codon). Top and bottom oligos were obtained for each guide with the bases 5'-CACC -3'added to the top oligo and 5'-AAAC-3' added to the complement of the bottom oligo.

Guides were cloned into a Cas9 containing plasmid (PX459) Addgene #62988, (Cambridge, MA) following methods previously outlined (Moyer and Holland, 2015). Top and bottom oligos were annealed and then phosphorylated by T4 PNK (NEB, Ipswich, MA). Guides were then cloned into PX459 that was cut using BbsI (NEB, Ipswich, MA) and ligated using T4

ligase (NEB Ipswich, MA). Guide-Cas9 containing plasmids were then sequenced using the U6

promotor primer (Ran et al., 2013) and purified using either endotoxin free mini-preps or midi-

preps according to manufacturer protocol (Promega, Madison, WI). Repair cassettes were

amplified using primers designed to be homologous to the C-terminal genomic DNA surrounding

the STOP codon. In all cases, the guide target sequence was mutated to prevent Cas9 from

recognizing the repair cassette.

Cells were grown in DMEM medium (ThermoFisher Scientific) with 10% Fetal bovine

serum (FBS, Atlanta Biologicals, Flowery Branch, GA) and 0.5X antibiotic/antimycotic solution

(final concentrations 50 U/mL penicillin, 0.05 mg/mL streptomycin, 0.125 ug/mL amphotericin B;

Sigma-Aldrich, St. Louis, MO) at 37 °C and 5% carbon dioxide (CO_2). For long term storage cells

were frozen in DMEM medium with 5% FBS and 0.5X antibiotic/antimycotic solution and 15%

DMSO and held at -80°C for 1-2 days before moving to liquid nitrogen.

To generate CRISPR modified cell lines, parental cells were nucleofected using an Amaxa

Nucleofector (Lonza, Portsmouth, NH) program I-013 and Mirus nucleofection reagent (Mirus Bio

LLC, Madison, WI) according to the manufacturer's recommendations. Plasmids and Repair

cassettes were used at ratio of 1:1 at a concentration of 1 µg DNA each. Following nucleofection,

cells were grown in regular growth media in 100mm dishes for 48-72 Hrs. and then 0.2g/L

Neomycin/G418 (InvivoGen, San Diego, CA) selection media was added. Media was then

changed daily for 10-14 days and colonies of green cells were picked using cloning rings and

returned to regular media for further screening and experiments.

Genotyping

Genomic DNA was isolated from clonal CRISPR tagged cells using Genomic DNA Mini Kit

(Invitrogen) according to manufacturer's recommendations. DNA was then amplified by PCR

using genomic primers targeting the Neomycin cassette- and downstream of the Stop codon.

Phusion polymerase (NEB) was used to amplify 1 µL of isolated genomic DNA in a 50 µL reaction

for 30 cycles. The resulting PCR products were analyzed using 0.8% agarose gel

electrophoresis. The expected band was excised, and gel extracted using Qiagen gel extraction

kit and sequenced to verify proper integration of the tag.

Live-cell TIRF microscopy (TIRFM)

Cells were seeded onto 35 mm glass bottom petri dishes (Cellvis) 2-3 days prior to imaging. With

the exception of the visualization of tip-tracking DHC-EGFP, cells were incubated in MEDIA

supplemented with 1 μM SiR-Tubulin (Cytoskeleton, Inc) for 30-60 minutes prior to washing out

the SiR-Tubulin with fresh DMEM before imaging. The cells were visualized on a Nikon Ti-E

microscope equipped with a TIRF illuminator, a 100X 1.49-NA differential interference contrast

TIRF Apochromat oil-immersion objective, Nikon perfect focus system (PFS), a Hamamatsu

ORCA-Flash 4.0 LT digital complementary metal-oxide semiconductor camera (C11440), four

laser lines (447, 488, 561, and 641 nm), and MetaMorph software (Molecular Devices). Flat

interphase cells were identified and the initial focal plane was set by focusing on the far-red SiR

tubulin signal with a particular emphasis on finding cells in which individual microtubules could be

seen between the nucleus and the glass-adhered plasma membrane. Individual snap shots were

then taken on the EGFP channel to refine the focal plane on the DHC or p50 puncta. Once

individual puncta were well-resolved the PFS was engaged and the cell was subjected to

streaming TIRFM for 12 seconds at an acquisition rate of 5 frames per second (200 ms exposure)

and 2x2 camera binning.

Quantifications of motility parameters and fluorescence intensities

High temporal (5 fps) TIRFM time-lapses were examined by eye to identify evident motile puncta,

the paths of which were traced manually in MetaMorph using the Multi-Line drawing function. A

kymograph of the line segment with a 5-pixel line width was then generated and the velocity was

determined by measuring the slope of the motility event on the kymograph from start to end using

the Single Line drawing function. The run time was measured by defining the start and end frame

of evident movement in the TIRFM time-lapse and the run length was then calculated by

multiplying the run time by the velocity measured in the kymograph of the motility event. The run

times and run lengths are likely underestimates of what dynein-dynactin can achieve due to the

fact that the puncta sometimes bleached or exited the TIRFM evanescent field.

The number of EGFP molecules per motile puncta of DHC and p50 was determined by

comparing their background-corrected fluorescence intensities to that of a known dimer standard:

human kinesin-1 tagged with EGFP. As previously described (Ye et al., 2018) *Drosophila* S2 cells

expressing inducible human Kinesin-1 (in this case tagged with EGFP) were induced with 500µM

CuSO4 for 16 hours to induce expression. The next day, the induced S2 cells were seeded onto

a concanavalin A coated glass bottom petri dish and allowed to adhere for ~1 hour prior to

visualization by TIRFM. The DHC-EGFP and p50-EGFP HeLa clones and the human Kinesin-1-

EGFP-expressing S2 cells were each visualized sequentially via live-cell TIRF microscopy using

identical imaging parameters (5 fps acquisition rate, 200 ms exposure, 2X2 binning) within the

same region of the camera chip. The region-in-region method (Ye and Maresca, 2018) was then

used to background correct and quantify the integrated fluorescence intensities of motile puncta

of DHC-, p50-, and kinesin-1-EGFP in the same TIRFM time-lapses that were used to measure

the motility parameters. The intensity of each motile puncta was measured for a single time-point

from each run in which the spot was most clearly resolved and did not have significant local

background signal from nearby EGFP puncta.

Western blotting

Twenty μg of total protein from cell lysates prepared from parental HeLa cells or the p50-EGFP

clone were loaded onto a 10% SDS-PAGE gel, run out, and transferred to a nitrocellulose

membrane on the Trans-Blot® Turbo™ transfer system (Bio-Rad Laboratories) using the

preprogrammed "HIGH MW" 10-minute protocol. After blocking for 1 hour in TBS with 0.1% Tween

and 5% milk, the membrane was incubated overnight at 4°C with mouse anti-dynactin p50 (BD

Transduction Laboratories, Cat. # 611002) diluted at 1:1000 in the block. The membrane was

washed 3X 5 minutes in TBS + 0.1% Tween and then incubated for 1 hour in milk containing

donkey-anti-mouse IgG secondary antibodies conjugated with HRP (Jackson ImmunoResearch

Laboratories, Inc.) diluted at 1:5000 in block. Following 3X 5 minutes washes in TBS + 0.1%

Tween the blot was incubated with Immobilon Western Chemiluminescent HRP Substrate

(Millipore) according to the manufacturer's recommendations. Imaging of the membrane was

done on a G:Box system controlled by GeneSnap software (Syngene).

Statistical analyses

Statistical analyses reporting p-values from two-tailed Student's t-tests were done in excel while

p-values generated with the randomization method were done using the PlotsofDifferences web

tool at: https://huygens.science.uva.nl/PlotsOfDifferences (Goedhart, 2019). PlotsofDifferences

does not rely on assumptions about the distribution of the data (normal versus non-normal) to

calculate p-values.

ACKNOWLEDGMENTS

We are grateful to Barbara Mann for generating the CRISPR-engineered HeLa cell lines. This

work was supported by an NIH grant (GM107026) to T.J.M. and by an NSF grant (MCB1817926)

to P.W.

CONFLICT OF INTEREST

The authors declare no competing financial interests.

REFERENCES

Ali, F.A., A.J. Zwetsloot, C.E. Stone, T.E. Morgan, R.F. Wademan, A.P. Carter, and A. Straube. 2023. KIF1C activates and extends dynein movement through the FHF cargo adaptor. *bioRxiv*:2023.2010.2026.564242.

Barisic, M., and H. Maiato. 2016. The Tubulin Code: A Navigation System for Chromosomes during Mitosis. *Trends Cell Biol*. 26:766-775.

Barisic, M., R. Silva e Sousa, S.K. Tripathy, M.M. Magiera, A.V. Zaytsev, A.L. Pereira, C. Janke, E.L. Grishchuk, and H. Maiato. 2015. Mitosis. Microtubule detyrosination guides chromosomes during mitosis. *Science*. 348:799-803.

Belyy, V., M.A. Schlager, H. Foster, A.E. Reimer, A.P. Carter, and A. Yildiz. 2016. The mammalian dynein-dynactin complex is a strong opponent to kinesin in a tug-of-war competition. *Nat Cell Biol*. 18:1018-1024.

Bielska, E., M. Schuster, Y. Roger, A. Berepiki, D.M. Soanes, N.J. Talbot, and G. Steinberg. 2014. Hook is an adapter that coordinates kinesin-3 and dynein cargo attachment on early endosomes. *J Cell Biol*. 204:989-1007.

Canty, J.T., R. Tan, E. Kusakci, J. Fernandes, and A. Yildiz. 2021. Structure and Mechanics of Dynein Motors. *Annu Rev Biophys*. 50:549-574.

Carter, A.P., A.G. Diamant, and L. Urnavicius. 2016. How dynein and dynactin transport cargos: a structural perspective. *Curr Opin Struct Biol*. 37:62-70.

Cason, S.E., and E.L.F. Holzbaur. 2022. Selective motor activation in organelle transport along axons. *Nat Rev Mol Cell Biol*. 23:699-714.

Cassimeris, L., N.K. Pryer, and E.D. Salmon. 1988. Real-time observations of microtubule dynamic instability in living cells. *J Cell Biol*. 107:2223-2231.

Chaaban, S., and A.P. Carter. 2022. Structure of dynein-dynactin on microtubules shows tandem adaptor binding. *Nature*. 610:212-216.

Chowdhury, S., S.A. Ketcham, T.A. Schroer, and G.C. Lander. 2015. Structural organization of the dynein-dynactin complex bound to microtubules. *Nat Struct Mol Biol*. 22:345-347.

Derr, N.D., B.S. Goodman, R. Jungmann, A.E. Leschziner, W.M. Shih, and S.L. Reck-Peterson. 2012. Tug-of-war in motor protein ensembles revealed with a programmable DNA origami scaffold. *Science*. 338:662-665.

Eckley, D.M., S.R. Gill, K.A. Melkonian, J.B. Bingham, H.V. Goodson, J.E. Heuser, and T.A. Schroer. 1999. Analysis of dynactin subcomplexes reveals a novel actin-related protein associated with the arp1 minifilament pointed end. *J Cell Biol*. 147:307-320.

Fellows, A.D., M. Bruntraeger, T. Burgold, A.R. Bassett, and A.P. Carter. 2023. Dynein and dynactin move long-range but are delivered separately to the axon tip. *bioRxiv*:2023.2007.2003.547521.

Ferreira, L.T., B. Orr, G. Rajendraprasad, A.J. Pereira, C. Lemos, J.T. Lima, C. Guasch Boldú, J.G. Ferreira, M. Barisic, and H. Maiato. 2020. α-Tubulin detyrosination impairs mitotic error correction by suppressing MCAK centromeric activity. *J Cell Biol*. 219.

Gill, S.R., T.A. Schroer, I. Szilak, E.R. Steuer, M.P. Sheetz, and D.W. Cleveland. 1991. Dynactin, a conserved, ubiquitously expressed component of an activator of vesicle motility mediated by cytoplasmic dynein. *J Cell Biol*. 115:1639-1650.

Goedhart, J. 2019. PlotsOfDifferences – a web app for the quantitative comparison of unpaired data. *bioRxiv*:578575.

Grotjahn, D.A., S. Chowdhury, Y. Xu, R.J. McKenney, T.A. Schroer, and G.C. Lander. 2018. Cryo-electron tomography reveals that dynactin recruits a team of dyneins for processive motility. *Nat Struct Mol Biol*. 25:203-207.

Gundersen, G.G., S. Khawaja, and J.C. Bulinski. 1987. Postpolymerization detyrosination of alpha-tubulin: a mechanism for subcellular differentiation of microtubules. *J Cell Biol*. 105:251-264.

Hancock, W.O. 2014. Bidirectional cargo transport: moving beyond tug of war. *Nat Rev Mol Cell Biol*. 15:615-628.

Hendricks, A.G., E. Perlson, J.L. Ross, H.W. Schroeder, 3rd, M. Tokito, and E.L. Holzbaur. 2010. Motor coordination via a tug-of-war mechanism drives bidirectional vesicle transport. *Curr Biol*. 20:697-702.

Hirokawa, N., Y. Noda, Y. Tanaka, and S. Niwa. 2009. Kinesin superfamily motor proteins and intracellular transport. *Nat Rev Mol Cell Biol*. 10:682-696.

Ide, A.H., K.F. DeLuca, O. Wiggan, S.M. Markus, and J.G. DeLuca. 2023. The role of kinetochore dynein in checkpoint silencing is restricted to disassembly of the corona. *Mol Biol Cell*. 34:ar76.

Kobayashi, T., and T. Murayama. 2009. Cell cycle-dependent microtubule-based dynamic transport of cytoplasmic dynein in mammalian cells. *PLoS One*. 4:e7827.

Maeder, C.I., K. Shen, and C.C. Hoogenraad. 2014. Axon and dendritic trafficking. *Curr Opin Neurobiol*. 27:165-170.

Mann, B.J., and P. Wadsworth. 2018. Distribution of Eg5 and TPX2 in mitosis: Insight from CRISPR tagged cells. *Cytoskeleton (Hoboken)*. 75:508-521.

McKenney, R.J., W. Huynh, M.E. Tanenbaum, G. Bhabha, and R.D. Vale. 2014. Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. *Science*. 345:337-341.

McKenney, R.J., W. Huynh, R.D. Vale, and M. Sirajuddin. 2016. Tyrosination of α-tubulin controls the initiation of processive dynein-dynactin motility. *Embo j*. 35:1175-1185.

Moyer, T.C., and A.J. Holland. 2015. Generation of a conditional analog-sensitive kinase in human cells using CRISPR/Cas9-mediated genome engineering. *Methods Cell Biol*. 129:19-36.

Nirschl, J.J., M.M. Magiera, J.E. Lazarus, C. Janke, and E.L. Holzbaur. 2016. α-Tubulin Tyrosination and CLIP-170 Phosphorylation Regulate the Initiation of Dynein-Driven Transport in Neurons. *Cell Rep*. 14:2637-2652.

Peris, L., M. Thery, J. Fauré, Y. Saoudi, L. Lafanechère, J.K. Chilton, P. Gordon-Weeks, N. Galjart, M. Bornens, L. Wordeman, J. Wehland, A. Andrieux, and D. Job. 2006. Tubulin tyrosination is a major factor affecting the recruitment of CAP-Gly proteins at microtubule plus ends. *J Cell Biol*. 174:839-849.

Pfister, K.K., E.M. Fisher, I.R. Gibbons, T.S. Hays, E.L. Holzbaur, J.R. McIntosh, M.E. Porter, T.A. Schroer, K.T. Vaughan, G.B. Witman, S.M. King, and R.B. Vallee. 2005. Cytoplasmic dynein nomenclature. *J Cell Biol*. 171:411-413.

Ran, F.A., P.D. Hsu, J. Wright, V. Agarwala, D.A. Scott, and F. Zhang. 2013. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc*. 8:2281-2308.

Reck-Peterson, S.L., W.B. Redwine, R.D. Vale, and A.P. Carter. 2018. The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol*. 19:382-398.

Roberts, B., A. Haupt, A. Tucker, T. Grancharova, J. Arakaki, M.A. Fuqua, A. Nelson, C. Hookway, S.A. Ludmann, I.A. Mueller, R. Yang, R. Horwitz, S.M. Rafelski, and R.N. Gunawardane. 2017. Systematic gene tagging using CRISPR/Cas9 in human stem cells to illuminate cell organization. *Mol Biol Cell*. 28:2854-2874.

Roll-Mecak, A. 2020. The Tubulin Code in Microtubule Dynamics and Information Encoding. *Dev Cell*. 54:7-20.

Rusan, N.M., C.J. Fagerstrom, A.M. Yvon, and P. Wadsworth. 2001. Cell cycle-dependent changes in microtubule dynamics in living cells expressing green fluorescent protein-alpha tubulin. *Mol Biol Cell*. 12:971-980.

Schlager, M.A., H.T. Hoang, L. Urnavicius, S.L. Bullock, and A.P. Carter. 2014. In vitro reconstitution of a highly processive recombinant human dynein complex. *Embo j*. 33:1855-1868.

Schroer, T.A., and M.P. Sheetz. 1991. Two activators of microtubule-based vesicle transport. *J Cell Biol*. 115:1309-1318.

Schuster, M., R. Lipowsky, M.A. Assmann, P. Lenz, and G. Steinberg. 2011. Transient binding of
dynein controls bidirectional long-range motility of early endosomes. *Proc Natl Acad Sci*
U S A. 108:3618-3623.
Sheridan, R.M., and D.L. Bentley. 2016. Selectable one-step PCR-mediated integration of a
degron for rapid depletion of endogenous human proteins. *Biotechniques*. 60:69-74.
Sirajuddin, M., L.M. Rice, and R.D. Vale. 2014. Regulation of microtubule motors by tubulin
isotypes and post-translational modifications. *Nat Cell Biol*. 16:335-344.
Splinter, D., D.S. Razafsky, M.A. Schlager, A. Serra-Marques, I. Grigoriev, J. Demmers, N.
Keijzer, K. Jiang, I. Poser, A.A. Hyman, C.C. Hoogenraad, S.J. King, and A. Akhmanova.
2012. BICD2, dynactin, and LIS1 cooperate in regulating dynein recruitment to cellular
structures. *Mol Biol Cell*. 23:4226-4241.
Stewart-Ornstein, J., and G. Lahav. 2016. Dynamics of CDKN1A in Single Cells Defined by an
Endogenous Fluorescent Tagging Toolkit. *Cell Rep*. 14:1800-1811.
Urnavicius, L., C.K. Lau, M.M. Elshenawy, E. Morales-Rios, C. Motz, A. Yildiz, and A.P. Carter.
2018. Cryo-EM shows how dynactin recruits two dyneins for faster movement. *Nature*.
554:202-206.
Urnavicius, L., K. Zhang, A.G. Diamant, C. Motz, M.A. Schlager, M. Yu, N.A. Patel, C.V. Robinson,
and A.P. Carter. 2015. The structure of the dynactin complex and its interaction with
dynein. *Science*. 347:1441-1446.
Vale, R.D. 1987. Intracellular Transport Using Microtubule-Based Motors. *Annual Review of Cell*
Biology. 3:347-378.
Vaughan, K.T., S.H. Tynan, N.E. Faulkner, C.J. Echeverri, and R.B. Vallee. 1999. Colocalization
of cytoplasmic dynein with dynactin and CLIP-170 at microtubule distal ends. *J Cell Sci*.
112 (Pt 10):1437-1447.
Vaughan, P.S., P. Miura, M. Henderson, B. Byrne, and K.T. Vaughan. 2002. A role for regulated
binding of p150(Glued) to microtubule plus ends in organelle transport. *J Cell Biol*.
158:305-319.
Walker, R.A., E.T. O'Brien, N.K. Pryer, M.F. Soboeiro, W.A. Voter, H.P. Erickson, and E.D.
Salmon. 1988. Dynamic instability of individual microtubules analyzed by video light
microscopy: rate constants and transition frequencies. *J Cell Biol*. 107:1437-1448.
Webster, D.R., J. Wehland, K. Weber, and G.G. Borisy. 1990. Detyrosination of alpha tubulin
does not stabilize microtubules in vivo. *J Cell Biol*. 111:113-122.
Ye, A.A., and T.J. Maresca. 2018. Measuring mitotic forces. *Methods Cell Biol*. 144:165-184.
Ye, A.A., V. Verma, and T.J. Maresca. 2018. NOD is a plus end-directed motor that binds EB1
via a new microtubule tip localization sequence. *J Cell Biol*. 217:3007-3017.
Zhang, K., H.E. Foster, A. Rondelet, S.E. Lacey, N. Bahi-Buisson, A.W. Bird, and A.P. Carter.
2017. Cryo-EM Reveals How Human Cytoplasmic Dynein Is Auto-inhibited and Activated.
Cell. 169:1303-1314.e1318.

 FIGURE LEGENDS

Figure 1. Dynein tip-tracks on polymerizing MTs and walks processively at high velocities

in interphase HeLa cells. (A) Representative spinning disc confocal time-lapse of a DHC-EGFP-

expressing HeLa cell showing robust tip-tracking. Zoomed views of the numbered boxed regions

are shown to the right with kymographs of the tip-tracking events highlighted with yellow arrows

in the zoomed panels. **(B)** Representative TIRFM time-lapse of a DHC-EGFP-expressing HeLa

cell showing the tip-tracking population. Zoomed views of the numbered boxed regions are shown

with kymographs of the tip-tracking events highlighted with yellow arrows. **(C)** Still frame from a

representative TIRFM time-lapse of a DHC-EGFP-expressing HeLa cell treated with SiR-Tubulin.

In the merge image DHC is green and Sir-Tubulin labeled MTs are magenta. **(D)** Still frame from

a representative high temporal resolution (5 fps) TIRFM time-lapse of a DHC-EGFP-expressing

HeLa cell treated with SiR-Tubulin. Boxed region is shown as a zoomed inset in the lower panel

with the track of a motile DHC puncta highlighted in green. **(E)** Representative kymographs of

motile puncta spanning the range of measured velocities (Vel) and run lengths (RL). **(F)**

Distribution of DHC velocities (n = 100 puncta). **(G)** Distribution of DHC run lengths (n = 81

puncta). **(H)** Distribution of DHC run times (n = 81 puncta). Scale bars, 10 μm (A-D); 1 μm (all

insets), 10 μm (horizontal); 1 min (vertical) in the kymographs in A and B; and 1 μm (horizontal);

1 second (vertical) in E. Displayed times are min:sec.

Figure 2. The dynactin complex component p50 tip-tracks on polymerizing MTs and walks

processively at high velocities in interphase HeLa cells. (A) Still frames from a representative

spinning disc confocal time-lapse of a p50-EGFP-expressing HeLa cell. Zoomed view of the

boxed region is shown with a kymograph of the tip-tracking event highlighted with the yellow

arrow. **(B)** Still frames from a representative TIRFM time-lapse of a p50-EGFP-expressing HeLa

cell showing the tip-tracking population. Zoomed view of the boxed region is shown with a

kymograph of the tip-tracking event highlighted by the yellow arrow. **(C)** Still frame from a

representative TIRFM time-lapse of a p50-EGFP-expressing HeLa cell treated with SiR-Tubulin.

In the merge image p50 is green and Sir-Tubulin labeled MTs are magenta. **(D)** Representative

kymographs of motile p50 puncta spanning the range of measured velocities (Vel) and run lengths

(RL). **(E)** Distribution of p50 velocities (n = 44 puncta). **(F)** Distribution of p50 run lengths (n = 41

puncta). **(G)** Distribution of p50 run times (n = 41 puncta). Scale bars, 10 μm (A-C); 1 μm (all

insets), 5 μm (horizontal); 1 min (vertical) in the kymographs in A and B; and 1 μm (horizontal); 1

second (vertical) in D. Displayed times are min:sec.

Figure 3. Motile DHC and p50 puncta exhibit identical motility parameters but different

fluorescent intensities. (A-C) Box and whisker plots of DHC and p50 velocities (A), run lengths

(B), and run times (C). **(D-F)** Distributions of kinesin-1-EGFP (D), DHC-EGFP (E), and p50-EGFP

background corrected fluorescence intensities. The dashed line in each histogram denotes the

mean value of the kinesin-1-EGFP data set. **(G)** Box and whisker plots of the kinesin-1, DHC, and

p50 fluorescence intensities. **(H)** PCR of genomic DNA from the DCH-EGP clone used in this

study using PCR primers flanking the integration site of the repair cassette. The upper band was

extracted and subjected to sequencing the results of which are shown in Figure 1 – supplemental

figure 1. **(I)** Western blot for p50 of cell lysates from the parental HeLa cell line and the p50-EGFP

clone used in this study. The tagged p50 runs ~30 kDa larger than the untagged p50 and is

expressed at ~5-6-fold lower levels than the endogenous p50. In the box and whisker plots,

whiskers indicate min-max, box is 25^{th}-75^{th} percentile, and bar is median. The reported p-values

were determined by a randomization method: n.s. is not significant (p > 0.05).

SUPPLEMENTAL MATERIALS

Supplemental Figures

Figure 1 – figure supplement 1. Repair cassette for generating the DHC-EGFP CRISPR-

engineered HeLa cell line with sequencing trace showing its integration at the dynein

heavy chain (DYNC1H1) gene locus.

Supplemental Videos

Video 1. Spinning disk confocal time-lapse of a DHC-EGFP CRISPR-engineered interphase

HeLa cell. Zoomed views of the boxed regions in the upper left panel are shown in the right

panels. Displayed times are min:sec. Scale bars, 10 µm (upper left panel) and 1 µm (right panels).

Relates to Fig. 1A.

Video 2. TIRFM time-lapse of a DHC-EGFP CRISPR-engineered interphase HeLa cell.

Displayed times are min:sec. Scale bar, 10 µm. Relates to Fig. 1B.

Video 3. TIRFM time-lapse of a DHC-EGFP (green) CRISPR-engineered interphase HeLa

cell treated with SiR-Tubulin (magenta). Displayed times are min:sec. Scale bar, 10 µm.

Relates to Fig. 1C.

Video 4. High temporal-resolution (5 frames per second) TIRFM time-lapse of a DHC-EGFP

CRISPR-engineered interphase HeLa cell treated with SiR-Tubulin. The boxed regions

highlight motile DHC puncta. Displayed times are sec:millisec. Scale bar, 1 µm. Relates to Fig.

1C.

Video 5. Spinning disk confocal time-lapse of a p50-EGFP CRISPR-engineered interphase

HeLa cell. Displayed times are min:sec. Scale bar, 10 µm. Relates to Fig. 2A.

Video 6. TIRFM time-lapse of a p50-EGFP CRISPR-engineered interphase HeLa cell.

Displayed times are min:sec. Scale bar, 10 µm. Relates to Fig. 2B.

Video 7. TIRFM time-lapse of a p50-EGFP (green) CRISPR-engineered interphase HeLa cell

treated with SiR-Tubulin (magenta). Displayed times are min:sec. Scale bar, 10 µm. Relates to

Fig. 2C.

Figure 1

A **DHC-EGFP** — Spinning disc — 1 — 2 — 0:00 0:36 1:06 1:15 — 1 min — 10 μm — 0:00 0:24 0:36 0:42 0:51

B **DHC-EGFP** — TIRFM — 1 — 2 — 0:00 0:22 0:40 1:30 — 1 min — 10 μm — 0:00 0:08 0:22 0:30 0:42

C **Docetaxol** — DHC-EGFP — SiR-Tubulin — Merge

D DHC-EGFP + SiR Tub (TIRFM)

E **Kymographs**

Vel: Slower — Average — Faster — Average — Slower
RL: Shorter — Shorter — Shorter — Average — Longer

1 s / 1 μm

Vel: Average — Faster
RL: Longer — Longer

F **DHC Velocity** 1.2 ± 0.05 μm/sec — Percentage — microns/sec

G **DHC Run Length** 2.8 ± 0.2 μm — Percentage — microns

H **DHC Run Time** 2.6 ± 0.2 seconds — Percentage — seconds

Figure 2

Figure 3

Figure 1 - figure supplement 1