Human dynein-dynactin is a fast processive motor in living cells

Vikash Verma[1], Patricia Wadsworth[1,2], and Thomas J. Maresca[1,2,]

[1] Biology Department, University of Massachusetts, Amherst

[2] Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst

Correspondence to:

Thomas J. Maresca
Biology Department
University of Massachusetts, Amherst
Amherst, MA 01003
Phone: 413-545-0957
Fax: 413-545-3243
E-mail: tmaresca@umass.edu

Patricia Wadsworth
Biology Department
University of Massachusetts, Amherst
Amherst, MA 01003
Phone: 413-545-4877
Fax: 413-545-3243
E-mail: patw@umass.edu

Running title: Direct visualization of dynein-dynactin motility in human cells

ABSTRACT

Minus-end directed transport along microtubules in eukaryotes is primarily mediated by

cytoplasmic dynein and its cofactor dynactin. Significant advances have been made in recent

years characterizing human dynein-dynactin structure and function using in vitro assays,

however, there is limited knowledge about the motile properties and functional organization of

dynein-dynactin in living human cells. Total internal reflection fluorescence microscopy (TIRFM)

of CRISPR-engineered human cells is employed here to visualize fluorescently tagged dynein

heavy chain (DHC) and p50 with high spatio-temporal resolution. We find that p50 and DHC

exhibit indistinguishable motility properties in their velocities, run lengths, and run times. The

dynein-dynactin complexes are fast (~1.2 µm/s) and run for several microns (~2.7 µm).

Quantification of the fluorescence intensities of motile puncta reveals that dynein-dynactin runs

are mediated by at least one DHC dimer while the velocity is consistent with that measured for

double dynein (two DHC dimers) complexes in vitro.

INTRODUCTION

The eukaryotic microtubule cytoskeleton plays a critical role in the organization, positioning and motility of organelles, mRNA, and proteins. Intracellular motility is mediated by molecular motor proteins that move cargoes along polarized microtubule tracks (for reviews see (Cason and Holzbaur, 2022; Reck-Peterson et al., 2018). The kinesin superfamily includes motors that are responsible for plus-end directed transport, and others that contribute to minus-end motility and regulation of microtubule dynamics (Hirokawa et al., 2009). In contrast, cytoplasmic dynein 1 motor complexes (hereafter dynein) (Canty et al., 2021; Pfister et al., 2005) are exclusively minus- end directed. The dynein complex is composed of two catalytic heavy chains, and additional light, light intermediate and intermediate chains (Carter et al., 2016; Cason and Holzbaur, 2022). The isolated dynein motor complex displays predominantly diffuse motility along microtubules in vitro, an observation that led to the identification of the dynactin complex, which is important for dynein motility (Gill et al., 1991; McKenney et al., 2014; Schlager et al., 2014; Schroer and Sheetz, 1991). In addition, a number of adaptor complexes that link dynein and dynactin to specific cargoes and activate the motor have been identified and characterized (Canty et al., 2021; Cason and Holzbaur, 2022; Reck-Peterson et al., 2018; Splinter et al., 2012).

Vesicular transport has been directly visualized by live-cell imaging in diverse cells including highly polarized axons (Canty et al., 2021). Motility of vesicles is bidirectional, both plus- and minus-end directed motors co-purify with cargoes, and adaptor complexes have been shown to associate with both kinesin and dynein motors (Ali et al., 2023; Hancock, 2014; Hendricks et al., 2010; Maeder et al., 2014; Vale, 1987). Together these observations indicate that the number and activity of cargo associated motors must be regulated to ensure efficient delivery to appropriate cellular destinations (Cason and Holzbaur, 2022). In vitro studies in which the number of motors bound to artificial cargoes can be precisely controlled demonstrate that cargoes with both plus and minus motors often stall or pause, or show directed motion that

is distinct from either individual motor acting alone; however, once directed motion is initiated,

reversals are infrequent (Belyy et al., 2016; Derr et al., 2012). Cryo-EM of dynein-dynactin

complexes revealed that the stoichiometry of dynein relative to dynactin is determined by the

activating adaptors that link them. Specifically, the adaptor BICD2 tends to favor single dynein

and the BICDR1 and HOOK3 tend to favor two dyneins assembled into a "double dynein"

complex (Chaaban and Carter, 2022; Grotjahn et al., 2018; Urnavicius et al., 2018; Urnavicius

et al., 2015). Interestingly, the processivity and velocity of dynein-dynactin differs depending on

the number of dyneins such that the double dynein complexes assembled with BICDR1 and

HOOK3 had a higher frequency of processive motility events and moved faster than the single

dynein complexes assembled with BICD2 (Urnavicius et al., 2018).

In contrast to the relative ease of visualizing motor proteins in vitro, the crowded three-

dimensional intracellular environment poses a challenge for imaging and quantifying motility

events in living cells. Puncta exhibiting bidirectional movement on MTs and comet-like motile

events that co-localized with EB1 were observed in human cells over-expressing a GFP-tagged

dynein intermediate chain, supporting a role for dynein in MT plus-end tip-tracking and vesicle

motility (Kobayashi and Murayama, 2009). Quantification of organelle motility in the highly

polarized filamentous fungus *Ustilago maydis* expressing endogenously tagged fluorescent

dynein revealed that binding of a single dynein to an anterograde-directed early endosome was

sufficient to trigger directional reversal of its transport (Schuster et al., 2011).

In this study, HeLa cells expressing CRISPR/Cas9 modified dynein heavy chain (DHC)

or the p50 subunit of dynactin were visualized using high-resolution fluorescence microscopy

approaches to gain insights into the behaviors of dynein and dynactin in living cells. In doing so,

we were able to directly observe and quantify core motility parameters of dynein-dynactin while

analyses of their fluorescence intensities relative to a known standard was informative of the

stoichiometry of dynein in motile dynein-dynactin complexes in proliferating human cells.

RESULTS AND DISCUSSION

Human HeLa cells were engineered using CRISPR/Cas9 to insert a cassette encoding FKBP and EGFP tags in frame at the 3' end of the dynein heavy chain (DYNC1H1) gene (**Figure 1 - figure supplement 1 A**). A clonal DHC-EGFP-expressing HeLa cell line was generated and subjected to live-cell fluorescence imaging to directly visualize DHC activities in interphase cells (**Figure 1 - figure supplement 1 B**). The most striking dynein behavior was tip-tracking on polymerizing MTs as an abundance of DHC comets traveling at the speed of MT polymerization (Cassimeris et al., 1988; Rusan et al., 2001; Walker et al., 1988) were readily observed by both spinning disc confocal microscopy and total internal reflection fluorescence microscopy (TIRFM) (**Figure 1 A, B, Videos 1 and 2**). SiR-Tubulin was next introduced to visualize DHC localization and behaviors relative to MTs (**Figure 1 C**). Interestingly, because SiR-Tubulin is a docetaxel derivative, its addition suppressed plus-end MT polymerization resulting in a significant reduction in the DHC tip-tracking population and a much clearer view of a different population of MT-associated DHC puncta (**Figure 1 C**). The SiR-Tubulin-treated cells were subjected to two-color TIRFM and DHC-EGFP puncta were clearly observed streaming on Sir-Tubulin labeled MTs, which was especially evident on MTs that were pinned between the nucleus and the plasma membrane (**Video 3**). DHC puncta were next visualized with higher temporal resolution by acquiring single color TIRFM time-lapses of the EGFP channel at a rate of 5 frames per second (fps) (**Figure 1 D, Video 4**). The TIRFM time-lapses were assessed by eye and several parameters of the motile puncta were measured using kymographs (**Figure 1 E**). The average DHC puncta moved at 1.2 ± 0.05 (mean ± SEM) µm/sec over a distance of 2.8 ± 0.2 µm for 2.6 ± 0.2 seconds (**Figure 1 F-H**). While some motile puncta appeared to switch "tracks" at MT intersections, which was inferred from observations of sudden high angle turns, directional switches on the same MT were infrequent (~3% of runs).

Activation of dynein motility requires its interaction with the dynactin complex (Chowdhury et al., 2015; Grotjahn et al., 2018; Urnavicius et al., 2018; Zhang et al., 2017).

CRISPR/Cas9-based genomic engineering was used to insert the cassette encoding FKBP and

EGFP tags at the 3' end of the p50/dynamitin gene and a clonal p50-EGFP-expressing HeLa

cell line was subjected to live-cell fluorescence imaging to visualize dynactin in interphase cells

(**Figure 2 - figure supplement 1 A-C**). Consistent with prior observations of dynactin

localization (Vaughan et al., 1999; Vaughan et al., 2002) and like dynein, dynactin exhibited

robust tip-tracking activity on growing MT plus-ends that was clearly visualized via both spinning

disc confocal imaging and TIRFM (**Figure 2 A, B; Videos 5 and 6**). Suppression of plus-end

polymerization dynamics upon introduction of SiR-Tubulin caused a loss of the tip-tracking pool

of dynactin, which made a second pool of motile, MT-associated puncta of p50 more

pronounced especially when visualized with TIRFM (**Figure 2 C, Videos 7 and 8**). The p50-

EGFP-expressing cells were then subjected to single color TIRFM at 5 fps and kymograph

analysis to measure the motility parameters of the dynactin complex (**Figure 2 D**). The mean

velocity of the p50 puncta was 1.2 ± 0.07 μm/sec (**Figure 2 E**) while the mean run length was

2.6 ± 0.2 μm (**Figure 2 F**) and the mean run time was 2.2 ± 0.2 seconds (**Figure 2 G**). Similar to

dynein, dynactin was observed to switch MT "tracks" but infrequently switched direction. In

comparing the motility parameters of the motile dynein and dynactin puncta, their velocities

(**Figure 3 A**), run lengths (**Figure 3 B**), and run times (**Figure 3 C**) were statistically

indistinguishable. Thus, we inferred that the motile population of dynein was associated with the

dynactin complex.

The fluorescence intensities of motile DHC and p50 puncta were next compared to

motile EGFP-tagged kinesin-1 dimers (**Figure 3 - figure supplement 1 A-C)** as a standard to

assess the stoichiometries of the dynein-dynactin complexes. The intensities of motile DHC and

p50 puncta were compared to kinesin-1-EGFP expressed in either *Drosophila melanogaster* S2

cells (**Figure 3 – figure supplement 2 A-C**) or HeLa cells (**Figure 3 D-F**). Interestingly, the

intensity of motile DHC-EGFP puncta was not statistically significantly different from the

intensity of motile kinesin-1-EGFP dimers; however, the intensity of motile p50 puncta had a

mean fluorescence intensity that was about half that of the motile kinesin-1-EGFP puncta

(**Figure 3 G, Figure 3 - figure supplement 2 D**). Thus, the motile dynein and dynactin

complexes visualized here via high-speed TIRFM were comprised, on average, of two EGFP-

tagged DHC molecules and a single p50-EGFP molecule. When considering how these values

relate to physiological stoichiometries of the motile dynein-dynactin complexes, it is important to

note that both the DHC-EGFP and p50-EGFP HeLa cell line clones are heterozygotes (**Figure 3**

H, I). Dynein motility requires DHC dimerization and if there is an equal likelihood of EGFP-

tagged and untagged DHC incorporation into a functional dimer then the motile dynein puncta

we visualized would contain two DHC dimers. We were unable to assess the relative expression

levels of tagged versus untagged DHC by western blot due to the very large size of DHC (>500

kDa) relative to the EGFP tag. However, we can conclude from our data that there is at least

one DHC dimer in the motile puncta. Interestingly, western blotting of cell extract prepared from

the p50-EGFP clone revealed that the cells expressed a ~5-6-fold molar excess of the untagged

p50 compared to p50-EGFP (**Figure 3 I**). Differential allele regulation has been observed for

endogenously tagged proteins (Mann and Wadsworth, 2018; Roberts et al., 2017) suggesting

that regulation of gene expression may help avoid deleterious effects when cells can only

tolerate a fraction of the total protein pool being modified. The relative levels of EGFP-tagged

versus untagged p50 most likely explains why motile p50 puncta only had a single EGFP

molecule despite the fact that the dynactin complex is known to have four copies of p50 (Eckley

et al., 1999; Urnavicius et al., 2015). Thus, we conclude that the motile dynactin complexes

visualized here are typically comprised of one p50-EGFP molecule and 3 untagged copies of

p50.

While several groups have recently imaged endogenously tagged dynein in mitotic HeLa

cells and iNeurons (Fellows et al., 2024; Ide et al., 2023), to our knowledge this is the first direct

visualization of motile, endogenously tagged human dynein-dynactin complexes in proliferating,

non-neuronal cells. In human interphase cells, two distinct motile populations of dynein-dynactin

are observed: one that tip-tracks on polymerizing microtubule (MT) plus-ends, and another that

moves processively along MTs at high velocities - approximately seven times faster than the tip-

tracking population. The tip-tracking pool may be regulated in a cell cycle–dependent manner,

as dynein-dynactin tracking events appear less robust in mitotic cells compared to interphase

cells. We propose that the ~3 µm run lengths measured for the motile pool likely underestimate

the true processivity of dynein-dynactin. This is due to the relatively short and dynamic MT

tracks available in interphase HeLa cells, and the fact that motile puncta could only be

visualized while within the TIRF field and prior to photobleaching. Indeed, dynein was recently

shown to be highly processive in iNeurons, with a mean run length of ~35 µm and some runs

exceeding 100 µm (Fellows et al., 2024). In contrast, a separate study (Tirumala et al., 2024)

reported that dynein is not highly processive, typically exhibiting runs of very short duration

(~0.6 s) in HeLa cells. A notable technical difference that may account for this discrepancy is

that our study visualizes endogenously tagged human DHC, whereas Tirumala et al.

characterized over-expressed mouse DHC in HeLa cells. Over-expression of the DHC may

result in an imbalance of the subunits that comprise the active motor complex, leading to

inactive, or less active complexes. Similarly, mouse DHC may not have the ability to efficiently

assemble into active and processive dynein-dynactin-adaptor complexes to the same extent as

human DHC. Two-color imaging of SiR-Tubulin-labeled MTs and dynein-dynactin revealed a

population of p50-EGFP and DHC-EGFP puncta not evidently associated with MTs, suggesting

the presence of a cortical pool of dynein-dynactin. Cortical dynein-dynactin has been clearly

visualized in mitotic mammalian cells, where it functions in spindle positioning (Busson et al.,

1998; Collins et al., 2012; Faulkner et al., 2000; Kiyomitsu and Cheeseman, 2012; Kobayashi

and Murayama, 2009). In interphase mammalian cells, dynein-dynactin also contributes to

centrosome positioning (Etienne-Manneville and Hall, 2001; Palazzo et al., 2001), and has been

proposed to do so via pulling forces exerted on centrosomal MTs by a cortical pool (Burakov et

al., 2003; Dujardin and Vallee, 2002; Gundersen, 2002) that we likely visualized by TIRFM.

While a cortical pool of dynein-dynactin may exhibit lateral diffusion at the plasma membrane, it

would, by necessity, be considerably less motile than conventionally tip-tracking or running

dynein-dynactin complexes.

175   The dynein-dynactin pool that exhibited directional motility on MTs was capable of

switching MT tracks, which is a known behavior of dynein-dynactin (Ross et al., 2008), as

evidenced by sharp changes in direction sometimes at near perpendicular angles. Dynein-

dynactin rarely (~3% of motile puncta) switched directions suggesting that there is not a

constant tug-of-war between dynein-dynactin and kinesins in proliferating cells as compared to

observations of bidirectional vesicular transport in neurons (Hancock, 2014; Hendricks et al.,

2010; Maeder et al., 2014). The low directional switching frequency observed here is consistent

with in vitro studies reconstituting the tug-of-war phenomenon showing that one type of motor

dominates once directional movement begins and that reversal events are rare (Ali et al., 2023;

Belyy et al., 2016; Derr et al., 2012). The velocities measured here in HeLa cells were

comparable to those measured for unopposed dynein and; therefore, suggest that there was not

significant resistive drag from associated kinesin motors as has been observed in vitro. The

velocity and low switching frequency of motile puncta suggest that any kinesin motors

associated with cargos being transported by the dynein-dynactin visualized here are inactive

and/or cannot effectively bind the MT lattice during dynein-dynactin-mediated transport in

interphase HeLa cells. Interestingly, directional switching was also uncommon during retrograde

dynein-dynactin movements in iNeuron axons (Fellows et al., 2024) and in retrograde trafficking

of early endosomes in *Ustilago maydis* mediated by dynein-dynactin containing the HOOK3

adaptor (Bielska et al., 2014; Schuster et al., 2011).

194   Finally, our cell-based data are consistent with recent in vitro characterizations of

dynein-dynactin structure, function, and regulation (Belyy et al., 2016; Chaaban and Carter,

2022; Urnavicius et al., 2018). Based on our fluorescence intensity measurements, we favor the

interpretation that the motile dynein-dynactin complexes visualized here are typically comprised

of a single dynactin complex bound to a tandem array of two dyneins. Our measured dynein-

dynactin velocity of 1.2 µm/sec further supports this conclusion since this speed is consistent

with in vitro velocities of dynein-dynactin complexes containing activating adaptors that tend to

recruit two dyneins. However, we do not exclude that the range of DHC velocities and intensities

measured here may also include sub-populations of complexes containing a single dynein

dimer. It would be valuable to test whether dynein-dynactin regulation mechanisms that have

been characterized in vitro also apply to the physiological regulation of dynein-dynactin activity

in living cells by measuring DHC motility parameters and intensities in cells depleted of different

cargo adaptors. Ultimately, the direct visualization of human dynein-dynactin and quantification

of its motility parameters and stoichiometries in living cells should be an important physiological

complement to in vitro assays, which altogether will better inform mechanistic models of the

cellular processes that rely on dynein-dynactin function.

MATERIALS AND METHODS

Cell culture

Parental HeLa and CRISPR-edited HeLa clones expressing DHC-EGFP or p50-EGFP were grown in standard DMEM medium (Gibco, USA) supplemented with 10% non-heat-inactivated FBS (Gibco, USA) and 0.5× antibiotic/antimycotic cocktail (Sigma-Aldrich) and maintained at 37°C with 5% CO2. Human-kinesin-1-EGFP-expressing *Drosophila* S2 cells were grown in Schneider's medium (Life Technologies) supplemented with 10% heat inactivated fetal bovine serum (FBS) and 0.5x antibiotic/antimycotic cocktail (Sigma), and maintained at 25°C. Kinesin-1-EGFP expression in S2 cells was induced with 500μM $CuSO_4$ for 16-18 hours prior to imaging.

CRISPR-engineered cell line production

CRISPR gene editing.

FKBP-EGFP tags were added to the C-terminus of human dynein heavy chain (DHC) and p50 using methods described previously (Sheridan and Bentley, 2016; Stewart-Ornstein and Lahav, 2016). In brief, repair cassettes comprised of FKBP-EGFP linked to a cleavable peptide (T2A) followed by a selectable marker (Neomycin) were cloned into pCMV and used for PCR reactions. Glycine-Alanine linkers were included between proteins in the construct. Guide sequences were selected using the CRISPR design tool (http://crispr.mit.edu/) from the Zhang lab at MIT (Ran et al., 2013) using "other regions" and the human target genome (hg19). The search tool was used to select guides close to the C-terminus of the protein of interest (~100nt surrounding and including the stop codon). Top and bottom oligos were obtained for each guide with the bases 5'-CACC -3'added to the top oligo and 5'-AAAC-3' added to the complement of the bottom oligo.

Guides were cloned into a Cas9 containing plasmid (PX459) Addgene #62988, (Cambridge, MA) following methods previously outlined (Moyer and Holland, 2015). Top and

bottom oligos were annealed and then phosphorylated by T4 PNK (NEB, Ipswich, MA). Guides

were then cloned into PX459 that was cut using BbsI (NEB, Ipswich, MA) and ligated using T4

ligase (NEB Ipswich, MA). Guide-Cas9 containing plasmids were then sequenced using the U6

promotor primer (Ran et al., 2013) and purified using either endotoxin free mini-preps or midi-

preps according to manufacturer protocol (Promega, Madison, WI). Repair cassettes were

amplified using primers designed to be homologous to the C-terminal genomic DNA surrounding

the STOP codon. In all cases, the guide target sequence was mutated to prevent Cas9 from

recognizing the repair cassette.

Cells were grown in DMEM medium (ThermoFisher Scientific) with 10% Fetal bovine

serum (FBS, Atlanta Biologicals, Flowery Branch, GA) and 0.5X antibiotic/antimycotic solution

(final concentrations 50 U/mL penicillin, 0.05 mg/mL streptomycin, 0.125 ug/mL amphotericin B;

Sigma-Aldrich, St. Louis, MO) at 37 $^\circ$C and 5% carbon dioxide (CO_2). For long term storage

cells were frozen in DMEM medium with 5% FBS and 0.5X antibiotic/antimycotic solution and

15% DMSO and held at -80°C for 1-2 days before moving to liquid nitrogen.

To generate CRISPR modified cell lines, parental cells were nucleofected using an

Amaxa Nucleofector (Lonza, Portsmouth, NH) program I-013 and Mirus nucleofection reagent

(Mirus Bio LLC, Madison, WI) according to the manufacturer's recommendations. Plasmids and

Repair cassettes were used at ratio of 1:1 at a concentration of 1 µg DNA each. Following

nucleofection, cells were grown in regular growth media in 100mm dishes for 48-72 Hrs. and

then 0.2g/L Neomycin/G418 (InvivoGen, San Diego, CA) selection media was added. Media

was then changed daily for 10-14 days and colonies of green cells were picked using cloning

rings and returned to regular media for further screening and experiments.

Genotyping

Genomic DNA was isolated from clonal CRISPR tagged cells using Genomic DNA Mini Kit

(Invitrogen) according to manufacturer's recommendations. DNA was then amplified by PCR

using genomic primers targeting the Neomycin cassette- and downstream of the Stop codon.

Phusion polymerase (NEB) was used to amplify 1 µL of isolated genomic DNA in a 50 µL

reaction for 30 cycles. The resulting PCR products were analyzed using 0.8% agarose gel

electrophoresis. The larger band was excised, and gel extracted using Qiagen gel extraction kit

and sequenced to verify proper integration of the tag.

Live-cell microscopy

Cells were seeded onto 35 mm glass bottom petri dishes (Cellvis) 2-3 days prior to imaging.

With the exception of the visualization of tip-tracking DHC-EGFP, cells were incubated in media

supplemented with 1 µM SiR-Tubulin (Cytoskeleton, Inc) for 30-60 minutes prior to washing out

the SiR-Tubulin with fresh DMEM before imaging. The cells were visualized on a Nikon Ti-E

microscope equipped with a TIRF illuminator, Borealis (Andor) retrofitted CSU-10 (Yokogawa)

spinning disc head, a 100X 1.49-NA differential interference contrast TIRF Apochromat oil-

immersion objective, Nikon perfect focus system (PFS), two ORCA-Flash4.0 LT Digital CMOS

cameras (Hamamatsu), four laser lines (447, 488, 561, and 641 nm) in a dual-output laser

launch system (Andor), and MetaMorph software (Molecular Devices). The imaging mode

(spinning disc versus TIRFM) of the system is set by selecting the appropriate fiber optic output

and corresponding CMOS camera. For TIRFM imaging, interphase cells were identified and the

initial focal plane was set by focusing on the far-red SiR tubulin signal with a particular emphasis

on finding cells in which individual microtubules could be seen between the nucleus and the

glass-adhered plasma membrane. Individual snap shots were then taken on the EGFP channel

to refine the focal plane on the DHC or p50 puncta. Once individual puncta were well-resolved

the PFS was engaged and the cell was subjected to streaming TIRFM for 12 seconds at an

acquisition rate of 5 frames per second (200 ms exposure) and 2x2 camera binning. The

population of dynein-dynactin on MTs positioned between the plasma membrane and nucleus

would fall within the 100-200 nm excitable range of the evanescent wave produced by TIRF

since the plasma membrane is typically ~5 nm and a MT is ~25 nm. While TIRFM is capable of

effectively visualizing dynein-dynactin motility in living cells, in some time-lapses the excitation

laser may have been angled to propagate as a highly inclined and laminated optical sheet

(HILO) (Tokunaga et al., 2008). The experimental design and analyses would be unaffected by

whether the motile punta were visualized via TIRFM or HILO microscopy. For consistency, the

imaging mode employed in this study is referred to as TIRFM.

Quantifications of motility parameters and fluorescence intensities

High temporal (5 fps) TIRFM time-lapses were examined by eye to identify evident motile

puncta, the paths of which were traced manually in MetaMorph using the Multi-Line drawing

function. A kymograph of the line segment with a 5-pixel line width was then generated and the

velocity was determined by measuring the slope of the motility event on the kymograph from

start to end using the Single Line drawing function. The run time was measured by defining the

start and end frame of evident movement in the TIRFM time-lapse and the run length was then

calculated by multiplying the run time by the velocity measured in the kymograph of the motility

event. The run times and run lengths are likely underestimates of what dynein-dynactin can

achieve due to the fact that the puncta sometimes bleached or exited the TIRFM evanescent

field.

The number of EGFP molecules per motile puncta of DHC and p50 was determined by

comparing their background-corrected fluorescence intensities to that of a known dimer

standard: human kinesin-1 tagged with EGFP in both *Drosophila* S2 cells and in HeLa cells. As

previously described (Ye et al., 2018), *Drosophila* S2 cells expressing inducible human kinesin-

1 (in this case tagged with EGFP) were induced with 500µM CuSO4 for 16 hours to induce

expression. The next day, the induced S2 cells were seeded onto a concanavalin A coated

glass bottom petri dish and allowed to adhere for ~1 hour prior to visualization by TIRFM. The

DHC-EGFP and p50-EGFP HeLa clones and the human kinesin-1-EGFP-expressing S2 cells

were each visualized sequentially via live-cell TIRF microscopy using identical imaging

parameters (5 fps acquisition rate, 200 ms exposure, 2X2 binning) within the same region of the

camera chip.

To compare DHC-EGFP and p50-EGFP intensities to human kinesin-1-EGFP in HeLa

cells, HeLa cells were seeded in a 35 mm dish in 2 mL of complete HeLa medium (DMEM

supplemented with 10% FBS and 0.5X antibiotic-antimycotic mix) 18–24 hours prior to

transfection at an initial density (0.25×10^6) to achieve ~50% confluency on the day of

transfection. On the day of transfection, two sterile 1.5 mL microcentrifuge tubes were prepared

and labeled A and B. Each tube received 125 µL of OPTI-MEM medium. Tube A was

supplemented with 7.5 µL of Lipofectamine 3000 reagent and vortexed briefly. Tube B was

supplemented with 250 ng of endotoxin-free plasmid pB80-hsKIF5B(1-560)-L-GFP (human

kinesin-1-EGFP) (Addgene plasmid # 193716) and 4 µL of P3000 reagent, followed by brief

vortexing. The contents of tube B were then transferred to tube A, vortexed gently, and

incubated at room temperature for 15 minutes to allow complex formation. The resulting DNA-

lipid complexes were added dropwise to the HeLa cells, which were then incubated at 37°C in a

humidified CO_2 incubator for 24 hours. After incubation, the medium was replaced with 2 mL of

fresh HeLa medium, and cells were subjected to imaging over the next 1-2 days (2-3 days post

transfection). The kinesin-1-EGFP transfected cells, and DHC-EGFP and p50-EGFP HeLa

clones were each visualized sequentially on the same day via live-cell TIRF microscopy using

identical imaging parameters within the same region of the camera chip. The region-in-region

method (Ye and Maresca, 2018) was used to background correct and quantify the integrated

fluorescence intensities of motile puncta of DHC-, p50-, and kinesin-1-EGFP (for both the S2

cell and HeLa cell comparisons). The intensity of each motile puncta was measured for a single

time-point from each run in which the spot was most clearly resolved and did not have

significant local background signal from nearby EGFP puncta. Cytoplasmic levels of DHC-EGFP

and p50-EGFP were quantified from max projections of spinning disk confocal Z-stacks by

subtracting the integrated intensity of a 25x25 pixel square ROI in the nucleus from the

integrated intensity of a 25x25 pixel ROI placed in a representative region of the cytoplasm.

Western blotting

Twenty µg of total protein from cell lysates prepared from parental HeLa cells or the p50-EGFP

clone were loaded onto a 10% SDS-PAGE gel, run out, and transferred to a nitrocellulose

membrane on the Trans-Blot® Turbo™ transfer system (Bio-Rad Laboratories) using the

preprogrammed "HIGH MW" 10-minute protocol. After blocking for 1 hour in TBS with 0.1%

Tween and 5% milk, the membrane was incubated overnight at 4°C with mouse anti-dynactin

p50 (BD Transduction Laboratories, Cat. # 611002) diluted at 1:1000 in the block. The

membrane was washed 3X 5 minutes in TBS + 0.1% Tween and then incubated for 1 hour in

milk containing donkey-anti-mouse IgG secondary antibodies conjugated with HRP (Jackson

ImmunoResearch Laboratories, Inc.) diluted at 1:5000 in block. Following 3X 5 minutes washes

in TBS + 0.1% Tween the blot was incubated with Immobilon Western Chemiluminescent HRP

Substrate (Millipore) according to the manufacturer's recommendations. Imaging of the

membrane was done on a G:Box system controlled by GeneSnap software (Syngene).

Statistical analyses

359 Reported p-values were generated with a randomization method using the PlotsofDifferences

360 web tool at: https://huygens.science.uva.nl/PlotsOfDifferences (Goedhart, 2019).

361 PlotsofDifferences does not rely on assumptions about the distribution of the data (normal

362 versus non-normal) to calculate p-values.

ACKNOWLEDGMENTS

We are grateful to Barbara Mann for generating the CRISPR-engineered HeLa cell lines. Thank

you to Thomas Laskarzewski for help with R and data visualization. pB80-hsKIF5B(1-560)-L-

GFP was a gift from Lukas Kapitein (Addgene plasmid # 193716 ;

http://n2t.net/addgene:193716 ; RRID:Addgene_193716). This work was supported by NIH

grants (GM107026 and GM156188) to T.J.M. and by an NSF grant (MCB1817926) to P.W.

CONFLICT OF INTEREST

The authors declare no competing financial interests.

REFERENCES

Ali, F.A., A.J. Zwetsloot, C.E. Stone, T.E. Morgan, R.F. Wademan, A.P. Carter, and A. Straube. 2023. KIF1C activates and extends dynein movement through the FHF cargo adaptor. *bioRxiv*:2023.2010.2026.564242.

Belyy, V., M.A. Schlager, H. Foster, A.E. Reimer, A.P. Carter, and A. Yildiz. 2016. The mammalian dynein-dynactin complex is a strong opponent to kinesin in a tug-of-war competition. *Nat Cell Biol*. 18:1018-1024.

Bielska, E., M. Schuster, Y. Roger, A. Berepiki, D.M. Soanes, N.J. Talbot, and G. Steinberg. 2014. Hook is an adapter that coordinates kinesin-3 and dynein cargo attachment on early endosomes. *J Cell Biol*. 204:989-1007.

Burakov, A., E. Nadezhdina, B. Slepchenko, and V. Rodionov. 2003. Centrosome positioning in interphase cells. *J Cell Biol*. 162:963-969.

Busson, S., D. Dujardin, A. Moreau, J. Dompierre, and J.R. De Mey. 1998. Dynein and dynactin are localized to astral microtubules and at cortical sites in mitotic epithelial cells. *Curr Biol*. 8:541-544.

Canty, J.T., R. Tan, E. Kusakci, J. Fernandes, and A. Yildiz. 2021. Structure and Mechanics of Dynein Motors. *Annu Rev Biophys*. 50:549-574.

Carter, A.P., A.G. Diamant, and L. Urnavicius. 2016. How dynein and dynactin transport cargos: a structural perspective. *Curr Opin Struct Biol*. 37:62-70.

Cason, S.E., and E.L.F. Holzbaur. 2022. Selective motor activation in organelle transport along axons. *Nat Rev Mol Cell Biol*. 23:699-714.

Cassimeris, L., N.K. Pryer, and E.D. Salmon. 1988. Real-time observations of microtubule dynamic instability in living cells. *J Cell Biol*. 107:2223-2231.

Chaaban, S., and A.P. Carter. 2022. Structure of dynein-dynactin on microtubules shows tandem adaptor binding. *Nature*. 610:212-216.

Chowdhury, S., S.A. Ketcham, T.A. Schroer, and G.C. Lander. 2015. Structural organization of the dynein-dynactin complex bound to microtubules. *Nat Struct Mol Biol*. 22:345-347.

Collins, E.S., S.K. Balchand, J.L. Faraci, P. Wadsworth, and W.L. Lee. 2012. Cell cycle-regulated cortical dynein/dynactin promotes symmetric cell division by differential pole motion in anaphase. *Mol Biol Cell*. 23:3380-3390.

Derr, N.D., B.S. Goodman, R. Jungmann, A.E. Leschziner, W.M. Shih, and S.L. Reck-Peterson. 2012. Tug-of-war in motor protein ensembles revealed with a programmable DNA origami scaffold. *Science*. 338:662-665.

Dujardin, D.L., and R.B. Vallee. 2002. Dynein at the cortex. *Curr Opin Cell Biol*. 14:44-49.

Eckley, D.M., S.R. Gill, K.A. Melkonian, J.B. Bingham, H.V. Goodson, J.E. Heuser, and T.A. Schroer. 1999. Analysis of dynactin subcomplexes reveals a novel actin-related protein associated with the arp1 minifilament pointed end. *J Cell Biol*. 147:307-320.

Etienne-Manneville, S., and A. Hall. 2001. Integrin-mediated activation of Cdc42 controls cell polarity in migrating astrocytes through PKCzeta. *Cell*. 106:489-498.

Faulkner, N.E., D.L. Dujardin, C.Y. Tai, K.T. Vaughan, C.B. O'Connell, Y. Wang, and R.B. Vallee. 2000. A role for the lissencephaly gene LIS1 in mitosis and cytoplasmic dynein function. *Nat Cell Biol*. 2:784-791.

Fellows, A.D., M. Bruntraeger, T. Burgold, A.R. Bassett, and A.P. Carter. 2024. Dynein and dynactin move long-range but are delivered separately to the axon tip. *J Cell Biol*. 223.

Gill, S.R., T.A. Schroer, I. Szilak, E.R. Steuer, M.P. Sheetz, and D.W. Cleveland. 1991. Dynactin, a conserved, ubiquitously expressed component of an activator of vesicle motility mediated by cytoplasmic dynein. *J Cell Biol*. 115:1639-1650.

Goedhart, J. 2019. PlotsOfDifferences – a web app for the quantitative comparison of unpaired data. *bioRxiv*:578575.

Grotjahn, D.A., S. Chowdhury, Y. Xu, R.J. McKenney, T.A. Schroer, and G.C. Lander. 2018. Cryo-electron tomography reveals that dynactin recruits a team of dyneins for processive motility. *Nat Struct Mol Biol*. 25:203-207.

Gundersen, G.G. 2002. Evolutionary conservation of microtubule-capture mechanisms. *Nat Rev Mol Cell Biol*. 3:296-304.

Hancock, W.O. 2014. Bidirectional cargo transport: moving beyond tug of war. *Nat Rev Mol Cell Biol*. 15:615-628.

Hendricks, A.G., E. Perlson, J.L. Ross, H.W. Schroeder, 3rd, M. Tokito, and E.L. Holzbaur. 2010. Motor coordination via a tug-of-war mechanism drives bidirectional vesicle transport. *Curr Biol*. 20:697-702.

Hirokawa, N., Y. Noda, Y. Tanaka, and S. Niwa. 2009. Kinesin superfamily motor proteins and intracellular transport. *Nat Rev Mol Cell Biol*. 10:682-696.

Ide, A.H., K.F. DeLuca, O. Wiggan, S.M. Markus, and J.G. DeLuca. 2023. The role of kinetochore dynein in checkpoint silencing is restricted to disassembly of the corona. *Mol Biol Cell*. 34:ar76.

Kiyomitsu, T., and I.M. Cheeseman. 2012. Chromosome- and spindle-pole-derived signals generate an intrinsic code for spindle position and orientation. *Nat Cell Biol*. 14:311-317.

Kobayashi, T., and T. Murayama. 2009. Cell cycle-dependent microtubule-based dynamic transport of cytoplasmic dynein in mammalian cells. *PLoS One*. 4:e7827.

Maeder, C.I., K. Shen, and C.C. Hoogenraad. 2014. Axon and dendritic trafficking. *Curr Opin Neurobiol*. 27:165-170.

Mann, B.J., and P. Wadsworth. 2018. Distribution of Eg5 and TPX2 in mitosis: Insight from CRISPR tagged cells. *Cytoskeleton (Hoboken)*. 75:508-521.

McKenney, R.J., W. Huynh, M.E. Tanenbaum, G. Bhabha, and R.D. Vale. 2014. Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. *Science*. 345:337-341.

Moyer, T.C., and A.J. Holland. 2015. Generation of a conditional analog-sensitive kinase in human cells using CRISPR/Cas9-mediated genome engineering. *Methods Cell Biol*. 129:19-36.

Palazzo, A.F., H.L. Joseph, Y.J. Chen, D.L. Dujardin, A.S. Alberts, K.K. Pfister, R.B. Vallee, and G.G. Gundersen. 2001. Cdc42, dynein, and dynactin regulate MTOC reorientation independent of Rho-regulated microtubule stabilization. *Curr Biol*. 11:1536-1541.

Pfister, K.K., E.M. Fisher, I.R. Gibbons, T.S. Hays, E.L. Holzbaur, J.R. McIntosh, M.E. Porter, T.A. Schroer, K.T. Vaughan, G.B. Witman, S.M. King, and R.B. Vallee. 2005. Cytoplasmic dynein nomenclature. *J Cell Biol*. 171:411-413.

Ran, F.A., P.D. Hsu, J. Wright, V. Agarwala, D.A. Scott, and F. Zhang. 2013. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc*. 8:2281-2308.

Reck-Peterson, S.L., W.B. Redwine, R.D. Vale, and A.P. Carter. 2018. The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol*. 19:382-398.

Roberts, B., A. Haupt, A. Tucker, T. Grancharova, J. Arakaki, M.A. Fuqua, A. Nelson, C. Hookway, S.A. Ludmann, I.A. Mueller, R. Yang, R. Horwitz, S.M. Rafelski, and R.N. Gunawardane. 2017. Systematic gene tagging using CRISPR/Cas9 in human stem cells to illuminate cell organization. *Mol Biol Cell*. 28:2854-2874.

Ross, J.L., H. Shuman, E.L. Holzbaur, and Y.E. Goldman. 2008. Kinesin and dynein-dynactin at intersecting microtubules: motor density affects dynein function. *Biophys J*. 94:3115-3125.

Rusan, N.M., C.J. Fagerstrom, A.M. Yvon, and P. Wadsworth. 2001. Cell cycle-dependent changes in microtubule dynamics in living cells expressing green fluorescent protein-alpha tubulin. *Mol Biol Cell*. 12:971-980.

Schlager, M.A., H.T. Hoang, L. Urnavicius, S.L. Bullock, and A.P. Carter. 2014. In vitro
reconstitution of a highly processive recombinant human dynein complex. *Embo j*.
33:1855-1868.
Schroer, T.A., and M.P. Sheetz. 1991. Two activators of microtubule-based vesicle transport. *J*
Cell Biol. 115:1309-1318.
Schuster, M., R. Lipowsky, M.A. Assmann, P. Lenz, and G. Steinberg. 2011. Transient binding
of dynein controls bidirectional long-range motility of early endosomes. *Proc Natl Acad*
Sci U S A. 108:3618-3623.
Sheridan, R.M., and D.L. Bentley. 2016. Selectable one-step PCR-mediated integration of a
degron for rapid depletion of endogenous human proteins. *Biotechniques*. 60:69-74.
Splinter, D., D.S. Razafsky, M.A. Schlager, A. Serra-Marques, I. Grigoriev, J. Demmers, N.
Keijzer, K. Jiang, I. Poser, A.A. Hyman, C.C. Hoogenraad, S.J. King, and A.
Akhmanova. 2012. BICD2, dynactin, and LIS1 cooperate in regulating dynein
recruitment to cellular structures. *Mol Biol Cell*. 23:4226-4241.
Stewart-Ornstein, J., and G. Lahav. 2016. Dynamics of CDKN1A in Single Cells Defined by an
Endogenous Fluorescent Tagging Toolkit. *Cell Rep*. 14:1800-1811.
Tirumala, N.A., G.M.I. Redpath, S.V. Skerhut, P. Dolai, N. Kapoor-Kaushik, N. Ariotti, K. Vijay
Kumar, and V. Ananthanarayanan. 2024. Single-molecule imaging of stochastic
interactions that drive dynein activation and cargo movement in cells. *J Cell Biol*. 223.
Tokunaga, M., N. Imamoto, and K. Sakata-Sogawa. 2008. Highly inclined thin illumination
enables clear single-molecule imaging in cells. *Nat Methods*. 5:159-161.
Urnavicius, L., C.K. Lau, M.M. Elshenawy, E. Morales-Rios, C. Motz, A. Yildiz, and A.P. Carter.
2018. Cryo-EM shows how dynactin recruits two dyneins for faster movement. *Nature*.
554:202-206.
Urnavicius, L., K. Zhang, A.G. Diamant, C. Motz, M.A. Schlager, M. Yu, N.A. Patel, C.V.
Robinson, and A.P. Carter. 2015. The structure of the dynactin complex and its
interaction with dynein. *Science*. 347:1441-1446.
Vale, R.D. 1987. Intracellular Transport Using Microtubule-Based Motors. *Annual Review of Cell*
Biology. 3:347-378.
Vaughan, K.T., S.H. Tynan, N.E. Faulkner, C.J. Echeverri, and R.B. Vallee. 1999. Colocalization
of cytoplasmic dynein with dynactin and CLIP-170 at microtubule distal ends. *J Cell Sci*.
112 (Pt 10):1437-1447.
Vaughan, P.S., P. Miura, M. Henderson, B. Byrne, and K.T. Vaughan. 2002. A role for regulated
binding of p150(Glued) to microtubule plus ends in organelle transport. *J Cell Biol*.
158:305-319.
Walker, R.A., E.T. O'Brien, N.K. Pryer, M.F. Soboeiro, W.A. Voter, H.P. Erickson, and E.D.
Salmon. 1988. Dynamic instability of individual microtubules analyzed by video light
microscopy: rate constants and transition frequencies. *J Cell Biol*. 107:1437-1448.
Ye, A.A., and T.J. Maresca. 2018. Measuring mitotic forces. *Methods Cell Biol*. 144:165-184.
Ye, A.A., V. Verma, and T.J. Maresca. 2018. NOD is a plus end-directed motor that binds EB1
via a new microtubule tip localization sequence. *J Cell Biol*. 217:3007-3017.
Zhang, K., H.E. Foster, A. Rondelet, S.E. Lacey, N. Bahi-Buisson, A.W. Bird, and A.P. Carter.
2017. Cryo-EM Reveals How Human Cytoplasmic Dynein Is Auto-inhibited and
Activated. *Cell*. 169:1303-1314.e1318.

 FIGURE LEGENDS

Figure 1. Dynein tip-tracks on polymerizing MTs and walks processively at high

velocities in interphase HeLa cells. (A) Representative spinning disc confocal time-lapse of a

DHC-EGFP-expressing HeLa cell showing robust tip-tracking. Zoomed views of the numbered

boxed regions are shown to the right with kymographs of the tip-tracking events highlighted with

yellow arrows in the zoomed panels. **(B)** Representative TIRFM time-lapse of a DHC-EGFP-

expressing HeLa cell showing the tip-tracking population. Zoomed views of the numbered boxed

regions are shown with kymographs of the tip-tracking events highlighted with yellow arrows.

(C) Still frame from a representative TIRFM time-lapse of a DHC-EGFP-expressing HeLa cell

treated with SiR-Tubulin. In the merge image DHC is green and Sir-Tubulin labeled MTs are

magenta. **(D)** Still frame from a representative high temporal resolution (5 fps) TIRFM time-

lapse of a DHC-EGFP-expressing HeLa cell treated with SiR-Tubulin. Boxed region is shown as

a zoomed inset in the lower panel with the track of a motile DHC puncta highlighted in green.

(E) Representative kymographs of motile puncta spanning the range of measured velocities

(Vel) and run lengths (RL). **(F)** Distribution of DHC velocities (n = 100 puncta). **(G)** Distribution of

DHC run lengths (n = 81 puncta). **(H)** Distribution of DHC run times (n = 81 puncta). Scale bars,

10 μm (A-D); 1 μm (all insets), 10 μm (horizontal); 1 min (vertical) in the kymographs in A and B;

and 1 μm (horizontal); 1 second (vertical) in E. Displayed times are min:sec.

Figure 2. The dynactin complex component p50 tip-tracks on polymerizing MTs and

walks processively at high velocities in interphase HeLa cells. (A) Still frames from a

representative spinning disc confocal time-lapse of a p50-EGFP-expressing HeLa cell. Zoomed

view of the boxed region is shown with a kymograph of the tip-tracking event highlighted with

the yellow arrow. **(B)** Still frames from a representative TIRFM time-lapse of a p50-EGFP-

expressing HeLa cell showing the tip-tracking population. Zoomed view of the boxed region is

shown with a kymograph of the tip-tracking event highlighted by the yellow arrow. **(C)** Still frame

from a representative TIRFM time-lapse of a p50-EGFP-expressing HeLa cell treated with SiR-

Tubulin. In the merge image p50 is green and Sir-Tubulin labeled MTs are magenta. **(D)**

Representative kymographs of motile p50 puncta spanning the range of measured velocities

(Vel) and run lengths (RL). **(E)** Distribution of p50 velocities (n = 44 puncta). **(F)** Distribution of

p50 run lengths (n = 41 puncta). **(G)** Distribution of p50 run times (n = 41 puncta). Scale bars,

10 μm (A-C); 1 μm (all insets), 5 μm (horizontal); 1 min (vertical) in the kymographs in A and B;

and 1 μm (horizontal); 1 second (vertical) in D. Displayed times are min:sec.

Figure 3. Motile DHC and p50 puncta exhibit identical motility parameters but different

fluorescent intensities. (A-C) Scatter plots of (A) velocities (DHC, n = 100; p50, n = 44), (B)

run lengths (DHC, n = 81; p50, n 41), and (C) run times (DHC, n = 81; p50, n 41). **(D-F)**

Distributions of the background corrected fluorescence intensities of motile puncta of (D)

kinesin-1-EGFP transiently expressed in HeLa cells, (E) DHC-EGFP, and (F) p50-EGFP. The

dashed line in each histogram denotes the mean value of the kinesin-1-EGFP data set. **(G)**

Scatter plots of the kinesin-1, DHC, and p50 fluorescence intensities (kinesin-1, n = 90 puncta;

DHC, n = 84 puncta; p50, n = 74 puncta). **(H)** PCR of genomic DNA from the DHC-EGP clone

used in this study using PCR primers flanking the integration site of the repair cassette. The

upper band was extracted and subjected to sequencing the results of which are shown in Figure

1 – supplemental figure 1 A. **(I)** Western blot for p50 of cell lysates from the parental HeLa cell

line and the p50-EGFP clone used in this study. The tagged p50 runs ~30 kDa larger than the

untagged p50 and is expressed at ~5-6-fold lower levels than the endogenous p50. Error bars

are mean values +/- standard deviations. The reported p-values were determined by a

randomization method: n.s. is not significant ($p > 0.05$).

 SUPPLEMENTAL MATERIALS

Supplemental Figures

Figure 1 – figure supplement 1. (A) Repair cassette for generating the DHC-EGFP CRISPR-

engineered HeLa cell line with sequencing trace showing its integration at the dynein heavy

chain (DYNC1H1) gene locus. **(B)** Montage of representative fields of view of the CRISPR-

engineered DHC-EGFP cell line used in this study. Expression levels are consistent across the

population and do not exhibit significant cell-to-cell variation. Scale bar, 20 µm.

Figure 2 – figure supplement 1. (A) Montage of representative fields of view of the CRISPR-

engineered p50-EGFP clone in comparison to the montage of DHC-EGFP fields from Figure 1 -

figure supplement 1 B. For the purpose of comparison, the two cells lines were visualized with

identical imaging parameters on the same day and displayed identically in the figure. **(B)** Scatter

plots of background corrected fluorescence intensities of cytoplasmic pools of DHC-EGFP (n =

31 cells) and p50-EGFP (n = 13 cells) from (A). **(C)** Montage of fields of view p50-EGFP clonal

cells visualized with longer exposure times (500 ms) than in (A). Error bars are mean values +/-

standard deviations. Scale bars, 20 µm.

Figure 3 – figure supplement 1. (A) Representative kymographs of motile kinesin-1-EGFP

puncta from transiently transfected HeLa cells. **(B)** Distribution of kinesin-1 velocities (mean +/-

SEM; n = 76 puncta). **(C)** Distribution of kinesin-1 run lengths (mean +/- SEM; n = 60 puncta).

Kinesin-1 expressed in HeLa cells exhibited a slower mean velocity and shorter mean run

length than the HeLa dynein-dynactin. Scale bars, 1 µm (horizontal); 1 second (vertical).

Figure 3 – figure supplement 2. (A-C) Distributions of background corrected fluorescence of

(A) kinesin-1-EGFP expressed in *Drosophila melanogaster* S2 cells, (B) DHC-EGFP, and (C)

p50-EGFP. The dashed line in each histogram denotes the mean value of the kinesin-1-EGFP

data set. **(D)** Box and whisker plots of the kinesin-1, DHC, and p50 fluorescence intensities

(kinesin-1, n = 100 puncta; DHC, n = 71 puncta; p50, n = 38 puncta). The reported p-values

were determined by a randomization method: n.s. is not significant (p > 0.05).

Supplemental Videos

Video 1. Spinning disk confocal time-lapse of a DHC-EGFP CRISPR-engineered

interphase HeLa cell. Zoomed views of the boxed regions in the upper left panel are shown in

the right panels. Displayed times are min:sec. Scale bars, 10 µm (upper left panel) and 1 µm

(right panels). Relates to Fig. 1A.

Video 2. TIRFM time-lapse of a DHC-EGFP CRISPR-engineered interphase HeLa cell.

Displayed times are min:sec. Scale bar, 10 µm. Relates to Fig. 1B.

Video 3. TIRFM time-lapse of a DHC-EGFP (green) CRISPR-engineered interphase HeLa

cell treated with SiR-Tubulin (magenta). Displayed times are min:sec. Scale bar, 10 µm.

Relates to Fig. 1C.

Video 4. High temporal-resolution (5 frames per second) TIRFM time-lapse of a DHC-

EGFP CRISPR-engineered interphase HeLa cell treated with SiR-Tubulin. The boxed

regions highlight motile DHC puncta. Displayed times are sec:millisec. Scale bar, 1 µm. Relates

to Fig. 1D.

Video 5.   Spinning disk confocal time-lapse of a p50-EGFP CRISPR-engineered

interphase HeLa cell. Displayed times are min:sec. Scale bar, 10 µm. Relates to Fig. 2A.

Video 6. TIRFM time-lapse of a p50-EGFP CRISPR-engineered interphase HeLa cell.

Displayed times are min:sec. Scale bar, 10 µm. Relates to Fig. 2B.

Video 7. TIRFM time-lapse of a p50-EGFP (green) CRISPR-engineered interphase HeLa

cell treated with SiR-Tubulin (magenta). Displayed times are min:sec. Scale bar, 10 µm.

Relates to Fig. 2C.

Video 8. TIRFM time-lapse of a p50-EGFP (green) CRISPR-engineered interphase HeLa

cell treated with SiR-Tubulin (magenta). Displayed times are min:sec. Scale bar, 10 µm.

Figure 1

A DHC-EGFP Spinning disc
1
2
0:00 0:36 1:06 1:15 1 min 10 µm
0:00 0:24 0:36 0:42 0:51

B DHC-EGFP TIRFM
1
2
0:00 0:22 0:40 1:30 1 min 10 µm
0:00 0:08 0:22 0:30 0:42

C Docetaxel
DHC-EGFP SiR-Tubulin Merge

D DHC-EGFP + SiR Tub (TIRFM)

E Kymographs
Vel: Slower Average Faster Average Slower
RL: Shorter Shorter Shorter Average Longer

1 s
1 µm

Vel: Average Faster
RL: Longer Longer

F DHC Velocity
1.2 ± 0.05 µm/sec
n=100
Percentage
Microns/sec

G DHC Run Length
2.8 ± 0.2 µm
n=81
Percentage
Microns

H DHC Run Time
2.6 ± 0.2 seconds
n=81
Percentage
Seconds

Figure 2

Figure 3