Fibroblast-derived IL-11 is a driver and therapeutic target in Peutz-Jeghers syndrome polyposis

Eva Domènech-Moreno [1,2*], Wei-Wen Lim [3,4*] , Anders Brandt [1,2], Toni T Lemmetyinen[6], Emma Viitala[6], Tomi P Mäkelä[1,2], Stuart A Cook[3,4,5*], Saara Ollila[6*]

1. HiLIFE-Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland.
2. iCAN Digital Precision Cancer Medicine Flagship, University of Helsinki, Helsinki, Finland.
3. National Heart Research Institute Singapore, National Heart Centre Singapore, 169609, Singapore
4. Cardiovascular and Metabolic Disorders Program, Duke-National University of Singapore Medical School
5. MRC-London Institute of Medical Sciences, Hammersmith Hospital Campus, London, UK
6. Translational Cancer Medicine Program, University of Helsinki, Helsinki, Finland.
* Equal contribution

Abstract

Inactivating germline mutations in the tumor suppressor kinase *LKB1* (*STK11*) predispose to Peutz-Jeghers Syndrome (PJS) with increased cancer risk and early-onset development of gastrointestinal polyps requiring regular surveillance. Studies using PJS mouse models have indicated fibroblasts as drivers of polyp formation. Here, we use single-cell RNA sequencing to investigate the fibroblast heterogeneity and tumorigenic mechanisms in a PJS mouse model. We identify five distinct gastric fibroblast subsets, including a polyp-enriched ST2 (*Il1rl1*)-expressing crypt top fibroblast (CTF) cluster, and show that the polyps arise from *Foxl1*-expressing CTFs. The transcriptional signature of the ST2+ fibroblasts overlapped with the effects seen by LKB1 loss *in vitro* and *in vivo*. The ST2+ fibroblasts also shared similarities with inflammation-associated fibroblasts, and inflammation exacerbated polyposis. Interestingly, the tumorigenic ST2+ fibroblasts showed a distinct signaling pattern, including high expression of interleukin 11 (IL-11). Notably, IL-11 was required for the LKB1 loss-induced transcriptional changes in fibroblasts, and treatment with an IL-11 neutralizing antibody substantially reduced the tumor burden in the PJS mouse model, suggesting therapeutic value. In summary, our study uncovers a critical mechanism underlying PJS polyposis and suggests that PJS patients could benefit from anti-IL-11 therapy.

Introduction

Peutz-Jeghers syndrome (PJS) is a hereditary gastrointestinal (GI) tumorigenesis syndrome with an estimated prevalence of 1 in 50,000–200,000 live births (Giardiello & Trimbath, 2006). PJS is characterized by the emergence of multiple benign hamartomatous tumors throughout the GI tract and an increased risk of cancer (Jelsig et al., 2023). Regular removal of polyps using endoscopy or surgery is often required to avoid intussusception, resulting in some patients experiencing significantly reduced quality of life. PJS is inherited through inactivating mutations in the tumor suppressor gene *LKB1* (*STK11*) (Hemminki et al., 1998), and *Lkb1* heterozygous mice recapitulate the polyp formation with development of mainly gastric hamartomatous polyps with full penetrance, allowing mechanistic studies (Rossi et al., 2002). Previously, we and others have demonstrated that LKB1 deletion limited to the mesenchymal compartment, or specifically to fibroblasts, was sufficient for clonal expansion and polyp formation, suggesting that stromal fibroblasts drive the development of PJS tumors (Katajisto et al. 2008; Ollila et al. 2018; Cotton et al. 2022). Also T-cells have been implicated in PJS polyposis (Poffenberger et al., 2018).

Recent work using single-cell RNA sequencing (scRNAseq) of small intestinal and colonic tissues has revealed the existence of spatially and functionally distinct fibroblast populations (Brügger et al., 2020; Brügger & Basler, 2023; McCarthy, Manieri, et al., 2020). These include crypt top fibroblast (CTF) subsets expressing Bone morphogenic protein (BMP) ligand genes such as *Bmp2*, *Bmp5,* and *Bmp7* to support differentiation, and crypt bottom fibroblasts (CBFs) providing stemness-supporting factors such as Gremlin-1 (*Grem1*) and R-spondin 3 (*Rspo3*). Gastric fibroblast populations are less well characterized but display overall similarity to intestinal fibroblasts (Kim et al., 2020). GI fibroblasts promote tissue regeneration during inflammation, and abnormally and persistently activated fibroblasts are noted in fibrosis, chronic inflammatory diseases, and tumorigenesis (Wei et al., 2021). The fibroblast subsets driving PJS polyposis have not been characterized.

There have been several attempts to reduce tumor burden in PJS models. Despite some effects using long-term high-dose treatment of PJS mice with rapamycin (Robinson et al., 2009), the polyposis appears to develop independently of the AMPK-mTORC1 axis (Cotton et al., 2022; Ollila et al., 2018). COX2 expression is elevated in some polyps, and a COX2 inhibitor, celecoxib, was noted to reduce polyp burden (Udd et al., 2004), indicating importance for

inflammatory pathways. We (Ollila et al., 2018) and others (Poffenberger et al., 2018) have shown that the polyps display active JAK/STAT signaling and that JAK inhibition dramatically reduces the PJS polyp burden. However, inhibition of JAK signaling is associated with significant side effects hindering long-term use. The precise mechanisms through which LKB1-deficient stromal cells promote PJS polyposis are unclear, complicating the design of targeted therapies with acceptable safety profiles.

In this study, we compare fibroblast populations in normal tissue and polyps using scRNA-seq of the normal gastric antrum and antral polyps and investigate tumorigenic mechanisms. We report a tumor-enriched fibroblast subpopulation characterized by the expression of ST2 (encoded by *Il1rl1*) and Interleukin-11 (*Il11*) driving PJS polyposis. Importantly, we show that targeting IL-11 impedes the LKB1 loss-induced fibroblast activation and efficiently reduces tumorigenesis in PJS mouse models. Our findings shed light on the role of stromal cells in driving the development of PJS polyps and suggest IL-11 as a potent therapeutic target for PJS-associated tumors.

Results

Enrichment of *ST2*$^+$ fibroblasts in PJS polyps

To get insight into the distribution and signaling activity of different cell types in PJS tumorigenesis, we performed scRNAseq of polyps and normal tissue isolated from *Twist2-Cre;Lkb1$^{fl/+}$;R26R-tdTomato* mice. In these mice, *Lkb1* heterozygosity is restricted to mesenchymal lineages, yet they develop PJS-type polyps with similar kinetics compared to *Lkb1$^{+/-}$* mice (Ollila et al., 2018). The polyps and adjacent normal tissue were dissected from the distal part of the stomach, the antrum, structurally similar to the colon and intestine with epithelial crypts (glands) opening to the gastric lumen, and the bottom of the crypt hosting the epithelial stem cells marked by Lgr5 (Leushacke et al., 2013). In total, we sequenced 17933 cells that, after quality control, resulted in 14641 cells representing 6112 cells from normal antrum and 8529 cells from polyps, clustering into epithelial cells (*Epcam$^+$*), endothelial cells (*Pecam1$^+$*), immune cells (*Ptprc$^+$*), smooth muscle cells (*Hhip$^+$, Myh11$^+$*), pericytes (*Rgs5$^+$*), and fibroblasts (*Pdgfra$^+$, Pdpn$^+$*) (Figure 1A). All cell types were found in both normal tissue and polyps (Supplementary Figure 1A-B). The *tdTomato* reporter expression, indicating recombination, was

noted in mesenchymal populations but was absent in epithelial and immune cells, indicating that LKB1 loss in immune cells such as T-cells is not required for polyp formation (Supplementary Figure 1C).

As fibroblasts appear to represent important tumor-initiating cells in PJS polyposis (Ollila et al., 2018), the analysis initially focused on fibroblast populations. We identified five distinct fibroblast clusters positive for the pan-fibroblast markers *Pdpn*, *Pdgfra,* and *Col6a1* (Figure 1B-C)*.* We compared these with fibroblast populations characterized in the small intestine and colon (Brügger et al. 2020; Brügger and Basler 2023; Pærregaard et al. 2023), consisting of: i) the *Pdgfra*-low, $Cd34^+$, $Cd55^+$ crypt bottom fibroblasts (CBFs) forming the niche for the stem cells residing at the crypt bottom by secreting stemness-supporting factors such as R-Spondin 3 (encoded by *Rspo3*) as well as the Bmp inhibitor Gremlin 1 (encoded by *Grem1*); and ii) the *Pdgfra*-high, $Cd34^-$, $Cd55^-$ crypt top fibroblasts (CTFs), promoting epithelial differentiation in the upper parts of the crypts by expressing e.g. BMP ligands. In our gastric fibroblasts, clusters 1 and 2 were positive for *Cd55* and *Cd34* and expressed *Grem1* and *Rspo3* along with low levels of *Pdgfra,* suggesting that they represent CBFs. However, as opposed to colonic and intestinal fibroblasts where *Cd81* is specifically expressed in one subset of the *Pdgfra* low fibroblasts (Brügger et al., 2020; McCarthy, Manieri, et al., 2020), expression of *Cd81* was similar across all gastric fibroblast populations (Figure 1C). Instead, cluster 1 cells specifically expressed *Pi16*, *Ebf1, Col8a1*, *Ackr4,* and *C3* (Figure 1D), aligning with universal fibroblasts that mainly reside in GI submucosa (Buechler et al., 2021; Muhl et al., 2020; Thomson et al., 2018). We named these cells Pi16-CBFs. The second CBF cluster specifically expressed *Lepr* and *Ctgf* as well as high levels of *Aspn, Sfrp1*, and *Fbln1*, and we labeled them as Lepr-CBFs (Figure 1D). The characteristic CBF gene profile (Figure 1C), along with an absence of Pi16-CBFs markers, suggested Lepr-CBFs are located in the lamina propria close to the crypt/gland base as previously shown for intestinal fibroblasts expressing *Lepr* (Deng et al., 2022; Manieri et al., 2023).

CTFs in the colon and intestine express differentiation-promoting ligands such as *Bmp5*, *Bmp7, Wnt5a, Nrg1,* as well as the subepithelial telocyte marker *Foxl1* (Aoki et al. 2016; Shoshkes-Carmel et al. 2018; Brügger et al. 2020; McCarthy et al. 2020; Lemmetyinen et al. 2023). These ligands were highly expressed in our fibroblast clusters 3-5, indicating a CTF identity (Figure 1C). The cells in cluster 3 were, however, low in *Pdgfra* expression but

expressed high levels of the non-canonical Wnt ligand *Wnt4* (Figure 1C-D), and we named them Wnt4-CTFs. The Wnt4-CTFs resembled a fibroblast subset that was recently shown to be subject to BMP-mediated regional regulation of niche factor expression exerted by subepithelial fibroblasts in the colon and intestine, restricting *Wnt4* expression in the upper parts of the crypts (Kraiczy et al., 2023). Cells from cluster 4 specifically expressed *Sox6* and *F3* along with higher expression levels of *Acta2,* previously identified markers for subepithelial myofibroblasts (SEMFs) corresponding to CTFs (Kinchen et al., 2018; McCarthy, Kraiczy, et al., 2020; Pasztoi & Ohnmacht, 2022) as well as *Bmp3, Efnb2*, and *Procr* (encoding Cd201), previously reported as a CTF marker in the colon (Melissari et al., 2021). We named the cluster 4 cells Sox6-CTFs (Figure 1C-D)*.* Finally, cluster 5, in addition to the CTF-like expression pattern, specifically expressed *Il1rl1* (encoding for ST2, the IL-33 receptor) along with inflammation and tissue growth-associated genes such as *Ptgs2*, *Fgf7, Il11,* and *Stc1*. We named the cluster 5 ST2-CTFs (Figure 1C-D).

We have previously shown that Interleukin-11 (IL-11) expression is increased in polyps of PJS patients and mouse models (Ollila et al., 2018), and IL-11 is known to induce *Il1rl1* expression in fibroblasts (Widjaja, Chothani, et al., 2022). Moreover, increased expression of COX2, encoded by *Ptgs2,* has also been associated with PJS tumorigenesis (Udd et al., 2004), and stromal PTGS2 was previously shown to promote intestinal tumorigenesis (Roulis et al., 2020), suggesting functional importance for the ST2-CTFs. Indeed, the ST2$^+$ fibroblast subset was almost exclusively found in polyps (Figure 1E). Interestingly, also the Wnt4-CTF population was enriched in polyps and expressed low levels of the ST2-markers *Il11* and *Il1rl1* in the polyp context, suggesting that ST2$^+$ fibroblasts may represent an activated state of the Wnt4-CTFs (Figure 1F). The Pi16-CBF population, on the other hand, was primarily derived from control tissue (Figure 1E), consistently with the likely localization of the Pi16$^+$ universal fibroblasts in the submucosa. Tissue staining confirmed high and mesenchyme-specific expression of the ST2 protein in polyps of *Twist2-Cre;Lkb1$^{fl/+}$* as well as the *Lkb1$^{+/-}$* mice (Figure 1G). The staining was most prevalent in the stroma of polyp edges, verifying the crypt top identity of the ST2-CTFs (Figure 1G).

In summary, we found that the fibroblast subsets in antral mucosa can be classified similarly to previously characterized intestinal and colonic fibroblasts but with specific differences, such as the lack of specificity in the *Cd81* expression (Table 1). Notably, one of the

fibroblast clusters, the ST2-CTF population, was significantly enriched in tumors, suggesting these cells are tumor-promoting cells in PJS.

Table 1. Identified gastric fibroblast subpopulations, top markers, and corresponding populations in the colon and intestine.

Cluster	Name	Specific markers	Pdgfra	Niche factors	Corresponding population in colon/intestine	Refs
1	**Pi16-CBFs**	Pi16, Ebf1	low	Grem1, Rspo1	Trophocyte / Pdgfra low, Cd81+ / CBF1 / Interstitial fibroblast	(Brügger et al., 2020; Buechler et al., 2021; McCarthy, Manieri, et al., 2020; Muhl et al., 2020)
2	**Lepr-CBFs**	Ctgf, Lepr	low	Grem1, Rspo1	Pdgfra low, Cd81- / CBF2	(Deng et al., 2022)
3	**Wnt4-CTF**	none	low	Bmp5, Bmp7, Nrg1, Wnt4	Pdgfra low, Cd81- / CBF2	(Kraiczy et al., 2023)
4	**Sox6-CTF**	Sox6, Bmp3	high	Bmp3, Bmp5, Bmp7, Nrg1, Wnt5a	SEMF / Telocyte / Pdgfra high / CTF	(Kinchen et al., 2018)
5	**ST2-CTF**	Il1rl1, Il11	high	Bmp5, Bmp7, Nrg1, Wnt5a	IAF/ Telocyte / Pdgfra high / CTF	(Kinchen et al., 2018; Nishina et al., 2021; Smillie et al., 2019)

CTF-specific loss of LKB1 induces polyposis and early expression of *Il1rl1* and *Il11*.

Based on our finding that ST2-CTFs were enriched in PJS tumors, we aimed to investigate whether depleting Lkb1 in CTFs was sufficient to induce ST2-CTF genes and polyp formation. To address this, we crossed the *Lkb1^{fl}* mice with *Foxl1-Cre* mice that restrict the Cre activity to subepithelial fibroblasts, representing the CTFs (Aoki et al., 2016) (Figure 1C). The *Foxl1-Cre;Lkb1$^{fl/fl}$* mice started showing signs of discomfort at 4 months of age, and autopsies confirmed extensive gastric polyposis that severely obstructed the GI tract (Figure 2A-B). Interestingly, mice with heterozygous loss of Lkb1 (*Lkb1$^{+/-}$* or *Twist2-Cre;Lkb1$^{fl/+}$*) develop polyps with significantly slower kinetics and survive until about one year of age (Ollila et al., 2018; Rossi et al., 2002). The fast polyposis in *Foxl1-Cre;Lkb1$^{fl/fl}$* mice confirmed our previous finding that homozygous loss of *Lkb1* in fibroblasts dramatically accelerates tumor formation (Ollila et al., 2018). Importantly, lineage tracing demonstrated that stromal loss of *Lkb1* in

Foxl1-expressing cells resulted in an expansion of *Lkb1*-deficient fibroblasts in the polyp stroma (Figure 2C).

To investigate if LKB1 loss induced the expression of the polyp-enriched ST2-CTF marker genes independently of polyp formation, we sorted gastric fibroblasts from young healthy *Foxl1-Cre;Lkb1$^{fl/fl}$;R26R-tdTomato* and control mice prior to tumor formation. Loss of *Lkb1* expression and concomitant increase of the ST2 markers *Il11* and *Il1rl1* was observed in the *Foxl1-* lineage fibroblasts of the *Lkb1$^{fl/fl}$* mice, showing that the ST2-CTF-like signature is a primary result of LKB1 loss (Figure 2D-E).

LKB1 loss recapitulates the inflammatory ST2-CTF transcriptome.

The ST2-CTFs expressed inflammatory and repair-associated genes (Figure 1D), and we sought to compare the ST2-CTF signature (top 25 genes expressed in the ST2-CTFs, Supplementary Table 1) with publicly available datasets related to GI injury and inflammation. The ST2 signature was strongly induced in the scRNAseq analysis of mouse colon fibroblasts treated with dextran sodium sulfate (DSS), inducing acute colon inflammation (Kinchen et al., 2018) (Figure 3A). Significant enrichment was also observed when the ST2 signature was compared to lipopolysaccharide (LPS)-treated primary intestinal fibroblasts, as well as with DSS-treated colon tissue lysates (Figure 3B). The ST2 signature was also enriched in inflammation-associated fibroblasts of ulcerative colitis patients (Smillie et al., 2019) (Figure 3C-D), indicating the signature is induced in inflammatory/regenerative fibroblasts in both mouse and human GI tract.

Consistently with our observations linking the ST2$^+$ fibroblasts to inflammatory fibroblasts, it was recently reported that *Lkb1* loss sensitizes mouse embryonic fibroblasts (MEFs) to inflammatory stimuli, such as IL-1β and LPS, resulting in enhanced induction of cytokine and chemokine expression (Compton et al., 2023). Thus, we investigated the potential role of inflammation in *Lkb1*-deficient fibroblasts and PJS tumorigenesis. In PJS, hamartomatous polyps develop throughout the GI tract, indicating that the underlying mechanisms are shared, and small intestinal polyps also develop in *Lkb1$^{+/-}$* mice (Bardeesy et al., 2002) and *Twist2-Cre;Lkb1$^{fl/+}$* mice (our unpublished observations). We cultured adult primary intestinal fibroblasts from *Lkb1$^{fl/fl}$* mice and deleted *Lkb1* using adenoviral delivery of CRE recombinase, while control cells were transduced with a green fluorescent protein (eGFP)-encoding virus

(Figure 3E). The transcriptional effects were assessed by quantitative PCR and bulk RNA sequencing. As expected, *Lkb1*-deficiency resulted in robust up-regulation of *Il1rl1* and *Il11* (Figure 3F)*,* and the transcriptome was highly enriched for the ST2-CTF signature (Figure 3G). Similarly to the ST2-CTF signature (Figure 3B-C), the *Lkb1*-deficient fibroblast signature (Supplementary Table 1) was significantly enriched for inflammation-associated datasets, indicating that *Lkb1* deletion is sufficient to recapitulate the transcriptional features of inflammatory/regenerative fibroblasts (Figure 3H), consistently with the increased *Il11* and *Il1rl1* expression in *Lkb1*-deficient fibroblasts of tumor-free mice (Figure 2D-E). Next, we studied the inflammatory response of *Lkb1*-deficient intestinal fibroblasts by exposing them to LPS (Figure 3I-J). Interestingly, LPS treatment resulted in robust further induction of *Il11* and *Il1rl1* as well as other ST2 signature genes (*Hgf*, *Fgf7*, *Sct1* and *Ptgs2*), IL6-family cytokines (*Il6*, *Lif*) and the chemokine *Cxcl5* (Figure 3K), indicating that the *Lkb1* loss induced inflammatory phenotype can be further stimulated by exogenous inflammatory factors in primary intestinal fibroblast, similarly to previously shown in MEFs (Compton et al., 2023), but with a more robust response for *Il11* as compared to *Il6*. Of note, the expression of most investigated genes was significantly increased in *Lkb1*-deficient cells already without LPS treatment (Figure 3K).

Inflammation exacerbates the LKB1 loss-mediated tumor-promoting phenotype.

The higher expression of the LPS response genes in *Lkb1*-deficient cells suggested a functional role for inflammation in promoting PJS polyposis. To address if the inflammatory phenotype of fibroblasts has an effect on epithelial growth, we set up an organoid-fibroblast co-culture assay where wild-type (WT) or *Lkb1*-deficient primary intestinal fibroblasts were embedded into a 3D matrix (Matrigel) with WT intestinal epithelial organoids (Figure 4A-B). Intestinal epithelial organoids grow in Matrigel as three-dimensional structures recapitulating the homeostatic crypt-villus architecture of intestinal epithelium (Sato et al., 2009), but tumorigenic and regenerative organoids fail to form the correct 3D-structure and instead grow as hyperproliferative spheroids (Langlands et al., 2018; Nusse et al., 2018). As determined by eye and quantified using our automated organoid classification tool "Tellu" (Domènech-Moreno et al., 2023), *Lkb1*-deficient fibroblasts induced an increased frequency of spheroids as compared to organoids in the cocultures, indicating that a regenerative/tumorigenic phenotype of the WT intestinal epithelium is promoted by *Lkb1*-deficient fibroblasts (Figure 4C). This was consistent

with *Lkb1*-deficient fibroblasts promoting polyp formation *in vivo*. The spheroid formation was further induced by LPS treatment in both WT and *Lkb1*-deficient fibroblast cocultures, suggesting that inflammation-induced growth-promoting signals from fibroblasts to epithelium are recapitulated by LKB1 loss and further boosted by inflammatory stimuli (Figure 4A-C).

To address the potential impact of inflammation-induced fibroblast reprogramming in PJS polyposis *in vivo*, we subjected *Twist2-Cre;Lkb1*[fl/+] mice to DSS-induced chronic inflammation (Figure 4D). Notably, while DSS is best known to result in colonic inflammation, the entire GI tract, including the stomach, is affected (Elsheikh et al., 2012). We observed a significantly reduced survival of DSS-treated *Twist2-Cre;Lkb1*[fl/+] mice (Figure 4E), linked with an increased gastric tumor size and a trend towards increased tumor burden (Figure 4F). Furthermore, subjecting the mice to a carcinogen (Azoxymethane, AOM) followed by DSS inflammation led to significantly increased gastric tumor burden in *Twist2-Cre;Lkb1*[fl/+] mice (Supplementary Figure 2A-B). The colonic tumor burden induced by AOM-DSS was also more prominent in *Twist2-Cre;Lkb1*[fl/+] mice compared to WT controls, albeit the difference did not reach statistical significance (Supplementary Figure 2C). Altogether, our results indicate that while exogenous inflammation is not required for PJS polyposis, it boosts the tumorigenic potential of LKB1 deficient mesenchyme, suggesting inflammation as a contributing mechanism driving polyp development.

ST2-CTFs present a unique signaling pattern targeting fibroblasts and epithelial cells

Next, we set out to take a broader look into our scRNAseq data and also included the non-fibroblast cell populations. Epithelial (*Epcam*[+]) cells included surface cells (*Muc5ac*[+]), proliferative isthmus cells (*Mki67*[+]), gland and base cells (*Aqp5*[+], *Muc6*[+]), as well as tuft cells (*Dclk1*[+], *Trpm5*[+]), identified using previously established markers (Haber et al. 2017; Xiao and Zhou 2020; Tan et al. 2020) (Supplementary Figure 3A-B). The distribution of the cell types was similar between normal antrum and polyp, in agreement with hamartomatous polyp type (Supplementary Figure 3C). However, the differential expression analysis of the epithelial cell fraction revealed some differences, such as upregulation of the regenerative markers *Reg3b* and *Reg3g* in polyps, confirming the epithelial origin of these changes noted in previous analyses of polyps lysates (Ollila et al., 2018) (Supplementary Figure 3D). In addition, we observed

significantly reduced epithelial expression of *Pgc* (encoding for Pepsinogen C) and *Muc5ac* (encoding for the secreted Mucin 5AC) in the polyps, indicating that some of the previously observed Lkb1 loss-driven epithelial differentiation defects (Udd et al., 2010) are at least in part driven by stromal cells (Supplementary Figure 3D). Reduced expression of Pepsinogen C was confirmed using immunohistochemistry (Supplementary Figure 3E). Analysis of immune cells (Supplementary Figure 4A-C) and endothelial cells (Supplementary Figure 4D-E) identified all major cell types (James et al., 2022; Kalucka et al., 2020) and suggested a putative increase of neutrophils in the polyp-derived clusters as previously reported by Poffenberger et al., (2018) (Supplementary Figure 4C, D).

Next, we analyzed cell-cell communication across all cell clusters identified in the control and polyp datasets using CellChat (Jin et al., 2021). This analysis revealed increased interactions within the polyp compared to the control antrum, suggesting activated signaling (Supplementary Figure 5A). Next, we plotted the total signaling strength from all cell types in a 2D space. Interestingly, the strongest outgoing and incoming signaling was observed in fibroblasts and was highest in the CTF clusters, including ST2-CTFs, suggesting that CTFs represent the most active cell types regarding paracrine signaling in our dataset (Figure 5A). We then further analyzed the outgoing and incoming cell-cell communication patterns across clusters. Five outgoing communication patterns were identified, mostly clustering along the major cell types (epithelial cells, endothelial cells, myeloid and lymphoid lineage immune cells, and fibroblasts) (Figure 5B). Interestingly, the tumorigenic ST2$^+$ fibroblasts exhibited a unique outgoing communication pattern, Pattern 5, separating from the signaling pattern of other fibroblast subsets that signaled according to Pattern 1. The main differences were the active IL6 family signaling and parathyroid hormone (PTH) family signaling, as well as increased signaling from NRG, HGF, and FGF families, suggesting that emergence of the ST2-CTFs cells induces an altered stromal niche (Figure 5B). Interestingly, the only ligand contributing to the IL-6 family signaling was IL-11 (Supplementary Figure 5B). Upon closer examination, we identified distinct receivers of the ST2-CTF outgoing cell communication. The signals to epithelial cells included NRG and HGF family ligands, while PTH was received by pericytes and IL-11 signal by fibroblasts (Figure 5B-C). Interestingly, analysis of the ligand-receptor pairs constituting the signaling patterns predicted HGF-MET and NRG1-ERBB3 as the most active pathways emanating from ST2-CTFs to epithelial cells, suggesting functional relevance as noted before in

regenerative and tumor contexts (Joosten et al. 2017; Jardé et al. 2020; Lemmetyinen et al. 2023) (Supplementary Figure 5B-C). FGF7-FGFR1 signaling was predicted to mainly target fibroblasts, while EREG-EGFR signaling was active to both epithelial cells and fibroblasts. Interestingly, the activation of IL-6 family signaling was attributed to increased expression of IL-11 signaling from ST2-CTFs to IL11RA1-IL6ST (Gp130) receptor-expressing fibroblasts populations (Supplementary Figure 5B-D). This observation suggested an upstream role for IL-11 in fibroblast reprogramming.

Inhibition of IL-11 blocks the ST2$^+$ signature and reduces PJS polyposis.

The scRNAseq data suggested a role for IL-11 in fibroblast reprogramming (Figure 5C, Figure 6A). Consistently, the *Lkb1*-deficient intestinal fibroblast signature was enriched in the transcriptome of IL-11-expressing fibroblasts isolated from inflammation-induced mouse colon tumors (Nishina et al., 2021), indicating shared functional outcomes of IL-11 expression and LKB1 loss (Figure 6B). To explore whether the *Lkb1*-deficient ST2-CTF-like fibroblast transcriptome was dependent on IL-11, we engineered *Il11* knock-out (KO) immortalized intestinal fibroblasts generated from *Lkb1$^{fl/fl}$* mice using CRISPR-Cas9 (Supplementary Figure 6A). Remarkably, the induction of the ST2$^+$ signature observed upon Cre-mediated *Lkb1* loss was significantly reduced in the absence of *Il11* (Figure 6C, Supplementary Figure 6B), implying that the *Lkb1*-loss mediated reprogramming of intestinal fibroblasts into the ST2-CTF-like phenotype is at least partially dependent on IL-11.

Interestingly, previous work has indicated a crucial role for IL-11 in cardiorenal fibrosis (Schafer et al., 2017), and a blocking antibody to IL-11, X203, inhibits fibroblast activation *in vitro* and *in vivo* (Ng et al., 2019). We hypothesized that blocking IL-11 in mice modeling PJS could effectively inhibit fibroblast reprogramming and tumorigenesis following *Lkb1* inactivation. To address this, the *Lkb1$^{+/-}$* mice were treated with X203 or IgG intraperitoneally at 20 mg/kg dose weekly from 30 to 50 weeks of age (Figure 6D). Notably, X203 treatment significantly reduced the polyp number and polyp burden while not affecting individual polyp size, implicating that IL-11 signaling plays a significant role in PJS polyp initiation (Figure 6 E-H). These results indicate that the LKB1 loss-mediated fibroblast reprogramming leading to PJS tumorigenesis is driven by therapeutically targetable expression of IL-11.

Discussion

Here, we investigated fibroblast heterogeneity and cell-cell interactions driving PJS polyposis using mouse models. Interestingly, the polyps were enriched for a tumorigenic fibroblast population, ST2-CTFs, presenting with a high expression of growth factors and inflammatory-associated genes such as *Il11*, *Il1rl1*, *Fgf7*, *Hgf*, *Stc1*, and *Ptgs2*. Importantly, loss of LKB1 was sufficient to induce an ST2-type inflammatory phenotype in the fibroblasts, and inflammatory insults further enhanced this phenotype. Thus, while LKB1 loss alone altered the fibroblast function, further activation via local inflammation may be involved in the etiology of PJS polyp development, also supported by the recent report on *Lkb1*-deficient MEFs being sensitized to inflammatory stimuli (Compton et al., 2023) and other reports connecting LKB1 loss to increased inflammation (Ollila et al. 2018; Poffenberger et al. 2018). Interestingly, the ST2-CTFs represented only a subset of fibroblasts in the polyps, densely populating the polyp edges. Since all fibroblasts in the polyps were heterozygous for LKB1 in our model, this observation suggests that additional insults, perhaps originating from GI lumen, are required for the fully activated ST2-CTF phenotype. This poses an interesting question of whether other than genetic LKB1 inhibition, such as inhibitory phosphorylation (Widjaja, Viswanathan, et al., 2022), is associated with fibroblast activation in inflammatory conditions. Since PJS patients are heterozygous for LKB1, and the mouse models with homozygous stromal *Lkb1* loss present with faster polyp formation (Ollila et al., 2018, this study), inhibition of the wild-type LKB1 allele could contribute to further fibroblast activation also in PJS polyposis. This would be an interesting topic for a future study.

GI fibroblast heterogeneity has been under intensive investigation in recent years (Kinchen et al. 2018; McCarthy et al. 2020; Brügger et al. 2020; Deng et al. 2022; Jasso et al. 2022; Pærregaard et al. 2023; Kraiczy et al. 2023) with little data on gastric fibroblasts (Kim et al. 2020). We found that antral fibroblasts clustered similarly to intestinal and colonic fibroblasts, including the formation of Wnt and Bmp gradients (Brügger et al., 2020; McCarthy, Manieri, et al., 2020) but with differences in specific marker expression: for example, in the gastric mucosa *Cd81* was expressed across all fibroblast subsets, as opposed to intestinal and colonic fibroblasts where *Cd81* specifically marks one of the fibroblast population providing the trophic support for stem cells (McCarthy et al. 2020; Brügger et al. 2020). The corresponding population in our dataset was specifically marked by *Pi16*, also known as a marker of "universal fibroblasts"

(Buechler et al., 2021; Muhl et al., 2020), consistent with submucosal location away from the tissue parenchyma. We also found a Lepr[+] CBF population. Interestingly, the Lepr[+] fibroblasts provide stem cell niche factors for hematopoietic stem cells (Comazzetto et al. 2019) and intestinal stem cells (Deng et al., 2022). Expression of *Grem1* and *Rspo3* in antral Lepr-CBFs suggested they represent the primary stem cell supporting fibroblasts also in the gastric antrum. Our analysis also revealed an intermediate fibroblast population, Wnt4-CTFs, with similarities to the recently published Wnt4[+] fraction of the *Pdgfra*-low, *Cd81[-]* cells in the colon and intestine (Kraiczy et al., 2023), suggesting that the division to *Grem1*-expressing and *Wnt4*-expressing *Pdgfra*-low mucosal fibroblasts is conserved along the GI tract. Cell-cell communication analysis predicted active BMP signaling between these fibroblast subsets (Figure 5C), suggesting that division to Lepr-CBFs and Wnt4-CTFs is regulated by BMP signaling also in the gastric antrum. The relevance of Wnt4 signaling to GI homeostasis is unclear, but previous reports suggest functions related to promoting differentiation (Eliazer et al., 2019; Zhang et al., 2021). The Sox6-CTF cluster identified in our analysis presented with markers suggesting a subepithelial location away from the crypt/gland bottom, consistent with previous data (Kinchen et al., 2018). Interestingly, SOX6 has been previously implicated in suppressing B-catenin-mediated transcription, consistent with the localization away from the stem cell zone (Iguchi et al. 2007).

Our results indicate that Lkb1 loss-induced IL-11 signaling is critical for activating the ST2-CTF phenotype. IL-11, a proinflammatory IL-6 family cytokine, is upregulated primarily in stromal fibroblasts in several non-malignant inflammatory diseases (Fung et al., 2022). Additionally, transgenic overexpression of IL-11 in mesenchymal cells leads to spontaneous inflammation of the gastrointestinal tract and spontaneous death in mice, demonstrating its direct implications on inflammation (Lim et al., 2020). In patients with ulcerative colitis, inflammatory-associated fibroblasts (IAF) characterized by high expression of IL-11 are expanded by more than 100-fold in inflamed tissue (Smillie et al., 2019). IL-11-expressing fibroblasts have been shown to emerge during inflammation-associated colon tumorigenesis in mice and be functionally associated with tumor formation (Nishina et al., 2021). Furthermore, the IL11[+] fibroblasts harbor a unique gene signature correlating with poor prognosis in colorectal cancer (Nishina et al., 2021). Interestingly, activation of STAT3 through IL-6 or IL-11 in cancer-associated fibroblasts (CAFs) is associated with poor prognosis (Heichler et al., 2020).

Interestingly, a stronger correlation between elevated STAT3 activation and IL-11 has been noted in human GI cancers as compared to IL-6 (Putoczki et al., 2013). Thus, the observed therapeutic responses to JAK inhibitors in PJS models (Ollila et al., 2018; Poffenberger et al., 2018) could result from the inhibition of the JAK/STAT3-driven fibroblast activation. The JAK inhibition could also act on innate immune cells and reduce the production of inflammatory cytokines such as IL-1β, a potent activator of LKB1-deficient MEFs (Poffenberger et al., 2018).

Interestingly, our CellChat cell-cell communication analysis provided a detailed overview of the dynamic interplay between ST2-CTFs and the surrounding cellular components, including high IL-11 signaling to other fibroblast populations. Also, BMP, non-canonical WNT as well as FGF family signaling were predicted to affect other fibroblasts predominantly, and it would be interesting to address the contribution of these signals to the fibroblast activation, as well as to the spatial fibroblast distribution (Figure 5C, Supplementary Figure 5C). However, despite the critical role of fibroblasts, the PJS polyps consist mostly of highly proliferating epithelial cells (Ollila et al., 2018; Udd et al., 2010). Our finding that IL-11 signaling impacts the fibroblast phenotype raises the question of the mechanism driving the epithelial hyperproliferation. Interestingly, ST2-CTFs expressed a plethora of growth-promoting factors such as HGF, NRG1 as well as PTGS2 (the rate-limiting enzyme for Prostaglandin E2), known to promote the growth of GI epithelial cells (Takahashi et al. 1995; Ding et al. 2018; Roulis et al. 2020; Jardé et al. 2020; Lemmetyinen et al. 2023). Thus, ST2-CTFs appear to play a pivotal role in reprogramming the fibroblasts into an inflammatory/regenerative state, as well as driving the proliferative response observed in the epithelium, likely driven by a combination of growth-promoting factors. In addition, it remains possible that the upregulation of fibroblast-originating chemokines such as CXCL5, and the subsequent recruitment of immune cells such as neutrophils, plays a functional role in polyp formation. However, as the immune cells in our model are wild-type for LKB1, any putative polyp-promoting functions likely represent secondary changes.

Our discovery that IL-11 was required for the full inflammatory, ST2-fibroblast transcriptome induced by LKB1 loss points towards a feed-forward loop between LKB1 inhibition and IL-11 expression, along the lines of a previous report (Widjaja, Viswanathan, et al., 2022). Notably, the efficient blocking of polyp formation using a blocking antibody for IL-11 in PJS mouse models clearly indicates that IL-11 signaling is a key step in PJS tumorigenesis,

which can be exploited therapeutically. As IL-11 expression is very low under normal homeostasis (Nishina et al. 2021), its targeting is expected to pose minimal side effects, opening a possible avenue for continuous treatment to reduce the need for surgical polyp removal, thus contributing to the quality of life in PJS patients. This calls for the initiation of clinical trials addressing the efficiency of IL-11 targeting in PJS. In conclusion, by unraveling the interplay between LKB1 loss, IL-11 signaling, and the fibroblast phenotype, our study provides valuable insights into the tumorigenic potential and pathogenic processes associated with these specific fibroblasts. These results contribute to a deeper understanding of the complex stromal microenvironment in gastric mucosa and have implications for future therapeutic interventions targeting the ST2-CTF population by targeting IL-11.

Methods

Animal experiments. The following mouse strains were used in this study: Twist2-Cre (The Jackson Laboratory #008712, (Yu et al., 2003)), Foxl1-Cre (The Jackson Laboratory #017738, (Sackett et al., 2007), Lkb1 flox (*Lkb1^{fl}*) (The Jackson Laboratory #014143, (Bardeesy et al., 2002)), *Lkb1$^{+/-}$* KO (Cyagen #S-KO-04594). Rosa26-tdTomato (The Jackson Laboratory #007914)(Madisen et al., 2010). Animals were housed at the University of Helsinki Laboratory Animal Center and the SingHealth Experimental Medicine Centre according to national and international legislation and guidelines and experiment approvals (ESAVI/3816/2023, 2019/SHS/1481 and 2019/SHS/1483). Mice had free access to food and water. Male and female mice with a similar gender ratio were used in all experiments unless otherwise stated.

Antibody treatment in *Lkb1$^{+/-}$* mice. To evaluate putative beneficial effects of neutralizing IL-11 signaling, 30-week-old *Lkb1$^{+/-}$* mice (22 males and 24 females) were block randomized to receive either anti-IL11 (X203, Genovac) or an IgG isotype control (11E10, Genovac) at a dose of 20 mg/kg once per week IP. At 50 weeks, mice were euthanized by ketamine (100 mg/kg) and xylazine (10 mg/kg) mixture given IP, and the stomach extracted for polyp assessment. The stomach and proximal duodenum were opened, flushed with ice cold PBS and gently swapped with a cotton bud to remove debris, prior to image capture under the stereomicroscope (Olympus, SZ61). Gastric polyps were enumerated, and size quantified using ImageJ (version 1.53, NIH). Polyp size was reported as the average polyp area per stomach, and total polyp

burden was assessed as the summation of individual polyps' area per mouse. 30-week-old untreated $Lkb1^{+/-}$ mice (7 males and 12 females) were included as a reference group prior to antibody treatment to gauge therapy effectiveness. For statistics, we performed a two-way ANOVA with Sidak post-hoc analysis for the two factors, treatment (30w untreated vs. 50w X203 vs. 50w IgG) and sex (males vs. females).

GI inflammation and inflammation-induced tumorigenesis models. Male mice were used for the inflammation experiments. The acute GI inflammation was induced by replacing the drinking water of mice with 3% dextran sodium sulfate (DB001, TdBlabs) diluted in water for 7 days. The mice were euthanized after a 3-day recovery period with normal drinking water. The chronic inflammation was induced with three rounds of 7-day 3% DSS treatment and a 14-day recovery with normal drinking water. The mice's weight was monitored daily throughout the experiment to monitor weight loss. The survival cohort of DSS-treated and control mice was followed, and the mice were euthanized when they started to show signs of discomfort (abnormal posture and reduced activity) and were autopsied to confirm large gastric polyps as the cause of death. The mouse cohort for tumor burden analysis was euthanized two weeks after the last DSS cycle. For the AOM-DSS inflammation-induced colorectal cancer model, mice were injected intraperitoneally with 10 mg/kg azoxymethane (AOM) in PBS, and after 7 days followed by three cycles of 7 days if 3% DSS and 14 days of recovery as described above. Gastric and colonic polyps were counted and measured under stereomicroscope (Leica MZFLIII).

Gastric tissue dissociation and scRNAseq library preparation
Antral polyps and adjacent antral mucosa from a pair of mice were dissociated into single cells as follows: the tissue was flushed with PBS, minced into < 1 mm pieces with scissors, and placed in a 15 ml falcon tube filled with 5 ml of a collagenase mixed solution (collagenase II (Gibco 17101-019) + IV (ThermoFischer 17104019), 1:1 ratio at 2 mg/ml). The samples were incubated horizontally in a 37°C shaking incubator (220 rpm) for 40 min. During this incubation time, pieces were pipetted up and down 10 times every 10 min, and the collagenase solution was changed after 20 minutes, collecting the supernatant containing the single cells. The supernatant (~10 ml) was then transferred to a clean Falcon tube, and collagenases were inactivated with cold DMEM supplemented with 20% fetal bovine serum (FBS). The samples were passed through a

70 um cell strainer, centrifuged for 10 min at 1100 rpm, and single cells were resuspended in 1–2 ml of PBS. Single-cell suspensions were stained with 7 AAD (LifeTechnologies #A1310) at a 1:1000 dilution prior to FACS sorting to exclude dead cells. 10.000 live cells per sample were sorted at ultra-purity using a Sony SH800Z Cell Sorter at the Biomedicum Flow Cytometry unit. The 10X v3 library kit (Library Construction Kit from 10X Genomics #1000190) was used for scRNA-seq library preparation. Samples were sequenced using a drop-seq-based 10X Genomics machine with a target of 5000 cells per sample.

Isolation of primary intestinal fibroblasts

The small intestine was cleaned with cold PBS using a syringe and 21-gauge needle, cut longitudinally, and rewashed from intestinal contents with cold PBS. The tissue was then placed in a Petri dish and cut into 1 cm pieces with scissors and forceps, and tissue pieces were transferred into a 15 ml Falcon tube. To remove intestinal epithelial cells, 10 ml of pre-warmed EDTA solution (5 mM EDTA and 1 mM DTT) was added to the tissue and incubated for 20 min at 37°C in a shaking incubator with the tubes placed horizontally. The tissue pieces were washed four times with cold PBS by vigorously shaking the tubes for 15 sec and removing the supernatant each time. After the last wash, the supernatant was no longer cloudy. The tissue was further minced to <1 mm using scalpels or scissors. 10 ml of collagenase mix digestion solution (collagenase II + IV mixed at 1:1 ratio at 2 mg/ml - Thermo Fisher Scientific) was added to the tissue and incubated horizontally for 20 min at 37°C in a shaking incubator. The tissue was then pipetted up and down with a 10 ml pipette 15 times. 20 ml of cold fibroblast growth media (DMEM+20% FBS+1% penicillin/streptomycin+1% L-glutamine) was added to stop the reaction. The tissue was then centrifuged at 1400 rpm for 10 min, and the supernatant was removed. The pellet was resuspended in 10 ml of growth media and placed in a culture plate. The plate was washed three times with PBS after 1 h to remove floating cells and debris.

Cell sorting

The antrum tissue was dissected and dissociated into single cells as described above (for scRNAseq tissue sample preparation). Single-cell suspensions were stained with 7-AAD at a 1:1000 dilution. Fibroblasts were enriched by negative selection excluding epithelial, endothelial, immune and red blood cells by staining the cell suspensions with antibodies against

Epcam (eBioscience 11-5791-82), Pecam1 (eBioscience 11-0311-82), Cd45 (eBioscience 11-0451-82), and Tert19 (eBioscience 11-5921-82) conjugated to FITC at a 1:100 dilution for 30 minutes on ice. Sorting was performed using a Sony SH800Z Cell Sorter. Single cells were gated using forward and side scatter characteristics, and live cells were gated by excluding 7 AAD-positive cells. Fibroblasts were enriched by excluding the FITC-stained cells. The remaining cell suspension consisted mainly of fibroblasts, and fibroblasts from the Foxl1-Cre lineage were gated using the TdTomato channel. 10,000 TdTomato+ and TdTomato- cells of the fibroblast enriched fractions were sorted directly into Tri reagent® (TR118, Fisher Scientific), and RNA was extracted according to manufacturer's instructions.

Adenoviral transduction of primary intestinal fibroblasts

1.2 million primary intestinal fibroblasts were seeded in a 10 cm plate. The adenoviruses (AdCre (Iowa-5) or AdGFP (Iowa-4) at MOI 1000 were added to 6 mL of fibroblast growth media (see above), and the cells were incubated overnight with the virus. Following incubation, the media was removed, and the cells were washed once with the fibroblast growth media. The cells were used for experiments 4 days after the transduction.

Co-culture of intestinal epithelial organoids and fibroblasts

Intestinal crypts were isolated from the small intestine of euthanized mice and cultured as previously described (Sato et al., 2009) with some modifications. Briefly, the small intestine was removed and mesentery was removed. The lumen was rinsed with ice-cold PBS using a syringe and gavage needle. The intestine was open longitudinally and gently rubbed to remove mucus. The small intestine was cut into 2-3 mm fragments with small scissors and the fragments were collected in a 50 ml Falcon tube and washed with PBS. The tube was then filled with 10 mM EDTA in PBS, and the tissue pieces were incubated for 45 minutes horizontally on ice. The EDTA-PBS solution was changed 3 times during this period, and gently shaking the tube each time. After an additional 60 minute incubation, the tube was shaken vigorously for 15 seconds to detach the crypts, and the contents were filtered through a 70 μm mesh into a fresh 50 ml Falcon tube to exclude villi and tissue pieces. The crypts were pelleted by centrifugation at 200 g for 2 min, washed with PBS and spun again to collect the crypts. The pellet was resuspended in ENR crypt media consisting of composed of basal organoid media (Advanced Dulbecco's modified

Eagle medium (DMEM):F12 (12634010, Gibco) supplemented with 1 M HEPES pH 7.4 (H0887, Sigma-Aldrich), 1× GlutaMAX (35050061, Gibco), 1× penicillin-streptomycin (P4333, Sigma-Aldrich), 1× N-2 supplement (17502001, Gibco) and 1× B-27 supplement (17504044, Gibco)) supplemented with EGF (Gibco, PMG8041; 50 ng/ml), NOGGIN (Peprotech, 250-38; 100 ng/ml), R-Spondin 1 (R&D Systems 3474-RS; 250 ng/ml) and N-acetylcysteine (A0737, Sigma-Aldrich; 1 μM). Y-27632 (Y0503, Sigma-Aldrich; 10 μM) was included in the ENR medium for the first two days. 20 μl domes containing 75% Matrigel (356231, Corning) and 25% crypt suspension were plated in pre-warmed 48-well plates. In total, approximately 200 crypts were plated per dome together. The domes were overlaid with 350 ul of ENR media and cultured at 37°C in a humidified atmosphere of 5% CO2. For co-cultures with fibroblasts, domes were overlaid with basal organoid media containing 5000 primary fibroblasts from *Lkb1^{fl/fl}* mice previously adenotransducted with AdCre/AdGFP and incubated for 24 h with or without 1ug/ml LPS (Sigma L4516).

Immunohistochemistry and immunofluorescence

Paraffin-embedded tissue sections were subjected to deparaffinization using a series of xylene and ethanol washes. Antigen retrieval was performed at 95°C for 25 minutes using Dako citrate buffer pH 9 (S2367) except for Pepsinogen C, where antigen retrieval was done with 0.05% Trypsin (215240, Difco) in 37C for 10 minutes, followed by 10 min in room temperature. The sections were washed with TBS and permeabilized with 0.1% Triton-X/PBS for 10 minutes at room temperature. A PAP pen was used to outline the sections, and blocking was done using 5% NGS in TBS for an hour at room temperature. Antibodies were incubated overnight at +4°C. Primary antibodies used were anti-RFP (1:500, 600-401-376, Rockland), anti-Pdgfra (1:100, AF1062, R&D Systems) and anti-St2 (1:150 AF1004, R&D Systems) and anti-Pepsinogen C. The following day, sections were washed with TBS and then incubated with a secondary antibody (Alexa Fluor-conjugated antibodies at 1:500) for one hour at room temperature. After washing again with TBS, counterstaining was performed using Hoechst (0.01ng/ml) for one minute, followed by rinsing with water and mounting in Mowiol. The stained sections were scanned using the Pannoramic 250 FLASH III digital slide scanner. Confocal imaging was performed with Leica TCS SP8 CARS Confocal microscope.

Gene editing using CRISPR-Cas9

Mouse primary intestinal fibroblasts from passage 3 were transduced with a retrovirus encoding residue 302 to 390 of p53 (Klefstrom et al., 1997) and grown as a pool. gRNAs targeting mouse Il11 locus were designed using Benchling Software. (gRNA1: 5' GCAGCCATTGTACATGCCGG3' targeting exon 4, gRNA2: 5'AGGGGCAACGACTCTATCTG3' targeting exon 2) and cloned into LentiCRISPRv2 plasmid as described (Sanjana et al., 2014; Shalem et al., 2014). For lentivirus production, 6 million HEK 293FT cells were seeded and transfected after 24 hours with the LentiCRISPRV2-IL11g1 or LentiCRISPRV2-IL11g2 construct together with the pLp Δ8,9 packaging plasmid and VSVG envelope plasmid using lipofectamine™ 2000 transfection reagent (Life Technologies, 11668019) and Opti-MEM reduced serum medium (Gibco, 31985-047). Viruses were collected after 48 hours, filtered through a 0.22 μm filter and stored at -80 °C until use. For transduction, 200.000 immortalized intestinal fibroblasts were seeded in 2 mL growth media on a 6-well plate. The media was replaced after 24 hours with 1 mL growth media mixed with 1 mL of virus media and 8 μg/mL of Polybrene (Merck Millipore, TR-1003-G). One day later, cells were split in growth media and selected using 2 μg/mL puromycin (Gibco, A1113803). The success of Il11 targeting was verified using Sanger sequencing.

Quantitative PCR

qRT-PCR analysis was conducted to examine gene expression levels. For RNA extraction, the TRI Reagent (AM9738, Thermo Fisher Scientific) was used, following the manufacturer's protocols. Subsequently, cDNA synthesis was performed using the TaqMan reverse transcription kit (N8080234, Applied Biosystems), and reverse transcription reactions were conducted using the MJ Research PTC-200 Thermal Cycler (Marshall Scientific). For qRT-PCR, the KAPA SYBR FAST qPCR Master Mix Universal reagent (KK4618, KAPA Biosystems) was utilized. All primers were diluted to a final concentration of 500 nM in nuclease-free water, with a reaction volume of 20 μl. StepOnePlus Real-Time PCR System (Thermo Fisher Scientific) was employed for sample analysis. Actb or Gapdh were used for normalization. The relative fold change in gene expression was determined using the ΔΔCT method described by Livak & (Livak & Schmittgen, 2001) (Livak & Schmittgen, 2001).

scRNA-seq data analysis

Fastq files were aligned to the mouse reference genome (mm10), modified to include the TdTomato sequence, and demultiplexed using CellRanger.v3 (Zheng et al., 2017) setting a cell target number of 5000 cells per sample. Seurat V3 was used for further data processing. Pre-filtering was performed based on percent mitochondria (<15%), UMI count (<100000), and at least 400 features detected to exclude low-quality cells. SCT transformed setting *nfeatures* to 3000 was used for the selection of integration features, and further PrepSCTIntegration was used without reference sample. Based on an elbow plot, 30 principal components were used for RunUMAP and FindNeighbors functions. Clusters were identified using FindClusters, and for exploratory analysis, a range of different resolutions was used (0.3, 0.5, 0.7, 1.0, 1.3). Main clusters were manually mapped based on known markers (Figure 1, FigS1, S2). Data was further sub-clustered within the main cell category, taking the first cell cluster assignment and a further filter based on the expression of the primary marker genes: epithelial cells (Epcam >1 read), fibroblast (Pdpn >1 read), endothelial (Pecam >1 read), and immune (Ptprc >1 read). The subcluster dataset was re-analyzed using the same SCT transform-integration pipeline explained above. To assess the cell markers enriched in each cluster, the Seurat function FindAllMarkers was used with the default 'roc' method and a minimum percentage of cells threshold set at 0.2. For differential expression analysis of the overall differences between tumor and control, FindMarkers function was used with the method set to MAST (Finak et al., 2015) and *min.pct* of 0.2 with the active identities of the Seurat object set to be the tissue type (tumor vs control). ST2 signature of 25 genes was created by selecting the top significant up-regulated genes with a log fold-change above 1.5 against the rest of the fibroblast clusters (Supplementary Table 1).

Cell-cell communication analysis

To analyze cell-cell communication, we utilized CellChat (Jin et al., 2021) with the following settings. A CellChat object was created from a Seurat object that included the identified clusters of all cells. This was achieved using the built-in function createCellChat from the CellChat R package. To ensure a comprehensive analysis, we utilized the complete CellChat ligand-receptor database, encompassing factors related to secretion, extracellular matrix (ECM), and cell-cell contact. We computed the communication probability and inferred the cellular communication network by executing the computeCommunProb method, wherein we set the parameters *raw.use*

and *population.size* as False. This setting was chosen to prevent the size of the clusters from dominating the interaction strength in the dataset. Additionally, we applied a cell-cell communication filter with a minimum cell threshold of 10 (min.cells=10), thereby excluding communication pairs that accounted for only a few cells. For cell-cell communication pattern analysis, we used the CellChat built-in function *identifyCommunicationPatterns* with a rank of k=5 previously inferred by Cophenetic and Silhouette analysis as suggested in their documentation (https://github.com/sqjin/CellChat)

Bulk RNA sequencing

RNA sequencing was performed using 3′UTR RNA sequencing with a 'Pool Seq' method, which is based on the Dropseq approach for single-cell sequencing (Macosko et al., 2015). The method involves priming mRNA using an oligo-dT primer incorporating a 12 bp barcode and an 8 bp unique molecular identifier (UMI) sequence. The UMI enables the removal of PCR duplicates during data analysis. To generate double-stranded cDNA, the single-stranded cDNA was converted using the template-switch effect, followed by PCR amplification using SMART PCR primers. Subsequently, the samples were pooled, and PCR sequencing pools were created using Nextera i7 primers and the Dropseq P5 primers. Sequencing was performed on the NextSeq 500 (Illumina) using the NextSeq High Output 75 cycle flow cell. The obtained fastq files were aligned to the mouse reference genome (GRCm38) using the STAR-aligner (v2.7.8a) (Dobin et al., 2013). To demultiplex the samples and create a count matrix of UMI counts, the 'CreateDGEMatrix' command from the BRB-seqToolsv1.4 suite of tools (https://github.com/DeplanckeLab/BRB-seqTools) was employed. The input for this analysis consisted of R2 aligned BAM files and R1 fastq files. The R1 reads had a length of 21 nucleotides and included a 12-nucleotide sample barcode, an 8-nucleotide UMI, and an additional nucleotide. For publication purposes, individual files for demultiplexed raw fastq files and the processed UMI count matrix of each sample, were generated using the demultiplex function from the BRB-seqTools suite. The CSC - IT Center for Science, Finland, provided the computational resources required for all the data processing steps. The Lkb1 deficient fibroblast signature was created by selecting significant genes upregulated above 2 log fold-change in Lkb1 KO (AdCre transduced) fibroblasts against wild-type fibroblasts (AdGFP transduced) (Supplementary Table 1).

Data Availability

scRNA-seq and bulk RNA seq files will be uploaded to ArrayExpress upon publication.

Acknowledgments

This mouse work was carried out with the support of HiLIFE Laboratory Animal Centre Core Facility, University of Helsinki, Finland. The 3′ RNA sequencing analysis was conducted at the Biomedicum Functional Genomics Unit, located at the Helsinki Institute of Life Science and Biocenter, Finland, affiliated with the University of Helsinki. The gRNA constructs were generated by the Genome Biology Unit supported by HiLIFE and the Faculty of Medicine, University of Helsinki, and Biocenter Finland. LentiCRISPR v2 was a gift from Feng Zhang (Addgene plasmid #52961). Images were generated using 3DHISTECH Pannoramic 250 FLASH III digital slide scanner at the Genome Biology Unit supported by HiLIFE and the Faculty of Medicine, University of Helsinki, and Biocenter Finland. The flow cytometry was performed at the HiLIFE Flow Cytometry Unit, University of Helsinki. Roni Liimatta is thanked for help in gRNA design, and Melissa Montrose, Reeta Huhtala, Riikka Saikkonen and Myriam Sevigny for technical assistance. We are grateful to Chen Xie and Daryl Yeong for their technical assistance rendered to the colony maintenance, antibody treatments, and tissue harvests of the *Lkb1*$^{+/-}$ mice performed at National Heart Centre Singapore, Singapore.

Funding

This study was supported by grants from Academy of Finland (grant numbers 320145 to S.O. and 1320185 to T.P.M.), University of Helsinki Research Foundation, Cancer Foundation, Biomedicum Helsinki-säätiö and the K. Albin Johanssons Stiftelse (to E.D.M.). S.A.C is supported by the Ministry of Health (MOH) National Medical Research Council (NMRC) Singapore Translational Research Investigator Award (STaR21nov-003), Clinician Scientist Individual Research Grant (CIRG18nov-0007), Goh Cardiovascular Research Award (Duke-NUS-GCR/2015/0014), and the Tanoto Foundation. W.-W.L is supported by the Advanced Manufacturing and Engineering Young Individual Research Grant (AME YIRG) of Agency for Science, Technology and Research (A*STAR) award (A2084c0157).

Competing Interests

S.A.C is a co-inventor of the published patents related to IL-11 therapies: WO/2017/103108, WO/2018/109170, WO/2018/109174. S.A.C. and W.-W.L are co-inventors of the published patent WO/2019/073057. S.A.C. is a co-founder and shareholder of Enleofen Bio PTE LTD, a company that develops anti-IL11 therapeutics, whose preclinical IL-11 platform was acquired by Boehringer Ingelheim in 2019 for further development.

References

Aoki, R., Shoshkes-Carmel, M., Gao, N., Shin, S., May, C. L., Golson, M. L., Zahm, A. M., Ray, M., Wiser, C. L., Wright, C. V. E., & Kaestner, K. H. (2016). Foxl1-expressing mesenchymal cells constitute the intestinal stem cell niche. *Cellular and Molecular Gastroenterology and Hepatology*, *2*(2), 175–188. https://doi.org/10.1016/j.jcmgh.2015.12.004

Bardeesy, N., Sinha, M., Hezel, A. F., Signoretti, S., Hathaway, N. A., Sharpless, N. E., Loda, M., Carrasco, D. R., & DePinho, R. A. (2002). Loss of the Lkb1 tumour suppressor provokes intestinal polyposis but resistance to transformation. *Nature*, *419*(6903), 162–167. https://doi.org/10.1038/nature01045

Brügger, M. D., & Basler, K. (2023). The diverse nature of intestinal fibroblasts in development, homeostasis, and disease. *Trends in Cell Biology*, *33*(10), 834–849. https://doi.org/10.1016/j.tcb.2023.03.007

Brügger, M. D., Valenta, T., Fazilaty, H., Hausmann, G., & Basler, K. (2020). Distinct populations of crypt-associated fibroblasts act as signaling hubs to control colon homeostasis. *PLoS Biology*, *18*(12), e3001032. https://doi.org/10.1371/journal.pbio.3001032

Buechler, M. B., Pradhan, R. N., Krishnamurty, A. T., Cox, C., Calviello, A. K., Wang, A. W., Yang, Y. A., Tam, L., Caothien, R., Roose-Girma, M., Modrusan, Z., Arron, J. R., Bourgon, R., Müller, S., & Turley, S. J. (2021). Cross-tissue organization of the fibroblast lineage. *Nature*, *593*(7860), 575–579. https://doi.org/10.1038/s41586-021-03549-5

Compton, S. E., Kitchen-Goosen, S. M., DeCamp, L. M., Lau, K. H., Mabvakure, B., Vos, M., Williams, K. S., Wong, K.-K., Shi, X., Rothbart, S. B., Krawczyk, C. M., & Jones, R. G. (2023). LKB1 controls inflammatory potential through CRTC2-dependent histone acetylation. *Molecular Cell*. https://doi.org/10.1016/j.molcel.2023.04.017

Cotton, J. L., Dang, K., Hu, L., Sun, Y., Singh, A., Rajurkar, M. S., Li, Q., Wu, X., & Mao, J. (2022). PTEN and LKB1 are differentially required in Gli1-expressing mesenchymal cells to suppress gastrointestinal polyposis. *Cell Reports*, *40*(3), 111125. https://doi.org/10.1016/j.celrep.2022.111125

Deng, M., Guerrero-Juarez, C. F., Sheng, X., Xu, J., Wu, X., Yao, K., Li, M., Yang, X., Li, G., Xiao, J., Liu, X., Wu, K., Ren, F., Nie, Q., Plikus, M. V., Yu, Z., & Lv, C. (2022). Lepr+ mesenchymal cells sense diet to modulate intestinal stem/progenitor cells via Leptin-Igf1 axis. *Cell Research*. https://doi.org/10.1038/s41422-022-00643-9

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* , *29*(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Domènech-Moreno, E., Brandt, A., Lemmetyinen, T. T., Wartiovaara, L., Mäkelä, T. P., & Ollila, S. (2023). Tellu: an object detector algorithm for automatic classification of intestinal organoids. *Disease Models & Mechanisms*. https://doi.org/10.1242/dmm.049756

Eliazer, S., Muncie, J. M., Christensen, J., Sun, X., D'Urso, R. S., Weaver, V. M., & Brack, A. S. (2019). Wnt4 from the Niche Controls the Mechano-Properties and Quiescent State of Muscle Stem Cells.

Cell Stem Cell, *25*(5), 654–665.e4. https://doi.org/10.1016/j.stem.2019.08.007

Elsheikh, W., Flannigan, K. L., McKnight, W., Ferraz, J. G. P., & Wallace, J. L. (2012). Dextran sulfate sodium induces pan-gastroenteritis in rodents: implications for studies of colitis. *Journal of Physiology and Pharmacology: An Official Journal of the Polish Physiological Society*, *63*(5), 463–469. https://www.ncbi.nlm.nih.gov/pubmed/23211300

Finak, G., McDavid, A., Yajima, M., Deng, J., Gersuk, V., Shalek, A. K., Slichter, C. K., Miller, H. W., McElrath, M. J., Prlic, M., Linsley, P. S., & Gottardo, R. (2015). MAST: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biology*, *16*, 278. https://doi.org/10.1186/s13059-015-0844-5

Fung, K. Y., Louis, C., Metcalfe, R. D., Kosasih, C. C., Wicks, I. P., Griffin, M. D. W., & Putoczki, T. L. (2022). Emerging roles for IL-11 in inflammatory diseases. *Cytokine*, *149*, 155750. https://doi.org/10.1016/j.cyto.2021.155750

Giardiello, F. M., & Trimbath, J. D. (2006). Peutz-Jeghers syndrome and management recommendations. *Clinical Gastroenterology and Hepatology: The Official Clinical Practice Journal of the American Gastroenterological Association*, *4*(4), 408–415. https://doi.org/10.1016/j.cgh.2005.11.005

Haber, A. L., Biton, M., Rogel, N., Herbst, R. H., Shekhar, K., Smillie, C., Burgin, G., Delorey, T. M., Howitt, M. R., Katz, Y., Tirosh, I., Beyaz, S., Dionne, D., Zhang, M., Raychowdhury, R., Garrett, W. S., Rozenblatt-Rosen, O., Shi, H. N., Yilmaz, O., … Regev, A. (2017). A single-cell survey of the small intestinal epithelium. *Nature*, nature24489. https://doi.org/10.1038/nature24489

Heichler, C., Scheibe, K., Schmied, A., Geppert, C. I., Schmid, B., Wirtz, S., Thoma, O.-M., Kramer, V., Waldner, M. J., Büttner, C., Farin, H. F., Pešić, M., Knieling, F., Merkel, S., Grüneboom, A., Gunzer, M., Grützmann, R., Rose-John, S., Koralov, S. B., … Neufert, C. (2020). STAT3 activation through IL-6/IL-11 in cancer-associated fibroblasts promotes colorectal tumour development and correlates with poor prognosis. *Gut*, *69*(7), 1269–1282. https://doi.org/10.1136/gutjnl-2019-319200

Hemminki, A., Markie, D., Tomlinson, I., Avizienyte, E., Roth, S., Loukola, A., Bignell, G., Warren, W., Aminoff, M., Höglund, P., Järvinen, H., Kristo, P., Pelin, K., Ridanpää, M., Salovaara, R., Toro, T., Bodmer, W., Olschwang, S., Olsen, A. S., … Aaltonen, L. A. (1998). A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature*, *391*(6663), 184–187. https://doi.org/10.1038/34432

James, K. R., Elmentaite, R., Teichmann, S. A., & Hold, G. L. (2022). Redefining intestinal immunity with single-cell transcriptomics. *Mucosal Immunology*, *15*(4), 531–541. https://doi.org/10.1038/s41385-021-00470-y

Jardé, T., Chan, W. H., Rossello, F. J., Kaur Kahlon, T., Theocharous, M., Kurian Arackal, T., Flores, T., Giraud, M., Richards, E., Chan, E., Kerr, G., Engel, R. M., Prasko, M., Donoghue, J. F., Abe, S.-I., Phesse, T. J., Nefzger, C. M., McMurrick, P. J., Powell, D. R., … Abud, H. E. (2020). Mesenchymal Niche-Derived Neuregulin-1 Drives Intestinal Stem Cell Proliferation and Regeneration of Damaged Epithelium. *Cell Stem Cell*. https://doi.org/10.1016/j.stem.2020.06.021

Jasso, G. J., Jaiswal, A., Varma, M., Laszewski, T., Grauel, A., Omar, A., Silva, N., Dranoff, G., Porter, J. A., Mansfield, K., Cremasco, V., Regev, A., Xavier, R. J., & Graham, D. B. (2022). Colon stroma mediates an inflammation-driven fibroblastic response controlling matrix remodeling and healing. *PLoS Biology*, *20*(1), e3001532. https://doi.org/10.1371/journal.pbio.3001532

Jelsig, A. M., Wullum, L., Kuhlmann, T. P., Ousager, L. B., Burisch, J., & Karstensen, J. G. (2023). Risk of Cancer and Mortality in Peutz-Jeghers Syndrome and Juvenile Polyposis Syndrome-A Nationwide Cohort Study With Matched Controls. *Gastroenterology*. https://doi.org/10.1053/j.gastro.2023.08.040

Jin, S., Guerrero-Juarez, C. F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung, P., Plikus, M. V., & Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat. *Nature Communications*, *12*(1), 1–20. https://doi.org/10.1038/s41467-021-21246-9

Joosten, S. P. J., Zeilstra, J., van Andel, H., Mijnals, R. C., Zaunbrecher, J., Duivenvoorden, A. A. M., van de Wetering, M., Clevers, H., Spaargaren, M., & Pals, S. T. (2017). MET Signaling Mediates Intestinal Crypt-Villus Development, Regeneration, and Adenoma Formation and Is Promoted by Stem Cell CD44 Isoforms. *Gastroenterology*, *153*(4), 1040–1053.e4.

https://doi.org/10.1053/j.gastro.2017.07.008

Kalucka, J., de Rooij, L. P. M. H., Goveia, J., Rohlenova, K., Dumas, S. J., Meta, E., Conchinha, N. V., Taverna, F., Teuwen, L.-A., Veys, K., García-Caballero, M., Khan, S., Geldhof, V., Sokol, L., Chen, R., Treps, L., Borri, M., de Zeeuw, P., Dubois, C., … Carmeliet, P. (2020). Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell*, *180*(4), 764–779.e20. https://doi.org/10.1016/j.cell.2020.01.015

Katajisto, P., Vaahtomeri, K., Ekman, N., Ventelä, E., Ristimäki, A., Bardeesy, N., Feil, R., DePinho, R. A., & Mäkelä, T. P. (2008). LKB1 signaling in mesenchymal cells required for suppression of gastrointestinal polyposis. *Nature Genetics*, *40*(4), 455–459. https://doi.org/10.1038/ng.98

Kim, J.-E., Fei, L., Yin, W.-C., Coquenlorge, S., Rao-Bhatia, A., Zhang, X., Shi, S. S. W., Lee, J. H., Hahn, N. A., Rizvi, W., Kim, K.-H., Sung, H.-K., Hui, C.-C., Guo, G., & Kim, T.-H. (2020). Single cell and genetic analyses reveal conserved populations and signaling mechanisms of gastrointestinal stromal niches. *Nature Communications*, *11*(1), 334. https://doi.org/10.1038/s41467-019-14058-5

Kinchen, J., Chen, H. H., Parikh, K., Antanaviciute, A., Jagielowicz, M., Fawkner-Corbett, D., Ashley, N., Cubitt, L., Mellado-Gomez, E., Attar, M., Sharma, E., Wills, Q., Bowden, R., Richter, F. C., Ahern, D., Puri, K. D., Henault, J., Gervais, F., Koohy, H., & Simmons, A. (2018). Structural Remodeling of the Human Colonic Mesenchyme in Inflammatory Bowel Disease. *Cell*, *175*(2), 372–386.e17. https://doi.org/10.1016/j.cell.2018.08.067

Klefstrom, J., Arighi, E., Littlewood, T., Jäättelä, M., Saksela, E., Evan, G. I., & Alitalo, K. (1997). Induction of TNF-sensitive cellular phenotype by c-Myc involves p53 and impaired NF-kappaB activation. *The EMBO Journal*, *16*(24), 7382–7392. https://doi.org/10.1093/emboj/16.24.7382

Kraiczy, J., McCarthy, N., Malagola, E., Tie, G., Madha, S., Boffelli, D., Wagner, D. E., Wang, T. C., & Shivdasani, R. A. (2023). Graded BMP signaling within intestinal crypt architecture directs self-organization of the Wnt-secreting stem cell niche. *Cell Stem Cell*, *30*(4), 433–449.e8. https://doi.org/10.1016/j.stem.2023.03.004

Langlands, A. J., Carroll, T. D., Chen, Y., & Näthke, I. (2018). Chir99021 and Valproic acid reduce the proliferative advantage of Apc mutant cells. *Cell Death & Disease*, *9*(3), 255. https://doi.org/10.1038/s41419-017-0199-9

Lemmetyinen, T. T., Viitala, E. W., Wartiovaara, L., Kaprio, T., Hagström, J., Haglund, C., Katajisto, P., Wang, T. C., Domènech-Moreno, E., & Ollila, S. (2023). Fibroblast-derived EGF ligand neuregulin 1 induces fetal-like reprogramming of the intestinal epithelium without supporting tumorigenic growth. *Disease Models & Mechanisms*, *16*(4). https://doi.org/10.1242/dmm.049692

Leushacke, M., Ng, A., Galle, J., Loeffler, M., & Barker, N. (2013). Lgr5(+) gastric stem cells divide symmetrically to effect epithelial homeostasis in the pylorus. *Cell Reports*, *5*(2), 349–356. https://doi.org/10.1016/j.celrep.2013.09.025

Lim, W.-W., Ng, B., Widjaja, A., Xie, C., Su, L., Ko, N., Lim, S.-Y., Kwek, X.-Y., Lim, S., Cook, S. A., & Schafer, S. (2020). Transgenic interleukin 11 expression causes cross-tissue fibro-inflammation and an inflammatory bowel phenotype in mice. *PloS One*, *15*(1), e0227505. https://doi.org/10.1371/journal.pone.0227505

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* , *25*(4), 402–408. https://doi.org/10.1006/meth.2001.1262

Macosko, E. Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas, A. R., Kamitaki, N., Martersteck, E. M., Trombetta, J. J., Weitz, D. A., Sanes, J. R., Shalek, A. K., Regev, A., & McCarroll, S. A. (2015). Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell*, *161*(5), 1202–1214. https://doi.org/10.1016/j.cell.2015.05.002

Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter, R. D., Hawrylycz, M. J., Jones, A. R., Lein, E. S., & Zeng, H. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nature Neuroscience*, *13*(1), 133–140. https://doi.org/10.1038/nn.2467

Manieri, E., Tie, G., Seruggia, D., Madha, S., Maglieri, A., Huang, K., Fujiwara, Y., Zhang, K., Orkin, S.

H., He, R., McCarthy, N., & Shivdasani, R. A. (2023). Defining the structure, signals, and cellular elements of the gastric mesenchymal niche. In *bioRxiv* (p. 2023.02.11.527728). https://doi.org/10.1101/2023.02.11.527728

McCarthy, N., Kraiczy, J., & Shivdasani, R. A. (2020). Cellular and molecular architecture of the intestinal stem cell niche. *Nature Cell Biology*, *22*(9), 1033–1041. https://doi.org/10.1038/s41556-020-0567-z

McCarthy, N., Manieri, E., Storm, E. E., Saadatpour, A., Luoma, A. M., Kapoor, V. N., Madha, S., Gaynor, L. T., Cox, C., Keerthivasan, S., Wucherpfennig, K., Yuan, G.-C., de Sauvage, F. J., Turley, S. J., & Shivdasani, R. A. (2020). Distinct Mesenchymal Cell Populations Generate the Essential Intestinal BMP Signaling Gradient. *Cell Stem Cell*. https://doi.org/10.1016/j.stem.2020.01.008

Melissari, M.-T., Henriques, A., Tzaferis, C., Prados, A., Sarris, M. E., Chalkidi, N., Mavroeidi, D., Chouvardas, P., Grammenoudi, S., Kollias, G., & Koliaraki, V. (2021). Col6a1+/CD201+ mesenchymal cells regulate intestinal morphogenesis and homeostasis. *Cellular and Molecular Life Sciences: CMLS*, *79*(1), 1. https://doi.org/10.1007/s00018-021-04071-7

Muhl, L., Genové, G., Leptidis, S., Liu, J., He, L., Mocci, G., Sun, Y., Gustafsson, S., Buyandelger, B., Chivukula, I. V., Segerstolpe, Å., Raschperger, E., Hansson, E. M., Björkegren, J. L. M., Peng, X.-R., Vanlandewijck, M., Lendahl, U., & Betsholtz, C. (2020). Single-cell analysis uncovers fibroblast heterogeneity and criteria for fibroblast and mural cell identification and discrimination. *Nature Communications*, *11*(1), 3953. https://doi.org/10.1038/s41467-020-17740-1

Ng, B., Dong, J., D'Agostino, G., Viswanathan, S., Widjaja, A. A., Lim, W.-W., Ko, N. S. J., Tan, J., Chothani, S. P., Huang, B., Xie, C., Pua, C. J., Chacko, A.-M., Guimarães-Camboa, N., Evans, S. M., Byrne, A. J., Maher, T. M., Liang, J., Jiang, D., … Cook, S. A. (2019). Interleukin-11 is a therapeutic target in idiopathic pulmonary fibrosis. *Science Translational Medicine*, *11*(511). https://doi.org/10.1126/scitranslmed.aaw1237

Nishina, T., Deguchi, Y., Ohshima, D., Takeda, W., Ohtsuka, M., Shichino, S., Ueha, S., Yamazaki, S., Kawauchi, M., Nakamura, E., Nishiyama, C., Kojima, Y., Adachi-Akahane, S., Hasegawa, M., Nakayama, M., Oshima, M., Yagita, H., Shibuya, K., Mikami, T., … Nakano, H. (2021). Interleukin-11-expressing fibroblasts have a unique gene signature correlated with poor prognosis of colorectal cancer. *Nature Communications*, *12*(1), 2281. https://doi.org/10.1038/s41467-021-22450-3

Nusse, Y. M., Savage, A. K., Marangoni, P., Rosendahl-Huber, A. K. M., Landman, T. A., de Sauvage, F. J., Locksley, R. M., & Klein, O. D. (2018). Parasitic helminths induce fetal-like reversion in the intestinal stem cell niche. *Nature*, *559*(7712), 109–113. https://doi.org/10.1038/s41586-018-0257-1

Ollila, S., Domènech-Moreno, E., Laajanen, K., Wong, I. P., Tripathi, S., Pentinmikko, N., Gao, Y., Yan, Y., Niemelä, E. H., Wang, T. C., Viollet, B., Leone, G., Katajisto, P., Vaahtomeri, K., & Mäkelä, T. P. (2018). Stromal Lkb1 deficiency leads to gastrointestinal tumorigenesis involving the IL-11-JAK/STAT3 pathway. *The Journal of Clinical Investigation*, *128*(1), 402–414. https://doi.org/10.1172/JCI93597

Pærregaard, S. I., Wulff, L., Schussek, S., Niss, K., Mörbe, U., Jendholm, J., Wendland, K., Andrusaite, A. T., Brulois, K. F., Nibbs, R. J. B., Sitnik, K., Mowat, A. M., Butcher, E. C., Brunak, S., & Agace, W. W. (2023). The small and large intestine contain related mesenchymal subsets that derive from embryonic Gli1+ precursors. *Nature Communications*, *14*(1), 2307. https://doi.org/10.1038/s41467-023-37952-5

Pasztoi, M., & Ohnmacht, C. (2022). Tissue Niches Formed by Intestinal Mesenchymal Stromal Cells in Mucosal Homeostasis and Immunity. *International Journal of Molecular Sciences*, *23*(9). https://doi.org/10.3390/ijms23095181

Poffenberger, M. C., Metcalfe-Roach, A., Aguilar, E., Chen, J., Hsu, B. E., Wong, A. H., Johnson, R. M., Flynn, B., Samborska, B., Ma, E. H., -P. Gravel, S., Tonelli, L., Devorkin, L., Kim, P., Hall, A., Izreig, S., Loginicheva, E., Beauchemin, N., Siegel, P. M., … Jones, R. G. (2018). LKB1 deficiency in T cells promotes the development of gastrointestinal polyposis. *Science*, *361*(6400), 406–411. https://doi.org/10.1126/science.aan3975

Putoczki, T. L., Thiem, S., Loving, A., Busuttil, R. A., Wilson, N. J., Ziegler, P. K., Nguyen, P. M.,

H., He, R., McCarthy, N., & Shivdasani, R. A. (2023). Defining the structure, signals, and cellular elements of the gastric mesenchymal niche. In *bioRxiv* (p. 2023.02.11.527728). https://doi.org/10.1101/2023.02.11.527728

McCarthy, N., Kraiczy, J., & Shivdasani, R. A. (2020). Cellular and molecular architecture of the intestinal stem cell niche. *Nature Cell Biology*, *22*(9), 1033–1041. https://doi.org/10.1038/s41556-020-0567-z

McCarthy, N., Manieri, E., Storm, E. E., Saadatpour, A., Luoma, A. M., Kapoor, V. N., Madha, S., Gaynor, L. T., Cox, C., Keerthivasan, S., Wucherpfennig, K., Yuan, G.-C., de Sauvage, F. J., Turley, S. J., & Shivdasani, R. A. (2020). Distinct Mesenchymal Cell Populations Generate the Essential Intestinal BMP Signaling Gradient. *Cell Stem Cell*. https://doi.org/10.1016/j.stem.2020.01.008

Melissari, M.-T., Henriques, A., Tzaferis, C., Prados, A., Sarris, M. E., Chalkidi, N., Mavroeidi, D., Chouvardas, P., Grammenoudi, S., Kollias, G., & Koliaraki, V. (2021). Col6a1+/CD201+ mesenchymal cells regulate intestinal morphogenesis and homeostasis. *Cellular and Molecular Life Sciences: CMLS*, *79*(1), 1. https://doi.org/10.1007/s00018-021-04071-7

Muhl, L., Genové, G., Leptidis, S., Liu, J., He, L., Mocci, G., Sun, Y., Gustafsson, S., Buyandelger, B., Chivukula, I. V., Segerstolpe, Å., Raschperger, E., Hansson, E. M., Björkegren, J. L. M., Peng, X.-R., Vanlandewijck, M., Lendahl, U., & Betsholtz, C. (2020). Single-cell analysis uncovers fibroblast heterogeneity and criteria for fibroblast and mural cell identification and discrimination. *Nature Communications*, *11*(1), 3953. https://doi.org/10.1038/s41467-020-17740-1

Ng, B., Dong, J., D'Agostino, G., Viswanathan, S., Widjaja, A. A., Lim, W.-W., Ko, N. S. J., Tan, J., Chothani, S. P., Huang, B., Xie, C., Pua, C. J., Chacko, A.-M., Guimarães-Camboa, N., Evans, S. M., Byrne, A. J., Maher, T. M., Liang, J., Jiang, D., … Cook, S. A. (2019). Interleukin-11 is a therapeutic target in idiopathic pulmonary fibrosis. *Science Translational Medicine*, *11*(511). https://doi.org/10.1126/scitranslmed.aaw1237

Nishina, T., Deguchi, Y., Ohshima, D., Takeda, W., Ohtsuka, M., Shichino, S., Ueha, S., Yamazaki, S., Kawauchi, M., Nakamura, E., Nishiyama, C., Kojima, Y., Adachi-Akahane, S., Hasegawa, M., Nakayama, M., Oshima, M., Yagita, H., Shibuya, K., Mikami, T., … Nakano, H. (2021). Interleukin-11-expressing fibroblasts have a unique gene signature correlated with poor prognosis of colorectal cancer. *Nature Communications*, *12*(1), 2281. https://doi.org/10.1038/s41467-021-22450-3

Nusse, Y. M., Savage, A. K., Marangoni, P., Rosendahl-Huber, A. K. M., Landman, T. A., de Sauvage, F. J., Locksley, R. M., & Klein, O. D. (2018). Parasitic helminths induce fetal-like reversion in the intestinal stem cell niche. *Nature*, *559*(7712), 109–113. https://doi.org/10.1038/s41586-018-0257-1

Ollila, S., Domènech-Moreno, E., Laajanen, K., Wong, I. P., Tripathi, S., Pentinmikko, N., Gao, Y., Yan, Y., Niemelä, E. H., Wang, T. C., Viollet, B., Leone, G., Katajisto, P., Vaahtomeri, K., & Mäkelä, T. P. (2018). Stromal Lkb1 deficiency leads to gastrointestinal tumorigenesis involving the IL-11-JAK/STAT3 pathway. *The Journal of Clinical Investigation*, *128*(1), 402–414. https://doi.org/10.1172/JCI93597

Pærregaard, S. I., Wulff, L., Schussek, S., Niss, K., Mörbe, U., Jendholm, J., Wendland, K., Andrusaite, A. T., Brulois, K. F., Nibbs, R. J. B., Sitnik, K., Mowat, A. M., Butcher, E. C., Brunak, S., & Agace, W. W. (2023). The small and large intestine contain related mesenchymal subsets that derive from embryonic Gli1+ precursors. *Nature Communications*, *14*(1), 2307. https://doi.org/10.1038/s41467-023-37952-5

Pasztoi, M., & Ohnmacht, C. (2022). Tissue Niches Formed by Intestinal Mesenchymal Stromal Cells in Mucosal Homeostasis and Immunity. *International Journal of Molecular Sciences*, *23*(9). https://doi.org/10.3390/ijms23095181

Poffenberger, M. C., Metcalfe-Roach, A., Aguilar, E., Chen, J., Hsu, B. E., Wong, A. H., Johnson, R. M., Flynn, B., Samborska, B., Ma, E. H., -P. Gravel, S., Tonelli, L., Devorkin, L., Kim, P., Hall, A., Izreig, S., Loginicheva, E., Beauchemin, N., Siegel, P. M., … Jones, R. G. (2018). LKB1 deficiency in T cells promotes the development of gastrointestinal polyposis. *Science*, *361*(6400), 406–411. https://doi.org/10.1126/science.aan3975

Putoczki, T. L., Thiem, S., Loving, A., Busuttil, R. A., Wilson, N. J., Ziegler, P. K., Nguyen, P. M.,

Preaudet, A., Farid, R., Edwards, K. M., Boglev, Y., Luwor, R. B., Jarnicki, A., Horst, D., Boussioutas, A., Heath, J. K., Sieber, O. M., Pleines, I., Kile, B. T., … Ernst, M. (2013). Interleukin-11 Is the Dominant IL-6 Family Cytokine during Gastrointestinal Tumorigenesis and Can Be Targeted Therapeutically. *Cancer Cell*, *24*(2), 257–271. https://doi.org/10.1016/j.ccr.2013.06.017

Robinson, J., Lai, C., Martin, A., Nye, E., Tomlinson, I., & Silver, A. (2009). Oral rapamycin reduces tumour burden and vascularization in Lkb1(+/-) mice. *The Journal of Pathology*, *219*(1), 35–40. https://doi.org/10.1002/path.2562

Rossi, D. J., Ylikorkala, A., Korsisaari, N., Salovaara, R., Luukko, K., Launonen, V., Henkemeyer, M., Ristimaki, A., Aaltonen, L. A., & Makela, T. P. (2002). Induction of cyclooxygenase-2 in a mouse model of Peutz-Jeghers polyposis. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(19), 12327–12332. https://doi.org/10.1073/pnas.192301399

Roulis, M., Kaklamanos, A., Schernthanner, M., Bielecki, P., Zhao, J., Kaffe, E., Frommelt, L.-S., Qu, R., Knapp, M. S., Henriques, A., Chalkidi, N., Koliaraki, V., Jiao, J., Brewer, J. R., Bacher, M., Blackburn, H. N., Zhao, X., Breyer, R. M., Aidinis, V., … Flavell, R. A. (2020). Paracrine orchestration of intestinal tumorigenesis by a mesenchymal niche. *Nature*, *580*(7804), 524–529. https://doi.org/10.1038/s41586-020-2166-3

Sackett, S. D., Fulmer, J. T., Friedman, J. R., & Kaestner, K. H. (2007). Foxl1-Cre BAC transgenic mice: a new tool for gene ablation in the gastrointestinal mesenchyme. *Genesis* , *45*(8), 518–522. https://doi.org/10.1002/dvg.20315

Sanjana, N. E., Shalem, O., & Zhang, F. (2014). Improved vectors and genome-wide libraries for CRISPR screening. *Nature Methods*, *11*(8), 783–784. https://doi.org/10.1038/nmeth.3047

Sato, T., Vries, R. G., Snippert, H. J., van de Wetering, M., Barker, N., Stange, D. E., van Es, J. H., Abo, A., Kujala, P., Peters, P. J., & Clevers, H. (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature*, *459*(7244), 262–265. https://doi.org/10.1038/nature07935

Schafer, S., Viswanathan, S., Widjaja, A. A., Lim, W.-W., Moreno-Moral, A., DeLaughter, D. M., Ng, B., Patone, G., Chow, K., Khin, E., Tan, J., Chothani, S. P., Ye, L., Rackham, O. J. L., Ko, N. S. J., Sahib, N. E., Pua, C. J., Zhen, N. T. G., Xie, C., … Cook, S. A. (2017). IL-11 is a crucial determinant of cardiovascular fibrosis. *Nature*, *552*(7683), 110–115. https://doi.org/10.1038/nature24676

Shalem, O., Sanjana, N. E., Hartenian, E., Shi, X., Scott, D. A., Mikkelson, T., Heckl, D., Ebert, B. L., Root, D. E., Doench, J. G., & Zhang, F. (2014). Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science*, *343*(6166), 84–87. https://doi.org/10.1126/science.1247005

Shoshkes-Carmel, M., Wang, Y. J., Wangensteen, K. J., Tóth, B., Kondo, A., Massasa, E. E., Itzkovitz, S., & Kaestner, K. H. (2018). Subepithelial telocytes are an important source of Wnts that supports intestinal crypts. *Nature*, *557*(7704), 242–246. https://doi.org/10.1038/s41586-018-0084-4

Smillie, C. S., Biton, M., Ordovas-Montanes, J., Sullivan, K. M., Burgin, G., Graham, D. B., Herbst, R. H., Rogel, N., Slyper, M., Waldman, J., Sud, M., Andrews, E., Velonias, G., Haber, A. L., Jagadeesh, K., Vickovic, S., Yao, J., Stevens, C., Dionne, D., … Regev, A. (2019). Intra- and Inter-cellular Rewiring of the Human Colon during Ulcerative Colitis. *Cell*, *178*(3), 714–730.e22. https://doi.org/10.1016/j.cell.2019.06.029

Tan, S. H., Swathi, Y., Tan, S., Goh, J., Seishima, R., Murakami, K., Oshima, M., Tsuji, T., Phuah, P., Tan, L. T., Wong, E., Fatehullah, A., Sheng, T., Ho, S. W. T., Grabsch, H. I., Srivastava, S., Teh, M., Denil, S. L. I. J., Mustafah, S., … Barker, N. (2020). AQP5 enriches for stem cells and cancer origins in the distal stomach. *Nature*, *578*(7795), 437–443. https://doi.org/10.1038/s41586-020-1973-x

Thomson, C. A., van de Pavert, S. A., Stakenborg, M., Labeeuw, E., Matteoli, G., Mowat, A. M., & Nibbs, R. J. B. (2018). Expression of the Atypical Chemokine Receptor ACKR4 Identifies a Novel Population of Intestinal Submucosal Fibroblasts That Preferentially Expresses Endothelial Cell Regulators. *Journal of Immunology* , *201*(1), 215–229. https://doi.org/10.4049/jimmunol.1700967

Udd, L., Katajisto, P., Kyyrönen, M., Ristimäki, A. P., & Mäkelä, T. P. (2010). Impaired gastric gland differentiation in Peutz-Jeghers syndrome. *The American Journal of Pathology*, *176*(5), 2467–2476. https://doi.org/10.2353/ajpath.2010.090519

Udd, L., Katajisto, P., Rossi, D. J., Lepistö, A., Lahesmaa, A.-M., Ylikorkala, A., Järvinen, H. J., Ristimäki, A. P., & Mäkelä, T. P. (2004). Suppression of Peutz-Jeghers polyposis by inhibition of cyclooxygenase-2. *Gastroenterology*, *127*(4), 1030–1037. https://doi.org/10.1053/j.gastro.2004.07.059

Wei, K., Nguyen, H. N., & Brenner, M. B. (2021). Fibroblast pathology in inflammatory diseases. *The Journal of Clinical Investigation*, *131*(20). https://doi.org/10.1172/JCI149538

Widjaja, A. A., Chothani, S., Viswanathan, S., Goh, J. W. T., Lim, W.-W., & Cook, S. A. (2022). IL11 Stimulates IL33 Expression and Proinflammatory Fibroblast Activation across Tissues. *International Journal of Molecular Sciences*, *23*(16). https://doi.org/10.3390/ijms23168900

Widjaja, A. A., Viswanathan, S., Wei Ting, J. G., Tan, J., Shekeran, S. G., Carling, D., Lim, W.-W., & Cook, S. A. (2022). IL11 stimulates ERK/P90RSK to inhibit LKB1/AMPK and activate mTOR initiating a mesenchymal program in stromal, epithelial, and cancer cells. *iScience*, *25*(8), 104806. https://doi.org/10.1016/j.isci.2022.104806

Xiao, S., & Zhou, L. (2020). Gastric Stem Cells: Physiological and Pathological Perspectives. *Frontiers in Cell and Developmental Biology*, *8*, 571536. https://doi.org/10.3389/fcell.2020.571536

Yu, K., Xu, J., Liu, Z., Sosic, D., Shao, J., Olson, E. N., Towler, D. A., & Ornitz, D. M. (2003). Conditional inactivation of FGF receptor 2 reveals an essential role for FGF signaling in the regulation of osteoblast function and bone growth. *Development* , *130*(13), 3063–3074. https://doi.org/10.1242/dev.00491

Zhang, Q., Pan, Y., Ji, J., Xu, Y., Zhang, Q., & Qin, L. (2021). Roles and action mechanisms of WNT4 in cell differentiation and human diseases: a review. *Cell Death Discovery*, *7*(1), 287. https://doi.org/10.1038/s41420-021-00668-w

Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S. B., Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L., Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W., Hindson, C. M., … Bielas, J. H. (2017). Massively parallel digital transcriptional profiling of single cells. *Nature Communications*, *8*, 14049. https://doi.org/10.1038/ncomms14049

Figure 1. Enrichment of ST2+ fibroblasts in PJS polyps

Twist2-Cre;Lkb1$^{flox/+}$, gastric polyp and adjacent normal antrum (control) scRNAseq

Figure 1. ST2-expressing fibroblasts are enriched in PJS polyps. (A) UMAP clustering of all cells identified in the whole scRNAseq dataset including normal stomach and gastric polyp tissue. **(B)** UMAP clustering of the fibroblast subpopulations from A. **(C)** Violin plots of markers shared by all fibroblasts as well as established markers for crypt bottom and crypt top fibroblast subsets. **(D)** Violin plots of top five markers of the identified fibroblast subpopulations. **(E)** Ratio of fibroblast subtype representation in tumors as compared to control gastric mucosa. *, p<0.05, Chi-square test. **(F)** Split violin plots indicating expression of *Il1rl1* and *Il11* expression across all fibroblast populations split by sample (grey: control, red: tumor). **(G)** Immunofluorescent staining of ST2 in tumors (T) and adjacent control mucosa (C) of *Lkb1*$^{+/-}$ or *Twist2Cre;Lkb1*$^{fl/+}$ mice.

Figure 2. CTF-specific Loss of LKB1 induces polyposis and expression of ST2-CTF markers

Figure 2. CTF-specific Loss of LKB1 induces polyposis and expression of ST2-CTF markers. (A)
Survival curve of the Foxl1-Cre;Lkb1fl/fl mice (n=4). **(B)** Representative images of a stomach from *Foxl1-Cre;Lkb1fl/fl* mouse at 17 weeks. **(C)** Representative histology images of tumor and adjacent mucosa stained for PDGFRA (Fibroblasts) and RFP (Foxl1 lineage) from *Foxl1-Cre;Lkb1fl/fl;R26R-TdTomato* mouse at 17 weeks of age. **(D)** FACS strategy to isolate Foxl1-lineage fibroblasts from young mice of indicated genotypes. **(E)** Expression of Lkb1, Il1rl1 and Il11 mRNA in the indicated fibroblast fractions. *, p>0.05, one-tailed T-test.

Figure 3. LKB1 loss recapitulates the inflammatory ST2-CTF transcriptome

Figure 3. LKB1 loss recapitulates the inflammatory ST2-CTF transcriptome. (A) Feature plot of the ST2+ CTF gene signature in homeostatic (Control) and inflamed (DSS) mouse colon fibroblasts. Data from(Kinchen et al., 2018). **(B)** GSEA of ST2+ CTF gene set against LPS treated primary fibroblasts and DSS treated mouse colon tissue lysates. **(C-D)**. UMAP **(C)** and feature plot **(D)** showing the ST2+ CTF gene signature in human colon stroma from ulcerative colitis patients ((Smillie et al., 2019). **(E)** Outline of the acute *in vitro* deletion of *Lkb1* using adenovirus in primary intestinal fibroblasts derived from *Lkb1^{fl/fl}* mice. **(F)** mRNA expression of *Lkb1, Il1rl1 and Il11* primary intestinal fibroblasts treated with adenovirus (Ctrl, Adeno-GFP; Cre, Adeno-Cre). Data from four independent experiments with two technical replicates each is shown. Average of the control replicates per experiment is used for normalization. **(G)** GSEA of the ST2+ gene signature against the ranked transcriptome from *Lkb1^{fl/fl}* primary intestinal fibroblasts. **(H)** GSEA of LKB1 loss induced gene signature against LPS treated primary fibroblasts and DSS treated mouse colon tissue lysates. **(I)** Experimental setup and for assessing LPS-induced gene expression differences in WT and Lkb1 KO primary intestinal fibroblasts. Lkb1 mRNA levels as compared to Adeno-GFP in control conditions are shown. **(J)** mRNA expression of *Lkb1* in control and LPS-treated primary intestinal fibroblasts transduced with adenovirus (Ctrl, Adeno-GFP; Cre, Adeno-Cre). Data from four independent experiments with two technical replicates each is shown. **(K)** Expression levels of indicated genes and genotypes in control and LPS induced conditions. Fold change to WT Control cells is plotted. A representative of three independent experiments is shown. *, p<0.05, Two-tailed T-test. Box plots indicate mean and standard deviation.

Figure 4. Inflammation exacerbates the LKB1 loss mediated tumor-promoting phenotype

Figure 4. Inflammation exacerbates the LKB1 loss mediated tumor-promoting phenotype. (A) Experimental setup of the in vitro co-culture experiments using primary fibroblasts from *Lkb1fl/fl* mice and organoids isolated from wild-type mice. **(B)** Representative images of organoids after 24h of co-culture with Lkb1 WT (AdGFP) or Lkb1 KO (AdCre) fibroblasts with or without LPS. **(C)** Quantification of the ratio of spheroids vs organoids per well in each condition. The mean relative area per well, standard deviation and individual data points are shown (a.u, arbitrary units). **(D)** Experimental outline of the chronic inflammation induced in *Twist2-Cre;Lkb1fl/+* mice. **(E)** Kaplan-Meier curve depicting survival of *Twist2-Cre;Lkb1fl/+* mice in control conditions (black line, n=5) and upon DSS treatment (yellow line, n=6) Log-rank test used. **(F)** Tumor burden (left) and tumor size distribution (right) of the *Twist2-Cre;Lkb1fl/+* mice in control conditions (control, n=9) and upon DSS treatment (n=11).

Figure 5. ST2-CTFs present a strong and unique cell-cell signaling pattern

Figure 5: ST2-CTFs possess a strong and unique cell-cell signaling pattern. (A) 2D representation of overall incoming and outgoing interactions in the tumor gastric mucosa. **(B).** Heatmap of the cell communication pattern analysis of outgoing and receiving signals between clusters. **(C)** Heatmap of the cell communication outgoing signal pattern 5 found in B specific of ST2$^+$ fibroblast.

Figure 6. Inhibition of IL-11 signaling inhibits the ST2+ transcriptome and reduces PJS polyposis

Figure 6. Inhibition of IL-11 signaling inhibits the ST2+ transcriptome and reduces PJS polyposis. (A) IL11 cell-cell signaling network originates from ST2-CTFs and targets fibroblast populations in PJS tumors. **(B)** GSEA of Lkb1 and ST2+ fibroblast gene sets (Supplementary Table 1) against the transcriptome of IL11-EGFP+ fibroblasts isolated from inflammation-associated colon cancer mouse model (Nishina et al., 2021). **(C)** Experimental setup (left) and gene expression of indicated genes in fibroblasts transduced with Cas9 and non-targeting (NT) gRNA or gRNA against IL11 (right). gRNA1 is shown, see also Supplementary Figure 6. **(D)** Schematic diagram illustrating antibody treatment in *Lkb1*[+/-] mice. 30-week-old *Lkb1*[+/-] mice with established polyposis were treated with either X203 or IgG isotype antibodies (20 mg/kg/week IP) up to 50 weeks (n=11 males, 12 females per antibody group). **(E)** Representative images of stomachs of 50-week-old *Lkb1*[+/-] mice treated with either antibody compared to 50-week-old wildtype littermates and 30-week-old untreated *Lkb1*[+/-] mice. Red arrows indicate gastric polyps. **(F-H)** Average polyp number **(F)**, size **(G)**, and total polyp burden **(H)** of *Lkb1*[+/-] mice treated with either antibody compared to untreated 30-week-old *Lkb1*[+/-] mice (n=7 males, 12 females).*, $p<0.05$, **, $p<0.01$, ***, $p<0.001$, ****, $p<0.0001$, two-way ANOVA with Sidak post-hoc test. ns, not significant.

Supplementary Figure 1. Single-cell sequencing of PJS type gastric tumors

*Twist2-Cre;Lkb1^flox/+^*, gastric polyp and adjacent normal antrum (control) scRNAseq

Supplementary Figure 1. Single-cell sequencing of PJS type gastric tumors. (A) UMAP clustering of all cells identified in the scRNAseq dataset from normal antrum (Control) and antral polyp. **(B)** Expression of tdTomato expression across the identified clusters. **(C)** Violin plots of the indicated marker genes as example markers used to identify the cell types.

Supplementary Figure 2. Inflammation leads to enhanced PJS polyposis

Supplementary Figure 2. Inflammation leads to enhanced PJS polyposis. (A) Outline of the inflammation-induced (AOM-DSS) model of colon tumorigenesis. AOM injection was performed at 20 weeks of age. **(B)** Gastric tumor burden in control or AOM-DSS treated Twist2-Cre;Lkb1fl/+ mice at 30 weeks of age. Ctrl, no treatment (n=9); AOM-DSS, n=8. **(C)** Colon tumor burden in control (Lkb1fl/+, n=7) and Mutant (Mut, Twist2-Cre;Lkb1fl/+, n=8) mice. **, p<0.01, Two-tailed T-test. n.s., not significant (p>0.05).

Supplementary Figure 3. Decreased epithelial Pepsinogen C in the stroma-specific PJS model.

Supplementary Figure 3. Decreased epithelial Pepsinogen C in the stroma-specific PJS model. (A) UMAP clustering of all epithelial cells identified in the scRNAseq dataset. **(B)** Markers used to classify epithelial cell types. **(C)** UMAP clustering of the epithelial cell clusters in control antrum and polyp. Equal number of cells was plotted. **(D)** Volcano plot of differentially expressed genes between tumor cells and control antrum epithelial cells. Red dots depict significant genes with a log2 fold change above 1. **(E)** Immunohistochemical staining of Pepsinogen C in *Twist2-Cre;Lkb1fl/+* antral polyp sections and adjacent normal antrum.

Supplementary Figure 4.Analysis of immune cells and endothelial cells in gastric polyps

Supplementary Figure 4. Analysis of immune cells and endothelial cells in gastric polyps. (A) UMAP clustering of all immune cells identified in the scRNAseq dataset. **(B)** Markers used to classify the immune cell types. **(C)** UMAP clustering of the immune cell clusters in control antrum and polyp. Equal number of cells was plotted. **(D)** UMAP clustering of all endothelial cells identified in control antrum and polyp. **(E)** Markers used to classify the endothelial cell types.

Supplementary Figure 5. ST2-fibroblast derived signaling pathways affecting epithelium and fibroblasts

Supplementary Figure 5. ST2-CTF-derived ligand-receptor signaling pairs. (A) Total number of interactions within fibroblasts and epithelial cells in scRNAseq of control and polyp datasets. **(B)** Main ligand and receptor contributors for each family of ligands. **(C)** The most active ligand-receptor signaling interactions from ST2-CTFs to epithelial cell clusters. **(D)** The most active ligand-receptor signaling interactions from ST2-CTFs to other fibroblast clusters. **(C-D)** The line color represents the signaling source, and the line width the strength of the signaling. Only fibroblast and epithelial clusters were included in the circo plots.

Supplementary Figure 6. Generation of IL-11 deficient fibroblasts

Supplementary Figure 6. Generation of IL-11 deficient fibroblasts. (A) Sanger sequencing of puromycin selected cell pools transduced with non-targeting (NT) gRNA or two independent gRNAs targeting Il11. The gRNA binding sites are highlighted, note mixed sequence and deletions (red arrows) in the targeted sequence. **(B)** Experimental setup (left) and gene expression of indicated genes in fibroblasts transduced with Cas9 and non-targeting (NT) gRNA or gRNA against IL11 (right). gRNA2 is shown, see also Figure 6C.