A disengaging property of medial accumbens shell dopamine.

Milan D. Valyear[1,2], Noémie M-L. Eustachon[1,2], Irina Alymova[3], Bryanna N. Tremblay[1], Nastasia M. Mitrikeski[4], *Jonathan P. Britt[1,2,4]

[1] Department of Psychology, McGill University, Montréal QC, Canada
[2] Center for Studies in Behavioral Neurobiology, Concordia University, Montréal, QC, Canada
[3] Cell and Systems Biology, University of Toronto, Toronto ON, Canada
[4] Integrative Program in Neuroscience, McGill University, Montréal QC, Canada

*Corresponding Author:
Department of Psychology
McGill University
1205 Ave. Dr. Penfield,
Montréal, QC, H3A 1B1
Canada
Phone: +1 514.298.3625
Email: jonathan.britt@mcgill.ca

Abstract: Electrical stimulation of the medial forebrain bundle vigorously reinforces self-stimulation behaviour, yet rodents perform operant responses to terminate this stimulation. The accumbens shell emerged as a substrate subserving the reinforcing properties of electrical medial forebrain bundle stimulation, whereas disengaging properties were attributed to incidentally recruited substrates near the electrode. Here, we examine whether there are dissociable reinforcing and disengaging properties of medial accumbens shell dopamine and probe the substrates underlying these properties. Using a temporally delimited self-stimulation procedure, transgenic DAT-Cre mice expressing channelrhodopsin-II in ventral tegmental area dopamine neurons were trained to hold-down a lever to engage, and then release the lever to disengage, optogenetic stimulation of dopaminergic inputs to the medial accumbens shell through an implanted optic fiber. The cumulative and mean duration of hold-downs show divergent frequency responses identifying dissociable reinforcing and disengaging properties of medial accumbens shell dopamine. At higher stimulation frequencies the cumulative duration of hold-downs grows, whereas the mean duration of hold-downs wanes. Dopamine agonists reduced the cumulative duration of self-stimulation hold-downs, but only a D1 agonist produced this reduction through decreases in the mean duration of hold-downs, which were lengthened with a D2 antagonist. Thus, reinforcing and disengaging properties of electrical medial forebrain stimulation may arise from the downstream activation of dopamine receptors, uncovering a disengaging property of medial accumbens shell dopamine.

Key words: Dopamine, Ventral Striatum, Addiction, Optogenetics, Valence, Reward

Significance Statement: Dopamine is thought to promote behaviour by acting as a reinforcer or error signal. Here, we show that mice vigorously self-stimulate dopamine inputs to the medial accumbens shell but control the duration of duration of these stimulations and prefer them to be brief. This disengaging property of medial accumbens shell dopamine depends on downstream neurotransmission at dopamine type 1 and 2 receptors. Thus, a single dopaminergic substrate, inputs to the medial accumbens shell, reinforces and disengages self-stimulation behaviour, highlighting the complexity and regional specificity of striatal dopamine function.

1. Introduction.

The discovery that rodents [1] and people [2,3] performed operant responses to electrically stimulate the medial forebrain bundle ignited speculation about the psychological consequences of this stimulation: Did it substitute for natural goals, positive memories, or utility – generate hunger, reward, or drive [4]? A parallel, search for the neural substrates recruited by electrical stimulation of the medial forebrain bundle spurred [5], ever complicated by the swath of cells and transgressing fibers characterizing this large bundle [5]. It was soon realized that the quality of this stimulation was not owed to a single psychological process. Rodents [6,7] and people [2] would perform responses to terminate ongoing stimulation. The reinforcing and disengaging properties of electrical medial forebrain bundle stimulation were presumed to arise from distinct neural substrates indiscriminately recruited by the electrode [8]. An imprecise but indelible link, between substrate and psychology was established.

A description of the neural substrates underlying the reinforcing properties of electrical brain stimulation focused on an area of the ventral striatum called the nucleus accumbens. Diverse patterns of electrical medial forebrain bundle stimulation similarly activated ventral tegmental area dopamine neurons [9] and their release of dopamine in the accumbens, but not dorsal striatum [10,11]. Within the accumbens, the medial shell subregion aligned with the reinforcing properties of medial forebrain bundle stimulation because dopamine levels here closely tracked maximally reinforcing stimulation parameters [12]. Further, pharmacologically elevating dopamine in the medial accumbens shell, but not adjacent striatal subregions, amplified the reinforcing efficacy of medial forebrain bundle stimulation [13]. Thus, the reinforcing properties of medial forebrain bundle stimulation are partly attributable to elevations in medial accumbens shell dopamine occurring through transsynaptic activation of ventral tegmental area dopamine neurons [14–16].

Optogenetics [17] and transgenic rodents [18,19] confirmed that ventral tegmental area dopamine neurons robustly supported self-stimulation, particularly at high stimulation frequencies, suggesting that these neurons underpinned the reinforcing properties of electrical medial forebrain bundle stimulation [18,20]. Once again however, questions were raised about diverse psychological processes engaged by optogenetic stimulation of ventral tegmental area dopamine neurons. When mice could choose between operant responses that delivered brief or prolonged bursts of high frequency optogenetic stimulation of ventral tegmental area dopamine neurons, brief bursts were preferred [21]. The preferential activation of ventral tegmental area dopamine neurons for brief durations is akin to behavioural responses that terminate ongoing electrical brain stimulation. Differently, however, the substrate serving to reinforce and disengage optogenetic self-stimulation of ventral tegmental area dopamine neurons is unequivocal.

Self-stimulation studies highlight the capacity for dopamine to reinforce behaviour, but the dominant view of dopamine as an error signal nuances the interpretation of such studies. The firing of ventral tegmental area dopamine neurons, and consequent striatal release of dopamine, is thought to encode a prediction error [22,23]. Operant responses for stimulation of dopamine neurons creates an error such that the receipt of stimulation is *better than expected*, and the response is repeated with increasing vigour. Attempts have been made to disentangle the hypotheses that optogenetic stimulation of ventral tegmental area dopamine neurons acts as a

reinforcer or error signal [15,24], but these procedures overlook separable roles of dopamine neurons with distinct striatal targets [25–27]. Notably, dopamine activity in the medial accumbens shell is inconsistent with error coding [28], modulated by the value of natural taste stimuli [29], and directly supports self-stimulation [27], highlighting a role for this substrate as a reinforcer.

A key feature of how animals engage with natural reinforcers is the capacity to control the duration of behaviour. Mammals [30], including wild mice [31], tightly control the duration of bouts of consummatory behaviour around taste [32]. In lab studies mice increase the duration of their bouts of licking a solution as it becomes sweeter and decrease this duration as the solution becomes bitter [33]. Changes in bout duration are often divorced from changes in bout frequency which relate more to the postingestive consequences of consumption, like satiety or malaise [34]. That is, the taste of a positively valenced sweet solution can support long bouts of licking but reinforce few bouts in total. It is unclear if, given the requisite control, mice would modulate the duration for which a non-sensory neural substrate was stimulated, much like they do with sensory taste receptors activated by natural reinforcers, like food.

We developed a temporally constrained self-stimulation procedure wherein mice are trained to hold-down a lever to engage, and then release the lever to disengage, optogenetic stimulation of dopaminergic inputs to the medial accumbens shell. As such, two principal measures describe self-stimulation: the cumulative and mean duration of lever hold-downs. The cumulative duration of hold-downs reflects the reinforcing properties of stimulation, arising either from changes in the mean duration or number of hold-downs, or both. Differently, the mean duration of hold-downs reflects a quality of the stimulation that arises as it is being experienced, allowing mice to disengage stimulation by releasing the lever. We use the temporally constrained self-stimulation procedure to dissociate reinforcing and disengaging properties of medial accumbens shell dopamine by examining whether the mean and cumulative durations of hold-downs diverge across stimulation frequences. Then, by modulating neurotransmission at dopamine type-1 (D1) and type-2 (D2) receptors with pharmacology we determined whether distinct downstream substrates impact the cumulative or mean duration of self-stimulation hold-downs. Selective recruitment of a genetically defined and input-specific substrate clarifies speculation about separable or overlapping substrates subserving the reinforcing and disengaging properties of electrical brain stimulation which we argue both arise at the level of dopamine inputs to the medial accumbens shell.

2. Materials and methods.

2.1. Subjects.

Subjects were twelve experimentally naïve DAT-Cre (7F, 5M) C57BL/6 mice (25-30 g, 8 weeks old; bred in-house) that were single or pair-housed in standard shoe-box cages (31 cm x 16.5 cm x 12 cm) under a reversed 12-hour light/dark cycle (Lights ON 9 AM, OFF 9 PM) with unrestricted access to standard chow and water. All procedures were conducted in the dark phase and in accordance with the Canadian Council on Animal Care and the McGill Animal Care Committee.

2.2. Surgery.

DAT-Cre transgenic mice were anesthetized with ketamine (Ventoquinol, 100mg/kg) and xylazine (Bayer, 10mg/kg) before being secured in a stereotaxic apparatus. The viral construct AAV5-DIO-CaMKIIa-hChR2(H134R)-eYFP (Neurophotonics, 5.0×10^{12} GC/ml) was microinfused using a Nanoject II injector (Drummond Scientific, 3-000-203-G/X, 10 μm tip) in the ventral tegmental area (10° angle, AP 1.14 mm, ML ±0.35 mm, DV -3.75 to -2.75 mm; 5 min infusion, 10 min diffusion). In the same surgery optic fibers (200 μm) were implanted bilaterally in the medial accumbens shell (10° angle, AP 1.5 mm, ML ±1.4 mm, DV -4.32 mm). Mice were left to recover for 2-3 months in their home-cage allowing sufficient viral expression in dopaminergic terminals.

2.3. Behavioral apparatus.

Behavioral training occurred in six conditioning chambers (ENV-307W-CT; Med-Associates Inc.), enclosed in sound-attenuating, fan-ventilated (ENV-025F) melamine cubicles (42.5 x 62.5 x 42 cm). Each chamber had a stainless-steel bar floor, paneled aluminum sidewalls, and a clear polycarbonate rear wall, ceiling, and front door. The upper left wall of the chamber featured a central white house-light (ENV-215M). The right wall contained two retractable levers (ENV-312-3) located on each side of a central fluid-port (ENV-303LP2-3). All experimental events were controlled and recorded using Med PC-V software.

2.4. Temporally constrained self-stimulation.

Mice received 40 self-stimulation training sessions (36 min) consisting of trials (30s; 48 trials/session) initiated and terminated with the extension and retraction, respectively, of active and inactive levers. In each trial, the active lever was pseudorandomly assigned to deliver pulsed laser stimulation (5 ms; 473 nm; 10 mW) at 2.5, 10, or 40 Hz such that one third of all trials represented each frequency. Trials occurred around a variable time inter-trial interval that was drawn from an exponential distribution from 2-18 s. When mice performed hold-downs by depressing the active lever, pulsed laser-light was delivered through an implanted optic fiber to the medial accumbens shell, which continued so long as the lever remained depressed and terminated when the lever was released. Inactive lever hold-downs had no consequence.

After training, the contributions of neurotransmission at D1 and D2 receptors to self-stimulation of medial accumbal shell dopamine inputs was tested. Prior to each test session mice received an intraperitoneal injection of Raclopride (D2 antagonist, 0.03 mg/kg), Quinpirole (D2 agonist, 0.125 mg/kg), SCH23390 (D1 antagonist, 0.02 mg/kg), SKF38393 (D1 agonist, 0.25 mg/kg) or saline. Three sessions for each treatment (15 total) occurred according to a Latin-square design. Retraining sessions without injections intervened each test session to mitigate potential treatment carryover effects.

2.5. Histology.

Mice were deeply anesthetized with sodium pentobarbital (Euthanyl™, 270 mg/kg) and perfused with a 4% paraformaldehyde 0.1 M phosphate-buffered saline solution. Brains were extracted and cryoprotected in a 4% paraformaldehyde 30% sucrose solution for 1-2 days before being frozen at -80°C. Brain tissue was sliced into 50 μm coronal sections on a Lecia™ cryostat, thaw-mounted on microscope slides, and cover-slipped with a MOWIOL + DAPI solution, then imaged using a

Leica^{TM} epifluorescent microscope to identify viral expression in the ventral tegmental area and optical fiber tracts in the medial accumbens shell.

2.6. Data and Statistical Analysis.
MED-PC V software controlled and recorded the timing of all experimental events. The cumulative and mean duration, number and latency of hold-downs were analyzed using repeated measures analysis of variance (RMANOVA) in Prism^{TM} v9.5.1 or SPSS v27. The Kolmogorov-Smirnov test was used to compare distributions. Conditioning sessions were blocked into groups of five (training) or three (test) sessions to ensure that enough hold-downs occurred to calculate mean hold-down duration. All analyses were conducted on raw data, although difference scores are shown to decompose significant interactions. Two mice did not perform a hold-down at one set of frequency-treatment conditions during the test phase and their data was excluded from test analyses. Post-hoc t-tests were Bonferroni-corrected (one family) and multiplicity-adjusted [35]. All analyses used α=.05.

3. Results.

3.1. Temporally constrained self-stimulation of dopamine inputs to the medial accumbens shell.
DAT-Cre mice were trained (**Fig 1a**) to hold-down an active lever to optogenetically stimulate dopamine inputs to the medial accumbens shell (**Fig 1b-e**). Across training blocks of five sessions [Session, $F_{(7, 77)}$=14.392, p<.001], mice increased the cumulative time spent holding down the active, but not inactive lever [Lever x Session, $F_{(7, 77)}$=14.332, p<.001; Lever, $F_{(1, 11)}$=34.470, p<.001], particularly so for higher frequency trials [Lever x Frequency x Session, $F_{(14, 154)}$=2.323, p=.006; Frequency x Session, $F_{(14, 154)}$=1.011, p=.445; Lever x Frequency, $F_{(2, 22)}$=4.470, p=.023; Frequency, $F_{(2, 22)}$=6.042, p=.008; **Fig 1f**]. A selective and frequency-dependent increase in the cumulative time spent holding down the active lever is consistent with dopamine in the medial accumbens shell reinforcing self-stimulation. While mice learned to lengthen the mean duration of active, but not inactive, hold-downs across sessions [Lever x Session, $F_{(7, 77)}$=4.509, p<.001; Lever, $F_{(1, 11)}$=37.598, p<.001; Session, $F_{(7, 77)}$=4.125, p=.001] the trajectory of this lengthening was not frequency-dependent [Lever x Frequency x Session, $F_{(14, 154)}$=.301, p=.993; Frequency x Session, $F_{(14, 154)}$=.671, p=.8; Lever x Frequency, $F_{(2, 22)}$=2.390, p=.115; Frequency, $F_{(2, 22)}$=1.802, p=.188; **Fig 1g**]. Consistently, a frequency-dependent increase in the cumulative, but not mean, duration of active hold-downs was accompanied by a frequency-dependent increase in the number of active, but not inactive, hold-downs across sessions [Lever x Session, $F_{(7, 77)}$=8.541, p<.001; Lever, $F_{(1, 11)}$=20.890, p=.001; Session, $F_{(7, 77)}$=8.126, p<.001; Lever x Frequency x Session, $F_{(14, 154)}$=2.679, p=.002; Frequency x Session, $F_{(14, 154)}$=1.088, p=.373; Lever x Frequency, $F_{(2, 22)}$=5.376, p=.013; Frequency, $F_{(2, 22)}$=3.450, p=.05; **Fig 1h**]. The cumulative duration and number of hold-downs were averaged across trials to compare against an intuitive trial duration of 30 s, however, the total number of hold-downs showed similar trajectories (**Fig S1a**).

3.2. Cumulative and mean duration of self-stimulation hold-downs diverge across frequencies.
During the last block of training sessions, the cumulative and mean duration of active hold-downs showed divergent frequency responses that were absent early in training (**Fig S2**). The cumulative duration of active, relative to inactive, hold-downs increased [Lever x Frequency, $F_{(2, 22)}$=3.594,

$p=.044$; Lever, $F_{(1, 22)}=72.57$, $p<.0001$; Frequency, $F_{(2, 22)}=3.789$, $p=.0385$; **Fig 2a**] whereas the mean duration of active, relative to inactive, hold-downs decreased across stimulation frequencies [Lever x Frequency, $F_{(2, 22)}=5.3$, $p=.0132$; Lever, $F_{(1, 22)}=17.54$, $p=.0015$; Frequency, $F_{(2, 22)}=.9435$, $p=.045$; **Fig 2b**]. Specifically, the cumulative active hold-down duration during 40 Hz trials was longer than during 10 [$t_{(11)}=3.367$, $p=.0083$] or 2.5 Hz [$t_{(11)}=3.79$, $p=.003$] trials, which did not differ [$t_{(11)}=.4339$, $p>.999$]. The increase in cumulative active hold-down duration across stimulation frequencies is consistent with reinforcing properties of medial accumbens shell dopamine. Differently, the mean duration of active hold-downs was briefer when the active lever was armed with 40 relative to 10 [$t_{(11)}=3.022$, $p=.0188$] or 2.5 Hz stimulation [$t_{(11)}=3.061$, $p=.0172$], which did not differ [$t_{(11)}=.03845$, $p>.999$]. The decrease in mean lever hold-down duration at higher stimulation frequencies is consistent with a disengaging property of medial accumbens shell dopamine. Although the requisite interaction was not observed [Lever x Frequency, $F_{(2, 22)}=2.607$, $p=.0963$; Frequency, $F_{(2, 22)}=1.928$, $p=.1692$; **Fig 2c**] more active than inactive lever hold-downs were performed [Lever, $F_{(1, 22)}=19.40$, $p<.0011$] and a corrected comparison revealed that more active hold-downs occurred during 40 than 2.5 Hz trials [$t_{(11)}=2.993$, $p=.0201$]. Neither the cumulative duration [all $t_{(11)}<.5358$, $p>.999$; **Fig 2d**], mean duration [all $t_{(11)}<.9620$, $p>.999$; **Fig 2e**], or number [all $t_{(11)}<.1787$, $p>.999$; **Fig 2f**] of inactive hold-downs varied across stimulation frequencies.

Relative frequency distributions (**Fig 2g-i**) revealed that long duration active hold-downs for high frequency stimulation seldom occurred as mice performed sub ~1 s hold-downs more often when the active lever was armed with 40 relative to 10 or 2.5 Hz stimulation [40 vs 2.5 Hz, D=.1517, $p<.001$; 40 vs 10 Hz, D=.1785, $p<.001$; 10 vs 2.5 Hz, D=.0482, $p=.2695$; **Fig 2j**].

3.3. Modulation of reinforcing and disengaging properties of medial accumbens shell dopamine.
At test, the impact of modulating neurotransmission at dopamine receptors on the temporally constrained self-stimulation of dopamine inputs to the medial accumbens shell was examined. The D1 agonist SKF38393 (.25 mg/kg) and D2 agonist quinpirole (.125 mg/kg) were used to augment neurotransmission, whereas the D1 antagonist SCH23390 (.02 mg/kg) and the D2 antagonist raclopride (.03 mg/kg) were used to blunt neurotransmission at dopamine receptors. Doses of dopamine receptor agonists and antagonists were chosen because they were extensively shown to not affect motor behaviour in mice [36–45].

During test sessions, the cumulative duration of active hold-downs varied across stimulation frequencies [Frequency, $F_{(2, 18)}=11.04$, $p=.0007$] and treatments [Treatment, $F_{(4, 36)}=73.3$, $p<.00001$], but these factors did not interact [Frequency x Treatment, $F_{(8, 72)}=1.302$, $p=.2564$; **Fig 3a,d**]. As during training, the cumulative duration of active hold-downs was elevated during 40 relative to 10 Hz trials [$t_{(9)}=3.313$, $p=.0116$] or 2.5 Hz [$t_{(9)}=4.543$, $p=.0008$; **Fig S3a,c**]. Augmenting neurotransmission with the D1 agonist SKF38393 [$t_{(9)}=12.94$, $p<.0001$] reduced the cumulative duration of active hold-downs whereas blunting neurotransmission with the D1 antagonist SCH23390 had no effect [$t_{(9)}=1.193$, $p>.999$; **Fig 3a**]. Augmenting neurotransmission with the D2 agonist quinpirole reduced the cumulative duration of active hold-downs [$t_{(9)}=6.013$, $p<.00001$] whereas blunting neurotransmission with the D2 antagonist raclopride had no effect [$t_{(9)}=2.118$,

p=.4118; **Fig 3d**]. Thus, the reinforcing properties of medial accumbens shell dopamine are reduced by augmenting neurotransmission at D1 and D2 receptors.

The mean duration of active hold-downs was modulated by treatment [Treatment, $F_{(4, 36)}$=11.33, *p*<.00001] in a frequency-dependent manner [Frequency x Treatment, $F_{(8, 72)}$=4.948, *p*=.0002; Frequency, $F_{(2, 18)}$=5.721, *p*<.0119; **Fig 3b,e**]. As during training, the mean duration of active hold-downs was briefer at 40 than 2.5 Hz [$t_{(9)}$=3.359, *p*=.0105; **Fig S3b,d**]. Augmenting neurotransmission with the D1 receptor agonist SKF38393 reduced the mean duration of active hold-downs relative to the D1 antagonist SCH23390 [2.5 Hz, $t_{(9)}$=6.396, *p*<.0001; 10 Hz, $t_{(9)}$=5.744, *p*<.0001; 40 Hz, $t_{(9)}$=4.815, *p*=.0002] and saline [2.5 Hz, $t_{(9)}$=6.22, *p*<.0001; 10 Hz, $t_{(9)}$=6.041, *p*<.0001; 40 Hz, $t_{(9)}$=5.125, *p*<.0001]; this effect was most pronounced at lower stimulation frequencies (**Fig 3b**). Blunting neurotransmission with the D2 receptor antagonist raclopride lengthened the mean duration of active hold-downs relative to the D2 agonist quinpirole [2.5 Hz, $t_{(9)}$=9.342, *p*<.0001; 10 Hz, $t_{(9)}$=9.480, *p*<.0001; 40 Hz, $t_{(9)}$=4.254, *p*<.0019] and saline [2.5 Hz, $t_{(9)}$=11.01, *p*<.0001; 10 Hz, $t_{(9)}$=12.32, *p*<.0001; 40 Hz, $t_{(9)}$=6.826, *p*=.0002]; this effect was most pronounced at lower stimulation frequencies (**Fig 3e**). Neither the D1 antagonist SCH23390 [2.5 Hz, $t_{(9)}$=.5261, *p*>.999; 10 Hz, $t_{(9)}$=.2973, *p*>.999; 40 Hz, $t_{(9)}$=.3093, *p*>.999] nor the D2 agonist quinpirole [2.5 Hz, $t_{(9)}$=1.670, *p*>.999; 10 Hz, $t_{(9)}$=2.841, *p*=.1754; 40 Hz, $t_{(9)}$=2.573, *p*=.3646] affected the mean duration of active hold-downs relative to saline. Thus, the disengaging property of medial accumbens shell dopamine was modulated by augmenting neurotransmission at D1 receptors and blunting neurotransmission at D2 receptors.

The number of active hold-downs did not vary across stimulation frequencies [Frequency, $F_{(2, 18)}$=1.773, *p*=.1983; Frequency x Treatment, $F_{(8, 72)}$=1.471, *p*=.1829] but was affected by treatment [Treatment, $F_{(4, 36)}$=3.84, *p*=.0106]. After statistical correction, the reduction in the number of active lever hold-downs relative to saline remained only marginally significant for the D1 agonist SKF28393 [$t_{(9)}$=2.954, *p*=.055] and other treatments were similar to saline [SCH23390, $t_{(9)}$=.4919, *p*>.9909; Quinpirole, $t_{(9)}$=1.050, *p*>.9099; Raclopride, $t_{(9)}$=1.809, *p*=.7873; **Fig 3c,f**]. The cumulative duration, mean duration, and number of inactive hold-downs were largely unaffected by stimulation frequency and treatments (**see Fig S4**).

3.4. The first self-stimulation hold-down.

By design, mice could not predetermine the stimulation frequency on offer thus the latency to initiate the first active hold-down was similar across frequencies [Frequency, $F_{(2, 18)}$=.04887, *p*=.9524; Frequency x Treatment, $F_{(8, 72)}$=.4328, *p*=.8976; **Fig S5a,c**]. Mice were slower to initiate the first active hold-down following treatment [Treatment, $F_{(4, 36)}$=40.94, *p*<.001] with the D1 agonist SKF38393 [$t_{(9)}$=10.18, *p*<.001; **Fig 4a**] and D2 agonist quinpirole [$t_{(9)}$=4.395, *p*<.001; **Fig 4c**]; the D1 antagonist SCH23390 [$t_{(9)}$=.129, *p*>.999] and D2 antagonist raclopride [$t_{(9)}$=.3429, *p*>.999] had no effect.

Random assignment of stimulation frequency at trial onset ensured that the duration of the first active lever hold-down occurred in response to stimulation as it was being experienced. The mean duration of the first active hold-down was briefer at 40 than 2.5 Hz [$t_{(9)}$=3.039, *p*=.0212; **Fig S5b,d**] and modulated by the D1 agonist SKF38393, the D2 agonist quinpirole, and the D2 antagonist

raclopride in a frequency-dependent manner [Frequency x Treatment, $F_{(8, 72)}$=2.611, p=.0144; **Fig 4b,d**]. The D1 agonist SKF38393 shortened the duration of the first active hold-down relative to SCH23390 [2.5 Hz, $t_{(9)}$=6.844, p<.0001; 10 Hz, $t_{(9)}$=6.215, p<.0001; 40 Hz, $t_{(9)}$=5.705 p<.0001] and saline [2.5 Hz, $t_{(9)}$=8.205, p<.0001; 10 Hz, $t_{(9)}$=6.869, p<.0001; 40 Hz, $t_{(9)}$=6.625 p<.0001; **Fig 4b**]. The D2 agonist quinpirole shortened [2.5 Hz, $t_{(9)}$=3.856, p=.0074; 10 Hz, $t_{(9)}$=3.614, p=0.0167; 40 Hz, $t_{(9)}$=4.137, p=.0028] whereas the D2 antagonist raclopride lengthened the duration of the first active hold-down [2.5 Hz, $t_{(9)}$=8.154, p<.0001; 10 Hz, $t_{(9)}$=9.841, p<.0001; 40 Hz, $t_{(9)}$=5.077, p<.0001] relative to saline (**Fig 4d**).

While the latency to initiate the first active hold-down was unaffected by stimulation frequency, the latency to initiate the second hold-down depended on the stimulation received on the first hold-down [Frequency, $F_{(2, 18)}$=4.018, p=.0361; **Fig S6a,c**]. Mice were quicker to perform a second active hold-down on 40 relative to 2.5 Hz trials [$t_{(9)}$=2.794, p=.036], confirming that high-frequency stimulation was strongly reinforcing and demonstrating that mice could update the vigour of their self-stimulation behavior after a single experience. Dopamine receptor agonists and antagonists did not impact the latency to initiate the second active hold-down confirming that treatment effects slowing initiation of the first active hold-down were thereafter overcome [Treatment, $F_{(4, 36)}$=.98, p=.4306; Frequency x Treatment, $F_{(8, 72)}$=.8784, p=.5389; **Fig S6b,d**]. The latency to initiate the second active hold-down on 10 Hz trials did not differ from 40 [$t_{(9)}$=.9798, p>.999] or 2.5 [$t_{(9)}$=1.814, p=.2592] Hz trials.

4. Discussion

The medial accumbens shell emerged as a substrate subserving the reinforcing properties of electrical medial forebrain bundle stimulation [12,13] and early studies attributed diverse psychological consequences of this stimulation to the indiscriminate recruitment of neural substrates near the electrode. Here, by allowing mice to control the duration for which they optogenetically self-stimulate a genetically defined and input-specific substrate we describe reinforcing and disengaging properties of medial accumbens shell dopamine. The principal evidence for this claim is that the cumulative and mean duration for which mice hold-down a lever to optogenetically stimulate dopaminergic inputs to the medial accumbens shell show divergent frequency responses. At higher stimulation frequencies the cumulative duration of hold-downs grows, consistent with more reinforcement, whereas the mean duration of hold-downs wanes, consistent with disengagement. Using pharmacology, we begin identifying substrates underpinning the reinforcing and disengaging properties of medial accumbens shell dopamine. Dopamine agonists reduced the cumulative duration of self-stimulation hold-downs, but only a D1 agonist produced this reduction through decreases in the mean duration of hold-downs, which were lengthened by a D2 antagonist. Thus, reinforcing and disengaging properties of electrical medial forebrain stimulation may arise from the downstream activation of dopamine receptors, uncovering a disengaging property of medial accumbens shell dopamine.

Across training, mice increased the cumulative duration, mean duration, and number of active relative to inactive lever hold-downs, consistent with the acquisition of temporally constrained self-stimulation. One previous study reported that mice could not learn to hold-down a lever to self-stimulate dopaminergic cell bodies in the substantia nigra [46], which may relate to an

impoverished role of nigral, relative to ventral tegmental area dopamine neurons, in reinforcing behaviour [27]. Additionally, nigral projection targets in the dorsal striatum [26] differ from ventral tegmental area dopamine inputs to the medial accumbens shell and likely engage circuitries that serve different functions. Also, the previous attempt to use hold-down duration as a self-stimulation operant may have been unsuccessful because of the high frequency of stimulation used, 50 Hz, which generally supported sub half-second hold-downs, consistent with the disengaging property of high frequency dopamine activity discovered in the current work. The effects of stimulation frequency detected here are smaller than in procedures where rodents control only the number, not duration, of trains of dopamine stimulation. Importantly, most previous work directly stimulated dopamine cell bodies in the ventral tegmental area [21,47], rather than accumbal inputs and few studies varied stimulation frequency. Nevertheless, mice demonstrate sophisticated and considerable (~25%) control over the duration with which dopamine inputs to the medial accumbens shell are stimulated.

We attribute decreases in mean hold-down duration during high frequency stimulation trials to the emergence of a disengaging property of medial accumbens shell dopamine. This interpretation is incongruent with a paradigm in contemporary neuroscience describing a relationship between striatal cells and motivated behaviour. In the striatum, the activity of neurons expressing D1 receptors is thought to encode positive valence and encourage behaviour, whereas neurons expressing D2 receptors oppose this influence, encoding negative valence when active and discouraging behaviour [48,49]. Thus, input from the ventral tegmental area to the striatum, including the accumbens, is thought to reinforce behaviour because dopamine elevates the activity of D1 neurons and mutes the activity of D2 neurons, serving as a strong positive valence signal. Certainly, we observed increases in the cumulative time spent self-stimulating dopamine inputs to the medial accumbens shell, consistent with reinforcing and positively valenced properties of driving this substrate. Differently, the mean duration for which mice hold-down a lever to stimulate dopaminergic inputs to the medial accumbens shell wanes at high stimulation frequencies, consistent with the emergence of a disengaging property of this substrate. It is unclear whether this disengaging property is inherently valenced, although mice also reduce the duration with which they interact with negatively valenced taste stimuli [33,34]. Ultimately, it is difficult to reconcile a disengaging property of medial accumbens shell dopamine with conventional views of striatum-wide dopamine as a reinforcer or error signal. Rather, the psychological consequences of striatal dopamine likely reflect subregion specificity and the modulation of downstream neuron activity, which is convergently shaped by corticostriatal inputs expressing dopamine receptors [50].

The disengaging property of medial accumbens shell dopamine may arise from the co-release of glutamate by ventral tegmental area dopamine neurons that innervate the accumbens shell. When mice could nose-poke at five concurrently available apertures armed with 40 Hz optogenetic stimulation of different durations (i.e., 1, 5, 20, 40 s), ventral tegmental area dopamine cell bodies predominantly supported responses for 5 s of stimulation. Differently, a subpopulation of ventral tegmental area dopamine neurons that co-release glutamate predominantly supported responses for 1 s of stimulation [21] and mice would perform an operant response to terminate noncontingent stimulation of these inputs to the medial accumbens shell [51]. Thus, glutamate co-release from dopaminergic inputs to the medial accumbens shell may contribute to the current behavior, where high-frequency stimulation supported the briefest mean duration of self-

stimulation hold-downs; however, the modulation of self-stimulation hold-downs with dopamine receptor agonists and antagonists, and a recent report that dopamine release absent glutamate is aversive [52], disfavours this possibility.

Dopamine receptor agonists and antagonists produced dissociable effects on the cumulative and mean durations for which mice self-stimulated dopaminergic inputs to the medial accumbens shell. While dopamine receptor agonists reduced the cumulative duration of self-stimulation hold-downs, only the D1 agonist SKF38393 did so by reducing the mean duration of hold-downs which was unaffected by the D2 agonist quinpirole. Augmenting neurotransmission at D2 receptors with quinpirole diminished the reinforcing properties of medial accumbens shell dopamine, which likely manifested through subtle coincident reductions in the mean duration and number of hold-downs. Although dopamine antagonists did not affect the cumulative duration of self-stimulation hold-downs, blunting neurotransmission at D2 receptors with raclopride lengthened self-stimulation hold-downs, as if to mitigate the disengaging property of medial accumbens shell dopamine. Differently, augmenting neurotransmission at D1 receptors with SKF38393 shortened self-stimulation hold-downs, consistent with amplification of the disengaging property of medial accumbens shell dopamine.

In the temporally constrained self-stimulation procedure, the first active hold-down of a trial is a clear measure of the disengaging properties of medial accumbens dopamine because mice cannot predetermine the available stimulation frequency. The first active hold-down on high-frequency trials was briefer than on low-frequency trials – this result is only explained by mice disengaging the lever to terminate stimulation in response to some quality of the stimulation arising in real time. Despite experiencing a disengaging property of medial accumbens shell dopamine on the first high-frequency hold-down, mice were quicker to initiate a second hold-down than during lower frequency trials. A single experience of high-frequency dopamine activity in the medial accumbens shell is sufficient to reinforce, and disengage, self-stimulation behaviour more strongly than lower frequency stimulation of the same substrate. Thus, the effects of stimulation frequency and pharmacological treatments on mean hold-down duration were observed on the first hold-down. Additionally, the D2 agonist quinpirole reduced the duration of the first active hold-down, which explains the companion reduction in cumulative active hold-down duration. Neurotransmission at D2 receptors thus partially underpins a disengaging property of medial accumbens shell dopamine, however, this effect diminished on subsequent hold-downs. The modulation of mean, or first, active hold-down duration with changes in neurotransmission at D2 receptors suggests that these neurons may play a principal role in determining whether, or when, animals disengage behavior.

5. **Conclusion**

Dopamine has varied and complex actions on inputs to and cells within the ventral striatum, but the consequence of these actions are simple: to reinforce behaviour. This conjecture is supported by decades of preclinical research manipulating and observing dopamine activity during behaviour and particularly the indiscriminate capacity for drugs that elevate dopamine levels to reinforce behaviour [53]. However, much of this work relied on changes in behavioural frequency to describe the reinforcing properties of dopamine. Even still, some findings hinted a disengaging property of dopamine. For example, psychostimulants paired with food consumption [54,55], even when administered volitionally [56], reduced food consumption while encouraging food approach

[57]. Further, the self-administration of psychostimulants is reduced, not amplified, by microinfusion of dopamine agonists into the accumbens [58]. These findings point to an inherent role of dopamine receptor activation in disengaging behaviour.

Here, using a temporally constrained self-stimulation procedure we report behaviour consistent with dissociable reinforcing and disengaging properties of medial accumbens shell dopamine. Specifically, intense dopamine activity in the medial accumbens shell is powerfully reinforcing but produces short, numerous, self-stimulation hold-downs. The reinforcing and disengaging properties of medial accumbens shell dopamine are differentially affected by neurotransmission at dopamine receptors. Notably, augmenting neurotransmission at D1 receptors serves to enhance the disengaging property, whereas blunting neurotransmission at D2 receptors amplifies the reinforcing property, of medial accumbens shell dopamine – effects which may arise from unintuitive striatal responses to dopamine activity [59,60]. The current finding that medial accumbens shell dopamine is both reinforcing and disengaging runs counter to conventional views of striatal dopamine but is well accompanied by recent work diversifying the role of dopamine in psychology [61–63].

Acknowledgements
M. D. V. was supported by a postdoctoral fellowship from the Fonds de recherche du Québec – Santé (FRQS) and a Brain Star Award from the Institute of Neurosciences, Mental Health, and Addiction of the Canadian Institutes of Health Research (CIHR). N. M-L. E. was supported by a graduate fellowship from the FRQS. J. P. B. was funded by the Natural Sciences and Engineering Research Council (05100-2020), the CIHR (PT-74038), the Canadian Research Council (950-233013), and the G. W. Stairs fund at McGill University (jointly held by M. D. V. and J. P. B.). The authors thank Dr. Bruno Giros for providing some of the mice used in this study.

Declarations
The authors have no competing interests to declare.

References

[1] Olds J, Milner P. Positive reinforcement produced by electrical stimulation of septal area and other regions of rat brain. Journal of Comparative and Physiological … 1954;47:419–27.

[2] Sem-Jacobsen CW, Torkildsen A. Depth recording and electrical stimulation in the human brain. In: Ramey ER, D S O, editors. Electrical Studies on the Unanesthetized Brain, P. B. Hoeber; 1960, p. 275–90.

[3] Bishop MP, Elder ST, Heath RG. Intracranial Self-Stimulation in Man. Science (1979) 1963;140:394–6. https://doi.org/10.1126/science.140.3565.394.

[4] Shizgal P. Brain Stimulation Reward. International Encyclopedia of the Social & Behavioral Sciences 2001;1:1358–62. https://doi.org/10.1007/978-3-642-36172-2_200188.

[5] Gallistel CR, Shizgal P, Yeomans JS. A portrait of the substrate for self-stimulation. Psychol Rev 1981;88:228–73. https://doi.org/10.1037/0033-295X.88.3.228.

[6] Olds ME, Olds J. Approach-avoidance analysis of rat diencephalon. Journal of Comparative Neurology 1963;120:259–95.

[7] Margules DL. Separation of Positive and Negative Reinforcing Systems in the Diencephalon of the Rat. Am J Psychol 1966;79:205–16.

[8] Olds J. Approach-avoidance dissociations in rat brain. Am J Physiol 1960;199:965–8. https://doi.org/10.1152/ajplegacy.1960.199.6.965.

[9] Moisan J, Rompré PP. Electrophysiological evidence that a subset of midbrain dopamine neurons integrate the reward signal induced by electrical stimulation of the posterior mesencephalon. Brain Res 1998;786:143–52. https://doi.org/10.1016/S0006-8993(97)01457-1.

[10] Hernandez L, Hoebel BG. Feeding and hypothalamic stimulation increase dopamine turnover in the accumbens. Physiol Behav 1988;44:599–606. https://doi.org/10.1016/0031-9384(88)90324-1.

[11] Hernandez L, Hoebel BG. Food reward and cocaine increase extracellular dopamine in the nucleus accumbens as measured by microdialysis. Life Science 1988;42:1705–12.

[12] Cossette MP, Conover K, Shizgal P. The neural substrates for the rewarding and dopamine-releasing effects of medial forebrain bundle stimulation have partially discrepant frequency responses. Behavioural Brain Research 2016;297:345–58. https://doi.org/10.1016/j.bbr.2015.10.029.

[13] Duchesne V, Boye SM. Differential contribution of mesoaccumbens and mesohabenular dopamine to intracranial self-stimulation. Neuropharmacology 2013;70:43–50. https://doi.org/10.1016/j.neuropharm.2013.01.005.

[14] Wise RA. Dual roles of dopamine in food and drug seeking: the drive-reward paradox. Biol Psychiatry 2013;73:819–26. https://doi.org/10.1016/j.biopsych.2012.09.001.

[15] Trujillo-Pisanty I, Conover K, Solis P, Palacios D, Shizgal P. Dopamine neurons do not constitute an obligatory stage in the final common path for the evaluation and pursuit of brain stimulation reward. PLoS One 2020;15:e0226722. https://doi.org/10.1371/journal.pone.0226722.

[16] Ikemoto S. Brain reward circuitry beyond the mesolimbic dopamine system: A neurobiological theory. Neurosci Biobehav Rev 2010;35:129–50. https://doi.org/10.1016/j.neubiorev.2010.02.001.

[17] Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K. Millisecond-timescale, genetically targeted optical control of neural activity. Nat Neurosci 2005;8:1263–8. https://doi.org/10.1038/nn1525.

[18] Witten IB, Steinberg EE, Lee SY, Davidson TJ, Zalocusky KA, Brodsky M, et al. Recombinase-driver rat lines: Tools, techniques, and optogenetic application to dopamine-mediated reinforcement. Neuron 2011;72:721–33. https://doi.org/10.1016/j.neuron.2011.10.028.

[19] Tsai H-C, Zhang F, Adamantidis AR, Stuber GD, Bonci A, de Lecea L, et al. Phasic Firing in Dopaminergic Neurons is Sufficient for Behavioral Conditioning. Science (1979) 2009;324:1080–4. https://doi.org/10.1126/science.1168878.

[20] Adamantidis AR, Tsai HC, Boutrel B, Zhang F, Stuber GD, Budygin EA, et al. Optogenetic interrogation of dopaminergic modulation of the multiple phases of reward-seeking behavior. Journal of Neuroscience 2011;31:10829–35. https://doi.org/10.1523/JNEUROSCI.2246-11.2011.

[21] Yoo JH, Zell V, Gutierrez-Reed N, Wu J, Ressler R, Shenasa MA, et al. Ventral tegmental area glutamate neurons co-release GABA and promote positive reinforcement. Nat Commun 2016;7:1–13. https://doi.org/10.1038/ncomms13697.

[22] Schultz W, Dayan P, Montague R. A Neural Substrate of Prediction and Reward. Science (1979) 1997;275:1593–9. https://doi.org/10.1126/science.275.5306.1593.

[23] Montague PR, Dayan P, Sejnowski TJ. A framework for mesencephalic dopamine systems based on predictive hebbian learning. The Journal of Neuroscience 1996;16:1936–47.

[24] Usypchuk AA, Maes EJ, Lozzi M, Gardner MPH, Schoenbaum G, Esber GR, et al. Rewarding Value or Prediction Error: Settling the debate over the role of dopamine in reward learning n.d. https://doi.org/10.1101/2022.11.06.515338.

[25] Valyear MD, Glovaci I, Zaari A, Lahlou S, Trujillo-pisanty I, Chapman CA, et al. Dissociable mesolimbic dopamine circuits control responding triggered by alcohol-predictive discrete cues and contexts. Nat Commun 2020;11:1–14. https://doi.org/10.1038/s41467-020-17543-4.

[26] Howe MW, Dombeck DA. Rapid signalling in distinct dopaminergic axons during locomotion and reward. Nature 2016;535:505–10. https://doi.org/10.1038/nature18942.

[27] Saunders BT, Richard JM, Margolis EB, Janak PH. Dopamine neurons create Pavlovian conditioned stimuli with circuit-defined motivational properties. Nat Neurosci 2018;21:1. https://doi.org/10.1038/s41593-018-0191-4.

[28] Saddoris MP, Cacciapaglia F, Wightman RM, Carelli RM. Differential dopamine release dynamics in the nucleus accumbens core and shell reveal complementary signals for error prediction and incentive motivation. The Journal of Neuroscience 2015;35:11572–82. https://doi.org/10.1523/JNEUROSCI.2344-15.2015.

[29] Roitman MF, Wheeler RA, Wightman RM, Carelli RM. Real-time chemical responses in the nucleus accumbens differentiate rewarding and aversive stimuli. Nature Neuroscience 2008 11:12 2008;11:1376–7. https://doi.org/10.1038/nn.2219.

[30] Foley WJ, Iason GR, Mcarthurr C. Role of plant secondary metabolites in the nutritional ecology of mammalian herbivores: how far have we come in 25 years? In: Jung HG, Fahey GC, editors. Nutritional ecology of herbivores: Proceedings of the Vth Internaional Symposium on the Nutrition of Herbivores, Savoy, IL: American society of animal science; 1999, p. 130–209.

[31] Weinstein SB, Dearing MD. Harvest mice (Reithrodontomys megalotis) consume monarch butterflies (Danaus plexippus). Ecology 2022;103:1–4. https://doi.org/10.1002/ecy.3607.

[32] Torregrossa A, Dearing MD. Nutritional Toxicology of Mammals: Regulated Intake of Plant Secondary Compounds. Funct Ecol 2009;23:48–56.

[33] Smith JC. Microstructure of the rat's intake of food, sucrose and saccharin in 24- hour tests. Neurosci Biobehav Rev 2000;24:199–212. https://doi.org/10.1016/S0149-7634(99)00073-1.

[34] Dwyer DM. Licking and liking: The assessment of hedonic responses in rodents. Quarterly Journal of Experimental Psychology 2012;65:371–94. https://doi.org/10.1080/17470218.2011.652969.

[35] Wright SP. Adjusted p-values for simultaneous inference. Biometrics 1992;48:1005–13. https://doi.org/10.1126/science.87.2249.103-a.

[36] Moore NA, Axton MS. Production of climbing behaviour in mice requires both D1 and D2 receptor activation. Psychopharmacology (Berl) 1988;94:263–6. https://doi.org/10.1007/BF00176857.

[37] Akhisaroglu M, Kurtuncu M, Manev H, Uz T. Diurnal rhythms in quinpirole-induced locomotor behaviors and striatal D2/D3 receptor levels in mice. Pharmacol Biochem Behav 2005;80:371–7. https://doi.org/10.1016/j.pbb.2004.11.016.

[38] Luque-Rojas MJ, Galeano P, Suárez J, Araos P, Santín LJ, De Fonseca FR, et al. Hyperactivity induced by the dopamine D2/D3 receptor agonist quinpirole is attenuated by inhibitors of endocannabinoid degradation in mice. International Journal of Neuropsychopharmacology 2013;16:661–76. https://doi.org/10.1017/S1461145712000569.

[39] Ponnusamy R, Nissim HA, Barad M. Systemic blockade of D2-like dopamine receptors facilitates extinction of conditioned fear in mice. Learning and Memory 2005;12:399–406. https://doi.org/10.1101/lm.96605.

[40] Zarrindast MR, Eliassi A. Differential effects of dopamine agonists on locomotion in intact and reserpine-treated mice. Gen Pharmacol 1991;22:1027–31. https://doi.org/10.1016/0306-3623(91)90573-O.

[41] Eshel G, Ross SB, Kelder D, Edis LEM, Jackson DM. α1 (but not α2)-Adrenoceptor Agonists in Combination with the Dopamine D2 Agonist Quinpirole Produce Locomotor Stimulation in Dopamine-Depleted Mice. Pharmacol Toxicol 1990;67:123–31. https://doi.org/10.1111/j.1600-0773.1990.tb00797.x.

[42] O'Neill MF, Shaw G. Comparison of dopamine receptor antagonists on hyperlocomotion induced by cocaine, amphetamine, MK-801 and the dopamine D1 agonist C-APB in mice. Psychopharmacology (Berl) 1999;145:237–50. https://doi.org/10.1007/s002130051055.

[43] Karper PE, De La Rosa H, Newman ER, Krall CM, Nazarian A, McDougall SA, et al. Role of D1-like receptors in amphetamine-induced behavioral sensitization: A study using d1a receptor knockout mice. Psychopharmacology (Berl) 2002;159:407–14. https://doi.org/10.1007/s00213-001-0936-7.

[44] Cabib S, Castellano C, Cestari V, Filibeck U, Puglisi-Allegra S. D1 and D2 receptor antagonists differently affect cocaine-induced locomotor hyperactivity in the mouse. Psychopharmacology (Berl) 1991;105:335–9. https://doi.org/10.1007/BF02244427.

[45] Womer DE, Jones BC, Gene Erwin V. Characterization of dopamine transporter and locomotor effects of cocaine, GBR 12909, epidepride, and SCH 23390 in C57BL and DBA mice. Pharmacol Biochem Behav 1994;48:327–35. https://doi.org/10.1016/0091-3057(94)90534-7.

[46] Rossi MA, Sukharnikova T, Hayrapetyan VY, Yang L, Yin HH. Operant Self-Stimulation of Dopamine Neurons in the Substantia Nigra. PLoS One 2013;8:1–7. https://doi.org/10.1371/journal.pone.0065799.

[47] Humburg BA, Jordan CJ, Zhang H, Shen H, Han X, Bi G, et al. Optogenetic brain-stimulation reward: A new procedure to re-evaluate the rewarding *versus* aversive effects of cannabinoids in dopamine transporter-Cre mice. Addiction Biology 2021;26. https://doi.org/10.1111/adb.13005.

[48] Cox J, Witten IB. Striatal circuits for reward learning and decision-making. Nat Rev Neurosci 2019;20:482–94. https://doi.org/10.1038/s41583-019-0189-2.

[49] Kravitz A V, Tye LD, Kreitzer AC. Distinct roles for direct and indirect pathway striatal neurons in reinforcement. Nat Neurosci 2012;15:816–8. https://doi.org/10.1038/nn.3100.

[50] Wang W, Dever D, Lowe J, Storey GP, Bhansali A, Eck EK, et al. Regulation of prefrontal excitatory neurotransmission by dopamine in the nucleus accumbens core. Journal of Physiology 2012;590:3743–69. https://doi.org/10.1113/jphysiol.2012.235200.

[51] Qi J, Zhang S, Wang H-L, Barker DJ, Miranda-Barrientos J, Morales M. VTA glutamatergic inputs to nucleus accumbens drive aversion by acting on GABAergic interneurons. Nat Neurosci 2016;19. https://doi.org/10.1038/nn.4281.

[52] Warlow SM, Singhal SM, Hollon NG, Hunker AC, Zweifel LS, Hnasko TS. Mesoaccumbal glutamate neurons drive reward via glutamate release but aversion via dopamine co-release. Neuron 2023. https://doi.org/10.1016/j.neuron.2023.11.002.

[53] Wise RA, Robble MA. Dopamine and Addiction. Annu Rev Psychol 2020;71:79–106. https://doi.org/10.1146/annurev-psych-010418-103337.

[54] Reicher MA, Holman EW. Location preference and flavor aversion reinforced by amphetamine in rats. Anim Learn Behav 1977;5:343–6. https://doi.org/10.3758/BF03209576/METRICS.

[55] Parker LA. Taste avoidance and taste aversion: Evidence for two different processes. Learn Behav 2003;31:165–72.

[56] Wise RA, Yokel RA, de Wit H. Both Positive Reinforcement and Conditioned Aversion from Amphetamine and from Apomorphine in Rats. Science (1979) 1976;191:1273–5.

[57] White NM, Sklar L, Amit Z. The reinforcing action of morphine and its paradoxical side effect. Psychopharmacology (Berl) 1977;52:63–6.

[58] Suto N, Wise RA. Satiating effects of cocaine are controlled by dopamine actions in the nucleus accumbens core. The Journal of Neuroscience 2011;31:17917–22. https://doi.org/10.1523/JNEUROSCI.1903-11.2011.

[59] Parker JG, Marshall JD, Ahanonu B, Wu YW, Kim TH, Grewe BF, et al. Diametric neural ensemble dynamics in parkinsonian and dyskinetic states. Nature 2018;557:177–82. https://doi.org/10.1038/s41586-018-0090-6.

[60] Yun S, Yang B, Martin MM, Yeh N, Contractor A, Parker JG. Modulating D1 rather than D2 receptor-expressing spiny-projection neurons corresponds to optimal antipsychotic effect. BioRxiv 2021:1–55.

[61] Sharpe MJ, Chang CY, Liu MA, Batchelor HM, Mueller LE, Jones JL, et al. Dopamine transients are sufficient and necessary for acquisition of model-based associations. Nat Neurosci 2017;20:735–42. https://doi.org/10.1038/nn.4538.

[62] Engelhard B, Finkelstein J, Cox J, Fleming W, Jang HJ, Ornelas S, et al. Specialized coding of sensory, motor, and cognitive variables in midbrain dopamine neurons. Nature 2019:456194. https://doi.org/10.1101/456194.

[63] Carter F, Cossette M, Trujillo-Pisanty I, Pallikaras V, Breton Y, Conover K, et al. Does phasic dopamine release cause policy updates? European Journal of Neuroscience 2023. https://doi.org/10.1111/EJN.16199.

Fig. 1 | Temporally constrained self-stimulation of dopamine inputs to the medial accumbens shell. a, Active and inactive levers were inserted into the conditioning chamber for 30 s trials (48 trials/session) which were separated by intertrial intervals (ITI; ~10 s). Active lever hold-downs triggered the delivery of patterned laser light (473 nm) to the medial accumbens shell at 2.5, 10, or 40 Hz which terminated when the lever was released. **b**, DAT-Cre mice (n=12, 7F, 5M) received microinfusion of the viral construct AAV2/9-EF1a-fDIO-hChR2(H134R)-eYFP in the ventral tegmental area and **c**, implantation of an optical fibre in the medial accumbens shell which are shown on modified panels from the atlas of Franklin and Watson (2007). Representative microscopy (5x) images with overlaid anatomical delineations showing viral expression (eYFP, green) and nuclei (DAPI, blue) in the **d**, ventral tegmental area and **e**, medial accumbens shell. **f**, Across blocks of 5 sessions, mice increased the cumulative time spent holding down the active lever during 30 s trials whereas the cumulative time spent holding down the inactive lever remained low. Higher-frequency trials came to support more cumulative active hold-down time than did lower-frequency trials [Lever x Frequency x Session, $F_{(14, 154)}$=2.323, p=.006]. **g**, The mean duration of active, but not inactive, lever hold-downs increased over sessions [Lever x Session, $F_{(7, 77)}$=4.509, p<.001]. **h**, Mice increased the number of active, but not inactive, lever hold-downs performed during 30 s trials across sessions particularly so for high-frequency relative to low-frequency trials [Lever x Frequency x Session, $F_{(14, 154)}$=2.679, p=.002]. Averaged data are mean ± s.e.m.

Figure 1

Fig. 2 | Cumulative and mean duration of self-stimulation hold-downs diverge across frequencies. a, Mice spent significantly more cumulative time holding-down the active lever during 40 relative to 10 [$t_{(11)}$=3.367, p=.0083] or 2.5 [$t_{(11)}$=3.79, p=.003] Hz trials [Lever x Frequency, $F_{(2, 22)}$=3.594, p=.044]. **b**, Differently, the mean duration of individual hold-downs, were briefer at 40 relative to 10 [$t_{(11)}$=3.022, p=.0188] or 2.5 [$t_{(11)}$=3.061, p=.0172] Hz [Lever x Frequency, $F_{(2, 22)}$=5.3, p=.0132]. **c**, Consistently, mice performed more hold downs during 40 relative to 2.5 Hz trials [$t_{(11)}$=2.993, p=.020; Lever x Frequency, $F_{(2, 22)}$=2.607, p=.0963]. The **d**, cumulative **e**, mean and **f**, number of inactive lever hold-downs did not vary across stimulation frequencies. The distribution of the durations of **g**, 2.5, **h**, 10, and **i**, 40 Hz active hold-downs were positively skewed. The **j**, relative cumulative distribution of hold-downs during 40 Hz trials was significantly leftward shifted relative to 2.5 [D=.1517, p<.001] and 10 Hz [D=.1785, p<.001] trials. Data in a-f are shown as within-subjects differences from the lowest stimulation frequency (2.5 Hz). Averaged data are mean ± s.e.m. Symbols represent data from individual mice. Asterisks indicate Bonferroni-corrected and multiplicity-adjusted post-hoc comparisons p<.01**, p<.05*.

Figure 2

Fig. 3| Modulation of reinforcing and disengaging properties of medial accumbens shell dopamine. Mice performed longer cumulative active lever hold-downs during 40 Hz than 10 Hz [$t_{(9)}$=3.313, p=.0116] or 2.5 Hz [$t_{(9)}$=4.543, p=.0008] trials [Frequency, $F_{(2, 18)}$=11.04, p=.0007]. Simple effects reveal that the **a**, D1 agonist SKF38393 [$t_{(9)}$=12.94, p<.0001] and **d**, D2 agonist quinpirole [$t_{(9)}$=6.013, p<.00001] reduced the cumulative duration of hold-downs irrespective of stimulation frequency [Treatment, $F_{(72, 30)}$=73.3, p<.00001]. The D1 and D2 antagonists, SCH23390 [$t_{(9)}$=1.193, p>.9999] and Raclopride [$t_{(9)}$=2.118, p=.4118], did not affect cumulative hold-down duration. The mean duration of active lever hold-downs were briefer at 40 Hz than at 2.5 Hz [$t_{(9)}$=3.359, p=.0105] and modulated by the D1 agonist SKF38393 and the D2 antagonist Raclopride in a frequency-dependent manner [Frequency x Treatment, $F_{(8, 72)}$=4.948, p=.0002]. **b**, The D1 agonist SKF38393 shortened [2.5 Hz, $t_{(9)}$=6.22, p<.0001; 10 Hz, $t_{(9)}$=6.041, p<.0001; 40 Hz, $t_{(9)}$=5.125, p<.0001] whereas the **e**, D2 antagonist Raclopride lengthened [2.5 Hz, $t_{(9)}$=11.01, p<.0001; 10 Hz, $t_{(9)}$=12.32, p<.0001; 40 Hz, $t_{(9)}$=6.826, p=.0002] mean hold-down duration relative to Saline. The D1 antagonist SCH23390 and D2 agonist Quinpirole did not affect mean hold-down duration. Neither the **c**, D1 antagonist SCH23390 and D1 agonist SKF38393 or the **f**, D2 agonist Quinpirole and D2 antagonist raclopride affected the number of hold-downs during trials at any frequency [Frequency x Treatment, $F_{(8, 72)}$=1.471, p=.1829]. Data are shown as within-subjects differences from the Saline treatment condition. Averaged data are mean ± s.e.m. Symbols represent data from individual mice. Asterisks indicate Bonferroni-corrected and multiplicity-adjusted post-hoc comparisons p<.001***, p<.01**, p<.05*. Crosses indicate similarly corrected simple effects of treatment relative to saline p<.0001⊦.

Figure 3

Fig. 4│ The first self-stimulation hold-down. The latency to initiate an active hold-down after trial onset did not vary across stimulation frequencies [Frequency, $F_{(2, 18)}$=.04887, p=.9524], but was significantly increased [Treatment, $F_{(4, 36)}$=40.94, p<.001] following treatment with the **a**, D1 agonist SKF38393 [$t_{(9)}$=10.18, p<.001] and **c**, D2 agonist Quinpirole [$t_{(9)}$=4.395, p<.001], relative to saline. The mean duration of the first active hold-down during a trial was briefer at 40 than 2.5 Hz [$t_{(9)}$=3.039, p=.0212] and modulated by the D1 agonist SKF38393, the D2 agonist Quinpirole, and the D2 antagonist Raclopride in a frequency-dependent manner [Frequency x Treatment, $F_{(8, 72)}$=2.611, p=.0144]. **b**, The D1 agonist SKF38393 shortened [2.5 Hz, $t_{(9)}$=8.205, p<.0001; 10 Hz, $t_{(9)}$=6.869, p<.0001; 40 Hz, $t_{(9)}$=6.625 p<.0001] the duration of the first active hold-down relative to saline. **d**, The D2 agonist Quinpirole shortened [2.5 Hz, $t_{(9)}$=3.856, p=.0074; 10 Hz, $t_{(9)}$=3.614, $p=0.0167$; 40 Hz, $t_{(9)}$=4.137, p=.0028] whereas the D2 antagonist Raclopride lengthened the duration of the first active hold-down [2.5 Hz, $t_{(9)}$=8.154, p<.0001; 10 Hz, $t_{(9)}$=9.841, p<.0001; 40 Hz, $t_{(9)}$=5.077, p<.0001] relative to saline. Data are shown as within-subjects differences from the Saline treatment condition. Averaged data are mean ± s.e.m. Symbols represent data from individual mice. Asterisks indicate Bonferroni-corrected and multiplicity-adjusted post-hoc comparisons, p<.001***, p<.001***, p<.05*.

Figure 4