Coevolution-induced stabilizing and destabilizing selection shapes species richness in clade co-diversification

Yichao Zeng[1,*] & David H. Hembry[2]

[1] *Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721, USA*

[2] *Department of Biology, University of Texas Permian Basin, Odessa, TX 79762, USA*

* Correspondence: yichaozeng44@gmail.com

ABSTRACT

Coevolution can occur as a result of species interactions. However, it remains unclear how coevolutionary processes translate into the accumulation of species richness over macroevolutionary timescales. Assuming speciation occurs in a metacommunity as a result of genetic differentiation across communities due to dispersal limitation, we examine the effects of coevolution-induced stabilizing and destabilizing selection of a single quantitative trait on species diversification. We propose and test two hypotheses. (1) Stabilizing selection within communities enhances species diversification through strengthened dispersal limitation. (2) Destabilizing selection within communities impedes species diversification through weakened dispersal limitation. Here, we simulate clade co-diversification using an individual-based model, considering scenarios where phenotypic evolution is shaped by neutral dynamics, mutualistic coevolution, or antagonistic coevolution, where coevolution operates through trait matching or trait difference, and where the strength of coevolutionary selection is symmetrical or asymmetrical. Our assumption that interactions occur between an independent party (whose individuals can establish or persist in a community independently, e.g. hosts) and a dependent party (whose individuals cannot establish or persist in a community without the independent party, e.g. parasites or obligate mutualists) yields two contrasting results. Stabilizing selection within communities enhances species diversification in the dependent clade but not in the independent clade. Conversely, destabilizing selection within communities impedes species diversification in the independent clade but not in the dependent clade. These results are partially

28   corroborated by empirical dispersal data, suggesting that these mechanisms might explain the

29   diversification of some of the most species-rich clades in the Tree of Life.

30   **Keywords:** Species interaction, symbiosis, coevolution, stabilizing selection, destabilizing

31   selection, dispersal limitation, diversification, metacommunity

INTRODUCTION

The natural world is filled with interactions between different species. Many such interactions take the form of a bipartite partnerships in which two interaction parties play two distinct roles, e.g. the interactions between herbivores and plants, parasites and hosts, pollinators and angiosperms, and predators and prey. Bipartite interactions can potentially result in coevolution, that is, reciprocal evolutionary change in two or more interacting lineages driven by natural selection (Thompson 2005). Coevolutionary studies have focused on topics including phenotypic evolution across geographic space (Parchman and Benkman 2002), partnership specificity (Cook and Rasplus 2003), and signatures of coevolution in community assembly (Endara et al. 2017). Two coevolving lineages can co-diversify, i.e. diversify simultaneously, when they are allowed enough time. This has been an active area of research since the "escape-and-radiate" hypothesis, which states that rapid diversification of phytophagous insects or their independent plants often follows a significant shift in interaction partners (Ehrlich and Raven 1964; Cogni, Quental & Guimarães 2022). The ways by which macroevolutionary dynamics can be generated by processes at the local scale can be understood in the light of the geographic mosaic theory of coevolution, which emphasizes that coevolutionary dynamics can occur between geographically connected populations (Thompson 2005). It is therefore plausible that speciation could arise out of coevolutionary mosaics (Hembry, Yoder, and Goodman 2014; Thompson, Segraves, and Althoff 2017). Specifically, local coadaptation could potentially decrease the chance of mating between populations, paving the way for reproductive isolation and consequently speciation (Thompson, Segraves, and Althoff 2017). However, whether speciation is more likely to be enhanced or impeded by coevolution remains unclear (Janz 2011; Harmon et al. 2019; Hembry and Weber 2020).

Coevolution is shaped by the fitness outcomes of species interactions, which can vary in at least three ways. First, species interactions may benefit both parties or benefits one party at the cost of the other, affecting differently the fitness and, consequently, the coevolutionary outcome. Second, coevolution can be characterized by whether they are mediated by trait difference or trait matching (Yoder and Nuismer 2010). Under the trait-difference scenario, a species' fitness is maximized when its trait value is very different from that of its interaction partner. Examples of the trait-difference scenario (in the sense of Yoder and Nuismer 2010) include antagonistic

interactions in which predators and prey evolve increasingly strong weaponry and defense

against each other (Vermeij 1994) as well as mutualistic arms races in which both partners gain

more benefits from their partner when they have a greater trait value than that of their interaction

partner, e.g., a longer tongue in moth pollinators and a longer nectary spur in flowers (Whittall

and Hodges 2007). Under the trait-matching scenario (in the sense of Yoder and Nuismer 2010),

on the other hand, a species' coevolutionary fitness is maximized when its trait value more

closely matches that of its interaction partner. Antagonistic examples of this scenario include

brood parasitism (Langmore, Hunt, and Kilner 2003) whereas mutualistic examples include

obligate pollination mutualisms (Althoff and Segraves 2022). Third, the strength of natural

selection imposed by a species interaction is not always symmetrical for the interacting parties,

with the selection often being much stronger on one party than on the other (Brodie and Brodie

1999; Andreazzi, Thompson, and Guimaraes 2017; Endara et al. 2017).

Figure 1. Our model describes the interplay between processes at three scales: within-community, between
community, and across a metacommunity. Within community, species interactions can result in coevolution and thus
generate selective pressure for individuals of belonging to both interacting parties. Between communities,
individuals disperse and exchange genes. Across an entire metacommunity, genetic differentiation occurs between
different communities and eventually results in speciation.

Figure 2. Hypothesized connection between within-community and between-community dynamics. A
metacommunity is composed of multiple communities (circles) connected through dispersal (straight arrows).
Species interactions (curved arrows) occur within each community. The size of a straight arrow indicates the level of
dispersal. Compared to neutral drift as the baseline, stabilizing selection is hypothesized to impede dispersal whereas
destabilizing selection is hypothesized to enhance dispersal.

Different selective regimes (i.e. modes of phenotypic evolution) can potentially arise from the

different fitness outcomes of species interactions mentioned above. Prior literature has focused

on the microevolutionary effects of different selective regimes on coevolution (Yoder and

Nuismer 2010; Hembry et al. 2014). However, a theory is lacking on whether and how different

modes of phenotypic selection induced by coevolution drive or impede the accumulation of

species richness. The geographic mosaic theory of coevolution points to the possibility of

coevolutionary diversification of clades while leaving the exact mechanisms largely unexplored

(Thompson 2005). Given that certain types of species interactions are central to the biology of

many organisms (e.g. herbivory to the biology of phytophagous insects), a better understanding

of how coevolution shapes diversification through phenotypic evolution along this line could

potentially explain the diversification of many clades. Here we consider a general mode of

speciation where dispersal limitation causes genetic differentiation across space, eventually

leading to speciation (Moran 1962; Etienne & Alonso 2005; Rosindell, Harmon & Etienne 2015;

Manceau, Lambert, Morlon 2015; Hubert et al. 2015; Maliet, Loeuille & Morlon 2020). Based

on this mode of speciation, our model considers how within-community coevolutionary

dynamics shapes between-community dispersal dynamics and eventually shapes speciation

dynamics across an entire metacommunity (Fig. 1).

We propose two novel hypotheses regarding the different ways by which within-community

coevolution shapes between-community dispersal and consequently shapes species across an

entire metacommunity (Fig. 2). Over temporal scales, stabilizing selection is characterized by

increased temporal trait stability compared to neutral drift, and destabilizing selection can

conversely be defined as a selective regime that decreases temporal trait stability (Gingerich

2019a; Gingerich 2019b). The idea that within-population stabilizing selection limits between-

population dispersal has been supported both empirically and theoretically (Tufto 2000; Tufto

2001; Lopez el al. 2008; Scheepens, Frei & Stöcklin 2010; Yeaman & Whitlock 2011; Huisman

and Tufto 2012; Zacchello, Vinyeta, and Ågren 2020). This leads to the intuitive notion that, the

stronger temporal trait stability a selective regime causes, the more strongly dispersal is limited –

and a stronger degree of dispersal limitation causes stronger genetic differentiation across space

and eventually a higher species richness accumulation. Following this intuition, we propose a

first hypothesis termed the *stabilizing selection hypothesis*: stabilizing selection impedes

dispersal (and thus increases the degree of dispersal limitation), which results in an increase in

genetic differentiation across space, eventually resulting in an increase in the species richness

accumulation. The same intuition can lead to another hypothesis, which is termed the

destabilizing selection hypothesis: destabilizing selection enhances dispersal (and thus reduces

the degree of dispersal limitation), which results in a reduction in genetic differentiation across

space, eventually resulting in a reduction in the species richness accumulation.

To test these hypotheses in the context of two coevolving clades, we built an individual-based

model for the coevolutionary accumulation of species richness across two-dimensional space.

We considered different coevolutionary scenarios, including those where the interaction is

mutualistic versus antagonistic, where the interaction is mediated by trait differences versus trait

matching, and where coevolutionary selection is symmetrically versus asymmetrically strong for

the interacting parties. In bipartite interactions such as antagonisms and mutualisms, one partner

is often more dependent on the other than the other way around. Specifically, many antagonistic

interactions occur between antagonistic symbionts (e.g. parasites, pathogens) and their hosts

without which they cannot survive (Schmid-Hempel 2013). Even in mutualistic interactions, the

mutual dependence of interaction partners can be highly asymmetric (Bascompte, Jordano, and

Olesen 2006; Bronstein 2015). Given this, we built into our model the differences between

independence and dependence in partnership, assuming that the independent party's individuals,

e.g. hosts, can establish or persist in a community independently, whereas a dependent party's

individuals cannot establish or persist in a community without the independent party, e.g.

parasites or obligate mutualists. Our simulation results suggest mechanisms for the

diversification of some of the most species-rich groups in the Tree of Life (Table 1).

Table 1. Examples of species interactions that may be described as independent-dependent partnerships and the
species richness estimates for the independent and dependent parties.

Interaction	Independent party (species richness)	Dependent party (species richness)	Reference(s)
Host plants and herbivorous insects	Plants (ca. 374,000)	Herbivorous insects (over 500,000)	Bernays (2009); Christenhusz and Byng (2016)
Host and parasites	Hosts (NA)	Parasites (ca. 6,000,000)	Dobson et al. (2008);
Host plants and mutualistic mites	Some plants (NA)	Mutualistic mites (NA)	Romero and Benson (2005)
Corals and *Symbiodinium*	Corals (ca. 10,661 estimated for Anthozoa)	*Symbiodinium* (NA)	Bánki et al. (2023)
Insects and gut microbes	Insects (ca. 5,500,000)	Gut microbes (NA)	Stork (2018)

METHODS

Model description

Partnership between independent and dependent individuals

In the model, we consider the partnership between independent and dependent individuals. An

independent individual can partner with multiple dependent individuals, but a dependent

individual can only partner with one independent individual. An independent individual does not

need to partner with any dependent individual for survival, but a dependent individual needs the

one and only independent individual that it partners with for survival. This is modeled in two

separate ways. First, during dispersal, an independent individual could survive where no

dependent individuals are present; however, a dependent individual dies where no independent

individuals are present. Second, during competition, when an independent individual dies, so do

all of its dependent partners; however, when a dependent individual dies, its one and only

independent partner does not die. It is important to note that, in nature, the party that interacts

with multiple individuals are not necessarily the independent (less dependent) party.

Dispersal

The model starts with two populations each belonging to one of the interacting parties (the

independent and dependent) placed at the central site of a $n \times n$ grid, where each cell represents

a geographic site. Each individual, independent or dependent, has the opportunity to disperse

only at birth and moves to one of the four neighboring sites or else remain at the same site (each

with an equal probability of 0.2). As previously mentioned, if a dependent individual moves to a

site where no independent individuals are present, it dies; however, if it moves to a site where at

least one independent individual is present, the dependent individual randomly chooses an

independent individual to partner with among the independent individuals available at that site.

The entire grid forms a metacommunity composed of $n \times n$ communities (sites) connected

through dispersal.

Sexual reproduction and speciation

The genetic model largely follows a previous model built for the diversification of a single clade

(Aguilée et al. 2018), but with simplifications. Each independent or dependent individual is

diploid and has a given number of L_{dist} loci determining its genetic distance from another

individual. A new mutation can arise at any of these loci at a fixed probability μ_{dist}. To

determine whether two individuals are genetically incompatible enough to be considered

reproductively isolated, we calculated the genetic distance between two individuals based on

how many loci carry completely different alleles between the two individuals. The genetic

distance is considered sufficient for reproductive isolation if greater than a fixed genetic distance

threshold T_{dist}. These features allowed speciation through genetic differentiation across the grid.

Both independent and dependent individuals are hermaphrodites and reproduce sexually. For

independent individuals or dependent individuals, each mating attempt occurs within each site

following these steps: (1) two individuals with the highest fitness are chosen regardless of

whether they have reproduced before (see "Fitness outcomes of species interactions" for how

fitness is decided; for simplicity, we did not choose a more complicated reproduction model); (2)

the two parents successfully reproduce n_{offspr} offspring if they are not reproductively isolated.

These steps are repeated until n_{mat} mating attempts are made, regardless of whether mating

attempts result in successful reproduction. For each offspring, the genotypes of the genetic

distance loci are determined by Mendelian independent assortment of parental alleles. Given that

dependent individuals die along with their independent partner but not vice versa, it is necessary

for n_{offspr} and n_{mat} to be greater for the dependent individuals than for the independent

individuals so that the dependent individuals do not die out.

Genetics of phenotype

193   Each independent or dependent individual has one locus determining its ecological phenotype

194   which consists of a single quantitative trait. Mutation occurs at each of these loci, with the

195   mutated allele value drawn from a normal distribution with a mean equal to the parent allele

196   value and a standard deviation equal to σ_{eco}. For each offspring, the genotype of the ecological

197   phenotype locus is determined by a Mendelian random segregation of parental alleles.

Competitive death

Many types of antagonisms and mutualisms can occur between a consumer and a resource, such

as herbivory, parasitism, or pollination (Bronstein 2015). Given that populations cannot grow

infinitely in consumer-resource systems due to resource competition (Abrams 2009), we

considered there to be a growth rate of zero when population size reached carrying capacity. In

the model, resource competition occurs among dependent individuals partnering with the same

independent individual (e.g., parasites on the same host) and among independent individuals

within the same site. The number of mutually competing individuals n cannot grow above

carrying capacities $K_{independent}$ or $K_{dependent}$. We ensured this by assigning $n_{independent} -$

$K_{independent}$ or $n_{dependent} - K_{dependent}$ individuals to death (where $n_{independent}$ and

$n_{dependent}$ are the numbers of mutually competing individuals, for independents and dependents,

respectively), on a lowest-fitness-first basis (see "Fitness outcomes of species interactions" for

how fitness is decided). Again, when an independent individual dies, so do all of the dependent

individuals that partner with it; however, when a dependent individual dies, its one and only

independent partner does not die.

Fitness outcomes of species interactions

We allowed different modes of phenotypic evolution to arise from species interactions with

different fitness outcomes (Fig. 3). Antagonisms (interactions that benefit one party at the cost of

the other) and mutualisms vary in whether each of the two interacting parties receives a benefit

or pays a cost (Bronstein 2015). For either antagonisms or mutualisms, the fitness of phenotypes

was modeled as the result of either (i) trait difference: how different the two interacting

phenotypes were (the direction of phenotypic difference between interacting parties does matter),

or (ii) trait matching: how closely two interacting trait values matched each other (the direction

of phenotypic difference between interacting parties does not matter). Following Yoder and

Nuismer (2010), the fitness of an individual i interacting with an antagonist or mutualist

individual j can be expressed as

224
$$w_i\left(z_i, z_{j1}, z_{j2}, \dots, z_{jn}\right) = w_0 + \sum_{k=1}^{n} \frac{\xi}{1 + e^{-\alpha(z_i - z_{jk})}} \tag{1}$$

225   under trait difference, or

226
$$w_i\left(z_i, z_{j1}, z_{j2}, \dots, z_{jn}\right) = w_0 + \sum_{k=1}^{n} \xi e^{-\alpha(z_i - z_{jk})^2} \tag{2}$$

under trait matching, where z_i and z_j are phenotypic trait values of interacting individuals, n is

the number of partners that individual i has, w_0 is the individual i's fitness in the absence of the

interaction, ξ is the cost or benefit to the individual (positive if beneficial, negative if costly), and

α determines the sensitivity of the fitness outcome to the difference between interacting trait

values (i.e. deviation from fitness neutrality where trait values do not affect fitness). The strength

of coevolutionary selection can often be highly asymmetrical, i.e., much weaker on one

interacting party than on the other (Brodie and Brodie 1999; Andreazzi, Thompson, and

Guimaraes 2017; Endara et al. 2017). Here we considered the extreme case of asymmetrical

coevolutionary selection by allowing either the independents' or dependents' individual fitness

to be unaffected by trait values (Function II in Fig. 3). A fully factorial design would contain

4*4=16 scenarios given the all the combinations of the 4 fitness functions. We simulated a total

of 9 possible scenarios (Table 2) to include all possible combinations except 7 combinations of

which no or few empirical cases are known, i.e., those where one clade's fitness follows trait

difference while the other's follows trait matching (Yoder and Nuismer 2010) or where the

interaction is costly for the dependent clade.

Figure 3. In species interactions, individual fitness is a function of the traits of interacting individuals. Each surface
is the fitness of individual i (w_i) as a function of z_i, which denotes the trait value of individual i itself, and z_j, which
denotes the trait value for individual j (individual i's interaction partner). Function I: Equation 1, $\xi = 0.1, \alpha = 1$.
Function II: $w_i = 0$. Function III: Equation 2, $\xi = 0.1, \alpha = 1$. Function IV: Equation 2, $\xi = -0.1, \alpha = 1$. For this
model, Function II, where individual fitness is held constant and independent from trait values, is equivalent to
Equation 1 or 2 when $\alpha = 0$ regardless of the value of ξ. Following Yoder and Nuismer (2010), the fitness function
does not dependent on whether the interaction is beneficial or costly when coevolution operates through trait
difference (Function I). In all panels, the ranges of the z_i and z_j axes are between 0 and 2 and the range of the w_i
axis is between -0.1 and 0.1.

Table 2. The 9 scenarios considered in this study and their corresponding fitness functions (Figure 2). 4 fitness
functions (I-IV) necessitate a total of 9 combinations after unrealistic combinations are excluded.

Scenario	Trait difference or trait matching	Fitness function for the independent	Fitness function for the dependent	Outcome of interaction
a	Difference	I	I	Mutualism or antagonism
b	Difference	I	II	Mutualism or antagonism
c	Difference	II	I	Mutualism or antagonism
d	Matching	III	III	Mutualism
e	Matching	III	II	Mutualism
f	Matching	II	III	Mutualism or antagonism
g	Matching	IV	III	Antagonism
h	Matching	IV	II	Antagonism
i	Unaffected	II	II	Mutualism or antagonism

Simulations

The model was implemented and the simulation results were analyzed in the R language (v4.0.0;
R Core Team, 2020). All simulations were run on the University of Arizona High-Performance
Computing clusters. We ran 96 replicates for each of the 9 scenarios, totaling 864 simulations.
The large number of replicate simulations were made possible by custom code that can record
the progress of a simulation and resume where that simulation is terminated by the workload
manager. Each simulation was run for 1500 generations. For each of the 9 simulated scenarios,
t-tests comparing the distributions of mean species richness during the 1401^{th}-1450^{th} generations
versus the 1451^{th}-1500^{th} generations found no statistically significant difference ($P > 0.05$, $n = 96$
replicates for each of the two distributions being compared). This confirmed that the duration of
simulation (1500 generations) was enough for simulations to reach a stationary state in terms of
species richness, despite the ongoing fluctuations in species richness near the end of simulation
(Fig. S1-S18). The large number of replicate simulations made the analyses computationally
intense. This computational challenge was overcome though parallel computation using the R
package doParallel (v1.0.17; Microsoft Corporation and Weston). All custom code used in this
work is available as a supplementary file (File S1) and on GitHub
(https://github.com/Dragonfly4412/Macro_Coevolution). All the constants used in the
simulations are provided in Table S1.

Identifying selective regimes based on temporal trait (in)stability

To identify the modes of phenotypic evolution for each scenario, we recorded the phenotypic

values of all independent and dependent individuals at the central site of the $n \times n$ grid during

the entire duration of each simulation (1500 generations). We then quantified the change in mean

trait value during each generation, Δz. Δz is conventionally referred to as step difference and is

measured in haldanes, i.e., standard deviations per generation on a timescale of one generation

(Gingerich 2019a; Gingerich 2019b). For example, a Δz value of 2 haldanes means that the

change in mean trait value from Generation 1 to Generation 2 is two times the standard deviation

of the trait distribution at Generation 1. We further took the absolute value of Δz to get $|\Delta z|$. We

then averaged $|\Delta z|$ across the entire duration of simulation to get the mean step difference $\overline{|\Delta z|}$

as a measure of temporal trait instability. We determined the minimum and maximum for the

$\overline{|\Delta z|}$ of the entirely neutral scenario (Scenario i, where both the independents' and dependents'

fitness are held constant and do not depend on trait values). We treated simulations in which $\overline{|\Delta z|}$

is greater than the neutral maximum as destabilizing selection and those in which $\overline{|\Delta z|}$ is less

than the neutral minimum as stabilizing selection.

Degree of dispersal limitation, genetic distance between sites, and species richness

We quantified the degree of dispersal limitation as the proportion of native individuals among all

individuals, with higher proportions of native individuals indicating stronger degrees of dispersal

limitation. We define a native individual as an individual that inhabits the same site at which it

was born. Specifically, what we measured was realized dispersal rather than dispersal *per se*,

because we are interested in the contribution of dispersal to local gene pools, which is contingent

on successful establishment of the dispersers following dispersal. The degree of dispersal

limitation we refer to is the degree to which dispersal success is limited. Then, we calculated

genetic distance between any two sites (i.e., between the two populations inhabiting the two

sites) by taking the mean genetic distance between any two individuals from the two sites (i.e.,

the two populations). The genetic distance between all sites was then quantified as the mean

genetic distance between any two sites. To quantify the species richness that each clade had

accumulated at the end of the simulation, we considered populations to belong to a single species

if their genetic distance did not exceed the genetic distance threshold for speciation T_{dist} (for

details, see "Sexual reproduction and speciation").

For the degree of dispersal limitation (the proportion of native individuals), we took the mean

over the entire 1500 generations of simulation because it is the cumulative effect of dispersal that

is of interest. For genetic distance between sites and species richness accumulation, we took the

mean over the last 10 generations of simulation because we were interested in them as the

eventual results of multiple generations of dispersal and speciation.

Figure 4. Coevolutionary trajectories within communities in each of the nine simulated scenarios (*a-i*). Solid lines indicate the mean trait value calculated for individuals at the central community (site) of the $n \times n$ grid in one of the 96 replicates for each scenario. Blue and yellow are used to indicate the independent and dependent trait values, respectively.

RESULTS

Each of the 9 simulated scenarios generated a unique coevolutionary trajectory within

communities (Fig. 4). The 9 result coevolutionary trajectories were each characterized by rapid

and continuous reciprocal escalation of trait values in the two clades characteristic of an arms

race (Scenario *a*; Fig. 4), trait escalation in the independent clade and an initial decrease in trait

value followed by a random walk (in the sense that it does not substantially differ from a

trajectory typical of the neutral scenario) in the dependent clade (Scenario *b*; Fig. 4), trait

escalation followed by a random walk in the dependent clade and random walk in the

independent clade (Scenario *c*; Fig. 4), varying degrees of temporal trait stationarity in the two

clades (Scenarios *d-f*; Fig. 4), dramatic fluctuation in trait value in both clades characteristic of

coevolutionary cycling (Scenario *g*; Fig. 4), a tendency for the host trait value to not overlap with

that of the dependent (Scenario *h*; Fig. 4), and random walks underlain by neutral dynamics in

both clades (Scenario *i*; Fig. 4).

Analyses of the effects of selective regime on the degree of dispersal limitation, genetic distance

between sites, and species richness accumulation revealed the mechanisms by which coevolution

shaped species richness (Fig. 5). Quantification of the mean step difference $\overline{|\Delta z|}$ as a measure of

temporal trait instability showed that the 9 scenarios generated all three modes of phenotypic

evolution, i.e. neutral drift, stabilizing selection, and destabilizing selection in the two clades.

For each of the relationships of interest (Fig. 5, A-C), we performed a loess regression to

visualize non-linearities, but also two localized simple linear regressions to quantify the average

effects of stabilizing and destabilizing selection (Fig. S19).

In the independent clade (Fig. 5, D-F; Fig. S19, D-F), compared to neutral drift, stabilizing

selection resulted in relatively weak increases in the degree of dispersal limitation (slope = -1.47

percentage/haldane, $P = 0.0013$, left line in Fig. S19D), genetic distance between sites (slope = -

0.1664 loci/haldane, $P = 0.208$, left line in Fig. S19E), and species richness accumulation (slope

= -16.4918 species/haldane, $P = 0.00117$, left line in Fig. S19F). However, compared to neutral

drift, destabilizing selection resulted in relatively strong reductions in degree of dispersal

limitation (slope = -12.11 percentage/haldane, $P < 2e-16$, right line in Fig. S19D), genetic

distance between sites (slope = -7.60415 loci/haldane, $P < 2e-16$, right line in Fig. S19E), and

species richness accumulation (slope = -181.4629 species/haldane, $P < 2e-16$, right line in Fig.

S19F). These results suggest that the predominant mechanism shaping species richness in the

independent clade is one where destabilizing selection reduces the degree of dispersal limitation,

which reduces genetic distance between sites, eventually resulting in reduced species richness

accumulation.

In the dependent clade (Fig. 5, G-I; Fig. S19, G-I), compared to neutral drift, stabilizing selection

resulted in relatively strong increases in the degree of dispersal limitation (slope = -3.755

percentage/haldane, $P < 2e\text{-}16$, left line in Fig. S19G), genetic distance between sites (slope = -

1.0355 loci/haldane, $P = 2.87e\text{-}12$, left line in Fig. S19H), and species richness accumulation

(slope = -35.7923 species/haldane, $P < 2e\text{-}16$, left line in Fig. S19I). The effect of stabilizing

selection showed strong non-linearity, with the highest degree of dispersal limitation, genetic

distance between sites, and species richness accumulation being achieved by intermediately

rather than extremely strong stabilizing selection (Fig. 5, G-I). However, compared to neutral

drift, destabilizing selection resulted in relatively week reductions or even a slight increase in the

degree of dispersal limitation (slope = -0.00114 percentage/haldane, $P = 0.993$, right line in Fig.

S19G), genetic distance between sites (slope = -0.08698 loci/haldane, $P = 0.853$, right line in

Fig. S19H), and species richness accumulation (slope = 6.4800 species/haldane, $P = 0.375$, right

line in Fig. S19I). These results suggest that the predominant mechanism shaping species

richness in the dependent clade is one where stabilizing selection increased the degree of

dispersal limitation, which increases genetic distance between sites, eventually resulting in

increased species richness accumulation.

Specifically, the destabilizing selection hypothesis explained how species richness accumulation

was reduced for the independent clade in the coevolutionary scenarios of mutualistic or

antagonistic arms races where selective pressure is comparable between the independent and

dependent individuals, i.e., a classical arms race, and in the scenario of antagonistic trait

matching where selective pressure is comparable between the independent and dependent

individuals (Fig. 5, D-F; Fig. S19, D-F; Scenarios a & g in Table 2). The variable $\overline{|\Delta z|}$, as a

measure of temporal trait instability, is turned out to be key in shaping species richness

accumulation. This is because trait instability is a direct result of coevolution and in turn shapes

dispersal and consequently speciation. The stabilizing selection hypothesis explained how

species richness was increased for the dependent clade in the coevolutionary scenario of

mutualistic trait matching where selective pressure is comparable between the independent and

dependent individuals, the scenario of mutualistic or antagonistic trait matching where selective

pressure is weak for independent individuals, and the scenario of antagonistic trait matching

where selective pressure is comparable between the independent and dependent individuals (Fig.

5, G-I; Fig. S19, G-I; Scenarios d, f & g in Table 2).

Figure 5. Observed relationships between with-community dynamics (trait instability), between-community dynamics (degree of dispersal limitation), and metacommunity dynamics (genetic distance between sites and species richness accumulation). Selective regimes (neutral drift, stabilizing selection, or destabilizing selection) are determined by trait instability measured as the mean step difference $\overline{|\Delta z|}$. (A)-(C): Expected relationships – a significantly negative slope is predicted based on the stabilizing and destabilizing selection hypotheses. (D)-(F): Observed differences between selection regimes in the independent clade. (G)-(I): Observed differences between selection regimes in the dependent clade. In (D)-(I), black lines with gray ribbons show loess regressions with 95% confidence intervals.

DISCUSSION

Here we have shown that coevolution shapes species richness through two different mechanisms depending on whether the clade of interest is independent or dependent. In the independent clade, destabilizing selection enhances dispersal, which results in a reduction in the genetic distance across space, eventually resulting in a reduction in the species richness accumulation (i.e. the destabilizing selection hypothesis). In the dependent clade, stabilizing selection impedes dispersal, which results in an increase in the genetic distance cross space, eventually resulting in an increase in the species richness accumulation (i.e. the stabilizing selection hypothesis). These results arose through two simple rules governing the dynamics of independent-dependent interactions: one independent individual can pair with multiple dependent individuals but not vice versa; a dependent individual dies in the absence of an independent partner but not vice versa. These rules should be applicable to a wide variety of systems including most host-phytophagous-insect interactions, most host-parasite interactions, most host-pathogen interactions, and some obligate interactions between mutualists and their hosts, although exceptions inevitably exist because the biology of species interactions is highly diverse..

Empirical data generally corroborate our results that coevolution-induced stabilizing and destabilizing selection differentially affect the dispersal of independent and dependent individuals, suggesting that our model provides a potential mechanistic explanation for these patterns of dispersal. Previous models have explored the conditions for coevolutionary stable states or strategies to be achieved, yet these model did not consider dispersal or speciation across space (Vasco, Nazarea & Richardson 1987; Gilchrist & Sasaki 2002; Day & Burns 2003). Empirical examples of coevolution-induced destabilizing selection include classical arms race where there is reciprocal escalation of trait values (Dawkins & Krebs 1979; Vermeij 1994; Whittall & Hodges 2007) and coevolutionary cycling where there is persistent alternation between different trait states such between high and low trait values (Prado et al. 2009; Ashby & Gupta 2014; Ashby & Boots 2015). Whether coevolution causes stabilizing or destabilizing selection is usually unknown except in a limited number of empirical systems, but the results of our model tend to match several empirical observations pertaining to ant-plant, ant-bacterium, and plant-fungus mutualisms. First, our model shows that coevolution-induced stabilizing selection impedes the dispersal of dependent individuals only. Although this conclusion is based

upon correlation, the direction of causation is inambiguous because the model took a bottom-up

approach where dispersal, a between-community process, arose as a result of species interaction

and coevlution within communities. This matches large-scale observations that only specialized

mutualism (high dependence on mutualistic partners) is associated with a reduced likelihood of

successful establishment beyond native ranges (Nathan et al. 2023). Second, our model shows

that coevolution-induced destabilizing selection enhances the dispersal of independent

individuals only. This matches large-scale observations that only generalized mutualism (low

dependence on mutualistic partners) is associated with an increased likelihood of successful

establishment beyond native ranges (Nathan et al. 2023). We show that coevolution-induced

stabilizing and destabilizing selection can generate these widely observable patterns. This

suggests that coevolution-induced stabilizing and destabilizing selection, as well as their

macroevolutionary consequences as shown in our model, may be prevalent in nature.

It is useful to understand these results in the light of environmental selection and trait multi-

dimensionality. Some traits under coevolutionary selection may be simultaneously under

selection from other factors in the environment. Selection imposed by the environment can be

either stabilizing or destabilizing – environment-induced stabilizing selection can operate

through selection against phenotypes that deviate from a fixed environmental optimum (Kopp &

Matuszewski 2014), whereas fluctuating environment can cause potentially destabilizing

selection (De Villemereuil 2020). Disruptive selection along an environmental gradient has itself

been shown as a mechanism driving speciation (Doebeli & Dieckmann 2003). It would be

interesting for future research to explore the interactive effects of coevolutionary selection and

environmental selection on co-diversification, as has been pointed out in cases where biotic

interactions drive speciation in tandem with spatial isolation (Kay and Sargent 2009, Hembry et

al. 2014). In addition, species interactions are sometime better modeled using multi-dimensional

traits (Eklöf et al. 2013). The trait difference mechanism can have similar effects on phenotypic

evolution to those of the trait matching mechanism when the trait of interest is multidimensional

(Yamamichi, Lyberger & Patel 2019). Therefore, destabilizing selection underlain by the trait

difference mechanism might be less common in systems with higher trait dimensionalities.

These results may provide mechanistic insights into the diversification of some extremely

diverse clades that fit the description of a dependent clade. Specifically, explaining the origins of

the diversity of phytophagous insects and parasites has long been an active area of research

(Ehrlich and Raven 1964; Poulin and Morand 2000; Hardy and Otto 2014; Weinstein and Kuris

2016; Kawahara et al. 2023), and our model provided a potential explanation for their staggering

diversity: stabilizing selection induced by coevolution selects against immigrants, reducing

dispersal success and consequently enhancing genetic differentiation across space that is

necessary for speciation under our assumption. This is conceptually similar to the notion that

local coadaptation can reduce the chance of mating between populations and this reduction in

gene flow can promote reproductive isolation and consequently speciation (Thompson, Segraves

& Althoff 2017). However, we furthered this notion by showing that this is likely to be the case

only for clades that fit our assumption about dependence such as phytophagous insects, parasites,

pathogens, or obligate mutualists that depend on facultative hosts, but is unlikely to be the case

for clades that fit our assumption about independence such as various clades of hosts as

mutualists or victims of phytophagous insects, parasites, and pathogens. Overall, the mechanisms

from this model provide potential mechanisms by which coevolution may have had a profound

impact on the diversification of phytophagous insects and parasites, although the prevalence of

coevolution in insect herbivory and parasitism has yet to be confirmed by empirical evidence.

Our results suggest that the diversity of clades that fit our description of an independent clade

(e.g., plants, hosts of parasites and pathogens, or the facultative hosts of obligate mutualists)

could also be explained in the light of coevolution. The effect of coevolution on an independent

clade's diversity, under our model assumptions, is likely negative for diversification as shown in

the destabilizing selection hypothesis. It is interesting in this light that independent clades, e.g.,

clades of angiosperms are often more species-poor than their dependent clades, e.g., clades of

phytophagous insects (Bernays 2009; Christenhusz & Byng 2016), although many other

mechanisms likely contribute to this asymmetry in richness. This decelerating effect of

coevolution on diversification that we show here has not been explored before, as classic

coevolutionary hypotheses such as the escape-and-radiate hypothesis tended to focus on the

accelerating effects of coevolution on diversification (Ehrlich and Raven 1964). The escape-and-

radiate hypothesis is less interested in speciation events per se than in bursts of diversification in

lineages that escape herbivory and is not specifically accounted for in our model.

Comparing our model to previous models provides some additional implications. We show that a

classical arms race impedes the accumulation of species richness in the independent clade but

has no effect in the dependent clade. Our results focusing on species richness is in contrast with

the results of a previous study showing that the classical arms race neither promotes nor inhibits

phenotypic diversification (Yoder and Nuismer 2010). We also show that destabilizing selection

also predicts a decrease in species richness in the independent clade in the scenario of

antagonistic trait matching where selective pressure is comparable between the independent and

dependent individuals. This also contrasts with the previous model which showed that

coevolution promotes phenotypic diversification when trait matching is costly, e.g., as in

competition or antagonisms. These contrasts suggest that phenotypic diversification can be

decoupled from species diversification during coevolution. The idea that having multiple

partners poses a constraint on diversification when the interaction is mutualistic and based on

trait matching has been shown in a non-spatially explicit model previously (Raimundo et al.

2014). This agrees with the results of our spatially explicit model and suggests that our results

could potentially arise through similar mechanisms. Overall, the new model contributes to an

ongoing effort to integrate metacommunity ecology with macroevolution in general (McPeek

2008; Reijenga, Murrell & Pigot 2021) and for independent-dependent systems in particular

(Forister & Jenkins 2017).

Our model provides an alternative or supportive explanation for some long-standing hypotheses.

In the seminal escape-and-radiate hypothesis (Ehrlich & Raven 1964), a lineage, independent or

dependent, that acquires a new defense or counter-defense (i.e. key innovations) may then

rapidly radiate into a new adaptive zone. We show a different picture here – coevolutionary

spatial dynamics can continuously enhance or impede speciation without involving sporadic key

innovations. There are also long-standing hypotheses that dependents should be more specialized

the more intimate their interactions with their independents are (Ollerton 2006; Thompson 1994).

These hypotheses generally predict that dependents should be more species-rich than

independents, which is consistent with our results suggesting that coevolution generally impedes

diversification for an independent clade and enhances diversification for a dependent clade.

However, the intimacy hypotheses are agnostic to the geography of speciation. It is also clear

that a single independent can contain a multitude of niches for different dependents (Farrell &

Sequeira 2004), so the asymmetry in richness is less surprising regardless of the specific

mechanisms of diversification.

In conclusion, here we show that there are two general mechanisms of coevolutionary

diversification: coevolution-induced stabilizing selection enhances the accumulation of species

richness in dependent clades, whereas coevolution-induced destabilizing selection impedes the

accumulation of species richness in independent clades. The model provides a new line of

thinking in bridging symbiotic biology, coevolution, metacommunity ecology, and

macroevolution. Given that symbiotic relationships between a more dependent party and a less

dependent party permeate the natural world (Margulis 1998), these general mechanisms of

coevolutionary diversification can potentially explain the diversification of many clades in the

Tree of Life.

Authorship statement

522   **Yichao Zeng**: Conceptualization, Methodology, Software, Validation, Formal analysis,

523   Investigation, Data cutation, Writing – Original draft, Review & Editing, Visualization. **David

524   H. Hembry**: Writing – Original draft, Review & Editing.

Acknowledgements

We thank Judie Bronstein, Régis Ferrière, Paulo R. Guimarães Jr., and John Wiens for

discussing and reviewing early versions of this manuscript. John Wiens edited early versions of

this manuscript. High-performance computing resources were provided by the University of

Arizona. These high-performance computing resources were funded by central funding and were

free to use for all university affiliates

(https://public.confluence.arizona.edu/display/UAHPC/Compute+

Resources). We thank Sara Willis for technical support with cluster computing.

REFERENCECS

Abrams, Peter A. 2009. "Determining the Functional Form of Density Dependence: Deductive

Approaches for Consumer-Resource Systems Having a Single Resource." *The American*

Naturalist 174 (3): 321–30.

Aguilée, Robin, Fanny Gascuel, Amaury Lambert, and Regis Ferriere. 2018. "Clade

diversification dynamics and the biotic and abiotic controls of speciation and extinction rates."

Nature Communications 2018 9:1 9 (1): 1–13.

Althoff, David M, and Kari A Segraves. 2022. "Evolution of Antagonistic and Mutualistic Traits

in the Yucca-Yucca Moth Obligate Pollination Mutualism." *Journal of Evolutionary Biology* 35

(1): 100–108.

Andreazzi, Cecilia S, John N Thompson, and Paulo R Guimaraes Jr. 2017. "Network Structure

and Selection Asymmetry Drive Coevolution in Species-Rich Antagonistic Interactions." *The*

American Naturalist 190 (1): 99–115.

Ashby, B., & Boots, M. (2015). Coevolution of parasite virulence and host mating

strategies. *Proceedings of the National Academy of Sciences*, *112*(43), 13290-13295.

Ashby, B., & Gupta, S. (2014). Parasitic castration promotes coevolutionary cycling but also

imposes a cost on sex. *Evolution*, *68*(8), 2234-2244.

Bánki, Olaf, Yury Roskov, Markus Döring, Geoff Ower, Diana Raquel Hernández Robles,

Camila Andrea Plata Corredor, Thomas Stjernegaard Jeppesen, et al. 2023. "Catalogue of Life

Checklist." Leiden, Netherlands: Catalogue of Life.

Bascompte, Jordi, Pedro Jordano, and Jens M. Olesen. 2006. "Asymmetric Coevolutionary

Networks Facilitate Biodiversity Maintenance." *Science* 312 (5772): 431–33.

Bernays, Elizabeth A. 2009. "Chapter 201 - Phytophagous Insects." In *Encyclopedia of Insects*

(Second Edition), edited by Vincent H. Resh and Ring T. Cardé, Second Edition, 798–800. San

Diego: Academic Press.

Brodie III, Edmund D, and Edmund D Brodie Jr. 1999. "Predator-Prey Arms Races:

Asymmetrical Selection on Predators and Prey May Be Reduced When Prey Are Dangerous."

Bioscience 49 (7): 557–68.

Bronstein, Judith L. 2015. "The study of mutualism." In *Mutualism*, 3–19. Oxford University
Press, Oxford.

Christenhusz, Maarten JM, and James W Byng. 2016. "The Number of Known Plants Species in
the World and Its Annual Increase." *Phytotaxa* 261 (3): 201–17.

Cogni, R., Quental, T. B., & Guimarães Jr, P. R. (2022). Ehrlich and Raven escape and radiate
coevolution hypothesis at different levels of organization: Past and future
perspectives. *Evolution*, *76*(6), 1108-1123.

Cook, James M., and Jean-Yves Rasplus. 2003. "Mutualists with Attitude: Coevolving Fig
Wasps and Figs." *Trends in Ecology & Evolution* 18 (5): 241–48.

Dawkins, R., & Krebs, J. R. (1979). Arms races between and within species. *Proceedings of the
Royal Society of London. Series B. Biological Sciences*, *205*(1161), 489-511.

Day, T., & Burns, J. G. (2003). A consideration of patterns of virulence arising from host-
parasite coevolution. *Evolution*, *57*(3), 671-676.

De Villemereuil, P., Charmantier, A., Arlt, D., Bize, P., Brekke, P., Brouwer, L., ... & Chevin, L.
M. (2020). Fluctuating optimum and temporally variable selection on breeding date in birds and
mammals. *Proceedings of the National Academy of Sciences*, *117*(50), 31969-31978.

Dobson, Andy, Kevin D. Lafferty, Armand M. Kuris, Ryan F. Hechinger, and Walter Jetz. 2008.
"Homage to Linnaeus: How Many Parasites? How Many Hosts?" *Proceedings of the National
Academy of Sciences* 105 (supplement_1): 11482–89.

Doebeli, M., & Dieckmann, U. (2003). Speciation along environmental
gradients. *Nature*, *421*(6920), 259-264.

Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N. P., ... & Allesina, S.
(2013). The dimensionality of ecological networks. *Ecology Letters*, *16*(5), 577-583.

Ehrlich, Paul R., and Peter H. Raven. 1964. "Butterflies and Plants: A Study in Coevolution."
Evolution 18: 586–608.

Endara, María José, Phyllis D. Coley, Gabrielle Ghabash, James A. Nicholls, Kyle G. Dexter,
David A. Donoso, Graham N. Stone, R. Toby Pennington, and Thomas A. Kursar. 2017.
"Coevolutionary arms race versus host defense chase in a tropical herbivore–plant system."
Proceedings of the National Academy of Sciences of the United States of America 114 (36):
E7499–7505.

Etienne, R. S., & Alonso, D. (2005). A dispersal-limited sampling theory for species and
alleles. *Ecology letters*, *8*(11), 1147-1156.

Farrell, B. D., & Sequeira, A. S. (2004). Evolutionary rates in the adaptive radiation of beetles on
plants. *Evolution*, *58*(9), 1984-2001.

Forister, M. L., & Jenkins, S. H. (2017). A neutral model for the evolution of diet breadth. *The*
American Naturalist, *190*(2), E40-E54.

Gilchrist, M. A., & Sasaki, A. (2002). Modeling host–parasite coevolution: a nested approach
based on mechanistic models. *Journal of Theoretical Biology*, *218*(3), 289-308.

Gingerich, P. (2019a). Temporal Scaling and Evolutionary Mode. In *Rates of Evolution: A*
Quantitative Synthesis (pp. 79-109). Cambridge: Cambridge University Press.

Gingerich, P. (2019b). Phenotypic Change in Experimental Lineages. In *Rates of Evolution: A*
Quantitative Synthesis (pp. 124-156). Cambridge: Cambridge University Press.

Hardy, Nate B., and Sarah P. Otto. 2014. "Specialization and generalization in the diversification
of phytophagous insects: tests of the musical chairs and oscillation hypotheses." *Proceedings of*
the Royal Society B: Biological Sciences 281 (1795): 20132960.

Hardy, N. B., Kaczvinsky, C., Bird, G., & Normark, B. B. (2020). What we don't know about
diet-breadth evolution in herbivorous insects. *Annual Review of Ecology, Evolution, and*
Systematics, *51*, 103-122.

Harmon, Luke J., Cecilia S. Andreazzi, Florence Débarre, Jonathan Drury, Emma E. Goldberg,
Ayana B. Martins, Carlos J. Melián, et al. 2019. "Detecting the macroevolutionary signal of
species interactions." *Journal of Evolutionary Biology* 32 (8): 769–82.

Hembry, David H., and Marjorie G. Weber. 2020. "Ecological interactions and macroevolution:
a new field with old roots." *Annual Review of Ecology, Evolution, and Systematics* 51: 215–43.

Hembry, David H., Jeremy B. Yoder, and Kari Roesch Goodman. 2014. "Coevolution and the
Diversification of Life." *The American Naturalist* 184 (4): 425–38.

Hubert, N., Calcagno, V., Etienne, R. S., & Mouquet, N. (2015). Metacommunity speciation
models and their implications for diversification theory. *Ecology Letters*, *18*(8), 864-881.

Huisman, Jisca, and Jarle Tufto. 2012. Comparison of non-gaussian quantitative genetic models
for migration and stabilizing selection. *Evolution* 66 (11): 3444–61.

Janz, Niklas. 2011. "Ehrlich and Raven Revisited: Mechanisms Underlying Codiversification of
Plants and Enemies." *Annual Review of Ecology, Evolution, and Systematics* 42: 71–89.

Kawahara, Akito Y, Caroline Storer, Ana Paula S Carvalho, David M Plotkin, Fabien L
Condamine, Mariana P Braga, Emily A Ellis, et al. 2023. "A Global Phylogeny of Butterflies
Reveals Their Evolutionary History, Ancestral Hosts and Biogeographic Origins." *Nature*
Ecology & Evolution, 1–11.

Kay, K. M., & Sargent, R. D. (2009). The role of animal pollination in plant speciation:
integrating ecology, geography, and genetics. *Annual Review of Ecology, Evolution, and*
Systematics, *40*, 637-656.

Kopp, M., & Matuszewski, S. (2014). Rapid evolution of quantitative traits: theoretical
perspectives. *Evolutionary Applications*, *7*(1), 169-191.

Langmore, Naomi E, Sarah Hunt, and Rebecca M Kilner. 2003. "Escalation of a Coevolutionary
Arms Race Through Host Rejection of Brood Parasitic Young." *Nature* 422 (6928): 157–60.

Lopez, S., Rousset, F., Shaw, F. H., Shaw, R. G., & Ronce, O. (2008). Migration load in plants:
role of pollen and seed dispersal in heterogeneous landscapes. *Journal of evolutionary*
biology, *21*(1), 294-309.

Margulis, L. (1998). *Symbiotic planet: a new look at evolution*. Basic books.

Maliet, O., Loeuille, N., & Morlon, H. (2020). An individual-based model for the eco-
evolutionary emergence of bipartite interaction networks. *Ecology Letters*, *23*(11), 1623-1634.

Manceau, M., Lambert, A. & Morlon, H. (2015). Phylogenies support out-of-equilibrium models
of biodiversity. Ecol. Lett., 18, 347–356.

McPeek, M. A. (2008). The ecological dynamics of clade diversification and community
assembly. *The American Naturalist*, *172*(6), E270-E284.

Microsoft Corporation and Steve Weston (2022). doParallel: Foreach Parallel Adaptor for the
'parallel' Package. R package version 1.0.17. https://CRAN.R-project.org/package=doParallel

Moran, P.A.P. (1962). *Statistical Processes of Evolutionary Theory*. Clarendon Press, Oxford,
UK.

Nathan, P., Economo, E. P., Guénard, B., Simonsen, A. K., & Frederickson, M. E. (2023).
Generalized mutualisms promote range expansion in both plant and ant partners. *Proceedings of*
the Royal Society B, *290*(2006), 20231083.

Ollerton, J. (2006). Biological barter": patterns of specialization compared across different
mutualisms. *Plant-pollinator interactions: from specialization to generalization. University of*
Chicago Press, Chicago, 411-435.

Parchman, Thomas L., and Craig W. Benkman. 2002. Diversifying coevolution between
crossbills and black spruce on Newfoundland. *Evolution* 56 (8): 1663–72.

Poulin, Robert, and Serge Morand. 2000. The diversity of parasites. *The Quarterly Review of*
Biology 75 (3): 277–93.

Prado, F., Sheih, A., West, J. D., & Kerr, B. (2009). Coevolutionary cycling of host sociality and
pathogen virulence in contact networks. *Journal of Theoretical Biology*, *261*(4), 561-569.

Raimundo, Rafael L. G., Jean P. Gibert, David H. Hembry, and Paulo R. Guimarães. 2014.
"Conflicting Selection in the Course of Adaptive Diversification: The Interplay Between
Mutualism and Intraspecific Competition." *The American Naturalist* 183 (3): 363–75.

Reijenga, B. R., Murrell, D. J., & Pigot, A. L. (2021). Priority effects and the macroevolutionary
dynamics of biodiversity. *Ecology Letters*, *24*(7), 1455-1466.

Romero, G. Q., & Benson, W. W. (2005). Biotic interactions of mites, plants and leaf
domatia. *Current Opinion in Plant Biology*, *8*(4), 436-440.

Rosindell, J., Harmon, L.J. & Etienne, R.S. (2015). Unifying ecology and macroevolution with
individual-based theory. Ecol. Lett., 18, 472–482.

Scheepens, J. F., Frei, E. S., & Stöcklin, J. (2010). Genotypic and environmental variation in
specific leaf area in a widespread Alpine plant after transplantation to different
altitudes. *Oecologia*, *164*, 141-150.

Schmid-Hempel, Paul. 2013. "18The diversity and natural history of parasites." In *Evolutionary*
Parasitology: The Integrated Study of Infections, Immunology, Ecology, and Genetics. Oxford
University Press.

Stork, Nigel E. 2018. "How Many Species of Insects and Other Terrestrial Arthropods Are There
on Earth?" *Annual Review of Entomology* 63 (1): 31–45.

Thompson, J. N. (1994). *The coevolutionary process*. University of Chicago press.

Thompson, John N. 2005. *The Geographic Mosaic of Coevolution*. University of Chicago Press.

Thompson, John N., Kari A. Segraves, and David M. Althoff. 2017. "Coevolution and
Macroevolution." In *Evolutionary Developmental Biology: A Reference Guide*, edited by Laura
Nuno de la Rosa and Gerd Müller, 1–13. Cham: Springer International Publishing.

Tufto, Jarle. 2000. "Quantitative Genetic Models for the Balance Between Migration and
Stabilizing Selection." *Genetics Research* 76 (3): 285–93.

Tufto, J. (2001). Effects of releasing maladapted individuals: a demographic-evolutionary
model. *The American Naturalist*, *158*(4), 331-340.

Vasco, D. A., Nazarea, A. D., & Richardson, R. H. (1987). Dynamics and stability in
coevolutionary ecological systems I. Community stability and coevolutionarily stable
states. *Theoretical population biology*, *31*(2), 273-305.

Vermeij, Geerat J. 1994. "The Evolutionary Interaction Among Species: Selection, Escalation,
and Coevolution." *Annual Review of Ecology and Systematics*, 219–36.

Weinstein, Sara B., and Armand M. Kuris. 2016. "Independent origins of parasitism in
Animalia." *Biology Letters* 12 (7): 20160324.

Whittall, Justen B, and Scott A Hodges. 2007. "Pollinator Shifts Drive Increasingly Long Nectar
Spurs in Columbine Flowers." *Nature* 447 (7145): 706–9.

Yamamichi, M., Lyberger, K., & Patel, S. (2019). Antagonistic coevolution between multiple
quantitative traits: Matching dynamics can arise from difference interactions. *Population
Ecology*, *61*(4), 362-370.

Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation under migration–
selection balance. *Evolution*, *65*(7), 1897-1911.

Yoder, Jeremy B., and S. L. Nuismer. 2010. "When does coevolution promote diversification?"
American Naturalist 176 (6): 802–17.

Zacchello, Giulia, Mariona Vinyeta, and Jon Ågren. 2020. "Strong Stabilizing Selection on
Timing of Germination in a Mediterranean Population of Arabidopsis Thaliana." *American
Journal of Botany* 107 (11): 1518–26.