OCA-B promotes autoimmune demyelination through control of stem-like CD4+ T cells

Erik P. Hughes[1,3], Amber R. Syage[1,3], Elnaz Mirzaei Mehrabad[2,3,4], Thomas E. Lane[5], Benjamin T. Spike[2,3], Dean Tantin[1,3]

[1]Department of Pathology, University of Utah School of Medicine, Salt Lake City, UT 84112 U.S.A.

[2]Department of Oncological Sciences, University of Utah School of Medicine, Salt Lake City, UT 84112 U.S.A.

[3]Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT 84112 U.S.A.

[4]School of Computing, University of Utah, Salt Lake City, UT, 84112 U.S.A

[5]Department of Neurobiology and Behavior, School of Biological Sciences, University of California, Irvine, CA 92697, U.S.A.

e-mail dean.tantin@path.utah.edu

Short title: OCA-B promotes CD4+ T cell mediated demyelinating disease

Keywords OCA-B/Bob1/Pou2af1; autoimmunity; multiple sclerosis; EAE; stem-like CD4+ T cells; relapsing-remitting

Words in abstract: 166

Words main text: 4,648

Total main display items: 6

Summary

Stem-like T cell populations can selectively contribute to autoimmunity, but the activities that

promote and sustain these populations are incompletely understood. Here, we show that T cell-

intrinsic loss of the transcription cofactor OCA-B protects mice from experimental autoimmune

encephalomyelitis (EAE) while preserving responses to CNS infection. In adoptive transfer EAE

models driven by multiple antigen encounters, OCA-B deletion nearly eliminates CNS infiltration,

proinflammatory cytokine production and clinical disease. OCA-B-expressing CD4$^+$ T cells within

the CNS of mice with EAE comprise a minority of the population but display a memory phenotype

and preferentially confer disease. In a relapsing-remitting EAE model, OCA-B T cell deficiency

specifically protects mice from relapse. During remission, OCA-B promotes the expression of

Tcf7, *Slamf6*, and *Sell* in proliferating T cell populations. At relapse, OCA-B loss results in both

the accumulation of an immunomodulatory CD4$^+$ T cell population expressing *Ccr9* and *Bach2*,

and the loss of pro-inflammatory gene expression from Th17 cells. These results identify OCA-B

as a driver of pathogenic stem-like T cells.

Introduction

2.5 million individuals worldwide suffer from multiple sclerosis (MS), with almost 1 million cases in the United States[1]. Despite recent advances, the development of new therapies that block MS while sparing immunosurveillance mechanisms needed to prevent infection and viral recrudescence remains a critical goal. Genome-wide association studies (GWAS) identify the *HLA* loci as the most important determinant of MS pathogenesis, emphasizing the impact of T cells in MS[2,3]. The association with the major histocompatibility complex (MHC) class II locus in particular indicates that myelin-specific CD4$^+$ T cells are key regulators of disease.

The predominant preclinical model of MS is experimental autoimmune encephalomyelitis (EAE), in which laboratory animals are immunized with myelin-specific proteins or peptides to activate antigen specific CD4$^+$ T cells that infiltrate the central nervous system (CNS) and mediate disease. EAE within the C57BL/6 mouse background establishes chronic disease. In these models, Th1 and Th17 effector CD4$^+$ T cell subsets are pathogenetic mediators[4,5]. After accessing the CNS, these cells promote neuroinflammation through the secretion of interferon-gamma (IFNγ), interleukin-17 (IL-17), granulocyte macrophage stimulating-factor (GM-CSF) and other inflammatory mediators and chemoattractants that recruit the cell types responsible for driving demyelination[6–8].

Myelin-reactive T cells from the blood of MS patients show stem/memory-like properties, with reduced need for co-stimulation, suggesting signs of previous activation[9–12]. Consistent with these findings, autoreactive and bystander memory-phenotype CD4$^+$ T cells preferentially transfer demyelinating disease in mouse models[13–15]. While these studies have identified vital CD4$^+$ T cell populations in EAE, further research on the mechanisms creating and sustaining these populations is required to develop more precise therapies.

Oct1, a POU domain transcription factor encoded by *Pou2f1*, has been linked to autoimmune states. GWAS studies have identified Oct1 binding site polymorphisms that associate with a predisposition for autoimmune diseases, including MS[16–19]. In CD4$^+$ T cells, Oct1 promotes the development of both CD4$^+$ memory[20] and EAE pathogenesis and demyelination[21]. In contrast, Oct1 loss preserves immune responses to the neurotrophic viral infection[21]. These observations highlight a promising therapeutic window in which targeting this pathway may limit autoimmune CNS demyelination without the broad immune suppression elicited by many current therapeutics. However, Oct1 is widely expressed and regulates both embryonic development and somatic stem cell function[22–24], making it a poor therapeutic target.

The lymphoid-restricted transcription cofactor OCA-B regulates transcription by docking with Oct1, and in B cells an additional related factor, Oct2[25]. In contrast to Oct1, OCA-B whole-

body knockouts are viable and fertile[26]. Polymorphisms in the *Ocab* (*Pou2af1*) locus are

associated with autoimmune diseases, including MS[27,28]. Although its expression is 50-100 fold

lower in T cells compared to B cells[29], targeting either OCA-B by genetic deletion in T cells or its

downstream pathways with a membrane-permeable peptide mimic suppresses type-1 diabetes[31].

OCA-B interacts with target genes such as *Il2, Ifng, Il21, Csf2 (Gmcsf), Tnfrsf4 (Ox40), Icos* and

Ctla4, but is dispensable for their expression following primary activation of T cells. Instead, OCA-

B regulates these targets under a narrow range of conditions, for example following secondary

stimulation in culture[20]. Many of these targets are directly implicated in MS pathogenesis[6–8,31–37].

OCA-B controls expression of *Tbx21* (*Tbet*), which encodes a key regulator of Th1 effector

responses, but also promotes Th17 differentiation and interacts with RORγt, a master regulator

of Th17 cells[20,38,39]. In the context of MS, the only published OCA-B work used whole body

knockouts and showed measurable but incomplete protection in chronic C57BL/6 EAE models[39].

Cumulatively, these findings indicate that OCA-B plays poorly defined but possibly significant

roles in MS pathogenesis.

Here, we show that OCA-B within the T cell compartment drives EAE by promoting

pathogenic maturation of stem-like CD4[+] T cells. In C57BL/6 mice, T cell OCA-B loss significantly

protects mice from chronic EAE, while preserving responses to the neurotrophic coronavirus

JHMV. Using Th1 and Th17 passive transfer assays in which reactive T cells encounter antigen

in donor mice, in polarizing culture, and again in naïve recipients, we show that OCA-B is essential

for the infiltration of pathogenic cells into the CNS and the manifestation of clinical disease. Using

an OCA-B reporter mouse, we find that OCA-B expressing CD4[+] T cells taken from MOG peptide

primed mice display pathogenic stem-like gene expression and preferentially transfer

demyelinating disease, highlighting OCA-B as a marker and promoter of encephalitogenic

autoimmune CD4[+] T cell activity. Intriguingly, T cell conditional OCA-B knockout In EAE models

on the autoimmune-prone non-obese diabetic (NOD) background, which develops a relapsing-

remitting disease[40], specifically protects against relapse. Single-cell transcriptomic analysis of

CD3ε[+] CNS-infiltrating T cells at remission and relapse indicates that OCA-B promotes disease

relapse through control of stem-like T cell populations, driving them to pathogenic Th17

differentiation. Cumulatively, these findings mark OCA-B as a critical mediator of the pathogenic

maturation of autoreactive T cells, and a promising potential therapeutic target.

Results

OCA-B promotes chronic EAE in C57BL/6 mice

To understand the impact of OCA-B in T cell mediated autoimmunity, we utilized an *Ocab*

(*Pou2af1*) conditional knockout allele crossed to the CD4-Cre driver which efficiently deletes

OCA-B in T cells[30]. We induced EAE in C57BL/6 female *Ocab*[fl/fl];CD4-Cre mice and littermate

Ocab[fl/fl] controls through subcutaneous (s.c.) injection of myelin oligodendrocyte glycoprotein

(MOG)$_{35-55}$ peptide in Complete Freund's Adjuvant (CFA) followed by intraperitoneal (i.p.)

injections of *Bordetella pertussis* toxin. *Ocab*[fl/fl];CD4-Cre mice showed reduced chronic EAE

clinical scores and weight loss compared to *Ocab*[fl/fl] littermate controls (Fig. 1a, b). Spinal cords

were taken at peak disease (day 15) for immunofluorescence and histological analysis. Myelin

basic protein (MBP) immunofluorescence showed a significant decrease in demyelinated white

matter in spinal sections from Ocab[fl/fl];CD4-Cre mice compared to littermate Ocab[fl/fl] controls while

DAPI showed decreased immune cell infiltration (Fig. 1c, d). Similarly, Ocab[fl/fl];CD4-Cre mice

showed a small but significant decrease in demyelination as measured by luxol fast blue (LFB) in

combination with hematoxylin and eosin (H&E) staining (Extended Data Fig. 1a, b). Peak disease

flow cytometric profiling showed decreased CD4+ T cell infiltration into the CNS of *Ocab*[fl/fl];CD4-

Cre mice, however no differences in other infiltrating cell populations or in the relative proportions

of the proinflammatory cytokines IFNγ, IL-17, GM-CSF, or CD44 were observed (Fig. 1e, f, and

Extended Data Fig. 1c-h). As no differences in the frequency of cytokine expression were

observed across total T cell populations, we studied MOG-specific CD4+ T cells using the MHC

class II I-A^b MOG$_{38-49}$ tetramers. At peak disease, *Ocab*[fl/fl];CD4-Cre mice showed a trending

reduction in CD44+ MOG tetramer-specific cell frequency (p=0.0637) and a significant reduction

in numbers (p=0.003) compared to controls (Fig. 1g, h). These results indicate that the protection

conferred by OCA-B loss in T cells is marked by modest decreases in MOG-specific CD4+ T cells

within the CNS, resulting in fewer pathogenic T cells driving demyelination.

OCA-B is essential for EAE mediated by adoptive transfer of Th1- and Th17-polarized T

cells

MS is driven by autoreactive CD4+ Th1 and Th17 cells within the CNS that recruit CD8+ T cells,

macrophages, microglia and neutrophils, which together mediate white-matter damage[41]. To

investigate the impact of OCA-B on autoimmune Th1 and Th17 populations, we used adoptive

transfer EAE models with female *Ocab*[fl/fl];CD4-Cre and *Ocab*[fl/fl] littermate controls. Mice were

primed with MOG$_{35-55}$ in CFA for 10-14 days. Following MOG priming, no differences between

genotypes in CD4[+] T cell viability, count, or effector cytokine (IFNγ and IL-17) expression were

observed (Extended Data Fig. 2a-e). Interestingly, CD4[+] T cells from Ocab[fl/fl];CD4-Cre mice

showed a modest increase in the percentage of cells expressing Ki67 compared to Ocab[fl/fl]

littermate controls (Extended Data Fig. 2f). To determine differences in adoptive transfer viability

and proliferation, CD4[+] T cells from MOG$_{35-55}$ primed Ocab[fl/fl] control and Ocab[fl/fl];CD4-Cre

experimental donors were labeled with fluorescent quantum dots (Ocab[fl/fl]-QTracker 655 and

Ocab[fl/fl];CD4-Cre-QTracker 525), combined 1:1 and transferred into naïve recipients. Three days

following mixed transfer, cell were recovered from the peritoneal cavities. The frequency of

Ocab[fl/fl];CD4-Cre donor CD4[+] T cells in the peritoneal cavity was decreased and showed

diminished viability compared to control donor CD4[+] T cells (Extended Data Fig. 2g-j). Similar to

cells collected before transfer, CD4[+] T cells from Ocab[fl/fl];CD4-Cre donors showed a significant

increase in Ki67 expression (Extended Data Fig. 2k, l). The combination of decreased numbers

and increased proliferation in transferred OCA-B knockout T cells may reflect preferential homing

of these cells to tissues such as the gut.

To investigate if OCA-B expression impacts Th1 or Th17 mediated adoptive transfer EAE,

CD4[+] T cells were purified from the lymph nodes of primed experimental or control donors and

polarized in vitro into Th1 and Th17 cells as previously described[42,43]. 3×10^6 Th1- or Th17-

polarized CD4[+] cells were transferred into naïve age-matched wild-type male C57BL/6 recipients

by i.p. injection. EAE onset in mice receiving either Th1- or Th17-polarized control cells was rapid

and severe (Fig. 2a, b). In contrast to the mild protection observed with OCA-B deficiency in

standard C57BL/6 EAE, mice adoptively transferred with T cells lacking OCA-B were almost

completely protected from clinical disease regardless of their polarization state. Interestingly, the

ability to initially polarize CD4[+] T cells into Th1 and Th17 effectors in vitro was minimally affected

by OCA-B loss (Extended Data Fig. 2m-o), whereas the same cells taken from the CNS of Th17-

engrafted recipient mice at day 15 not only showed decreased CNS infiltration but also poor IL-

17 expression (Fig. 2c-f). This effect appeared to be specific to IL-17, as no significant difference

in IFNγ expression was observed (Extended Data Fig. 2p) These results indicate a central role for

OCA-B in promoting robust neuroinflammatory T cell responses upon antigen reencounter in

adoptive transfer EAE models.

OCA-B is dispensable for neurotropic viral clearance

To investigate if T cell OCA-B deletion impacts the ability to mount immune responses and control

viral infection in the CNS, male *Ocab*fl/fl;CD4-Cre and *Ocab*fl/fl littermate controls were intracranially

inoculated with 1500 plaque-forming units (PFU) of the John H. Muller strain of murine hepatitis

virus (JHMV), a β-coronavirus adapted to infect glial cells (astrocytes, microglia and

oligodendrocytes) in the mouse brain and spinal cord. Inoculation with JHMV results in acute

encephalomyelitis and chronic demyelination similar to autoantigen-driven EAE but without

persistent autoantigen. Disease severity was determined by evaluating JHMV clinical scores over

21 days post-infection (dpi). *Ocab*fl/fl;CD4-Cre animals showed negligible differences in clinical

scores compared to *Ocab*fl/fl littermates (Fig. 3a), suggesting that OCA-B loss preserves baseline

T cell responses to acute infection. Spinal cords were taken on 12 and 21 dpi for

immunofluorescence and histological analysis to investigate differences in demyelination. MBP

immunofluorescence at 12 dpi showed similar demyelination and immune infiltration in both

Ocabfl/fl and Ocabfl/fl;CD4-Cre spinal sections (Fig. 3b, c). Similarly, LFB histology of brain tissue

at 21 dpi showed no significant differences in demyelination between OCA-B deficient and control

infected animals (Extended Data Fig. 3a, b). Flow cytometric analysis of brain tissue showed no

significant differences in infiltrating CD4$^+$ T cell numbers or their ability to express IFNγ at 7, 12,

and 21 dpi (Fig. 3d-f, and Extended Data Fig. 3c). Similarly, no differences in either the total

numbers of CD8$^+$ T cells, or the percent expression or numbers of IFNγ expressing CD8$^+$ T cells

was observed (Extended Data Fig. 3d-f). Brain homogenates were collected on 7, 12 and 21 dpi

to measure viral titers and clearance. *Ocab*fl/fl;CD4-Cre and *Ocab*fl/fl littermate control animals

showed similar titers at dpi 7 and 12, and virus was undetectable at day 21 regardless of genotype

(Fig. 3g). This result suggests that OCA-B is dispensable for immune responses to and clearance

of neurotropic viruses.

OCA-B expressing CD4$^+$ T cells display pathogenic stem-like properties and preferentially

transfer demyelinating disease

We used a recently described an OCA-B reporter (OCA-B-mCherry) mouse which selectively

labels OCA-B expressing B and T cells with mCherry[29] to evaluate OCA-B expression within CNS

infiltrating T cells. EAE was induced in female C57BL/6 homozygous OCA-B-mCherry reporter

mice with MOG$_{35-55}$, and CNS-infiltrating CD4$^+$ T cells assessed by flow cytometry at peak clinical

disease. mCherry-positive cells showed increased CD44 and CD62L co-expression, indicative of

memory-like properties in the OCA-B expressing T cell population (Fig. 4a). The association of

OCA-B expression with a memory-like profile was observed in the brain, spine and draining cervical lymph nodes (Fig. 4b). Interestingly, the frequency of CD44$^+$CD62L$^-$CD4$^+$ T cells was elevated within the OCA-B expressing population specifically within the cervical lymph nodes (Extended Data Fig. 4a).

Memory-like CD4$^+$ T cells have been implicated as major orchestrators of autoimmune demyelination in mice and humans[9–12,14,15,44]. To see if this memory-like profile of OCA-B expressing CD4$^+$ T cells is specific to CNS infiltrating cells, we performed flow cytometry using draining lymph node and splenic T cells from MOG-primed OCA-B-mCherry reporter mice. Similar to OCA-B$^+$ CNS infiltrating cells, OCA-B expressing CD4$^+$ T cells taken from the cervical lymph node and spleen showed increased CD44 and CD62L co-expression (Fig. 4c, d).

We next evaluated OCA-B-mCherry positive and negative populations by RNA sequencing of CD4$^+$ T cells isolated from the lymph nodes of 14-day MOG-primed mice. ~1500 genes were upregulated and ~350 downregulated in OCA-B expressing compared to OCA-B negative cells (Supplementary Table 1). Upregulated genes in OCA-B expressing cells included *Cxcr6*, *Slamf6*, *Il23r*, *Rorc*, *Ifng*, and *Il17a* while downregulated genes included *Foxp3*, *Il7r* and *Ccr9* (Fig. 4e, f). Expression of *Tcf7* (which encodes TCF1) was high within both OCA-B positive and negative populations (Supplementary Table 1). Gene ontology (GO) and pathway analysis of upregulated genes show enrichment for terms associated with autoimmune diseases including lupus, autoimmune thyroid disease, and type-1 diabetes (Fig. 4g, and Supplementary Table 2).

To determine if OCA-B expression can be used to prospectively identify viable pathogenic CD4$^+$ T cell populations, we used OCA-B-mCherry reporter mice as donors in passive transfer EAE. 11-week-old male and female littermates homozygous for the OCA-B-mCherry reporter were primed for 14 days with MOG$_{35-55}$ in CFA as previously described[43]. CD4$^+$ T cells were isolated from spleen and lymph nodes and sorted based on mCherry expression (Extended Data Fig. 4b). 9×10^5 mCherry negative and positive cells were cultured in Th1 polarizing conditions and subsequently transferred into age-matched male C57BL/6 recipients as previously described[14]. Disease progression was measured by clinical score and normalized weight. Recipients of OCA-B-expressing mCherry positive cells showed increased clinical disease severity and weight loss compared to mCherry negative recipients (Fig. 4h, and Extended Data Fig. 4c). These findings indicate that OCA-B marks CD4$^+$ T cells with pathogenic stem-like gene expression and can be used to prospectively identify peripheral CD4$^+$ T cells with elevated encephalitogenic properties.

OCA-B promotes neuroinflammation in relapsing-remitting EAE models via control of stem/memory-like T cell populations

In contrast to the chronic form of EAE that develops in C57BL/6 mouse models, EAE on the NOD background exhibits a relapsing-remitting pattern that more closely mimics human disease[40]. To study the impact of OCA-B loss in T cells on mouse models that more closely resemble human MS disease patterns, we utilized *Ocab* T cell conditional knockout mice crossed to the NOD background[30]. EAE was induced in 15-18 week old female NOD.*Ocab*^{fl/fl};CD4-Cre and littermate control NOD.*Ocab*^{fl/fl} mice with MOG$_{35-55}$. Clinical scores and weights were recorded for 47 days to observe initial disease onset, relapse and relapse recovery. No significant differences in clinical scores were observed between groups at disease onset (day 10-19), however following the primary remission, NOD.*Ocab*^{fl/fl};CD4-Cre animals were almost completely protected against disease relapse (Fig. 5a). Individual NOD mouse clinical scores are shown in Extended Data Fig. 5a-b. Normalized weights of animals also reflected the protective effect of OCA-B loss in T cells at relapse (Fig. 5b).

To understand differences in CNS-infiltrating T cells underlying relapse protection at a population and transcriptional level, we performed single-cell RNA sequencing (scRNAseq) of CD3ε^+ T cell isolates from CNS (brain + spinal cord) at remission (day 24) and peak relapse (day 33). Cells from 3-6 NOD.*Ocab*^{fl/fl};CD4-Cre and NOD.*Ocab*^{fl/fl} littermate controls were collected and combined for analysis. 7799 (remission) and 9224 (relapse) *Ocab* knockout, and 6009 (remission) and 7525 (relapse) control cells were used for analysis, with an average per cell read depth ranging from 90,000 to 190,000. For remission, Uniform Manifold Approximation and Projection (UMAP) clustering using both genotypes revealed 16 clusters corresponding to a broad T cell repertoire consisting of natural-killer T (NKT) cells (the largest cluster) and different CD4$^+$ and CD8$^+$ subsets (Fig. 5c). Noteworthy clusters based on mRNA expression included proliferating (*Ki67*, *Pcna*, *Mcm2*), Th1-like (*Tbx21*, *Ifng*, *Tnf*, *Il12rb2*) and Treg (*Foxp3*, *Ctla4*) (Supplementary Table 3). Feature plots for *Foxp3* and *Mki67* are shown in Fig. 5d. Additional feature plots for *Cxcr6*, *Il23r* and *Bach2* are shown in Extended Data Fig. 5c. Notably there was a large decrease in NKT cell numbers in knockout mice (36% control vs. 14% experimental). Within this cluster, changes in gene expression between control and conditional knockout groups were minimal. Most NKT cells from both groups showed a limited inflammatory cytokine response, consistent with the lack of clinical disease at remission (Extended Data Fig. 5d). The decrease in the large NKT population in the experimental group proportionally increases the percentage contribution of the

other clusters, and therefore similar or decreased percentages of these cells identifies reduced

populations which correlate with protection from relapse, e.g. proliferating T cells. Other clusters

such as Tregs increased disproportionately. Differential gene expression analysis of key clusters

(Treg, Proliferating, and Th1-like) revealed increased expression of *Slamf6* in Tregs lacking OCA-

B, while OCA-B deficient proliferating and Th1-like cells showed decreased expression of genes

associated with quiescence (*Btg1* and *Samhd1*), memory/stemness (*Slamf6*, *Tcf7*, and *Sell*) and

pathogenicity (*Tcf7* and *Tox* co-expression, Fig. 5e). Spectral cytometry showed no significant

difference in the frequency of CNS infiltrating CD4$^+$ T cells between groups at remission (Fig. 5f).

However, OCA-B deficient CD4$^+$ T cells did show a significant reduction in CXCR6 and IFNγ/GM-

CSF co-expression, indicative of reduced Th17 pathogenicity (Fig. 5g-i). These results indicate

that loss of OCA-B in T cells selectively reduces proliferation and differentiation of pathogenic

stem/memory-like CD4$^+$ T cells during disease remission.

At relapse, multiple CNS-infiltrating stem-like and effector CD4$^+$ and CD8$^+$ control T cell

populations were present (Fig. 6a and Extended Data Fig. 6a; Supplementary Table 4). TCR

sequencing revealed an expansion of a subset of TCR clones in the OCA-B deficient CD8 clusters

while most other clusters were highly polyclonal (Fig. 6b). Comparing the 10 most expanded TCR

clonotypes between knockouts and controls, there were several clonotypes that were shared but

differentially expanded (Extended Data Fig. 6b and Supplementary Table 5). A "Ccr9 stem-like"

cluster expressing the gut homing chemokine receptor *Ccr9*, as well as *Ccr7*, *Bach2*, and *Tcf7*

notably appeared at relapse (Fig. 6c, Supplementary Table 4). TCR clonotype analysis indicated

that this population was highly polyclonal (Fig. 6b). Additionally, a "Stem-like CD4-2" cluster

expressing *Tcf7*, Ccr7, *Bach2, Cd44* and low-level *Cxcr6*, and a prominent "Th17-like" cluster

expressing *Tcf7, Cd44*, *Il1r1, Lgals3*, and higher levels of *Cxcr6* was identified (Fig. 6c). This latter

cluster likely contains pathogenic effector cells.

In the absence of OCA-B, NKT cell populations were normalized compared to the

remission timepoint, and several clusters showed major changes in the percentage of cells

including most prominently the Ccr9-stem-like cells which were much more prevalent (Fig. 6d).

Feature plots comparing OCA-B T cell deficient mice to littermate controls for *Ccr9* and the

transcription factors *Tcf7* and *Bach2* are shown in Fig. 6d. The cells were also marked by the

expression of *Cd4*, *Tox, Ccr7, Ccr4* and limited expression of *Sell* (Extended Data Fig. 6c).

RNA velocity analysis of spliced and unspliced transcripts[45] identified clusters of T cells,

most bearing static gene expression profiles consistent with terminal differentiation, but some with

differentiating or transitional gene expression (Fig. 6e). Notably, OCA-B knockout cells showed

increased transcriptional stasis within the Ccr9-stem-like and proliferating clusters, and a

decreased transcriptional directionality of Th17 cells towards gd17 cells, the latter of which

express high levels of pathogenic markers including *Il1r1*, *Cxcr6*, and *Lgals3* (Extended Data Fig.

6a). Pseudotime analysis identified a transcriptional capacity in the control *Ccr9* expressing

population to differentiate into pathogenic Th17-like cells, whereas cells lacking OCA-B were

limited in this capacity, possibly resulting in the accumulation of these cells (Extended Data Fig.

6d). These changes culminated in the Th17-like effector knockout cells expressing reduced

pathogenicity-associated genes such as *Pdcd1*, *Tigit* and *Il1r1* (Fig. 6f). Together these results

suggest OCA-B mediates the transition of non-pathogenic stem-like cells to pathogenic Th17 cells

which ultimately drive disease relapse.

Discussion

The development of therapies that inhibit autoimmunity while minimizing impacts on normal

immune function remains a critical unmet goal in the field. In this study, we show that expression

of the transcriptional co-regulator OCA-B promotes a pathogenic, stem/memory-like function in T

cells, resulting in the establishment of CNS autoimmunity, and in particular episodes of relapse.

Loss of OCA-B attenuates these populations in chronic EAE, in adoptive transfer EAE, and in

NOD.EAE during episodes of remission. During relapse, OCA-B T cell knockout mice are almost

completely protected, and accumulate a population of CD4+ T cells expressing *Ccr9*, *Bach2*, and

Tcf7. OCA-B expression also marks viable pathogenic stem/memory-like CD4+ T cells that

preferentially transfer EAE. Furthermore, the effects of modulating OCA-B are selective, as

responses to infection with a neurotropic coronavirus that generates similar clinical symptoms but

lacks an autoantigen were largely unaffected.

OCA-B is a member of a small family of transcription co-regulators that also includes OCA-

T1, OCA-T2 and IκBζ[46,47]. It docks with Oct1 upon CD4+ T cell activation to directly regulate

immunomodulatory genes such as *Il2*, *Ifng*, *Il17a*, Tbx21 (T-bet) and *Csf2* (*Gmcsf*). Rather than

functioning as a primary inducer of these genes, it is required for their robust expression upon

antigen reencounter[20,30]. Notably, unlike transcription factors such as NF-AT, AP-1 and NF-κB,

Oct1 and OCA-B are dispensable for normal primary T cell responses[20]. These findings predict

important roles for OCA-B in CD4+ T cells specifically in situations of antigen reencounter, which

is a necessary feature of autoimmunity. Conditional T cell knockout of Oct1 reduces EAE severity,

with reduced proinflammatory cytokine expression and increased anergy[21]. Deletion of Oct1 in T

cells was dispensable for neurotropic viral clearance[21], however the ubiquitous expression of Oct1

in adult tissue limits its translational potential. In contrast, OCA-B expression is largely confined

to B and T cells and whole-body knockout mice are viable and fertile[48]. OCA-B is highly expressed

in B cells and required for germinal center formation[49,50]. It is expressed in CD4+ T cells upon

activation at levels 50-100-fold less than in B cells[30,51,52]. This suggests that with proper

therapeutic dosing, OCA-B could be inhibited selectively in T cells with minimal impact to the B

cell compartment. Short-term administration of an OCA-B peptide mimic that inhibits OCA-B's

downstream effector functions reverses spontaneous type-1 diabetes in NOD mice[30]. However,

toxicity limits its efficacy, including within EAE models. Further research is required for the

development of safe and effective OCA-B inhibitors.

OCA-B whole body knockout animals are partially protected from EAE[39], however, the
known activity of OCA-B in B cells made it unclear if the protection is T cell-derived. Here we
show that OCA-B promotes EAE pathogenesis through T cells. Opportunistic infections and
reactivation of latent CNS-tropic viruses are significant barriers for MS therapeutics that focus on
global immune suppression as a therapeutic mechanism. When OCA-B T cell conditional
knockout mice are challenged with the neurotropic coronavirus JHMV, there is little difference in
their ability to mount immune responses and control virus replication. This finding provides a
potential therapeutic window in which targeting the low levels of OCA-B in T cells could be used
to blunt autoimmunity while sparing baseline immune responses.

Unlike the significant but modest protection observed in chronic C57BL/6 EAE, OCA-B
loss in adoptive transfer EAE almost completely protects recipient animals from disease. This
may indicate that adoptive transfer EAE models involve more autoantigen re-exposures, as
transferred T cells must re-encounter antigen in a naïve host before promoting disease. Protection
was observed regardless of whether donor cells were Th1- or Th17-polarized, indicating that
OCA-B functions through a mechanism that parallels rather than enforces the Th1/Th17
paradigm. We therefore sought other mechanisms that might explain why OCA-B T cell knockouts
are so strongly protected.

Recent findings indicate an emerging role for stem/memory-like T cells in promoting
autoimmunity[44,53]. For example, myelin-specific CD4$^+$ T cells in MS patients show signs of a
previously activated, memory-like phenotype compared to healthy controls[9,10,12]. Furthermore,
memory-like Th1 and Th17 cells are increased in the peripheral blood of MS patients and display
increased pathogenicity compared to conventional Th17 Cells[54]. Similarly, in EAE models,
peripheral central memory-like CD4$^+$CD44^{hi}CD62L^{hi} T cells confer increased disease severity
upon transfer[14]. Both Oct1 and OCA-B promote the functionality of long-lived CD4$^+$ memory T
cells following antigenic rechallenge[29,55]. An OCA-B-mCherry reporter mouse allele, which
effectively labels memory progenitor CD4$^+$ T cells during acute infection[29], also labels CD4$^+$ T
cells that preferentially confer EAE. OCA-B^{hi} cells taken from the CNS at peak EAE selectively
express stem/memory markers. Further profiling OCA-B^{hi} CD4$^+$ T cells suggests that OCA-B
promotes the transition of non-pathogenic stem-like cells to a pathogenic state associated with
markers such as *Cxcr6*, *Slamf6*, *Ifng*, and *Il17a*. The connection between OCA-B expression and
pathogenic memory/stem-like T cells likely has not been appreciated before because OCA-B is
expressed at levels too low to be reproducibly captured with many current single-cell RNA-seq
technologies.

Relapsing-remitting EAE models have been developed using the Swiss Jim Lambert (SJL)

and autoimmune prone non-obese diabetic (NOD) backgrounds[40,56]. In NOD mice, MOG$_{35-55}$ EAE

has been described as relapsing-remitting with minimal initial disease that increases in severity

upon relapse[57,58]. Using this model with the conditional *Pou2af1* allele backcrossed to the NOD

background[30], we found that OCA-B T cell loss minimally affects initial clinical presentation, but

instead protects animals specifically at relapse. The finding that OCA-B loss in conventional

C57BL/6 EAE results in partial protection intermediate between the initial and relapse phases of

NOD.EAE suggests that chronic C57BL/6 EAE may not model either initial presentation or relapse

accurately, but instead represents a superposition of the two.

Single-cell RNA expression profiling at remission identifies a significant loss of *Slamf6*,

Tcf7 (encoding TCF1) and *Sell* from proliferative stem-like CD4$^+$ T cells in the OCA-B deficient

condition. While the majority of research on TCF1-expressing stem-like T cells focuses on CD8$^+$

populations with the capacity for self-renewal and differentiation into effector cells, analogous

descriptions have been made for CD4$^+$ populations[44,59,60]. Indeed, autoimmune CD4$^+$ T cells

expressing these memory/stem-like markers have been reported as an encephalitogenic

reservoir that play a key role in EAE pathogenicity[44]. In the same study, bulk RNA sequencing

associated OCA-B expression within this stem-like reservoir[44].

Single-cell RNA-seq at relapse identified multiple emergent populations of effector T cells

compared to remission. Notably, there was also an accumulation of a unique population of CD4$^+$

cells in OCA-B T cell deficient mice. These cells are conspicuously marked by expression not

only of *Ccr4*, but also of *Ccr9*, which encodes a gut-homing chemokine receptor. CCR9$^+$ memory-

like CD4$^+$ T cells have been linked to MS and EAE protection[61]. The cells also express *Bach2*, a

transcription regulator which was recently associated with non-pathogenic immunomodulatory

Th17-polarized cells in the context of C57BL/6 EAE[62]. *Bach2* expression was largely concentrated

in the "CCR9 stem-like" cluster, which is more numerous in the knockout condition. Consistently,

in bulk RNA-seq *Bach2* mRNA was upregulated in OCA-B$^-$ cells ($p<10^{-13}$), although with a small

(+1.74) fold change (Supplementary Table 1). In B cells, OCA-B is known to regulate *Bach2*[63].

Velocity and pseudotime analysis indicates that these cells are normally transcriptionally inclined

to progress towards an highly pathogenic Th17 population, but that progression stalls in mice

whose T cells lack OCA-B. Consistently, the Th17 effector population in OCA-B T cell knockout

mice lacks expression of key genes such as *Il1r1* and *Pdcd1*, both of which are associated with

pathogenicity in CNS-infiltrating Th17 cells[48,64]. The cumulative alterations in these populations in

OCA-B knockouts likely explains why these mice are strongly protected against disease relapse.

In conclusion, this study indicates a prominent role for OCA-B in driving stem-like CD4$^+$ T cell

populations towards a pathogenic state while playing a minimal role in acute antiviral responses

to a CNS infection.

Figure legends

Fig.1 l OCA-B loss protects animals from chronic EAE

a, *Ocab*$^{fl/fl}$ (n=16) and *Ocab*$^{fl/fl}$;CD4-Cre (n=22) mice were injected with MOG$_{35-55}$ peptide in CFA and *pertussis* toxin to induce EAE. Clinical scores were evaluated after EAE induction to determine disease progression. **b**, Animal weights were recorded to evaluate weight loss as a measurement of disease progression. **c**, Representative DAPI/MBP immunostaining of thoracic sections taken from mice 15 days after EAE induction. Areas of demyelination are marked by decreased MBP staining (green) and often coincide with increased cellular infiltration (blue). **d**, Quantification of demyelination in experimental (n=6) and control mice (n=5). **e**, Brain and spinal cords were isolated on day 15 of EAE and analyzed by spectral cytometry. Representative flow cytometry plots of CD45$^+$TCRβ$^+$ CNS infiltrating T cells showing frequency of CD4$^+$ and CD8$^+$ T cells. **f**, Quantification of the number of CNS infiltrating CD4$^+$ T cells. *Ocab*$^{fl/fl}$ (n=13) and *Ocab*$^{fl/fl}$;CD4-Cre (n=9). **g**, Representative plots showing frequency of CD44$^+$ MOG$_{38-49}$ tetramer-positive CD4$^+$ T cells. **h**, Quantification of the frequency and count of CD44$^+$ MOG$_{38-49}$ tetramer-positive CD4$^+$ T cells. *Ocab*$^{fl/fl}$ (n=12) and *Ocab*$^{fl/fl}$;CD4-Cre (n=9). For clinical scores and normalized weights, data represent mean ±SEM. All other data represents mean ±SD.

Fig.2 l OCA-B promotes Th1 and Th17 adoptive transfer EAE through recall response

a,**b**, *Ocab*$^{fl/fl}$ and *Ocab*$^{fl/fl}$;CD4-Cre mice were primed with MOG$_{35-55}$ peptide in CFA for 10-14 days. Cells from the spleens and lymph nodes of primed mice were isolated and cultured for 4 days in Th1 or Th17 polarizing conditions. 3.0x10^6 Th1- or Th17-polarized cells were injected i.p. into C57BL/6 wildtype recipient mice (Th1: *Ocab*$^{fl/fl}$ n=5, *Ocab*$^{fl/fl}$;CD4-Cre n=5) (Th17: *Ocab*$^{fl/fl}$ n=3, *Ocab*$^{fl/fl}$;CD4-Cre n=4). Clinical scores were assessed for 14-15 days. **c**, At Th17 adoptive transfer EAE endpoint, brains and spinal cords were analyzed by flow cytometry. Quantification of brain- and spinal cord-infiltrating CD4$^+$ T cells following Th17 adoptive transfer. **d**, Representative flow cytometry plots showing the frequency of IFNγ and IL-17 expressing CD4$^+$ T cells within the spines of Th17 recipient mice. **e**, Quantification of IL-17 frequency in spinal cord infiltrating CD4$^+$ T cells. **f**, Quantification of the total count of IL-17$^+$ CD4$^+$ T cells within the spinal cord of recipient mice. For clinical scores, data represent mean ±SEM. All other data represents mean ±SD.

Fig.3 l OCA-B is dispensable for T cell response to CNS infection with a neurotropic virus

a, *Ocab^{fl/fl}* (n=19) and *Ocab^{fl/fl}*;CD4-Cre (n=17) mice were injected intracranially with 1500 PFU

JHMV. Clinical scores were recorded for 21 days post infection (dpi). **b**, Representative

DAPI/MBP immunofluorescence staining of spinal cord sections at 12 dpi. Demyelinated areas

are marked by decreased MBP staining (green) and often coincide with increased cellular

infiltration (blue). **c**, Quantification of % demyelination in spinal cord sections. (n=7 per group).

Half brains were taken at 7, 12, and 21 dpi for flow cytometric and viral titer analysis. **d**,

Quantification of CD4$^+$ T cell counts within half brain at 7, 12, 21 dpi. *Ocab^{fl/fl}* (n=7-9 per time

point) and *Ocab^{fl/fl}*;CD4-Cre (n=6-7 per time point). **e**, Representative plots showing the frequency

of IFNγ expressing CD4$^+$ T cells 7, 12, and 21 dpi. **f**, Quantification of frequency of IFNγ expressing

CD4$^+$ T cells. *Ocab^{fl/fl}* (n=7-9 per time point) and *Ocab^{fl/fl}*;CD4-Cre (n=6-7 per time point). **g**, Half

brain viral titers at dpi 7, 12, and 21 (n.d., not detected). *Ocab^{fl/fl}* (n=7-9 per time point) and

Ocab^{fl/fl};CD4-Cre (n=6-7 per time point). These data represent the combined results of two

independent experiments. For clinical scores, data represent mean ±SEM. All other data

represents mean ±SD.

Fig.4 | OCA-B expression marks encephalitogenic stem-like CD4$^+$ T cells

a, OCA-B-mCherry mice were injected with MOG$_{35-55}$ peptide in CFA and *pertussis* toxin to induce

EAE. Cervical lymph nodes (CLNs), brains, and spines were isolated at peak disease (day 15)

and analyzed by flow cytometry for mCherry, CD44 and CD62L expression. **b**, CD44 and CD62L

expressing CD4$^+$ T cells were quantified based on mCherry expression. **c**, OCA-B-mCherry mice

were injected with MOG$_{35-55}$ peptide in CFA for 14 days. CD4$^+$ T cells within CLNs and spleens

were profiled by flow cytometry for mCherry, CD44, and CD62L expression. **d**, Quantification of

CD44 and CD62L expression of OCA-B negative and positive CD4$^+$ T cells from MOG-primed

OCA-B-mCherry mice. **e**, Bulk RNA sequencing was performed on OCA-B positive and negative

CD4$^+$ T cells from MOG$_{35-55}$ primed OCA-B-mCherry reporter mice. Heatmap showing relative

expression of the top 40 differentially expressed genes from OCA-B positive and negative CD4$^+$

T cells. **f**, Volcano plot of differentially expressed genes between OCA-B positive and negative T

cells (p value > 0.05 and Log2 fold change > 1 are colored). **g**, Gene Ontology (GO) terms

associated with OCA-B positive and negative groups. **h**, EAE clinical scores of C57BL/6 wild-type

mice following passive transfer of OCA-B-positive or -negative Th1 polarized CD4$^+$ T cells. Clinical

score values represent mean ±SEM. All other data represent mean ±SD.

Fig.5 l OCA-B promotes relapsing-remitting EAE through stem-like CD4⁺ T cells

a, NOD.*Ocab^{fl/fl}* (n=19) and NOD.*Ocab^{fl/fl}*;CD4-Cre (n=15) mice were injected with MOG$_{35-55}$ peptide in CFA and *pertussis* toxin to induce EAE. Clinical scores of animals representing relapsing-remitting disease progression over time. **b**, Animal weights were recorded to determine weight loss throughout initial disease and relapse. **c**, Singe cell RNA sequencing was performed on pooled CD3ε⁺ cells isolated from the brain and spine of NOD.*Ocab^{fl/fl}* (n=3) and NOD.*Ocab^{fl/fl}*;CD4-Cre (n=4) 24 days after EAE induction (remission timepoint). Cell populations were plotted in a UMAP using the Seurat R package and percentages are shown for each cluster. Clusters were identified through differential gene expression analysis. **d**, Feature plots comparing the expression of *Foxp3* and *Mki67* amongst clusters. **E,** Violin plots showing the expression of *Btg1*, *Samhd1*, *Slamf6*, *Tcf7, Sell* and *Tox* within the Treg, proliferating, and Th1-like clusters. **f**, CNS cells isolated at remission timepoint (day 24) were analyzed by spectral cytometry. Representative flow cytometry plots and quantification showing the frequency of CD4⁺ T cells between control and experimental groups. **g**, Representative flow cytometry plots and quantification showing the frequency of CXCR6 expressing CD4⁺ T cells. **h,** Representative flow cytometry plots showing the frequency of CD4⁺ T cells expressing IFNγ and GM-CSF. **i**, Quantification of CD4⁺ T cells expressing IFNγ and GM-CSF. For clinical scores, data represents mean ±SEM. All other data represents mean ±SD.

Fig.6 l OCA-B promotes disease relapse through control of pathogenic stem-like Th17 differentiation

a, Singe cell RNA and TCR sequencing were performed on pooled CD3ε⁺ cells isolated from the brain and spine of NOD.*Ocab^{fl/fl}* (n=5) and NOD.*Ocab^{fl/fl}*;CD4-Cre (n=6) 33 days after EAE induction (relapse timepoint). Cell populations were plotted in a UMAP using the Seurat R package and percentages are shown for each cluster. Clusters were identified through differential gene expression analysis. **b**, UMAP TCR clonotype expansion between *Ocab^{fl/fl}* and *Ocab^{fl/fl}*;CD4-Cre amongst clusters. **c**, Violin plots showing expression of *Ccr9*, *Bach2*, *Ccr7*, *Tcf7*, *Cd44*, *Cxcr6*, *Il1r1* and *Lgals3* in the Ccr9 Stem-like CD4, Stem-like CD4-2, and Th17-like clusters. **d**, Feature plots comparing the expression of *Ccr9*, *Tcf7*, and *Bach2* amongst clusters. **e**, UMAPs showing RNA velocity of spliced and unspliced transcripts by experimental condition. **f**, Violin plots showing expression of *Pdcd1*, *Tigit*, and *Il1r1* in the Th17-like cluster between control and experimental groups.

Extended data legends

Extended Data Fig.1. Minimal differences in cytokine expressing CD4$^+$ T cell, CD8$^+$ T cell, B cell, macrophage or microglial cell counts at peak chronic EAE in OCA-B knockouts

a, Representative spinal cord LFB/H&E staining of thoracic sections taken from mice 15 days after EAE induction. Areas of demyelination are outlined in black and marked by increased loss of LFB (blue) staining and visibility of H&E (pink) within the white matter. **b**, Quantification of the % demyelination from LFB/H&E histology. **c**, Quantification of the frequency of CNS infiltrating CD4$^+$ T cells between Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **d**, Quantification of frequency and count of CD8α positive T cells between Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **e**, Quantification of the number of B cells, macrophages and microglia. **f**, Representative flow cytometry plots comparing the expression of CD44$^+$ within CNS infiltrating CD8$^+$ T cells from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **g**, Quantification of the frequency of CNS infiltrating CD44$^+$ CD8$^+$ T cells within Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **h**, Quantification of frequency and count of IFNγ, IL-17, GM-CSF, and CD44 expressing CD4$^+$ T cells from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. All data represents mean $\pm$SD.

Extended Data Fig.2. Loss of OCA-B does not strongly alter initial Th1 or Th17 polarization in vitro

a, Experimental schematic for assessing CD4$^+$ T cell differences between Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$CD4-Cre prior to *in vitro* Th1 or Th17 polarization. Lymph nodes (cervical, brachial, axillary, and inguinal) were harvested from Ocab$^{fl/fl}$ (n=4) and Ocab$^{fl/fl}$;CD4-Cre (n=4) mice following 14 days of MOG$_{35-55}$/CFA priming and analyzed by flow cytometry **b**, Frequency of viable cells isolated from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **c**, Total CD4$^+$ T cell numbers isolated from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **d**, Quantification of the frequency of IFNγ expressing CD4$^+$ T cells isolated from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **e**, Quantification of the frequency of IL-17 expressing CD4$^+$ T cells derived from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. **f**, Frequency of Ki67 expressing CD4$^+$ T cells from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. CD4$^+$ T cells were isolated from Ocab$^{fl/fl}$ (n=8) and Ocab$^{fl/fl}$;CD4-Cre (n=8) mice following 14 days of MOG$_{35-55}$/CFA priming, fluorescently labeled using QTracker nanocrystals (Ocab$^{fl/fl}$-QTracker 655) (Ocab$^{fl/fl}$;CD4-Cre-Qtracker 525), injected i.p. into naïve C57BL/6 recipients and assessed after 3 days by flow cytometry. **g**, Representative flow cytometry plot showing QTracker 655 and QTracker 525 labeled CD4$^+$ T cells isolated from the peritoneal cavity of recipient mice. **h**,

Quantification of the frequency of QTracker 655 and QTracker 525 labeled CD4[+] T cells isolated

from the peritoneal cavity of recipient mice. **i**, Representative flow cytometry plots comparing

viability of peritoneal QTracker 655 and QTracker 525 labeled CD4[+] T cells by Live/Dead Blue

staining. **j**, Peritoneal QTracker 655 and QTracker 525 labeled CD4[+] T cell viability

quantification. **k**, Representative flow cytometry plots showing Ki67 expression within QTracker

655 and QTracker 525 labeled CD4[+] T cells isolated from the peritoneal cavity. **l**, Quantification

of the frequency of Ki67 expressing QTracker 655 and QTracker 525 labeled CD4[+] T cells. **m**,

Representative flow cytometry plots showing IFNγ and IL-17 expressing CD4[+] T cells from

Ocab[fl/fl] (n=5) and Ocab[fl/fl];CD4-Cre (n=5) mice after Th1 and Th17 *in vitro* polarization. **n**,

Quantification of IFNγ expressing CD4[+] T cells after Th1 polarization. **o**, Quantification of IL-17

expressing CD4[+] T cells after Th17 polarization. **p**, Quantification of the frequency of IFNγ

expressing CD4[+] T cells isolated from the spinal cords of Th17 adoptive transfer EAE recipient

mice. All data represents mean ±SD.

Extended Data Fig.3. Normal immune responses to intracranial JHMV infection in the

absence of OCA-B

a, Representative LFB/H&E histology of spinal cord sections at dpi 21. **b**, Demyelination

quantification of LFB/H&E histology at dpi 21. **c**, Quantification of the number of IFNγ expressing

CD4[+] T cells at dpi 7, 12 and 21. **d**, Quantification of the number of CD8[+] T cells at dpi 7, 12 and

21. **e**, Quantification of the frequency of IFNγ expressing CD8[+] T cells at dpi 7, 12 and 21. **f,**

Quantification of the number of IFNγ expressing CD8[+] T cells at dpi 7, 12 and 21. All data

represents mean ±SD.

Extended Data Fig.4. Sorting OCA-B-mCherry reporter T cells and additional analysis of

the effects of transferring OCA-B[hi] versus OCA-B[lo] CD4[+] cells

a, Quantification of the frequency of CD44[+] CD62[-] CD4[+] T cells within the cervical lymph nodes

of OCA-B-mCherry reporter mice at peak EAE by OCA-B-mCherry expression. **b**, Gating

strategy for sorting of OCA-B[+] and OCA-B[-] CD4[+] T cells. **c**, Normalized weights of NOD.Ocab[fl/fl]

and NOD.Ocab[fl/fl];CD4-Cre mice following EAE induction. Normalized weight is represented by

mean ±SEM. All other data represents mean ±SD.

Extended Data Fig.5. Individual NOD mouse EAE clinical scores, additional remission

scRNA-seq cluster annotation genes, and heat map of gene enrichments by cluster

a, Clinical scores of individual NOD.Ocab$^{fl/fl}$ mice following EAE induction. **b**, Clinical scores of

individual NOD.Ocab$^{fl/fl}$;CD4-Cre mice following EAE induction. **c**, UMAP feature plots showing

expression of *Cxcr6*, *Il23r*, and *Bach2* amongst clusters at remission timepoint. **d**, Heatmap

showing the top 10 genes expressed by cluster at remission EAE timepoint.

Extended Data Fig.6. Heat map of NOD.EAE relapse gene enrichments by cluster, TCR

clonotype overlap between condition, example relapse feature plots, and pseudotime

analysis of relapse single cell RNA seq

a, Heatmap showing the top 10 genes expressed by cluster at relapse EAE timepoint. **b,** Alluvial

plot showing top 10 TCR clonotype gene overlap between Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre groups.

c, UMAP feature showing the expression of *CD4*, *CD8a, Tox, Sell*, *Ccr7*, and *Ccr4* amongst

clusters at relapse timepoint. **d**, Pseudotime analysis of relapse single cell from root starting

node at or near Ccr9 stem-like cluster.

Extended Data Fig.7. OCA-B promotes pathogenic maturation of stem-like Th17 cells in

EAE

OCA-B expression promotes and selectively labels pathogenic stem-like Th17 cells which

preferentially transfer EAE and instigate disease relapse.

Methods

Chronic C57BL/6 EAE

All animal work was approved by the University of Utah Institutional Animal Care and Use

Committee. 11-12 week-old female C57BL/6 littermate animals were used for all experiments.

For adoptive transfer experiments using the OCA-B-mCherry reporter allele[29], 11 week-old

females were used for chronic EAE and 9 week-old males were used as MOG_{35-55} primed adoptive

transfer donors. EAE was induced on day 0 by two 100 μL s.c. injections into both hind flanks of

0.5 μmol/mL MOG_{35-55} peptide (University of Utah Peptide Core) emulsified in complete Freud's

adjuvant (CFA), comprised of incomplete Freund's adjuvant (Cat# 77145;ThermoFisher) together

with 4 mg/mL heat-killed *M. tuberculosis* (Cat# DF3114338; Fisher Scientific). Animals were given

two 400 ng i.p. injections of *B. pertussis* toxin (Cat# 181; List Biological Laboratories) on days 0

and 2.

Clinical Scoring

EAE clinical scores were determined using a 0-5 scale with 0.5 scoring intervals: 0.5, tip of tail is

limp or tail muscle strain observed; 1, complete limp tail; 1.5, complete limp tail and at least one

hind limb consistently falls through wire rack when animal is dropped; 2, complete limp tail and

one hind leg dragging without movement beyond hip; 2.5, complete limp tail and dragging of both

hind limbs without movement beyond hip; 3, complete hind limb paralysis; 3.5, complete hind limb

paralysis with flat laying hind quarters and hump in upper torso; 4, complete hind limb and partial

forelimb paralysis, minimal movement around cage; 4.5, complete hind limb and partial forelimb

paralysis, no movement around cage; 5, seizure or death. Animals scoring a $\geq$4 for more than 48

hours were euthanized and scored 5 for the remainder of the experiment. JHMV clinical disease

was evaluated using previously described 0-4 point scale[65].

Immunofluorescence and Histology

Spinal columns were isolated from C57BL/6 Ocab^fl/fl and Ocab^fl/fl;CD4-Cre animals after EAE or

JHMV induced demyelination and fixed overnight in in 4% paraformaldehyde at 4C. Spinal cords

were subsequently isolated and cryoprotected in 30% sucrose for 3-5 days. Thoracic and lumbar

sections were embedded in optimum cutting temperature (OCT) compound (ThermoFisher

Scientific) and frozen and stored at -80C. 8 μm spinal sections were cut by cryostat and stained

with Luxol fast blue (Cat# 212170250; ThermoFisher) along with hematoxylin (Cat# 8495; ENG

Scientific) and eosin (Cat# E511-25; Fisher Chemical). Demyelination was assessed by dividing

the area of demyelinated white matter by total white matter as previously described[66,67]. For

immunofluorescence, spinal cord sections were blocked in 5% normal goat serum (NGS; Cat#

005-000-121, Jackson ImmunoResearch) and 0.1% Tween-20 for 1 hour at room temperature.

Sections were stained overnight at 4°C with primary antibody rabbit anti-MBP (1:1000; Cat#

PA1050, Boster), followed by secondary antibody goat anti-rabbit conjugated with Alexa Fluor

488 (1:750; Cat# A11008, Invitrogen) for 2 hours at room temperature. Primary and secondary

antibodies were diluted in 1% bovine serum albumin (BSA) and 0.1% Tween-20. Slides were

cover-slipped using Fluoromount-G with DAPI (Cat# 00495952, Invitrogen). Sections were

imaged with a Zeiss Axioscan 7 Microscope Slide Scanner using a 20× objective.

Flow cytometry and spectral cytometry

Spinal cords and brains were harvested at peak disease or specified timepoint and mechanically

dissociated by passing through a 100 µm cell strainer. A 30/70% Percoll gradient (Cat# 17544501;

Cytiva) was used to enrich mononuclear cells. Cells from collected spleens and lymph nodes

(cervical, brachial, axillary and inguinal) were passed through a 70 µm strainer and erythrocytes

lysed by ammonium-chloride-potassium lysis buffer (150 mM NH_4Cl, 10 mM $KHCO_3$, 0.1 mM

Na_2EDTA). For intracellular staining, cells were cultured in complete media (RPMI 1640, 10%

FBS, 1% Pen/Strep, Glutamax) supplemented with 1 µL/mL brefeldin A (Cat# 555029; Golgiplug,

BD), 50 ng/mL phorbol myristate acetate (Cat# P1585; Sigma-Aldrich) and 1 µg/mL ionomycin

(Cat# I0634; Sigma-Aldrich) for 4 hr. Cultured cells were fixed using cytofix/cytoperm (Cat#

554714; BD) according to manufacturer's protocol and stained for intracellular cytokines in

perm/wash buffer. Flow cytometry samples were profiled using a BD Fortessa (BD Biosciences,

Fig. 2, 3, 4) or an Aurora spectral flow cytometer (Cytek, Fig. 1, 2, 5). FlowJo software (BD

Biosciences) was used for data analysis.

Th1 and Th17 polarization and adoptive transfer C57BL/6 EAE

10-12 week-old female C57BL/6 littermate animals were used for all experiments. Donor animals

were immunized with MOG_{35-55} in CFA as above. 10-14 days post-immunization, spleen and

lymph node (cervical, axillary, brachial and inguinal) cells were collected and placed into mixed

culture with MOG peptide and cytokines at 5.0×10^6 cells/mL. For Th1 conditions, complete media

was supplemented with 50 µg/mL MOG_{35-55}, 6 ng/mL rmIL-6 (Cat# 216-16; Peprotech) and 2

ng/mL rmIFNγ (Cat# 315-05; Peprotech). For Th17 conditions, complete media was

supplemented with 50 μg/mL MOG$_{35-55}$, 8 ng/mL rmIL-23 (Cat# 1887-ML-010; R&D Systems), 10

ng/mL rmIL-1α (Cat# 211-11A; Peprotech), and 10 μg/mL anti-IFNγ (Cat# BE0055; BioXcell

XMG1.2). After 4 days of respective culture, CD4$^+$ T cells were purified using negative magnetic

selection and resuspended in PBS at 1.5$\times$10^6 cells/mL. 100 μL cell suspension was injected i.p.

into age-matched male C57BL/6 recipients (~13 week-old at time of disease induction). Clinical

disease was scored for 15 days.

Pre-polarization profiling and mixed adoptive transfer

10 week-old C57BL/6 *Ocab$^{fl/fl}$* and *Ocab$^{fl/fl}$;CD4-Cre* littermates were immunized with MOG$_{35-55}$ in

CFA as described above. After 14 days, spleen and lymph node (cervical, axillary, brachial and

inguinal) cells were isolated as described above. For the pre-polarization analysis (Extended Data

Fig. 2b-f), isolated lymph node cells were cultured in complete media (RPMI 1640, 10% FBS, 1%

Pen/Strep, Glutamax) supplemented with 1 μL/mL brefeldin A (Cat# 555029; Golgiplug, BD), 50

ng/mL phorbol myristate acetate (Cat# P1585; Sigma-Aldrich) and 1 μg/mL ionomycin (Cat#

I0634; Sigma-Aldrich) for 4 hr. Cells were stained for surface markers, fixed using the

Foxp3/Transcription Factor Staining Buffer Kit (Cat#00-5523-00, ThermoFisher), and stained for

intracellular markers (IFNγ, IL-17, and Ki67). For the mixed adoptive transfer (Extended Data Fig.

2g-l), CD4$^+$ cells were enriched from isolated splenocytes by magnetic negative selection (Miltenyi

Bio kit and LS columns). Enriched CD4$^+$ T cells from *Ocab$^{fl/fl}$* donors were labeled with QTracker

655 (Cat#Q25021MP, ThermoFisher) and enriched CD4$^+$ T cells from *Ocab$^{fl/fl}$;CD4-Cre* donors

were labeled with QTracker 525 (Cat#Q25041MP, ThermoFisher) according to manufactures

instructions. Labeled cells from both genotypes were mixed at a 1:1 ratio and injected i.p. into

naïve age-match C56BL/6 recipients at 6.0x10^6 cells per mouse. After 72 hours, cells were

isolated from the peritoneal cavity of mixed adoptive transfer recipients as previously described[68].

Isolated cells were stimulated in the presence of brefeldin A, fixed and stained for surface and

intracellular markers as described above. Cells were analyzed by spectral cytometry for presence

of QTracker nanocrystals, viability by Live/Dead Blue staining (Cat#L23105, ThermoFisher) and

Ki67 expression.

JHMV

Ocab^{fl/fl} and *Ocab^{fl/fl}*;CD4-Cre C57BL/6 mice were anesthetized using isoflurane. Mice were

intracranially (i.c) injected with 1500 plaque-forming units (PFU) of JHMV (strain V34) suspended

in 30 μL Hank's balanced salt solution (ThermoFisher). Clinical disease was assessed for 21 days

using a previously described scale[69]. Mice were sacrificed at days 7, 12, and 21 post-infection to

assess viral titers within brain homogenates using a plaque assay previously described[70]. Spinal

cords were taken on day 12 and 21 post-infection to assess demyelination by DAPI/MBP and

luxol fast blue and hematoxylin/eosin histology respectively.

OCA-B-mCherry adoptive transfer EAE

OCA-B-mCherry reporter allele animals[29] were primed for 14 days with MOG_{35-55} in CFA as

described above. After priming, spleen and lymph nodes were collected and CD4$^+$ T cells were

purified by magnetic negative selection (Miltenyi Bio kit and LS columns). Non-CD4$^+$ T cells were

removed from the LS column and placed on ice for subsequent in vitro culture. Purified CD4$^+$ T

cells were separated by OCA-B-mCherry expression (PE-Texas Red) using flow automated cell

sorting (FACS). OCA-B-mCherry positive and negative CD4$^+$ T cells were placed into mixed

culture in complete medium with the previously purified non-CD4$^+$ T cells at a 1:10 ratio (CD4$^+$ to

non-CD4$^+$) at $1.0×10^6$ cells/mL. The culture was supplemented with 20 μg/mL MOG_{35-55} and 0.5

ng/mL IL-12p70 (Cat# 210-12; Peprotech). After two days of culture, CD4$^+$ T cells were again

purified through negative magnetic selection. Purified CD4$^+$ T cells from OCA-B-mCherry positive

and negative cultures were resuspended in PBS at $8.8×10^6$ cells/mL. 100 μL cell suspension was

injected i.p. into age-matched male C57BL/6 recipients. Clinical disease was assessed for 24

days.

Bulk RNAseq

OCA-B-mCherry reporter mice (n=9) were primed for 14 days with MOG_{35-55} in CFA as described

above. After priming, lymph nodes (cervical, axillary, brachial and inguinal) were isolated and

mCherry positive and negative CD4$^+$ T cells were collected by FACS. Total RNA, three groups

(pooled 3 mice each), was isolated using the Quick RNA micro kit (Cat# R1050; Zymo Research)

with DNase treatment (Cat# 79254; Qiagen). RNA concentration was measured with a Qubit RNA

HS Assay Kit (Cat# #Q32855; Fisher Scientific). RNA quality was evaluated with an RNA

ScreenTape Assay (Cat# 5067-5579, 5067-5580; Agilent Technologies). Total RNA samples (5-

500 ng) were hybridized with NEBNext rRNA Depletion Kit v2 (Cat# E7400; New England Biolabs)

to diminish rRNA from the samples. Stranded RNA sequencing libraries were prepared as
described using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (Cat# E7760L;
New England Biolabs). Purified libraries were qualified on an 4150 TapeStation (Agilent
Technologies) using a D1000 ScreenTape assay (Cat# 5567-5582, 5067-5583; Agilent
Technologies). The molarity of adapter-modified molecules was defined by quantitative PCR
using the Kapa Biosystems Kapa Library Quant Kit (Cat# KK4824Roche). Individual libraries were
normalized to 5 nM. NovaSeq 150×150 bp sequencing libraries were chemically denatured and
applied to an Illumina NovaSeq flow cell using the XP workflow (20043131). Following transfer to
an Illumina NovaSeq 6000 instrument, 150×150 cycle paired-end sequencing was performed
using a S4 reagent Kit v1.5 (Cat# 20028312; Illumina). 8-11 million reads were generated per
sample and were aligned to the *GRCm38* mouse reference genome. Differential gene expression
and pathway analysis were conducted using the DESeq2 and Enrichr R packages.
730
NOD-EAE
15-18 week-old female littermate *Ocab^{fl/fl}* and *Ocab^{fl/fl}*;CD4-Cre animals backcrossed to a NOD
strain background[30] were used for all experiments. EAE was induced by two 100 μL s.c. injections
into both hind flanks of 0.5 μmol/mL MOG$_{35-55}$ peptide emulsified in incomplete Freuds adjuvant
(Cat# 77145; ThermoFisher) supplemented with 4 mg/mL heat-killed *M. tuberculosis* (Cat#
DF3114338; Fisher Scientific) to form CFA. Animals were given two 400 ng i.p. injections of
Bordetella pertussis toxin (Cat# 181; List Biological Laboratories) on days 0 and 2. Clinical
disease was assessed over 47 days using a 5-point scale described below. Remission timepoints
were determined to be between days 22-26 while relapse was determined to be between days
26-39.
741
Single-cell RNA-seq
NOD.*Ocab^{fl/fl}* and NOD.*Ocab^{fl/fl}*;CD4-Cre animals were induced with EAE and monitored for
disease progression. Brains and spinal cords were isolated on day 24 (remission) and day 33
(relapse), and processed for FACS sorting. Viable cells were sorted by CD3ϵ using a FACSAria
(BD Bioscience). Cells from each condition were isolated from 3-4 mice and combined. Cells were
processed using the 5' 10X Genomics Chromium platform according to the manufacturer's
instructions. Paired-end high-throughput sequencing (125 cycles) was performed using a
NovaSeq instrument (Illumina). Sequencing reads were processed by using the 10X Genomics

CellRanger pipeline and further analyzed using the Seurat R package. Analysis of cells used a

standard filtering protocol removing cells with unique feature counts of >5,000 or <100, as well as

cells with >10% mitochondrial counts (indicative of dead cells). No more than 10% of total cells

were removed by this process. Cells were subjected to unsupervised hierarchical clustering

followed by uniform manifold approximation and projection (UMAP) to visualize clusters with

similar gene expression and their relative positions. RNA velocity analysis was performed using

velocyto.R version 0.6 to model cell state transitions using the guidelines provided in Manno et

al.[45] and the associated repository at https://github.com/velocyto-team/velocyto.R. High

dimensional velocity vectors were graphed on the predefined Seurat UMAP projection using

localized average grid projections. Pseudotime analysis was conducted using Seurat and

Monocle3 packages. Integrated data was divided into subset based on genotype and Monocle3

was used to calculate trajectory, order cells and plot pseudotime for each condition.

Statistics

Multiple two-tailed student's t-tests were used to determine statistical differences in EAE clinical

scores and all error bars denote ±SEM unless otherwise noted. Two-way ANOVA was used to

determine differences in weight loss throughout disease progression. GraphPad Prism software

was used for all statistics and graphing. Two-tailed student's t-tests were used to determine

statistical differences in cell populations observed by flow cytometry and all error bars represent

±SD unless otherwise noted. For all figures, ns = $p>0.05$, * = $p\leq0.05$, ** = $p\leq0.01$, *** = $p\leq0.001$,

**** = $p\leq0.0001$.

Reporting summary

Information regarding research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

All RNA sequencing, single-cell sequencing, and single-cell T cell V(D)J sequencing data associated with the findings of this study have been deposited to NCBI's GEO database under accession code GSE243727.

Acknowledgements

We thank R.M. O'Connell and M. Bettini for critical reading of the manuscript. We thank J. Marvin and University of Utah Health Sciences Center Flow Cytometry core, and B. Dalley, O. Allen and the High-Throughput Genomics core. Peptides were synthesized by the DNA/Peptide facility. We thank T. Lane for assistance with JHMV constructs, protocols and reagents. This work was supported by grants from the National Institutes of Health/ National Institute of Allergy and Infectious Diseases (R01AI162929 and R01AI100873) and the Praespero Foundation (Canada) to DT, and training grants from the National Institute of Neurological Disorders and Stroke (T32NS115664) to EPH and National Institute of Allergy and Infectious Diseases (T32AI055434) to ARS.

Author contributions

E.P.H. and D.T. conceived the study. E.P.H. and A.S. performed experiments. E.P.H, A.S., E.M.M. and B.T.S. analyzed data. T.E.L., B.T.S. and D.T. provided critical resources and supervision for the study. E.P.H. and D.T. wrote, reviewed and revised the manuscript.

Competing interests

The authors declare that they have no conflicts of interest

References

1.   Wallin, M. T. *et al.* The prevalence of MS in the United States. *Neurology* **92**, e1029–
e1040 (2019).

2.   Wtccc, C. C. *et al.* Genetic risk and a primary role for cell-mediated immune mechanisms
in multiple sclerosis. *Nature* **476**, 214–219 (2011).

3.   IMSGC. A high-density screen for linkage in multiple sclerosis. *Am. J. Hum. Genet.* **77**,
454–467 (2005).

4.   Adorini, L., Guery, J.-C. & Trembleau, S. Manipulation of the Th1/Th2 Cell Balance: An
Approach to Treat Human Autoimmune Diseases? *Autoimmunity* **23**, 53–68 (1996).

5.   Langrish, C. L. *et al.* IL-23 drives a pathogenic T cell population that induces autoimmune
inflammation. *J. Exp. Med.* **201**, 233–240 (2005).

6.   Brucklacher-Waldert, V., Stuerner, K., Kolster, M., Wolthausen, J. & Tolosa, E.
Phenotypical and functional characterization of T helper 17 cells in multiple sclerosis.
Brain **132**, 3329–3341 (2009).

7.   Montes, M. *et al.* Oligoclonal myelin-reactive T-cell infiltrates derived from multiple
sclerosis lesions are enriched in Th17 cells. *Clin. Immunol.* **130**, 133–144 (2009).

8.   Hartmann, F. J. *et al.* Multiple sclerosis-associated IL2RA polymorphism controls GM-
CSF production in human TH cells. *Nat. Commun.* **5**, 5056 (2014).

9.   Scholz, C., Patton, K. T., Anderson, D. E., Freeman, G. J. & Hafler, D. A. Expansion of
Autoreactive T Cells in Multiple Sclerosis Is Independent of Exogenous B7 Costimulation.
J. Immunol. **160**, 1532–1538 (1998).

10.   Burns, J. B., Bartholomew, B. D. & Lobo, S. T. Isolation of CD45RO+, Memory T Cells
Recognizing Proteolipid Protein from Neurologically Normal Subjects. *Cell. Immunol.* **212**,
44–50 (2001).

11.   Pelfrey, C. M. *et al.* Quantification of Self-Recognition in Multiple Sclerosis by Single-Cell
Analysis of Cytokine Production. *J. Immunol.* **165**, 1641–1651 (2000).

12.   Lovett-Racke, A. E. *et al.* Decreased dependence of myelin basic protein-reactive T cells
on CD28-mediated costimulation in multiple sclerosis patients. A marker of
activated/memory T cells. *J. Clin. Invest.* **101**, 725–730 (1998).

13.   Lee, H.-G. *et al.* Pathogenic function of bystander-activated memory-like CD4+ T cells in
autoimmune encephalomyelitis. *Nat. Commun.* **10**, 709 (2019).

14.   Williams, J. L. *et al.* Memory cells specific for myelin oligodendrocyte glycoprotein (MOG)
govern the transfer of experimental autoimmune encephalomyelitis. *J. Neuroimmunol.*

234, 84–92 (2011).

15.   Elyaman, W. *et al.* Distinct Functions of Autoreactive Memory and Effector CD4+ T Cells

in Experimental Autoimmune Encephalomyelitis. *Am. J. Pathol.* **173**, 411–422 (2008).

16.   Maurano, M. T. *et al.* Systematic Localization of Common Disease-Associated Variation

in Regulatory DNA. *Science (80-.).* **337**, 1190–1195 (2012).

17.   Graham, D. S. C., Wong, A. K., McHugh, N. J., Whittaker, J. C. & Vyse, T. J. Evidence for

unique association signals in SLE at the CD28–CTLA4–ICOS locus in a family-based

study. *Hum. Mol. Genet.* **15**, 3195–3205 (2006).

18.   van Heel, D. A. Inflammatory bowel disease is associated with a TNF polymorphism that

affects an interaction between the OCT1 and NF-kappaB transcription factors. *Hum. Mol.*

Genet. **11**, 1281–1289 (2002).

19.   Farh, K. K. *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease

variants. *Nature* **518**, 337–343 (2015).

20.   Shakya, A. *et al.* Oct1 and OCA-B are selectively required for CD4 memory T cell

function. *J. Exp. Med.* **212**, 2115–2131 (2015).

21.   Kim, H. *et al.* T cell-selective deletion of Oct1 protects animals from autoimmune

neuroinflammation while maintaining neurotropic pathogen response. *J.*

Neuroinflammation **16**, 133 (2019).

22.   Vázquez-Arreguín, K. *et al.* Oct1/Pou2f1 is selectively required for colon regeneration and

regulates colon malignancy. *PLOS Genet.* **15**, e1007687 (2019).

23.   Perovanovic, J. *et al.* Oct1 cooperates with the Smad family of transcription factors to

promote mesodermal lineage specification. *Sci. Signal.* **16**, (2023).

24.   Sebastiano, V. *et al.* Oct1 regulates trophoblast development during early mouse

embryogenesis. *Development* **137**, 3551–3560 (2010).

25.   Hwang, S. S., Kim, L. K., Lee, G. R. & Flavell, R. A. Role of OCT-1 and partner proteins

in T cell differentiation. *Biochim. Biophys. Acta - Gene Regul. Mech.* **1859**, 825–831

(2016).

26.   Kim, U. *et al.* The B-cell-specific transcription coactivator OCA-B/OBF-1/Bob-1 is

essential for normal production of immunoglobulin isotypes. *Nature* **383**, 542–547 (1996).

27.   The GAMES Collaborative Group. Linkage disequilibrium screening for multiple sclerosis

implicates JAG1 and POU2AF1 as susceptibility genes in Europeans. *J. Neuroimmunol.*

179, 108–116 (2006).

28.   Nakamura, M. *et al.* Genome-wide Association Study Identifies TNFSF15 and POU2AF1

as Susceptibility Loci for Primary Biliary Cirrhosis in the Japanese Population. *Am. J.*

Hum. Genet. **91**, 721–728 (2012).

29.   Sun, W. *et al.* OCA-B/Pou2af1 is sufficient to promote CD4+ T cell memory and

prospectively identifies memory precursors. *Proc. Natl. Acad. Sci. U. S. A.* **121**,

e2309153121 (2024).

30.   Kim, H. *et al.* Targeting transcriptional coregulator OCA-B/Pou2af1 blocks activated

autoreactive T cells in the pancreas and type 1 diabetes. *J. Exp. Med.* **218**, 133 (2021).

31.   Tzartos, J. S. *et al.* IL-21 and IL-21 Receptor Expression in Lymphocytes and Neurons in

Multiple Sclerosis Brain. *Am. J. Pathol.* **178**, 794–802 (2011).

32.   Ndhlovu, L. C., Ishii, N., Murata, K., Sato, T. & Sugamura, K. Critical Involvement of

OX40 Ligand Signals in the T Cell Priming Events During Experimental Autoimmune

Encephalomyelitis. *J. Immunol.* **167**, 2991–2999 (2001).

33.   Hurwitz, A. A., Sullivan, T. J., Sobel, R. A. & Allison, J. P. Cytotoxic T lymphocyte

antigen-4 (CTLA-4) limits the expansion of encephalitogenic T cells in experimental

autoimmune encephalomyelitis (EAE)-resistant BALB/c mice. *Proc. Natl. Acad. Sci.* **99**,

3013–3017 (2002).

34.   Arellano, G., Ottum, P. A., Reyes, L. I., Burgos, P. I. & Naves, R. Stage-Specific Role of

Interferon-Gamma in Experimental Autoimmune Encephalomyelitis and Multiple

Sclerosis. *Front. Immunol.* **6**, (2015).

35.   Rottman, J. B. *et al.* The costimulatory molecule ICOS plays an important role in the

immunopathogenesis of EAE. *Nat. Immunol.* **2**, 605–611 (2001).

36.   Rouse, M., Nagarkatti, M. & Nagarkatti, P. S. The role of IL-2 in the activation and

expansion of regulatory T-cells and the development of experimental autoimmune

encephalomyelitis. *Immunobiology* **218**, 674–682 (2013).

37.   Duncker, P. C., Stoolman, J. S., Huber, A. K. & Segal, B. M. GM-CSF Promotes Chronic

Disability in Experimental Autoimmune Encephalomyelitis by Altering the Composition of

Central Nervous System–Infiltrating Cells, but Is Dispensable for Disease Induction. *J.*

Immunol. **200**, 966–973 (2018).

38.   Yosef, N. *et al.* Dynamic regulatory network controlling TH17 cell differentiation. *Nature*

496, 461–468 (2013).

39.   Ikegami, I., Takaki, H., Kamiya, S., Kamekura, R. & Ichimiya, S. Bob1 enhances RORγt-

mediated IL-17A expression in Th17 cells through interaction with RORγt. *Biochem.*

Biophys. Res. Commun. **514**, 1167–1171 (2019).

40.   Baker, D. *et al.* Autoimmune encephalomyelitis in NOD mice is not initially a progressive

898   multiple sclerosis model. *Ann. Clin. Transl. Neurol.* **6**, acn3.792 (2019).

899   41.   Rumble, J. M. *et al.* Neutrophil-related factors as biomarkers in EAE and MS. *J. Exp.*

900   *Med.* **212**, 23–35 (2015).

901   42.   Kroenke, M. A., Carlson, T. J., Andjelkovic, A. V. & Segal, B. M. IL-12– and IL-23–

902   modulated T cells induce distinct types of EAE based on histology, CNS chemokine

903   profile, and response to cytokine inhibition. *J. Exp. Med.* **205**, 1535–1541 (2008).

904   43.   Grifka-Walk, H. M., Lalor, S. J. & Segal, B. M. Highly polarized Th17 cells induce EAE via

905   a T-bet independent mechanism. *Eur. J. Immunol.* **43**, 2824–2831 (2013).

906   44.   Schnell, A. *et al.* Stem-like intestinal Th17 cells give rise to pathogenic effector T cells

907   during autoimmunity. *Cell* **184**, 6281-6298.e23 (2021).

908   45.   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

909   46.   Alpsoy, A. *et al.* IκBζ is a dual-use coactivator of NF-κB and POU transcription factors.

910   *Mol. Cell* (2024). doi:10.1016/j.molcel.2024.01.007

911   47.   Wu, X. S. *et al.* OCA-T1 and OCA-T2 are coactivators of POU2F3 in the tuft cell lineage.

912   *Nature* **607**, 169–175 (2022).

913   48.   Schubart, D. B., Rolink, A., Kosco-Vilbois, M. H., Botteri, F. & Matthias, P. B-cell-specific

914   coactivator OBF-1/OCA-B/Bob1 required for immune response and germinal centre

915   formation. *Nature* **383**, 538–42 (1996).

916   49.   Schubart, D. B., Rolink, A., Kosco-Vilbois, M. H., Botteri, F. & Matthias, P. B-cell-specif ic

917   coactivator OBF-1/OCA-B/Bob1 required for immune response and germinal centre

918   formation. *Nature* **383**, 538–542 (1996).

919   50.   Qin, X. F., Reichlin, A., Luo, Y., Roeder, R. G. & Nussenzweig, M. C. OCA-B integrates B

920   cell antigen receptor-, CD40L- and IL 4-mediated signals for the germinal center pathway

921   of B cell development. *EMBO J.* **17**, 5066–75 (1998).

922   51.   Heng, T. S. P. *et al.* The Immunological Genome Project: networks of gene expression in

923   immune cells. *Nat. Immunol.* **9**, 1091–1094 (2008).

924   52.   Zwilling, S., Dieckmann, A., Pfisterer, P., Angel, P. & Wirth, T. Inducible Expression and

925   Phosphorylation of Coactivator BOB.1/OBF.1 in T Cells. *Science (80-.).* **277**, 221–225

926   (1997).

927   53.   Gearty, S. V. *et al.* An autoimmune stem-like CD8 T cell population drives type 1

928   diabetes. *Nature* **602**, 156–161 (2022).

54.   Paroni, M. *et al.* Recognition of viral and self-antigens by T H 1 and T H 1/T H 17 central

memory cells in patients with multiple sclerosis reveals distinct roles in immune

surveillance and relapses. *J. Allergy Clin. Immunol.* **140**, 797–808 (2017).

55. Johns, T. G. *et al.* Myelin oligodendrocyte glycoprotein induces a demyelinating

encephalomyelitis resembling multiple sclerosis. *J. Immunol.* **154**, 5536–41 (1995).

56. McRae, B. L. *et al.* Induction of active and adoptive relapsing experimental autoimmune

encephalomyelitis (EAE) using an encephalitogenic epitope of proteolipid protein. *J.*

Neuroimmunol. **38**, 229–240 (1992).

57. Bernard, C. C. A. *et al.* Myelin oligodendrocyte glycoprotein: a novel candidate

autoantigen in multiple sclerosis. *J. Mol. Med.* **75**, 77–88 (1997).

58. Anderson, A. C. *et al.* A Transgenic Model of Central Nervous System Autoimmunity

Mediated by CD4+ and CD8+ T and B Cells. *J. Immunol.* **188**, 2084–2092 (2012).

59. Nish, S. A. *et al.* CD4+ T cell effector commitment coupled to self-renewal by asymmetric

cell divisions. *J. Exp. Med.* **214**, 39–47 (2017).

60. Mammadli, M., Suo, L., Sen, J. M. & Karimi, M. TCF-1 Is Required for CD4 T Cell

Persistence Functions during AlloImmunity. *Int. J. Mol. Sci.* **24**, 4326 (2023).

61. Kadowaki, A., Saga, R., Lin, Y., Sato, W. & Yamamura, T. Gut microbiota-dependent

CCR9+CD4+ T cells are altered in secondary progressive multiple sclerosis. *Brain* **142**,

916–931 (2019).

62. Thakore, P. I. *et al.* The chromatin landscape of Th17 cells reveals mechanisms of

diversification of regulatory and pro-inflammatory states. *BioRxiv* (2022).

63. Song, S. *et al.* OBF1 and Oct factors control the germinal center transcriptional program.

Blood **137**, 2920–2934 (2021).

64. Sha, Y. & Markovic-Plese, S. Activated IL-1RI Signaling Pathway Induces Th17 Cell

Differentiation via Interferon Regulatory Factor 4 Signaling in Patients with Relapsing-

Remitting Multiple Sclerosis. *Front. Immunol.* **7**, (2016).

65. Lane, T. E. *et al.* A Central Role for CD4 + T Cells and RANTES in Virus-Induced Central

Nervous System Inflammation and Demyelination. *J. Virol.* **74**, 1415–1424 (2000).

66. Blanc, C. A. *et al.* Sphingosine-1-Phosphate Receptor Antagonism Enhances

Proliferation and Migration of Engrafted Neural Progenitor Cells in a Model of Viral-

Induced Demyelination. *Am. J. Pathol.* **185**, 2819–2832 (2015).

67. Plaisted, W. C. *et al.* Remyelination Is Correlated with Regulatory T Cell Induction

Following Human Embryoid Body-Derived Neural Precursor Cell Transplantation in a

Viral Model of Multiple Sclerosis. *PLoS One* **11**, e0157620 (2016).

68.   Ray, A. & Dittel, B. N. Isolation of mouse peritoneal cavity cells. *J. Vis. Exp.* (2010).

964   doi:10.3791/1488

965   69.   Lane, T. E. *et al.* A central role for CD4(+) T cells and RANTES in virus-induced central

966   nervous system inflammation and demyelination. *J. Virol.* **74**, 1415–24 (2000).

967   70.   Dickey, L. L., Worne, C. L., Glover, J. L., Lane, T. E. & O'Connell, R. M. MicroRNA-155

968   enhances T cell trafficking and antiviral effector function in a model of coronavirus-

969   induced neurologic disease. *J. Neuroinflammation* **13**, 240 (2016).

a OCA-B-mCherry EAE Peak Disease

Live, CD19⁻, CD4⁺ | OCA-B (-) | OCA-B (+)

b

c OCA-B-mCherry MOG₃₅₋₅₅ Primed

d

e

f

g OCA-B⁺ Enriched Pathways

KEGG Pathway	*p*-value
Systemic Lupus Erythematosus	1.02E-04
Cytokine Cytokine Receptor Interaction	5.04E-04
Autoimmune Thyroid Disease	1.16E-03
Type I Diabetes Mellitus	2.06E-03

h

a NOD.EAE relapse

Ocab^fl/fl   Ocab^fl/fl;CD4-Cre

NKT cluster 1
CD8 Teff
Stem-like CD4-1
Treg
Th17-like

Stem-like CD4/CD8
NKT cluster 2
gd17
Stem-like CD4-2

Ccr9 stem-like
Migratory CD4/CD8
Proliferating
Memory-like CD8

b TCR Clonotype - Relapse

Ocab^fl/fl

Ocab^fl/fl;CD4-Cre

Large (20 < X ≤ 100)
Medium (5 < X ≤ 20)
Small (1 < X ≤ 5)
Single (X= 1)

c *Ccr9* *Bach2* *Ccr7* *Tcf7*

Cd44 *Cxcr6* *Il1r1* *Lgals3*

Ccr9 Stem-like / Stem-like CD4-2 / Th17-like

d Ocab^fl/fl Ocab^fl/fl;CD4-Cre Ocab^fl/fl Ocab^fl/fl;CD4-Cre Ocab^fl/fl Ocab^fl/fl;CD4-Cre

Ccr9 *Tcf7* *Bach2*

e Ocab^fl/fl Ocab^fl/fl;CD4-Cre

f Th17-like

Pdcd1 *Tigit* *Il1r1*

Ocab^fl/fl   Ocab^fl/fl;CD4-Cre