OCA-B promotes pathogenic maturation of stem-like CD4+ T cells and autoimmune demyelination

Erik P. Hughes[1,3], Amber R. Syage[1,3], Elnaz Mirzaei Mehrabad[2,3,4], Thomas E. Lane[5], Benjamin T. Spike[2,3], Dean Tantin[1,3]

[1]Department of Pathology, University of Utah School of Medicine, Salt Lake City, UT, USA. [2]Department of Oncological Sciences, University of Utah School of Medicine, Salt Lake City, UT, USA. [3]Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT, USA. [4]School of Computing, University of Utah, Salt Lake City, UT, USA. [5]Department of Neurobiology and Behavior, School of Biological Sciences, University of California, Irvine, CA, USA.

Correspondence to Dean Tantin:

Huntsman Cancer Institute, 2000 Circle of Hope, Research South Room 3707

Salt Lake City, UT 84112-5550

dean.tantin@path.utah.edu

801-587-3035

Short title: OCA-B promotes CD4+ T cell mediated demyelinating disease

Keywords Adaptive immunity; OCA-B/Bob1/Pou2af1; autoimmunity; multiple sclerosis; experimental autoimmune encephalomyelopathy; CD4+ T cells; relapsing-remitting

Conflicts of interest statement: The authors have declared that no conflict of interest exists

Words in abstract: 195

Words main text: 11,124

Total main display items: 6

Abstract

Stem-like T cells selectively contribute to autoimmunity, but the activities that promote their

pathogenicity are incompletely understood. Here, we identify the transcription coregulator OCA-

B as a driver of the pathogenic maturation of stem-like CD4$^+$ T cell to promote autoimmune

demyelination. Using a human multiple sclerosis (MS) patient dataset, we show that *POU2AF1*,

the gene encoding OCA-B, is elevated in CD4$^+$ T cells from secondary-progressive MS patient

lesions. We show that T cell-intrinsic OCA-B loss protects mice from experimental autoimmune

encephalomyelitis (EAE) while preserving responses to viral CNS infection. In EAE models driven

by antigen reencounter, OCA-B deletion nearly eliminates CNS infiltration, proinflammatory

cytokine production and clinical disease. CNS OCA-B-expressing CD4$^+$ T of mice with EAE

express an encephalitogenic gene program and preferentially confer disease. In a relapsing-

remitting EAE model, OCA-B loss protects mice specifically at relapse. During remission, OCA-B

promotes the expression of *Tcf7*, *Slamf6*, and *Sell* in proliferating CNS T cell populations. At

relapse timepoints, OCA-B loss results in both the accumulation of an immunomodulatory CD4$^+$

T cell population expressing *Ccr9* and *Bach2*, and loss of pro-inflammatory gene expression from

Th17 cells. These results identify OCA-B as a driver of pathogenic CD4$^+$ T cells.

Introduction

2.8 million individuals worldwide suffer from multiple sclerosis (MS), with almost 1 million cases in the United States (1). Despite recent advances, the development of new therapies that block MS while sparing immunosurveillance mechanisms needed to prevent infection and viral recrudescence remains a critical goal. Genome-wide association studies (GWAS) identify the *HLA* loci as the most important determinant of MS pathogenesis, emphasizing the impact of T cells (2, 3). The association with the major histocompatibility complex (MHC) class II locus in particular indicates that myelin-specific CD4[+] T cells are key regulators of disease.

The predominant preclinical model of MS is experimental autoimmune encephalomyelitis (EAE), in which laboratory animals are immunized with myelin proteins or peptides to activate antigen-specific CD4[+] T cells that infiltrate the central nervous system (CNS) and mediate disease. EAE within the C57BL/6 mouse background establishes chronic disease. In these models, Th1 and Th17 effector CD4[+] T cell subsets are pathogenetic mediators (4, 5). After accessing the CNS, these cells promote neuroinflammation through the secretion of interferon-gamma (IFNγ), interleukin-17 (IL-17), granulocyte macrophage stimulating-factor (GM-CSF) and other inflammatory mediators and chemoattractants that recruit the cell types responsible for driving demyelination (6–8).

Myelin-reactive T cells from the blood of MS patients show stem/memory-like properties, with reduced need for co-stimulation, suggesting signs of previous activation (9–12). Consistent with these findings, autoreactive and bystander memory-phenotype CD4[+] T cells preferentially transfer demyelinating disease in mouse models (13–15). While these studies have identified vital CD4[+] T cell populations in EAE, further research on the mechanisms sustaining these populations and driving their pathogenic activity is required to develop more precise therapies.

Oct1, a POU domain transcription factor encoded by *Pou2f1*, has been linked to autoimmune disease. GWAS studies have identified Oct1 binding site polymorphisms that associate with a predisposition for autoimmune diseases, including MS (16–19). In CD4[+] T cells, Oct1 promotes the development of both CD4[+] memory (20) and EAE pathogenesis and demyelination (21). In contrast, Oct1 loss preserves immune responses to the neurotrophic viral infection (21). These observations highlight a promising pathway which when targeted may limit autoimmune CNS demyelination without the broad immune suppression elicited by many current therapeutics. However, Oct1 itself is widely expressed and regulates both embryonic development and somatic stem cell function (22–24), making it a poor therapeutic target.

The lymphoid-restricted transcription cofactor OCA-B regulates transcription by docking with Oct1, and in B cells an additional related factor, Oct2 (25). In contrast to Oct1, OCA-B whole-

body knockouts are viable and fertile (26). Polymorphisms in the *Ocab* (*Pou2af1*) locus are

associated with autoimmune diseases, including MS (27, 28). Although its expression is 50-100

fold lower in T cells compared to B cells (29), targeting either OCA-B by genetic deletion in T cells

or its downstream pathways with a membrane-permeable peptide mimic suppresses type-1

diabetes (30). OCA-B interacts with target genes such as *Il2, Ifng, Il21, Csf2 (Gmcsf), Tnfrsf4*

(Ox40), Icos and *Ctla4*, but is dispensable for their expression following primary activation of T

cells. Instead, OCA-B regulates these targets under a narrow range of conditions, for example

following secondary stimulation in culture (20). Many of these targets are directly implicated in

MS pathogenesis (6–8, 31–37). OCA-B controls expression of *Tbx21* (*Tbet*), which encodes a

key regulator of Th1 effector responses, but also promotes Th17 differentiation and interacts with

RORγt, a master regulator of Th17 cells (20, 38, 39). In the context of MS, the only prior OCA-B

work used whole body knockouts and showed partial protection in chronic C57BL/6 EAE models

(39). Cumulatively, these findings indicate that OCA-B plays poorly defined but possibly significant

roles in MS pathogenesis.

Here, we show that OCA-B within the T cell compartment drives EAE by promoting

pathogenic maturation of stem-like CD4[+] T cells. In C57BL/6 mice, T cell OCA-B loss significantly

protects mice from chronic EAE while preserving responses to the neurotrophic coronavirus

JHMV. Using Th1 and Th17 passive transfer assays in which reactive T cells are primed with

antigen in donor mice and in polarizing culture, and reencounter antigen in naïve recipients, we

show that OCA-B is critical for the infiltration of pathogenic cells into the CNS and the

manifestation of clinical disease. Using an OCA-B reporter mouse, we find that OCA-B expressing

CD4[+] T cells taken from MOG peptide primed mice preferentially display pathogenic stem-like

gene expression and preferentially transfer demyelinating disease, highlighting OCA-B as a

marker and promoter of encephalitogenic autoimmune CD4[+] T cell activity. Intriguingly, T cell

conditional OCA-B knockout in EAE models on the autoimmune-prone non-obese diabetic (NOD)

background, which develop relapsing-remitting disease (40), specifically protects against relapse.

Single-cell transcriptomic analysis of CD3ε[+], CNS-infiltrating T cells at remission and relapse

indicates that OCA-B promotes disease relapse through control of stem-like T cell populations,

driving them to pathogenic Th17 differentiation. Cumulatively, these findings mark OCA-B as a

critical mediator of the pathogenic maturation of autoreactive T cells, and a promising potential

therapeutic target.

Results

Elevated expression of the gene encoding OCA-B in CD4+ T cells from secondary-progressive human MS lesions. To determine if *OCAB* (*POU2AF1*) is elevated in human MS samples, we mined high-throughput single-nucleus RNA-seq data of glial and immune cells (including T cells) from a recent study using single-nucleus RNA-seq to evaluate gene expression in brain tissue from 54 MS patients and 26 non-MS controls (41). The number of relapsing-remitting (RRMS) samples was small. The N=1 RRMS sample showed far higher *POU2AF1* expression compared to CTR (Figure 1A). There were more primary-progressive (PPMS) samples, and interestingly only one sample showed significant expression, perhaps reflecting imperfections in diagnoses. Most dramatically, a subset of secondary-progressive (SPMS) samples showed highly elevated OCA-B expression (Figure 1A). We then reclustered CD4$^+$ cell nuclei from this dataset. Uniform Manifold Approximation and Projection (UMAP) feature plots revealed that the elevated *POU2AF1* found in T cell nuclei found from SPMS patients was confined to one specific cluster that also expresses the pathogenic marker *IL1R1* (Figure 1B).

OCA-B promotes chronic EAE in C57BL/6 mice. To understand the impact of OCA-B in T cell mediated autoimmunity, we utilized an *Ocab* conditional knockout allele crossed to the CD4-Cre driver which efficiently deletes OCA-B in T cells (30). We induced EAE in C57BL/6 female *Ocab$^{fl/fl}$*;CD4-Cre mice and littermate *Ocab$^{fl/fl}$* controls through subcutaneous (s.c.) injection of myelin oligodendrocyte glycoprotein (MOG)$_{35-55}$ peptide in Complete Freund's Adjuvant (CFA) followed by intraperitoneal (i.p.) injections of *Bordetella pertussis* toxin. *Ocab$^{fl/fl}$*;CD4-Cre mice showed reduced chronic EAE clinical scores and weight loss compared to *Ocab$^{fl/fl}$* littermate controls (Figure 1C-D). Spinal cords were taken at peak disease (day 15) for immunofluorescence and histological analysis. Myelin basic protein (MBP) immunofluorescence showed a significant decrease in demyelinated white matter in spinal sections from Ocab$^{fl/fl}$;CD4-Cre mice compared to littermate Ocab$^{fl/fl}$ controls while DAPI showed simultaneously decreased immune cell infiltration (Figure 1E-F). Ocab$^{fl/fl}$;CD4-Cre mice also showed a significant decrease in demyelination as measured by luxol fast blue (LFB) in combination with hematoxylin and eosin (H&E) staining (Supplemental Figure 1A-B). Flow cytometric profiling at peak disease showed decreased CD4$^+$ T cell CNS infiltration using *Ocab$^{fl/fl}$*;CD4-Cre mice, however no differences in other infiltrating cell populations or in the relative frequencies of the proinflammatory cytokines IFNy, IL-17, GM-CSF, or CD44 were observed (Figure 1G-H and Supplemental Figure 1C-H). We studied MOG-specific CD4$^+$ T cells using the MHC class II I-A^b MOG$_{38-49}$ tetramers. At peak

disease, *Ocab*^{fl/fl};CD4-Cre mice showed a trending reduction in MOG tetramer-specific cell

frequency (*p*=0.055) and a significant reduction in numbers (*p*=0.003) compared to controls

(Figure 1I-J). These results indicate that the protection conferred by OCA-B loss in T cells is

marked by modest decreases in MOG-specific CD4$^+$ T cells within the CNS, resulting in fewer

pathogenic T cells driving demyelination.

OCA-B is essential for EAE mediated by adoptive transfer of Th1- and Th17-polarized T

cells. MS is driven by autoreactive CD4$^+$ Th1 and Th17 cells within the CNS that recruit CD8$^+$ T

cells, macrophages, microglia and neutrophils, which together mediate white-matter damage (42).

To investigate the impact of OCA-B on autoimmune Th1 and Th17 populations, we used adoptive

transfer EAE models with female *Ocab*^{fl/fl};CD4-Cre and *Ocab*^{fl/fl} littermate controls. Mice were

primed with MOG$_{35-55}$ in CFA for 14 days. Following MOG priming, CD4$^+$ T cells were purified

from the lymph nodes of primed experimental or control donors. No differences between

genotypes in CD4$^+$ T cell viability, count, or effector cytokine (IFNγ and IL-17) expression were

observed (Supplemental Figure 2A-E).

To investigate if OCA-B expression impacts Th1 or Th17 mediated adoptive transfer EAE,

CD4$^+$ T cells purified from donors were polarized in vitro into Th1 and Th17 cells as previously

described (43, 44). 3$\times$10^6 Th1- or Th17-polarized CD4$^+$ cells were transferred into naïve age-

matched wild-type male C57BL/6 recipients by i.p. injection. EAE onset in mice receiving either

Th1- or Th17-polarized control cells was rapid and severe (Figure 2A-B). In contrast to the mild

protection observed with OCA-B deficiency in standard C57BL/6 EAE, mice adoptively transferred

with T cells lacking OCA-B were almost completely protected from clinical disease regardless of

their polarization state. Interestingly, the ability to initially polarize CD4$^+$ T cells into Th1 and Th17

effectors in vitro was only minimally affected by OCA-B loss (Supplemental Figure 2F-H), whereas

cells taken from the CNS of engrafted recipient mice at day 15 not only showed decreased CNS

infiltration but also reduced IL-17 expression (Figure 2C-F). This effect appeared to be specific to

IL-17, as no significant difference in IFNγ expression was observed (Supplemental Figure 2I).

These results indicate a central role for OCA-B in promoting robust neuroinflammatory T cell

responses upon antigen reencounter in adoptive transfer EAE models.

OCA-B is dispensable for neurotropic viral clearance. To investigate if T cell OCA-B

deletion impacts the ability to mount immune responses and control CNS viral infection, male

Ocab^{fl/fl};CD4-Cre and *Ocab*^{fl/fl} littermate controls were intracranially inoculated with 1500 plaque-

forming units (PFU) of the John H. Muller strain of murine hepatitis virus (JHMV), a β-coronavirus

178 adapted to infect glial cells (astrocytes, microglia and oligodendrocytes) in the mouse brain and

179 spinal cord. Inoculation with JHMV results in acute encephalomyelitis and chronic demyelination

180 similar to autoantigen-driven EAE. Disease severity was determined by evaluating clinical scores

181 over 21 days post-infection (dpi). *Ocab*[fl/fl];CD4-Cre animals showed negligible differences in

182 clinical scores compared to *Ocab*[fl/fl] littermates (Figure 3A), suggesting that OCA-B loss preserves

183 baseline T cell responses to acute infection. Spinal cords were taken on 12 and 21 dpi for

184 immunofluorescence and histological analysis to investigate differences in demyelination. MBP

185 immunofluorescence at 12 dpi showed similar demyelination and immune infiltration in both

186 Ocab[fl/fl] and Ocab[fl/fl];CD4-Cre spinal sections (Figure 3B-C). Similarly, LFB histology of brain

187 tissue at 21 dpi showed no significant differences in demyelination between OCA-B deficient and

188 control infected animals (Supplemental Figure 3A-B). Flow cytometric analysis of brain tissue

189 showed no significant differences in infiltrating CD4$^+$ T cell numbers or IFNγ expression at 7, 12,

190 or 21 dpi (Figure 3D-F, and Supplemental Figure 3C). Similarly, no differences in either total CD8$^+$

191 T cell numbers, or the percentage or numbers of IFNγ-expressing CD8$^+$ T cells was observed

192 (Supplemental Figure 3D-F). Half brain homogenates were collected on 7, 12 and 21 dpi to

193 measure viral titers and clearance. *Ocab*[fl/fl];CD4-Cre and *Ocab*[fl/fl] littermate control animals

194 showed similar titers at dpi 7 and 12, and virus was undetectable at day 21 regardless of genotype

195 (Figure 3G). This result suggests that OCA-B is dispensable for immune responses to and

196 clearance of neurotropic viruses.

197   *OCA-B expressing CD4$^+$ T cells display pathogenic stem-like properties and preferentially*

198 *transfer demyelinating disease.* We used a recently described OCA-B-mCherry reporter mouse

199 (29) to evaluate OCA-B expression within CNS infiltrating T cells. EAE was induced in female

200 C57BL/6 homozygous OCA-B-mCherry reporter mice with MOG$_{35-55}$, and CNS-infiltrating CD4$^+$ T

201 cells assessed by flow cytometry at peak clinical disease. mCherry-positive T cells showed

202 increased CD44 and CD62L co-expression, indicative of increased memory-like cells in the OCA-

203 B expressing population (Figure 4A). The association of OCA-B expression with a memory-like

204 profile was observed in the brain, spine and draining cervical lymph nodes (Figure 4B).

205 Interestingly, the frequency of CD44$^+$CD62L$^-$CD4$^+$ T cells was elevated within the OCA-B

206 expressing population specifically within the cervical lymph nodes (Supplemental Figure 4A).

207   Memory-like CD4$^+$ T cells have been implicated as major orchestrators of autoimmune

208 demyelination in mice and humans (9–12, 14, 15, 45). To see if this memory-like profile of OCA-

209 B expressing CD4$^+$ T cells is specific to CNS infiltrating cells, we performed flow cytometry using

draining lymph node and splenic T cells from MOG-primed OCA-B-mCherry reporter mice. Similar

to OCA-B[+] CNS infiltrating cells, OCA-B expressing CD4[+] T cells taken from the cervical lymph

node and spleen showed increased CD44 and CD62L co-expression (Figure 4C-D).

We next evaluated OCA-B-mCherry positive and negative populations by RNA

sequencing of CD4[+] T cells isolated from the lymph nodes of 14-day MOG-primed mice. ~1500

genes were upregulated and ~350 downregulated in OCA-B expressing compared to OCA-B

negative cells (Supplemental Table 1). Upregulated genes in OCA-B expressing cells included

Cxcr6, *Slamf6*, *Il23r*, *Rorc*, *Ifng*, *Bcl6*, and *Il17a* while downregulated genes included *Foxp3*, *Il7r*

and *Ccr9* (Figure 4E-F). Expression of *Tcf7* (which encodes TCF1) was high within both OCA-B

positive and negative populations (Supplemental Table 1). Gene ontology (GO) and pathway

analysis of upregulated genes show enrichment for terms associated with autoimmune diseases

including lupus, autoimmune thyroid disease, and type-1 diabetes (Figure 4G, and Supplemental

Table 2).

To determine if OCA-B expression can be used to prospectively identify viable pathogenic

CD4[+] T cell populations, we used OCA-B-mCherry reporter mice as donors in passive transfer

EAE. 11-week-old male and female littermates homozygous for the OCA-B-mCherry reporter

were primed for 14 days with MOG$_{35-55}$ in CFA as previously described (44). CD4[+] T cells were

isolated from spleen and lymph nodes and sorted based on mCherry expression (Supplemental

Figure 4B). 9×10^5 mCherry negative and positive cells were cultured in Th1 polarizing conditions

and subsequently transferred into age-matched male C57BL/6 recipients as previously described

(14). Disease progression was measured by clinical score and normalized weight. Recipients of

OCA-B-expressing mCherry positive cells showed increased clinical disease severity and weight

loss compared to mCherry negative recipients (Figure 4H and Supplemental Figure 4C). These

findings indicate that OCA-B marks CD4[+] T cells with pathogenic stem-like gene expression and

can be used to prospectively identify peripheral CD4[+] T cells with elevated encephalitogenic

properties.

OCA-B promotes neuroinflammation in relapsing-remitting EAE models via control of

stem/memory-like T cell populations. In contrast to the chronic form of EAE that develops in

C57BL/6 mouse models, EAE on the NOD background exhibits a relapsing-remitting pattern that

more closely mimics human disease (40). To study the impact of OCA-B loss in T cells on mouse

models that more closely resemble human MS disease patterns, we utilized *Ocab* T cell

conditional knockout mice crossed to the NOD background (30). EAE was induced in 15-18 week

old female NOD.*Ocab*^{fl/fl};CD4-Cre and littermate control NOD.*Ocab*^{fl/fl} mice with MOG$_{35-55}$. Clinical

scores and weights were recorded for 47 days to observe initial disease onset, relapse and

relapse recovery. No significant differences in clinical scores were observed between groups at

disease onset (day 10-19), however following the primary remission, NOD.*Ocab*^{fl/fl};CD4-Cre

animals were almost completely protected against disease relapse (Figure 5A). Individual NOD

mouse clinical scores are shown in Supplemental Figure 5A-B. Normalized weights of animals

also reflected the protective effect of OCA-B loss in T cells at relapse (Figure 5B).

To understand differences in CNS-infiltrating T cells underlying relapse protection at a

population and transcriptional level, we performed single-cell RNA sequencing (scRNA-seq) of

CD3ε$^+$ T cell isolates from CNS (brain + spinal cord) at remission (day 24) and peak relapse (day

33). Cells from 3-6 NOD.*Ocab*^{fl/fl};CD4-Cre and NOD.*Ocab*^{fl/fl} littermate controls were collected and

combined for analysis. 7799 (remission) and 9224 (relapse) *Ocab* knockout, and 6009 (remission)

and 7525 (relapse) control cells were used for analysis, with an average per cell read depth

ranging from 90,000 to 190,000. For remission, UMAP clustering using both genotypes revealed

16 clusters corresponding to a broad T cell repertoire consisting of natural-killer T (NKT) cells (the

largest cluster) and different CD4$^+$ and CD8$^+$ subsets (Figure 5C). Noteworthy clusters based on

mRNA expression included proliferating (*Ki67*, *Pcna*, *Mcm2*), Th1-like (*Tbx21*, *Ifng*, *Tnf*, *Il12rb2*)

and Treg (*Foxp3*, *Ctla4*) (Supplemental Table 3). Feature plots for *Foxp3* and *Mki67* are shown

in Figure 5D. Additional feature plots for *Cxcr6*, *Il23r* and *Bach2* are shown in Supplemental Figure

5C. Notably there was a large decrease in NKT cell numbers in knockout mice (36% control vs.

14% experimental). Within this cluster, changes in gene expression between control and

conditional knockout groups were minimal. Most NKT cells from both groups showed a limited

inflammatory cytokine response, consistent with the lack of clinical disease at remission

(Supplemental Figure 5D). The decrease in the large NKT population in the experimental group

proportionally increases the percentage contribution of the other clusters, and therefore similar or

decreased percentages of these cells identifies reduced populations which correlate with

protection from relapse, e.g. proliferating T cells. Other clusters such as Tregs increased

disproportionately. Differential gene expression analysis of key clusters (Treg, Proliferating, and

Th1-like) revealed increased expression of *Slamf6* in Tregs lacking OCA-B, while OCA-B deficient

proliferating and Th1-like cells showed decreased expression of genes associated with

quiescence (*Btg1* and *Samhd1*), memory/stemness (*Slamf6*, *Tcf7*, and *Sell*) and pathogenicity

(*Tcf7* and *Tox* co-expression, Figure 5E). Spectral cytometry showed no significant difference in

the frequency of CNS infiltrating CD4$^+$ T cells between groups at remission (Figure 5F). However,

OCA-B deficient CD4$^+$ T cells did show a significant reduction in CXCR6 and IFNγ/GM-CSF co-

expression, indicative of reduced Th17 pathogenicity (Figure 5G-I). These results indicate that

loss of OCA-B in T cells selectively reduces proliferation and differentiation of pathogenic

stem/memory-like CD4$^+$ T cells during disease remission.

At relapse, multiple CNS-infiltrating stem-like and effector CD4$^+$ and CD8$^+$ control T cell

populations were present (Figure 6A and Supplemental Figure 6A, Supplemental Table 4). TCR

sequencing revealed an expansion of a subset of TCR clones in the OCA-B deficient CD8 clusters

while most other clusters were highly polyclonal (Figure 6B). Comparing the 10 most expanded

TCR clonotypes between knockouts and controls, there were several clonotypes that were shared

but differentially expanded (Figure S6B and Supplemental Table 5). A "Ccr9 stem-like" cluster

expressing the gut homing chemokine receptor *Ccr9*, as well as *Ccr7*, *Bach2*, and *Tcf7* notably

appeared at relapse (Figure 6C and Supplemental Table 4). TCR clonotype analysis indicated

that this population was highly polyclonal (Figure 6B). Additionally, a "Stem-like CD4-2" cluster

expressing *Tcf7*, Ccr7, *Bach2, Cd44* and low-level *Cxcr6*, and a prominent "Th17-like" cluster

expressing *Tcf7, Cd44, Il1r1, Lgals3*, and higher levels of *Cxcr6* was identified (Figure 6C). This

latter cluster likely contains pathogenic effector cells.

In the absence of OCA-B, NKT cell populations were normalized compared to the

remission timepoint, and several clusters showed major changes in the percentage of cells

including most prominently the Ccr9-stem-like cells which were much more prevalent (Figure 6D).

Feature plots comparing OCA-B T cell deficient mice to littermate controls for *Ccr9* and the

transcription factors *Tcf7* and *Bach2* are shown in Figure 6D. The cells were also marked by the

expression of *Cd4*, *Tox, Ccr7, Ccr4* and limited expression of *Sell* (Figure 6C).

RNA velocity analysis of spliced and unspliced transcripts (46) identified clusters of T cells,

most bearing static gene expression profiles consistent with terminal differentiation, but some with

differentiating or transitional gene expression (Figure 6E). Notably, OCA-B knockout cells showed

increased transcriptional stasis within the Ccr9-stem-like and proliferating clusters, and a

decreased transcriptional directionality of Th17 cells towards gd17 cells, the latter of which

express high levels of pathogenic markers including *Il1r1*, *Cxcr6*, and *Lgals3* (Supplemental

Figure 6A). Pseudotime analysis identified a transcriptional capacity in the control *Ccr9*

expressing population to differentiate into pathogenic Th17-like cells, whereas cells lacking OCA-

B were limited in this capacity, possibly resulting in the accumulation of these cells (Supplemental

Figure 6D). These changes culminated in the Th17-like effector knockout cells expressing

reduced pathogenicity-associated genes such as *Pdcd1*, *Tigit* and *Il1r* (Figure 6F). Together

these results suggest OCA-B mediates the transition of non-pathogenic stem-like cells to

pathogenic Th17 cells which ultimately drive disease relapse.

Discussion

The development of therapies that inhibit autoimmunity while minimizing impacts on normal immune function remains a critical unmet goal in the field. In this study, we show that expression of mRNA encoding the transcriptional co-regulator OCA-B is elevated in CD4$^+$ T cells from SPMS brain tissue, and that OCA-B promotes the pathogenic maturation of T cells and establishment of CNS autoimmunity, in particular episodes of relapse. Loss of OCA-B attenuates these populations in chronic EAE, in adoptive transfer EAE, and in NOD.EAE during episodes of remission. During relapse, OCA-B T cell knockout mice are almost completely protected, and accumulate a population of CD4$^+$ T cells expressing *Ccr9*, *Bach2*, and *Tcf7*. OCA-B expression also marks viable pathogenic stem/memory-like CD4$^+$ T cells that preferentially transfer EAE. Furthermore, the effects of modulating OCA-B are selective, as responses to infection with a neurotropic coronavirus that generates similar clinical symptoms but lacks an autoantigen were largely unaffected.

OCA-B is a member of a small family of transcription co-regulators that also includes OCA-T1, OCA-T2 and IκBζ (47, 48). It docks with Oct1 upon CD4$^+$ T cell activation to directly regulate immunomodulatory genes such as *Il2*, *Ifng*, *Il17a*, Tbx21 (T-bet) and *Csf2* (*Gmcsf*). Rather than functioning as a primary inducer of these genes, it is required for their robust expression upon antigen reencounter (20, 30). Notably, unlike transcription factors such as NF-AT, AP-1 and NF-κB, Oct1 and OCA-B are dispensable for normal primary T cell responses (20). These findings predict important roles for OCA-B in CD4$^+$ T cells specifically in situations of antigen reencounter, which is a necessary feature of autoimmunity. Conditional T cell knockout of Oct1 reduces EAE severity, with reduced proinflammatory cytokine expression and increased anergy (21). Deletion of Oct1 in T cells was dispensable for neurotropic viral clearance (21), however the ubiquitous expression of Oct1 in adult tissue limits its translational potential. In contrast, OCA-B expression is largely confined to B and T cells and whole-body knockout mice are viable and fertile (49). OCA-B is highly expressed in B cells and required for germinal center formation (49, 50). It is expressed in CD4$^+$ T cells upon activation at levels 50-100-fold less than in B cells (30, 51, 52). This suggests that with proper therapeutic dosing, OCA-B could be inhibited selectively in T cells with minimal impact to the B cell compartment. Short-term administration of an OCA-B peptide mimic that inhibits OCA-B's downstream effector functions reverses spontaneous type-1 diabetes in NOD mice (30). However, toxicity limits its efficacy, including within EAE models. Further research is required for the development of safe and effective OCA-B inhibitors.

OCA-B whole body knockout animals are partially protected from EAE (39), however, the
known activity of OCA-B in B cells made it unclear if the protection is T cell-derived. Here we
show that OCA-B promotes EAE pathogenesis through T cells. Opportunistic infections and
reactivation of latent CNS-tropic viruses are significant barriers for MS therapeutics that focus on
global immune suppression as a therapeutic mechanism. When OCA-B T cell conditional
knockout mice are challenged with the neurotropic coronavirus JHMV, there is little difference in
their ability to mount immune responses and control virus replication. These data are consistent
with previous findings that OCA-B loss in T cells preserves lymphocytic choriomeningitis virus
(LCMV) acute viral infection response, instead selectively impacting memory recall responses
(20, 29). This finding provides a potential therapeutic window in which targeting the low levels of
OCA-B in T cells could be used to blunt autoimmunity while sparing baseline immune responses.

Unlike the significant but modest protection observed in chronic C57BL/6 EAE, OCA-B
loss in adoptive transfer EAE almost completely protects recipient animals from disease. This
may indicate that adoptive transfer EAE models involve more autoantigen re-exposures, as
transferred T cells must re-encounter antigen in a naïve host before promoting disease. Protection
was observed regardless of whether donor cells were Th1- or Th17-polarized, indicating that
OCA-B functions through a mechanism that parallels rather than enforces the Th1/Th17
paradigm. We therefore sought other mechanisms that might explain why OCA-B T cell knockouts
are so strongly protected.

Recent findings indicate an emerging role for stem/memory-like T cells in promoting
autoimmunity (45, 53, 54). For example, myelin-specific CD4+ T cells in MS patients show signs
of a previously activated, memory-like phenotype compared to healthy controls (9, 10, 12).
Furthermore, memory-like Th1 and Th17 cells are increased in the peripheral blood of MS patients
and display increased pathogenicity compared to conventional Th17 Cells (55). Similarly, in EAE
models, peripheral central memory-like CD4+CD44hiCD62Lhi T cells confer increased disease
severity upon transfer (14). Both Oct1 and OCA-B promote the functionality of long-lived CD4+
memory T cells following antigenic rechallenge (29, 56). An OCA-B-mCherry reporter mouse
allele, which effectively labels memory progenitor CD4+ T cells during acute infection (29), also
labels CD4+ T cells that preferentially confer EAE. OCA-Bhi cells taken from the CNS at peak EAE
selectively express stem/memory markers. Further profiling OCA-Bhi CD4+ T cells suggests that
OCA-B promotes the transition of non-pathogenic stem-like cells to a pathogenic state associated
with markers such as *Cxcr6*, *Slamf6*, *Ifng*, and *Il17a*. Elevated *Pou2af1* (*Ocab*) transcripts are
also found in pathogenic Th17 cell populations (57). The connection between OCA-B expression

and pathogenic memory/stem-like T cells likely has not been appreciated before because OCA-

B is expressed at levels too low to be reproducibly captured with many current single-cell RNA-

seq technologies.

Relapsing-remitting EAE models have been developed using the Swiss Jim Lambert (SJL)

and autoimmune prone non-obese diabetic (NOD) backgrounds (40, 58). In NOD mice, MOG$_{35\text{-}55}$

EAE has been described as relapsing-remitting with minimal initial disease that increases in

severity upon relapse (59, 60). Using this model with the conditional *Pou2af1* allele backcrossed

to the NOD background (30), we found that OCA-B T cell loss minimally affects initial clinical

presentation, but instead protects animals specifically at relapse. The finding that OCA-B loss in

conventional C57BL/6 EAE results in partial protection intermediate between the initial and

relapse phases of NOD.EAE suggests that chronic C57BL/6 EAE may not model either initial

presentation or relapse accurately, but instead represents a superposition of the two.

Single-cell RNA expression profiling at remission identifies a significant loss of *Slamf6*,

Tcf7 (encoding TCF1) and *Sell* from proliferative stem-like CD4$^+$ T cells in the OCA-B deficient

condition. While the majority of research on TCF1-expressing stem-like T cells focuses on CD8$^+$

populations with the capacity for self-renewal and differentiation into effector cells, analogous

descriptions have been made for CD4$^+$ populations (45, 61, 62). Indeed, autoimmune CD4$^+$ T

cells expressing these memory/stem-like markers have been reported as an encephalitogenic

reservoir that play a key role in EAE pathogenicity (45). In the same study, bulk RNA sequencing

associated OCA-B expression within this stem-like reservoir (45).

Single-cell RNA-seq at relapse identified multiple emergent populations of effector T cells

compared to remission. Notably, there was also an accumulation of a unique population of CD4$^+$

cells in OCA-B T cell deficient mice. These cells are conspicuously marked by expression not

only of *Ccr4*, but also of *Ccr9*, which encodes a gut-homing chemokine receptor. CCR9$^+$ memory-

like CD4$^+$ T cells have been linked to MS and EAE protection (63). The cells also express *Bach2*,

a transcription regulator which was recently associated with non-pathogenic immunomodulatory

Th17-polarized cells in the context of C57BL/6 EAE (64). *Bach2* expression was largely

concentrated in the "CCR9 stem-like" cluster, which is more numerous in the knockout condition.

Consistently, in bulk RNA-seq *Bach2* mRNA was upregulated in OCA-B$^-$ cells (p<10^{-13}), although

with a small (+1.74) fold change (Supplemental Table 1). Similarly, recent work associates

Pou2af1 (*Ocab*) with pathogenic Th17/Th1 cells compared to the non-pathogenic *Bach2*

expressing cells (64). In B cells, OCA-B is known to regulate *Bach2* (65), however the interplay

between OCA-B and Bach2 and how this influences the pathogenic maturation of Th17 cells

merits additional research. Velocity and pseudotime analysis indicates that these cells are

normally transcriptionally inclined to progress towards an highly pathogenic Th17 population, but

that progression stalls in mice whose T cells lack OCA-B. Consistently, the Th17 effector

population in OCA-B T cell knockout mice lacks expression of key genes such as *Il1r1* and *Pdcd1*,

both of which are associated with pathogenicity in CNS-infiltrating Th17 cells[48,64]. The cumulative

alterations in these populations in OCA-B knockouts likely explains why these mice are strongly

protected against disease relapse. In conclusion, this study indicates a prominent role for OCA-B

in driving stem-like CD4[+] T cell populations towards a pathogenic state while playing a minimal

role in acute antiviral responses to a CNS infection.

Methods

Sex as a biological variable. The objective of this study was to determine the role of the

transcriptional regulator OCA-B in demyelinating disease. OCA-B T cell conditional knockout mice

and OCA-B reporter mice were utilized in multiple mouse demyelinating disease models to

evaluate the impact of T cell OCA-B expression on clinical disease manifestation and the

phenotypes of neuroinflammatory cells, in particular T cells. Both male and female animals were

used for experiments except for active EAE experiments where only female animals were used

as these are the better model spinal cord demyelination and inflammation.

Analysis of publicly available human control and MS single nuclei RNAseq dataset. Initial

evaluation of *POU2AF1* expression in T cells was conducted in the interactive web browser

created by the authors of the dataset available at https://malhotralab.shinyapps.io/MS_broad/

(41). Briefly, *POU2AF1* differential gene expression was filtered to show T cell gene expression

by diagnosis with a minimal threshold cutoff of 10 cells per sample. Cleaned aligned matrix files

were then downloaded from https://zenodo.org/records/8338963 and subsequently processed by

using the 10X Genomics CellRanger pipeline and further analyzed using the 10X Genomics

Loupe Browser 7 software. 19,906 CD4$^+$ cells were then subclustered and evaluated for

POU2AF1 and *IL1R1* gene expression by diagnosis.

Chronic C57BL/6 EAE. 11-12 week-old female C57BL/6 littermate animals were used for

all experiments. For adoptive transfer experiments using the OCA-B-mCherry reporter allele (29),

11 week-old females were used for chronic EAE and 9 week-old males were used as MOG$_{35-55}$

primed adoptive transfer donors. EAE was induced on day 0 by two 100 μL s.c. injections into

both hind flanks of 0.5 μmol/mL MOG$_{35-55}$ peptide (University of Utah Peptide Core) emulsified in

complete Freud's adjuvant (CFA), comprised of incomplete Freund's adjuvant (Cat#

77145;ThermoFisher) together with 4 mg/mL heat-killed *M. tuberculosis* (Cat# DF3114338; Fisher

Scientific). Animals were given two 400 ng i.p. injections of *B. pertussis* toxin (Cat# 181; List

Biological Laboratories) on days 0 and 2.

Clinical Scoring. EAE clinical scores were determined using a 0-5 scale with 0.5 scoring

intervals: 0.5, tip of tail is limp or tail muscle strain observed; 1, complete limp tail; 1.5, complete

limp tail and at least one hind limb consistently falls through wire rack when animal is dropped; 2,

complete limp tail and one hind leg dragging without movement beyond hip; 2.5, complete limp

tail and dragging of both hind limbs without movement beyond hip; 3, complete hind limb paralysis;

3.5, complete hind limb paralysis with flat laying hind quarters and hump in upper torso; 4,

complete hind limb and partial forelimb paralysis, minimal movement around cage; 4.5, complete

hind limb and partial forelimb paralysis, no movement around cage; 5, seizure or death. Animals

scoring a $\geq$4 for more than 48 hours were euthanized and scored 5 for the remainder of the

experiment. JHMV clinical disease was evaluated using previously described 0-4 point scale (67).

Immunofluorescence and Histology. Spinal columns were isolated from C57BL/6 Ocab[fl/fl]

and Ocab[fl/fl];CD4-Cre animals after EAE or JHMV induced demyelination and fixed overnight in in

4% paraformaldehyde at 4C. Spinal cords were subsequently isolated and cryoprotected in 30%

sucrose for 3-5 days. Thoracic and lumbar sections were embedded in optimum cutting

temperature (OCT) compound (ThermoFisher Scientific) and frozen and stored at -80C. 8 μm

spinal sections were cut by cryostat and stained with Luxol fast blue (Cat# 212170250;

ThermoFisher) along with hematoxylin (Cat# 8495; ENG Scientific) and eosin (Cat# E511-25;

Fisher Chemical). Demyelination was assessed by dividing the area of demyelinated white matter

by total white matter as previously described (68, 69). For immunofluorescence, spinal cord

sections were blocked in 5% normal goat serum (NGS; Cat# 005-000-121, Jackson

ImmunoReseach) and 0.1% Tween-20 for 1 hour at room temperature. Sections were stained

overnight at 4°C with primary antibody rabbit anti-MBP (1:1000; Cat# PA1050, Boster), followed

by secondary antibody goat anti-rabbit conjugated with Alexa Fluor 488 (1:750; Cat# A11008,

Invitrogen) for 2 hours at room temperature. Primary and secondary antibodies were diluted in

1% bovine serum albumin (BSA) and 0.1% Tween-20. Slides were cover-slipped using

Fluoromount-G with DAPI (Cat# 00495952, Invitrogen). Sections were imaged with a Zeiss

Axioscan 7 Microscope Slide Scanner using a 20× objective.

Flow cytometry and spectral cytometry. Spinal cords and brains were harvested at peak

disease or specified timepoint and mechanically dissociated by passing through a 100 μm cell

strainer. A 30/70% Percoll gradient (Cat# 17544501; Cytiva) was used to enrich mononuclear

cells. Cells from collected spleens and lymph nodes (cervical, brachial, axillary and inguinal) were

passed through a 70 μm strainer and erythrocytes lysed by ammonium-chloride-potassium lysis

buffer (150 mM NH_4Cl, 10 mM $KHCO_3$, 0.1 mM Na_2EDTA). For intracellular staining, cells were

cultured in complete media (RPMI 1640, 10% FBS, 1% Pen/Strep, Glutamax) supplemented with

1 μL/mL brefeldin A (Cat# 555029; Golgiplug, BD), 50 ng/mL phorbol myristate acetate (Cat#

P1585; Sigma-Aldrich) and 1 μg/mL ionomycin (Cat# I0634; Sigma-Aldrich) for 4 hr. Cultured

cells were fixed using cytofix/cytoperm (Cat# 554714; BD) according to manufacturer's protocol

and stained for intracellular cytokines in perm/wash buffer. Flow cytometry samples were profiled

using a BD Fortessa (BD Biosciences, Figure 2, 3, 4) or an Aurora spectral flow cytometer (Cytek,

Fig 1 and 5). FlowJo software (BD Biosciences) was used for data analysis.

Pre-polarization profiling and mixed adoptive transfer. 10 week-old C57BL/6 *Ocab$^{fl/fl}$* and

Ocab$^{fl/fl}$;CD4-Cre littermates were immunized with MOG$_{35-55}$ in CFA as described above. After 14

days, spleen and lymph node (cervical, axillary, brachial and inguinal) cells were isolated and

cultured in complete media (RPMI 1640, 10% FBS, 1% Pen/Strep, Glutamax) supplemented with

1 μL/mL brefeldin A (Cat# 555029; Golgiplug, BD), 50 ng/mL phorbol myristate acetate (Cat#

P1585; Sigma-Aldrich) and 1 μg/mL ionomycin (Cat# I0634; Sigma-Aldrich) for 4 hr. Cells were

stained for surface markers, fixed using the Foxp3/Transcription Factor Staining Buffer Kit

(Cat#00-5523-00, ThermoFisher), and stained for intracellular markers (IFNγ and IL-17).

Th1 and Th17 polarization and adoptive transfer C57BL/6 EAE. 10-12 week-old female

C57BL/6 littermate animals were used for all experiments. Donor animals were immunized with

MOG$_{35-55}$ in CFA as above. 10-14 days post-immunization, spleen and lymph node (cervical,

axillary, brachial and inguinal) cells were collected and placed into mixed culture with MOG

peptide and cytokines at 5.0×10^6 cells/mL. For Th1 conditions, complete media was

supplemented with 50 μg/mL MOG$_{35-55}$, 6 ng/mL rmIL-6 (Cat# 216-16; Peprotech) and 2 ng/mL

rmIFNγ (Cat# 315-05; Peprotech). For Th17 conditions, complete media was supplemented with

50 μg/mL MOG$_{35-55}$, 8 ng/mL rmIL-23 (Cat# 1887-ML-010; R&D Systems), 10 ng/mL rmIL-1α

(Cat# 211-11A; Peprotech), and 10 μg/mL anti-IFNγ (Cat# BE0055; BioXcell XMG1.2). After 4

days of respective culture, CD4$^+$ T cells were purified using negative magnetic selection and

resuspended in PBS at 1.5×10^6 cells/mL. 100 μL cell suspension was injected i.p. into age-

matched male C57BL/6 recipients (~13 week-old at time of disease induction). Clinical disease

was scored for 15 days.

JHMV. Ocab$^{fl/fl}$ and *Ocab$^{fl/fl}$*;CD4-Cre C57BL/6 mice were anesthetized using isoflurane.

Mice were intracranially (i.c) injected with 1500 plaque-forming units (PFU) of JHMV (strain V34)

suspended in 30 μL Hank's balanced salt solution (ThermoFisher). Clinical disease was assessed

for 21 days using a previously described scale (67). Mice were sacrificed at days 7, 12, and 21

post-infection to assess viral titers within brain homogenates using a plaque assay previously

described (70). Spinal cords were taken on day 12 and 21 post-infection to assess demyelination

by DAPI/MBP and luxol fast blue and hematoxylin/eosin histology respectively.

OCA-B-mCherry adoptive transfer EAE. OCA-B-mCherry reporter allele animals (29) were

primed for 14 days with MOG$_{35-55}$ in CFA as described above. After priming, spleen and lymph

nodes were collected and CD4+ T cells were purified by magnetic negative selection (Miltenyi Bio

kit and LS columns). Non-CD4+ T cells were removed from the LS column and placed on ice for

subsequent in vitro culture. Purified CD4+ T cells were separated by OCA-B-mCherry expression

(PE-Texas Red) using flow automated cell sorting (FACS). OCA-B-mCherry positive and negative

CD4+ T cells were placed into mixed culture in complete medium with the previously purified non-

CD4+ T cells at a 1:10 ratio (CD4+ to non-CD4+) at 1.0×10^6 cells/mL. The culture was

supplemented with 20 μg/mL MOG$_{35-55}$ and 0.5 ng/mL IL-12p70 (Cat# 210-12; Peprotech). After

two days of culture, CD4+ T cells were again purified through negative magnetic selection. Purified

CD4+ T cells from OCA-B-mCherry positive and negative cultures were resuspended in PBS at

8.8×10^6 cells/mL. 100 µL cell suspension was injected i.p. into age-matched male C57BL/6

recipients. Clinical disease was assessed for 24 days.

Bulk RNA-seq. OCA-B-mCherry reporter mice (n=9) were primed for 14 days with MOG$_{35-}$

$_{55}$ in CFA as described above. After priming, lymph nodes (cervical, axillary, brachial and inguinal)

were isolated and mCherry positive and negative CD4+ T cells were collected by FACS. Total

RNA, three groups (pooled 3 mice each), was isolated using the Quick RNA micro kit (Cat#

R1050; Zymo Research) with DNase treatment (Cat# 79254; Qiagen). RNA concentration was

measured with a Qubit RNA HS Assay Kit (Cat# #Q32855; Fisher Scientific). RNA quality was

evaluated with an RNA ScreenTape Assay (Cat# 5067-5579, 5067-5580; Agilent Technologies).

Total RNA samples (5-500 ng) were hybridized with NEBNext rRNA Depletion Kit v2 (Cat# E7400;

New England Biolabs) to diminish rRNA from the samples. Stranded RNA sequencing libraries

were prepared as described using the NEBNext Ultra II Directional RNA Library Prep Kit for

Illumina (Cat# E7760L; New England Biolabs). Purified libraries were qualified on an 4150

TapeStation (Agilent Technologies) using a D1000 ScreenTape assay (Cat# 5567-5582, 5067-

5583; Agilent Technologies). The molarity of adapter-modified molecules was defined by

quantitative PCR using the Kapa Biosystems Kapa Library Quant Kit (Cat# KK4824Roche).

Individual libraries were normalized to 5 nM. NovaSeq 150×150 bp sequencing libraries were

chemically denatured and applied to an Illumina NovaSeq flow cell using the XP workflow

(20043131). Following transfer to an Illumina NovaSeq 6000 instrument, 150×150 cycle paired-

end sequencing was performed using a S4 reagent Kit v1.5 (Cat# 20028312; Illumina). 8-11

million reads were generated per sample and were aligned to the *GRCm38* mouse reference

genome. Differential gene expression and pathway analysis were conducted using the DESeq2

and Enrichr R packages.

NOD-EAE. 15-18 week-old female littermate *Ocab^{fl/fl}* and *Ocab^{fl/fl}*;CD4-Cre animals
backcrossed to a NOD strain background (30) were used for all experiments. EAE was induced
by two 100 μL s.c. injections into both hind flanks of 0.5 μmol/mL MOG$_{35-55}$ peptide emulsified in
incomplete Freuds adjuvant (Cat# 77145; ThermoFisher) supplemented with 4 mg/mL heat-killed
M. tuberculosis (Cat# DF3114338; Fisher Scientific) to form CFA. Animals were given two 400 ng
i.p. injections of *Bordetella pertussis* toxin (Cat# 181; List Biological Laboratories) on days 0 and
2. Clinical disease was assessed over 47 days using a 5-point scale described below. Remission
timepoints were determined to be between days 22-26 while relapse was determined to be
between days 26-39.

Single-cell RNA-seq. NOD.*Ocab^{fl/fl}* and NOD.*Ocab^{fl/fl}*;CD4-Cre animals were induced with
EAE and monitored for disease progression. Brains and spinal cords were isolated on day 24
(remission) and day 33 (relapse), and processed for FACS sorting. Viable cells were sorted by
CD3ϵ using a FACSAria (BD Bioscience). Cells from each condition were isolated from 3-4 mice
and combined. Cells were processed using the 5' 10X Genomics Chromium platform according
to the manufacturer's instructions. Paired-end high-throughput sequencing (125 cycles) was
performed using a NovaSeq instrument (Illumina). Sequencing reads were processed by using
the 10X Genomics CellRanger pipeline and further analyzed using the Seurat R package.
Analysis of cells used a standard filtering protocol removing cells with unique feature counts of
>5,000 or <100, as well as cells with >10% mitochondrial counts (indicative of dead cells). No
more than 10% of total cells were removed by this process. Cells were subjected to unsupervised
hierarchical clustering followed by uniform manifold approximation and projection (UMAP) to
visualize clusters with similar gene expression and their relative positions. RNA velocity analysis
was performed using velocyto.R version 0.6 to model cell state transitions using the guidelines
provided in Manno et al. (46) and the associated repository at https://github.com/velocyto-
team/velocyto.R. High dimensional velocity vectors were graphed on the predefined Seurat
UMAP projection using localized average grid projections. Pseudotime analysis was conducted
using Seurat and Monocle3 packages. Integrated data was divided into subset based on genotype
and Monocle3 was used to calculate trajectory, order cells and plot pseudotime for each condition.

Statistics. Multiple two-tailed student's t-tests were used to determine statistical
differences in EAE clinical scores and all error bars denote $\pm$SEM unless otherwise noted. Two-
way ANOVA was used to determine differences in weight loss throughout disease progression.
GraphPad Prism software was used for all statistics and graphing. Two-tailed student's t-tests

were used to determine statistical differences in cell populations observed by flow cytometry and

all error bars represent ±SD unless otherwise noted. For all figures, ns = $p > 0.05$, * = $p \leq 0.05$, ** =

$p \leq 0.01$, *** = $p \leq 0.001$, **** = $p \leq 0.0001$.

Study approval. All animal experiments were performed in strict accordance with the NIH

guide for the Care and Use of Laboratory animals and institutional guidelines for animal care at

the University of Utah under approved protocol #00001553.

Data availability. All RNA sequencing, single-cell sequencing, and single-cell T cell V(D)J

sequencing data associated with the findings of this study have been deposited to NCBI's GEO

database under accession code GSE243727.

Figure legends

Figure 1. OCA-B loss protects animals from chronic EAE. (**A**) CD4[+] T cell nuclei from single-
nucleus RNAseq data of primary-progressive (PPMS), relapsing-remitting MS (RRMS), and
secondary-progressive MS (SPMS) patient lesions and controls (41) were identified and tested in
silico for *POU2AF1* expression, which is displayed as counts per million. (**B**) The CD4[+] T cell
nuclei were reclustered and displayed as UMAP feature plots showing *POU2AF1* (left) and *Il1R1*
(right) expression in CTR and SPMS brain tissue. Red arrow highlights a likely pathogenic CD4[+]
population expressing high levels of *IL1R1*. (**C**) *Ocab^{fl/fl}* (n=16) and *Ocab^{fl/fl}*;CD4-Cre (n=22) mice
were injected with MOG$_{35-55}$ peptide in CFA and *pertussis* toxin to induce EAE. Clinical scores
were evaluated after EAE induction to determine disease progression. (**D**) Animal weights were
recorded to evaluate weight loss as a measurement of disease progression. (**E**) Representative
DAPI/MBP immunostaining of thoracic sections taken from mice 15 days after EAE induction.
Areas of demyelination are marked by decreased MBP staining (green) and often coincide with
increased cellular infiltration (blue). (**F**) Quantification of demyelination in *Ocab^{fl/fl}* (n=5) and
Ocab^{fl/fl};CD4-Cre mice (n=6). (**G**) Brain and spinal cords were isolated on day 15 of EAE and
analyzed by spectral cytometry. Representative flow cytometry plots of CD45[+]TCRβ[+] CNS
infiltrating T cells showing frequency of CD4[+] and CD8[+] T cells. (**H**) Quantification of the number
of CNS infiltrating CD4[+] T cells. *Ocab^{fl/fl}* (n=9) and *Ocab^{fl/fl}*;CD4-Cre (n=13). (**I**) Representative
plots showing frequency of MOG$_{38-49}$ tetramer-positive CD4[+] T cells. (**J**) Quantification of the
frequency and count of MOG$_{38-49}$ tetramer-positive CD4[+] T cells. *Ocab^{fl/fl}* (n=9) and *Ocab^{fl/fl}*;CD4-
Cre (n=13). For clinical scores and normalized weights, values represent mean ±SEM. All other
values represent mean ±SD.

Figure 2. OCA-B promotes Th1 and Th17 adoptive transfer EAE through recall response.
(**A,B**) *Ocab^{fl/fl}* and *Ocab^{fl/fl}*;CD4-Cre mice were primed with MOG$_{35-55}$ peptide in CFA for 10-14
days. Cells from the spleens and lymph nodes of primed mice were isolated and cultured for 4
days in Th1 or Th17 polarizing conditions. 3.0x10[6] Th1- or Th17-polarized cells were injected i.p.
into C57BL/6 wildtype recipient mice (Th1: *Ocab^{fl/fl}* n=5, *Ocab^{fl/fl}*;CD4-Cre n=5) (Th17: *Ocab^{fl/fl}*
n=3, *Ocab^{fl/fl}*;CD4-Cre n=4). Clinical scores were assessed for 14-15 days. (**C**) At Th17 adoptive
transfer EAE endpoint, brains and spinal cords were analyzed by flow cytometry. Quantification
of brain- and spinal cord-infiltrating CD4[+] T cells following Th17 adoptive transfer. (**D**)
Representative flow cytometry plots showing the frequency of IFNγ and IL-17 expressing CD4[+] T

cells within the spines of Th17 recipient mice. (**E**) Quantification of IL-17 frequency in spinal cord

infiltrating CD4$^+$ T cells. (**F**) Quantification of the total count of IL-17$^+$ CD4$^+$ T cells within the spinal

cord of recipient mice. For clinical scores, values represent mean ±SEM. All other values

represent mean ±SD.

Figure 3. OCA-B is dispensable for T cell response to CNS infection with a neurotropic

virus. (**A**) *Ocab$^{fl/fl}$* (n=19) and *Ocab$^{fl/fl}$*;CD4-Cre (n=17) mice were injected intracranially with 1500

PFU JHMV. Clinical scores were recorded for 21 days post infection (dpi). (**B**) Representative

DAPI/MBP immunofluorescence staining of spinal cord sections at 12 dpi. Demyelinated areas

are marked by decreased MBP staining (green) and often coincide with increased cellular

infiltration (blue). (**C**) Quantification of % demyelination in spinal cord sections. (n=7 per group).

Half brains were taken at 7, 12, and 21 dpi for flow cytometric and viral titer analysis. (**D**)

Quantification of CD4$^+$ T cell counts within half brain at 7, 12, 21 dpi. *Ocab$^{fl/fl}$* (n=7-9 per time

point) and *Ocab$^{fl/fl}$*;CD4-Cre (n=6-7 per time point). (**E**) Representative plots showing the

frequency of IFNγ expressing CD4$^+$ T cells 7, 12, and 21 dpi. (**F**) Quantification of frequency of

IFNγ expressing CD4$^+$ T cells. *Ocab$^{fl/fl}$* (n=7-9 per time point) and *Ocab$^{fl/fl}$*;CD4-Cre (n=6-7 per

time point). (**G**) Viral titers of half-brains at dpi 7, 12, and 21 (n.d., not detected). *Ocab$^{fl/fl}$* (n=7-9

per time point) and *Ocab$^{fl/fl}$*;CD4-Cre (n=6-7 per time point). These data represent the combined

results of two independent experiments. For clinical scores, values represent mean ±SEM. All

other values represent mean ±SD.

Figure 4. OCA-B expression marks encephalitogenic stem-like CD4$^+$ T cells. (**A**) OCA-B-

mCherry mice were injected with MOG$_{35-55}$ peptide in CFA and *pertussis* toxin to induce EAE.

Cervical lymph nodes (CLNs), brains, and spines were isolated at peak disease (day 15) and

analyzed by flow cytometry for mCherry, CD44 and CD62L expression. (**B**) CD44$^+$ and CD62L$^+$

expressing CD4$^+$ T cells were quantified based on mCherry expression. Mean values are shown

from n=5 biological replicates. (**C**) OCA-B-mCherry mice were injected with MOG$_{35-55}$ peptide in

CFA, and after 14 days CD4$^+$ T cells within CLNs and spleens were profiled by flow cytometry for

mCherry, CD44, and CD62L expression. (**D**) Quantification of CD44 and CD62L expression of

OCA-B negative and positive CD4$^+$ T cells from MOG-primed OCA-B-mCherry mice. n=3

biological replicates are shown. (**E**) Heatmap showing relative expression of the top 40

differentially expressed genes from OCA-B positive and negative CD4+ T cells from bulk RNA

sequencing of OCA-B positive and negative CD4+ T cells from MOG$_{35-55}$ primed OCA-B-mCherry

reporter mice. (F) Volcano plot of differentially expressed genes between OCA-B positive and

negative T cells (*p*-value < 0.05 and Log2 fold change > 1 are colored). (**G**) Gene Ontology (GO)

terms associated with OCA-B positive and negative groups. (**H**) EAE clinical scores of C57BL/6

wild-type mice following passive transfer of OCA-B-positive or -negative Th1 polarized CD4+ T

cells. Clinical score values represent mean ±SEM. All other values represent mean ±SD.

Figure 5. OCA-B promotes relapsing-remitting EAE through stem-like CD4+ T cells. (**A**)

NOD.*Ocab$^{fl/fl}$* (n=19) and NOD.*Ocab$^{fl/fl}$*;CD4-Cre (n=15) mice were injected with MOG$_{35-55}$ peptide

in CFA and *pertussis* toxin to induce EAE. Clinical scores of animals representing relapsing-

remitting disease progression are shown as a function of time. (**B**) Animal weights were recorded

to determine weight loss throughout initial disease and relapse. (**C**) Singe cell RNA sequencing

was performed on pooled CD3ε+ cells isolated from the brain and spine of NOD.*Ocab$^{fl/fl}$* (n=3) and

NOD.*Ocab$^{fl/fl}$*;CD4-Cre (n=4) 24 days after EAE induction (remission timepoint). Cell populations

were plotted in a UMAP using the Seurat R package and percentages are shown for each cluster.

Clusters were identified through differential gene expression analysis. (**D**) Feature plots

comparing the expression of *Foxp3* and *Mki67* amongst clusters. (**E**) Violin plots showing the

expression of *Btg1*, *Samhd1*, *Slamf6*, *Tcf7, Sell* and *Tox* within the Treg, proliferating, and Th1-

like clusters. (**F**) CNS cells isolated at remission timepoint (day 24) were analyzed by spectral

cytometry. Representative flow cytometry plots and quantification showing the frequency of CD4+

T cells between control and experimental groups. (**G**) Representative flow cytometry plots and

quantification showing the frequency of CXCR6 expressing CD4+ T cells. (**H**) Representative flow

cytometry plots showing the frequency of CD4+ T cells expressing IFNγ and GM-CSF. (**I**)

Quantification of CD4+ T cells expressing IFNγ and GM-CSF. For clinical scores, data represent

mean ±SEM. All other data represents mean ±SD.

Figure 6. OCA-B promotes disease relapse through control of pathogenic stem-like Th17

differentiation. (**A**) Singe cell RNA and TCR sequencing were performed on pooled CD3ε+ cells

isolated from the brain and spine of NOD.*Ocab$^{fl/fl}$* (n=5) and NOD.*Ocab$^{fl/fl}$*;CD4-Cre (n=6) 33 days

after EAE induction (relapse timepoint). Cell populations were plotted in a UMAP using the Seurat

R package and percentages are shown for each cluster. Clusters were identified through

differential gene expression analysis. (**B**) UMAP TCR clonotype expansion between $Ocab^{fl/fl}$ and

$Ocab^{fl/fl}$;CD4-Cre amongst clusters. (**C**) Violin plots showing expression of *Ccr9*, *Bach2*, *Ccr7*,

Tcf7, *Cd44*, *Cxcr6*, *Il1r1* and *Lgals3* in the Ccr9 Stem-like CD4, Stem-like CD4-2, and Th17-like

clusters. (**D**) Feature plots comparing the expression of *Ccr9*, *Tcf7*, and *Bach2* amongst clusters.

(**E**) UMAPs showing RNA velocity of spliced and unspliced transcripts by experimental condition.

(**F**) Violin plots showing expression of *Pdcd1*, *Tigit*, and *Il1r1* in the Th17-like cluster between

control and experimental groups.

Supplemental figure legends

Supplemental Figure 1. Minimal differences in cytokine expressing CD4$^+$ T cell, CD8$^+$ T cell,

B cell, macrophage or microglial cell counts at peak chronic EAE in OCA-B knockouts. (**A**)

Representative spinal cord LFB/H&E staining of thoracic sections taken from mice 15 days after

EAE induction. Areas of demyelination are outlined in black and marked by increased loss of LFB

(blue) staining and visibility of H&E (pink) within the white matter. (**B**) Quantification of the %

demyelination from LFB/H&E histology. (**C**) Quantification of the frequency of CNS infiltrating

CD4$^+$ T cells between Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. (**D**) Quantification of frequency and

count of CD8α positive T cells between Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. (**E**) Quantification of

the number of B cells, macrophages and microglia. (**F**) Representative flow cytometry plots

comparing the expression of CD44$^+$ within CNS infiltrating CD8$^+$ T cells from Ocab$^{fl/fl}$ and

Ocab$^{fl/fl}$;CD4-Cre mice. (**G**) Quantification of the frequency of CNS infiltrating CD44$^+$ CD8$^+$ T cells

within Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. (**H**) Quantification of frequency and count of IFNγ, IL-

17, GM-CSF, and CD44 expressing CD4$^+$ T cells from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. All data

represent mean $\pm$SD.

Supplemental Figure 2. Loss of OCA-B does not strongly alter initial Th1 or Th17

polarization in vitro. (**A**) Experimental schematic for assessing CD4$^+$ T cell differences between

Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$CD4-Cre prior to *in vitro* Th1 or Th17 polarization. Lymph nodes (cervical,

brachial, axillary, and inguinal) were harvested from Ocab$^{fl/fl}$ (n=4) and Ocab$^{fl/fl}$;CD4-Cre (n=4)

mice following 14 days of MOG$_{35-55}$/CFA priming and analyzed by flow cytometry. (**B**) Frequency

of viable cells isolated from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. (**C**) Total CD4$^+$ T cell numbers

isolated from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. (**D**) Quantification of the frequency of IFNγ

expressing CD4$^+$ T cells isolated from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice. (**E**) Quantification of

the frequency of IL-17 expressing CD4$^+$ T cells derived from Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre mice.

(**F**) Representative flow cytometry plots showing IFNγ and IL-17 expressing CD4$^+$ T cells from

Ocab$^{fl/fl}$ (n=5) and Ocab$^{fl/fl}$;CD4-Cre (n=5) mice after Th1 and Th17 *in vitro* polarization. (**G**)

Quantification of IFNγ expressing CD4$^+$ T cells after Th1 polarization. (**H**) Quantification of IL-17

expressing CD4$^+$ T cells after Th17 polarization. (**I**) Quantification of the frequency of IFNγ

expressing CD4$^+$ T cells isolated from the spinal cords of Th17 adoptive transfer EAE recipient

mice. All data represent mean $\pm$SD.

Supplemental Figure 3. Normal immune responses to intracranial JHMV infection in the absence of OCA-B. (**A**) Representative LFB/H&E histology of spinal cord sections at dpi 21. (**B**) Demyelination quantification of LFB/H&E histology at dpi 21. (**C**) Quantification of the number of IFNγ expressing CD4$^+$ T cells at dpi 7, 12 and 21. (**D**) Quantification of the number of CD8$^+$ T cells at dpi 7, 12 and 21. (**E**) Quantification of the frequency of IFNγ expressing CD8$^+$ T cells at dpi 7, 12 and 21. (**F**) Quantification of the number of IFNγ expressing CD8$^+$ T cells at dpi 7, 12 and 21. All data represent mean ±SD.

Supplemental Figure 4. Sorting OCA-B-mCherry reporter T cells and additional analysis of the effects of transferring OCA-B^{hi} versus OCA-B^{lo} CD4$^+$ cells. (**A**) Quantification of the frequency of CD44$^+$ CD62$^-$ CD4$^+$ T cells within the cervical lymph nodes of OCA-B-mCherry reporter mice at peak EAE by OCA-B-mCherry expression. (**B**) Gating strategy for sorting of OCA-B$^+$ and OCA-B$^-$ CD4$^+$ T cells. (**C**) Normalized weights of NOD.Ocab$^{fl/fl}$ and NOD.Ocab$^{fl/fl}$;CD4-Cre mice following EAE induction. Normalized weight is represented by mean ±SEM. All other data represent mean ±SD.

Supplemental Figure 5. Individual NOD mouse EAE clinical scores, additional remission scRNA-seq cluster annotation genes, and heat map of gene enrichments by cluster. (**A**) Clinical scores of individual NOD.Ocab$^{fl/fl}$ mice following EAE induction. (**B**) Clinical scores of individual NOD.Ocab$^{fl/fl}$;CD4-Cre mice following EAE induction. (**C**) UMAP feature plots showing expression of *Cxcr6*, *Il23r*, and *Bach2* amongst clusters at remission timepoint. (**D**) Heatmap showing the top 10 genes expressed by cluster at remission EAE timepoint.

Supplemental Figure 6. Heat map of NOD.EAE relapse gene enrichments by cluster, TCR clonotype overlap between condition, example relapse feature plots, and pseudotime analysis of relapse single cell RNA seq. (**A**) Heatmap showing the top 10 genes expressed by cluster at relapse EAE timepoint. (**B**) Alluvial plot showing top 10 TCR clonotype gene overlap between Ocab$^{fl/fl}$ and Ocab$^{fl/fl}$;CD4-Cre groups. (**C**) UMAP feature showing the expression of *CD4*, *CD8a, Tox*, *Sell*, *Ccr7*, and *Ccr4* amongst clusters at relapse timepoint. (**D**) Pseudotime analysis of relapse single cell from root starting node at or near the *Ccr9*-expressing stem-like cluster.

Acknowledgements

We thank R.M. O'Connell and M. Bettini for critical reading of the manuscript. We thank J. Marvin

and University of Utah Health Sciences Center Flow Cytometry core, and B. Dalley, O. Allen and

the High-Throughput Genomics core. Peptides were synthesized by the DNA/Peptide facility. We

thank T. Lane for assistance with JHMV constructs, protocols and reagents. This work was

supported by grants from the National Institutes of Health/ National Institute of Allergy and

Infectious Diseases (R01AI162929 and R01AI100873) and the Praespero Foundation (Canada)

to DT, and training grants from the National Institute of Neurological Disorders and Stroke

(T32NS115664) to EPH and National Institute of Allergy and Infectious Diseases (T32AI055434)

to ARS.

Author contributions: E.P.H. and D.T. conceived the study. E.P.H. and A.S. performed

experiments. E.P.H, A.S., E.M.M. and B.T.S. analyzed data. T.E.L., B.T.S. and D.T. provided

critical resources and supervision for the study. E.P.H. and D.T. wrote, reviewed and revised the

manuscript.

Conflict of interest

The authors declare that they have no conflicts of interest.

References

References

1. Walton C, et al. Rising prevalence of multiple sclerosis worldwide: Insights from the Atlas of
MS, third edition. *Mult Scler J.* 2020;26(14):1816–1821.

2. Wtccc CC, et al. Genetic risk and a primary role for cell-mediated immune mechanisms in
multiple sclerosis. *Nature.* 2011;476(7359):214–219.

3. IMSGC. A high-density screen for linkage in multiple sclerosis. *Am J Hum Genet.*
2005;77(3):454–467.

4. Adorini L, Guery J-C, Trembleau S. Manipulation of the Th1/Th2 Cell Balance: An Approach
to Treat Human Autoimmune Diseases? *Autoimmunity.* 1996;23(1):53–68.

5. Langrish CL, et al. IL-23 drives a pathogenic T cell population that induces autoimmune
inflammation. *J Exp Med.* 2005;201(2):233–240.

6. Brucklacher-Waldert V, et al. Phenotypical and functional characterization of T helper 17 cells
in multiple sclerosis. *Brain.* 2009;132(12):3329–3341.

7. Montes M, et al. Oligoclonal myelin-reactive T-cell infiltrates derived from multiple sclerosis
lesions are enriched in Th17 cells. *Clin Immunol.* 2009;130(2):133–144.

8. Hartmann FJ, et al. Multiple sclerosis-associated IL2RA polymorphism controls GM-CSF
production in human TH cells. *Nat Commun.* 2014;5(1):5056.

9. Scholz C, et al. Expansion of Autoreactive T Cells in Multiple Sclerosis Is Independent of
Exogenous B7 Costimulation. *J Immunol.* 1998;160(3):1532–1538.

10. Burns JB, Bartholomew BD, Lobo ST. Isolation of CD45RO+, Memory T Cells Recognizing
Proteolipid Protein from Neurologically Normal Subjects. *Cell Immunol.* 2001;212(1):44–50.

11. Pelfrey CM, et al. Quantification of Self-Recognition in Multiple Sclerosis by Single-Cell
Analysis of Cytokine Production. *J Immunol.* 2000;165(3):1641–1651.

12. Lovett-Racke AE, et al. Decreased dependence of myelin basic protein-reactive T cells on
CD28-mediated costimulation in multiple sclerosis patients. A marker of activated/memory T
cells. *J Clin Invest.* 1998;101(4):725–730.

13. Lee H-G, et al. Pathogenic function of bystander-activated memory-like CD4+ T cells in
autoimmune encephalomyelitis. *Nat Commun.* 2019;10(1):709.

14. Williams JL, et al. Memory cells specific for myelin oligodendrocyte glycoprotein (MOG)
govern the transfer of experimental autoimmune encephalomyelitis. *J Neuroimmunol.*
2011;234(1–2):84–92.

15. Elyaman W, et al. Distinct Functions of Autoreactive Memory and Effector CD4+ T Cells in
Experimental Autoimmune Encephalomyelitis. *Am J Pathol.* 2008;173(2):411–422.

16. Maurano MT, et al. Systematic Localization of Common Disease-Associated Variation in

Regulatory DNA. *Science (80-)*. 2012;337(6099):1190–1195.

17. Graham DSC, et al. Evidence for unique association signals in SLE at the CD28–CTLA4–

ICOS locus in a family-based study. *Hum Mol Genet*. 2006;15(21):3195–3205.

18. van Heel DA. Inflammatory bowel disease is associated with a TNF polymorphism that

affects an interaction between the OCT1 and NF-kappaB transcription factors. *Hum Mol Genet*.

2002;11(11):1281–1289.

19. Farh KK, et al. Genetic and epigenetic fine mapping of causal autoimmune disease variants.

Nature. 2015;518(7539):337–343.

20. Shakya A, et al. Oct1 and OCA-B are selectively required for CD4 memory T cell function. *J

Exp Med*. 2015;212(12):2115–2131.

21. Kim H, et al. T cell-selective deletion of Oct1 protects animals from autoimmune

neuroinflammation while maintaining neurotropic pathogen response. *J Neuroinflammation*.

2019;16(1):133.

22. Vázquez-Arreguín K, et al. Oct1/Pou2f1 is selectively required for colon regeneration and

regulates colon malignancy. *PLOS Genet*. 2019;15(5):e1007687.

23. Perovanovic J, et al. Oct1 cooperates with the Smad family of transcription factors to

promote mesodermal lineage specification. *Sci Signal.* 2023;16(781).

https://doi.org/10.1126/scisignal.add5750.

24. Sebastiano V, et al. Oct1 regulates trophoblast development during early mouse

embryogenesis. *Development*. 2010;137(21):3551–3560.

25. Hwang SS, et al. Role of OCT-1 and partner proteins in T cell differentiation. *Biochim

Biophys Acta - Gene Regul Mech*. 2016;1859(6):825–831.

26. Kim U, et al. The B-cell-specific transcription coactivator OCA-B/OBF-1/Bob-1 is essential

for normal production of immunoglobulin isotypes. *Nature*. 1996;383(6600):542–547.

27. The GAMES Collaborative Group. Linkage disequilibrium screening for multiple sclerosis

implicates JAG1 and POU2AF1 as susceptibility genes in Europeans. *J Neuroimmunol*.

2006;179(1–2):108–116.

28. Nakamura M, et al. Genome-wide Association Study Identifies TNFSF15 and POU2AF1 as

Susceptibility Loci for Primary Biliary Cirrhosis in the Japanese Population. *Am J Hum Genet*.

2012;91(4):721–728.

29. Sun W, et al. OCA-B/Pou2af1 is sufficient to promote CD4+ T cell memory and

prospectively identifies memory precursors. *Proc Natl Acad Sci U S A*.

2024;121(9):e2309153121.

30. Kim H, et al. Targeting transcriptional coregulator OCA-B/Pou2af1 blocks activated

autoreactive T cells in the pancreas and type 1 diabetes. *J Exp Med*. 2021;218(3):133.

31. Tzartos JS, et al. IL-21 and IL-21 Receptor Expression in Lymphocytes and Neurons in

Multiple Sclerosis Brain. *Am J Pathol*. 2011;178(2):794–802.

32. Ndhlovu LC, et al. Critical Involvement of OX40 Ligand Signals in the T Cell Priming Events

During Experimental Autoimmune Encephalomyelitis. *J Immunol*. 2001;167(5):2991–2999.

33. Hurwitz AA, et al. Cytotoxic T lymphocyte antigen-4 (CTLA-4) limits the expansion of

encephalitogenic T cells in experimental autoimmune encephalomyelitis (EAE)-resistant BALB/c

mice. *Proc Natl Acad Sci*. 2002;99(5):3013–3017.

34. Arellano G, et al. Stage-Specific Role of Interferon-Gamma in Experimental Autoimmune

Encephalomyelitis and Multiple Sclerosis. *Front Immunol*. 2015;6(SEP).

https://doi.org/10.3389/fimmu.2015.00492.

35. Rottman JB, et al. The costimulatory molecule ICOS plays an important role in the

immunopathogenesis of EAE. *Nat Immunol*. 2001;2(7):605–611.

36. Rouse M, Nagarkatti M, Nagarkatti PS. The role of IL-2 in the activation and expansion of

regulatory T-cells and the development of experimental autoimmune encephalomyelitis.

Immunobiology. 2013;218(4):674–682.

37. Duncker PC, et al. GM-CSF Promotes Chronic Disability in Experimental Autoimmune

Encephalomyelitis by Altering the Composition of Central Nervous System–Infiltrating Cells, but

Is Dispensable for Disease Induction. *J Immunol*. 2018;200(3):966–973.

38. Yosef N, et al. Dynamic regulatory network controlling TH17 cell differentiation. *Nature*.

2013;496(7446):461–468.

39. Ikegami I, et al. Bob1 enhances RORγt-mediated IL-17A expression in Th17 cells through

interaction with RORγt. *Biochem Biophys Res Commun*. 2019;514(4):1167–1171.

40. Baker D, et al. Autoimmune encephalomyelitis in NOD mice is not initially a progressive

multiple sclerosis model. *Ann Clin Transl Neurol*. 2019;6(8):acn3.792.

41. Macnair W et al. Single nuclei RNAseq stratifies multiple sclerosis patients into three distinct

white matter glia responses. *BioRxiv*. [published online ahead of print: 2022].

https://doi.org/https://doi.org/10.1101/2022.04.06.487263.

42. Rumble JM, et al. Neutrophil-related factors as biomarkers in EAE and MS. *J Exp Med*.

2015;212(1):23–35.

43. Kroenke MA, et al. IL-12– and IL-23–modulated T cells induce distinct types of EAE based

on histology, CNS chemokine profile, and response to cytokine inhibition. *J Exp Med*.

2008;205(7):1535–1541.

44. Grifka-Walk HM, Lalor SJ, Segal BM. Highly polarized Th17 cells induce EAE via a T-bet

independent mechanism. *Eur J Immunol*. 2013;43(11):2824–2831.

45. Schnell A, et al. Stem-like intestinal Th17 cells give rise to pathogenic effector T cells during

autoimmunity. *Cell*. 2021;184(26):6281-6298.e23.

46. La Manno G, et al. RNA velocity of single cells. *Nature*. 2018;560(7719):494–498.

47. Alpsoy A, et al. IκBζ is a dual-use coactivator of NF-κB and POU transcription factors. *Mol*

Cell. [published online ahead of print: February 24, 2024].

https://doi.org/10.1016/j.molcel.2024.01.007.

48. Wu XS, et al. OCA-T1 and OCA-T2 are coactivators of POU2F3 in the tuft cell lineage.

Nature. 2022;607(7917):169–175.

49. Schubart DB, et al. B-cell-specif ic coactivator OBF-1/OCA-B/Bob1 required for immune

response and germinal centre formation. *Nature*. 1996;383(6600):538–542.

50. Qin XF, et al. OCA-B integrates B cell antigen receptor-, CD40L- and IL 4-mediated signals

for the germinal center pathway of B cell development. *EMBO J*. 1998;17(17):5066–75.

51. Heng TSP, et al. The Immunological Genome Project: networks of gene expression in

immune cells. *Nat Immunol*. 2008;9(10):1091–1094.

52. Zwilling S, et al. Inducible Expression and Phosphorylation of Coactivator BOB.1/OBF.1 in T

Cells. *Science (80-)*. 1997;277(5323):221–225.

53. Gearty S V., et al. An autoimmune stem-like CD8 T cell population drives type 1 diabetes.

Nature. 2022;602(7895):156–161.

54. Li Y, et al. Stem-like T cells are associated with the pathogenesis of ulcerative colitis in

humans. *Nat Immunol*. [published online ahead of print: June 19, 2024].

https://doi.org/10.1038/s41590-024-01860-7.

55. Paroni M, et al. Recognition of viral and self-antigens by T H 1 and T H 1/T H 17 central

memory cells in patients with multiple sclerosis reveals distinct roles in immune surveillance and

relapses. *J Allergy Clin Immunol*. 2017;140(3):797–808.

56. Johns TG, et al. Myelin oligodendrocyte glycoprotein induces a demyelinating

encephalomyelitis resembling multiple sclerosis. *J Immunol*. 1995;154(10):5536–41.

57. Gao Y, et al. Transcription factor EGR2 controls homing and pathogenicity of TH17 cells in

the central nervous system. *Nat Immunol*. 2023;24(8):1331–1344.

58. McRae BL, et al. Induction of active and adoptive relapsing experimental autoimmune

encephalomyelitis (EAE) using an encephalitogenic epitope of proteolipid protein. *J*

Neuroimmunol. 1992;38(3):229–240.

59. Bernard CCA, et al. Myelin oligodendrocyte glycoprotein: a novel candidate autoantigen in

multiple sclerosis. *J Mol Med*. 1997;75(2):77–88.

60. Anderson AC, et al. A Transgenic Model of Central Nervous System Autoimmunity Mediated

by CD4+ and CD8+ T and B Cells. *J Immunol*. 2012;188(5):2084–2092.

61. Nish SA, et al. CD4+ T cell effector commitment coupled to self-renewal by asymmetric cell

divisions. *J Exp Med*. 2017;214(1):39–47.

62. Mammadli M, et al. TCF-1 Is Required for CD4 T Cell Persistence Functions during

AlloImmunity. *Int J Mol Sci*. 2023;24(5):4326.

63. Kadowaki A, et al. Gut microbiota-dependent CCR9+CD4+ T cells are altered in secondary

progressive multiple sclerosis. *Brain*. 2019;142(4):916–931.

64. Thakore PI, et al. BACH2 regulates diversification of regulatory and proinflammatory

chromatin states in TH17 cells. *Nat Immunol*. [published online ahead of print: July 15, 2024].

https://doi.org/10.1038/s41590-024-01901-1.

65. Song S, et al. OBF1 and Oct factors control the germinal center transcriptional program.

Blood. 2021;137(21):2920–2934.

66. Sha Y, Markovic-Plese S. Activated IL-1RI Signaling Pathway Induces Th17 Cell

Differentiation via Interferon Regulatory Factor 4 Signaling in Patients with Relapsing-Remitting

Multiple Sclerosis. *Front Immunol*. 2016;7. https://doi.org/10.3389/fimmu.2016.00543.

67. Lane TE, et al. A central role for CD4(+) T cells and RANTES in virus-induced central

nervous system inflammation and demyelination. *J Virol*. 2000;74(3):1415–24.

68. Blanc CA, et al. Sphingosine-1-Phosphate Receptor Antagonism Enhances Proliferation and

Migration of Engrafted Neural Progenitor Cells in a Model of Viral-Induced Demyelination. *Am J*

Pathol. 2015;185(10):2819–2832.

69. Plaisted WC, et al. Remyelination Is Correlated with Regulatory T Cell Induction Following

Human Embryoid Body-Derived Neural Precursor Cell Transplantation in a Viral Model of

Multiple Sclerosis. *PLoS One*. 2016;11(6):e0157620.

70. Dickey LL, et al. MicroRNA-155 enhances T cell trafficking and antiviral effector function in a

model of coronavirus-induced neurologic disease. *J Neuroinflammation*. 2016;13(1):240.